Dysregulation of Neuroprotective Lipoxin Pathway in Astrocytes in Response to Cytokines

and Ocular Hypertension

Authors: Shruthi Karnam[1], Shubham Maurya[1], Elainna Ng, Amodini Choudhary[1], Arzin

Thobani[1], John G Flanagan[1]*, Karsten Gronert[1,2]*

1. Herbert Wertheim School of Optometry and Vision Science, University of California Berkeley,

Berkeley, California, United States.

2. Infectious Disease and Immunity Program, Herbert Wertheim School of Optometry and Vision

Science, University of California Berkeley, CA, United States

*Corresponding Authors:

*Karsten Gronert, Herbert Wertheim School of Optometry and Vision Science, Minor Hall,

University of California Berkeley, Berkeley, CA 94720, USA. Email: kgronert@berkeley.edu

*John Flanagan, Herbert Wertheim School of Optometry and Vision Science, Minor Hall,

University of California Berkeley, Berkeley, CA 94720, USA. Email: jgflanagan@berkeley.edu

Abstract

Glaucoma leads to vision loss due to retinal ganglion cell death. Astrocyte reactivity contributes

to neurodegeneration. Our recent study found that lipoxin B_4 (LXB_4), produced by retinal

astrocytes, has direct neuroprotective actions on retinal ganglion cells. In this study, we aimed to

investigate how the autacoid LXB_4 influences astrocyte activity in the retina under inflammatory

cytokine-induced activation and during ocular hypertension. The protective activity of LXB_4 was

investigated *in vivo* using the mouse silicone-oil model of chronic ocular hypertension (n=40).

By employing a range of analytical techniques, including bulk RNA-seq, RNAscope in-situ

hybridization, qPCR, and lipidomic analyses, we discovered the formation of lipoxins and

expression of the lipoxin pathway in rodents (including the retina and optic nerve), primates

(optic nerve), and human brain astrocytes, indicating the presence of this neuroprotective

pathway across various species. Findings in the mouse retina identified significant dysregulation

of the lipoxin pathway in response to chronic ocular hypertension, leading to an increase in 5-

lipoxygenase (5-LOX) activity and a decrease in 15-LOX activity. This dysregulation was

coincident with a marked upregulation of astrocyte reactivity. Reactive human brain astrocytes

also showed a significant increase in 5-LOX. Treatment with LXB_4 amplified the lipoxin

biosynthetic pathway by restoring and amplifying the generation of another member of the

lipoxin family, LXA_4, and mitigated astrocyte reactivity in mouse retinas and human brain

astrocytes. In conclusion, the lipoxin pathway is functionally expressed in rodents, primates, and

human astrocytes, and is a resident neuroprotective pathway that is downregulated in reactive

astrocytes. Novel cellular targets for LXB_4's neuroprotective action are inhibition of astrocyte

reactivity and restoration of lipoxin generation. Amplifying the lipoxin pathway is a potential

target to disrupt or prevent astrocyte reactivity in neurodegenerative diseases, including retinal

ganglion cell death in glaucoma.

Keywords: Glaucoma, astrocyte reactivity, lipoxygenase, lipoxin B_4, neurodegeneration, retina,

optic nerve

Introduction

Glaucoma is a progressive neurodegenerative disease characterized by the degeneration of retinal

ganglion cells (RGCs) and their axons [34, 44]. It is a leading cause of irreversible blindness

worldwide, and it is estimated that the number of people affected by glaucoma will increase to

112 million worldwide by 2040 [24]. Although intraocular pressure (IOP) is the only modifiable

risk factor, most glaucoma patients experience continued but reduced progression. Proposed

reasons for this continued degeneration include noncompliance with medication, insufficient IOP

reduction, and non-IOP-related causes [6, 15, 21]. There is an unresolved need to develop

neuroprotective strategies to prevent or delay the death of RGCs, independent of IOP, in

glaucoma [20].

61   Lipoxins (LXA_4 and LXB_4) are eicosanoids generated from arachidonic acid. Their

canonical protective actions in various pathological processes include regulating the initial

response of innate and adaptive immune cells and promoting the resolution of acute

inflammation [32, 38, 52]. We recently identified a noncanonical protective action of lipoxins,

namely their direct regulation of RGCs [30]. The human and mouse receptor for LXA_4, formyl

peptide receptor 2 (FPR2/ALX) [39], is expressed in RGCs. A receptor for LXB_4, the lipoxin

with more potent neuroprotective actions in the retina [30], has yet to be identified. The

biosynthesis of lipoxins requires the coordinated interaction of 5-lipoxygenase (LOX) with

12/15-LOX enzymes [13, 49] (An overview of lipoxin biosynthesis is included in supplementary

Fig. 1). Primary astrocytes generate lipoxins *in vitro* and acute exocytotoxic stress downregulates

lipoxin formation and lipoxygenase expression in the retina [30]. However, the regulation of the

lipoxin pathway in ocular hypertension has yet to be investigated. More importantly, it is unclear

whether retinal astrocyte reactivity, a primary response to ocular hypertension, regulates the

astrocyte lipoxin pathway and whether lipoxin formation is a unique phenotype of mouse retinal

astrocytes. These knowledge gaps are of interest as therapeutic amplification of the retinal

lipoxin pathway is a potential neuroprotective target.

Astrocytes are glial cells in the central nervous system and retina and play a crucial role

in maintaining homeostasis. They provide metabolic substrates, growth factors, and trophic

support to neurons and are also involved in synapse formation and plasticity [42]. Astrocytes also

work closely with the immune system, as they can detect danger signals, respond to secreted

cytokines and chemokines, and activate adaptive immune responses [8]. In rodent models of

ocular hypertension, astrocyte reactivity in the retina and optic nerve head is a key feature of the

retinal injury response [27, 46]. There is also an initial rise in astrocyte reactivity within the

distal optic nerve during the degeneration process [43]. Reactive astrocytes can release

proinflammatory molecules and oxidative stress-inducing factors, which initiate or amplify

neuroinflammation and neuronal damage [9]. Recently, it has been suggested that microglial

release of proinflammatory cytokines can induce the formation of neurotoxic reactive astrocytes

in the retina and brain [27].

In this study, we aimed to investigate the regulation and functional expression of lipoxin

pathways in the mouse retina, optic nerves of mice and primates, and human astrocytes. We also

examined whether the neuroprotective action of the autacoid LXB_4 includes reducing astrocyte

reactivity in response to inflammatory cytokine-induced activation and ocular hypertension.

Materials and Methods

Mice

C57BL/6J male mice (Jackson Laboratory, Bar Harbor, ME) were allowed to acclimatize to the

housing facility for 2 weeks before experimentation. Mouse studies were performed in

compliance with the ARVO Statement for the Use of Animals in Ophthalmic and Vision

Research, and all experimental procedures were approved by the Institutional Animal Care and

Use Committee (IACUC) at the University of California, Berkeley. Mice were maintained in a

pathogen-free vivarium under a 12-hour dark and light cycle with *ad libitum* food and water.

Macaque optic nerves

The macaque optic nerves used for the lipidomic study were from the eyes of a 9-year and 306-

day-old female from the Oregon National Primate Research Center (NPRC). For the RNAseq

studies, optic nerves were from a 12-year-old male acquired at UC Berkeley. In both cases, the

eyes were collected postmortem for approved studies by different research groups, and the optic

nerves were donated to the Gronert/Flanagan Lab as discarded tissue.

Induction of ocular hypertension by injection of silicone oil

Ocular hypertension was induced in the 8-week-old C57BL/6J mice by injection of

pharmaceutical-grade silicon oil (Alcon silikon 1000, Geneva, Switzerland) in the anterior

chamber of both eyes, a model of chronic ocular hypertension that was established by Jie Zhang

and colleagues [11, 50, 51]. For these experiments the volume of silicone oil was reduced to 1.2

µL to enable a slower onset of mild ocular hypertension induced retinal injury. Briefly, 8-week-

old mice were anesthetized by an intraperitoneal injection of ketamine/xylazine (100 mg/kg and

10 mg/kg, respectively). One drop of topical anesthetic (proparacaine hydrochloride, 0.5%,

Akorn, Lake Forest, IL) was applied to the cornea of both eyes before the injection. Silicone oil

(1.2 µL) was injected into the anterior chamber of both eyes through a 30 G Hamilton glass

syringe. After injection, a drop of antibiotic (tobramycin ophthalmic solution 0.3%) was placed

on the eye. Sterile PBS (phosphate-buffered saline) was injected into another set of mice and

used as a control. Mice were euthanized at 2, 4, and 8-week time points following injections for

various experimental procedures.

IOP Measurements/Pupil Dilation

IOP was monitored once every 2 weeks until 8 weeks after silicon oil injection using the

TonoLab tonometer (Colonial Medical Supply, Londonderry, NH). In this model, the pupil needs

to be dilated sufficiently to expose it beyond the coverage of the silicone oil droplet, which leads

to the reconnection of the front and back chambers of the eye [11]. Each measurement of

intraocular pressure (IOP) can be seen as a form of pupil dilation treatment. The dilation of the

pupil was accomplished following the protocol outlined in the published literature [11, 50, 51].

Cell Culture

Cultured primary human cortical astrocytes (HA1800) were purchased from ScienCell (Carlsbad,

CA) and treated with the cytokines: IL-1α, TNF- α, and C1q for 24 hours [19]. Astrocytes were

treated with LXB_4 (Cayman Chemicals, Ann Arbor, MI) (1 µM) or vehicle for 20 mins before

adding cytokines. After 24 hours, the cells were collected in TRIzol (Invitrogen, Waltham, MA)

for RNA isolation or ice-cold methanol for lipidomic analysis. For primary retinal astrocytes and

optic nerve astrocytes, four mouse retinas were pooled and digested using papain (Worthington,

Lakewood, NJ) and plated on poly-D-lysine coated flasks with astrocyte growth media (Lonza,

Basel, Switzerland), changing the media every 3 days. Over 95% of the cultured cells were

identified as astrocytes through immunofluorescence staining for GFAP and vimentin and were

negative for RGCs, microglia, and oligodendrocyte markers.

Bulk RNA-seq

Total RNA from macaque optic nerves (n=2) was extracted using TRIzol extraction. Sequencing

libraries were prepared using the SMARTer v4 Ultra Low Input RNA Kit (Clontech, Mountain

View, CA). Sequencing was performed on NovaSeq S4 150PE platform (Illumina, San Diego,

CA). Library preparation and RNA-seq were performed at QB3 Genomics, UC Berkeley,

Berkeley, CA, RRID: SCR_022170. A quality check of FASTQ files was performed with FastQC

v.0.11.9, and adapter sequences were trimmed using Trim Galore! v.0.6.6. Trimmed reads were

mapped on mouse genome version mm39 (University of California Santa Cruz) with the STAR

alignment tool v.2.7.1a using RefGene annotation. The total number of read counts mapped for

each gene was extracted from BAM files using FeatureCount v.1.5.3. The count matrix was

imported into Rstudio v.4.2.0. It was normalized using the Transcripts Per Million method by

NormalizeTPM. Mean values of normalized counts for each gene were plotted for visualization.

Electroretinography

Before ERG, the mice underwent overnight dark adaptation. They were then anesthetized and

received tropicamide (1%, Akorn, Lake Forest, IL) and phenylephrine (2.5%, Paragon BioTeck,

Portland, OR) to dilate their pupils. The Celeris system (Diagnosys LLC, Lowell, MA) was used

to conduct the ERG measurement, with stimulus intensities ranging from -5.90 to 2.25 log

cd.s/m^2. The tail subdermal needle electrode served as ground. A positive scotopic threshold

response (pSTR) elicited with an intensity of -2.50 log cd.s/m^2 was used to represent RGC

function (average of 20 repeats, interstimulus interval of 2 seconds).

Optical Coherence Tomography

Before imaging, mice received anesthesia and were given lubricant eye drops. We used a

Bioptigen spectral domain OCT system (Envisu R2300, Durham, NC) for imaging. Our previous

studies described the OCT image acquisition and analysis protocols [28, 29]. We captured an en-

face retinal fundus image with the optic nerve head in the center, using 1.8 x 1.8 mm rectangular

scanning. Each replication involved 100 B-scan images (1536 A-scans for each B-scan). Retinal

layer B-scan images were analyzed by masked observers using ImageJ software. We quantified

the thickness of the retinal layer at both left and right locations from the center of the optic nerve

head in each retinal quadrant. The average data from both locations represented the thickness

value.

Retinal Flat Mount and RGC Quantification

Eyes were removed at 8 weeks and fixed with 4% paraformaldehyde (PFA). Posterior eyecups

were incubated overnight in blocking buffer (10% normal donkey serum and 2% PBS-Triton-X-

100). Eyecups were stained with primary antibodies, GFAP (Abcam, Cambridge, UK), and

RBPMS (Phosphosolutions, Aurora, CO) (1:1000 dilution each) and were incubated on a shaker

for 48-72 hours at 4 °C. After incubation, eyecups were washed with 0.25% PBST and further

incubated with either Alexa Fluor 488- or 594- conjugated secondary antibodies (Invitrogen; at

1:1000 dilution) under dark conditions for 2 hours at room temperature. After washing with

PBST, the retinas were gently detached, and four perpendicular clover cuts were made.

ProLong™ Gold Antifade Mountant (#P36930, Invitrogen, Eugene, OR) was added to mount the

retinas. Retinas were imaged on a Nikon Ti Eclipse microscope using a 20x objective. A

minimum of 5-6 independent specimens (biological replicates) were analyzed for each flat

mount. There was no observed RGC loss in the central region. To count peripheral RGCs, eight

fields were sampled from the peripheral areas of each retina using a 20x lens on a Nikon Ti

Eclipse microscope. (Supplementary Figure 2 displays a flat-mounted retina that defines the

central and peripheral regions). RGCs were counted using ImageJ software. The percentage of

RGC survival was determined by calculating the ratio of surviving RGC numbers in the injured

eyes compared to the control eyes.

Immunohistochemistry

The eyes were fixed in 4% PFA and transferred into 30% sucrose. Tissues were then embedded

in OCT media and cut into 10-μm-thick sections using a CM1900 Cryostat (Leica, Wetzlar,

Germany). Primary human astrocytes were fixed with 4% PFA for 20 mins. The cells were

permeabilized with 0.3% Triton®-X-100 PBS for 5 mins at room temperature. Sections and cells

were blocked using a blocking buffer (5% donkey serum in 0.3% Triton X-100 containing PBS)

for 30 mins. Primary antibodies against GFAP (Abcam), RBPMS (Phosphosolutions), LCN2

(Invitrogen), 5-LOX (Abcam), Pax2 (Abcam), Vimentin (Invitrogen), and CD44 (Proteintech)

were added to blocking buffer and incubated overnight at 4 °C. Sections and cells were washed

and incubated with Alexa Fluor 488- and 594- conjugated secondary antibodies (Invitrogen; at

1:1000 dilution) for double labeling for 2 hours at room temperature under dark conditions. They

were then washed and mounted using ProLong™ Gold Antifade Mountant, and images were

captured using a Nikon Ti Eclipse microscope.

RNAscope in situ hybridization

RNA in situ hybridization for *Alox5*, *Alox15*, and *Fpr2* mRNA was performed manually using

RNAscope Multiplex Fluorescent Reagent Kit v2 Assay (Advanced Cell Diagnostics, Inc.,

Newark, CA) according to the manufacturer's instructions. Briefly, 10 μm of PFA fixed frozen

retina and optic nerve sections were pretreated with heat and protease (30 mins, 40 °C) before

hybridization with the target probes. Tissue was incubated in primary amplification probes and

Opal^{TM} 690 dye fluorophore (2-hour primary probe, 30 mins AMP1, 30 mins AMP2, 15 mins

AMP3, and 30 mins fluorophore at 40 °C) and washed between steps with RNAscope washing

buffer. Each sample was quality controlled for RNA integrity with an RNAscope® probe specific

to PPIB RNA and for background with a probe specific to bacterial dapB RNA. Specific RNA

staining signals were identified as red, punctate dots. The tissue was counterstained with DAPI.

After mounting in ProLong™ Gold Antifade Mountant, images were acquired on a Nikon Ti

Eclipse microscope using a 40x objective. The RNAscope probes used were as follows: *Alox15*

(Cat. No: 539781-C1), *Alox5* (Cat. No: 436101-C2), and *Fpr2* (Cat. No: 509281-C3).

qPCR

RNA was isolated from mouse retinal tissue and human brain astrocytes and quantified

according to our published protocol [30]. *Gapdh* was used as the reference gene. The mouse

primers used in this study are listed in Table. Primers for the human genes used in this study are

listed in Table 2 (please refer to supplementary material for primer details).

Liquid Chromatography-tandem mass spectrometry (LC☐MS/MS)

Mouse retinas and macaque optic nerves were immediately snapped, frozen on dry ice, and

stored at -80 °C before processing for lipidomic analyses. Eicosanoids and PUFAs in the cells

and conditioned media from astrocyte cultures were identified and quantified by LC☐MS/MS

according to our published protocol [30]. Briefly, deuterated internal standards (15-HETE-d_8,

LXA$_4$-d_5, and AA-d_8) were added to all samples before processing to calculate class-specific

recoveries. Frozen tissues were placed in MeOH and processed in a refrigerated beat

homogenizer. Supernatants were extracted using C18 solid-phase columns. Extracted lipids were

analyzed by LC☐MS/MS using an AB SCIEX 4500 QTRAP mass spectrometer. Analysis was

carried out in negative ion mode, and eicosanoids and PUFAs were quantitated using scheduled

multiple reaction monitoring (MRM) using from 3 to 4 specific transition ions for each analyte.

The peak identification and integration criterion was a signal-to-noise ratio above 5:1. For

quantification, calibration curves and HPLC retention times for each analyte were established

with synthetic standards (Cayman Chemicals).

Statistical Analysis

All values are presented as the mean ± standard error of the mean (SEM) of n observations,

where n represents the number of animals studied or independent *in vitro* experiments. Statistical

analysis was performed using GraphPad Prism 9.0 (GraphPad Software, San Diego, CA). One-

way ANOVA followed by post hoc Tukey's multiple comparison test or unpaired Student's t-test

was used to compare intergroup differences. The results were considered statistically significant

when $P<0.05$.

Results

The lipoxin pathway is expressed in mouse retina, optic nerve, and retinal astrocytes.

The expression of lipoxin pathways was investigated in the retina, optic nerve, and retinal

astrocytes of C57BL/6J mice. The genes in question are known to be expressed at levels that

cannot be detected using single-cell transcriptomics, a known limitation of this method.

Additionally, since there are no specific antibodies for detecting lipoxin pathways in mouse

retina and optic nerve, we used RNAscope and qPCR for our investigations. RNAscope in situ

hybridization was performed on the retina and optic nerve region cross-sections to determine the

cell-specific expression of *Alox5, Alox15, and Fpr2*. The data showed that the lipoxin circuits

were localized in the ganglion cell layer (GCL), inner nuclear layer (INL), and outer nuclear

layer (ONL) of the retina, and in the optic nerve region (Fig. 1A). The quality of the RNA was

confirmed using positive and negative controls, as illustrated in Supplementary Figure 3. To

confirm the expression of the lipoxin circuits in retinal astrocytes, qPCR was performed on

cultured primary mouse retinal astrocytes. The results showed that *Alox5, Alox15, and Fpr2* were

expressed with average Ct values of 28.23±0.3, 29.07±0.3, and 28.58±0.2, respectively (n=3).

Notably, higher RNA expression was observed in the optic nerves with average Ct values of

27.75±0.2 (Alox5), 24.65±0.6 (Alox15), and 25.94±0.7 (Fpr2) (n=3) (Fig. 1B). These results

confirm expression of the lipoxin pathway in the retina and optic nerve of mice, including retinal

astrocytes.

Lipoxin pathway is expressed in macaque optic nerves and human primary brain

astrocytes.

To explore expression of the lipoxin pathway in primates, we conducted RNA-seq analyses on

macaque optic nerves. Results established high expression of *ALOX15*, moderate expression of

ALOX5 and *ALOX5AP*, and lower expression of *ALOX12, ALOX15B, FPR2,* and *FPR3* genes

(Fig. 2A). Consistent with gene expression, lipidomic analysis confirmed active eicosanoid

pathways and lipoxin formation in primate optic nerves as evidenced by the presence of released

arachidonic acid and significant levels of 5-HETE, 12-HETE, 15-HETE and LXB$_4$ (Fig. 2B).

The cell identity of the primary human brain astrocyte line was validated by positive staining

with astrocyte-specific markers GFAP, Vimentin, Pax2, and CD44 (Fig. 2C). Primary human

astrocytes, like primate optic nerves, also expressed the lipoxin biosynthetic pathway, namely

ALOX5 and *ALOX15,* and the LXA$_4$ receptor *FPR2*, and lipidomic analyses confirmed

endogenous formation of eicosanoids and lipoxins. Interestingly, cultured human brain astrocytes

generated both LXA$_4$ and LXB$_4$ (Fig. 2D). These data demonstrate that neuroprotective lipoxins

are formed within non-human primate optic nerves and can be generated by primary human brain

astrocytes.

Validation of the mild chronic model of ocular hypertension

IOP and RGC density changes were assessed in control and silicone oil-injected eyes of mice.

Fig. 3A shows that IOP was significantly elevated in the silicone oil-injected eyes compared to

control eyes. After 8 weeks mice were euthanized, and RGC density was assessed by staining

flat-mounted retinas with the RGC marker, RBPMS (Fig. 3B). Peripheral RGC loss of 17% was

observed in silicon oil-injected eyes compared to control eyes (Fig. 3C). Changes in RGC

function were measured longitudinally using ERG measurements of pSTR. The average

waveforms of pSTR were elicited by a flash intensity of -2.5 log cd.s/m^2 at 8 weeks (Fig. 3D).

The results indicated a loss of RGC function in hypertensive eyes compared to control

normotensive eyes. Additionally, loss in ganglion cell complex (GCC) thickness, a clinically

relevant biomarker of nerve fiber loss, was measured longitudinally by OCT (Fig. 3E). Mild

ocular hypertension caused a significant 10% relative loss ($p < 0.05$) compared to normotensive

control eyes (Fig. 3F). Overall, these results confirm that the injection of 1.2 µl of silicone oil

established mild, chronic ocular hypertension and mild damage to the RGCs after 8 weeks. The

sustained elevated intraocular pressure (IOP) resulted in mild morphological and functional loss

of retinal ganglion cells (RGCs).

Dysregulation of lipoxin pathway in response to chronic mild ocular hypertension

Lipoxin pathway gene expression was analyzed in retinas from normotensive and hypertensive

eyes at 2, 4, and 8 weeks to investigate the regulation of lipoxin pathway in response to chronic

mild ocular hypertension. The results demonstrate that *Alox5AP* (5-lipoxygenase activating

protein, FLAP), which is a cofactor required for full 5-LOX enzymatic activity, was significantly

upregulated at 2 and 4 weeks, while there was no change in the levels of *Alox5* in response to

ocular hypertension. Interestingly, the levels of *Alox15* and *Fpr2* were significantly upregulated

initially at 2 and 4 weeks but significantly downregulated at 8 weeks (Fig. 4A). To investigate

the functional dysregulation of the lipoxin pathway, four retinas were pooled at 4 weeks and

analyzed by LC☐MS/MS-based lipidomics to establish formation and changes in eicosanoid and

PUFA profiles. Lipidomic analysis demonstrated that arachidonic acid (AA) and the 15-LOX

pathway marker, 15-HETE, were significantly decreased in response to ocular hypertension. In

contrast, the 5-LOX pathway markers, 5-HETE, and 4-HDHA were significantly increased (Fig.

4B). These findings indicate that expression of the homeostatic lipoxin pathway is regulated

during ocular hypertension and that activity of 5-LOX and 15-LOX is dysregulated.

Activation of Müller glia and astrocyte reactivity markers in response to ocular

hypertension

The impact of mild ocular hypertension on astrocyte and Müller glial reactivity was assessed by

immunofluorescence staining of retinal cross-sections with cell-specific reactivity markers from

normotensive and hypertensive eyes. Expression of GFAP, an intermediate filament protein that

is a marker of macroglial reactivity, was induced and upregulated in Müller glia and astrocytes,

as early as 2 weeks after ocular hypertension (Fig. 5A) and remained upregulated in the retina for

8 weeks and was coincident with loss of RGCs as evidenced by RBPMS staining in the GCL

layer (Fig. 5B). GFAP staining of retinal flat mounts also established marked reactive astrogliosis

after 8 weeks of mild ocular hypertension (Fig. 5C). To investigate macroglia reactivity further,

we measured glutamate aspartate transporter (GLAST) gene expression, which is highly

expressed in Müller cells as part of their homeostatic function to support RGCs. GLAST

expression was significantly decreased after 2 and 4 weeks of ocular hypertension (Fig. 5D).

This dysregulation has also been previously observed in rat models of optic nerve injury [33].

Gene expression of signature astrocyte reactivity markers (*Gfap, Lcn2, Steap4, Gbp2, Serping1,*

and Fbln5) were significantly upregulated in response to ocular hypertension at 2, 4, and 8

weeks compared to that in normotensive controls (Fig. 5E). These findings confirm that the early

occurrence of Müller glia and astrocyte reactivity is a characteristic feature in this mouse model

of mild ocular hypertension-induced retinal injury.

Therapeutic LXB$_4$ treatment reduces astrocyte and Muller glial reactivity in the mouse

retina

We next assessed whether the neuroprotective actions of lipoxins in ocular hypertension [30]

include regulation of macroglia reactivity. Therapeutic LXB$_4$ methyl ester treatment (n=8, 5 ng/g

IP; 0.5 ng/g topical) was initiated after 1 week of ocular hypertension and administered every

other day until the 4th week. The dosing and preparation of lipoxins were based on previous

studies [5, 30]. qPCR analysis of markers of macroglia reactivity established that treatment with

LXB$_4$ resulted in a significant upregulation of GLAST. In addition, LXB$_4$ treatment significantly

downregulated astrocyte reactivity markers such as *Lcn2* and *Steap4*. The levels of *Gbp2*,

Serping1, and Fbln5 remained unchanged (Fig. 6). These findings identify a potentially new

mechanism for the neuroprotective actions of LXB$_4$ treatment, namely reversing macroglia

reactivity and restoring their homeostatic function.

LXB$_4$ treatment modulates the lipoxin pathway in primary human brain astrocytes

To establish macroglia-specific actions for LXB$_4$, we used primary human brain astrocytes that

express a fully functional lipoxin pathway identical to that found in the mouse primary retinal

astrocytes (Fig. 2D). To induce astrocyte reactivity, cells were treated with established

inflammatory cytokines, which are also generated in the retina in response to ocular hypertension

[19]. qPCR analysis confirmed that cytokine treatment significantly upregulated reactive

astrocyte markers, including *LCN2, SERPING1,* and *C3*, in primary human brain astrocytes.

LXB$_4$ treatment significantly downregulated these astrocyte reactivity markers (Fig. 7A).

Immunofluorescence analysis of LCN2 was consistent with the qPCR findings. These results

establish direct regulation of astrocyte reactivity as a new bioaction of LXB$_4$.

368   Reactive astrocytes also showed increased gene expression of 5-LOX (ALOX5)

compared to controls. LXB$_4$ treatment reduced 5-LOX expression in reactive astrocytes (Figs.

7A and B). Hence, we examined the functional consequence of LXB$_4$ regulation of 5-LOX

expression in reactive astrocytes. Consistent with changes in 5-LOX gene expression (Fig 7B),

5-HETE formation was significantly increased in reactive astrocytes, and LXB$_4$ treatment

reduced 5-HETE generation. More importantly, reactive astrocytes generated less LXA$_4$, but

LXB$_4$ treatment restored and amplified LXA$_4$ formation in reactive astrocytes (Fig. 7C).

Concentrations of added LXB$_4$ in media remained high after 24 hours, so we were unable to

assess whether LXB$_4$ treatment also amplified endogenous astrocyte LXB$_4$ formation. These

findings suggest LXB$_4$ restores the homeostatic phenotype, and both recovers and amplifies

neuroprotective lipoxin formation in cytokine-challenged primary human astrocytes.

Discussion

Glaucoma results in the gradual loss of retinal ganglion cells (RGCs), leading to irreversible

vision loss [18]. Astrocyte reactivity is a key pathological mechanism that contributes to the

neurodegeneration of RGCs in glaucoma, as astrocytes play a crucial role in maintaining the

health and function of RGCs [2]. Recently, we identified that lipoxins (LXA$_4$ and LXB$_4$) derived

from astrocytes promote neuroprotection against acute and chronic injury in the retina by directly

acting on RGCs [30]. However, the regulation of lipoxin formation in the retina and the extent of

its neuroprotective activity in glaucoma remains to be defined.

Our study revealed that astrocytes express 5-LOX and 15-LOX, two enzymes essential

for lipoxin generation by a single cell [13, 48, 49]. We assessed the functional activity of the

lipoxin pathway through LC☐MS/MS-based lipidomics and confirmed the presence of

endogenous lipoxin formation and activity of 5-LOX and 15-LOX in rodent and primate

astrocytes, retina, and optic nerves. Notably, we observed lipoxin formation in the optic nerves of

nonhuman primates, which, to our knowledge, has not been previously reported. Additionally,

we demonstrated formation of both LXA_4 and LXB_4 by primary human brain astrocytes, aligning

with a recent study reporting lipoxin formation in astrocytes of human brain organoids using

ELISA [35]. Astrocytes, the most abundant cell type in the central nervous system, play vital

regulatory roles in neurogenesis, synaptogenesis, blood☐brain barrier permeability, and

extracellular homeostasis, impacting various aspects of brain function [41]. These results

emphasize the potential significance of the resident neuroprotective lipoxin pathway not only in

the retina but also in the brain and optic nerve [25, 30].

5-LOX is required to generate leukotrienes, which are proinflammatory lipid mediators,

and this enzyme has been implicated in promoting proinflammatory lipid mediator synthesis in

various neurodegenerative diseases [36]. However, 5-LOX is also required for generating

lipoxins and most of the proresolving lipid mediators (SPMs) [25, 38, 49]. Hence, the role of 5-

LOX is context- and cell-dependent. In the retina, 5-LOX has been identified as the essential

enzyme for the protective actions of dietary DHA supplementation in retinopathy of prematurity

[37]. Moreover, in 5-LOX knockout mice, autoimmune uveitis pathogenesis is markedly

amplified, and in a retinal model of glutamate excitotoxicity, 5-LOX inhibition amplifies retinal

ganglion cell death [30]. These findings indicate that in the retina, homeostatic 5-LOX is

protective.

The function of 5-LOX, namely the products this pathway generates, depends on the

balanced interaction with 15-LOX, an interaction that leads to lipoxin formation. 15-LOX is

highly expressed in epithelial cells of healthy mucosal tissues, retinal pigmented epithelial cells,

tissue-resident homeostatic macrophages, immune regulatory macrophages and

polymorphonuclear leukocytes (PMNs) [3, 14, 16, 47]. 15-LOX has immunoregulatory and

wound healing roles important for maintaining tissue homeostasis, especially in the eye [3, 14,

17, 47]. A key feature of an immune-regulatory and pro-resolving phenotype of PMN is high

expression of 15-LOX to enable generation of lipoxins [14, 26]. Dysregulation of the balanced

expression and function of 5- and 15-lipoxygenases has been implicated in several

neurodegenerative diseases [25] and deletion of either 5-LOX or 15-LOX amplifies experimental

allergic encephalomyelitis in mice [10]. Our findings reveal a shift in the homeostatic balance of

5-LOX and 15-LOX activities in the retina and astrocytes as a feature of reactivity and ocular

hypertension-induced pathogenesis. Astrocyte reactivity and retinal response to ocular

hypertension correlated with an increase in 5-LOX and a decrease in 15-LOX activity.

Interestingly, our findings of dysregulated lipoxygenase activity are consistent with previous

studies that have reported increased levels of 5-LOX activity in Alzheimer's disease (AD) [7, 12,

22]. Additionally, a separate study reported decreased levels of 15-LOX in the CA1 region of the

hippocampus in patients with moderate to severe stages of AD [31]. In contrast, several studies

have demonstrated that inhibition of mouse 12/15-LOX enzymes can reduce ischemic injury and

diabetic pathogenesis [1, 45]. In short, there is a complex interplay between 5- and 15- LOX, and

their regulation and dysregulation are a key feature of neurodegenerative diseases such as

Alzheimer's disease and glaucoma.

Our study identified a novel target for the neuroprotective effects of LXB_4, demonstrating

its ability to inhibit astrocyte reactivity both *in vitro* and *in vivo*. Astrocytes play a crucial role in

maintaining the health and function of RGCs and the vasculature in the retina and optic nerve.

Astrocyte reactivity has been implicated as a key pathological feature in various

neurodegenerative diseases, including glaucoma [4, 23]. Our findings establish, for the first time,

a direct action of LXB_4 on macroglia by downregulating key reactive markers in primary human

astrocytes *in vitro*. This finding highlights the significance of LXB_4 as an autocrine signal in

astrocytes, similar to its established role in the PMN, the primary cell type that generates

lipoxins, but also the primary cell target to downregulate their proinflammatory function during

acute inflammation [14, 40, 49]. Thus, our results suggest that LXB_4 serves as an autocrine

signal in astrocytes, downregulating their reactive phenotype.

Previous studies have demonstrated the anti-inflammatory effects of lipoxins in various

neuroinflammatory diseases, indicating their therapeutic potential [52]. However, the source of

lipoxins in the central nervous system and the potential targets for their protective actions have

not been thoroughly investigated. Our study reveals that LXB_4 treatment restores and

significantly amplifies the generation of LXA_4, another lipoxin with distinct bioactions, in

reactive astrocytes. The specific receptor for LXB_4, distinct from the LXA_4 receptor Fpr2/ALX,

remains to be identified, as does the mechanism underlying the upregulation of lipoxin formation

in astrocytes. LXB_4 treatment reduced amplified 5-LOX activity in reactive human astrocytes,

suggesting that restoring the balanced interaction between 5-LOX and 15-LOX may partly

explain the restoration of lipoxin production in astrocytes.

Conclusions: The lipoxin pathway is functionally expressed in retina and brain astrocytes, and

optic nerves of rodents and primates, and is a resident neuroprotective pathway that is

downregulated in reactive astrocytes. Novel cellular targets for LXB_4's neuroprotective action is

inhibition of astrocyte reactivity and restoration of lipoxin generation. Amplifying the lipoxin

pathway is a potential target to disrupt or prevent astrocyte reactivity in neurodegenerative

diseases.

Abbreviations: LXB_4- Lipoxin B_4, LXA_4- Lipoxin A_4, IOP- Intraocular pressure, RGCs- Retinal

ganglion cells, LOX- Lipoxygenase, ERG- Electroretinogram, OCT- Optical coherence

tomography, LC-MS/MS- liquid chromatography-tandem mass spectrometry, AA- Arachidonic

acid, PUFA- Polyunsaturated fatty acids, HPLC- High-performance liquid chromatography,

ONL- Outer nuclear layer, INL- Inner nuclear layer, GCL- Ganglion cell layer, GCC- Ganglion

cell complex, SPM- Specialized pro-resolving lipid mediators, PMN- Poly morphonuclear

leukocytes.

Declarations:

Ethics approval and consent to participate

Animal studies were performed in compliance with the ARVO Statement for the Use of Animals

in Ophthalmic and Vision Research, and all experimental procedures were approved by the

Institutional Animal Care and Use Committee (IACUC) at the University of California, Berkeley.

Consent for publication

Not applicable

Availability of data and material

The RNA sequencing data from the macaque study conducted in this research has been deposited

in the GEO under the accession code GSE241131.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was partly supported by the NIH grants R01EY030218 (KG, JGF) and P30EY003176.

Acknowledgments

We thank Dr. Teresa Puthussery (UC Berkeley), Dr. Joni D. Wallis (UC Berkeley) and the tissue

distribution program at Oregon National Primate Research Center (Grant

number: P51OD011092) for providing post-mortem macaque optic nerve samples.

Authors contribution

SK: Conceptualization, Methodology, Validation, Investigation, Formal analysis, Data Curation,

Writing- Original Draft, Visualization; SM: Methodology, Validation, Investigation, Formal

analysis, Data Curation; EN, AC, AT: performed lipidomic analyses; JF, KG: Conceptualization,

Supervision, Resources, Writing-Review & Editing, Funding Acquisition. All authors read and

approved the final manuscript.

Figure legends

Figure 1. Expression of lipoxin pathway in mouse retina and optic nerve. (A) Representative

images of RNAscope ISH (red) (indicated with white arrows) of retina and optic nerve using

mRNA probes for *Alox5*, *Alox15*, and *Fpr2*. Nuclei were labeled with DAPI to distinguish

nuclear layers of the retina. All images were obtained using a 40x objective. Scale bar = 50 µm.

(n=3). Labels: ganglion cell layer (GCL), inner nuclear layer (INL), outer nuclear layer (ONL).

(B) Bar graph represents Ct values of lipoxin pathway genes in retinal astrocytes and optic

nerves analyzed by qPCR (n=3).

Figure 2. Expression of lipoxin pathway in optic nerve of primates and primary human

brain astrocytes. (A) RNA-seq of the macaque optic nerve showing lipoxin pathway genes

(n=2). **(B)** LC/MS/MS-based lipidomic quantification of endogenous lipid mediators in primate

optic nerve (n=2). **(C)** Representative immunofluorescence images of primary human brain

astrocytes stained with GFAP, Pax2, Vimentin, and CD44. **(D)** LC–MS/MS-based lipidomic

quantification of endogenous lipid mediators in human brain astrocytes (n=2).

Figure 3. Validation of the silicone oil model. (A) IOPs were increased after silicone oil (SO)

injection (n=20). **(B)** Representative confocal images of flat-mounted retina sections showing

surviving RBPMS-positive (red) RGCs at 8 weeks. Scale bar = 50 µm. **(C)** Bar graph indicates

% of peripheral RGC loss. **(D)** Average ERG waveforms of RGC responses (pSTR, -2.5 log

cd.s.m^{-2}) at 8 weeks for each group. **(E)** Representative OCT images of mouse retina at 8 weeks.

GCC: ganglion cell complex, including RNFL, GCL, and IPL layers, indicated by double-ended

yellow arrows. **(F)** Bar graph indicating the % reduction in GCC thickness at 8 weeks. (n=4-6).

****p< 0.001. (Ctrl- Control; OHT- ocular hypertension).

Figure 4. Regulation of lipoxin pathway in response to ocular hypertension. (A) The time

course response of lipoxin pathway genes was analyzed by qPCR at 2, 4, and 8 weeks. The

results were normalized to GAPDH and age-matched controls. n=3-4 (2 retinas were pooled for

each replicate). **(B)** LC/MS/MS-based lipidomic quantification of endogenous lipid mediators in

the mouse retina at 4 weeks, n=3 (4 retinas were pooled for each replicate). *p< 0.05, **p<0.01,

**p<0.001. (Ctrl- Control; OHT- ocular hypertension).

Figure 5. Regulation of astrocyte reactivity in response to ocular hypertension. (A)

Representative immunofluorescence images of GFAP (green) at 2 weeks, and **(B)** GFAP (green)

RBPMS (red) at 8 weeks of ocular hypertension. Nuclei were labeled with DAPI. All images

were obtained using a 40x objective. Scale bar = 50 µm. Labels: ganglion cell layer (GCL), inner

nuclear layer (INL), outer nuclear layer (ONL). **(C)** Representative images of a portion of flat-

mounted retinas stained with GFAP (green) at 8 weeks. **(D)** Muller glia reactivity gene, GLAST,

was analyzed by qPCR at 2 and 4 weeks. The results were normalized to GAPDH. **(E)** Astrocyte

markers were significantly upregulated at 2, 4, and 8 weeks when analyzed by qPCR. The results

were normalized to GAPDH and controls. n=4 (2 retinas were pooled for each replicate). *p<

0.05, **p<0.01, **p<0.001. (Ctrl- Control; OHT- ocular hypertension).

Figure 6. Effect of LXB$_4$ treatment on astrocyte reactivity markers in the mouse retina.

LXB$_4$ methyl ester treatment (n=8, 5 ng/g IP; 0.5 ng/g topical) was started after one week of

silicone oil injection every other day until the 4th week. Treatment with LXB$_4$ modulates

astrocyte reactivity genes in the ocular hypertension model, n=4 (2 retinas were pooled for each

replicate). (Ctrl- Control; OHT- ocular hypertension). *p< 0.05, **p<0.01, ***p<0.001.

Figure 7. Effect of LXB$_4$ treatment on lipoxin pathway in human brain astrocytes. (A)

Astrocyte reactivity genes analyzed by qPCR. The results were normalized to GAPDH. Data are

presented as the mean±SEM, n=3-4. **(B)** Representative immunofluorescence images of LCN2

and 5-LOX. Nuclei were labeled with DAPI. **(C)** LC☐MS/MS-based lipidomic quantification of

endogenous lipid mediators, n=4-5. (Ctrl-untreated control astrocytes, RA- reactive astrocytes).

*p< 0.05, **p<0.01, ***p<0.001.

References

1   Al-Shabrawey M, Zhang W, McDonald D (2015) Diabetic retinopathy: mechanism,
diagnosis, prevention, and treatment. Biomed Res Int 2015: 854593 Doi
10.1155/2015/854593
2   Alqawlaq S, Flanagan JG, Sivak JM (2019) All roads lead to glaucoma: Induced retinal
injury cascades contribute to a common neurodegenerative outcome. Exp Eye Res 183:
88-97 Doi 10.1016/j.exer.2018.11.005
3   Bazan NG, Calandria JM, Serhan CN (2010) Rescue and repair during photoreceptor cell
renewal mediated by docosahexaenoic acid-derived neuroprotectin D1. J Lipid Res 51:
2018-2031 Doi 10.1194/jlr.R001131
4   Ben Haim L, Carrillo-de Sauvage MA, Ceyzériat K, Escartin C (2015) Elusive roles for
reactive astrocytes in neurodegenerative diseases. Front Cell Neurosci 9: 278 Doi
10.3389/fncel.2015.00278
5   Borgeson E, Johnson AM, Lee YS, Till A, Syed GH, Ali-Shah ST, Guiry PJ, Dalli J, Colas RA,
Serhan CNet al (2015) Lipoxin A4 Attenuates Obesity-Induced Adipose Inflammation and
Associated Liver and Kidney Disease. Cell Metab 22: 125-137 Doi
10.1016/j.cmet.2015.05.003
6   Chauhan BC, Malik R, Shuba LM, Rafuse PE, Nicolela MT, Artes PH (2014) Rates of
glaucomatous visual field change in a large clinical population. Invest Ophthalmol Vis Sci
55: 4135-4143 Doi 10.1167/iovs.14-14643
7   Chen F, Ghosh A, Lin J, Zhang C, Pan Y, Thakur A, Singh K, Hong H, Tang S (2020) 5-
lipoxygenase pathway and its downstream cysteinyl leukotrienes as potential
therapeutic targets for Alzheimer's disease. Brain Behav Immun 88: 844-855 Doi
10.1016/j.bbi.2020.03.022
8   Colombo E, Farina C (2016) Astrocytes: Key Regulators of Neuroinflammation. Trends
Immunol 37: 608-620 Doi 10.1016/j.it.2016.06.006
9   Ding ZB, Song LJ, Wang Q, Kumar G, Yan YQ, Ma CG (2021) Astrocytes: a double-edged
sword in neurodegenerative diseases. Neural Regen Res 16: 1702-1710 Doi
10.4103/1673-5374.306064
10   Emerson MR, LeVine SM (2004) Experimental allergic encephalomyelitis is exacerbated
in mice deficient for 12/15-lipoxygenase or 5-lipoxygenase. Brain Res 1021: 140-145 Doi
10.1016/j.brainres.2004.06.045
11   Fang F, Zhang J, Zhuang P, Liu P, Li L, Huang H, Webber HC, Xu Y, Liu L, Dalal Ret al (2021)
Chronic mild and acute severe glaucomatous neurodegeneration derived from silicone
oil-induced ocular hypertension. Sci Rep 11: 9052 Doi 10.1038/s41598-021-88690-x

12 Firuzi O, Zhuo J, Chinnici CM, Wisniewski T, Praticò D (2008) 5-Lipoxygenase gene
disruption reduces amyloid-beta pathology in a mouse model of Alzheimer's disease.
FASEB J 22: 1169-1178 Doi 10.1096/fj.07-9131.com
13 Flitter BA, Fang X, Matthay MA, Gronert K (2021) The potential of lipid mediator
networks as ocular surface therapeutics and biomarkers. Ocul Surf 19: 104-114 Doi
10.1016/j.jtos.2020.04.008
14 Gao Y, Su J, Zhang Y, Chan A, Sin JH, Wu D, Min K, Gronert K (2018) Dietary DHA amplifies
LXA. Mucosal Immunol 11: 1674-1683 Doi 10.1038/s41385-018-0070-z
15 Giammaria S, Hutchison DM, Rafuse PE, Shuba LM, LeBlanc RP, Nicolela MT, Chauhan BC
(2022) Rates of Visual Field Change in Patients With Glaucoma and Healthy Individuals:
Findings From a Median 25-Year Follow-up. JAMA Ophthalmol 140: 504-511 Doi
10.1001/jamaophthalmol.2022.0671
16 Gronert K (2005) Lipoxins in the eye and their role in wound healing. Prostaglandins
Leukot Essent Fatty Acids 73: 221-229 Doi 10.1016/j.plefa.2005.05.009
17 Gronert K, Maheshwari N, Khan N, Hassan IR, Dunn M, Laniado Schwartzman M (2005) A
role for the mouse 12/15-lipoxygenase pathway in promoting epithelial wound healing
and host defense. J Biol Chem 280: 15267-15278 Doi 10.1074/jbc.M410638200
18 Gupta N, Yücel YH (2007) Glaucoma as a neurodegenerative disease. Curr Opin
Ophthalmol 18: 110-114 Doi 10.1097/ICU.0b013e3280895aea
19 Guttenplan KA, Weigel MK, Prakash P, Wijewardhane PR, Hasel P, Rufen-Blanchette U,
Münch AE, Blum JA, Fine J, Neal MCet al (2021) Neurotoxic reactive astrocytes induce
cell death via saturated lipids. Nature 599: 102-107 Doi 10.1038/s41586-021-03960-y
20 Guymer C, Wood JP, Chidlow G, Casson RJ (2019) Neuroprotection in glaucoma: recent
advances and clinical translation. Clin Exp Ophthalmol 47: 88-105 Doi
10.1111/ceo.13336
21 Heijl A, Buchholz P, Norrgren G, Bengtsson B (2013) Rates of visual field progression in
clinical glaucoma care. Acta Ophthalmol 91: 406-412 Doi 10.1111/j.1755-
3768.2012.02492.x
22 Ikonomovic MD, Abrahamson EE, Uz T, Manev H, Dekosky ST (2008) Increased 5-
lipoxygenase immunoreactivity in the hippocampus of patients with Alzheimer's disease.
J Histochem Cytochem 56: 1065-1073 Doi 10.1369/jhc.2008.951855
23 Inman DM, Horner PJ (2007) Reactive nonproliferative gliosis predominates in a chronic
mouse model of glaucoma. Glia 55: 942-953 Doi 10.1002/glia.20516
24 Jonas JB, Aung T, Bourne RR, Bron AM, Ritch R, Panda-Jonas S (2017) Glaucoma. Lancet
390: 2183-2193 Doi 10.1016/S0140-6736(17)31469-1
25 Kim C, Livne-Bar I, Gronert K, Sivak JM (2020) Fair-Weather Friends: Evidence of Lipoxin
Dysregulation in Neurodegeneration. Mol Nutr Food Res 64: e1801076 Doi
10.1002/mnfr.201801076
26 Levy BD, Clish CB, Schmidt B, Gronert K, Serhan CN (2001) Lipid mediator class switching
during acute inflammation: signals in resolution. Nat Immunol 2: 612-619 Doi
10.1038/89759
27 Liddelow SA, Guttenplan KA, Clarke LE, Bennett FC, Bohlen CJ, Schirmer L, Bennett ML,
Münch AE, Chung WS, Peterson TCet al (2017) Neurotoxic reactive astrocytes are
induced by activated microglia. Nature 541: 481-487 Doi 10.1038/nature21029

28 Liu HH, Flanagan JG (2017) A Mouse Model of Chronic Ocular Hypertension Induced by
Circumlimbal Suture. Invest Ophthalmol Vis Sci 58: 353-361 Doi 10.1167/iovs.16-20576

29 Liu HH, Zhang L, Shi M, Chen L, Flanagan JG (2017) Comparison of laser and circumlimbal
suture induced elevation of intraocular pressure in albino CD-1 mice. PLoS One 12:
e0189094 Doi 10.1371/journal.pone.0189094

30 Livne-Bar I, Wei J, Liu HH, Alqawlaq S, Won GJ, Tuccitto A, Gronert K, Flanagan JG, Sivak
JM (2017) Astrocyte-derived lipoxins A4 and B4 promote neuroprotection from acute
and chronic injury. The Journal of clinical investigation 127: 4403-4414 Doi
10.1172/jci77398

31 Lukiw WJ, Cui JG, Marcheselli VL, Bodker M, Botkjaer A, Gotlinger K, Serhan CN, Bazan
NG (2005) A role for docosahexaenoic acid-derived neuroprotectin D1 in neural cell
survival and Alzheimer disease. The Journal of clinical investigation 115: 2774-2783 Doi
10.1172/jci25420

32 Maderna P, Godson C (2009) Lipoxins: resolutionary road. Br J Pharmacol 158: 947-959
Doi 10.1111/j.1476-5381.2009.00386.x

33 Martin KR, Levkovitch-Verbin H, Valenta D, Baumrind L, Pease ME, Quigley HA (2002)
Retinal glutamate transporter changes in experimental glaucoma and after optic nerve
transection in the rat. Invest Ophthalmol Vis Sci 43: 2236-2243

34 Nucci C, Martucci A, Giannini C, Morrone LA, Bagetta G, Mancino R (2018)
Neuroprotective agents in the management of glaucoma. Eye (Lond) 32: 938-945 Doi
10.1038/s41433-018-0050-2

35 Pamplona FA, Vitória G, Sudo FK, Ribeiro FC, Isaac AR, Moraes CA, Chauvet MG, Ledur
PF, Karmirian K, Ornelas IMet al (2022) Age-linked suppression of lipoxin A4 associates
with cognitive deficits in mice and humans. Transl Psychiatry 12: 439 Doi
10.1038/s41398-022-02208-1

36 Rådmark O, Werz O, Steinhilber D, Samuelsson B (2015) 5-Lipoxygenase, a key enzyme
for leukotriene biosynthesis in health and disease. Biochim Biophys Acta 1851: 331-339
Doi 10.1016/j.bbalip.2014.08.012

37 Sapieha P, Stahl A, Chen J, Seaward MR, Willett KL, Krah NM, Dennison RJ, Connor KM,
Aderman CM, Liclican Eet al (2011) 5-Lipoxygenase metabolite 4-HDHA is a mediator of
the antiangiogenic effect of ω-3 polyunsaturated fatty acids. Sci Transl Med 3: 69ra12
Doi 10.1126/scitranslmed.3001571

38 Serhan CN (2014) Pro-resolving lipid mediators are leads for resolution physiology.
Nature 510: 92-101 Doi 10.1038/nature13479

39 Serhan CN, Chiang N, Dalli J, Levy BD (2014) Lipid mediators in the resolution of
inflammation. Cold Spring Harb Perspect Biol 7: a016311 Doi
10.1101/cshperspect.a016311

40 Serhan CN, Hamberg M, Samuelsson B (1984) Lipoxins: novel series of biologically active
compounds formed from arachidonic acid in human leukocytes. Proc Natl Acad Sci U S A
81: 5335-5339 Doi 10.1073/pnas.81.17.5335

41 Siracusa R, Fusco R, Cuzzocrea S (2019) Astrocytes: Role and Functions in Brain
Pathologies. Front Pharmacol 10: 1114 Doi 10.3389/fphar.2019.01114

42 Sofroniew MV (2009) Molecular dissection of reactive astrogliosis and glial scar
formation. Trends Neurosci 32: 638-647 Doi 10.1016/j.tins.2009.08.002

43 Son JL, Soto I, Oglesby E, Lopez-Roca T, Pease ME, Quigley HA, Marsh-Armstrong N
(2010) Glaucomatous optic nerve injury involves early astrocyte reactivity and late
oligodendrocyte loss. Glia 58: 780-789 Doi 10.1002/glia.20962
44 Tham YC, Li X, Wong TY, Quigley HA, Aung T, Cheng CY (2014) Global prevalence of
glaucoma and projections of glaucoma burden through 2040: a systematic review and
meta-analysis. Ophthalmology 121: 2081-2090 Doi 10.1016/j.ophtha.2014.05.013
45 van Leyen K, Klötzsch C, Harrer JU (2011) Brain tumor imaging with transcranial
sonography: state of the art and review of the literature. Ultraschall Med 32: 572-581
Doi 10.1055/s-0031-1273443
46 Wang R, Seifert P, Jakobs TC (2017) Astrocytes in the Optic Nerve Head of Glaucomatous
Mice Display a Characteristic Reactive Phenotype. Invest Ophthalmol Vis Sci 58: 924-932
Doi 10.1167/iovs.16-20571
47 Wang SB, Hu KM, Seamon KJ, Mani V, Chen Y, Gronert K (2012) Estrogen negatively
regulates epithelial wound healing and protective lipid mediator circuits in the cornea.
FASEB J 26: 1506-1516 Doi 10.1096/fj.11-198036
48 Wei J, Gronert K (2019) Eicosanoid and Specialized Proresolving Mediator Regulation of
Lymphoid Cells. Trends Biochem Sci 44: 214-225 Doi 10.1016/j.tibs.2018.10.007
49 Wei J, Gronert K (2017) The role of pro-resolving lipid mediators in ocular diseases. Mol
Aspects Med 58: 37-43 Doi 10.1016/j.mam.2017.03.006
50 Zhang J, Fang F, Li L, Huang H, Webber HC, Sun Y, Mahajan VB, Hu Y (2019) A Reversible
Silicon Oil-Induced Ocular Hypertension Model in Mice. J Vis Exp: Doi 10.3791/60409
51 Zhang J, Li L, Huang H, Fang F, Webber HC, Zhuang P, Liu L, Dalal R, Tang PH, Mahajan
VBet al (2019) Silicone oil-induced ocular hypertension and glaucomatous
neurodegeneration in mouse. Elife 8: Doi 10.7554/eLife.45881
52 Zhang J, Li Z, Fan M, Jin W (2022) Lipoxins in the Nervous System: Brighter Prospects for
Neuroprotection. Front Pharmacol 13: 781889 Doi 10.3389/fphar.2022.781889
699