Dark side of the honeymoon:
reconstructing the Asian x European rose breeding
history through the lens of genomics

Thibault Leroy[1,2,3], Elise Albert[1,4], Tatiana Thouroude[1], Sylvie Baudino[5], Jean-Claude Caissard[5], Annie Chastellier[1], Jérôme Chameau[1], Julien Jeauffre[1], Thérèse Loubert[6], Saretta Nindya Paramita[5], Alix Pernet[1], Vanessa Soufflet-Freslon[1], Cristiana Oghina-Pavie[7], Fabrice Foucher[1], Laurence Hibrand-Saint Oyant[1]*, Jérémy Clotault[1]*

[1] Univ Angers, Institut Agro, INRAE, IRHS, SFR QUASAV, F-49000 Angers, France
[2] Department of Botany and Biodiversity Research, University of Vienna, Vienna, Austria
[3] GenPhySE, Université de Toulouse, INRAE, ENVT, Castanet Tolosan, France
[4] Syngenta Seeds SAS, Saint-Sauveur, France
[5] LBVpam-UMR5079, Université Jean Monnet Saint-Etienne, CNRS, Saint-Etienne, France
[6] Roseraie Loubert, Les Rosiers-sur-Loire, Gennes-Val-de-Loire, France
[7] Université d'Angers, CNRS, TEMOS, SFR CONFLUENCES, Angers, France

* These authors contributed equally to this work.

Corresponding author: Thibault Leroy, GenPhySE, Université de Toulouse, INRAE, INPT, INP-ENVT, Castanet Tolosan, France thibault.leroy@inrae.fr

ORCIDs:
Thibault Leroy: 0000-0003-2259-9723
Elise Albert: 0000-0001-5060-4505
Tatiana Thouroude: 0000-0001-7908-7353
Sylvie Baudino: 0000-0002-3367-172X
Jean-Claude Caissard: 0000-0001-7938-3101
Annie Chastellier: none
Jérôme Chameau: none
Julien Jeauffre: 0000-0001-6770-0552
Thérèse Loubert: none
Saretta Nindya Paramita: 0000-0003-4013-5009
Alix Pernet: none
Vanessa Soufflet-Freslon: 0000-0002-1264-7595
Cristiana Oghina-Pavie: none
Fabrice Foucher: 0000-0002-3693-7183
Laurence Hibrand-Saint Oyant : 0000-0002-7604-3811
Jérémy Clotault : 0000-0002-7678-1929

Summary

- Roses hold significant symbolic value in Western cultural heritage, often serving as a symbol of love and romance. Despite their ancient cultivation, the appreciation for the phenotypic diversity of roses emerged relatively recently, notably during the 19th century. This period is characterized by a remarkable expansion in the number of varieties, from around 100 to over 8,000, representing a golden age for roses.

- To trace the history of rose breeding in Europe and unveil genetic changes during this period, we gathered phenotypic and genetic data from 204 accessions. These included botanical roses and varieties cultivated between 1800 and 1910. Whole-genome sequences from 32 accessions were also included.

- Our analysis revealed a temporal shift in the genetic makeup, transitioning from a historical European to a near-Asian genetic background within a few generations. This shift was accompanied by a notable reduction in genetic diversity, attributed to the backcrossing with the less diverse Asian genepool, plus some genomic signatures of selection.

- We have generated the largest GWAS catalog for rose to date, offering a valuable resource for future breeding initiatives. We emphasize the critical importance of preserving ancient rose collections to safeguard diversity and ensure a sustainable breeding for the long term.

Introduction

The neolithic revolution, the move from nomadic hunter-gatherers to rooted agrarian societies, has profoundly changed the history of humankind (Diamond, 2002). This transition has allowed some populations to have larger food surpluses, favoring rapid population growth, a division of labor, technological innovations and, at the end, the establishment of dominant societies and colonizers (Diamond, 2005). Domestication is expected to generate drastic reductions in the effective population sizes due to the subsampling of the wild progenitor species and the selection pressures that have then further reduced population sizes, which have long been assumed to lead to sudden losses of diversity (Nei *et al.,* 1975; see Gaut *et al.,* 2015 for empirical evidences). Recent investigations thanks to ancient DNA samples, in sorghum in particular, are however more consistent with a long-term gradual, linear or nearly linear, decline rather than a rapid drop, requestioning in part this bottleneck scenario (Allaby *et al.,* 2019; Brown, 2019; Smith *et al.*, 2019).

Unlike species domesticated for food production, the domestication of floral plants for aesthetic purposes is assumed to be far more recent, which can therefore even more question the existence of reductions of genetic diversity over these short periods of time. For most of the ornamentals, domestication indeed occurred in the last 500 years (Purugganan, 2022). Economically secure groups, especially the bourgeoisie in search of luxuries or aesthetic

pleasure, have played a major role in this advent, leading to an exploding number of domesticated ornamental species, in such a way that ornamentals have been described as outnumbering all other domesticated species combined today (Gessert, 1993, 2010; Chowdhuri & Deka, 2019). A typical example of this new passion for flowers in Western countries is tulips. Initially appreciated by Ottoman sultans and elites, the tulips imported to the Netherlands have led to a tulipmania, an irrational frenzy during the 17th century. Garden roses represent another emblematic example of domesticated flowers (Altman *et al.,* 2022), with a unique symbolic charge (Goody, 1993). Although rose domestication can be traced back as 5000 years BP (Altman *et al.*, 2022), the process of domestication is poorly understood. Its golden age is assumed to be even more recent than tulips, dating back to the 19th century (Leus *et al.,* 2018).

Despite roses being cultivated since antiquity, both independently in China and the Mediterranean region, the number of varieties has remained particularly limited for a long time. In ancient Rome, six cultivated species, assumed to correspond to cultivated forms of *Rosa alba*, *R. gallica*, *R. damascena*, and *R. moschata* were described (Pline l'Ancien, 2013). During the Middle Ages, roses were cultivated not only for their ornamental uses but also as religious symbols and for their medicinal properties (Touw, 1982). In Asia, roses were cultivated as early as the Han Dynasty, over 2,000 years ago, with selection of roses with new traits such as continuous flowering. The introduction of Asian roses to Europe at the end of the 18th century revolutionized rose breeding bringing traits such as recurrent blooming, novel scents and new colors (Wylie, 1954).

Thanks to considerable literature review efforts focusing on the 19th century, the history of a spectacular increase - by a factor 80 or so - in the number of rose varieties has been revealed (~100 in 1800, 6,000-8,000 in 1900, 30,000-35,000 today; Marriott, 2003; Oghină-Pavie, 2021). The diversification during the 19th century is associated with crosses between the two previously isolated genetic backgrounds, the Asian and European gene pools, giving rise to a large number of horticultural groups, including hybrid tea roses from the late 19th century (Martin *et al.,* 2001). Hybrid tea roses are considered as the parents of the modern roses and are known to be phenotypically diverse, harboring traits that originate from Chinese roses, especially the capacity of recurrent flowering, a particularly targeted trait that led to an extension of the flowering period in Europe (Marriott, 2003; Soufflet-Freslon *et al.,* 2021). Large-scale genetic investigation based on more than a thousand varieties genotyped at 32 SSR markers have indeed identified 16 genetic groups, most associated with a gradient of population structure from a European (groups 1-3) to an Asian (group 9) genetic background during the 19th century, with more recent modern roses (>1914) exhibiting a close to Asian genetic structure (group 8, Liorzou *et al.,* 2016). Despite the progress made regarding the description of the genetic structure, there are many unknowns, including the number of generations of breeding, the levels of diversity and their evolution or the proportion of the respective European and Asian genomes. Similarly, while progress has been made regarding the description of genetic variation at some key genes, including *RoKSN* or *AP2*, controlling the duration of the blooming period and the doubling of the petal number, respectively (Iwata *et al.,* 2012; François *et al.,* 2018; Gattolin *et al.,* 2018; Soufflet-Freslon *et al.,* 2021), the genetic bases of most targeted traits in roses remain unknown.

Roses represent an excellent ornamental model to reconstruct the past history of breeding and investigate the genomic footprints of artificial selection. First, and most importantly, roses are maintained and reproduced through vegetative culture, in particular grafting, allowing the continuous maintenance of ancient varieties. This allows the phenotyping

of roses that were bred at different periods of time in a single environment. In addition, it allows easy access to fresh material and therefore modern DNA from these varieties, without requiring expensive and challenging strategies such as the use of ancient DNA samples. Second, several reference genome assemblies are available for roses, in particular from the variety *Rosa chinensis* 'Old Blush' (Hibrand-Saint Oyant *et al.,* 2018; Raymond *et al.,* 2018), one of the four main ancient Asian roses used in plant breeding history (Liorzou *et al.,* 2016). Third, roses have relatively small genomes, around 500-550 Mbp - which for instance contrasts with the 34-Gbp of the tulip genome - allowing the resequencing of some rose accessions at affordable cost. Fourth, other molecular resources are available for roses, including a high density SNP array (68,893 markers, Koning-Boucoiran *et al.,* 2015), making high resolution mapping and Genome-Wide Association Studies (GWAS) possible (SchulzDietmar *et al.,* 2016; Hibrand-Saint Oyant *et al.,* 2018). However, one challenge associated with the model is the variable ploidy level among roses, ranging from diploid to decaploid, even if most roses are either diploid or tetraploid. GWAS methods were initially developed for diploid models, but more recent developments, especially for polyploids (*e.g.* Rosyara *et al.,* 2016), provide new opportunities to uncover the genetic bases for important traits in roses. One additional difficulty associated with this model is the complexity of the genetic relationships in the *Rosa* genus, with a hundred or more wild species (Wissemann & Ritz, 2007; Debray *et al.,* 2021), for which approximately fifteen of them may have genetically contributed to present-day rose diversity.

In this study, we reconstruct the history of the rose garden breeding in Europe. To do so, we assembled a unique dataset, composed of large phenotypic data, genetic data from 204 roses genotyped with a 69k SNP array, plus whole-genome sequence data from 32 roses. The sequenced accessions are either botanical roses or varieties bred in Europe between 1800 and 1910, and include star varieties at that time. Botanical roses are considered as wild species, but they are maintained in rose collections and are assumed to be non-intentionally selected. By combining population genomics and quantitative genetic analyses, we (i) described the genomic makeup of the rose varieties, (ii) provided an approximate number of generations of selection, (iii) estimated the levels of genetic diversity and investigated their evolution throughout the period, (iv) identified potential footprints of artificial selection, and (v) performed the largest GWAS to date to uncover the genetic bases of many horticulturally important traits.

Materials and methods

Sampling & DNA extraction

Roses suspected to be diploid and tetraploid, based on prior knowledge (Liorzou *et al.*, 2016), were prioritized during our sampling campaigns (see below). Our sampling also had the objective to encompass a range of rose groups covering the 19th century, including the one that are hereafter referred to as ancient Asian and European roses, early European × Asian hybrids, hybrid tea and botanical roses.

A total of 288 samples were collected in the 'Rose Loubert' rose collection and nursery (Roseraie Loubert, Gennes-Val-de-Loire, Pays de-la Loire, France) in summers 2020 and

2021, except for *Rosa* ✕ *wichurana* and 'Old Blush', both collected in an experimental
collection at INRAE (UE Horticulture, Angers, Pays de la Loire, France).

In complement, we performed whole-genome sequencing for fifteen roses. Newly sequenced
samples were chosen, based on their presumed importance during the 19th century
(considered as "star varieties" based on extensive literature searches on historical catalogs of
varieties; Cristiana Oghina-Pavie, personal communication) and the within-group diversity
previously observed at 32 SSR markers (Liorzou et al., 2016). Fresh plant material was
collected in the Roseraie Loubert in summer 2021, with the notable exception of 'Jin Ou Fan
Lu', a cultivar that was previously collected by Shubin Li in China (Liorzou *et al.*, 2016). It
should be noted that we had access to only a small part of the ancient roses, both because of
the investigated collection, but also because a part of the ancient varieties have not been
preserved down to this day. As a consequence, our sampling was potentially impacted by a
conservation bias.

For each rose, 100 mg of leaves were collected and DNA was extracted from either lyophilised
or fresh leaves using the Qiagen 96 Plant Kit (Qiagen, Hilden, Germany). DNA quality and
quantity were checked with the Nanodrop technologies and Qubit kit prior to sending samples
to the different facilities.

Genotyping and sequencing data

Genotyping was performed on 3*96-plates at the Gentyane facility (INRAE Clermont-Ferrand,
France) using the Affymetrix Axiom technology. To further check the quality of the genotyping
(see below), we included technical replicates (21 duplicate samples, as well as a triplicate one,
shown in yellow in Table S1).

Regarding sequenced samples, library preparations and sequencing were performed at the
Vienna BioCenter Core Facilities (VBCF, Vienna, Austria). For the library preparations, the
facility used the Ultra II DNA kit for Illumina (New England Biolabs), which included an
enzymatic shearing as a first step. Quality control was performed prior to sequencing. Quality
control of the obtained NGS libraries was then assessed prior to the sequencing using the
Fragment Analyzer technology. The 15 genotypes were then sequenced on a single NovaSeq
SP flow cell (300 cycles, 2*150 reads). Demultiplexing was also performed by the VBCF
facility.

Phenotyping

218
All phenotyping data were collected in the rose garden (Roseraie Loubert, Rosiers sur Loire,
Gennes-Val-de-Loire, France). The number of petals was scored on at least two and at most
five flowers. We used averaged petal counts across the different flowers. We also recorded
the number of flowers per inflorescence and categorized roses by the primary color of their
flowers. Additionally, we measured the number of prickles and acicles, along with several
architectural traits: height (meters), circumference (meters), growth form, and bushiness
(binary). While pruning is performed within the rose garden, it is important to note that it
respects the natural form of roses and is not expected to considerably alter their architecture

or growth patterns. However, this limitation associated with the size of the roses in the rose
garden should be kept in mind when interpreting these results.

Plants were scored in the rose garden every week during the blooming period (from spring to
autumn) for the recurrent blooming. This flowering monitoring was conducted over a period of
three years (2014-2016). We have previously developed a statistical method on the flowering
curve using Gaussian mixture models and created indicators, as Reb.Mag for the magnitude
of reblooming (for details, see Proïa *et al.*, 2016). A value of 0 means that the rose flowers
only in spring (once-flowering roses), the higher the value (above 0), the higher the reblooming
capacity (see Supplementary Note 1). For the sake of clarity, the index shown in Fig. 1B
corresponds to a simplified version, with values corresponding to three phenotypic classes (1:
non-recurrent flowering, 2: partly recurrent, 3: recurrent flowering), an average value of the
index across the three years of scoring was then computed for each individual.

Flowers were also collected during two years (2017 and/or 2018) to analyze scent
components. Concretely, from two to six flowers per sample and year (2017: mean=2.34,
sd=0.78; 2018: mean=2.33, sd=0.81) were harvested around 10:00 am at stage 4 of flower
development, as previously described (Bergougnoux *et al.,* 2007) and volatile compounds
were extracted following Roccia *et al.* (2019). Per-genotype averaged values of each
compound per year were used for all the analyses.

Susceptibility to black spot disease was also scored in the field over three years (2014-2016).
Each cultivar was scored, between July and September, depending on the development of
disease and according to the scale indicated in Marolleau *et al.* (2020). The observed black
spot symptoms were due to natural infection with natural inoculum of fungal strains. It should
be noticed that we were scoring roses in the field, with present-day strains. The strains present
at the time of breeding (*i.e.* 19th century) might have been different, including their virulence
gene repertoires and aggressiveness factors.

252
Data analysis

The analyses performed in the manuscript are described below, considering (i) the analyses
based on the SNP array and (ii) those on the whole-genome sequencing data. All the scripts
used have been made available on a Zenodo repository (10.5281/zenodo.14450242).

1/ SNP array

<u>*Genotyping*</u>

We used a two-round genotyping procedure to ensure highly accurate calls. First, we
performed a QC analysis under Axiom Analysis Suite based on all the genotypes (288). We
used the Dish Quality Control (DQC), the recommended QC metric for the Axiom SNP array
to capture the extent to which the distribution of signal values is separated from background
values, with 0 indicating no separation and 1 indicating perfect separation. Based on this
genotyping, four individuals exhibited very low DQC (<0.25) and were excluded. Based on all
the other individuals (with DQC > 0.85, 284 samples), we generated a first genotype matrix
with FitPoly (Voorrips *et al.*, 2011) assuming a ploidy=4 for all samples.

Given that roses are grafted, all samples with a close genetic proximity with the rootstock
cultivar used in the Loubert's rose garden were considered as sampling errors and were
excluded from the dataset (seven in total). In addition to these previously excluded genotypes,
we excluded seven additional genotypes with a DQC < 0.94 and a QC Calling rate < 0.89,
resulting in a dataset of 270 individuals. For this new dataset, we performed detailed QC and
then converted the output in a FitPoly format thanks to FitPolyTools (version 1.1; unpublished
R package by Roeland Voorrips). To take into account the variable ploidy level between
individuals, with the vast majority of the accessions used expected to be either diploid or
tetraploid, we generated a final genotyping matrix using FitPoly assuming ploidy=4 for all
individuals, which means that diploid individuals with heterozygous alleles are generally
expected to be called AABB. The rose SNP array targets 68,893 SNPs, with probes for both
strands, leading to a total number of 137,786 probes (for details, see Koning-Bourcoiran *et al.,*
2015). After SNP filtering, we obtained a dataset of 92,007 probes, corresponding to 56,467
of the 68,893 unique SNPs. Given all the difficulties associated with the genotyping of the SNP
array, especially in a species with a variable ploidy level, we made the choice of using all
probes satisfying our QC criteria in the subsequent analyses.
283

QC checking and clone exclusion
Given that roses are clonally propagated and used commercially worldwide, several varieties
can be registered under different cultivar names. In addition, spontaneous somatic mutants
(also called sport) exhibiting different traits (*e.g.* flower color or shape) are frequent in roses
and could have therefore been registered with two different names. To exclude potential clones
in the dataset that could bias our quantitative and population genomics analyses, we computed
the proportion of similar calls between all pairs of individuals and considered all samples
exhibiting a value higher than 0.965 as potential clones. To define this value, we considered
the observed distribution of the proportion of similar calls based on all pairwise comparisons
and the lowest observed value for a replicate of the analysis (0.973 for '*Persian yellow*'), see
Fig. S1 and Table S1 for a list. As a consequence, our "clone-corrected" dataset contains 204
unique samples. All subsequent analyses are based on this dataset.
296

Population structure
We used the R package adegenet (Jombart, 2008) to explore the population structure among
the 204 individuals. We first used plink (v2.00, Purcell *et al.,* 2007, with --index-pairwise 20 5
0.2) to generate a pruned SNP set of 11,848 SNPs among the 92,007 markers. We then used
the df2genind (with ploidy=4) to generate the input files before using the dudi.pca function of
the adegenet package. PCA results were then plotted using both the ggplot2 and ggrepel
packages (Wickham, 2016; Slowikowski, 2023). To confirm the population structure observed
with the PCA, we also used faststructure (Raj *et al.,* 2014), with K ranging from two to five
groups.
306

GWAS
Before performing GWAS analyses, we further filtered the dataset in order to exclude poorly
informative markers. SNPs with a missing rate greater than 10% and/or a Minor Allele
Frequency (MAF) < 0.05 were discarded. GWAS was then performed with GWASpoly
(Rosyara *et al.*, 2016) for all traits. Concretely, we used the Q + K linear mixed model, providing
our own Q matrix based on the Principal Coordinates Analysis (dudi.pco, adegenet, Jombart,
2008) as covariates. Considering both the tetraploidy and the absence of strong expectations

regarding the additive *vs.* dominant effects of the genes for the studied traits, we considered
all the potential models of gene action in polyploids and implemented in different models under
GWASpoly (*i.e.* general, additive, diplo-additive, diplo-general, simplex and duplex-dominant
models, for details regarding the models, see Rosyara *et al.,* 2016). Concretely, by utilizing
GWASpoly to analyze the genetic basis of traits in European roses, our objective was to
address the complexity of the polyploid gene action as much as possible. To explore the
results, we first generated the QQplot and the Manhattan plot using R base, but we then
generated linear or circular representations of the results under the R packages qqman (Turner
et al., 2018) or circlize (Gu *et al.,* 2014), respectively. In addition to the Manhattan plots, these
circular views include density curves in order to identify some potentially interesting regions,
even when the empirical background noise is high, generally associated with a general inflation
of p-values (as tracked with the QQplots). Given the number of traits analyzed, as well as the
number of models explored for each, all the 'circlize' visualizations were made available thanks
to a dedicated website (https://roseGWASbrowser.github.io/). The website was successfully
tested on various web browsers, including Firefox (*e.g.* v. 133.0) and Chromium (*e.g.* v.
131.0.6778.85). The rest of the results including the p- and q-values, as well as the QQplots
and the linear Manhattan plots were made available on a Zenodo repository (see the data
availability section).
332
333
334   ***2/ Whole-genome sequences***
335
Mapping and calling
A total of 32 genotypes are used as part of the whole-genome analyses, corresponding to a
set of 17 rose sequences publicly available and 15 new sequences generated in this study.
The new sequences have been made available on the Sequence Read Archive (SRA) under
the BioProject PRJNA997103 (see Table S2 for all SRA accession numbers).
Briefly, we used Trimmomatic (v.0.38, Bolger *et al.,* 2014) to remove adapters, trim and filter
reads using the following set of parameters: LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15
MINLEN:50. All trimmed reads were then mapped against a high-quality rose reference
genome (Hibrand-Saint Oyant *et al.,* 2018) with BWA mem2 (v. 2.2, Vasimuddin *et al.,* 2019)
using default settings. Most of the mapped data corresponds to paired ends reads but we also
mapped single end reads (*i.e.* corresponding to unpaired after trimming). MarkDuplicates
(Picard v. 2.20.7; Picard toolkit, 2019) was then used to remove potential PCR duplicates.
Variant calling was performed on GATK4 (v.4.2.2.0, Poplin *et al.,* 2018), following the GATK
best practices guidelines. In brief, we performed a first round of HaplotypeCaller to generate
the GVCFs, followed by a CombineGVCFs and a GenotypeGVCFs with the "--all-sites true"
option. Variants were then filtered with VariantFiltration, discarding variants with SOR>4.0,
MQ<30, QD<2, FS>60, MQRankSum< -20, and ReadPosRankSum either <- 10 or >10.
353
Ploidy inference
The ploidy level of each sample was empirically determined by considering the observed allelic
balance at heterozygous sites. Specifically, for each individual, we subsampled 2% of the total
number of SNPs and then considered, for each individual, the heterozygous calls exhibiting a
coverage greater or equal to 20 and then generated the per-individual density curves of the
allelic balance based on all these SNPs. Because most of the polymorphisms were expected
to be rare in the population, we could observe the first neat peak to be near f=0.25, 0.33 and
0.5 for tetraploids, triploids and diploids (see Fig. S2). Similar methods have been developed

over the last decade to empirically estimate the ploidy level (*e.g.* ploidyNGS, Augusto Corrêa dos Santos *et al.,* 2017). To ensure that the inferred ploidy level was sufficiently robust for our subsequent analyses, we checked the overall consistency of the results with previous in-house estimates of the ploidy level (for details, see Liorzou *et al.,* 2016 and http://dx.doi.org/10.5281/zenodo.56704).

Population structure & kinship

Population genetic structure among the 32 varieties was performed using Principal Component Analysis (PCA). Analyses were run on a set of 17,669 SNPs, which was generated by first randomly selecting 50,369 biallelic SNPs across the genome and then pruning this set for linkage disequilibrium with plink (v1.90, parameters: "--index-pairwise 20 5 0.2"). We used the snpgdsPCA function from the SNPRelate R package (Zheng *et al.,* 2012) to generate the PCA. We then used ggplot2 (Wickham, 2016) to generate the final plots. Given that the reference corresponds to an ancient Asian rose, we also empirically checked that the observed population structure could not have been generated by a strong mapping bias (Fig. S3). Similarly, given that the first round of calling assumed diploid individuals which is not correct for all individuals (see ploidy inference), we also performed PCA directly based on allele counts (AD field from the vcf) from a list of 10.5 million SNPs randomly selected among the biallelic SNPs to evaluate the robustness of our population structure (i) to a larger SNP set and (ii) to deviation from diploidy. PCA was performed with the function 'randomallele.pca' from the R package 'poolfstat' (Hivert *et al.*, 2018; Gautier *et al.*, 2022; Supplementary Note 2). In addition, as a control of the population structure observed using PCAs, we also used faststructure (Raj *et al.,* 2014) on the same 50k SNP set. To do so, we first used plink (v2.00, Purcell *et al.,* 2007) to generate the input files and then inferred the population structure for K=1 to K=6. Barplots of the inferred individual ancestry memberships were also generated using ggplot2.

We performed a family relationship inference by estimating kinship coefficients under KING (Manichaikul *et al.,* 2010) based on the whole SNP set (77,862,879 filtered SNPs). The initial objective of this analysis was to ensure that no clones were present in our sequencing dataset (see also above). But given that KING is able to infer relationships up to the third-degree (*i.e.* great grandparents and great grandchildren, first cousins, or similar), we used KING to infer some potential family relationships. We first generated the input files with plink (v2.00) and then estimated kinship coefficients. Kinship values higher than 0.354 correspond to clones, while coefficients coefficients within the intervals [0.177, 0.354], [0.0884, 0.177], and [0.0442, 0.0884] correspond to 1st-degree, 2nd-degree, and 3rd-degree relationships, respectively. Since KING analysis was conducted on the initial SNP calling round, resulting in diploid calls, we cannot rule out a potential impact of variable ploidy levels among the roses in the accuracy of the number of generations (Supplementary Note 2). We used the R package igraph (Csárdi & Nepusz, 2006) to generate a comprehensive network based on all the inferred family relationships.

Nucleotide diversity estimates

As compared to population structure and kinship inferences, estimates of diversity are highly dependent on the final quality of the SNP set. To ensure unbiased estimates, we used the filtered set of SNPs based on the 32 samples (77,862,879 filtered SNPs) to perform a Base Quality Score Recalibration (BQSR) on the 19 retained samples, as recommended by GATK in the absence of a highly-reliable SNP catalog for the focal species. Then, we used GATK's

ApplyBQSR function. HaplotypeCaller was then performed assuming the inferred ploidy (see previous section) using option "--ploidy", before using CombineGVCFs, GenotypeGVCFs and VariantFiltration as previously described to generate a final SNP set of 54,481,222 filtered SNPs. We then reconstructed whole-genome fasta sequences based on the complete VCF (vcf with variant and non-variant positions; to know more about the importance of using non-variant positions, see Korunes & Samuk, 2021) following Leroy and collaborators (2021a, 2021b). The approach was adapted to the rose specificities to reconstruct up to 4 sequences per individual, by explicitly considering the inferred ploidy (per-individual number of alleles in the GT field). Briefly, the pipeline reconstructs the sequence by considering the coverage at each position. All positions that are not in between the 5th and 95th centiles of the individual coverage, as well as those with coverage lower than 3 were hard-masked in the reconstructed sequences. For positions satisfying the coverage criteria, the reference allele is added to the reconstructed sequences, except at PASS positions, where alleles are added according to the genotyping calls (*i.e.* the GT field). Triallelic (or more) variants were discarded. Similarly, indels were ignored to keep the sequence length the same across individuals and therefore obtain perfectly aligned sequence blocks. Nucleotide diversity and Tajima's D were then computed on non-overlapping 100-kbp sliding windows following Leroy *et al.* (2021a) for each group, based on the individuals falling in the four following groups: ancient Asian, ancient European, early European x Asian and hybrid tea roses (Table S2). Botanical roses were not included in this part of the study because their population structure is not consistent with a single group. A Reduction of Diversity (RoD) index was computed by $1 - \frac{\pi Group1}{\pi Group2}$, with *Group1* and *Group2* corresponding to hybrid tea roses and ancient European, respectively, for a comparison aiming at estimating the evolution of the diversity in hybrid tea roses as compared to the ancient European samples. A positive value of the RoD index therefore indicates a net reduction of the nucleotide diversity between the two groups. Circular visualization was generated using the R package circlize (Gu *et al.,* 2014).

Diagnostic alleles

We then considered allele frequency at diagnostic alleles between the ancient Asian and European genetic backgrounds as a source of information regarding the local ancestry of early European x Asian and hybrid tea roses. For each filtered SNP among the final list of 54,481,222 SNPs, allele frequencies of the four groups were computed directly based on the filtered joint vcf thanks to a home-made script (script_freqgroup4localancestryprop.py), which was made available on a Zenodo repository (DOI: 10.5281/zenodo.14450242). Given that the number of genotypes available per group is limited (4-7 individuals, corresponding to a total of 16 chromosome sets per group), allele frequency was only computed if the number of missing alleles was lower or equal to 4 (*i.e.* corresponding to a maximum of 1 tetraploid individual or 2 diploid individuals without genotyping calls). We then subsampled SNPs exhibiting diagnostic alleles, defined as those for which the reference allele frequency is 0 in one group and 1 in the other. In this specific example, all diagnostic SNPs identified - at the notable exception of 2 SNPs that were subsequently excluded - exhibit an allele frequency of 0 in the European and of 1 in the Asian group. Indeed, all the raw sequencing data were mapped against the genome of an ancient Asian genotype ('Old Blush'; Hibrand-Saint Oyant *et al.,* 2018). Consequently, the reference alleles are expected to be more associated with the ancient Asian background. The match is almost perfect here given that we also used the 'Old Blush' variety as a member of the ancient Asian group in our study. As a consequence, all diagnostic alleles are expected to have an allele frequency of 1 in the ancient Asian group and, consequently, 0 in the ancient

European group. Importantly, the diagnostic SNPs are based on our limited panel of varieties.
Consequently, all are not expected to be truly diagnostic based on a larger panel.
Local ancestry of early European x Asian or hybrid tea groups was then estimated by
computing the median of the observed allele frequencies at diagnostic markers. Estimates
were based on non-overlapping 100-kbp sliding windows spanning the whole genome for
windows containing at least 5 diagnostic SNPs.
463
Local footprints of selection
To know more about the genomic regions targeted by 19th century breeders, we identified
regions that could be consistent with selective sweeps. One limitation associated with the
model is the recent history of the rose breeding and the associated limited number of
generations of recombination (see results). As a consequence, explicit selective sweep
methods could not be used here to identify potential footprints of selection. As an alternative,
we identified regions simultaneously exhibiting the lowest π, the most negative Tajima's D and
the highest RoD values. Given that this strategy does not fully circumvent the problem of the
lack of generations of recombination, we have decided to focus our detection on hybrid tea
roses, the latest group in the history of breeding among those investigated in this study, and
therefore expected to have experienced the highest number of generations of recombination,
in order to deliver more interpretable results. We considered two sets of candidate windows.
The stringent set corresponds to windows falling in the last centile of the three metrics
supporting a potential selective sweep (*i.e.* bottom 1% of nucleotide diversity and Tajima's D
values, top 1% of the RoD comparing hybrid tea roses with either Ancient European or Ancient
Asian), and a less stringent set considering the last five centiles (Table S4).
480
481

Results

482
483
484 **19th century: directed or unintended evolution of some key rose traits**
485
Asexual rose propagation through grafting or cutting allows the maintenance of ancient
roses in gardens and therefore the direct comparisons of phenotypes obtained at different
periods of the breeding history. We explored the phenotypic variation of a large collection of
botanical and ancient accessions, representing more than 200 roses bred between 1800 and
1910 (Fig. 1A), a period which is considered as the golden age for rose breeding. In this study,
we investigated about 30 traits in 200+ varieties, including several presumably crucial traits for
breeding, such as the blooming period, the color and number of petals, the number of prickles,
the resistance to black spot disease, plus some flower and plant architecture phenotypes (Fig.
1B). A particularly massive effort was made to analyze the floral scent by quantifying the
volatile compounds of the collection using a GC-MS strategy, including the 2-phenylethanol
and the geraniol, which represent two main components of the floral rose scent (Spiller *et al.,*
2010; Magnard *et al.,* 2015; Roccia *et al.,* 2019; Caissard *et al.,* 2022).
498

499
Figure 1: Main phenotypic changes along the 19th century plant breeding history. A.
Pictures highlighting the phenotypic variation of flowers at different periods of the rose
breeding history among the different accessions used in this study. Pictures were taken by T.
Leroy in the Loubert rose collection in 2021 and 2022 and selected to be representative
regarding the number of petals and the variation of colors observed among all the varieties of
a given period. Pictures are not to scale. ***B.*** *Phenotypic variation for some of the studied traits:*
number of petals (dark pink), recurrent flowering (light pink, estimated by an index), scent
components (2-phenylethanol and geraniol contents, light beige and light yellow, respectively)
and susceptibility to black spot disease (yellow). Averages are shown with light blue lines.
Tukey's honestly significant difference (HSD) tests at α=0.05 are indicated with letters.
510

Among all these traits, those associated with flowering are expected to be the main
target of selection during the 19th century, including the production of roses containing tens of
petals and the ability to have an extended blooming period. Most botanical accessions indeed
produce flowers containing only five petals (Fig. 1B) and during a limited period of time in
spring. Based on our sampling, the mean number of petals has strongly increased during the
first half of the 19th century (including roses with up to 210 petals), emphasizing the
importance of selecting for this trait at the beginning of the 19th century. From 1870, the trend
reversed with a reduction of the mean number of petals. Phenotypic variation also supports a
relatively constant increase of the repeat-flowering phenotype based on our rose collection,
with a progressive increase of the mean of the recurrent blooming index during the 19th
century (Fig. 1B).
A main criticism associated with modern roses is the popular impression of a reduction
of the flower fragrance. Based on our collection of roses, a slight reduction was indeed
observed during the 19th century based on the mean contents of both 2-phenylethanol and
geraniol, two main components of the rose scent. Changes during this period are however
either weakly significant or non-significant depending on the components, owing mainly to the
large observed variance among roses of each period (Fig. 1B).
Temporal changes in the resistance to *Diplocarpon rosae*, the pathogen responsible
for black spot disease - one of the most serious diseases on garden roses today -, were also
investigated. As compared to the previous examples, resistance to black spot was likely not
directly selected by breeders at that time since rose growers started to show interests for

Diplocarpon rosae (also called *Marssonina* ou *Marsonia rosae* Briosi & Cavara) at the
beginning of the 20[th] century (Ducomet, 1903; Cristiana Oghina-Pavie, personal
communication), even though today the resistance to this fungal pathogen represents one of
the main breeding targets. Botanical roses from our collection exhibit exceptional resistance
levels to *D. rosea* (*i.e.* low susceptibility score) but this resistance has progressively
plummeted during the rose breeding history, at least until 1890 (Fig. 1B, see also Fig. S4),
which therefore questions the origin of this erosion of resistance (loss of resistance alleles,
evolution of the pathogen, etc). Yellow and orange-colored roses are often suspected by
breeders to be more susceptible to black spot, hypothetically leading to a trade-off between
colors and resistance for breeders. No significant differences in susceptibility were however
observed between varieties with different main color petals (Fig. S5). The hypothesis regarding
the higher susceptibility of yellow flower varieties remains open since our sampling is biased
toward pink and white flowers, largely underrepresenting the genotypes with yellow flowers,
mostly because roses with yellow- or orange-colored flowers were relatively rare during the
19th century (Leus *et al.,* 2018).
547
548
549 **A rapid shift from a European to an Asian genetic background**
550
The same large collection of rose varieties was genotyped at 68,893 SNPs using the
Rose Genotyping Array (WagRhSNP Axiom Array, Koning-Boucoiran *et al.,* 2015), a SNP
array developed from modern cut and garden roses. After the removal of clonal genotypes, a
final set of 204 rose varieties was used for the subsequent analyses (see Materials and
methods, Fig. S1 and Table S1). The first axis of a Principal Component Analysis (PCA)
explains more than a fifth of the total variance (21.4%) and isolates the two ancient genetic
backgrounds: an Asian one (in yellow) and an European one (in blue, Fig. 2A). The whole
dataset forms a gradient of population structure between the two backgrounds, which is
consistent with the frequent crosses between European and Asian roses throughout the 19th
century. The second axis isolates some samples, including most of the botanical samples,
from the Asian-European gradient. This population structure is mostly consistent with a
previous report based on the genotyping at 32 microsatellite markers of an even larger
collection of varieties and the identification of 16 genetic groups (Fig. 2A, Liorzou *et al.,* 2016).
Independently from this first dataset, we resequenced 15 varieties and collected publicly
available rose genomes to generate a dataset of 32 samples, composed of ancient Asian,
ancient European, early European x Asian, hybrid tea and botanical samples (Table S2). After
joint-genotyping and SNP filtering, we generated a dataset of 77,862,879 filtered SNPs on the
32 samples. PCA based on a LD-pruned set of 17,669 variants among 50,369 randomly
selected SNPs (Fig. 2B) is consistent with an Asian-European gradient of population structure,
as previously identified with the SNP array (Fig. 2A). The proportion of variance explained by
the first axes is however lower based on the sequencing data, likely due to SNP ascertainment
bias associated with the design of the array, since the SNP set was established based on the
analysis of more modern varieties, therefore emphasizing more on the European-Asian
gradient (*i.e.* the axis 1). Based on the WGS data, some axes can be built due to a single
botanical individual (*e.g.* axes 2 and 4, Fig. S8), suggesting that a large proportion of the
diversity is indeed only present in the botanical accessions.

Figure 2: Population structure and levels of genetic diversity in several rose groups used in the 19th century rose breeding history. A. *Population structure of the 204 varieties genotyped with the rose SNP array based on the two first axes of a Principal Component Analysis (PCA). To check the consistency with previous works, the membership of the samples to one of the 16 genetic groups defined by Liorzou and collaborators (2016) based on the analysis of SSR markers are shown (for details, see Liorzou et al., 2016). Roses with ancient European and ancient Asian backgrounds are shown in blue and yellow, respectively (typically corresponding to groups 1-3 and 9, respectively; see Figs. S6-S7). **B.** Population structure based on the whole-genome sequences used in this study. Axes 1 and 3 of a Principal Component Analysis (PCA) based on a random sampling of 50k SNPs. Ancient European, ancient Asian, early European x Asian, hybrid tea varieties and botanical accessions are shown in blue, yellow, green, khaki and red, respectively. Results from other components as well as more detailed information about the cultivar IDs are shown in Figs. S8 and Table S2, respectively (but see also Fig. S9 and Supplementary Note 2). **C.** Network of the relationships inferred based on the kinship coefficients estimated under KING based on the whole SNP set of 77,862,879 SNPs. First-, second- and third-degree relationships are represented with red, orange and yellow lines, respectively. See Table S2 for a correspondence table between accession numbers and cultivar IDs. **D.** Genetic diversity estimates based on the different groups. Distribution of the nucleotide diversity levels based on non-overlapping 100 kb sliding windows spanning the genome (each dot corresponds to a window). Genomic windows on chr00 are not included. For ease of comparison, a vertical line is shown representing the mean π value per group.*

Family relationships were then inferred using KING (Manichaikul *et al.,* 2010) based
on the pairwise kinship values calculated on the sequencing data. Among the 32 samples,
only a few individuals are not connected to at least another individual of the dataset through a
third degree relationship (*i.e.* kinship coefficient > 0.0442) (Fig. 2C). These individuals
correspond to all botanical samples, plus two ancient Asian samples (*R. odorata gigantea* and
R. chinensis spontanea, numbers 13 and 14 in Fig. 2C, respectively). The rest of the
individuals are connected through a large network, linking the ancient European and the
ancient Asian backgrounds through the early European x Asian and hybrid Tea varieties (Fig.
2C). This result is remarkably consistent with the expected history of selection during the 19th
century, with (1) the intercrossing of the Asian genepool with the European one in the
European breeding practices, (2) leading to the obtention of early European x Asian varieties,
and (3) the backcrossing of these early hybrids with the ancient Asian genepool to obtain the
hybrid tea varieties. Obtaining such a network with a so limited number of samples, and without
integrating prior knowledge about the potential pedigrees, suggests that the selection was
based on a quite narrow diversity, in terms of the number of varieties used. In addition, it
suggests that the number of generations of selection was also very limited (but see also some
limits in Supplementary Note 2). Hybrid roses from the 19th century are indeed connected to
Asian and European gene pools by less than 10 generations of breeding (Fig. 2C, but see also
Supplementary Note 2).
To investigate the contribution of Asian or European gene pools to early European x
Asian and hybrid tea roses, we identified SNPs with diagnostic alleles between the ancient
Asian and ancient European samples. In total, 170,637 diagnostic SNPs were identified
among the final list of 54,481,222 SNPs (0.31%), distributed on all chromosomes (Table S3).
At these markers, the early European x Asian samples remarkably have a 1:1 ratio, *i.e.* an
almost equal share between the two gene pools (50.9% and 49.1% of Asian and European
alleles, respectively; Fig. S10). One might propose that this 1:1 ratio implies that all early
European x Asian samples used in the study are first-generation (F1) hybrids between the
ancient Asian and European groups of samples. However, detailed analyses at the individual
level are inconsistent with this hypothesis (Supplementary Note 3). If we can exclude the F1
hypothesis, we can then subsequently exclude the BC1 hypothesis, despite the similarly
noticeable 3:1 ratio observed for hybrid tea roses (74.8% and 25.2% of Asian and European
alleles, respectively; Fig. S10). To gain a more accurate understanding of the 19th-century
rose breeding history, especially concerning the number of generations leading to the "star
varieties" in our study, additional research reconstructing pedigrees based on a larger number
of whole-genome sequenced accessions will be needed.
635
636
637   **Genetic diversity erosion**
638
Nucleotide diversity was estimated over non-overlapping 100-kbp sliding windows on
the four groups (namely, ancient European, ancient Asian, early European x Asian and hybrid
tea roses), each containing the same number of complete chromosome sets (16) after
considering the inferred ploidy based on the distribution of allelic balance (Table S2, *e.g.* in
Ancient Asian the 16 complete chromosome sets corresponds to 6 individuals inferred as
diploid individuals, plus a tetraploid). Among all the dataset, the genetic diversity in roses was
relatively high with an average π of 1.47×10^{-2} (Fig. 2D). Significant differences in diversity were
observed between groups, with higher diversity in the ancient European (1.67×10^{-2}) than in the
ancient Asian samples (1.20×10^{-2}).

The "early European x Asian" group exhibits an averaged nucleotide diversity of
1.52×10^{-2}, a value that slightly exceeds the expectation assuming both the previous estimates
and the genetic makeup of this group (50.1%/49.1%, expectation: 1.43×10^{-2}). This slight burst
of genetic diversity in the first generations of hybridization could be explained by the mixing of
parental alleles from the two quite divergent genetic backgrounds.
The hybrid tea group has a low nucleotide diversity (average $\pi_{observed}=1.21 \times 10^{-2}$; Fig.
2D), a level that is roughly similar to those of the Asian background ($\pi_{observed}=1.20 \times 10^{-2}$). This
diversity is much lower than expected assuming the inferred genomic composition (74.8%
Asian, 25.2% European, $\pi_{expected}=1.32 \times 10^{-2}$), corresponding to an average Reduction of
Diversity index (RoD) of 8.1%. Given that this breeding history occurred in Europe and that
hybrid tea roses have then been preferred to the early European x Asian hybrids in the gardens
of residential homes, the evolution between hybrid tea roses and the other European groups
is informative about the genetic loss during this period. Comparing these two groups, hybrid
teas exhibit an important reduction of diversity as compared to the early European x Asian
roses (average RoD=18.5%) and even more as compared to the ancient European accessions
(average RoD=27.5%). One could argue that hybrid teas were exported worldwide, including
in Asia and therefore that this history of selection could have contributed to an increase of the
genetic diversity in Asia. Considering this hypothesis, we compared ancient Asian and hybrid
teas and observed such a potential - albeit marginal - gain (RoD= -1.04%). Our results are
therefore overall consistent with a net and rapid erosion of nucleotide diversity in Europe during
the rose breeding of the 19th century.
The pattern of reduction of diversity between groups is consistent across the
chromosomes, even if it varies in intensity (Fig. S11). Comparing the diversity of hybrid teas
as compared to ancient European roses, 2.3% of the windows exhibit a reduction of at least
half of the diversity (105 among 4,628 windows with RoD exceeding 50%), a value that
appears particularly strong given the extremely limited number of generations (Fig. 2C). The
maximum observed RoD value on a window is 87.3%, but corresponds to an unanchored
scaffold (Chr00) and should therefore be interpreted cautiously. The maximum value for a
window anchored on a chromosome is 77.4% (Chr6: 11.2-11.3 Mb). Among the 60 windows
with a ROD>0.5 and that are not located on unanchored scaffolds, 35% (21) and 43% (26) are
located on chromosomes 3 and 5, respectively, suggesting that these two chromosomes were
particularly targeted by artificial selection.
680
681
682   **Detecting local genomic footprints of artificial selection**
683
To know more about the history of rose breeding, we investigated how the genomic
landscapes of diversity and Tajima's D have evolved during the rose breeding history (Fig. 3).
All landscapes of nucleotide diversity are highly correlated between groups (Fig. S12), even
those of the ancient European and ancient Asian groups (Pearson's r=0.301, p=3.57e^{-87}).
However, the Tajima's D landscapes of these two groups are independent (Pearson's r=-0.02,
p=0.20), suggesting that the correlations of the genomic landscape of diversity are not
associated with similar selection targets, but rather more likely shaped by some long-term
shared genomic features (*e.g.* the recombination landscape, Shang *et al.,* 2023). The genomic
landscapes of diversity become more and more correlated with time, especially when
compared to the ancient Asian background (ancient Asian vs. early European x Asian
samples, Pearson's r=0.408, p=1.44e^{-165}; ancient Asian vs. hybrid tea samples, Pearson's,
r=0.566, p=0). Consistently, the Tajima's D landscapes of the hybrid tea is more correlated

with the one of the ancient Asian (Pearson's, r=0.339, p9.26e^{-112}) than with the one of ancient
European samples (Pearson's, r=0.170, p=2.70e^{-28}). However the high level and significance
of the latter correlation also suggest that artificial selection targeted alleles from European
origin in some genomic regions.

Interestingly, the genomic landscapes of nucleotide diversity and Tajima's D are
significantly correlated in ancient European (Pearson's r=0.061, p=9.78e^{-5}), as well as in
ancient Asian samples (Pearson's r=0.430, p=2.79e^{-185}). However this correlation largely
differs in intensity between the two groups. This result is consistent with the lower diversity
(Fig. 2D) and the larger variance in Tajima's D in the ancient Asian group (Fig. S13). All
together, these results suggest that ancient Asian samples were more impacted by artificial
selection prior to their use in the European plant breeding programs than the ancient European
samples. Our genomic evidence could be interpreted with regards to the documented
knowledge of the long-term history of rose cultivation in China. Chinese roses have been
widely cultivated since the Han Dynasty, and selection is known to have occurred during this
period, at least for some important traits, including the extension of the flowering period.

In order to identify some local genomic footprints of selection, we used hybrid tea roses
as references. Indeed, given the limited number of generations of breeding detected (Fig. 2C),
the use of hybrid teas allows us to maximize the probability of identifying local footprints of
artificial selection. Hybrid tea genomes were scanned for windows exhibiting a shared signal
of (i) low π, (ii) negative Tajima's D and (iii) high RoD π$_{Hybrid Tea}$ *vs.* either π$_{Asian}$ or π$_{European}$
(Figs. 3; see Materials & Methods for details). The three summary statistics do not fully overlap
in their detection capabilities, each offering distinct specificities for detection (Fig S14).
Eighteen windows were detected using the most stringent criteria (last centile of the three
metrics, purple lines in Fig. 3), including 15 on a 4-Mb region of the chromosome 3 (from 24.1
and 28.2 Mb, see Table S4). This region is in the immediate vicinity of a previously
characterized 5Mb-inversion containing the *RoKSN* gene (28.8 - 33.1 Mb, Kawamura *et al.,*
2022), a presumably main breeding target to obtain recurrent-blooming roses with an extended
period of blooming. In addition, to these 18 windows, we detected 81 additional windows (last
five centiles of the three metrics, pink lines in Fig. 3). Remarkably, five of these additional
windows are located at the very end of the inversion (32.5 - 33.1Mb, see Table S4). The three
additional windows detected using the most stringent filtering criteria are located on
chromosome 5 (two adjacent windows, between 22.9 and 33.1 Mb) and on chromosome 1
(33.6-33.7 Mb, Table S4). Most of the additional regions based on the less stringent criteria
(last five centiles, see Table S4) are consistent with these three previously detected genomic
regions (*i.e.* on chromosomes 1, 3 and 5), but also highlighted a fourth region on chromosome
7 (11 windows between 15.9 and 21.1 Mb, Fig. 3 and Table S4).
732
733
734

Fig. 3: Detection of regions exhibiting footprints of artificial selection in hybrid tea roses. *From external to internal: nucleotide diversity, Tajima's D, RoD Hybrid tea vs. Ancient European, and local ancestry as estimated at diagnostic SNPs. Blue and yellow refer to ancient European and ancient Asian backgrounds respectively, while dark and light green refer to early European x Asian and hybrid tea groups, respectively. Pink and purple lines correspond to windows that simultaneously exhibit one of the lowest nucleotide diversity (bottom 5% or bottom 1%, respectively), one of the lowest Tajima's D (bottom 5% or bottom 1%, respectively) and one of the highest RoD (either in RoD European vs. Hybrid tea or RoD Asian vs Hybrid tea, top 5% or top 1%, respectively). Results from chr00 (corresponding to all unanchored scaffolds in Hibrand-Saint Oyant et al., 2018) are not shown.*

The largest rose GWAS catalog to date

In addition to reconstructing the history of rose breeding, we conducted a large Genome-Wide Association Study (GWAS) using GWASpoly (Rosyara *et al.,* 2016) with the aim of identifying associations between the genotypes at the 204 varieties and the different phenotypes, accounting for the levels of population structure and kinship among individuals. All GWAS was performed to identify associations at 46,691 markers, which corresponds to the fraction of markers from the SNP array with a missing rate lower than 10% and a MAF exceeding 0.05. Given the large number of phenotypes we investigated, as well as the variety of scenarios possible regarding dominance in polyploids (Rosyara *et al.,* 2016), a dedicated

website portal has been created in order to explore all the results (hereafter referred to as the rose GWAS browser, https://roseGWASbrowser.Z.io/). The online browser allows to select a category of traits (*e.g.* petal counts or colors, plant architecture, scent), then a specific trait (*e.g.* 2-phenylethanol content), and select a specific scenario regarding the polyploid gene action, among additive, simplex dominant and duplex dominant scenarios. By default, the rose GWAS browser reports the most general type of genetic model ("general model"), a model that has the advantages of making no assumptions regarding the underlying gene action, since the effect for each genotype class can be arbitrary (*e.g.* associations due to ABBB genotypes as compared to all others). General models have the advantage of providing an overview of the results for the other models, explaining also their use for Fig. 4. However, as previously indicated, results from the other models, which are more explicit regarding the gene action should be investigated in complement. Given that the number of SNPs also vary along the genome, we also use a local enrichment in associated SNPs to more easily identify associated regions. All the results from the GWAS were also made available on the Zenodo repository, including QQplots, linear Manhattan plots, as well as all p-values and q-values. For some traits, the GWAS explorer not only reports the results of the GWAS when the trait was analyzed quantitatively, but also when analyzed qualitatively, which can be meaningful when distribution of the trait values was observed to be more bimodally distributed.

The rose GWAS explorer is a tool designed to accelerate research on rose breeding. As a textbook example, we have decided to focus on the identification of genomic regions that correlated with the resistance to the blackspot disease, which is one of the most impactful diseases in garden roses. To identify some regions, we performed GWAS for three years of scoring (Fig. 4, see also Fig. S15). Based on the general model of GWASpoly, the number of significantly associated SNPs greatly varies depending on the year of scoring (red dots, Fig. 4). Additional associated regions are also observed on some other chromosomes for some of the years, including on a region of chromosome 3 that collocates with a meta-QTL recently identified in three different experimentally segregating populations (Lopez Arias *et al.,* 2020). It should be noted that the associated QQplots (Fig. S16) can also highlight an inflation of p-values, which varies in intensity from one model to another, but that can be substantial for the general model (Fig. S16). To facilitate interpretation of the results in this context, we included local enrichments of associated SNPs (*i.e.* number of associated SNPs relative to the total number of SNPs within a 1-Mb window, inner track, Fig. 4). However, this limitation related to p-value inflation should be considered when interpreting the results for this trait, as well as for the other traits displayed on the Rose GWAS browser.

Discussion

Evolution of the genetic diversity in roses

Genetic diversity is one fundamental unit of biodiversity since sufficient within-species diversity is needed to face different threats, including diseases, pests, changes in climate, as well as more intrinsic factors associated with evolution under low effective population sizes, and therefore reduce mid- and long-term extinction risks. The adopted post-2020 framework of the Convention on Biological Diversity (CBD, 2022) includes central ambitions regarding the maintenance and the restoration of the genetic diversity in wild species, but also in domesticated species, in order to safeguard their adaptive potential (Goal A & Target 4). It has

been recently reported a sharp reduction of the genetic diversity in bread wheat, with a loss of diversity of around 30% over the 20th century (Pont *et al.,* 2019). For ornamentals, for which domestication is far more recent (Purugganan, 2022), little is known about the large-scale genomic impacts of plant breeding.

Figure 4: GWAS for black spot resistance. *1: Per-window SNP array density and gene density based on the reference genome (Hibrand-Saint Oyant et al., 2018). 2-4: Manhattan plots of the p-values along the genome for the three years of scoring (2: 2014, 3: 2015, 4: 2016) under the general model of GWASpoly. SNPs with q-values < 0.20, 0.10 and 0.05 are shown in yellow, orange and red, respectively. 5: Local density in highly associated SNPs (p-value < 0.01) for each of the three years of scoring (2014, 2015 and 2016 are shown with blue, green and purple lines). QTLs and metaQTLs from Lopez-Arias et al., 2020 based on three segregating populations (connected by male resistant parent Rosa x wichurana) are also shown. The candidate R genes correspond to an expert annotation of NBS-LRR encoding genes (Lopez-Arias et al., 2020). Results for SNPs located on chr00 (corresponding to all unanchored scaffolds in Hibrand-Saint Oyant et al., 2018) are not shown here (but see Fig. S15 for non-circular Manhattan plots of the p-values including chr00).*

In the present study, we have questioned how the genetic diversity has evolved during
the 19th century, a century corresponding to the golden age for European rose breeding, with
an exploding number of varieties (estimated to be around 100 in 1800 and to be 8,000 in 1900;
Marriott, 2003, Oghină-Pavie, 2021). Our result is consistent with a substantial reduction of
diversity, of around 27.5% in a century in Europe (Fig. 2D), comparing the ancient European
and the hybrid tea roses. This reduction has likely taken place across a very limited number
of generations (< 10, Fig. 2C, but see also Supplementary Note 2) and is associated with both
the backcrossing with a pre-selected Asian genetic background (Figs. 2 & S6-9) and with
additional footprints of artificial selection (Fig. 3).
832
Although this observed reduction of genetic diversity could be considered as more
minor regarding a horticultural species as compared to a crop species such as bread wheat,
it still holds implications for humans. Indeed, roses represent a common heritage of mankind
and serve as a significant cornerstone in the ornamental sector, in addition to being used as a
source of food and medicine in some societies. In roses, grafting however makes the long-
term maintenance of ancient varieties possible, allowing the backcrossing of modern varieties,
improved regarding some specific traits, with more ancient and genetically diverse varieties.
This restoration of this genetic diversity is however only possible if the ancient garden roses
are maintained over the long term. In roses, two main action targets should be therefore
proposed: (i) increasing effort, especially regarding state funding, for safeguarding gardens
containing botanical and ancient varieties, and (ii) alert the stakeholders of the importance of
maintaining and restoring the genetic diversity in order to ensure a sustainable breeding for
long-term, which could be associated with new future breeding targets (*e.g.* new emerging
diseases, anthropogenic disturbances or ecosystemic services in urban areas).
847
848

849   **Looking back in time to identify selection targets**
850
Another tremendous advantage of the use of grafted garden roses as domesticated
plant models is our ability to phenotype varieties bred during more than a century all together
at the same time in a common rose garden. Phenotypically, we indeed have reported important
temporal changes in the number of petals, the ability of roses to bloom several times in a year
and the susceptibility to the black spot disease (Fig. 1), even if the latter trait evolution was
likely unintended. The temporal increase in the number of petals is relatively continuous until
1870, before a slight reduction on average. This is consistent with a change in aesthetic
preferences, since (i) varieties with up to 60 petals are nowadays preferred to obtain high-
centered buds when the rose flowers are opened, (ii) the rosomania had led to the
establishment of rose connoisseur communities from 1880, interested by new targets of
selection, including wild-like phenotypes such as single flower (five petals) and spring roses
(Oghina-Pavie, 2020). Indeed, at the end of the 19th century, additional rose species were
used as breeding materials, such as *R. multiflora* and *R. rugosa*, with single flower and non-
recurrent blooming phenotypes. This would be consistent with the apparent progression
toward a bimodal distribution for this specific trait and questions the diversification of the
consumer tastes at that time, including the return to simple flowers for some varieties (Fig. 1;
see also Oghina-Pavie, 2020). Roses of a given period should not be considered as a
homogeneous ensemble, they are rather representative of different tastes and breeding
objectives among breeders, in such a way that the history of breeding should not be
oversimplified. Similarly to petal counts, the susceptibility to black spot has increased until

1890, before a reduction on average, again associated with a marked bimodal distribution (Fig.
1).

873   The extension of the flowering period by selecting recurrent blooming varieties was
often assumed to be the main target of selection during the 19th century. Consistently, during
this period, we indeed observed a continuous increase of an index that measures the ability
of a rose to rebloom (Fig. 1). Such a continuous selection has likely contributed to generate a
marked footprint of selection in rose genomes, since the sharpest signature of the selective
sweep is observed above one of the two breaking points (chromosome 3, 24.6-28.2) of a
previously characterized 5-Mb inversion of the *RoKSN* gene, *RoKSN*[NULL] allele (chromosome
3, 28.8 - 33.1 Mb, Hibrand-Saint Oyant *et al.,* 2018; Kawamura *et al.,* 2022). However, using
the less stringent criteria for detecting sweeps, five additional windows were detected near the
second breaking point of the inversion (32.5 - 33.1Mb, see Table S4), therefore providing
additional support that this footprint is associated with the artificial selection of Asian alleles at
RoKSN. Consistent signal was also observed at the second breaking point (Fig. 3), in
agreement with the selection in this region. Several alleles at the *RoKSN* gene leading to the
extended period were described, including a *RoKSN*[copia] and a *RoKSN*[NULL]. Previously, it was
demonstrated a progressive selection of the *RoKSN*[copia] allele during the 19th century (Soufflet-
Freslon *et al.,* 2021). This allele is not present in the reference genome we used, this genome
contains a large rearrangement at the *RoKSN* locus leading to the *RoKSN*[NULL] allele (Hibrand-
Saint Oyant *et al.,* 2018). In the immediate vicinity of the inversion, we find also an homologue
of *AP2/TOE* (RC3G0243000, 33.2Mb), another crucial gene controlling the double flower
phenotype and therefore partly the number of petals (François *et al.,* 2018; Hibrand-Saint
Oyant *et al.*, 2018). However, *AP2/TOE* is unlikely to be the gene responsible for this footprint
(see Supplementary Note 1), suggesting that recurrent-flowering alleles of RoKSN were the
main targets of artificial selection. This rearrangement may explain the complexity of the local
footprints associated with the nucleotide diversity and Tajima's D landscapes. Additional
footprints have been also uncovered, but currently were not associated with any previously
reported genes. In the future, thanks to more numerous whole-genome sequences available,
as well as the new technological and methodological developments allowing the access to
phased polyploid genomes, a much clearer picture could be anticipated. Extending these
investigations to the 20th century could be of particular interest since more modern roses are
characterized by a clearer distinction between the genetics used for garden *vs.* cut roses. Cut
roses - especially red ones - have potentially experienced a more intense reduction of global
genetic diversity and more pronounced local footprints of selection, as typical expected based
on the huge difference of the two respective markets (*e.g.* in France in 2018, garden roses:
50M€, cut roses: 376M€, Clotault *et al.*, 2022).

907
908
909 **Advances and challenges for GWAS in rose**
910
911   As a tool for the future of rose breeding, we have released the largest GWAS catalog
for roses (https://roseGWASbrowser.github.io/). Our GWAS browser allows the website
visitors to explore the association with more than 25 traits, on the basis of our panel of 204
accessions. The objective is to make future breeding more effective for some traits, including
the resistance to the black spot disease (Fig. 4). This trait represents a typical example of an
important target need in order to find sources of genetic resistances in roses, in the context of
a tightening environmental legislation associated with the use of chemicals due to their
deleterious environmental impacts and health risks.

Despite the great advance made with the release of the GWAS catalog, the direct
insights appear relatively limited to date. Indeed, the GWAS strategy in our panel has
generated few repeatable and easily interpretable signatures of associations. Overall, the
relative lack of resolution is likely associated with the difficulty of performing GWA studies in
populations for which phenotypes still highly correlate with the population structure gradient
as a result of the limited number of generations (*e.g.* see Fig. S4). Indeed, similar results were
observed for most of the investigated traits. As a consequence, all the results should be
interpreted with some caution, requiring further validation through independent analyses.
Regarding the black score disease, some potential genomic regions of interest were detected,
including a genomic region at the very end of chromosome 4 that is associated with the three
years of scoring (Fig. 4). Additional locations are also observed on chromosome 3 for some of
the years, including in a region that collocates with a meta-QTL recently identified in three
different experimentally segregating populations, which were phenotyped for black spot
disease over several years and locations (Lopez Arias *et al.,* 2020).
As compared with the use of QTL mapping in experimental crosses, extensive GWAS
analyses seem to have a more limited benefit in rose than on some other domesticated or non-
domesticated species (*e.g.* Korte & Farlow, 2013; Alqudah *et al.,* 2020). Several non-mutually
exclusive reasons could explain this paradox: (i) a too limited number of varieties used in our
study, (ii) the limited number of generations of recombination in the investigated collection (Fig.
2C) that could contribute to maintaining a predominant association with the population
structure, leading to a low statistical power in the model. Combining GWAS and QTL
approaches (Fig. 4), when available, offers the possibility to increase the statistical support for
some regions of interest, extends the allelic diversity by considering other alleles than those
segregating between the parents, describes the genomic locations of the associated regions
and prioritizes the research effort. As a typical example, the last third of chromosome 3 could
appear as a more promising region for selection, therefore guiding the research efforts in this
direction. Remarkably, chromosome 3 has been associated with many traits in rose
(resistance to black spot, but also number of petals, fragrance, prickles, recurrent blooming,
self-incompatibility; *e.g.* Hibrand-Saint Oyant *et al.,* 2018, Kawamura *et al.,* 2022), which could
therefore suggests the presence of tradeoffs between some traits of interest. A more detailed
genomic characterization of the genomic landscapes, including local linkage disequilibrium
patterns, based on more modern varieties, is especially needed on this chromosome.
951
952

Conclusion

954
Combining large phenotypic data and genetic investigations using 204 individuals
genotyped using a high density SNP array, plus 32 whole-genome sequences, we found
support for both phenotypic and genomic changes during the 19th century. With an
unprecedented accuracy, we confirmed the progressive shift from a European to a more Asian-
like genetic background, with Hybrid tea roses exhibiting a three-quarter Asian genetic
background. This genetic makeup likely corresponds to several generations of backcrossing
of the early European x Asian varieties with the ancient genetic background, as it was
previously suggested (Liorzou *et al.,* 2016, see also Supplementary Note 3). As a result of the
lower resident diversity in the ancient Asian varieties, partly due to a more ancient and intense
history of selection in Asia, as well as to some specific genomic footprints of artificial selection
- especially in the region of *RoKSN* -, rose breeding in Europe has contributed to a substantial

loss of genetic diversity during the 19th century (up to 27.5% in hybrid tea roses). In addition
to providing a fine description of the history of rose breeding, we have generated the largest
GWAS catalog to date, with the aim of providing fundamental knowledge for the future of rose
breeding.
970
971

Acknowledgments

This present article is dedicated to the memory of our cherished colleague, Laurence Hibrand-
Saint Oyant, who passed away in November 2024 (see our dedication section, Supporting
Information). We acknowledge the Roseraie Loubert (Gennes-Val-de-Loire, France) and
Biological Resource Center Pome Fruits and Roses (Angers, France) for maintaining the
genetic resources and their associated data. We also acknowledge the Gentyane facility
(INRAE Clermont-Ferrand, France) and the ANAN platform of the SFR Quasav (Angers,
France; including Muriel Bahut) for the genotyping data. We would like to thank the Next
Generation Sequencing Facility at Vienna BioCenter Core Facilities (VBCF), member of the
Vienna BioCenter (VBC), Austria. We are grateful to the genotoul bioinformatics platform
Toulouse Occitanie (Bioinfo Genotoul, https://doi.org/10.15454/1.5572369328961167E12) for
providing computing and storage resources. We also thank Jérôme Chené (Pépinières Roses
Loubert, Gennes-Val-de-Loire, Pays-de-la-Loire, France) for providing fresh plant material of
a few whole-genome resequenced individuals. We also thank Roeland Voorrips and Giorgio
Tumino for their valuable advice regarding fitPoly and GWASpoly at the beginning of the
project.

989
990

Funding

This research was conducted in the framework of the regional programme "Objectif Végétal,
Research, Education and Innovation in Pays-de-la-Loire", supported by the French Region
Pays-de-la-Loire, Angers Loire Métropole and the European Regional Development Fund. A
part of the phenotypic data was supported by the FLORHIGE (French Region Pays-de-la-
Loire), the SIFLOR (BAP division of INRAE) and Rosascent projects (ANR, grant number
ANR-16-CE20-0024-01). Thibault Leroy is also grateful to the university of Vienna, Austria for
its financial support regarding the whole-genome sequencing of the 15 rose varieties.

1000
1001

Competing interests

1004 The authors declare no competing interest. Although E.A. is a Syngenta seeds SAS employee,
1005 her contribution to the project predates her recruitment by the company and does not pertain
1006 to her current activities or the interests of the company.
1007
1008

Author contributions

1011 T.Le., E.A., S.B., J-C.C., C.O.P, F.F., L.H.S.O & J.Cl. designed research; T.Le., E.A., T.T.,
1012 A.C, J.Ch, T.Lo., A.P., V.S.F, F.F., L.H.S.O & J.Cl contributed to the sampling; A.C. & L.H.S.O.

performed DNA extractions; J.J. performed the *RoKSN* and *RoAP2 genes* genotyping; T.T., J.C., T.Lo., A.P. & V.S.F contributed to the phenotyping; S.B., J-C.C. & S.N.P. contributed to the GC-mass spectrometry data; T.Le & E.A. analyzed data under the supervision of C.O.P, F.F., L.H.S.O. & J.Cl.; T.Le. drafted the manuscript. All authors have subsequently read, edited and approved the manuscript.

Data availability

Raw sequencing data of the newly sequenced roses are publicly available on the Sequence Read Archive (SRA) under the BioProject PRJNA997103. Scripts were made available on Zenodo (https://doi.org/10.5281/zenodo.14450241). The GWAS rose browser is available at https://roseGWASbrowser.github.io.

References

Allaby RG, Ware RL, Kistler L. 2019. A re-evaluation of the domestication bottleneck from archaeogenomic evidence. *Evolutionary Applications* 12: 29–37.

Alqudah AM, Sallam A, Stephen Baenziger P, Börner A. 2020. GWAS: Fast-forwarding gene identification and characterization in temperate Cereals: lessons from Barley – A review. *Journal of Advanced Research* 22: 119–135.

Altman A, Shennan S, Odling-Smee J. 2022. Ornamental plant domestication by aesthetics-driven human cultural niche construction. *Trends in Plant Science* 27: 124–138.

Augusto Corrêa Dos Santos R, Goldman GH, Riaño-Pachón DM. 2017. ploidyNGS: visually exploring ploidy with Next Generation Sequencing data. *Bioinformatics (Oxford, England)* 33: 2575–2576.

Bergougnoux V, Caissard J-C, Jullien F, Magnard J-L, Scalliet G, Cock JM, Hugueney P, Baudino S. 2007. Both the adaxial and abaxial epidermal layers of the rose petal emit volatile scent compounds. *Planta* 226: 853–866.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics (Oxford, England)* 30: 2114–2120.

Brown TA. 2019. Is the domestication bottleneck a myth? *Nature Plants* 5: 337–338.

Caissard J-C, Adrar I, Conart C, Paramita SN, Baudino S. 2023. Do we really know the scent of roses? *Botany Letters* 170: 77–88.

Chowdhuri TK, Deka K. 2019. Biodiversity and Conservation of Ornamental Crops. *Conservation and Utilization of Horticultural Genetic Resources*.

Clotault J, Foucher F, Grapin A, Hibrand-Saint Oyant L, Malécot V, Paillard S, Pernet A, Soufflet-Freslon V, Oghina-Pavie C. 2022. Rose selection in France: a long and beautiful history. *Chronica Horticulturae* 62: 28–37.

Convention on Biological Diversity. 2022. COP15: Final text of Kunming-Montreal Global Biodiversity Framework.

Csárdi G, Nepusz T. 2006. The igraph software package for complex network research.

Debray K, Le Paslier M-C, Bérard A, Thouroude T, Michel G, Marie-Magdelaine J, Bruneau A, Foucher F, Malécot V. 2022. Unveiling the Patterns of Reticulated Evolutionary Processes with Phylogenomics: Hybridization and Polyploidy in the Genus Rosa. *Systematic Biology* 71: 547–569.

Diamond J. 2002. Evolution, consequences and future of plant and animal domestication. *Nature* 418: 700–707.

Diamond JM. 2005. *Guns, germs, and steel: the fates of human societies*. New York: Norton.

Ducomet, V. 1903. Une maladie des feuilles du Rosier *Marsonia Rosae* Briosi et cav. in Le Jardin: 199.

François L, Verdenaud M, Fu X, Ruleman D, Dubois A, Vandenbussche M, Bendahmane A, Raymond O, Just J, Bendahmane M. 2018. A miR172 target-deficient AP2-like gene correlates with the double flower phenotype in roses. *Scientific Reports* 8: 12912.

Gattolin S, Cirilli M, Pacheco I, Ciacciulli A, Da Silva Linge C, Mauroux J-B, Lambert P, Cammarata E, Bassi D, Pascal T, *et al.* 2018. Deletion of the miR172 target site in a TOE-type gene is a strong candidate variant for dominant double-flower trait in Rosaceae. *The Plant Journal* 96: 358–371.

Gaut BS, Díez CM, Morrell PL. 2015. Genomics and the Contrasting Dynamics of Annual and Perennial Domestication. *Trends in Genetics* 31: 709–719.

Gautier M, Vitalis R, Flori L, Estoup A. 2022. f-Statistics estimation and admixture graph construction with Pool-Seq or allele count data using the R package poolfstat. *Molecular Ecology Resources* 22: 1394–1416.

Gessert G. 1993. Flowers of Human Presence: Effects of Esthetic Values on the Evolution of Ornamental Plants. *Leonardo* 26: 37–44.

Gessert G. 2010. *Green Light: Toward an Art of Evolution*. The MIT Press.

Goody J. 1993. *The Culture of Flowers*. Cambridge University Press.

Gu Z, Gu L, Eils R, Schlesner M, Brors B. 2014. circlize implements and enhances circular visualization in R. *Bioinformatics* 30: 2811–2812.

Hibrand Saint-Oyant L, Ruttink T, Hamama L, Kirov I, Lakhwani D, Zhou NN, Bourke PM, Daccord N, Leus L, Schulz D, *et al.* 2018. A high-quality genome sequence of Rosa chinensis to elucidate ornamental traits. *Nature Plants* 4: 473–484.

Hivert V, Leblois R, Petit EJ, Gautier M, Vitalis R. 2018. Measuring Genetic Differentiation from Pool-seq Data. *Genetics* 210: 315–330.

Iwata H, Gaston A, Remay A, Thouroude T, Jeauffre J, Kawamura K, Oyant LH-S, Araki T, Denoyes B, Foucher F. 2012. The TFL1 homologue KSN is a regulator of continuous flowering in rose and strawberry. *The Plant journal: for cell and molecular biology* 69: 116–125.

Jombart T. 2008. adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics* 24: 1403–1405.

Kawamura K, Ueda Y, Matsumoto S, Horibe T, Otagaki S, Wang L, Wang G, Hibrand-Saint
Oyant L, Foucher F, Linde M, *et al.* 2022. The identification of the Rosa S-locus provides new
insights into the breeding and wild origins of continuous-flowering roses. *Horticulture Research*
9: uhac155.

Koning-Boucoiran CFS, Esselink GD, Vukosavljev M, van 't Westende WPC, Gitonga VW,
Krens FA, Voorrips RE, van de Weg WE, Schulz D, Debener T, *et al.* 2015. Using RNA-Seq
to assemble a rose transcriptome with more than 13,000 full-length expressed genes and to
develop the WagRhSNP 68k Axiom SNP array for rose (Rosa L.). *Frontiers in plant science*
6: 249.

Korte A, Farlow A. 2013. The advantages and limitations of trait analysis with GWAS: a review.
Plant methods 9: 29.

Korunes KL, Samuk K. 2021. pixy: Unbiased estimation of nucleotide diversity and divergence
in the presence of missing data. Molecular Ecology Resources 21:1359-1368.

Leroy T, Anselmetti Y, Tilak M-K, Bérard S, Csukonyi L, Gabrielli M, Scornavacca C, Milá B,
Thébaud C, Nabholz B. 2021a. A bird's white-eye view on avian sex chromosome evolution.
Peer Community Journal 1.

Leroy T, Rousselle M, Tilak M-K, Caizergues AE, Scornavacca C, Recuerda M, Fuchs J, Illera
JC, De Swardt DH, Blanco G, *et al.* 2021b. Island songbirds as windows into evolution in small
populations. *Current Biology* 31: 1303-1310.e4.

Leus L, Van Laere K, De Riek J, Van Huylenbroeck J. 2018. Rose. In: Van Huylenbroeck J,
ed. Ornamental Crops. Cham: Springer International Publishing, 719–767.

Liorzou M, Pernet A, Li S, Chastellier A, Thouroude T, Michel G, Malécot V, Gaillard S, Briée
C, Foucher F, *et al.* 2016. Nineteenth century French rose (Rosa sp.) germplasm shows a shift
over time from a European to an Asian genetic background. *Journal of Experimental Botany*
67: 4711–4725.

Lopez Arias DC, Chastellier A, Thouroude T, Bradeen J, Van Eck L, De Oliveira Y, Paillard S,
Foucher F, Hibrand-Saint Oyant L, Soufflet-Freslon V. 2020. Characterization of black spot
resistance in diploid roses with QTL detection, meta-analysis and candidate-gene
identification. *Theoretical and Applied Genetics* 133: 3299–3321.

Magnard J-L, Roccia A, Caissard J-C, Vergne P, Sun P, Hecquet R, Dubois A, Hibrand-Saint
Oyant L, Jullien F, Nicolè F, *et al.* 2015. Biosynthesis of monoterpene scent compounds in
roses. *Science* 349: 81–83.

Manichaikul A, Mychaleckyj JC, Rich SS, Daly K, Sale M, Chen W-M. 2010. Robust
relationship inference in genome-wide association studies. *Bioinformatics (Oxford, England)*
26: 2867–2873.

Marolleau B, Petiteau A, Bellanger M-N, Sannier M, Le Pocreau N, Porcher L, Paillard S,
Foucher F, Thouroude T, Serres-Giardi L, *et al.* 2020. Strong differentiation within Diplocarpon
rosae strains based on microsatellite markers and greenhouse-based inoculation protocol on
Rosa. *Plant Pathology* 69: 1093–1107.

Marriott M. 2003. Encyclopedia of Rose Science.

Martin M, Piola F, Chessel D, Jay M, Heizmann P. 2001. The domestication process of the
Modern Rose: genetic structure and allelic composition of the rose complex. *Theoretical and*
Applied Genetics 102: 398–404.

Moyers BT, Morrell PL, McKay JK. 2018. Genetic Costs of Domestication and Improvement.
Journal of Heredity 109: 103–116.

Nei M, Maruyama T, Chakraborty R. 1975. The bottleneck effect and genetic variability in
populations. *Evolution* 29: 1–10.

Oghină-Pavie C. 2020. La raison des rosomanes. Les sciences naturelles dans le Journal des
Roses à la fin du XIXe siècle. *Romantisme* 190: 64-78.

Oghină-Pavie C. 2021. Les rosiers et la nouveauté horticole en France au début du XIXe
siècle, in J. Baudry, J. Blanc, L. Hilaire-Pérez, M. Ratcliff, S. Wenger (eds), Produire du
nouveau ? Arts-techniques-sciences en Europe (1400-1900), Paris, CNRS Éditions: 83-94.

Picard toolkit. 2019. *Broad Institute, GitHub* repository https://broadinstitute.github.io/picard/

Pline l'Ancien. 2013. *Histoire naturelle*. Paris: Gallimard (traduction).

Pont C, Leroy T, Seidel M, Tondelli A, Duchemin W, Armisen D, Lang D, Bustos-Korts D, Goué
N, Balfourier F, *et al.* 2019. Tracing the ancestry of modern bread wheats. *Nature Genetics*
51: 905–911.

Poplin R, Ruano-Rubio V, DePristo M, Fennell J, Carneiro M, Auwera G, Kling D, Gauthier L,
Levy-Moonshine A, Roazen D, *et al.* 2018. Scaling accurate genetic variant discovery to tens
of thousands of samples. *bioRxiv*: 201178.

Proïa F, Pernet A, Thouroude T, Michel G, Clotault J. 2016. On the characterization of
flowering curves using Gaussian mixture models. *Journal of Theoretical Biology* 402: 75–88.

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, de
Bakker PIW, Daly MJ, *et al.* 2007. PLINK: a tool set for whole-genome association and
population-based linkage analyses. *American journal of human genetics* 81: 559–575.

Purugganan MD. 2022. What is domestication? *Trends in ecology & evolution* 37: 663–671.

Raj A, Stephens M, Pritchard JK. 2014. fastSTRUCTURE: variational inference of population
structure in large SNP data sets. *Genetics* 197: 573–589.

Raymond O, Gouzy J, Just J, Badouin H, Verdenaud M, Lemainque A, Vergne P, Moja S,
Choisne N, Pont C, *et al.* 2018. The *Rosa* genome provides new insights into the domestication
of modern roses. *Nature Genetics* 50: 772–777.

Roccia A, Hibrand-Saint Oyant L, Cavel E, Caissard J-C, Machenaud J, Thouroude T, Jeauffre
J, Bony A, Dubois A, Vergne P, *et al.* 2019. Biosynthesis of 2-Phenylethanol in Rose Petals Is
Linked to the Expression of One Allele of RhPAAS. *Plant physiology* 179: 1064–1079.

Rosyara UR, De Jong WS, Douches DS, Endelman JB. 2016. Software for Genome-Wide
Association Studies in Autopolyploids and Its Application to Potato. *The Plant Genome* 9:
plantgenome2015.08.0073.

Schulz DF, Schott RT, Voorrips RE, Smulders MJM, Linde M, Debener T. 2016. Genome-
Wide Association Analysis of the Anthocyanin and Carotenoid Contents of Rose Petals.
Frontiers in plant science 7: 1798.

Shang H, Field DL, Paun O, Rendón-Anaya M, Hess J, Vogl C, Liu J, Ingvarsson PK, Lexer
C, Leroy T. 2023. Drivers of genomic landscapes of differentiation across a *Populus*
divergence gradient. *Molecular Ecology*, in press.

Slowikowski K. 2023. ggrepel: Automatically Position Non-Overlapping Text Labels with 'ggplot2'.

Smith O, Nicholson WV, Kistler L, Mace E, Clapham A, Rose P, Stevens C, Ware R, Samavedam S, Barker G, *et al.* 2019. A domestication history of dynamic adaptation and genomic deterioration in Sorghum. *Nature Plants* 5: 369–379.

Soufflet-Freslon V, Araou E, Jeauffre J, Thouroude T, Chastellier A, Michel G, Mikanagi Y, Kawamura K, Banfield M, Oghina-Pavie C, *et al.* 2021. Diversity and selection of the continuous-flowering gene, RoKSN, in rose. *Horticulture Research* 8: 76.

Spiller M, Berger RG, Debener T. 2010. Genetic dissection of scent metabolic profiles in diploid rose populations. *Theoretical and Applied Genetics* 120: 1461–1471.

Touw M. 1982. Roses in the Middle Ages. *Economic Botany* **36**: 71–83.

Turner S. 2018. qqman: an R package for visualizing GWAS results using Q-Q and manhattan plots. Journal of Open Source Software 3:731

Vasimuddin M, Misra S, Li H, Aluru S. 2019. Efficient Architecture-Aware Acceleration of BWA-MEM for Multicore Systems. In: 314–324.

Voorrips R.E., Gort G. Vosman B. 2011. Genotype calling in tetraploid species from bi-allelic marker data using mixture models. BMC Bioinformatics 12, 172.

Wickham H. 2016. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

Wissemann V, Ritz CM. 2007. Evolutionary patterns and processes in the genus Rosa (Rosaceae) and their implications for host-parasite co-evolution. *Plant Systematics and Evolution* 266: 79–89.

Wylie A. 1954. The history of garden roses. *Journal of the Royal Horticultural Society* 79: 555–571.

Zheng X, Levine D, Shen J, Gogarten SM, Laurie C, Weir BS. 2012. A high-performance computing toolset for relatedness and principal component analysis of SNP data. *Bioinformatics (Oxford, England)* 28: 3326–3328.