Flagellar switch inverted repeat sequence variation impacts *Clostridioides difficile* RT027/MLST1

virulence.

Nguyen T. Q. Nhu[a,b,#], Huaiying Lin[b], Jonathan K. Sia[c], Evan S. Snitkin[d], Vincent Young[d,e], Mini Kamboj[f], Eric

G. Pamer[a,b], Phoebe A. Rice[g], Qiwen Dong[a,b,#].

[a]Department of Medicine, University of Chicago, Chicago, Illinois, USA

[b]Duchossois Family Institute, University of Chicago, Chicago, Illinois, USA

[c]Immunology Program, Memorial Sloan Kettering Cancer Center, New York City, New York, USA

[d]Division of Infectious Diseases, Department of Internal Medicine, University of Michigan, Ann Arbor,

Michigan, USA

[e]Department of Microbiology & Immunology, University of Michigan, Ann Arbor, MI, USA

[f]Infection Control, Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, New

York, USA

[g]Department of Biochemistry and Molecular Biology, University of Chicago, Chicago, IL, USA

Running Head: Flagellar switch inverted repeats impact *C. difficile* ST1 virulence.

[#]Address correspondence to Qiwen Dong, dongq@uchicago.edu; Nguyen T. Q. Nhu, nhun@umich.edu

*Present address: Nguyen T. Q. Nhu, Division of Infectious Diseases, Department of Internal Medicine,

University of Michigan, Ann Arbor, Michigan, USA

ABSTRACT The impact of *Clostridioides difficile* strain diversity on the severity of intestinal infection and

diarrhea remains unclear. The RT027/ST1 strain is highly prevalent and has been associated with more

severe disease and increased toxin production. Previous studies using antibiotic-treated mice, however,

have demonstrated that the severity of diarrhea and colonic inflammation caused by clinical RT027/ST1

C. difficile isolates cultured from different patients is isolate-specific and varies from avirulent to lethal.

Herein we demonstrate that increased flagellar expression is associated with increased in vivo virulence.

While sequencing the flagellar operon did not identify differences between high and low virulence

isolates, high virulence was associated with strains encoding higher proportions of the phase variable

flagellar switch in the ON mode. We found that the proportion of bacteria with the switch in the ON mode

correlates with maximum weight loss, a readout for virulence in mice. Further analysis of flagellar switch

sequences identified variant inverted repeat (IR) sequences, with 40% of isolates having lost one A or T in

the IR compared to the genome of *C. difficile* R20291. The isolates with the R20291 IR sequence have

increased virulence compared to those with variant IR (p = 3.2e-05) and these isolates have markedly

reduced ability to invert the flagellar switch. Our results suggest that a restricted capacity to invert the

flagellar switch during infection is associated with reduced virulence of *C. difficile* RT027/ST1 strains and

may account for some of the inconsistent associations between *C. difficile* infection and disease severity

in patients.

IMPORTANCE *C. difficile* RT027/ST1 has been associated with high virulence. However, variable virulence

among ST1 isolates has also been observed. Recent studies have shed light on the impacts of the gut

microbiota and host immunity during *Clostridioides difficile* infection (CDI). Beyond that, determining the

genetic factors of *C. difficile* regulating virulence is essential for a successful CDI diagnosis and treatment.

Here, we report a strong association between the flexibility of the flagellar switch and the virulence of *C.*

difficile. Specifically, the inverted repeats of the flagellar switch correlate with switch flexibility. Overall,

our study explores variations in *C. difficile* virulence and highlights the potential of using flagellar switch

inverted repeats as an easily accessible genetic trait to predict pathogen virulence.

KEYWORDS *Clostridioides difficile*, RT027, virulence, flagellar switch, inverted repeats, recombinase

INTRODUCTION

Clostridioides difficile (*C. difficile*) is the leading cause of hospital-acquired infection in the U.S., with an

estimated incidence of approximately 224.000 cases per year (1). The incidence of community-acquired

C. difficile infection (CDI) is also increasing, with up to 51.2 cases /100.000 population (2). Antibiotic-

mediated depletion of commensal bacterial species increases the risk of CDI and prolonged hospitalization

increases exposure to *C. difficile* spores. The spectrum of disease associated with *C. difficile* infection

ranges from asymptomatic colonization to mild/moderate diarrhea to severe pseudomembranous colitis

and death. It has been estimated that CDI causes approximately 12.800 deaths and costs the US Health

care system one billion U.S. dollars per year (1).

C. difficile virulence is mediated by two glucosylating toxins, TcdA and TcdB (3). These toxins are

internalized by epithelial cells by endocytosis (4-6) and are released into the cytosol where they

glucosylate and inactivate intracellular GTPases (7-9). These activities can disrupt cell signaling pathways

and cytoskeletal structure, leading to cell death (10). The impact of TcdA and TcdB on disease progression

and severity can be varied, depending on the strain of *C. difficile* and the host's innate and adaptive

immune defenses (11, 12).

The extent to which *C. difficile* strains vary in virulence is unclear. Because every patient with *C. difficile*

infection has a distinct residual microbiota and constellation of risk and resistance factors that impact the

severity of disease, it has been challenging to determine the impact of the pathogen's intrinsic virulence

on disease course (13, 14). The multi-locus sequence type 1 (ST1) strain, a.k.a. as NAP1/B1/ribotype 027,

is highly transmissible and has increased multidrug resistance (15, 16) and was initially reported to be

hyper-virulent (17, 18). Subsequent studies, however, suggested its virulence was similar to other strains

(19, 20). It emerged in the early 2000s in the USA and Canada then spread to European and South

American countries (21). Today, ST1 can be found worldwide. The virulence of ST1 has been attributed to

higher levels of toxin production (22). Subsequent studies, however, reported heterogeneity in toxin

production and virulence of ST-1 strains (23, 24).

In addition to toxin expression, bacterial motility mediated by flagella has been implicated in *C. difficile*

virulence. Similar to other gut-bacterial intestinal pathogens such as *Salmonella typhi* and *Escherichia coli,*

C. difficile is a peritrichous bacterium. In fact, the flagellar expression has been shown to contribute to

virulence by enhancing host cell invasion, adhesion to epithelial cells, and systemic inflammation of *S.*

enterica and *E. coli* (25-27). Deletion of flagellar structural genes or inactivation of the flagellar motor

resulted in distinct impacts on *in vivo* virulence, depending on the *C. difficile* strain (28). In *C. difficile*, the

deletion of *fli*C, which encodes flagellin, reduces the virulence of *C. difficile* R20291, an ST1 strain, in a

mouse model (29). In contrast, in *C. difficile* strain CD630, the deletion of *fli*C did not decrease virulence

in the hamster model of CDI (30). Recent studies demonstrated that flagellar expression is coupled with

TcdA/TcdB production, suggesting that toxin-mediated virulence is coupled to flagellin expression (31).

Whether flagellar expression contributes to the relative virulence of different *C. difficile* ST1 strains is

unclear.

In this study, we investigated the heterogeneity of *C. difficile* ST1 strains isolated by fecal culture from

patients with CDI. Testing of 22 ST1 clinical isolates in mice revealed marked differences in virulence. We

find that sequence differences in the flagellar switch region contribute to variations in ST1 strain virulence.

Specifically, we demonstrate strain diversity in the inverted repeat (IR) region of the flagellar switch

among ST1 strains. The proportion of ON versus OFF in the flagellar switch is determined by the sequence

of the IR. The published IR sequence in *C. difficile* R20291 enhances flexibility in switching between ON

and OFF but this IR sequence is not shared by all ST1 strains, many of which have reduced ability to alter

flagellar expression. Our results demonstrate that flexibility in flagellar switching mediated by the IR

contributes to heterogeneity in *C. difficile* ST1 virulence.

RESULTS

Highly virulent isolates express more flagella and demonstrate higher motility.

Previous studies characterizing *in vivo* virulence of 23 *C. difficile* ST1 isolates revealed disease severities in

infected C57BL/6 mice that ranged from mortality to absence of any weight loss or diarrhea (Dong et al.

in press). While the avirulence of 2 ST1 strains was attributed to a deletion in the cdtR gene (Dong et al.

in press), the pathogenicity and CDT loci that encode TcdA and TcdB glucosylating toxins and the CDT

binary toxin CDT were identical in the remaining 21 ST1 strains. To begin to identify additional mechanisms

for variable virulence among ST1 strains, we selected two strains that induced >10% weight loss (ST1-12

and ST1-53) and two strains that resulted in < 10% weight loss (ST1-6 and ST1-27) for further study (Fig.

S1). Transcriptomic profiling of these strains in cecal contents obtained 24 hours following infection of

antibiotic treated C57BL/6 mice demonstrated similar transcription of both glucosylating toxin and binary

toxin genes (Fig. 1 & Fig. S3). Similar *in vivo* toxin transcription among these strains was supported by toxin

profiling of fecal samples (Fig. S2).

Transcription of flagellar-related genes, on the other hand, was increased in highly virulent compared to

less virulent isolates (Fig. 1A), which was further validated by RT-qPCR (Fig. S3). To determine whether

flagellar-related gene transcription correlates with flagellar expression, we obtained transmission

electron microscopy (TEM) images of the four ST1 *C. difficile* strains grown on hard agar plates. High

virulence ST1 strains were found to be flagellated, with 4-5 peritrichous flagella, while low virulence ST1

isolates lacked detectable flagella (Fig. 1B).

To test the motility of ST1 *C. difficile* strains, we cultured bacteria in a hydrogel environment and measured

the spread of *C. difficile* colonies on 0.3% agar buffered with BHIS pH 7.0. Highly virulent ST1-12 and ST1-

53 strains produced larger colony sizes after one day of incubation than the reduced-virulence ST1-6 and

ST1-27 strains (Fig. 1C). We did not detect mutations in genes encoding flagellar proteins in high- or low-

virulence ST1 isolates.

Although low virulence ST1 strains cells have reduced motility and flagellar expression, we were able to

select strain variants that had upregulated flagellin expression by selectively culturing bacteria from the

periphery of swim colonies, which we refer to as E strains (i.e., ST1-6E and ST1-27E). TEM images

demonstrate that ST1-6E and ST1-27E strains are highly flagellated (Fig. 2A). Cecal contents of mice

infected with the E strains had higher Tcd toxins compared to the parental isolates (Fig. 2B).

The shift in flagellar expression and toxin production directed us to examine the flagellar switch, which is

in the upstream region of *flg*B operon (31) and has a length of approximately 200bp with a 154bp switch

region and two flanking inverted repeats. The switch can flip between the ON and OFF orientation but the

factors regulating its position remain unclear. We used the reads obtained by Illumina sequencing to

identify ON and OFF sequences in ST1 strains and discovered that both are present in all of the clinical

isolates. We quantified the fraction of ON and OFF sequences in *C. difficile* liquid culture more precisely

by flagellar switch qPCR (31) and found that high-virulence isolates have 70-80% flagellar switches in the

ON position, whereas less virulent isolates either have >90% in either the ON or OFF position. A higher

proportion of flagellar switches in the ON position weakly correlates with maximum weight loss by linear

regression (R^2 = 0.39) (Fig. 2C), suggesting that alternative factors regulate the virulence.

The virulence has a strong association with the IR type.

As we expanded the alignment to all 68 isolates in our collection, we found that 65 isolates have the

flagellar switch sequence identical to *C. difficile* R20291 while the remaining three isolates have the

flagellar switch identical to *C. difficile* CD630. To examine whether the local genetic structure might impact

the flagellar switch directions, we compared the left and right IR of the flagellar switch. While most isolates

have IRs identical to the reference strain *C. difficile* R20291, 40% of isolates has at least one less A or T in

either IRs. We identified four types of IR flanking regions in our strain collection and named them as

Common1 (C1-59.38% 6A/6T), Common2 (C2-32.81%, 5A/6T or 6A/5T), Rare1 (R1-6.25%, 6A/5T or 5A/6T),

and Rare2 (R2-1.56%, 5A/5T) (Fig. 3A). More specifically, C1 has the sequence as seen in *C. difficile* R20291

with 6A on the left IR (LIR) and 6T on the right IR (RIR) (Fig. 4A, sequence in the box). The number of A and

T on both IRs is the same when C1 switches between OFF and ON. On the other hand, C2 has LIR-5A and

RIR-6T when OFF and LIR-6A and RIR-5T when ON. R1 has LIR-6A and RIR-5T when OFF and LIR-5A and

RIR-6T when ON. R2 has LIR-5A and RIR-5T either when it is ON or OFF. Depending on the biased direction

of the flagellar switch, we can further classify C2 and R1 types as C2-ON and C2-OFF, R1-ON and R1-OFF.

R2 appears in only one strain, ST1-67, when it is in the OFF direction. Comparing to other IR types, we

found a strong correlation between IR type C1 and more severe weigh loss in infected mice (p = 3.2e-05)

(Fig. 3B). We speculate that the IR type C1 may contribute more to the virulence of *C. difficile* than the

other types.

The inverted repeat type affects the flexibility of the flagella switch.

To further investigate whether different IR types bias the flagellar switch direction, we expanded the

flagellar switch qPCR analysis to all our isolates, 64 of which yielded a successful amplification. 3-4

biological replicates were examined per isolate. To contrast the proportion of ON and OFF switch in each

culture, we determined the ΔCt value from qPCR results.

$\Delta Ct = |Ct_{ON} - Ct_{OFF}|$

in which Ct_{ON} is the cycle threshold from qPCR amplifying the ON sequence, and Ct_{OFF} is the cycle threshold

from qPCR amplifying the OFF sequence.

159 We found that isolates with IR type C1 have the lowest average ΔCt, with 2.09 (0.54 to 5.00) cycles,

160 indicating a flexible flagellar switch that can flip between ON and OFF positions. When calculating the

161 proportion, the C1 culture is biased towards ON fraction, with approximately 72.62% (54.61-99.18) (Fig.

162 4B). On the other hand, IR type C2 has an average of ΔCt 8.88 (7.08 to 15.30) cycles when it is at OFF

163 direction. The difference is significant larger when the C2 switch is ON, with 14.87 (14.46 -15.30) cycles.

164 Other rarer IR types like R1 and R2 have significantly larger ΔCt (Fig. 4A), with 15.36 (13.62-16.57) cycles

165 for R1-OFF and 16.28 for R1-ON. The rarest type, R2-OFF with only one isolate, has the largest ΔCt, with

166 20.99 cycles. Because of the large ΔCt, the proportion of ON and OFF in isolates with C2, R1, and R2 are

167 very skewed toward one direction. In average, the biased-OFF isolates have 95.58% (88.71-100.00) OFF

168 genomes, whereas the biased-ON isolates have 96.67% (92.60-100.00) ON genomes.

169 To test if the division of ON and OFF fraction of each isolate displays a similar pattern in an *in vivo* model,

170 we performed qPCR from DNA extracted from fecal pellets collected from mice 24 hours after infection

171 with *C. difficile* isolates. Isolates from each switch type, including *C. difficile* R20291 yielded that same

172 proportions of ON and OFF detected in *in vitro* cultures, with IR type C1 having the lowest ΔCt 1.10 (0.09-

173 3.00) for three distinct strains (Fig. 4A) with 76.69% ON (65.62-85.86%) and 23.31% OFF (Fig. 4B). The

174 other switch types maintained larger ΔCt which resulted in biased to ON or OFF status. These results

175 suggest that the C1 switch type enables more flexible switching of flagellar expression which may

176 contribute to *in vivo* virulence of *C. difficile*.

177 **The IR type C1 shows more variation in the switch direction in *in vivo* model.**

178 We infected mice with *C. difficile* R20291 and clinical isolates with different switch types and monitored

179 the ON and OFF fractions in fecal samples and cecal contents over the course of eight days (Fig. 5). Isolates

180 with IR type C2, R1, and R2 remain largely locked in terms of the proportion of ON and OFF over the course

181 of eight days. In contrast, we observed more variation in switch status in mice infected with IR type C1

strains. In some mice, C1 isolates are predominantly in the OFF status, with switching occurring

predominantly in the recovery phase after resolution of diarrhea and weight loss.

***C. difficile* ST1 strains in the US have variable IR sequences.**

To further confirm IR diversity, we screened 1359 ST1 isolates with whole-genome sequences downloaded

from NCBI Biosamples. These samples are from the CDC HAI-Seq *C. difficile* collection (PRJNA629351) (32),

Texas and Michigan medical centers (PRJNA595724, PRJNA561087). The remaining samples come from

more recent publications (33-36). All *C. difficile* samples were collected from 1988 to 2020. Most

(1318/1359) are from Continental Europe, UK, Ireland and the U.S., with a small number (20/1359) of

samples from East Asia and Australia. Most *C. difficile* ST1 samples have flagellar switch with IR type C1,

accounting for 96.47% (1311/1359) of samples. Type C2 appears with only 2.43% (33/1359) of samples,

of which 27/33 originate from the U.S between 2007 and 2020. Other IR types are present at reduced

frequency of only 0.66% of samples (Fig. 6A). Focusing on the more recent CDC HAI-Seq collection, which

were predominantly submitted from Minnesota between 2009 and 2020, IR type C1 remains dominant,

but C2 accounts for 30.16% (19/63) samples, similar to our collection (32.80%). Interestingly, the

proportion of C2 increased in recent years, and accounts for up to 52.94% (9/17) of samples collected in

2020 (Fig. 6B). Our analysis on the prevalence of different IR types, especially in the U.S., suggests that the

heterogeneity in ST1 virulence maybe related to diversity of IR types.

Proposed model of flagellar switch inversion

RecV, a tyrosine recombinase, has been reported as necessary for the inversion of several switches in *C.*

difficile (37). However, the preferred DNA sequence for RecV binding had not been identified, nor was it

known how RecV avoids recombining incorrect pairs of sites – e.g., the ends of two different invertible

regions. Here we propose the mode of action of recV on the flagellar switch sequence.

The general mechanism for tyrosine recombinase-mediated recombination has been well-studied (38, 39)

and can be used as a basis for understanding RecV action. We propose that the inverted repeats which

flank each invertible switch are themselves degenerate inverted repeats consisting of two copies of a

RecV-binding motif flanking a central 6-bp spacer region. Recombination would occur within a tetramer

of RecV that pairs two such sites, with cleavage and strand exchange occurring at the outer edges of the

spacers. Recombinase binding does not depend on spacer sequence. However, because the spacer

segments become heteroduplex in the product (1 strand from each partner site), recombination of two

binding sites can only proceed to completion if spacers have identical sequences.

First, we used AlphaFold2 to predict the structure of a recV tetramer (Fig. 7A). The predicted structure is

highly similar to those of the catalytic domains of other tyrosine recombinases such as Cre (which

recombines loxP DNA sites), including the C-terminal helix that mediates protein – protein contacts. (38,

39). However, RecV lacks the N-terminal DNA binding domain found in many tyrosine recombinases. We

speculate that RecV's catalytic domain may bind more tightly and specifically than that of many tyrosine

recombinases, alleviating the need for an additional domain.

Second, we compared the inverted repeats of all six switches catalyzed at least in part by RecV in *C. difficile*

R20291, taking into account the approximate DNA cleavage points previously identified (37). We were

able to identify a conserved ~13bp potential RecV binding motif that is found in inverted repeat flanking

a 6bp spacer, as expected for canonical tyrosine recombinase binding sites (Fig. 7B). Searching the entire

C.   difficile   R20291   genome   for   a   pattern   based   on   this,

`AWAGTWNCCNTTWNNNNNNNWAANGGNWACTWT`, (W= A or T; N = any base; one mismatch allowed per 10

bp) produced 17 hits, including both ends of *C. difficile* inversions (cdi) 2-5 and one end each of cdis 1 and

6 (the other ends contain similar sequences but too many mismatches) and 7 additional sites that may be

false positives. In the future, more sophisticated position-weighted searches would be likely to yield more

accurate results. In further support of this motif, previous work has reported that inversion in *C. difficile*

R20291 is locked when the conserved nucleotides in the RIR are deleted, suggesting that the activity of

recV was impaired (40).

Further examination of the ends of the 6 switches reveals how the system avoids aberrant recombination

between incorrectly paired repeats (Fig. 7C). For each individual switch, the spacer sequences within the

RecV dimer-binding sites at each end are identical. However, each switch has its own "spacer code."

Therefore, even if a RecV tetramer were to pair two ends of different switch regions, it would be difficult

to complete recombination between them.

The length of this variable spacer region is similar to the length of the spacer in other tyrosine recombinase

systems, and we propose that variations in its length explain our findings. We predict the model for recV

cleavage as in Figure 7D in which the tyrosine cleaves the inverted repeat at the 5' end of the last

conserved nucleotide. After the cleavage, two strands exchange the spacer region and form a Holliday

structure to initiate the inversion of the switch. In our collection, type C1 has the spacer sequence as LIR-

tacaaa(6)/RIR-tttgta(6). The other types have less bases on either left or right of the spacer region. For

example, C2 has sequence as LIR-tacaa(5)/RIR-tttgta(6), R1 as LIR-tacaaa(6)/RIR-ttgta(5), and R2 as LIR-

tacaa(5)/RIR-ttgta(5). Previous work on xerCD/ dif site system has reported a less efficient recombination

when there is a change in number of bps in the spacer region (41). With our model, C1 has the most

efficient spacer sequence, allowing it to switch back and forth easier. The flexibility helps it to maintain

an adequate proportion of both statuses of the switch. Other IR types have unequal or less bps in the

spacer sequences which may result in a stuck Holliday structure. Further research focusing on this spacer

should be done to validate the model.

DISCUSSION

C. difficile ST1 has attracted public attention as one of the hypervirulent and highly antibiotic-resistant

strains. However, its virulence relative to other *C. difficile* strains remains controversial. Our findings

suggest that the flagellar switch, specifically the number of A and T in the left and right IRs, account for

variability in *in vivo* virulence.

C. difficile virulence is mediated by the secretion of two glucosylating toxins TcdA and TcdB, and the binary

toxin CDT. In our collection, we did not find an association between toxin expression in fecal samples and

the degree of virulence. In fact, several studies reported that the toxin profile from stool samples is not

associated with severe CDI (23, 42). Toxin expression is partly regulated by the flagellar switch. We found

a weak correlation between the flagellar switch direction and virulence in the mouse model. The OFF

isolates secrete significantly more toxin when the flagellar switch inverts to ON. However, this increment

does not always increase the virulence at the strain level (data not shown). This result suggests that factors

in addition to toxin secretion contribute to virulence of *C. difficile* ST1 isolates.

The published flagellar switch for the R20291 ST1 strain contains a 154-bp switching region and a 21-bp

IR (31). The IR has a high content of A and T. Interestingly, we found four types of IRs with different

numbers of A and T among our clinical isolates. Community strains with different numbers of bp in the IR

were reported previously in ribotype 12 (43). In our collection, the most prevalent IR, C1, is present in

over half of the isolates. We found that the IR type C1 correlates with more severe weight loss and the

other IR types cause milder CDI symptoms. The difference in genotype between C1 and other IR types

may explain the heterogeneity in virulence of *C. difficile* ST1 (44, 45).

The strong association between C1 and high virulence suggest that this IR have some unique features. C1

has high percentage of ON status in *in vitro* and *in vivo* conditions. However, the higher proportion of ON

is unlikely to account for increased virulence as the highly biased ON isolates have reduced *in vivo*

virulence. In the mouse model, the IR type C1 has increased flexibility in switching between ON and OFF

which may contribute to virulence. In some mice, C1 isolates switch toward more OFF cells during the

recovery phase of CDI. On the other hand, the extremely biased ON isolates maintain the same percentage

over time. These results suggest that the capacity to switch between ON and OFF and maintain adequate

proportions of ON and OFF cells and may increase *C. difficile* virulence. Flagella production is a trade-off.

Other bacteria like *Pseudomonas aeruginosa* and *Vibrio cholera* use different strategies to switch between

sessile and motile phenotypes to reduce the metabolic cost of flagellar expression (46-48). In fact, in more

than 1359 ST1 isolates worldwide, more than 95% have type C1, a flexible type of switch. In *C. difficile*

ST1, ability to turn on and off the flagellar switch may reduce fitness costs and thereby impact virulence.

It is interesting that C2-OFF is the second most common type whereas it has a strong association with

reduced virulence. More specifically, the percentage of C2-OFF is higher in the U.S comparing to other

areas, potentially resulting from higher rates of testing and *C. difficile* cultivation in the U.S. In our

collection and in the CDC HAI-Seq *C. difficile* collection, C2-OFF accounts for approximately 30% of all

isolates. C2-OFF isolates are mostly non-flagellated. In the mouse model, C2-OFF remains mostly OFF over

the course of eight observation days. Flagellar synthesis and motility is costly (49). In addition, flagella are

targets for mucosal innate immune response. Saving the metabolic cost and evading the immune system

might be a strategy to increase persistence of *C. difficile*.

Our study correlates the flagellar switch and IR with the virulence of ST1 *C. difficile* clinical isolates. We

report here different types of IR in *C. difficile* ST1 and their association with disease severity in the mouse

model of CDI and propose a model to explain different ratios of ON and OFF status between IR types.

Further research on the flagellar switch regions of clinical ST1 and non-ST1 strains may provide insights

into the wide range of disease severities in patients infected with *C. difficile*.

METHODS

Sample collection

Clinical *C. difficile* isolates were collected from 2013 to 2017 at Memorial Sloan Kettering Cancer Center

(MSKCC) from patients receiving bone marrow transplants and cancer chemotherapy (50). *C. difficile*

isolates were classified using multilocus sequence typing (MLST) method (51). The isolates were

sequenced whole genome in an Illumina Hiseq platform (52) and circularized by Oxford Nanopore

Technologies.

Bacterial growth and spore collection

Frozen stock of *C. difficile* was streak on Brain heart infusion (BHI) agar plus 0.1% (w/v) sodium

taurocholate hydrate (Cat. 86339, Sigma-Aldrich). A single colony was picked and subcultured in BHI

medium (BD, 237500) with yeast extract and 0.1% L-Cysteine (BHIS) at 37°C in anaerobic chamber (Coy

lab).

To collect spore for mouse infection, *C. difficile* was either incubated in BHIS broth for approximately two

months or on 70:30 agar for 4-5 days to encourage sporulation (53). Next, spore is separated from cell

debris by gradient centrifuge with 20% (wt/vol) HistoDenz (D2158, Sigma-Aldrich) or 50%

(wt/vol) sucrose, then wash five times in sterile water. Spore solution was further incubated at 65°C for

20 minutes to omit vegetative cells. Spore purity is confirmed by plating on BHIS plate, checking for the

absence of vegetative cell growth.

Mouse experiment

C57BL/6 female mice from six- to eight-week-old were purchased from Jackson laboratory, room MP14.

Mice were randomized and administered *ad libitum* with a cocktail of metronidazole (0.25g/l), neomycin

(0.25g/l), and vancomycin (0.25g/l) for three days then rest for two days before being intraperitoneally

injected with 200μg clindamycin. After 24 hours, mice were oral gavage with approximately 200 spores of

C. difficile. All mice infected with *C. difficile* were single housed then monitored for weight loss and clinical

sign of disease for seven days.

CFU/ml counting, and toxin titer measurement was done from fecal pellet collected one day after

infection. Toxin assay was performed as described elsewhere (24).

RNA sequencing

Mice were infected with spore from four selected isolates (ST1-6, ST1-12, ST1-27, and ST1-53). After 24

hours, mice were sacrificed to collect cecal content for RNA extraction (Qiagen RNeasy PowerMicrobiome

kit). Total RNA was sent to Genewiz for rRNA removal and library preparation before sequencing. Adapters

were trimmed off from the raw reads, and their quality was assessed and controlled using Trimmomatic

(v.0.39) (54), then human genome was identified and removed by kneaddata (v0.7.10,

https://github.com/biobakery/kneaddata), while ribosomal RNA was removed by aligned the clean non-

host reads to silva database (138.1 SSURef). The remaining reads from each sample were mapped to their

corresponding circularized genome using bowtie2(55), and reads counts of each gene were obtained by

running featureCounts from Subread (v2.0.1) (56) ,and the core gene counts were normalized by DESeq2

(57).

Swim plates

C. difficile was grown in BHIS broth to late log phase then diluted to the same OD_{600nm} ~0.5. A 10μl tip is

immerged into diluted *C. difficile* culture and then stabbed into 0.3% agarose buffered with BHIS, pH 7.0.

The swim plates were then incubated at 37°C. After 24 hours, the colony size was measured and compared

between isolates.

Transmission electron microscope (TEM)

C. difficile was streak on BHIS agar plate and incubate overnight. To collect cells for TEM, 10μl of distilled

water was drop onto a colony. After two minutes, a copper grid (CF400-Cu, EMS) was put on top of the

soaked colony so that vegetative cells were passively transferred to the grid. To stain the cells, one drop

of uranyl acetate 1% was added to the grid and incubate for 30 seconds. TEM pictures of the stained cells

were then taken at 2900X1.4 or 5900X1.4 using FEI Tecnai F30 microscope at University of Chicago

Advanced Electron Microscopy Facility.

Quantitative PCR of the flagellar switch

C. difficile was grown in BHIS broth to late log phase (OD$_{600nm}$ ~ 0.5) with three to four replicates per

isolate. *C. difficile* DNA was extracted by Qiamp PowerFecal Pro DNA kit (Cat. 51804). Quantitative real

time PCR was done using 20ng template and flagellar switch primers (31). *C. difficile* R20291 mutated

strains with flagellar switch locked ON and OFF were used as the control (40). Adenosine kinase (*adk)* gene

is used as a reference gene. qPCR was set up using PowerUp™ SYBR™ Green Master Mix (Cat. A25742)

and run by QuantStudio Real-Time PCR Systems. Percentage of the switch direction was calculated using

the *ΔCt* method.

Flagella switch alignment and IR type survey in public databases

C.difficile isolates (N=131) from BioProject PRJNA595724 and PRJEB2318 (N=1198) were downloaded

from NCBI and assembled into contigs using SPAdes (58). Eleven of those isolates were lost during the

downloading and assembling process. In addition, a collection of 66 *C.difficile* genomes from Patric (date:

Mar. 15 2022) { https://www.bv-brc.org/} were also downloaded. MLST was determined on those contigs

by mlst {Seemann T, mlst Github https://github.com/tseemann/mlst}. Flagellar switch region and 50 bp

upstream and downstream of 68 isolates in our collection of both the ON and OFF sequences were used

as query to BLAST (59) against the assembled contigs, and hits with at least 85% identity and 85% coverage

of the query are considered a valid match.

RNA extraction, reverse transcription and RT-qPCR

Cecal RNA was extracted using Rneasy PowerMicrobiome Kit (Qiagen) according to the manufacturer's

instructions. Complementary DNA was generated using the QuantiTect reverse transcriptase kit (Qiagen)

according to the manufacturer's instructions. Quantitative PCR was performed on complementary DNA

using primers with PowerTrack SYBR Green Master Mix (Thermo Fisher). Reactions were run on a

QuantStudio 6 pro (Thermo Fisher). Relative abundance was normalized by $\Delta\Delta$Ct. TcdA_qFor 5'-

GTATGGATAGGTGGAGAAGTCA-3'; TcdA_qRev 5'-CTCTTCCTCTAGTAGCTGTAATGC-3' {Babakhani 2012 J

Antimicrobial Chemotherapy}; TcdB_qFor 5'-AGCAGTTGAATATAGTGGTTTAGTTAGAGTTG-3'; TcdB_qRev

5'-CATGCTTTTTTAGTTTCTGGATTGAA-3' {Wroblewski 2009 J Clin Microbiol.}; FliS1_qFor 5'-

TGCAGGACAATGGGCAAAGG-3'; FliS1_qRev 5'- CAGGCAACACATTATCTATTACCTGG-3'; FliE_qFor 5'-

AGGCGAAGATGTTTCTATGCA-3'; FliE_qRev 5'- ACCTTATTCATTTCTTGATATGCATCA-3'

Structure prediction and conserved sequence analysis

The RecV tetramer structure was predicted by Alphafold-multimer using a web-based Colab

implementation, and default parameters (60-62). Sequences of the inverted repeats flanking the switch

regions were located in the *Clostridium difficile* R20291 genome (accession number CP029423) using

Snapgene (https://www.snapgene.com/). After manual alignment, the sequence logo shown in Fig. 7B

was created using https://weblogo.berkeley.edu/logo.cgi (63)   An approximation of the proposed

consensus   sequence   for   binding   of   a   RecV   dimer   was   based   on   that   logo

(`AWAGTWNCCNTTWNNNNNNWAANGGNWACTWT`) and was tested by searching the full genome in

Snapgene for related sequences with 1 mismatch allowed per 10bp.

Statistical analysis

Kruskal-Wallis or Two Sample T tests were performed to test the difference in maximum weight loss, swim

plate and ΔCt results. Statistical significance was determined by using a *P* value of <0.05. Linear regression

analysis was used to estimate the correlation between IR type and weight loss.

ACKNOWLEDGEMENT

We would like to thank Dr. Rita Tamayo for providing the primers and reference strains for the flagellar switch qPCR. This work was supported by the National Institutes of Health R01 AI095706 and by the Duchossois Family Institute of the University of Chicago.. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

AUTHOR CONTRIBUTION

N.T.Q.N., Q.D. and E.G.P. conceived the project. Q.D., N.T.Q.N., H.L. and Q.D. analyzed the data. N.T.Q.N., Q.D., and J.K.S., performed experiments. M.K. isolated *C. difficile* isolates. V.B.Y. and E.S.S. sequenced clinical isolates. N.T.Q.N., H.L., Q.D., P.A.R. and E.G.P. interpreted the results and wrote the manuscript.

REFERENCES

1.   CDC. 2019. Antibiotic Resistance Threats in the United States Atlanta GUSDoHaHS, CDC; 2019.,
2.   CDC. 2020. 2020 Annual Report.
3.   Lyerly DM, Krivan HC, Wilkins TD. 1988. Clostridium difficile: its disease and toxins. Clin Microbiol Rev 1:1-18.
4.   Ciesielski-Treska J, Ulrich G, Rihn B, Aunis D. 1989. Mechanism of action of Clostridium difficile toxin B: role of external medium and cytoskeletal organization in intoxicated cells. Eur J Cell Biol 48:191-202.
5.   Florin I, Thelestam M. 1983. Internalization of Clostridium difficile cytotoxin into cultured human lung fibroblasts. Biochim Biophys Acta 763:383-92.
6.   Henriques B, Florin I, Thelestam M. 1987. Cellular internalisation of Clostridium difficile toxin A. Microb Pathog 2:455-63.
7.   Just I, Selzer J, Hofmann F, Green GA, Aktories K. 1996. Inactivation of Ras by Clostridium sordellii lethal toxin-catalyzed glucosylation. J Biol Chem 271:10149-53.
8.   Just I, Selzer J, Wilm M, von Eichel-Streiber C, Mann M, Aktories K. 1995. Glucosylation of Rho proteins by Clostridium difficile toxin B. Nature 375:500-3.
9.   Just I, Wilm M, Selzer J, Rex G, von Eichel-Streiber C, Mann M, Aktories K. 1995. The enterotoxin from Clostridium difficile (ToxA) monoglucosylates the Rho proteins. J Biol Chem 270:13932-6.
10.   Just I, Fritz G, Aktories K, Giry M, Popoff MR, Boquet P, Hegenbarth S, von Eichel-Streiber C. 1994. Clostridium difficile toxin B acts on the GTP-binding protein Rho. J Biol Chem 269:10706-12.
11.   Schnizlein MK, Young VB. 2022. Capturing the environment of the Clostridioides difficile infection cycle. Nat Rev Gastroenterol Hepatol 19:508-520.

12. Vedantam G, Clark A, Chu M, McQuade R, Mallozzi M, Viswanathan VK. 2012. Clostridium difficile
infection: toxins and non-toxin virulence factors, and their contributions to disease establishment
and host response. Gut Microbes 3:121-34.
13. Lesniak NA, Schubert AM, Flynn KJ, Leslie JL, Sinani H, Bergin IL, Young VB, Schloss PD. 2022. The
Gut Bacterial Community Potentiates Clostridioides difficile Infection Severity. mBio
13:e0118322.
14. Loo VG, Bourgault AM, Poirier L, Lamothe F, Michaud S, Turgeon N, Toye B, Beaudoin A, Frost EH,
Gilca R, Brassard P, Dendukuri N, Beliveau C, Oughton M, Brukner I, Dascal A. 2011. Host and
pathogen factors for Clostridium difficile infection and colonization. N Engl J Med 365:1693-703.
15. Tenover FC, Tickler IA, Persing DH. 2012. Antimicrobial-resistant strains of Clostridium difficile
from North America. Antimicrob Agents Chemother 56:2929-32.
16. Wieczorkiewicz JT, Lopansri BK, Cheknis A, Osmolski JR, Hecht DW, Gerding DN, Johnson S. 2016.
Fluoroquinolone and Macrolide Exposure Predict Clostridium difficile Infection with the Highly
Fluoroquinolone- and Macrolide-Resistant Epidemic C. difficile Strain BI/NAP1/027. Antimicrob
Agents Chemother 60:418-23.
17. Hubert B, Loo VG, Bourgault AM, Poirier L, Dascal A, Fortin E, Dionne M, Lorange M. 2007. A
portrait of the geographic dissemination of the Clostridium difficile North American pulsed-field
type 1 strain and the epidemiology of C. difficile-associated disease in Quebec. Clin Infect Dis
44:238-44.
18. Labbe AC, Poirier L, Maccannell D, Louie T, Savoie M, Beliveau C, Laverdiere M, Pepin J. 2008.
Clostridium difficile infections in a Canadian tertiary care hospital before and during a regional
epidemic associated with the BI/NAP1/027 strain. Antimicrob Agents Chemother 52:3180-7.
19. Morgan OW, Rodrigues B, Elston T, Verlander NQ, Brown DF, Brazier J, Reacher M. 2008. Clinical
severity of Clostridium difficile PCR ribotype 027: a case-case study. PLoS One 3:e1812.
20. Sirard S, Valiquette L, Fortier LC. 2011. Lack of association between clinical outcome of Clostridium
difficile infections, strain type, and virulence-associated phenotypes. J Clin Microbiol 49:4040-6.
21. Valiente E, Cairns MD, Wren BW. 2014. The Clostridium difficile PCR ribotype 027 lineage: a
pathogen on the move. Clin Microbiol Infect 20:396-404.
22. Warny M, Pepin J, Fang A, Killgore G, Thompson A, Brazier J, Frost E, McDonald LC. 2005. Toxin
production by an emerging strain of Clostridium difficile associated with outbreaks of severe
disease in North America and Europe. Lancet 366:1079-84.
23. Carlson PE, Jr., Walk ST, Bourgis AE, Liu MW, Kopliku F, Lo E, Young VB, Aronoff DM, Hanna PC.
2013. The relationship between phenotype, ribotype, and clinical disease in human Clostridium
difficile isolates. Anaerobe 24:109-16.
24. Qiwen Dong HL, Marie-Maude Allen, Julian R Garneau, Jonathan K Sia, Rita C Smith, Fidel Haro,
Tracy McMillen, Rosemary L Pope, Carolyn Metcalfe, Victoria Burgo, Che Woodson, Nicholas
Dylla, Claire Kohout, Anitha Sundararajan, Evan S Snitkin, Vincent B Young, Louis-Charles Fortier,
Mini Kamboj, Eric G Pamer. 2023. Virulence and genomic diversity among clinical isolates of ST1
(BI/NAP1/027) Clostridioides difficile. bioRxiv doi:doi: 10.1101/2023.01.12.523823.
25. La Ragione RM, Sayers AR, Woodward MJ. 2000. The role of fimbriae and flagella in the
colonization, invasion and persistence of Escherichia coli O78:K80 in the day-old-chick model.
Epidemiol Infect 124:351-63.
26. Eaves-Pyles T, Murthy K, Liaudet L, Virag L, Ross G, Soriano FG, Szabo C, Salzman AL. 2001.
Flagellin, a novel mediator of Salmonella-induced epithelial activation and systemic inflammation:
I kappa B alpha degradation, induction of nitric oxide synthase, induction of proinflammatory
mediators, and cardiovascular dysfunction. J Immunol 166:1248-60.
27. Horstmann JA, Lunelli M, Cazzola H, Heidemann J, Kuhne C, Steffen P, Szefs S, Rossi C, Lokareddy
RK, Wang C, Lemaire L, Hughes KT, Uetrecht C, Schluter H, Grassl GA, Stradal TEB, Rossez Y, Kolbe

M, Erhardt M. 2020. Methylation of Salmonella Typhimurium flagella promotes bacterial adhesion
and host cell invasion. Nat Commun 11:2013.

28.   Baban ST, Kuehne SA, Barketi-Klai A, Cartman ST, Kelly ML, Hardie KR, Kansau I, Collignon A,
Minton NP. 2013. The role of flagella in Clostridium difficile pathogenesis: comparison between a
non-epidemic and an epidemic strain. PLoS One 8:e73026.

29.   Batah J, Kobeissy H, Bui Pham PT, Deneve-Larrazet C, Kuehne S, Collignon A, Janoir-Jouveshomme
C, Marvaud JC, Kansau I. 2017. Clostridium difficile flagella induce a pro-inflammatory response
in intestinal epithelium of mice in cooperation with toxins. Sci Rep 7:3256.

30.   Dingle TC, Mulvey GL, Armstrong GD. 2011. Mutagenic analysis of the Clostridium difficile flagellar
proteins, FliC and FliD, and their contribution to virulence in hamsters. Infect Immun 79:4061-7.

31.   Anjuwon-Foster BR, Tamayo R. 2017. A genetic switch controls the production of flagella and
toxins in Clostridium difficile. PLoS Genet 13:e1006701.

32.   Paulick A, Adamczyk M, Anderson K, Vlachos N, Machado MJ, McAllister G, Korhonen L, Guh AY,
Halpin AL, Rasheed JK, Karlsson M, Lutgring JD, Gargis AS, Emerging Infections Program
Clostridioides difficile Pathogen G. 2021. Characterization of Clostridioides difficile Isolates
Available through the CDC & FDA Antibiotic Resistance Isolate Bank. Microbiol Resour Announc
10.

33.   Dingle KE, Didelot X, Quan TP, Eyre DW, Stoesser N, Golubchik T, Harding RM, Wilson DJ, Griffiths
D, Vaughan A, Finney JM, Wyllie DH, Oakley SJ, Fawley WN, Freeman J, Morris K, Martin J, Howard
P, Gorbach S, Goldstein EJC, Citron DM, Hopkins S, Hope R, Johnson AP, Wilcox MH, Peto TEA,
Walker AS, Crook DW, Modernising Medical Microbiology Informatics G. 2017. Effects of control
interventions on Clostridium difficile infection in England: an observational study. Lancet Infect
Dis 17:411-421.

34.   Eyre DW, Davies KA, Davis G, Fawley WN, Dingle KE, De Maio N, Karas A, Crook DW, Peto TEA,
Walker AS, Wilcox MH, Group ES. 2018. Two Distinct Patterns of Clostridium difficile Diversity
Across Europe Indicating Contrasting Routes of Spread. Clin Infect Dis 67:1035-1044.

35.   Eyre DW, Didelot X, Buckley AM, Freeman J, Moura IB, Crook DW, Peto TEA, Walker AS, Wilcox
MH, Dingle KE. 2019. Clostridium difficile trehalose metabolism variants are common and not
associated with adverse patient outcomes when variably present in the same lineage.
EBioMedicine 43:347-355.

36.   Stoesser N, Crook DW, Fung R, Griffiths D, Harding RM, Kachrimanidou M, Keshav S, Peto TE,
Vaughan A, Walker AS, Dingle KE. 2011. Molecular epidemiology of Clostridium difficile strains in
children compared with that of strains circulating in adults with Clostridium difficile-associated
infection. J Clin Microbiol 49:3994-6.

37.   Sekulovic O, Mathias Garrett E, Bourgeois J, Tamayo R, Shen A, Camilli A. 2018. Genome-wide
detection of conservative site-specific recombination in bacteria. PLoS Genet 14:e1007332.

38.   Grindley ND, Whiteson KL, Rice PA. 2006. Mechanisms of site-specific recombination. Annu Rev
Biochem 75:567-605.

39.   Meinke G, Bohm A, Hauber J, Pisabarro MT, Buchholz F. 2016. Cre Recombinase and Other
Tyrosine Recombinases. Chem Rev 116:12785-12820.

40.   Trzilova D, Warren MAH, Gadda NC, Williams CL, Tamayo R. 2022. Flagellum and toxin phase
variation impacts intestinal colonization and disease development in a mouse model of
Clostridioides difficile infection. Gut Microbes 14:2038854.

41.   Blakely G, Sherratt D. 1996. Determinants of selectivity in Xer site-specific recombination. Genes
Dev 10:762-73.

42.   Rao K, Micic D, Natarajan M, Winters S, Kiel MJ, Walk ST, Santhosh K, Mogle JA, Galecki AT, LeBar
W, Higgins PD, Young VB, Aronoff DM. 2015. Clostridium difficile ribotype 027: relationship to age,

detectability of toxins A or B in stool with rapid testing, severe infection, and mortality. Clin Infect
Dis 61:233-41.
43. Anjuwon-Foster BR, Maldonado-Vazquez N, Tamayo R. 2018. Characterization of Flagellum and
Toxin Phase Variation in Clostridioides difficile Ribotype 012 Isolates. J Bacteriol 200.
44. Gu T, Li W, Yang LL, Yang SM, He Q, He HY, Sun DL. 2022. Systematic review of guidelines for the
diagnosis and treatment of Clostridioides difficile infection. Front Cell Infect Microbiol 12:926482.
45. Johnson S, Lavergne V, Skinner AM, Gonzales-Luna AJ, Garey KW, Kelly CP, Wilcox MH. 2021.
Clinical Practice Guideline by the Infectious Diseases Society of America (IDSA) and Society for
Healthcare Epidemiology of America (SHEA): 2021 Focused Update Guidelines on Management of
Clostridioides difficile Infection in Adults. Clin Infect Dis 73:755-757.
46. Fernandez NL, Hsueh BY, Nhu NTQ, Franklin JL, Dufour YS, Waters CM. 2020. Vibrio cholerae
adapts to sessile and motile lifestyles by cyclic di-GMP regulation of cell shape. Proc Natl Acad Sci
U S A 117:29046-29054.
47. Lichtenberg M, Kragh KN, Fritz B, Kirkegaard JB, Tolker-Nielsen T, Bjarnsholt T. 2022. Cyclic-di-
GMP signaling controls metabolic activity in Pseudomonas aeruginosa. Cell Rep 41:111515.
48. Nadell CD, Bassler BL. 2011. A fitness trade-off between local competition and dispersal in Vibrio
cholerae biofilms. Proc Natl Acad Sci U S A 108:14181-5.
49. Schavemaker PE, Lynch M. 2022. Flagellar energy costs across the tree of life. Elife 11.
50. Kamboj M, McMillen T, Syed M, Chow HY, Jani K, Aslam A, Brite J, Fanelli B, Hasan NA, Dadlani M,
Westblade L, Zehir A, Simon M, Babady NE. 2021. Evaluation of a Combined Multilocus Sequence
Typing and Whole-Genome Sequencing Two-Step Algorithm for Routine Typing of Clostridioides
difficile. J Clin Microbiol 59.
51. Griffiths D, Fawley W, Kachrimanidou M, Bowden R, Crook DW, Fung R, Golubchik T, Harding RM,
Jeffery KJ, Jolley KA, Kirton R, Peto TE, Rees G, Stoesser N, Vaughan A, Walker AS, Young BC, Wilcox
M, Dingle KE. 2010. Multilocus sequence typing of Clostridium difficile. J Clin Microbiol 48:770-8.
52. Miles-Jay A, Young VB, Pamer EG, Savidge TC, Kamboj M, Garey KW, Snitkin ES. 2021. A multisite
genomic epidemiology study of Clostridioides difficile infections in the USA supports differential
roles of healthcare versus community spread for two common strains. Microb Genom 7.
53. Weldy M, Evert C, Dosa PI, Khoruts A, Sadowsky MJ. 2020. Convenient Protocol for Production
and Purification of Clostridioides difficile Spores for Germination Studies. STAR Protoc 1:100071.
54. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics 30:2114-20.
55. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods 9:357-
9.
56. Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features. Bioinformatics 30:923-30.
57. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. Genome Biol 15:550.
58. Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. 2020. Using SPAdes De Novo
Assembler. Curr Protoc Bioinformatics 70:e102.
59. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+:
architecture and applications. BMC Bioinformatics 10:421.
60. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R,
Zidek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes B,
Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger M,
Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW, Kavukcuoglu K, Kohli
P, Hassabis D. 2021. Highly accurate protein structure prediction with AlphaFold. Nature 596:583-
589.

61.   Mirdita M, Schutze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. 2022. ColabFold: making
562   protein folding accessible to all. Nat Methods 19:679-682.
563   62.   Richard Evans PON, Alexander Pritzel, Natasha Antropova, Andrew Senior, Tim Green, Augustin
564   Žídek, Russ Bates, Sam Blackwell, Jason Yim, Olaf Ronneberger, Sebastian Bodenstein, Michal
565   Zielinski, Alex Bridgland, Anna Potapenko, Andrew Cowie, Kathryn Tunyasuvunakool, Rishub Jain,
566   Ellen Clancy, Pushmeet Kohli, John Jumper, Demis Hassabis. 2022. Protein complex prediction
567   with AlphaFold-Multimer. bioRxiv doi:https://doi.org/10.1101/2021.10.04.463034.
568   63.   Crooks GE, Hon G, Chandonia JM, Brenner SE. 2004. WebLogo: a sequence logo generator.
569   Genome Res 14:1188-90.

FIGURE LEGENDS

Figure 1. Highly virulent isolates have more flagella. A) Heat map of RNA sequencing results from four ST1 isolates. The heat map is generated based on the number of transcripts per sample after normalization. RNA was extracted from cecal content of mice after one day infected with *C. difficile* ST1 isolates (n= 5 mice per isolate). B) TEM pictures of four selected isolates. C) Swim plates results of four isolates after 24 hours incubation (n = 3 replicates per isolate). Highly virulent isolates: pink. Less virulent isolate: black

Figure 2. The role of flagellar switch in *C. difficile* ST1 virulence. A) The swimming cells (E strains) collected from less virulent isolates have flagella and B) higher toxin production in fecal samples. C) Summary of the proportion of ON and OFF of each isolates and its correlation with the maximum weight loss. Data is generated from the isolates shown in Figure 1.

Figure 3. The flagellar-switch IR type has a strong association with the virulence. A) Flagellar switch alignment of some representative isolates with the reference sequence of *C. difficile* R20291. B) Comparison between IR type C1 and other non-C1 IR type regarding the maximum weight loss. Data is generated based on the collection of isolates shown in Figure 1.

Figure 4. Flagellar-switch IR type C1 is more flexible than other IR types. A) The plot summarizes the ΔCt of all IR types. ΔCt is calculated from flagellar-switch qPCR results. B) The percentage of ON genomes per IR type. For *in vitro* condition, the qPCR was performed using genomic DNA extracted from BHIS culture (each dot represents for one isolate, n = 3 to 4 replicates per isolate). For *in vivo* condition, genomic DNA was extracted from day-one-post-infection fecal pellet (each dot represents for one mouse, n = 4 to 10 mice per isolate). *In vivo* results for IR type C1 are generated from R20291, ST1-12, and ST1-53 (18 mice), C2-ON from ST1-19 (4 mice), C2-OFF from ST1-6, and ST1-27 (8 mice), R1-ON from ST1-51 (4 mice), R1-OFF from ST1-13, ST1-68 (8 mice), and R2 from ST1-67 (4 mice).

Figure 5. Flagellar-switch IR type C1 has larger variation in terms of the percentage of ON genomes. Data was generated from qPCR on DNA extracted from fecal pellets collected over the course of seven days. Day-eighth data was from DNA extracted from cecal content. Each dot represents for one mouse.

Figure 6. The diversity of IR types is mostly seen in the U.S. A) The contribution of each IR type across geographical areas. B) The contribution of each IR type in the U.S across years as comparing with our collection (MSKCC) and C) the recent CDC HAI *C. difficile* collection.

Figure 7. The proposed model for recV binding to the flagellar switch. A) The tetramer structure of recV
when binding to the IRs of the flagellar switch. B) The conserved sequence in the IR of six switches
mediated by recV in *C. difficile* R20291. C) The proposed spacer sequence in the flagellar switch IRs
(underlined by arrows). We are showing here the sequence of the upper strand. D) The binding of
tetramer recV to the flagellar switch IRs initiates the inversion of the switch. The model is drawn and
licensed by BioRender.

SUPPLEMENTAL FIGURE LEGENDS

SUPPLEMENTAL FIGURE LEGENDS

Figure S1. *C. difficile* ST1 isolates have variable virulence in antibiotic-treated mice. A) Boxplot showing
the CFU/g data from fecal pellet collected one day after infection (n = 2 to 13 mice per isolate). B) The
maximum weight loss percentage (n = 5 to 13 mice per isolate). In antibiotic-treated mice infected with
C. difficile ST1, the most severe weight loss happens at day two or day three after infection C) Toxin titer
from fecal pellets collected one day after infection (n = 2 to 13 mice per isolate, except for ST1-51). Each
black dot represents for a data point from one mouse. The red asterisk represents for an average value.
The dash line in panel B shows the 10% cutoff of weight loss.

Figure S2. Comparable toxin production and CFU between high and low virulence isolates. A) The toxin
titer and B) CFU/g generated from fecal content collected from mice one day post infection (n = 5 mice
per isolate).

Figure S3. RT_qPCR confirms the difference in flagellar gene expression between highly and less virulent
isolates. Fold change of flagellar genes fliE A), fliS1 B), and toxin genes tcdA C), and tcdB D) to adk to
ST1-6 (n=3 to 4 RT-qPCR for each isolate).

Figure S1. *C. difficile* ST1 isolates have variable virulence in antibiotic-treated mice.

Figure 1. Highly virulent isolates have more flagella.

Figure S2. Comparable toxin production and CFU between high and low virulence isolates.

Figure S3. RT_qPCR confirms the difference in flagellar gene expression between highly and less virulent isolates.

Figure 2. The role of flagellar switch in *C. difficile* ST1 virulence.

A)

Isolate	Virulence	Type	Left inverted repeat	Central region	Right inverted repeat
R20291	High	C1	AAGTTACTATTTTAC AAAAAA	AGGCAA……….AAGTCA	TTTTTTGTAATATAGCAACTT
ST1-12	High	C1	AAGTTACTATTTTAC AAAAAA	AGGCAA……….AAGTCA	TTTTTTGTAATATAGCAACTT
ST1-53	High	C1	AAGTTACTATTTTAC AAAAAA	AGGCAA……….AAGTCA	TTTTTTGTAATATAGCAACTT
ST1-6	Low	C2-OFF	AAGTTACTATTTTAC AAAAA	TGACTT……….. TTGCCT	TTTTTTGTAATATAGCAACTT
ST1-27	Low	C2-OFF	AAGTTACTATTTTAC AAAAA	TGACTT……….. TTGCCT	TTTTTTGTAATATAGCAACTT
ST1-19	Intermediate	C2-ON	AAGTTACTATTTTAC AAAAAA	AGGCAA……….AAGTCA	TTTTT GTAATATAGCAACTT
ST1_65	Low	C2-ON	AAGTTACTATTTTAC AAAAAA	AGGCAA……….AAGTCA	TTTTT GTAATATAGCAACTT
ST1-68	Low	R1-OFF	AAGTTACTATTTTAC AAAAAA	TGACTT……….. TTGCCT	TTTTT GTAATATAGCAACTT
ST1-51	Low	R1-ON	AAGTTACTATTTTAC AAAAA	AGGCAA……….AAGTCA	TTTTTTGTAATATAGCAACTT
ST1-67	Low	R2-OFF	AAGTTACTATTTTAC AAAAA	TGACTT……….. TTGCCT	TTTTT GTAATATAGCAACTT

B)

Figure 3. The flagellar-switch IR type has a strong association with the virulence.

Figure 4. Flagellar-switch IR type C1 is more flexible than other IR types.

Figure 5. Flagellar-switch IR type C1 has larger variation in terms of the percentage of ON genomes.

Figure 6. The diversity of IR types is mostly seen in the U.S.

Figure 7. The proposed model for recV binding to the flagellar switch.