IQCH regulates spermatogenesis by interacting with CaM to promote the expression of RNA-binding proteins

Tiechao Ruan[1,2]†, Ruixi Zhou[1,2]†, Yihong Yang[3]†, Junchen Guo[4,5], Chuan Jiang[1],

Xiang Wang[1], Gan Shen[1], Siyu Dai[1], Suren Chen[6*], Ying Shen[1,7*]

[1] Key Laboratory of Obstetrics, Gynecologic and Pediatric Diseases and Birth Defects of the Ministry of Education, Sichuan University, Chengdu, 610041, China;

[2] Department of Pediatrics, West China Second University Hospital, Sichuan University, Chengdu 610041 China;

[3] Reproduction Medical Center of West China Second University Hospital, Key Laboratory of Obstetric, Gynecologic and Pediatric Diseases and Birth Defects of Ministry of Education, Sichuan University, Chengdu 610041, China;

[4] Sichuan University-The Chinese University of Hong Kong (SCU-CUHK) Joint Laboratory for Reproductive Medicine, Key Laboratory of Obstetric, Gynaecologic and Paediatric Diseases and Birth Defects of Ministry of Education, West China Second University Hospital, Sichuan University, Chengdu 610041, China.

[5] Reproductive Endocrinology and Regulation Laboratory, Department of Obstetric and Gynaecologic, West China Second University Hospital, Sichuan University, Chengdu 610041, China.

[6] Education Key Laboratory of Cell Proliferation & Regulation Biology, College of Life Sciences, Beijing Normal University, Beijing 100875, China

[7] NHC Key Laboratory of Chronobiology, Sichuan University, Chengdu 610041, China;

†These authors contributed equally to this work

*Corresponding author:

Suren Chen

E-mail address: chensr@bnu.edu.cn;

Ying Shen

Phone: +86-15982083663.

E-mail address: yingcaishen01@163.com

Abstract

IQ motif-containing proteins can be recognized by calmodulin (CaM) and are

essential for many biological processes. However, the role of IQ motif-containing

proteins in spermatogenesis is largely unknown. In this study, we identified a

loss-of-function mutation in the novel gene IQ motif-containing H (*IQCH*) in a

Chinese family with male infertility characterized by a cracked flagellar axoneme and

abnormal mitochondrial structure. To verify the function of IQCH, *Iqch*-knockout

mice were generated via CRISPR-Cas9 technology, which reproduced human

phenotypes. Mechanistically, IQCH can bind to CaM and subsequently regulate the

expression of RNA-binding proteins (especially HNRPAB), which are indispensable

for spermatogenesis. Overall, this study revealed the function of IQCH, expanded the

role of IQ motif-containing proteins in reproductive processes, and provided

important guidance for genetic counseling and genetic diagnosis of male infertility.

Introduction

Introduction

Spermatogenesis is one of the most complex biological processes in male organisms

and functions to produce mature spermatozoa from spermatogonia in three phases: (i)

mitosis, (ii) meiosis, and (iii) spermiogenesis (Hess and Renato de Franca 2008). This

delicate process can be easily disturbed and further cause male reproductive disorders.

Male infertility affects 7% of men in the general population, and its causes vary and

include anatomical or genetic abnormalities, systemic diseases, infections, trauma,

iatrogenic injury, and gonadotoxins (Krausz et al. 2018). Approximately 10% of the

human genome is related to reproductive processes; thus, male infertility is often

predicted to be largely genetic in origin, whereas only 4% of infertile men are

diagnosed with a distinct genetic cause (Krausz and Riera-Escamilla 2018). Genetic

causes are highly heterogeneous and involve chromosomal abnormalities, point

mutations in single genes, copy number variations, microRNA dysfunction, and

polygenic defects (Meschede and Horst 1997, Traven et al. 2017). The highest

percentage of known genetic factors that account for up to 25% of male infertility are

karyotype anomalies, Y chromosome microdeletions, and *CFTR* mutations, which are

mostly associated with azoospermia (Krausz and Riera-Escamilla 2018). However, in

a relatively high proportion of infertile men (40%), the etiology cannot be recognized

and is also referred to as idiopathic. With the help of assisted reproductive techniques

(ART), some men have the chance to reproduce; however, there is a risk of passing on

undetermined genetic abnormalities. In addition, genetic defects leading to

fertilization failure and embryo development arrest cannot be effectively rescued by

ART; thus, discovering novel genetic factors and further confirming their molecular

mechanisms are of clinical importance.

Calmodulin (CaM) is defined as a major Ca^{2+} sensor that activates many kinds of

enzymes in response to an increase in intracellular Ca^{2+} through interactions with a

diverse group of cellular proteins (Klee et al. 1980, Means et al. 1982). CaM has a

completely conserved amino acid sequence across all vertebrates (Jensen et al. 2018).

Specifically, in the human genome, there are three independent CaM genes

(*CALM1-3*), while in the mouse genome, there are five CaM genes (*Calm1-5*) (Jensen

et al. 2018) (https://www.ncbi.nlm.nih.gov/gene/?term=calm+mouse). Previous

studies have reported that CaM is extraordinarily abundant in the mammalian testis

and brain (Smoake et al. 1974, Kakiuchi et al. 1982). Moreover, CaM is

predominantly expressed in the mammalian testis, from the pachytene to meiotic

division stages (Smoake et al. 1974, Kakiuchi et al. 1982, YAMAMOTO 1985, Sano

et al. 1987, Kägi et al. 1988, Moriya et al. 1995). CaM binds to proteins through

recognition motifs, including the short CaM-binding motif containing conserved Ile

and Gln residues (IQ motif) for Ca^{2+}-independent binding and two related motifs for

Ca^{2+}-dependent binding (Nie et al. 2009). The identified IQ motif-containing proteins

included a good range of proteins with biological functions, one of which is a sperm

surface protein, suggesting that IQ motif-containing proteins might play a potential

role in male reproductive processes (Wen et al. 1999, Dolmetsch et al. 2001).

Currently, 15 IQ motif-containing proteins have been identified in humans, namely,

IQCA1, IQCB1, IQCC, IQCD, IQCE, IQCF1, IQCF2, IQCF5, IQCF6, IQCG, IQCH,

IQCJ, IQCK, IQCM, and IQCN. With the exception of IQCF2, mice possess a similar

set of IQ proteins (https://www.ncbi.nlm.nih.gov/gene/?term= IQCF2). Notably,

IQCF4 is present in mice but absent in humans. However, among these IQ

motif-containing proteins, only a few have been reported to be responsible for male

fertility. It has been shown that the IQ motif-containing D (IQCD) primarily

accumulates in the acrosome area of spermatids during mouse spermiogenesis, and

the acrosome reaction is inhibited in human spermatozoa by the anti-IQCD antibody,

suggesting a potential function of IQCD in fertilization and the acrosome reaction

(Zhang et al. 2019). Moreover, *Iqcf1* knockout (KO) male mice showed significantly

less fertility, which was related to reduced sperm motility and acrosome reactions

(Fang et al. 2015). Importantly, *Iqcg* is required for mouse spermiogenesis, which is

attributable to its role in flagellar formation and/or function (Harris et al. 2014).

Humans and mice without IQCN presented failed fertilization rates related to

manchette assembly defects (Dai et al. 2022). Although reliable findings provide

substantial insight into IQ motif-containing proteins that participate in male

reproductive processes, the relevant regulatory mechanism and the function of many

other IQ motif-containing proteins in spermatogenesis have not been determined thus

far.

Here, we revealed a testis-specific IQ motif containing H gene, *IQCH*, which is

essential for spermiogenesis and fertilization. Disruption of IQCH leads to deficient

acrosome activity and abnormal structure of the axoneme and mitochondria in both

humans and mice. Moreover, it is suggested that the interaction of IQCH with CaM is

a prerequisite for IQCH function, which further regulates the expression of

RNA-binding proteins (especially HNRPAB) during spermatogenesis. Collectively,

our findings unveiled a novel genetic diagnostic indicator of male infertility, and the

uncovered mechanism of IQCH in spermatogenesis might shed new light on the

treatment of this disease.

Results

Identification of a novel homozygous variant of *IQCH* involved in male infertility

A 33-year-old man from a consanguineous family with primary infertility for four

years was recruited for our study (Figure 1A). Semen analysis of this patient revealed

inordinate decreases in sperm motility and count, as well as abnormal sperm

morphology (Table 1). In addition to the proband, his great uncle was also affected,

and he had never conceived during sexual intercourse without contraception in his

marriage. We further explored the possible genetic cause through whole-exome

sequencing (WES) analysis, and a homozygous deletion variant in *IQCH*

(c.387+1_387+10del, NM_0010317152) was identified in the proband. This variant is

rare in the general human population according to the 1000 Genomes Project

(0.007%), ExAC Browser (0.059%), and gnomAD databases (0.024%). We further

clarified the putative contribution of this variant in this family by Sanger sequencing.

Noticeably, this homozygous variant was confirmed in his infertile great uncle, and

the fertile parents of the proband carried the heterozygote of this variant (Figure 1A).

Moreover, we used a minigene splicing assay to examine the effect of this variant

on *IQCH* mRNA splicing. The electrophoresis results revealed that wild-type *IQCH*

(WT-*IQCH*) yielded one transcript with a size of 381 bp, whereas the mutant *IQCH*

(Mut-*IQCH*) resulted in one strong band of 263 bp (Figure 1B, i). Sanger sequencing

further showed that the variant resulted in the absence of the whole exon 4, which was

expected to translate into a truncated protein (Figure 1B, ii and iii). Therefore, we

constructed a plasmid containing the aberrant cDNA sequence of *IQCH* caused by the

splicing mutation to verify the effect of the variant on protein expression. The western

blotting results showed that the Mut-c*IQCH* plasmid did not express IQCH, while the

WT-c*IQCH* plasmid did (Figure 1B, iv). We further conducted immunofluorescence

staining of spermatozoa from the proband and the control. We detected very little

expression of the IQCH protein in spermatozoa from the proband compared to the

control (Figure 1C). Taken together, these results reveal that this identified *IQCH*

variant is responsible for splicing abnormalities and further causes a lack of IQCH

expression, which is likely the genetic cause of male infertility in this family.

IQCH deficiency leads to sperm with cracked axoneme structures accompanied

by defects in the acrosome and mitochondria

A comprehensive sperm morphology analysis was further conducted on the proband

(the semen was not available from his great uncle). Papanicolaou staining revealed

that the spermatozoa displayed multiple morphological flagellar abnormalities, such

as coiled, bent, irregular, and even fractured tails (Figure 2A). These morphological

anomalies were confirmed more precisely by scanning electron microscopy (SEM)

analysis (Figure 2B). Significantly, more subtle abnormalities, such as axoneme

exposure, bending, and cracking, were further identified between the midpiece and the

principal piece of the proband's spermatozoa (Figure 2B). As expected, a deranged or

incomplete '9+2' microtubule structure of the flagella was observed in the proband by

transmission electron microscopy (TEM) analysis (Figure 2—figure supplement 1). In

addition, we detected ultrastructural defects in the spermatozoa nucleus, including

irregular shapes, large vacuoles, and deformed acrosomes (Figure 2C). The

mitochondria of the spermatozoa had an abnormal arrangement and enlarged diameter

(Figure 2C).

Moreover, we performed immunofluorescence staining for the marker of the

acrosome (peanut agglutinin: PNA) as well as the mitochondrial marker (transcription

factor A, mitochondrial: TFAM) to confirm the deficiency of the acrosomes and

mitochondria in the proband's spermatozoa. The results of the PNA and TFAM

staining suggested that the spermatozoa acrosomes and mitochondria were severely

defective in the proband compared to the control (Figure 2D, E). In addition, we

performed staining for SEPTIN4 (SEPT4), a functional marker of the annulus, to

explore whether the flagellar fracture at the joint between the middle piece and

principal piece was caused by a nonfunctional annulus. There was no significant

difference in the SEPT4 signal in the spermatozoa between the proband and the

control (Figure 2—figure supplement 1).

Impairment of male fertility in mice without *Iqch*

To further elucidate the role of IQCH in male reproduction, we first explored the

pattern of *Iqch* expression in mice by quantitative real-time PCR (qPCR) and found

that *Iqch* was more highly expressed in the mouse testis than in other organs (Figure

3—figure supplement 1). Additionally, we investigated the temporal expression of

Iqch in mouse testes on different postnatal days. The results revealed that the

expression of *Iqch* significantly increased on postnatal day 21, peaked on postnatal

day 35, and then stabilized (Figure 3—figure supplement 1). To better understand the

role of IQCH in spermatogenesis, we performed immunofluorescence staining of

germ cells at different developmental stages in human and mouse testes. IQCHs

exhibited the same pattern in human and mouse spermatogenesis and were detected

mainly in the cytoplasm of spermatocytes and round spermatids and in the flagella of

late spermatids (Figure 3—figure supplement 1). The testis-enriched expression of

Iqch/IQCH suggested its potential role in spermatogenesis.

We next generated *Iqch* knockout (KO) mice using clustered regularly interspaced

short palindromic repeat (CRISPR)-Cas9 technology. We used a guide RNA targeting

exons 2 through 3 of *Iqch* to achieve the knockout (Figure 3—figure supplement 2).

Polymerase chain reaction (PCR), reverse transcription-PCR (RT□PCR), and western

blotting further confirmed the success of the *Iqch* KO mouse construction (Figure

3—figure supplement 2). The *Iqch* KO mice showed no overt abnormalities in their

development or behavior. The *Iqch* KO female mice showed normal fertility, and

hematoxylin-eosin (H&E) staining further showed that the *Iqch* KO female mice had

normal follicle development compared to that of the WT mice (Figure 3—figure

supplement 2). However, the fertility of the KO male mice was significantly lower

than that of the WT mice, as was the pregnancy rate and litter size (Figure 3—figure

supplement 2). There was no detectable difference in the testis or epididymis/body

weight ratio between the WT and *Iqch* KO mice (Figure 3—figure supplement 2), and

the histology of the testes and epididymides in the *Iqch* KO mice showed no obvious

abnormalities compared to those in the WT mice (Figure 3—figure supplement 3).

However, computer-assisted sperm analysis (CASA) confirmed that sperm

motility was significantly reduced in the KO mice, and the sperm count was slightly

decreased (Table 2; Movies S1 and S2). Moreover, morphological anomalies of the

sperm flagella, such as an unmasking, bending, or cracking axoneme, which

recapitulated the flagellar phenotype of the infertile patient, were easily observed

(Figure 3A, B). We further found that during the spermatogenic process,

abnormalities in axoneme exposure had already occurred when the spermatozoa

flagellum developed (Figure 3—figure supplement 4). Intriguingly, tail defects,

including bending and cracking between the middle piece and the principal piece,

were mainly present in the epididymal spermatozoa (Figure 3—figure supplement 4),

suggesting that more severe flagellum breakage might occur during sperm movement.

We further performed TEM analysis to investigate the ultrastructural defects in

the testicular and epididymal spermatozoa of the *Iqch* KO mice. In contrast to those of

the WT mice, the spermatozoa of the *Iqch* KO mice showed dilated intermembrane

spaces of mitochondria and even loss of some mitochondrial material (Figure 3C).

However, the annulus did not significantly differ between the *Iqch* KO and WT mice

(Figure 3C), indicating that the disruptive link between the flagellar middle piece and

the principal piece might result from mitochondrial defects rather than the annulus.

Immunofluorescence staining of mitochondrial markers (Solute carrier family 25

member 4: SLC25A4) and SEPT4 also revealed that the mitochondria of the

spermatozoa of the *Iqch* KO mice were severely defective, and there was no

significant difference in the annulus of the spermatozoa between the *Iqch* KO mice

and WT mice, which was consistent with the findings of the proband (Figure

3—figure supplement 5).

Poor in vitro fertilization (IVF) outcomes in *Iqch* KO male mice

To evaluate the cause of the impaired fertility of the *Iqch* KO male mice, mature

oocytes from WT female mice in their reproductive period were retrieved and placed

into culture dishes with sperm from the *Iqch* KO mice and WT mice for IVF treatment.

We found that 70% of the embryos had pronuclei in the WT group, while the

fertilization rate of the *Iqch* KO group was dramatically reduced, accounting for 47%

(Figure 4A). Consequently, the percentages of both two-cell embryos and blastocysts

were significantly lower in the *Iqch* KO male mice than in the WT male mice (Figure

4A). Noticeably, an inactive acrosome reaction was observed in most of the sperm

from the *Iqch* KO mice (Figure 4B), leading to the inability to cross the zona

pellucida (Figure 4—figure supplement 1). Furthermore, PNA staining revealed

abnormal acrosome development in the different development germ cells of the *Iqch*

KO mice (Figure 4—figure supplement 1). Similarly, the expression of PNA was

abnormal in the mature sperm of *Iqch* KO male mice (Figure 4C). In the sperm of the

Iqch KO male mice and the proband, the PLCζ fluorescence signal was attenuated and

abnormally localized (Figure 4D; Figure 4—figure supplement 1). Taken together,

these data suggested that *Iqch* might play an important role in acrosomal formation,

which is essential for fertilization.

RNA-binding proteins are the most relevant targets by which IQCH regulates

spermatogenesis

To elucidate the molecular mechanism by which IQCH regulates male fertility, we

performed liquid chromatography tandem mass spectrometry (LC□MS/MS) analysis

using mouse sperm lysates and detected 288 interactors of IQCH (Figure 5—source

data 1). Gene Ontology (GO) analysis of the IQCH-bound proteins revealed many

particularly enriched pathways related to fertilization, sperm axoneme assembly,

mitochondrion organization, calcium channel, and RNA processing (Figure 5A).

Intriguingly, 33 ribosomal proteins were identified (Figure 5B), indicating that IQCH

might be involved in protein synthesis. Using proteomic analysis of sperm from *Iqch*

KO mice, we further assessed key proteins that might be upregulated by IQCH. A

total of 1,993 differentially expressed proteins were quantified, including 807

upregulated proteins and 1,186 downregulated proteins (Figure 5C). GO analysis

revealed that the significantly downregulated proteins were enriched in RNA

processing, gene expression, mitochondrion biogenesis, and calcium ion regulation

(Figure 5D), which was consistent with the enrichment of the IQCH-bound proteins.

Importantly, cross-analysis revealed that 76 proteins were shared between the

IQCH-bound proteins and the downregulated proteins in *Iqch* KO mice (Figure 5E),

suggesting that IQCH might regulate their expression by the interaction. Among the

76 proteins, 21 were RNA-binding proteins (RBPs), 10 of which were suggested to be

involved in spermatogenesis (Kuroda et al. 2000, Yang et al. 2012, Chapman et al.

2013, Fukuda et al. 2013, Legrand et al. 2019, Sechi et al. 2019, Liang et al. 2021,

Tian and Petkov 2021, Wang et al. 2022, Xu et al. 2022); 8 were involved in

mitochondrial function; and 4 were calcium channel activity-related proteins (Figure

5F). We focused on SYNCRIP, HNRNPK, FUS, EWSR1, ANXA7, SLC25A4, and

HNRPAB, which play important roles in spermatogenesis. Specifically, SYNCRIP,

HNRNPK, FUS, EWSR1, and HNRPAB are RBPs that are linked to spermatogenesis

by controlling mRNA translation or spermatid postmeiotic transcription (Kuroda et al.

2000, Fukuda et al. 2013, Sechi et al. 2019, Tian and Petkov 2021, Xu et al. 2022).

ANXA7 is a calcium-dependent phospholipid-binding protein that is a negative

regulator of mitochondrial apoptosis (Du et al. 2015). Loss of SLC25A4 results in

mitochondrial energy metabolism defects in mice (Graham et al. 1997). We further

confirmed the binding of IQCH to these proteins by co-immunoprecipitation (co-IP)

(Figure 6A) and confirmed the downregulation of IQCH in the sperm of *Iqch* KO

mice by immunofluorescence staining and western blotting (Figure 6B; Figure

6—figure supplement 1 and Figure 6—figure supplement 2).

Among these interactors of IQCH, HNRPAB was the most significantly

downregulated protein according to proteomic analysis (Figure 5—source data 2),

implying that HNRPAB might be the main target of IQCH. As an RBP, HNRPAB has

been suggested to play an important role in spermatogenesis by regulating the

translation of testicular mRNAs (Fukuda et al. 2013). We thus employed RNA-seq

analysis of sperm from *Iqch* KO and WT mice to investigate the effects on RNA

levels in the absence of IQCH. Importantly, among the downregulated genes, most

were related to male fertility, such as axoneme assembly, spermatid differentiation,

flagellated sperm motility, and fertilization (Figure 6—figure supplement 3). We

hypothesized that this downregulation is linked to HNRPAB binding. Considering the

disrupted fertilization and axoneme assembly in *Igch* KO mice, we selected the

essential molecules involved in these two processes, *Catsper1*, *Catsper2*, *Catsper3*,

Ccdc40, *Ccdc39*, *Ccdc65*, *Dnah8*, *Irrc6,* and *Dnhd1*, to confirm our speculation. We

carried out RNA immunoprecipitation on formaldehyde-crosslinked sperm followed

by qPCR to evaluate the interactions between HNRPAB and *Catsper1*, *Catsper2*,

Catsper3, *Ccdc40*, *Ccdc39*, *Ccdc65*, *Dnah8*, *Irrc6,* and *Dnhd1* in KO and WT mice.

As expected, binding between HNRPAB and important molecules was detected in the

WT mice (Figure 6C), supporting the functional role of HNRPAB in testicular

mRNAs. Intriguingly, significantly decreased interactions between HNRPAB and

those molecules were observed in the KO mice (Figure 6C). Therefore, IQCH is

involved in spermatogenesis mainly by regulating RBPs, especially HNRPAB, to

further mediate essential mRNA expression in the testis.

The interaction of IQCH with CaM is a prerequisite for IQCH function

Given that IQCH is a calmodulin-binding protein, we hypothesized that IQCH

regulates these key molecules by interacting with CaM (Figure 6—figure supplement

4). As expected, CaM interacted with IQCH, as indicated by LC☐MS/MS analysis.

We initially confirmed the binding of IQCH and CaM in WT sperm by co-IP (Figure

6D). In addition, their colocalization was detected during spermatogenesis by

immunofluorescence staining (Figure 6—figure supplement 4).

We further confirmed the binding of IQCH to CaM in K562 cells, which express

both IQCH and CaM (Figure 6E). Moreover, a decrease in HNRPAB expression was

observed when IQCH or CaM was knocked down (Figure 6E; Figure 6—figure

supplement 2 and Figure 6—figure supplement 4), suggesting that IQCH or CaM

might regulate HNRPAB expression. We then overexpressed IQCH and knocked

down CaM or overexpressed CaM and knocked down IQCH in K562 cells and found

that the reduced HNRPAB expression could not be rescued in either of these two

situations (Figure 6F; Figure 6—figure supplement 2 and Figure 6—figure

supplement 4), indicating that CaM or IQCH alone cannot mediate HNRPAB

expression. These findings suggested that HNRPAB expression might be regulated by

the interaction of IQCH with CaM.

Using a microscale thermophoresis (MST) assay, we further confirmed that the IQ

motif of IQCH is required for CaM binding. The binding affinity between IQCH

(target) from cell lysates overexpressing GFP-*IQCH* plasmids and recombinant CaM

(ligand) increased with increasing concentrations of recombinant CaM (Figure 6G).

However, their binding was disrupted when the IQ motif of IQCH was deleted (Figure

6G). co-IP assays verified the above findings (Figure 6H; Figure 6—figure

supplement 2). Not surprisingly, the expression of HNRPAB in cells cotransfected

with the *IQCH* (△IQ)/*CaM* plasmid and the control/*CaM* plasmid was similar but was

lower than that in cells overexpressing the WT-*IQCH* and *CaM* plasmids (Figure 6H;

Figure 6—figure supplement 2). Collectively, our findings suggest that IQCH

interacts with CaM via the IQ motif to manipulate the expression of important

molecules, especially HNRPAB, to play a role in spermatogenesis (Figure 7).

Discussion

Throughout the evolution of divergent species, IQCH has been conserved, indicating

its fundamental role in organisms. As a novel CaM-binding protein, IQCH was first

identified in human and mouse testes and exclusively localized in spermatocytes and

spermatids, suggesting its potential activity in spermatogenesis (Yin et al. 2005).

However, since IQCH was identified, no other findings have supported its function in

male reproduction. In this study, we found that the *IQCH* mutation impaired male

fertility, as indicated by flagellar morphological abnormalities and axoneme cracking.

An aberrant ultrastructure in sperm with the *IQCH* mutation was associated with

severe defects in acrosomes and mitochondria. Furthermore, *Iqch* KO mice exhibited

similar irregularities in the flagellum, especially the axoneme and mitochondria. We

also found that *Iqch* KO mice exhibited different degrees of functional defects in

acrosomes. Thus, our work supported the vital role of IQCH in flagellar and acrosome

development.

To date, only four IQ motif-containing proteins, IQCD, IQCF1, IQCG, and IQCN,

have been suggested to participate in spermatogenesis (Harris et al. 2014, Fang et al.

2015, Zhang et al. 2019, Dai et al. 2022). Most IQ motif-containing proteins function

in Ca^{2+}-dependent biological processes by binding to CaM (Chen et al. 2014, Harris et

al. 2014, Fang et al. 2015); thus, IQCD, IQCF1, and IQCN are involved in

fertilization and are relevant to the acrosome reaction, sperm capacitation, or

manchette assembly (Fang et al. 2015, Zhang et al. 2019, Dai et al. 2022). In addition,

because CaM has an impact on the actin cytoskeleton, *Iqcg* KO mice exhibit

detachment of sperm heads from tails (Harris et al. 2014). In our study, the *Iqch* KO

mice also showed an impaired acrosome reaction, which caused reduced fertilization.

Intriguingly, axoneme breaks at the annulus and mitochondrial defects were

prominent in the flagella of the *Iqch* KO mice, whose phenotypes were unexplored in

the previously discovered IQ motif-containing proteins. Because CaM activation can

stimulate changes in the actin cytoskeleton, it is reasonable that flagellum formation is

defective when IQCH is absent. Ca^{2+} is a key player in the regulation of

mitochondrial functions (Bravo-Sagua et al. 2017). Thus, the absence of IQCH leads

to a failure of interaction with CaM and disruption of normal Ca^{2+} signaling, which

consequently causes mitochondrial defects in *Iqch* KO mice and patients with *IQCH*

variants. Our findings suggested that fertilization is the main action of IQ

motif-containing proteins, and each specific IQ motif-containing protein also has its

own distinct role in spermatogenesis.

However, there are few publications regarding the underlying mechanism by

which IQ motif-containing proteins manipulate reproductive processes, except for a

recent study showing that IQCN interacts with CaM to regulate the expression of

CaM-binding proteins, including manchette-related proteins (LZTFL1, KIF27, and

RSPH6A), IFT family proteins and their motor proteins (IFT22, IFT43, IFT74, IFT81,

IFT140, IFT172, WDR19, TTC21B, and DYNC2H1), and ribosomal protein family

proteins (RPS5, RPS25, RPS27, and RPSA) (Dai et al. 2022). In our study, IQCH was

shown to regulate the expression of RBPs via interactions with these proteins. RBPs

are a large class of proteins that assemble with RNAs to form ribonucleoproteins

(RNPs). RBPs function at various stages of RNA processing, including alternative

splicing, RNA modification, nuclear export, localization, stability, and translation

efficiency (Corley et al. 2020). RNA processing in male reproductive biology is

essential for the production of mature sperm, and many RBPs have been demonstrated

to be indispensable during spermatogenesis (Morgan et al. 2021). Among the RBPs,

HNRPAB, a heterogeneous nuclear RNP, showed the most significant reduction

among the downregulated interactors of IQCH, suggesting that HNRPAB might be

the most important downstream effector protein of IQCH. It has been reported that

HNRPAB plays a central role in spermatogenesis by regulating stage-specific

translation of testicular mRNAs (Fukuda et al. 2013). As expected, a RNA-binding

protein immunoprecipitation (RIP) assay using an anti-HNRPAB antibody revealed

binding between HNRPAB and the mRNAs of several important genes (*Catsper1*,

Catsper2, *Catsper3*, *Ccdc40*, *Ccdc39*, *Ccdc65*, *Dnah8*, *Irrc6*, and *Dnhd1*) involved in

spermatogenesis. We further found that the interaction between IQCH and CaM is a

prerequisite for regulating HNRPAB expression. We thus hypothesized that IQCH

regulates male fertility by binding to CaM and controlling HNRPAB to manipulate

the expression of key mRNAs involved in spermatogenesis.

In conclusion, our study identified a novel IQ motif-containing protein-coding

gene, *IQCH/Iqch*, which is responsible for spermatogenesis in humans and mice,

broadening the scope of known male infertility-associated genes and thus further

illustrating the genetic landscape of this disease. In addition, our study suggested an

unexplored mechanism in which IQCH regulates the expression of key RBPs

involved in spermatogenesis, especially HNRPAB, by interacting with CaM to play

crucial roles in fertilization and axoneme assembly. We believe that our findings will

provide genetic diagnostic markers and potential therapeutic targets for infertile males

with *IQCH* pathogenic variants.

Materials and Methods.

Study participants

A family with male infertility was recruited from West China Second University

Hospital, and healthy Chinese volunteers were enrolled as controls. The study was

conducted according to the tenets of the Declaration of Helsinki, and ethical approval

was obtained from the Ethical Review Board of West China Second University

Hospital, Sichuan University. Each subject signed an informed consent form.

WES and Sanger sequencing

Peripheral blood samples were obtained from all subjects, and genomic DNA was

extracted using a QIAamp DNA Blood Mini Kit (QIAGEN, 51126). The exomes of

the subjects were captured by Agilent SureSelect Human All Exon V6 Enrichment

kits (Agilent) and then sequenced on a HiSeq X-TEN system (Illumina, USA). All

reads were mapped to the human reference sequence (UCSC Genome Browser hg19)

using Burrows–Wheeler Alignment. After quality filtration with the Genome Analysis

Toolkit, functional annotation was performed using ANNOVAR through a series of

databases, including the 1000 Genomes Project, dbSNP, HGMD, and ExAC.

PolyPhen-2, SIFT, MutationTaster, and CADD were used for functional prediction.

Variant verification of *IQCH* in patients was confirmed by Sanger sequencing using

the primers listed in Figure 1-source data 1.

Minigene assay

A minigene splicing assay was used to explore the effect of the *IQCH* variant

(c.387+1_387+10del) on splicing. The WT-*IQCH* and Mut-*IQCH* sequences,

including intron 3, exon 4, and intron 4, were PCR-amplified separately from the

genomic DNA of the control and patient. The two amplified fragments were cloned

and inserted into the minigene vector pSPL3 between the EcoRI and BamHI sites

through the Basic Seamless Cloning and Assembly Kit (TransGen Biotech, China,

CU201-02). After transfection into HEK293T cells by DNA and siRNA transfection

reagent (Polypus, 101000046), the splicing patterns of the transcripts produced from

the WT-*IQCH* and Mut-*IQCH* plasmids were analyzed by RT⎕PCR, gel

electrophoresis, and Sanger sequencing. The primers used for plasmid genomic

amplification and RT□PCR are listed in Figure 1-source data 1.

Iqch **knockout mice**

The animal experiments were approved by the Experimental Animal Management and

Ethics Committee of West China Second University Hospital, Sichuan University. All

animal procedures complied with the guidelines of the Animal Care and Use

Committee of Sichuan University. The mouse *Iqch* gene has 6 transcripts and is

located on chromosome 9. According to the structure of the *Iqch* gene, exons 2~3 of

the *Iqch-201* (ENSMUST00000042322.10) transcript were recommended as the

knockout region. The region contains a 215 bp coding sequence, and deletion of this

region was expected to disrupt IQCH protein function. The sgRNA sequences

synthesized by Sangon Biotech are listed in Figure 3-source data 1. The two

complementary DNA oligos of each sgRNA target were annealed and ligated to the

pUC57-sgRNA plasmid (Addgene) for cloning. The recombinant plasmid was

transformed into DH5α competent cells, and positive clones were screened based on

kanamycin resistance and sequencing. The recombinant plasmid was linearized and

purified by phenol□chloroform extraction. In vitro transcription of the sgRNAs was

performed using the MEGAshortscript Kit (Ambion, AM1354), and the sgRNAs were

purified using the MEGAclear Kit (Ambion, AM1908). Cas9 mRNA was purchased

from TriLink BioTechnologies. The mouse zygotes were coinjected with an RNA

mixture of Cas9 mRNA (~50 ng/µl) and sgRNA (~30 ng/µl). The injected zygotes

were transferred into pseudopregnant recipients to obtain the F0 generation. DNA was

extracted from the tail tissues of the 7-day-old offspring, and PCR amplification was

carried out with genotyping primers. A stable F1 generation (heterozygous mice) was

obtained by mating the positive F0 generation mice with wild-type (WT)

C57BL/6JG-pt mice. The primers used for genotyping are listed in Figure 3-source

data 1.

Mouse fertility testing

To confirm the fertility of the *Iqch* KO male mice, natural mating tests were

conducted. Briefly, six *Iqch* KO and six littermate WT sexually mature male mice

(8-12 weeks old) were paired with two 6- to 8-week-old normal C57BL/6J females

(each male was mated with two female mice) for 6 months. Regarding the fertility of

Iqch KO female mice, WT (n=15) and *Iqch* KO (n=9) females were mated with WT

males for 6 months. The presence of vaginal plugs was used as indicator of successful

mating and reproductive behavior. The vaginal plugs of the mice were examined

every morning. Female mice with vaginal plugs were fed separately, and the number

of pups per litter was recorded on the day of birth.

IVF

Eight-week-old C57BL/6J female mice were superovulated by injecting 5 IU of

pregnant mare serum gonadotropin (PMSG) followed by 5 IU of human chorionic

gonadotropin (hCG) 48 h later. Sperm were released from the cauda epididymis of

10-week-old male mice, and sperm capacitation was performed for 50 min using test

yolk buffer with HEPES (TYH) solution. Cumulus-oocyte complexes (COCs) were

obtained from the ampulla of the uterine tube 14 h after hCG injection. The ampulla

was torn with a syringe needle, and the COCs were gently squeezed onto liquid drops

of human fallopian tube fluid (HTF) medium. The COCs were then incubated with ~5

µl of the sperm suspension in HTF liquid drops at 37 °C under 5% CO_2. After 6 h, the

eggs were washed several times using HTF medium to remove cumulus cells and then

transferred to liquid drops of potassium simplex optimized medium (KSOM).

Acrosome reaction analysis

The spermatozoa were collected from the cauda epididymis and capacitated for 50

min in TYH medium at 37 °C under 5% CO_2. Highly motile sperm were collected

from the upper portion of the medium, and 10 µM calcium ionophore A23187

(Sigma☐Aldrich, C7522) was added to induce the acrosome reaction. After 15 min,

the spermatozoa were spotted on a glass microscope slide, dried, and fixed with 4%

paraformaldehyde (PFA) for 10 min. The acrosomes were stained with Coomassie

brilliant blue.

H&E staining

For staining of the testicular tissues, samples were dissected from the adult mice and

fixed in 4% PFA overnight at 4 °C. The fixed tissues were embedded in paraffin,

sectioned (5 µm thick), dewaxed, and rehydrated. The testis sections were stained

with hematoxylin and eosin solution (Beyotime, C0105M) before imaging using a

microscope (Leica, Germany).

Immunofluorescence staining

For sperm immunostaining, fresh sperm samples were fixed with 4% PFA for 20 min

at room temperature. For the staining of the testis tissues, heat-induced antigen

retrieval was performed in citrate antigen retrieval solution (Beyotime, P0081)

according to the manufacturer's instructions. After permeabilization with 1% Triton

X-100 for 30 min, the sperm slides were blocked with 5% bovine serum albumin

serum for 1 h at room temperature. Primary antibodies were added to the slides, which

were then incubated overnight at 4 °C. After washing three times with PBS, the slides

were incubated with secondary antibodies for 1 h at room temperature. The nuclei

were counterstained with DAPI dye (Sigma□Aldrich, 28718-90-3) and mounted with

antifade mounting medium. Images were captured with a laser-scanning confocal

microscope (Olympus, Japan). Detailed information on the antibodies is provided in

Figure 6-source data 2.

qPCR

Total RNA was extracted from mouse testes using an RNA Easy Fast Tissue/Cell Kit

(Tiangen Biotech, China, 4992732). Approximately 0.3 mg of total RNA was

converted into cDNA with the PrimeScript RT Reagent Kit (Takara, RR037A)

according to the manufacturer's instructions. The cDNAs were individually diluted

10-fold for use as templates for subsequent qPCR with iTaq Universal SYBR Green

Supermix (Bio-Rad, 1725124). Mouse *Gapdh* was used as an internal control. The

mRNA expression of the detected targets was quantified according to the $2^{-\Delta\Delta Ct}$

method. The primers used for qPCR are listed in Figure 3-source data 1and Figure

6-source data 1.

Papanicolaou staining

Semen samples were fixed in 4% paraformaldehyde, and the coated slides were

air-dried before being rehydrated with 80%, 70%, and 50% ethanol and distilled water.

The samples were then stained with Lea's hematoxylin, rinsed with distilled water,

and stained with G-6 orange stain and EA-50. Following staining, the slides were

dehydrated with ethanol and mounted.

SEM

The spermatozoa were fixed in 2.5% phosphate-buffered glutaraldehyde (GA) at room

temperature for 30 min and then deposited on coverslips. The coverslips were

dehydrated via an ascending gradient of 50%, 70%, 95%, and 100% ethanol and

air-dried. The specimens were then attached to specimen holders and coated with gold

particles using an ion sputter coater before being viewed with a JSM-IT300 scanning

electron microscope (JEOL, Japan).

TEM

The precipitate of the spermatozoa was fixed with 2.5% (vol/vol) glutaraldehyde in

0.1 M phosphate buffer (PB) (pH 7.4) and washed two times in PB and two times in

ddH$_2$O. Then, the tissues were immersed in 1% (wt/vol) OsO$_4$ and 1.5% (wt/vol)

potassium ferricyanide aqueous solution at 4 °C for 2 h. After washing, the samples

were dehydrated through graded alcohol (30%, 50%, 70%, 80%, 90%, 100%, 10 min

each) in pure acetone (10 min twice). The samples were then infiltrated in a graded

mixture (3:1, 1:1, 1:3) of acetone and SPI-PON812 resin (21 ml of SPO-PON812, 13

ml of DDSA and 11 ml of NMA), and then the pure resin was changed. The

specimens were embedded in pure resin with 1.5% BDMA and polymerized for 12 h

at 45 °C and 48 h at 60 °C. Ultrathin sections (70 nm thick) were cut with a

microtome (Leica EM UC6, Germany), double-stained with uranyl acetate and lead

citrate, and examined by a transmission electron microscope (TECNAI G2 F20,

Philips, USA).

Proteomic quantification

Proteins were extracted from the sperm samples using radioimmunoprecipitation

assay (RIPA) lysis buffer (Applygen, C1053) supplemented with 1 mM

phenylmethanesulfonyl fluoride (PMSF) and protease inhibitors on ice. The

supernatants were collected following centrifugation at 12,000 × g for 20 min. Protein

concentrations were calculated by Bradford quantification and sodium dodecyl

sulfate□polyacrylamide gel electrophoresis (SDS□PAGE). An enzyme solution at a

ratio of 1:20 of trypsin enzyme (μg) to substrate protein (μg) was added to 100 μg of

the protein samples, which were subsequently vortexed, centrifuged at low speed for 1

min, and incubated at 37 °C for 4 h. The peptide liquid obtained after salt removal

was freeze-dried. The peptide sample was dissolved in 0.5 M TEAB, added to the

corresponding iTRAQ labeling reagent and stored at room temperature for 2 h. The

Shimadzu LC-20AB liquid phase system was used for purification, and the separation

column was a 5 μm 4.6 × 250 mm Gemini C18 column for liquid phase separation of

the sample. The dried peptide samples were reconstituted with mobile phase A (2%

can, 0.1% FA) and centrifuged at 20,000 × g for 10 min, after which the supernatant

was collected for injection. The separation was performed by an UltiMate 3000

UHPLC (Thermo Fisher, USA). The sample was first enriched in a trap column,

desalted and then placed into a self-packed C18 column. The peptides separated by

liquid phase chromatography were ionized by a nanoESI source and then passed to a

Q Exactive HF X tandem mass spectrometer (Thermo Fisher, USA) for DDA

(data-dependent acquisition) mode detection. The raw data were converted to mgf

files for bioinformatics analysis, and protein identification from tandem mass spectra

was performed by database searching (UniProt). The protein quantification process

included the following steps: protein identification, tag impurity correction, data

normalization, missing value imputation, protein ratio calculation, statistical analysis,

and results presentation. Proteins with a 1.5-fold change and a p value (using

Student's t test) less than 0.05 were defined as differentially expressed proteins.

RNA sequencing

Total RNA (~1 μg) was isolated from fresh spermatozoa from the WT mice (n=3) and

the *Iqch* KO mice (n=3) using the RNAsimple Total RNA Kit (Tiangen Biotech,

4992858). A Ribo-Zero™ rRNA Removal Kit (Ilumina, MRZPL1224) was used to

remove rRNA from the samples. RNA integrity was evaluated using an Agilent 2100

Bioanalyzer system (Agilent Technologies). An mRNA sequencing library for

RNA-seq was constructed using the TruSeq RNA Library Prep Kit v2 (Illumina,

RS-122-2001), followed by paired-end (2□×□100□bp) sequencing using the

Illumina HiSeq 4000 Sequencing System (Illumina, USA) at the Beijing Genomics

Institute. The raw paired-end reads were filtered through a FASTX-Toolkit filter. The

quality of the reads was confirmed by FastQC (ver: 0.11.3). The raw data (raw reads)

in the fastq format were processed through in-house Perl scripts. The feature counts

(ver: l.5.0-p3) were used to count the read numbers mapped to each gene. The

fragments per kilobase million (FPKM) of each gene were calculated based on the

length of the gene and the read count mapped to the gene. Signaling pathway

matching analysis for the list of differentially expressed genes (DEGs) was performed

using the Kyoto Encyclopedia of Genes and Genomes (KEGG) Mapper

(https://www.genome.jp/kegg/). The mapping of GO to DEGs was carried out using

Blast2GO (ver: 4.1.9).

LC□MS/MS analysis

Sperm from WT mice were lysed using RIPA buffer supplemented with 1 mM PMSF

and protease inhibitors (Applygen, P1265) on ice. The lysates were then centrifuged

at 14,000 × g for 15 min at 4°C, and the clear supernatants were incubated with 5 μg IQCH antibody (the rabbit polyclonal IQCH antibody was synthesized using the immunogen KAEAATKIQATWKSYKARSSFISYRQKKWA) at 4°C overnight. The next day, the incubated product was mixed with 50 μL protein A/G magnetic beads (Bimake, B23202) and incubated for 2 h at room temperature with rotation. Subsequent washes of the immunoprecipitates were performed three times using a wash buffer (50 mM Tris-HCl, 150 mM NaCl, 0.1% Triton X-100) and boiled in 1× SDS-PAGE buffer at 95 °C for 10 min. The samples were separated by SDS-PAGE and subjected to Coomassie brilliant blue staining. Protein bands were then excised from the gel for subsequent LC-MS/MS analysis. Subsequent LC-MS/MS analysis was performed using a Q Exactive^{TM} HF-X mass spectrometer (Thermo Fisher, USA) equipped with a nanoelectrospray ion source at Jingjie PTM BioLabs. The raw data files were queried against the Mus_musculus_10090_SP_20210721.fasta database, which contains 17,089 entries. The search employed MaxQuant (v.1.6.15.0) and its integrated Andromeda search engine, utilizing default settings, to identify proteins and ascertain their respective label-free quantification (LFQ) values. The LC–MS/MS results underwent two main processing steps: (i) exclusion of contaminant proteins, reverse sequences, and proteins identified solely by site and (ii) normalization of LFQ intensity. The normalization utilized the median of commonly identified proteins in the sample, with missing values imputed by the minimum value.

RIP

RIP assays were utilized to isolate RNA binding with HNRPAB, following a previous study (Fuentes-Iglesias et al. 2020). Briefly, testicular tissue was disaggregated using a cryogenic tissue homogenizer and subsequently crosslinked in 1% formaldehyde (w/v) for 10 min. Crosslinking was halted by 2.5 M glycine and 25 mM Tris-base buffer for 5 min, and the samples were lysed in RIP lysis buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1% NP-40, 1 mM DTT, 1 mM PMSF, and 100 U/ml RNase Inhibitor) on ice for 10 min. The lysates were then sonicated in an ice bath for 5 cycles (10 seconds on, 20 seconds off), followed by centrifugation at 14,000 g for 10 min at 4°C. The supernatant (10% volume was kept as input) was incubated overnight at 4°C with primary antibodies. Anti-HNRPAB antibody and mouse IgG were employed in the RIP assays. The lysates were subsequently incubated with 75 μL of protein A/G magnetic beads at 4°C for 4 h with rotation. The beads were washed three times with RIP washing buffer (20 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.1% NP-40, 0.1% SDS, 100 U/ml RNase inhibitor) and then incubated at 70°C for 45 min to reverse the crosslinks. Target RNAs from the input and eluate were extracted using TRIzol reagent (Invitrogen, 10296028CN) and reverse-transcribed with the PrimeScript RT Reagent Kit (Takara, RR037A), according to the manufacturer's instructions.

Western blotting and co-IP

Proteins were extracted from cultured cells and mouse sperm using lysis buffer on ice. The supernatants were collected following centrifugation at 14,000 $\times$ g for 20 min.

The proteins were electrophoresed on 10% SDS□PAGE gels and transferred to

nitrocellulose membranes (GE Healthcare). The blots were blocked in 5% milk and

incubated with primary antibodies overnight at 4 °C, followed by incubation with

anti-rabbit or anti-mouse IgG heavy and light chain (H&L) (horseradish peroxidase,

HRP) (Abmart, M21002 and M21001) at a 1/10,000 dilution for 1 h. The signals were

evaluated using Super ECL Plus western blotting Substrate (Applygen, P1050) and a

Tanon-5200 Multi chemiluminescence imaging system (Tanon, China).

For the co-IP assays, the extracted proteins were incubated with primary

antibodies overnight at 4 °C. The lysates were then incubated with 20 μl of Pierce™

Protein A/G-conjugated Agarose for 2 h at room temperature. The beads were washed

with washing buffer, eluted with 1.2×SDS loading buffer, and boiled for 5 min at

95 °C. Finally, the products were separated via SDS□PAGE and analyzed by

immunoblotting. Detailed information on the antibodies used in the western blotting

experiments is provided in Figure 6-source data 2.

Antibody production

The peptide used for raising the anti-IQCH antibody was derived from amino acid

residues 406–435 (KAEAATKIQATWKSYKARSSFISYRQKKWA) of mouse

IQCH. The peptide coupled with keyhole limpet hemocyanin (KLH) (Sigma□Aldrich,

H7017) was dissolved in saline, emulsified with 1□ml of Freund's complete adjuvant

(Beyotime, P2036), and injected at multiple sites on the backs of New Zealand white

rabbits. The antiserum was collected within 2 weeks after the final injection.

Cell culture and transfection

We purchased K562 cells (CRL-3344) and HEK293T cells (CRL-11268) from the

American Type Culture Collection. The K562 cell line is a human leukemia cell line

showing high expression levels of IQCH and CaM and was used for the knockdown

experiments. K562 cells were cultured in Basic Roswell Park Memorial Institute 1640

Medium (Gibco, C11875500BT) supplemented with 10% fetal bovine serum (Gibco,

12483020). HEK293T cells were cultured in Dulbecco's modified Eagle medium

(Gibco, 11965092) supplemented with 10% fetal bovine serum (Gibco, 12483020).

The expression plasmids pcDNA3.1-Flag-*CALM2*, pCMV-MCS-3* flag-WT-*IQCH,*

and pCMV-MCS-3* flag-*IQCH (*△IQ*)* and small interfering RNAs (siRNAs)

targeting *CALM2* and *IQCH* were constructed by Vigene Biosciences (Jinan, China).

The sequences of the siRNAs used are listed in Figure 6—source data 1. The plasmids

and siRNAs were transfected into cells with jetPRIME transfection reagents (Polypus,

101000046) according to the manufacturer's protocol.

MST assay

The MST experiments were conducted on a Monolith NT.115 system (NanoTemper

Technologies, Germany). Lysates of the HEK293T cells transfected with fluorescent

GFP-*IQCH* or GFP-*IQCH*(△IQ) were normalized by raw fluorescence (count), diluted

using MST buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 10 mM MgCl2, 0.05%

(v/v) Tween 20), and added to 16 PCR tubes (10 μl per tube). Then, the purified CaM

was diluted into 16 gradients using MST buffer. Ten microliters of different

concentrations of CaM were mixed with 10 μl of fluorescent GFP-IQCH protein and

GFP-*IQCH*(△IQ) and reacted in a dark box for 15 min at room temperature. The

samples were added to monolith capillaries (NanoTemper, MO-L022) and

subsequently subjected to MST analysis. The measurement protocol times were as

follows: fluorescence before 5 sec, MST after 30 sec, fluorescence after 5 sec, and

delay of 25 sec. The dissociation constant (Kd) was determined using a single-site

model to fit the curve.

Statistical analysis

The data were compared for statistical significance using GraphPad Prism version

9.0.0 (GraphPad Software). The unpaired, two-tailed Student's t test was used for the

statistical analyses. The data are presented as the mean ± SEM, and statistically

significant differences are represented as $*p<0.05$.

References

Bravo-Sagua, R., V. Parra, C. López-Crisosto, P. Díaz, A. F. Quest and S. Lavandero (2017).

"Calcium Transport and Signaling in Mitochondria." Compr Physiol **7**(2): 623-634.

Chapman, K. M., H. M. Powell, J. Chaudhary, J. M. Shelton, J. A. Richardson, T. E.

Richardson and F. K. Hamra (2013). "Linking spermatid ribonucleic acid (RNA) binding protein

and retrogene diversity to reproductive success." Mol Cell Proteomics **12**(11): 3221-3236.

Chen, L. T., W. X. Liang, S. Chen, R. K. Li, J. L. Tan, P. F. Xu, L. F. Luo, L. Wang, S. H. Yu, G.

Meng, K. K. Li, T. X. Liu, Z. Chen and S. J. Chen (2014). "Functional and molecular features of

the calmodulin-interacting protein IQCG required for haematopoiesis in zebrafish." Nat

Commun **5**: 3811.

Corley, M., M. C. Burns and G. W. Yeo (2020). "How RNA-Binding Proteins Interact with RNA:

Molecules and Mechanisms." Mol Cell **78**(1): 9-29.

Dai, J., Q. Li, Q. Zhou, S. Zhang, J. Chen, Y. Wang, J. Guo, Y. Gu, F. Gong, Y. Tan, G. Lu, W.

Zheng and G. Lin (2022). "IQCN disruption causes fertilization failure and male infertility due to

manchette assembly defect." EMBO Mol Med **14**(12): e16501.

Dolmetsch, R. E., U. Pajvani, K. Fife, J. M. Spotts and M. E. Greenberg (2001). "Signaling to

the nucleus by an L-type calcium channel-calmodulin complex through the MAP kinase

pathway." Science **294**(5541): 333-339.

Du, Y., Y. Huang, Y. Gao, B. Song, J. Mao, L. Chen, L. Bai and J. Tang (2015). "Annexin A7

modulates BAG4 and BAG4-binding proteins in mitochondrial apoptosis." Biomed

Pharmacother **74**: 30-34.

Fang, P., W. Xu, D. Li, X. Zhao, J. Dai, Z. Wang, X. Yan, M. Qin, Y. Zhang, C. Xu, L. Wang and

Z. Qiao (2015). "A novel acrosomal protein, IQCF1, involved in sperm capacitation and the

acrosome reaction." Andrology **3**(2): 332-344.

Fuentes-Iglesias, A., V. Garcia-Outeiral, J. A. Pardavila, J. Wang, M. Fidalgo and D. Guallar

(2020). "An Optimized Immunoprecipitation Protocol for Assessing Protein-RNA Interactions In

Vitro." STAR Protoc **1**(2).

Fukuda, N., T. Fukuda, J. Sinnamon, A. Hernandez-Hernandez, M. Izadi, C. S. Raju, K.

Czaplinski and P. Percipalle (2013). "The transacting factor CBF-A/Hnrnpab binds to the

A2RE/RTS element of protamine 2 mRNA and contributes to its translational regulation during

mouse spermatogenesis." PLoS Genet **9**(10): e1003858.

Graham, B. H., K. G. Waymire, B. Cottrell, I. A. Trounce, G. R. MacGregor and D. C. Wallace

(1997). "A mouse model for mitochondrial myopathy and cardiomyopathy resulting from a

deficiency in the heart/muscle isoform of the adenine nucleotide translocator." Nat Genet **16**(3):

226-234.

Harris, T. P., K. J. Schimenti, R. J. Munroe and J. C. Schimenti (2014). "IQ motif-containing G

(Iqcg) is required for mouse spermiogenesis." G3 (Bethesda) **4**(2): 367-372.

Hess, R. A. and L. Renato de Franca (2008). "Spermatogenesis and cycle of the seminiferous

epithelium." Adv Exp Med Biol **636**: 1-15.

Jensen, H. H., M. Brohus, M. Nyegaard and M. T. Overgaard (2018). "Human Calmodulin

Mutations." Front Mol Neurosci **11**: 396.

Kägi, U., J. G. Chafouleas, A. W. Norman and C. W. Heizmann (1988). "Developmental

appearance of the Ca2+-binding proteins parvalbumin, calbindin D-28K, S-100 proteins and

calmodulin during testicular development in the rat." Cell Tissue Res **252**(2): 359-365.

Kakiuchi, S., S. Yasuda, R. Yamazaki, Y. Teshima, K. Kanda, R. Kakiuchi and K. Sobue

(1982). "Quantitative determinations of calmodulin in the supernatant and particulate fractions

of mammalian tissues." J Biochem **92**(4): 1041-1048.

Klee, C. B., T. H. Crouch and P. G. Richman (1980). "Calmodulin." Annu Rev Biochem **49**:

489-515.

Krausz, C., F. Cioppi and A. Riera-Escamilla (2018). "Testing for genetic contributions to

infertility: potential clinical impact." Expert Rev Mol Diagn **18**(4): 331-346.

Krausz, C. and A. Riera-Escamilla (2018). "Genetics of male infertility." Nat Rev Urol **15**(6):

369-384.

Kuroda, M., J. Sok, L. Webb, H. Baechtold, F. Urano, Y. Yin, P. Chung, D. G. de Rooij, A.

Akhmedov, T. Ashley and D. Ron (2000). "Male sterility and enhanced radiation sensitivity in

TLS(-/-) mice." Embo j **19**(3): 453-462.

Legrand, J. M. D., A. L. Chan, H. M. La, F. J. Rossello, M. L. Änkö, F. V. Fuller-Pace and R. M.

Hobbs (2019). "DDX5 plays essential transcriptional and post-transcriptional roles in the

maintenance and function of spermatogonia." Nat Commun **10**(1): 2278.

Liang, J., Y. Zheng, W. Zeng, L. Chen, S. Yang, P. Du, Y. Wang, X. Yu and X. Zhang (2021).

"Comparison of proteomic profiles from the testicular tissue of males with impaired and normal

spermatogenesis." Syst Biol Reprod Med **67**(2): 127-136.

Means, A. R., J. S. Tash and J. G. Chafouleas (1982). "Physiological implications of the

presence, distribution, and regulation of calmodulin in eukaryotic cells." Physiol Rev **62**(1):

1-39.

Meschede, D. and J. Horst (1997). "The molecular genetics of male infertility." Mol Hum

Reprod **3**(5): 419-430.

Morgan, M., L. Kumar, Y. Li and M. Baptissart (2021). "Post-transcriptional regulation in

spermatogenesis: all RNA pathways lead to healthy sperm." Cell Mol Life Sci **78**(24):

8049-8071.

Moriya, M., K. Fujinaga, M. Yazawa and C. Katagiri (1995). "Immunohistochemical localization

of the calcium/calmodulin-dependent protein phosphatase, calcineurin, in the mouse testis: its

unique accumulation in spermatid nuclei." Cell Tissue Res **281**(2): 273-281.

Nie, D., X. Yang and Z. Yankai (2009). "Molecular cloning and expression profile analysis of a

novel mouse testis-specific expression gene mtIQ1." Mol Biol Rep **36**(5): 1203-1209.

Sano, M., A. S. Ohshima, N. Kawamura, S. Kitajima and A. Mizutani (1987).

"Immunohistochemical study of calmodulin in developing mouse testis." J Exp Zool **241**(1):

51-59.

Sechi, S., A. Frappaolo, A. Karimpour-Ghahnavieh, M. Gottardo, R. Burla, L. Di Francesco, E.

Szafer-Glusman, E. Schininà, M. T. Fuller, I. Saggio, M. G. Riparbelli, G. Callaini and M. G.

Giansanti (2019). "Drosophila Doublefault protein coordinates multiple events during male

meiosis by controlling mRNA translation." Development **146**(22).

Smoake, J. A., S. Y. Song and W. Y. Cheung (1974). "Cyclic 3',5'-nucleotide

phosphodiesterase. Distribution and developmental changes of the enzyme and its protein

activator in mammalian tissues and cells." Biochim Biophys Acta **341**(2): 402-411.

Tian, H. and P. M. Petkov (2021). "Mouse EWSR1 is crucial for spermatid post-meiotic

transcription and spermiogenesis." Development **148**(11).

Traven, E., A. Ogrinc and T. Kunej (2017). "Initiative for standardization of reporting genetics

of male infertility." Syst Biol Reprod Med **63**(1): 58-66.

Wang, X. L., J. M. Li and S. Q. Yuan (2022). "Characterization of the protein expression and

localization of hnRNP family members during murine spermatogenesis." Asian J Androl.

Wen, Y., R. T. Richardson and G. O'Rand M (1999). "Processing of the sperm protein Sp17

during the acrosome reaction and characterization as a calmodulin binding protein." Dev Biol

206(2): 113-122.

Xu, H., J. Guo, W. Wu, Q. Han, Y. Huang, Y. Wang, C. Li, X. Cheng, P. Zhang and Y. Xu

(2022). "Deletion of Hnrnpk Gene Causes Infertility in Male Mice by Disrupting

Spermatogenesis." Cells **11**(8).

YAMAMOTO, N. J. A. h. e. c. (1985). "Immunoelectron microscopic localization of calmodulin

in guinea pig testis and spermatozoa." **18**(2): 199-211.

Yang, F., Q. Wei, R. S. Adelstein and P. J. Wang (2012). "Non-muscle myosin IIB is essential

for cytokinesis during male meiotic cell divisions." Dev Biol **369**(2): 356-361.

Yin, L. L., J. M. Li, Z. M. Zhou and J. H. Sha (2005). "Identification of a novel testis-specific

gene and its potential roles in testis development/spermatogenesis." Asian J Androl **7**(2):

127-137.

Zhang, P., W. Jiang, N. Luo, W. Zhu and L. Fan (2019). "IQ motif containing D (IQCD), a new

acrosomal protein involved in the acrosome reaction and fertilisation." Reprod Fertil Dev **31**(5):

898-914.

Acknowledgments: We thank the patient and the family members for their support

during this research study.

Competing interests: The authors declare that they have no competing interests.

Author contributions: Y.S. designed and supervised the study experiments. Y.Y.

collected the data and conducted the clinical evaluations. T.R., R.Z., and Y.S.

performed the experiments and analyzed most of the data. S.C. generated the CRISPR

mice. J.G., C.J., X.W., G.S. and S.D. performed the cell experiments. Y.S. wrote the

manuscript with input from others. All authors revised and approved the article.

Ethics Approval and Consent to Participate: For human participants, this study was

performed in accordance with the principles of the Declaration of Helsinki. Approval

was granted by the Ethics Committee of the Second West China Hospital of Sichuan

University. Informed consent was obtained from all individual participants included in

the study. The animal experiments were performed in accordance with the

recommendations of the Guide for the Care and Use of Laboratory Animals of the

National Institutes of Health, and the animal experiments were approved by the

Experimental Animal Management and Ethics Committee of West China Second

University Hospital (IACUC no. 2021(070)).

Funding: This study was funded by the National Key Research and Development

Project (2019YFA0802101)

Data and materials availability: All data needed to evaluate the conclusions in the

paper are present in the paper and/or the Supplementary Materials.

 Figure legends

Figure 1. Identification of a homozygous splicing mutation in *IQCH* in a

consanguineous family with male infertility. (A) Pedigree analysis of the

consanguineous family with two infertile males (II-1 and IV-1), with the black arrow

pointing to the proband. Sanger sequencing revealed that the affected males exhibited

a homozygous variant in *IQCH*. Sequence chromatograms are shown below the

pedigree. **(B)** (i) The electrophoresis results of the minigene assay show a decrease in

the molecular weight of the RT□PCR products generated from Mut-*IQCH* (263 bp)

compared with those from WT-*IQCH* (381 bp). (ii) Sanger sequencing of the

complementary DNA of the splicing mutation showing the deletion of exon 4 in

IQCH. (iii) The pattern diagram demonstrating the splicing effects caused by the

IQCH mutation. (iv) Western blotting results showed that the Mut–c*IQCH* plasmids

did not express IQCH. NC, negative control. Three independent experiments were

performed. **(C)** Immunofluorescence staining showed that the expression of IQCH

was barely detected in the proband's sperm compared with that in the control (blue,

DAPI; green, α-Tubulin; red, IQCH; scale bars, 5 μm).

Figure 2. Abnormal flagellar morphology and defective acrosomes and

mitochondria in the infertile patient. (A and **B)** Papanicolaou staining (A) and

scanning electron microscopy (SEM) (B) results showing flagellar morphological

abnormalities (scale bars in A, 5 μm; scale bars in B, 2.5 μm). The dotted boxes and

red arrowheads denote axoneme cracking and exposure. **(C)** Transmission electron

microscopy (TEM) results showing deformed acrosomes and abnormal arrangement

and diameter of mitochondria (scale bars, 500 nm). The red arrowheads denote

abnormal mitochondria. N, nucleus; M, mitochondria. **(D** and **E)** Defects in the

acrosome and mitochondria were observed in the proband's sperm by PNA (D) and

TFAM (E) staining (blue, DAPI; green, α-Tubulin; red, PNA or TFAM; scale bars, 5

μm). PNA, peanut agglutinin; TFAM, transcription factor A, mitochondrial.

Figure 3. The absence of *Iqch* impaired spermatogenesis in mice. (A and **B)**

Papanicolaou staining (B) and SEM (C) results showing unmasking, bending, or

cracking of the axoneme in spermatozoa from *Iqch* KO mice (n = 3 biologically

independent WT mice and KO mice; scale bars in B, 5 μm; scale bars in C, 2.5 μm).

The red arrowheads point to axoneme abnormalities. **(C)** TEM revealing dilated

intermembrane spaces of mitochondria and a normal annulus in the testicular and

epididymal spermatozoa of *Iqch* KO mice (n = 3 biologically independent WT mice

and KO mice; scale bars, 500 nm). The red arrowheads point to the dilated

intermembrane spaces of mitochondria. The green arrowheads point to the normal

annulus. M, mitochondria.

Figure 4. Poor IVF treatment outcomes resulting from the use of sperm from

***Iqch* KO mice. (A)** Representative pronucleus embryos, two-cell embryos, and

blastocysts obtained from WT mice and *Iqch* KO mice. *Iqch* KO mice exhibited

significantly lower fertilization rates, cleavage rates, and blastocyst formation rates

than WT mice (n = 3 biologically independent WT mice and KO mice; scale bars, 100

μm; Student's t test; *p < 0.05; error bars, s.e.m.). **(B)** The acrosome reaction rates in

the capacitated spermatozoa from the WT mice and *Iqch* KO mice were determined

by Coomassie brilliant blue staining. The acrosome reaction rates were reduced in the

spermatozoa from the *Iqch* KO mice (n = 3 biologically independent WT mice and

KO mice; scale bars, 5 μm; Student's t test; *p < 0.05; error bars, s.e.m.). The green

arrowheads indicate the reacted acrosomes. The red arrowheads indicate intact

acrosomes. (**C** and **D**) PNA (C) and PLCζ (D) staining showing abnormal acrosome

morphology and aberrant PLCζ localization and expression in spermatozoa from *Iqch*

KO mice (n = 3 biologically independent WT mice and KO mice; scale bars, scale

bars, 5 μm). The dotted box indicates the typical pattern of PLCζ localization and

expression in spermatozoa from WT mice. PNA, peanut agglutinin; PLCζ,

phospholipase C zeta 1.

Figure 5. IQCH bound and upregulated male reproduction-related proteins in

mouse sperm **(A)** Bubble plots of the GO analysis showing that the IQCH-interacting

proteins are significantly enriched in spermatogenesis and RNA processing in three

categories: biological process, cellular component, and molecular function. GO, Gene

Ontology. **(B)** Thirty-three ribosomal proteins interacted with IQCH in mouse sperm.

(C) A heatmap showing the differential protein results from the proteomic analysis of

the sperm from the WT and *Iqch* KO mice. **(D)** Bubble plots showing the decreased

enrichment of proteins related to the spermatogenetic process and RNA processing

according to the GO analysis of the *Iqch* KO mice compared to the WT mice. **(E)**

Venn diagram depicting the 76 overlapping proteins among the 1186 downregulated

proteins in sperm from *Iqch* KO mice and the 288 proteins that bind to IQCH. **(F)**

Chord diagram showing 21 proteins involved in RNA binding, 8 proteins involved in

mitochondrial function, and 4 proteins involved in calcium channel activity among the

76 overlapping proteins. HNRPAB showed the greatest reduction in expression in the

Iqch KO mice.

Figure 6. IQCH interacted with CaM to regulate HNRPAB expression and

spermatogenesis. (A) co-IP of mouse sperm lysates revealed that the seven proteins

most relevant to the phenotype of the *Iqch* KO mice were associated with IQCH. **(B)**

Western blotting showing the reduced expression of the seven proteins in the sperm

from the *Iqch* KO mice compared to the WT mice. **(C)** qPCR analysis of RNA

immunoprecipitation (RIP) using HNRPAB antibodies and IgG antibodies on mouse

sperm lysates showed that HNRPAB interacted with several RNAs associated with

fertilization and axoneme assembly. qPCR analysis of RIP in the sperm lysates from

the *Iqch* KO mice revealed a decrease in the interaction between HNRPAB and the

RNA targets compared to that in the WT mice (Student's t test; *p < 0.05; error bars,

s.e.m.). **(D)** co-IP assays showing the binding of IQCH and CaM in WT sperm. **(E)**

co-IP assays showed that the decreased expression of HNRPAB was due to the

reduced binding of IQCH to CaM resulting from the knockdown of IQCH or CaM. **(F)**

The overexpression of IQCH and the simultaneous knockdown of CaM or the

overexpression of CaM and the simultaneous knockdown of IQCH in K562 cells

further confirmed that the downregulation of HNRPAB was due to the diminished

interaction between IQCH and CaM, as determined by western blotting analysis. **(G)**

Analysis of the binding affinity between GFP-IQCH from the cell lysates (target) and

recombinant CaM (ligand) by a microscale thermophoresis (MST) assay showing the

interaction between IQCH and CaM. Their interaction was disrupted after deletion of

the IQ motif within IQCH. **(H)** co-IP of HEK293 cells cotransfected with the

WT-*IQCH* and WT-*CaM* plasmids, cotransfected with the *IQCH* (△IQ) and WT-*CaM*

plasmids, or cotransfected with the control and WT-*CaM* plasmids showed that CaM

interacted with the IQ motif. The downregulation of HNRPAB was due to the

disrupted interaction between IQCH and CaM. Three independent experiments were

performed.

Figure 7. Proposed model for the mechanisms underlying the involvement of

IQCH in spermatogenesis. IQCH interacts with CaM via the IQ motif to regulate the

expression of RNA-binding proteins. RNA binding proteins, particularly HNRPAB,

bind and regulate several RNAs that influence the development of the acrosome,

mitochondria, and axoneme, thereby playing a critical role in spermatogenesis.

Table 1: Results of the semen analysis and sperm morphology examination of the patient

Parameter	Patient	Reference*
Sperm volume, ml	2.8	≥1.5
Sperm concentration, million/ml	10.0	≥15
Vitality, %	30	≥58
Motility, %	4	≥32
Abnormal morphology (%)	99.5	-

*: The reference values were based on the guidelines provided in the WHO manual 5th edition (2010).

971 **Table 2. Semen analysis using CASA in the mouse model of *Iqch* KO**

	Adult Male Mice		
	WT	KO	P[b] value
Semen parameters			
Sperm concentration (10^6/ml) [a]	94.51±10.68	67.75±3.70	0.038
Motility (%)	57.08±2.18	5.49±3.25	<0.001
Progressive motility (%)	57.00±2.05	5.49±3.25	<0.001
Sperm locomotion parameters			
Curvilinear velocity (VCL) (μm/s)	78.29±6.23	9.94±4.03	<0.001
Straight-line velocity (VSL) (μm/s)	36.18±2.09	2.90±1.98	<0.001
Average path velocity (VAP) (μm/s)	45.10±0. 19	4.17±2.49	<0.001
Amplitude of lateral head displacement (ALH) (μm)	0.77±0. 02	0.13±0.03	<0.001
Linearity (LIN)	0.46±0.01	0.27±0.08	0.047
Wobble (WOB, = VAP/VCL)	0.58±0.04	0.40±0.08	0.033
Straightness (STR, = VSL/VAP)	0.80±0.05	0.67±0.07	0.069
Beat-cross frequency (BCF) (Hz)	4.33±0.19	0.65±0.26	<0.001

[a] Epididymides and vas deferens, [b] two-tailed Student's t test

n = 3 biologically independent WT mice or KO mice.

Supplementary information

Figure 2—figure supplement 1. TEM and immunofluorescence analysis of the

abnormal phenotype of spermatozoa in the patient. (A) TEM results showing the

deranged or incomplete '9+2' microtubule structure of the flagella. The percentage of

abnormal '9+2' microtubule structures in the control and proband (scale bars, 500 nm;

Student's t test; *p < 0.05; error bars, s.e.m.). CP, central-pair microtubule; MTD,

peripheral microtubule doublet; ODF, outer dense fiber; M, mitochondrion. **(B)**

Immunofluorescence staining of SEPTIN4 showing no significant difference in the

annulus between spermatozoa from the control and the proband (blue, DAPI; green,

α-Tubulin; red, SEPTIN4; scale bars, 5 μm).

Figure 3—figure supplement 1. The expression pattern of IQCH in testes. (A)

qPCR analysis showing that *Iqch* was predominantly expressed in the testes, and *Iqch*

began to be significantly expressed on P21, reached its peak on P35, and then showed

a stable expression pattern (Student's t test; *p < 0.05; error bars, s.e.m.). P, postnatal

day. **(B)** Immunofluorescence staining of IQCH in isolated human germ cells (blue,

DAPI; red, IQCH; scale bars, 5 μm). **(C)** Immunofluorescence staining of IQCH in

isolated mouse germ cells (blue, DAPI; red, IQCH; scale bars, 5 μm). Three

independent experiments were performed.

Figure 3—figure supplement 2. Generation of *Iqch* KO mice. (A) A schematic

illustration of the strategy used to generate the *Iqch* KO mice. Exons 2 through 3 of

the *Iqch* gene were targeted by the CRISPR-Cas9 system. **(B)** PCR, RT☐PCR, and

western blotting were used to determine the genotype and KO efficiency (n = 3

biologically independent WT mice and KO mice). **(C)** Female fertility was unaffected

in the *Iqch* KO mice according to the mouse fertility test. *Iqch* KO female mice (n=9)

and WT female mice (n=15) were mated with WT male mice for 6 months at a

male/female ratio of 1:3 (Student's t test; ns, not significant; error bars, s.e.m.). **(D)**

Representative images of H&E-stained WT and *Iqch* KO ovary tissue sections (n = 3

biologically independent WT mice and KO mice; scale bars, 75 μm). **(E)** The number

of primordial follicles, primary follicles, secondary follicles, and antral follicles in the

ovaries of the WT and *Iqch* KO mice (n = 3 biologically independent WT mice and

KO mice; Student's t test; ns, not significant; error bars, s.e.m.). **(F)** Fertility of the

WT and *Iqch* KO males. KO male mice and littermate WT male mice were bred with

WT female mice. The average pregnancy rates and litter sizes were measured (n = 6

biologically independent WT mice and KO mice; Student's t test; *p < 0.05; error

bars, s.e.m.). preg, pregnancy. **(G)** The sizes of the epididymides and testes of the WT

and *Iqch* KO mice showing no significantly different. The testis and epididymis/body

weight ratio of the *Iqch* KO mice were not significantly different from those of the

WT male mice (n = 3 biologically independent WT mice and KO mice; Student's t

test; ns, not significant; error bars, s.e.m.).

1016

Figure 3—figure supplement 3. No obvious abnormalities in the histology of the

testes or epididymis were observed in the *Iqch* KO mice. (**A**) H&E staining of

testicular sections from adult *Iqch* KO mice show no obvious abnormalities at

different stages of spermatogenesis compared to those from WT mice (n = 3

biologically independent WT mice and KO mice; scale bars, 50 μm). (**B**) H&E

staining of epididymal sections from adult WT and *Iqch* KO mice reveal no

significant difference in the lumens of the epididymides (n = 3 biologically

independent WT mice and KO mice; scale bars, 50 μm).

1025

Figure 3—figure supplement 4. (**A**) Papanicolaou staining of the isolated germ cells

from the testes of the *Iqch* KO mice revealed cracking of the axoneme at steps 13-14

and 15-16 (n = 3 biologically independent WT mice and KO mice; scale bars, 5 μm).

(**B**) The cracking of the axoneme became more severe in the isolated epididymal

spermatozoa than in the testicular spermatozoa (n = 3 biologically independent WT

mice and KO mice; scale bars, 5 μm). The dotted boxes and red arrowheads denote

axoneme cracking and exposure.

1033

Figure 3—figure supplement 5. The spermatozoa from the *Iqch* KO mice

exhibited defective mitochondria. (**A**) Immunofluorescence staining of SEPTIN4

revealed no significant differences at the annulus between the spermatozoa from the

WT mice and those from the *Iqch* KO mice (n = 3 biologically independent WT mice

and KO mice; scale bars, 5 μm). (**B**) Immunofluorescence staining of TFAM revealed

defects in the mitochondria of the spermatozoa from the *Iqch* KO mice (n = 3

biologically independent WT mice and KO mice; scale bars, 5 μm). TFAM,

Transcription Factor A, Mitochondrial.

1042

Figure 4—figure supplement 1. The loss of function of IQCH resulted in

deficiencies in oocyte activation and acrosomes. (A) Representative images of the

IVF treatments showing that the spermatozoa from the *Iqch* KO mice did not

penetrate the zona pellucida (n = 3 biologically independent WT mice and KO mice;

scale bars, 50 μm). The red arrowheads indicate the spermatozoa penetrating the zona

pellucida. **(B)** Immunofluorescence staining of the germ cells at different

developmental stages revealed that the development of the PNA was impaired in *Iqch*

KO mice (n = 3 biologically independent WT mice and KO mice; blue, DAPI; purple,

PNA; scale bars, 50 μm). **(C)** Immunofluorescence staining suggested that the loss of

function of IQCH resulted in the abnormal location and expression of PLCζ in the

sperm of the proband (blue, DAPI; green, α-Tubulin; red, PLCζ; scale bars, 50 μm).

(D) Immunofluorescence staining of germ cells at different developmental stages

from *Iqch* KO mice revealed the aberrant expression of PLCζ compared to that in WT

mice (n = 3 biologically independent WT mice and KO mice; blue, DAPI; green,

α-Tubulin; red, PLCζ; scale bars, 50 μm). PNA, peanut agglutinin; PLCζ,

phospholipase C zeta 1.

1059

Figure 6—figure supplement 1. The reduced expression levels of the seven

proteins most relevant to the phenotype of the spermatozoa of the *Iqch* KO mice.

Immunofluorescence staining showing that the expression levels of the seven proteins

most relevant to the phenotype of the *Iqch* KO mice, including HNRNPK, HNRPAB,

EWSR1, SYNCRIP, FUS, and ANXA7, were reduced in the spermatozoa of the *Iqch*

KO mice compared to those of the WT mice (n = 3 biologically independent WT mice

and KO mice; blue, DAPI; green, α-Tubulin; red, HNRNPK, HNRPAB, EWSR1,

SYNCRIP, FUS, or ANXA7; scale bars, 5 μm).

1068

Figure 6—figure supplement 2. **(A)** Densitometry analysis performed on

corresponding Western blotting to assess the expression levels of seven proteins most

relevant to the *Iqch* KO mouse phenotype in WT and *Iqch* KO mice (n = 3

biologically independent WT mice and KO mice; Student's t test; *p < 0.05; error

bars, s.e.m.). **(B** and **C)** Statistical analysis was performed on the protein expression

levels of CaM, IQCH, and HNRPAB, as well as on the interaction between IQCH and

CaM in K562 cells with IQCH (B) or CaM (C) knockdown (Student's t test; *p < 0.05;

error bars, s.e.m.). siRNA, small interfering RNA. **(D** and **E)** Quantitative analysis

was conducted on the protein expression levels of CaM, IQCH, and HNRPAB, as well

as on the interaction between IQCH and CaM in K562 cells overexpressing IQCH

with simultaneous knockdown of CaM (D) or overexpressing CaM with simultaneous

knockdown of IQCH (E) (Student's t test; *p < 0.05; error bars, s.e.m.). siRNA, small

interfering RNA. **(F)** Statistical analysis was conducted on the protein expression

levels of CaM, IQCH, and HNRPAB, as well as on the interaction between IQCH and

CaM in HEK293 cells cotransfected with the WT-*IQCH* and WT-*CaM* plasmids,

cotransfected with the *IQCH* (△IQ) and WT-*CaM* plasmids, or cotransfected with the

control and WT-*CaM* plasmids (Student's t test; *p < 0.05; error bars, s.e.m.). Three

independent experiments were performed.

1087

1088

1089   **Figure 6—figure supplement 3. GO analysis of the downregulated genes of the**

1090   ***Iqch* KO mice through RNA sequencing.** Bubble plots of the GO analysis revealed

1091   the effects of IQCH deficiency on male fertility in three categories: BP, biological

1092   process; CC, cellular component; and MF, molecular function.

1093

Figure 6—figure supplement 4. IQCH interacted with CaM to regulate

spermatogenesis. (A) A schematic representation of IQCH showing an IQ motif

conserved between humans and mice. The domains were predicted via SMART

(http://smart.embl-heidelberg.de/). **(B)** The protein☐protein docking between IQCH

and CaM was predicted by the AlphaFold Protein Structure Database

(https://AlphaFold.ebi.ac.uk/) and LZerD website (https://lzerd.kiharalab.org/). **(C)**

Immunofluorescence staining of germ cells at different developmental stages

suggested the colocalization of CaM and IQCH in elongating spermatids in WT mice.

As expected, IQCH was not detected in the testes of the *Iqch* KO mice, and only CaM

was expressed in elongating spermatids in the *Iqch* KO mice (n = 3 biologically

independent WT mice and KO mice; scale bars, 50 μm). **(D)** qPCR analysis

demonstrated reduced *IQCH* and *CaM* expression levels in K562 cells following

transfection with *IQCH*-siRNA and *CaM*-siRNA, respectively, compared to those in

K562 cells transfected with the negative control siRNA. The downregulation of

HNRPAB resulted from decreased *IQCH* or *CaM* expression (Student's t test; *p <

0.05; error bars, s.e.m.). **(E)** qPCR analysis of K562 cells overexpressing *IQCH* with

simultaneous knockdown of *CaM* or overexpressing *CaM* with simultaneous

knockdown of *IQCH* revealed significant downregulation of *HNRPAB* (Student's t

test; *p < 0.05; error bars, s.e.m.). siRNA, small interfering RNA. Three independent

experiments were performed.

1114

A

III-2 Heterozygous mutation

III-3 Heterozygous mutation

II-1 Homozygous mutation

IV-1 Homozygous mutation

M=c.387+1_387+10del

WT allele

Exon 4   Intron 4

B

i Marker NC pSPL3 WT-*IQCH* Mut-*IQCH*
381bp
263bp

ii WT-*IQCH*
Exon V1   Exon4 deletion   Exon V2
Mut-*IQCH*
Exon V1 Exon V2

iii c.387+1_387+10del

V1 — Exon4 — V2

WT-*IQCH*   Mut-*IQCH*

V1 Exon4 V2   V1 V2
92bp 118bp 171bp   92bp 171bp

iv WT-c*IQCH* Mut-c*IQCH*
IB: IQCH
IB: GAPDH

C

Control   Proband

DAPI/α-Tubulin

DAPI/IQCH

Merge

Protein accession	Protein names	Gene names	Protein score
Q9D8E6	60S ribosomal protein L4	Rpl4	221
P47911	60S ribosomal protein L6	Rpl6	196
P14148	60S ribosomal protein L7	Rpl7	164
P47963	60S ribosomal protein L13	Rpl13	143
Q9CR57	60S ribosomal protein L14	Rpl14	109
P62908	40S ribosomal protein S3	Rps3	97
Q6ZWN5	40S ribosomal protein S9	Rps9	96
P35980	60S ribosomal protein L18	Rpl18	91
P27659	60S ribosomal protein L3	Rpl3	83
P62264	40S ribosomal protein S14	Rps14	83
P12970	60S ribosomal protein L7a	Rpl7a	73
P14131	40S ribosomal protein S16	Rps16	73
P19253	60S ribosomal protein L13a	Rpl13a	72
P62702	40S ribosomal protein S4, X isoform	Rps4x	66
P62301	40S ribosomal protein S13	Rps13	65
P47915	60S ribosomal protein L29	Rpl29	63
P14115	60S ribosomal protein L27a	Rpl27a	62
P61255	60S ribosomal protein L26	Rpl26	60
P84099	60S ribosomal protein L19	Rpl19	58
P25444	40S ribosomal protein S2	Rps2	53
P47962	60S ribosomal protein L5	Rpl5	53
Q9CPR4	60S ribosomal protein L17	Rpl17	51
P62754	40S ribosomal protein S6	Rps6	48
P14869	60S acidic ribosomal protein P0	Rplp0	47
P62751	60S ribosomal protein L23a	Rpl23a	47
P60867	40S ribosomal protein S20	Rps20	47
O09167	60S ribosomal protein L21	Rpl21	44
Q9CZM2	60S ribosomal protein L15	Rpl15	42
P35979	60S ribosomal protein L12	Rpl12	42
P97351	40S ribosomal protein S3a	Rps3a	36
P62855	40S ribosomal protein S26	Rps26	35
P62082	40S ribosomal protein S7	Rps7	34
P62849	40S ribosomal protein S24	Rps24	32

IQ motif

IQCH

CaM

HNRPAB

RNA binding proteins

Dnhd1   *Catsper2*

Catsper1   *Catsper3*

Ccdc40   *Ccdc39*

Ccdc65   *Irrc6*

RNAs

Acrosomes

Flagellar axoneme

Acrosome

Acroplaxome

Manchette

Mitochondria

Axoneme