IQCH regulates spermatogenesis by interacting with CaM to promote RNA-binding proteins' expression

Tiechao Ruan[1, 2]†, Ruixi Zhou[1, 2]†, Yihong Yang[3]†, Junchen Guo[4, 5], Chuan Jiang[1],

Xiang Wang[1], Gan Shen[1], Siyu Dai[1], Suren Chen[6*], Ying Shen[1*]

[1] Key Laboratory of Obstetrics, Gynecologic and Pediatric Diseases and Birth Defects of the Ministry of Education, Sichuan University, Chengdu, 610041, China;

[2] Department of Pediatrics, West China Second University Hospital, Sichuan University, Chengdu 610041 China;

[3] Reproduction Medical Center of West China Second University Hospital, Key Laboratory of Obstetric, Gynecologic and Pediatric Diseases and Birth Defects of Ministry of Education, Sichuan University, Chengdu 610041, China;

[4] Sichuan University-The Chinese University of Hong Kong (SCU-CUHK) Joint Laboratory for Reproductive Medicine, Key Laboratory of Obstetric, Gynaecologic and Paediatric Diseases and Birth Defects of Ministry of Education, West China Second University Hospital, Sichuan University, Chengdu 610041, China.

[5] Reproductive Endocrinology and Regulation Laboratory, Department of Obstetric and Gynaecologic, West China Second University Hospital, Sichuan University, Chengdu 610041, China.

[6] Education Key Laboratory of Cell Proliferation & Regulation Biology, College of Life Sciences, Beijing Normal University, Beijing 100875, China

†These authors contributed equally to this work

*Corresponding author:

Suren Chen

E-mail address: chensr@bnu.edu.cn;

Ying Shen

Phone: +86-15982083663.

E-mail address: yingcaishen01@163.com

Abstract

IQ motif-containing proteins can be recognized by calmodulin (CaM) and are

essential for many biological processes. However, the role of IQ motif-containing

proteins in spermatogenesis is largely unknown. In this study, we identified a

loss-of-function mutation in the novel gene IQ motif-containing H (*IQCH*) in a

Chinese family with male infertility, characterized by a cracked flagellar axoneme and

abnormal mitochondrial structure. To verify the function of IQCH, *Iqch*-knockout

mice were generated by CRISPR-Cas9 technology which reproduced the human

phenotypes. Mechanistically, IQCH can bind to CaM and then regulate the expression

of RNA-binding proteins (especially HNRPAB), which are indispensable for

spermatogenesis. Collectively, this study firstly unveiled the function of IQCH,

expanded the role of IQ motif-containing proteins in reproductive processes, and

provided important guidance for genetic counseling and gene diagnosis for male

infertility.

Introduction

Spermatogenesis is the most complex biological process in male organisms and functions to produce mature spermatozoa from spermatogonia in three phases: (i) spermatocytogenesis (mitosis), (ii) meiosis, and (iii) spermiogenesis (Hess and Renato de Franca 2008). This delicate process can be easily disturbed and further cause male reproductive disorders. Male infertility affects 7% of men in the general population, and its causes vary, ranging from anatomical or genetic abnormalities, systemic diseases, infections, trauma, iatrogenic injury, and gonadotoxins (Krausz et al. 2018). Approximately 10% of the human genome is related to reproductive processes; thus, male infertility is often predicted to be largely genetic in origin, whereas only 4% of infertile men are diagnosed with a distinct genetic cause (Krausz and Riera-Escamilla 2018). Genetic causes are highly heterologous and involve chromosomal abnormalities, point mutations in single genes, copy number variations, microRNA dysfunction, and polygenic defects (Meschede and Horst 1997, Traven et al. 2017). The highest percentage of known genetic factors that account for up to 25% of male infertility are karyotype anomalies, Y chromosome microdeletions, and *CFTR* mutations, which are mostly associated with azoospermia (Krausz and Riera-Escamilla 2018). However, in a relatively high proportion of infertile men (40%), the etiology cannot be recognized and is also referred to as idiopathic. With the help of assisted reproductive techniques (ART), some men have the chance to reproduce, however, there is a risk of passing on undetermined genetic abnormalities. In addition, genetic defects leading to fertilization failure and embryo development

arrest cannot be effectively rescued by ART; thus, discovering novel genetic factors

and further confirming their molecular mechanisms are of clinical importance.

Calcium (Ca) is an essential element that acts as a universal intracellular second

messenger. Ca is indispensable for many physiological processes in male reproduction,

including spermatogenesis, sperm movement, capacitation, hyperactivation, acrosome

reaction, chemotaxis, and fertilization (Valsa et al. 2015). Therefore, Ca deficiency is

mentioned to be a main contributor to male infertility (Beigi Harchegani et al. 2019).

Calmodulin (CaM) is defined as a major Ca sensor that activates many kinds of

enzymes in response to an increase in intracellular Ca^{2+} by interacting with a diverse

group of cellular proteins (Klee et al. 1980, Means et al. 1982). CaM binds to proteins

through recognition motifs, including the short CaM-binding motif containing

conserved Ile and Gln residues (IQ motif) for Ca^{2+}-independent binding and two

related motifs for Ca^{2+}-dependent binding (Nie et al. 2009). The identified IQ

motif-containing proteins include a good range of proteins with biological functions,

one of which is sperm surface proteins, suggesting that IQ motif-containing proteins

might play a potential role in male reproductive processes (Wen et al. 1999,

Dolmetsch et al. 2001). Moreover, it has been revealed that CaM is predominantly

expressed in the mammalian testis, from the pachytene to meiotic division stages

(Smoake et al. 1974, Kakiuchi et al. 1982, YAMAMOTO 1985, Sano et al. 1987, Kägi

et al. 1988, Moriya et al. 1995). However, limited IQ motif-containing proteins have

been reported to be responsible for male fertility. It has been shown that the IQ

motif-containing D (IQCD) is primarily accumulated in the acrosome area of

spermatids during mouse spermiogenesis, and the acrosome reaction was inhibited in

human spermatozoa by anti-IQCD antibody, suggesting a potential function of IQCD

in fertilization and the acrosome reaction (Zhang et al. 2019). Moreover, *Iqcf1*

knockout (KO) male mice showed significantly less fertility, which was related to

reduced sperm motility and acrosome reaction (Fang et al. 2015). Importantly, *Iqcg* is

required for mouse spermiogenesis, which is attributable to its role in flagellar

formation and/or function (Harris et al. 2014). Humans and mice without IQCN

presented failed fertilization rates related to manchette assembly defects (Dai et al.

2022). Although reliable findings provide substantial clues for IQ motif-containing

proteins participating in male reproductive processes, the relevant regulatory

mechanism and the function of many other IQ motif-containing proteins in

spermatogenesis have not been determined thus far.

Here, we revealed a novel IQ motif-containing protein, IQCH, which is essential

for spermiogenesis and fertilization. Disrupting IQCH leads to deficient acrosome

action and abnormal structure of the axoneme and mitochondria in both humans and

mice. Moreover, it is suggested that the interaction of IQCH and CaM is a prerequisite

for IQCH function, which further regulates the expression of RNA-binding proteins

(especially HNRPAB) during spermatogenesis. Collectively, our findings unveiled a

novel genetic diagnostic indicator of male infertility, and the uncovered mechanism of

IQCH in spermatogenesis might shed new light on the treatment of this disease.

Results

Identification of a novel IQ motif-containing protein, IQCH, involved in male fertility

A 33-year-old man from a consanguineous family with primary infertility for four years was recruited in our study (Fig. 1A). Semen analysis of this patient revealed an inordinately decreasing sperm motility and count, as well as abnormal sperm morphology (Table 1). In addition to the proband, his great uncle was also affected, he had never conceived during sexual intercourse without contraception in his marriage. We further explored the possible genetic cause through WES analysis, and a homozygous deletion variant in *IQCH* (c.387+1_387+10del) was identified in the proband. This variant is rare in the general human population according to the 1000 Genomes Project (0.007%), ExAC Browser (0.059%), and gnomAD databases (0.024%). We further clarified the putative contribution of this variant in this family by Sanger sequencing. Noticeably, this homozygous variant was affirmed in his infertile great-uncle, and the fertile parents of the proband carried the heterozygote of this variant (Fig. 1A).

Moreover, we used the minigene splicing assay to examine the effect of this variant on *IQCH* mRNA splicing. The electrophoresis results revealed that WT-*IQCH* yielded one transcript with a size of 381 bp, whereas the Mut-*IQCH* resulted in one strong band of 263 bp (Fig. 1B, i). Sanger sequencing further showed that the variant resulted in the absence of the whole exon 4, which was expected to translate into a truncated protein (Fig. 1B, ii and iii). Therefore, we further constructed a plasmid containing the aberrant cDNA sequence of *IQCH* caused by the splicing mutation to

verify the effect of the variant on protein expression. The western blotting results

showed that the Mut-c*IQCH* plasmid did not express IQCH while the WT-c*IQCH*

plasmid did (Fig. 1B, iv). We further conducted immunofluorescence staining of

spermatozoa from the proband and the control. We barely detected the expression of

IQCH protein in the spermatozoa from the proband compared to the control (Fig. 1C).

Taken together, these results reveal that this identified *IQCH* variant is responsible for

the splicing abnormality and further causes the lack of IQCH expression, which is

likely the genetic cause of male infertility in this family.

Nonfunctional IQCH leads to sperm with cracked axoneme structures

accompanied by defects in the acrosome and mitochondria

A comprehensive sperm morphology analysis was further conducted on the proband

(the semen was not available from his great-uncle). Papanicolaou staining revealed

that the spermatozoa displayed multiple flagellar morphological abnormalities, such

as coiled, bent, irregular, and even fractured tails (Fig. 2A). These morphological

anomalies were confirmed more precisely by SEM (Fig. 2B). Significantly, more

subtle abnormalities, such as axoneme exposure, bending, and cracking, were further

identified between the midpiece and the principal piece of the proband's spermatozoa

(Fig. 2B). As expected, a deranged or incomplete '9+2' microtubule structure of the

flagella were observed from the proband by TEM analysis (fig. S1, A). In addition, we

detected ultrastructural defects in the spermatozoa nucleus, including irregular shape,

large vacuoles, and deformed acrosomes (Fig. 2C). The mitochondria of the

spermatozoa had an abnormal arrangement and enlarged diameter (Fig. 2C).

Moreover, we performed immunofluorescence staining for the marker of the

acrosome (peanut agglutinin: PNA) as well as the mitochondrial marker

(Transcription Factor A, Mitochondrial: TFAM) to confirm the deficient of the

acrosomes and mitochondria in the proband's spermatozoa. The results of the PNA

and TFAM staining suggested that the spermatozoa acrosomes and mitochondria were

severely defective in the proband compared to the control (Fig. 2, D and E). In

addition, we performed SEPTIN4 (SEPT4) staining, a functional marker of the

annulus, to explore whether the flagellar fracture at the joint between the middle piece

and principal piece was caused by a nonfunctional annulus. The results showed that

there was no significant difference in the SEPT4 signal of the spermatozoa between

the proband and the control (fig. S1, B).

Impairment of male fertility in mice without *Iqch*

To further consolidate the role of IQCH in male reproduction, we first explored the

pattern of *Iqch* expression in mice by qPCR and found that *Iqch* was predominantly

expressed in the mouse testis compared to other organs (fig. S2, A). Additionally, we

investigated the temporal expression of *Iqch* in mouse testes on different postnatal

days. The results revealed that the expression of *Iqch* showed a significant increase on

postnatal Day 21, peaked on postnatal Day 35, and then presented stable expression

(fig. S2, A). To better understand the role of IQCH during spermatogenesis, we

performed immunofluorescence staining of the germ cells at different developmental

stages in human and mouse testes. The exact localization of IQCH shared the same

pattern in human and mouse spermatogenesis, being mainly detected in the cytoplasm

of spermatocytes and round spermatids and the flagella of late spermatids (fig. S2, B

and C). The testis-enriched expression of *Iqch*/IQCH suggested its potential role in

spermatogenesis.

We next generated *Iqch* knockout (KO) mice using clustered regularly interspaced

short palindromic repeat CRISPR-Cas9 technology. We used a guide RNA targeting

exons 2 through 3 of *Iqch* to achieve the knockout (fig. S3, A). PCR, RT-PCR, and

western blotting analysis further confirmed the success of the *Iqch* KO mouse

construction (fig. S3, B). The *Iqch* KO mice showed no overt abnormalities in their

development or behavior. The *Iqch* KO female mice showed normal fertility, and

hematoxylin-eosin (HE) staining further showed that the *Iqch* KO female mice had

normal follicle development compared to the WT mice (fig. S3, C-E). However, the

fertility of the KO male mice was significantly reduced compared to that of the WT

mice, including a reduction in the pregnancy rate and litter size (Fig. S3, F). There

was no detectable difference in the testis/body weight ratio between the WT and *Iqch*

KO mice (fig. S3, G), and the histology of testes and epididymides in the *Iqch* KO

mice showed no obvious abnormalities compared to the WT mice (fig. S4, A and B).

However, the results of the computer-assisted sperm analysis (CASA) affirmed

that sperm motility was significantly reduced in the KO mice, and the sperm count

was slightly decreased (Table 2, Movie S1 and S2). Moreover, morphological

anomalies of the sperm flagella were easily observed, such as an unmasking, bending,

or cracking axoneme, which recapitulated the flagellar phenotype of the infertile

patient (Fig. 3, A and B). We further found that during the spermatogenic process, the

abnormality of axoneme exposure had already occurred when the spermatozoa

flagellum developed (fig. S5, A). Intriguingly, tail defects, including bending and

cracking between the middle piece and the principal piece, were mainly present in the

epididymal spermatozoa (fig. S5, B), suggesting that more severe flagellum breakage

might occur during sperm movement.

We further performed TEM analysis to investigate the ultrastructural defects in

the testicular and epididymal spermatozoa of the *Iqch* KO mice. In contrast to the WT

mice, the spermatozoa of the *Iqch* KO mice showed dilated intermembrane spaces of

mitochondria and even loss of some mitochondrial material (Fig. 3C). However, the

annulus did not present significant differences between the *Iqch* KO and WT mice

(Fig. 3C), indicating that the disruptive link between the flagellar middle piece and

the principal piece might result from mitochondrial defects but not the annulus. The

immunofluorescence staining of the mitochondrial marker (Solute carrier family 25

member 4: SLC25A4) and SEPT4 also revealed that the mitochondria of the

spermatozoa of the *Iqch* KO mice were severely defective, and there was no

significant difference in the annulus of the spermatozoa between the *Iqch* KO mice

and WT mice, which was consistent with the proband (fig. S6, A and B).

Poor IVF outcomes in the *Iqch* KO male mice

To evaluate the cause of the damaged fertility of the *Iqch* KO male mice, mature

oocytes from WT female mice in their reproductive period were retrieved and placed

into culture dishes with sperm from the *Iqch* KO mice and WT mice for IVF treatment.

We found that 70% of the embryos had pronuclei in the WT group, while the

fertilization rate of the *Iqch* KO group was dramatically reduced, accounting for 47%

(Fig. 4A). Consequently, the rates of both two-cell embryos and blastocysts were

significantly lower in the *Iqch* KO male mice than in the WT male mice (Fig. 4A).

Noticeably, the inactive acrosome reaction was observed in most of the sperm from

the *Iqch* KO mice (Fig. 4B), leading to the inability to cross the zona pellucida (fig.

S7, A). Furthermore, abnormal acrosome development was evident in the testes of the

Iqch KO mice using PNA staining (fig. S7, B). Similarly, the expression of PNA was

abnormal in the mature sperm of the *Iqch* KO male mice (Fig. 4C). In the sperm and

testis of the *Iqch* KO male mice and the proband, the PLCζ fluorescence signal was

attenuated and abnormally localized (Fig. 4D, fig. S7, C and D). Taken together, these

data suggested that *Iqch* might play an important role in the acrosomal formation,

which is essential for fertilization.

RNA-binding proteins are the most relevant targets by which IQCH regulates

spermatogenesis

To elucidate the molecular mechanism by which IQCH regulates male fertility, we

performed LC-MS/MS analysis using mouse sperm lysates and detected 288

interactors of IQCH. Gene Ontology (GO) analysis of the IQCH-bound proteins

revealed a particular enrichment in fertilization, sperm axoneme assembly,

mitochondrial organization, calcium channel, and RNA processing (Fig. 5A).

Intriguingly, 33 ribosomal proteins were identified (Fig. 5B), indicating that IQCH

might be involved in protein synthesis. Using proteomic analysis on sperm from the

Iqch KO mice, we further assessed key proteins that might be activated by IQCH. A

total of 1,993 differential proteins were quantified, including 807 upregulated proteins

and 1,186 downregulated proteins (Fig. 5C). GO analysis revealed that the

significantly downregulated proteins were enriched in RNA processing, gene

expression, mitochondrion biogenesis, and calcium ion regulation (Fig. 5D), which

was consistent with the enrichment of the IQCH-bound proteins.

Importantly, the cross-analysis revealed that 76 proteins were shared between the

IQCH-bound proteins and the IQCH-activated proteins (Fig. 5E), implicating this

subset of genes as direct targets. Among the 76 proteins, 21 were RNA-binding

proteins (RBPs), 10 of which were suggested to be involved in

spermatogenesis(Kuroda et al. 2000, Yang et al. 2012, Chapman et al. 2013, Fukuda et

al. 2013, Legrand et al. 2019, Sechi et al. 2019, Liang et al. 2021, Tian and Petkov

2021, Wang et al. 2022, Xu et al. 2022); 8 were involved in mitochondrial function;

and 4 were calcium channel activity-related proteins (Fig. 5F). We focused on

SYNCRIP, HNRNPK, FUS, EWSR1, ANXA7, SLC25A4, and HNRPAB, the loss of

which showed the greatest influence on the phenotype of the *Iqch* KO mice.

Specifically, SYNCRIP, HNRNPK, FUS, EWSR1, and HNRPAB are RBPs that are

linked to spermatogenesis by controlling mRNA translation or spermatid post-meiotic

transcription (Kuroda et al. 2000, Fukuda et al. 2013, Sechi et al. 2019, Tian and

Petkov 2021, Xu et al. 2022). ANXA7 is a calcium-dependent phospholipid-binding

protein that is a negative regulator of mitochondrial apoptosis (Du et al. 2015). Loss

of SLC25A4 results in mitochondrial energy metabolism defects in mice (Graham et

al. 1997). We further confirmed the binding of IQCH between these proteins by the

Co-IP assay (Fig. 6A) and substantiated their downregulation in the sperm of the *Iqch*

KO mice by immunofluorescence staining and western blotting (Fig. 6B, fig. S8).

Among these interactors of IQCH, HNRPAB was the most significantly

downregulated protein seen by proteomic analysis (Data S1), implying that HNRPAB

might be the main target of IQCH. As an RBP, HNRPAB has been suggested to play

an important role in spermatogenesis by regulating the translation of testicular

mRNAs (Fukuda et al. 2013). We thus employed RNA-seq analysis using the sperm

from the *Iqch* KO and WT mice to investigate the effects of RNA levels when IQCH

is absent. Importantly, among the downregulated genes, most were related to male

fertility, such as axoneme assembly, spermatid differentiation, flagellated sperm

motility, and fertilization (fig. S9). We hypothesized that this downregulation is

linked to HNRPAB binding. Considering the disrupted fertilization and axoneme

assembly in the *Igch* KO mice, we chose the essential molecules involved in these

two processes, including *Catsper1*, *Catsper2*, *Catsper3*, *Ccdc40*, *Ccdc39*, *Ccdc65*,

Dnah8, *Irrc6,* and *Dnhd1*, to confirm our speculation. We carried out RNA

immunoprecipitation on formaldehyde cross-linked sperm followed by qPCR to

evaluate the interactions between HNRPAB and *Catsper1*, *Catsper2*, *Catsper3*,

Ccdc40, *Ccdc39*, *Ccdc65*, *Dnah8*, *Irrc6,* and *Dnhd1* in the KO and WT mice. As

expected, the binding between HNRPAB and those important molecules was detected

in the WT mice (Fig. 6C), supporting the functional role of HNRPAB in testicular

mRNAs. Intriguingly, significantly decreased interactions between HNRPAB and

those molecules were observed in the KO mice (Fig. 6C). Therefore, it is indicated

that IQCH is involved in spermatogenesis by mainly regulating RBPs, especially

HNRPAB, to further mediate essential mRNA expression in the testis.

The interaction of IQCH and CaM is a prerequisite for IQCH function

Given that IQCH is a calmodulin-binding protein, we hypothesized that IQCH

regulates these key molecules by interacting with CaM (fig. S10, A and B). As

expected, CaM interacted with IQCH, as indicated by LC-MS/MS analysis. We

initially confirmed the binding of IQCH and CaM in the WT sperm but not in the KO

sperm by the Co-IP assay (Fig. 6D). In addition, their colocalization was detected

during spermatogenesis by immunofluorescence staining (fig. S10, C).

To confirm that the interaction of IQCH and CaM is the prerequisite for regulating

HNRPAB expression, we downregulated IQCH or CaM in K562 cells, which express

both IQCH and CaM. As expected, reduced binding of IQCH and CaM was observed

in both cell lines with IQCH or CaM knocked down, with a concomitant decrease in

the expression of HNRPAB (Fig. 6E, fig. S10, D). We then overexpressed IQCH and

knocked down CaM or overexpressed CaM and knocked down IQCH in K562 cells to

check the change in HNRPAB levels. Consistently, the two situations also showed a

reduction in the expression of HNRPAB resulting from the diminished interaction

between IQCH and CaM (Fig. 6F, fig. S10, E).

We further confirmed that the IQ motif of IQCH is required for CaM binding

using a microscale thermophoresis (MST) assay. The binding affinity between IQCH

(target) from cell lysates overexpressing GFP-*IQCH* plasmids and recombinant CaM

(ligand) was enhanced with increasing concentrations of recombinant CaM (Fig. 6G).

However, their binding was disrupted when the IQ motif of IQCH was deleted (Fig.

6G). The Co-IP assay verified the above findings (Fig. 6H). Not surprisingly, the

expression of HNRPAB in cells co-transfected with *IQCH* (△IQ) and *CaM* plasmids

was lower than that in the cells overexpressing the WT-*IQCH* and *CaM* plasmids (Fig.

6H). Collectively, our findings suggest that IQCH interacts with CaM via IQ-motif to

manipulate the expression of important molecules, especially HNRPAB, to play a role

in spermatogenesis (Fig. 7).

Discussion

Throughout the evolution of divergent species, IQCH has been conserved, indicating

its fundamental role in organisms. As a novel CaM-binding protein, IQCH was first

identified in human and mouse testes and exclusively localized in spermatocytes and

spermatids, suggesting its potential activity in spermatogenesis (Yin et al. 2005).

However, since IQCH was identified, no other findings have supported its function in

male reproduction. In this study, we found that the mutation of *IQCH* impaired male

fertility, including flagellar morphological abnormalities and axoneme cracking.

Aberrant ultrastructure in sperm with the *IQCH* mutation was associated with

severely defective acrosomes and mitochondria. Furthermore, *Iqch* KO mice showed

similar irregularities in the flagellum, especially the axoneme and mitochondria. We

also identified that *Iqch* KO mice displayed different degrees of functional defects in

acrosomes. Thus, our work supported the vital role of IQCH in flagellar and acrosome

development.

To date, only four IQ motif-containing proteins, including IQCD, IQCF1, IQCG,

and IQCN, have been suggested to participate in spermatogenesis (Harris et al. 2014,

Fang et al. 2015, Zhang et al. 2019, Dai et al. 2022). Most IQ motif-containing

proteins function in Ca^{2+}dependent biological processes by binding to CaM (Chen et

al. 2014, Harris et al. 2014, Fang et al. 2015); thus, IQCD, IQCF1, and IQCN are

involved in fertilization and are relevant to the acrosome reaction, sperm capacitation,

or manchette assembly (Fang et al. 2015, Zhang et al. 2019, Dai et al. 2022). In

addition, because CaM has an impact on the actin cytoskeleton, *Iqcg* KO mice exhibit

a detachment of sperm heads from tails (Harris et al. 2014). In our study, the *Iqch* KO

mice also showed an impaired acrosome reaction, which caused reduced fertilization.

Intriguingly, the axoneme breaks at the annulus and the mitochondrial defects were

outstanding in the flagella of the *Iqch* KO mice, whose phenotypes were unexplored

in the previously discovered IQ motif-containing proteins. Because CaM activation

can stimulate actin cytoskeleton changes, it is reasonable that flagellum formation is

defective when IQCH is absent. Ca^{2+} is a key player in the regulation of

mitochondrial functions (Bravo-Sagua et al. 2017). Thus, the absence of IQCH leads

to a failed interaction with CaM and disrupts normal Ca^{2+} signaling, which

consequently causes mitochondrial defects in *Iqch* KO mice and patients with *IQCH*

variants. Our findings suggest that the fertilization function is the main action of IQ

motif-containing proteins, while the IQ motif-containing protein has its own role in

spermatogenesis.

However, there are few publications regarding the underlying mechanism of IQ

motif-containing proteins manipulating reproductive processes, except for a recent

study that stated that IQCN interacted with CaM to regulate the expression of

CaM-binding proteins, including manchette-related proteins (LZTFL1, KIF27, and

RSPH6A), IFT family proteins and their motor proteins (IFT22, IFT43, IFT74, IFT81,

IFT140, IFT172, WDR19, TTC21B, and DYNC2H1), and ribosomal protein family

proteins (RPS5, RPS25, RPS27, and RPSA) (Dai et al. 2022). In our study, it was

observed that IQCH regulates the expression of RBPs by their interactions. RBPs are

a large class of proteins that assemble with RNAs to form ribonucleoproteins (RNPs).

RBPs function at various stages of RNA processing, including alternative splicing,

RNA modification, nuclear export, localization, stability, and translation efficiency

(Corley et al. 2020). RNA processing in male reproductive biology is essential for the

production of mature sperm, and many RBPs have been demonstrated to be

indispensable during spermatogenesis (Morgan et al. 2021). Among the RBPs,

HNRPAB, a heterogeneous nuclear RNP, showed the most significant reduction

among the downregulated interactors of IQCH, suggesting that HNRPAB might be

the most important employee of IQCH. It has been reported that HNRPAB plays a

central role in spermatogenesis by regulating stage-specific translation of testicular

mRNAs (Fukuda et al. 2013). As expected, a RIP assay using an anti-HNRPAB

antibody revealed binding between HNRPAB and the mRNAs of several important

genes (*Catsper1*, *Catsper2*, *Catsper3*, *Ccdc40*, *Ccdc39*, *Ccdc65*, *Dnah8*, *Irrc6*, and

Dnhd1) involved in spermatogenesis. We further found that the interaction of IQCH

and CaM is a prerequisite for regulating HNRPAB expression. We thus hypothesized

that IQCH regulates male fertility by binding to CaM and controls HNRPAB to

manipulate the expression of key mRNAs involved in spermatogenesis.

In conclusion, our study identified a novel IQ motif-containing protein-coding

gene, *IQCH*/*Iqch*, which is responsible for spermatogenesis in humans and mice,

broadening the scope of known male infertility-associated genes thus further

illustrating the genetic landscape of this disease. In addition, our study suggested an

unexplored mechanism in which IQCH regulates the expression of key RBPs in

spermatogenesis, especially HNRPAB, by interacting with CaM to play crucial roles

in fertilization and axoneme assembly. We believe that our findings will provide

genetic diagnostic markers and potential therapeutic targets for infertile males with

IQCH pathogenic variants.

Materials and Methods.

Study participants

A male infertility family was recruited from West China Second University Hospital,

and healthy Chinese volunteers were enrolled as controls. The study was conducted

according to the tenets of the Declaration of Helsinki, and ethical approval was

obtained from the Ethical Review Board of West China Second University Hospital,

Sichuan University. Each subject signed an informed consent form.

WES and Sanger sequencing

Peripheral blood samples were obtained from all subjects, and genomic DNA was

extracted using a QIAamp DNA Blood Mini Kit (QIAGEN, Germany; 51126). The

exomes of the subject were captured by Agilent SureSelect Human All Exon V6

Enrichment kits (Agilent, CA, USA) and then sequenced on a HiSeq X-TEN system

(Illumina, CA, USA). All reads were mapped to the human reference sequence

(UCSC Genome Browser hg19) using Burrows-Wheeler Alignment. After quality

filtration with the Genome Analysis Toolkit, functional annotation was performed

using ANNOVAR through a series of databases, including the 1000 Genomes Project,

dbSNP, HGMD, and ExAC. PolyPhen-2, SIFT, MutationTaster, and CADD were used

for functional prediction. Variant verification of *IQCH* in patients was confirmed by

Sanger sequencing using the primers listed in table S1.

Minigene assay

The minigene splicing assay was used to explore the effect of the *IQCH* variant

(c.387+1_387+10del) on splicing. WT-*IQCH* and Mut-*IQCH* sequences, including

intron 3, exon 4, and intron 4, were PCR-amplified separately from the genomic DNA

of the control and patient. The two amplified fragments were cloned into the minigene

vector pSPL3 between the EcoRI and BamHI sites through the Basic Seamless

Cloning and Assembly Kit (TransGen Biotech, China, CU201-02). After transfection

into HEK293T cells by DNA and siRNA transfection reagent (Polypus, France,

101000046), the splicing patterns of the transcripts produced from the WT-*IQCH* and

Mut-*IQCH* plasmids were analyzed by RT-PCR, gel electrophoresis, and Sanger

sequencing. The primers used for the plasmid genomic amplification and the RT-PCR

are listed in table S1.

***Iqch* knockout mice**

The animal experiments were approved by the Experimental Animal Management and

Ethics Committee of West China Second University Hospital, Sichuan University. All

animal procedures complied with the Animal Care and Use Committee of Sichuan

University. The mouse *Iqch* gene has 6 transcripts and is located on chromosome 9.

According to the structure of the *Iqch* gene, exons 2~3 of the *Iqch-201*

(ENSMUST00000042322.10) transcript were recommended as the knockout region.

The region contains a 215 bp coding sequence, and deletion of this region was

expected to disrupt IQCH protein function. The sgRNA sequences, synthesized by

Sangon Biotech (Shanghai, China), are listed in table S2. The two complementary

DNA oligos of each sgRNA target were annealed and ligated to the pUC57-sgRNA

plasmid (Addgene, USA) for cloning. The recombinant plasmid was transformed into

DH5α competent cells, and the positive clone was screened based on kanamycin

resistance and sequencing. The recombinant plasmid was linearized and purified by

phenol-chloroform extraction. Transcriptions of the sgRNAs *in vitro* were performed

using the MEGAshortscript Kit (Ambion, USA, AM1354) and purified using the

MEGAclear Kit (Ambion, USA, AM1908). Cas9 mRNA was purchased from TriLink

BioTechnologies. The mouse zygotes were co-injected with the RNA mixture of Cas9

mRNA (~50 ng/µl) and sgRNA (~30 ng/µl). The injected zygotes were transferred

into pseudopregnant recipients to obtain the F0 generation. DNA was extracted from

the tail tissues of the 7-day-old offspring, and PCR amplification was carried out with

genotyping primers. A stable F1 generation (heterozygous mice) was obtained by

mating the positive F0 generation mice with wild-type C57BL/6JG-pt mice. Primers

used for the genotyping are listed in table S2.

Mouse fertility testing

To confirm the fertility of the *Iqch* KO male mice, natural mating tests were

conducted. Briefly, three *Iqch* KO and three littermate control sexually mature male

mice (8-12 weeks old) were paired with two 6- to 8-week-old normal C57BL/6J

females (each male was mated with two female mice) for 6 months. The vaginal plugs

of the mice were examined every morning. Female mice with vaginal plugs were fed

separately, and the number of pups per litter was recorded.

In vitro fertilization

Eight-week-old C57BL/6J female mice were superovulated by injecting 5 IU of

pregnant mare serum gonadotropin (PMSG), followed by 5 IU of human chorionic

gonadotropin (hCG) 48 h later. Sperm was released from the cauda epididymis of

10-week-old male mice, and sperm capacitation was performed for 50 min using a

TYH solution. The cumulus-oocyte complexes (COCs) were obtained from the

ampulla of the uterine tube 14 h after the hCG injection. The ampulla was torn with a

syringe needle, and the COCs were gently squeezed onto liquid drops of an HTF

medium. The COCs were then incubated with ~5 μl of the sperm suspension in HTF

liquid drops at 37 °C under 5% CO_2. After 6 h, the eggs were washed several times

using an HTF medium to remove the cumulus cells and then transferred to liquid

drops of a KSOM medium.

Acrosome reaction analysis

The spermatozoa were collected from the cauda epididymis and capacitated for 50

min in a TYH medium at 37 °C under 5% CO_2. Highly motile sperm were collected

from the upper portion of the medium, and 10 μM of calcium ionophore A23187

(Sigma-Aldrich, CA, USA, C7522) was added to induce the acrosome reaction. After

15 min, the spermatozoa were spotted on a glass microscope slide, dried, and fixed

with 4% PFA for 10 min. The acrosomes were stained with Coomassie brilliant blue

staining.

Hematoxylin-eosin (H&E) staining

For the staining of the testicular tissues, samples were dissected from the adult mice

and fixed in 4% PFA overnight at 4 °C. The fixed tissues were embedded in paraffin,

sectioned (5 μm thick), dewaxed, and rehydrated. The sections of the testis were

stained with hematoxylin and an eosin solution (Beyotime, Shanghai, China, C0105M)

before imaging using a microscope (Leica, Germany).

Immunofluorescence staining

For sperm immunostaining, fresh sperm samples were fixed with 4% PFA for 20 min

at room temperature. For the staining of the testis tissues, heat-induced antigen

retrieval was performed in a citrate antigen retrieval solution (Beyotime, Shanghai,

China, P0081) according to the manufacturer's requirements. After permeabilization

with 1% Triton X-100 for 30 min, the sperm slides were blocked with 5% bovine

serum albumin serum for 1 h at room temperature. Primary antibodies were added to

the slide and incubated overnight at 4 °C. After washing three times with PBS, the

slides were incubated with secondary antibodies for 1 h at room temperature. The

nuclei were counterstained with DAPI dye (Sigma☐Aldrich, CA, USA, 28718-90-3)

and mounted with an antifade mounting medium. Images were captured with a

laser-scanning confocal microscope (Olympus, Japan). Detailed information on the

antibodies is provided in table S3.

qPCR

Total RNA was extracted from the mouse testes using an RNA Easy Fast Tissue/Cell

Kit (Tiangen Biotech, China, 4992732). Approximately 0.3 mg of total RNA was

converted into cDNA with the PrimeScript RT Reagent Kit (Takara, Japan, RR037A)

according to the manufacturer's instructions. The cDNAs were individually diluted

10-fold to be used as templates for the subsequent real-time fluorescence quantitative

PCR (qPCR) with iTaq Universal SYBR Green Supermix (Bio-Rad Laboratories,

USA, 1725124). Mouse *Gapdh* was used as an internal control. The mRNA

expression of detected targets was quantified according to the $2^{-\Delta\Delta Ct}$ method. The

primers used for the qPCR are listed in table S1 and table S2.

Papanicolaou staining

Semen samples were fixed in 4% paraformaldehyde, and the coated slides were

air-dried before being rehydrated with 80%, 70%, and 50% ethanol and distilled water.

The samples were then stained with Lea's hematoxylin, rinsed with distilled water,

and stained with G-6 orange stain and EA-50. Following the staining, the slides were

dehydrated with ethanol and mounted.

Scanning electron microscopy (SEM)

The spermatozoa were fixed in 2.5% phosphate-buffered glutaraldehyde (GA)

(Zhongjingkeyi Technology, Beijing, China) at room temperature for 30 min and then

deposited on coverslips. The coverslips were dehydrated via an ascending gradient of

50%, 70%, 95%, and 100% ethanol and air-dried. The specimens were then attached

to specimen holders and coated with gold particles using an ion sputter coater before

being viewed with a JSM-IT300 scanning electron microscope (JEOL, Tokyo, Japan).

Transmission electron microscopy (TEM)

Precipitation of the spermatozoa was fixed with 2.5% (vol/vol) glutaraldehyde in a 0.1

M phosphate buffer (PB) (pH 7.4), washed two times in PB, and two times in ddH$_2$O.

Then, the tissues were immersed in 1% (wt/vol) OsO$_4$ and 1.5% (wt/vol) potassium

ferricyanide aqueous solution at 4 °C for 2 h. After washing, the samples were

dehydrated through graded alcohol (30%, 50%, 70%, 80%, 90%, 100%, 10 min each)

in pure acetone (10 min twice). The samples were then infiltrated in a graded mixture

(3:1, 1:1, 1:3) of acetone and SPI-PON812 resin (21 ml SPO-PON812, 13 ml DDSA

and 11 ml NMA), and then the pure resin was changed. The specimens were

embedded in pure resin with 1.5% BDMA and polymerized for 12 h at 45 °C and 48 h

at 60 °C. The ultrathin sections (70 nm thick) were sectioned with a microtome (Leica

EM UC6, Germany), double-stained with uranyl acetate and lead citrate, and

examined by a transmission electron microscope (TECNAI G2 F20, Philips, USA).

Proteomic quantification

Proteins were extracted from the sperm samples using a RIPA lysis buffer (Applygen,

Beijing, China, C1053) containing 1 mM PMSF and protease inhibitors on ice. The

supernatants were collected following centrifugation at 12,000 × g for 20 min. Protein

concentrations were calculated by Bradford quantification and SDS-PAGE. An

enzyme solution at a ratio of 1:20 of trypsin enzyme (μg) to substrate protein (μg) was

added to 100 μg of the protein samples, vortexed, centrifuged at low speed for 1

minute, and incubated at 37 °C for 4 h. The peptide liquid obtained after salt removal

was freeze-dried. The peptide sample was dissolved in 0.5 M TEAB and added to the

corresponding iTRAQ labeling reagent and stored at room temperature for 2 h. The

Shimadzu LC-20AB liquid phase system was used for purification, and the separation

column was a 5 μm 4.6 × 250 mm Gemini C18 column for liquid phase separation of

the sample. The dried peptide samples were reconstituted with mobile phase A (2%

can, 0.1% FA) and centrifuged at 20,000 × g for 10 min, and the supernatant was

taken for injection. The separation was performed by UltiMate 3000 UHPLC (Thermo

Fisher). The sample was first enriched in a trap column and desalted and then entered

into a self-packed C18 column. The peptides separated by the liquid phase

chromatography were ionized by a nanoESI source and then passed to a tandem mass

spectrometer Q-Exactive HF X (Thermo Fisher) for DDA (Data Dependent

Acquisition) mode detection. The raw data were converted to mgf files for

bioinformatics analysis, and protein identification from tandem mass spectra was

performed by database searching (UniProt). The protein quantification process

includes the following steps: protein identification, tag impurity correction, data

normalization, missing value imputation, protein ratio calculation, statistical analysis,

and results presentation. Proteins with a 1.5-fold change and a p-value (using

Student's *t*-test) less than 0.05 were defined as differentially expressed proteins.

RNA sequencing

Total RNA (~1 μg) was isolated from fresh spermatozoa from the WT mice (n=3) and

the *Iqch* KO mice (n=3) using the RNAsimple Total RNA Kit (Tiangen Biotech,

China, 4992858). A Ribo-Zero™ rRNA Removal Kit (MRZPL1224, llumina) was

used to remove rRNA from the samples. RNA integrity was evaluated using the

Agilent 2100 Bioanalyzer system (Agilent Technologies). An mRNA sequencing

library for the RNA-seq was constructed using the TruSeq RNA Library Prep kit v2

(RS-122-2001, Illumina), followed by paired-end (2 × 100 bp) sequencing using

the Illumina HiSeq 4000 Sequencing System (Illumina) at the Beijing Genomics

Institute. Raw paired-end reads were filtered by FASTX-Toolkit. The quality of the

reads was confirmed by FastQC (ver: 0.11.3). The raw data (raw reads) in the fastq

format were processed through in-house Perl scripts. Feature Counts (ver: l.5.0-p3)

were used to count the read numbers mapped to each gene. The FPKM of each gene

was calculated based on the length of the gene and the read count mapped to the gene.

Signaling pathway matching analysis for the differentially expressed gene (DEG) list

was performed using KEGG Mapper in the Kyoto Encyclopedia of Genes and

Genomes (KEGG, https://www.genome.jp/kegg/). The mapping of GO to DEGs was

carried out using Blast2GO (ver: 4.1.9).

Western blotting and Co-IP

Proteins were extracted from the cultured cells and mouse testes using a RIPA lysis

buffer containing 1 mM PMSF and protease inhibitors (Applygen, Beijing, China,

P1265) on ice. The supernatants were collected following centrifugation at 12,000 × g

for 20 min. The proteins were electrophoresed in 10% SDS-PAGE gels and

transferred to nitrocellulose membranes (GE Healthcare). The blots were blocked in 5%

milk and incubated with primary antibodies overnight at 4 °C, followed by incubation

with anti-rabbit or anti-mouse IgG H&L (HRP) (Abmart, Shanghai, China, M21002

and M21001) at a 1/10,000 dilution for 1 h. The signals were evaluated using a Super

ECL Plus Western Blotting Substrate (Applygen, Beijing, China, P1050) and a

Tanon-5200 Multi chemiluminescence imaging system (Tanon, Shanghai, China).

For the Co-IP assays, the extracted proteins were incubated with primary

antibodies overnight at 4 °C. The lysates were then incubated with 20 μl of Pierce™

Protein A/G-conjugated Agarose for 2 h at room temperature. The beads were washed

with a washing buffer [50 mM Tris-HCl (pH 7.4), 0.1% Triton X-100, and 500 mM

NaCl], eluted with a 1.2×SDS loading buffer, and boiled for 5 min at 95 °C. Finally,

the products were separated using SDS-PAGE and analyzed using immunoblotting

procedures. Detailed information on the antibodies used in the western blotting

experiments is provided in table S3.

Antibody production

The peptide used for raising the anti-IQCH antibody was derived from amino acid

residues 406–435 (KAEAATKIQATWKSYKARSSFISYRQKKWA) of mouse

IQCH. The peptide coupled with keyhole limpet hemocyanin (KLH) (Sigma-Aldrich,

USA, H7017) was dissolved in saline, emulsified with 1 ml of Freund's complete

adjuvant (Beyotime, China, P2036), and injected at multiple sites on the back of New

Zealand white rabbits. The antiserum was collected within 2 weeks after the final

injection.

Cell culture and transfection

We purchased the K562 cells (CRL-3344) and the HEK293T cells (CRL-11268) from

the American Type Culture Collection. The K562 cells were cultured in Basic

Roswell Park Memorial Institute 1640 Medium (Gibco, USA, C11875500BT)

supplemented with 10% fetal bovine serum (Gibco, USA, 12483020). The HEK293T

cells were cultured in Dulbecco's Modified Eagle Medium (Gibco, USA, 11965092)

supplemented with 10% fetal bovine serum (Gibco, USA, 12483020). The expression

plasmids, including pcDNA3.1-Flag-*CALM2*, pCMV-MCS-3* flag-WT-*IQCH,* and

pCMV-MCS-3* flag- *IQCH(*△IQ*)*, and small interfering RNAs (siRNAs) of *CALM2*

and *IQCH* were constructed by Vigene Biosciences (Jinan, China). The target

sequences of siRNA are listed in table S1. The plasmids and siRNA were transfected

into cells with a jetPRIME transfection reagent (Polypus, France, 101000046)

according to the manufacturer's protocol.

Microscale thermophoresis (MST) assay.

The MST experiments were conducted on a Monolith NT.115 system (NanoTemper

Technologies, Germany). Lysates of the HEK293T cells transfected with fluorescent

GFP-*IQCH* or GFP-*IQCH*(△IQ) were normalized by raw fluorescence (count), diluted

using an MST buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 10 mM MgCl2, 0.05%

(v/v) Tween 20), and added to 16 PCR tubes (10 μl per tube). Then, the purified CaM

was diluted using an MST buffer into 16 gradients. Ten microliters of different

concentrations of CaM were mixed with 10 μl of fluorescent GFP-IQCH protein and

GFP-*IQCH*(△IQ) and reacted in a dark box for 15 min at room temperature. The

samples were added to monolith capillaries (NanoTemper, MO-L022) and

subsequently subjected to MST analysis. The measurement protocol times were as

follows: fluorescence before 5 sec, MST on 30 sec, fluorescence after 5 sec, delay 25

sec. The dissociation constant (Kd) was determined using a single-site model to fit the

curve.

Statistical analysis

Data were compared for statistical significance using GraphPad Prism version

9.0.0 (GraphPad Software). The unpaired, two-tailed Student's t-test was used for the

statistical analyses. The data are presented as the mean ± SEM, and statistically

significant differences are represented as *p<0.05.

References

Beigi Harchegani, A., A. Irandoost, M. Mirnamniha, H. Rahmani, E. Tahmasbpour and A.

Shahriary (2019). "Possible Mechanisms for The Effects of Calcium Deficiency on Male

Infertility." Int J Fertil Steril **12**(4): 267-272.

Bravo-Sagua, R., V. Parra, C. López-Crisosto, P. Díaz, A. F. Quest and S. Lavandero (2017).

"Calcium Transport and Signaling in Mitochondria." Compr Physiol **7**(2): 623-634.

Chapman, K. M., H. M. Powell, J. Chaudhary, J. M. Shelton, J. A. Richardson, T. E.

Richardson and F. K. Hamra (2013). "Linking spermatid ribonucleic acid (RNA) binding protein

and retrogene diversity to reproductive success." Mol Cell Proteomics **12**(11): 3221-3236.

Chen, L. T., W. X. Liang, S. Chen, R. K. Li, J. L. Tan, P. F. Xu, L. F. Luo, L. Wang, S. H. Yu, G.

Meng, K. K. Li, T. X. Liu, Z. Chen and S. J. Chen (2014). "Functional and molecular features of

the calmodulin-interacting protein IQCG required for haematopoiesis in zebrafish." Nat

Commun **5**: 3811.

Corley, M., M. C. Burns and G. W. Yeo (2020). "How RNA-Binding Proteins Interact with RNA:

Molecules and Mechanisms." Mol Cell **78**(1): 9-29.

Dai, J., Q. Li, Q. Zhou, S. Zhang, J. Chen, Y. Wang, J. Guo, Y. Gu, F. Gong, Y. Tan, G. Lu, W.

Zheng and G. Lin (2022). "IQCN disruption causes fertilization failure and male infertility due to

manchette assembly defect." EMBO Mol Med **14**(12): e16501.

Dolmetsch, R. E., U. Pajvani, K. Fife, J. M. Spotts and M. E. Greenberg (2001). "Signaling to

the nucleus by an L-type calcium channel-calmodulin complex through the MAP kinase

pathway." Science **294**(5541): 333-339.

Du, Y., Y. Huang, Y. Gao, B. Song, J. Mao, L. Chen, L. Bai and J. Tang (2015). "Annexin A7

674 modulates BAG4 and BAG4-binding proteins in mitochondrial apoptosis." Biomed

675 Pharmacother **74**: 30-34.

676 Fang, P., W. Xu, D. Li, X. Zhao, J. Dai, Z. Wang, X. Yan, M. Qin, Y. Zhang, C. Xu, L. Wang and

677 Z. Qiao (2015). "A novel acrosomal protein, IQCF1, involved in sperm capacitation and the

678 acrosome reaction." Andrology **3**(2): 332-344.

679 Fukuda, N., T. Fukuda, J. Sinnamon, A. Hernandez-Hernandez, M. Izadi, C. S. Raju, K.

680 Czaplinski and P. Percipalle (2013). "The transacting factor CBF-A/Hnrnpab binds to the

681 A2RE/RTS element of protamine 2 mRNA and contributes to its translational regulation during

682 mouse spermatogenesis." PLoS Genet **9**(10): e1003858.

683 Graham, B. H., K. G. Waymire, B. Cottrell, I. A. Trounce, G. R. MacGregor and D. C. Wallace

684 (1997). "A mouse model for mitochondrial myopathy and cardiomyopathy resulting from a

685 deficiency in the heart/muscle isoform of the adenine nucleotide translocator." Nat Genet **16**(3):

686 226-234.

687 Harris, T. P., K. J. Schimenti, R. J. Munroe and J. C. Schimenti (2014). "IQ motif-containing G

688 (Iqcg) is required for mouse spermiogenesis." G3 (Bethesda) **4**(2): 367-372.

689 Hess, R. A. and L. Renato de Franca (2008). "Spermatogenesis and cycle of the seminiferous

690 epithelium." Adv Exp Med Biol **636**: 1-15.

691 Kägi, U., J. G. Chafouleas, A. W. Norman and C. W. Heizmann (1988). "Developmental

692 appearance of the Ca2+-binding proteins parvalbumin, calbindin D-28K, S-100 proteins and

693 calmodulin during testicular development in the rat." Cell Tissue Res **252**(2): 359-365.

694 Kakiuchi, S., S. Yasuda, R. Yamazaki, Y. Teshima, K. Kanda, R. Kakiuchi and K. Sobue

695 (1982). "Quantitative determinations of calmodulin in the supernatant and particulate fractions

of mammalian tissues." <u>J Biochem</u> **92**(4): 1041-1048.

Klee, C. B., T. H. Crouch and P. G. Richman (1980). "Calmodulin." <u>Annu Rev Biochem</u> **49**:

489-515.

Krausz, C., F. Cioppi and A. Riera-Escamilla (2018). "Testing for genetic contributions to

infertility: potential clinical impact." <u>Expert Rev Mol Diagn</u> **18**(4): 331-346.

Krausz, C. and A. Riera-Escamilla (2018). "Genetics of male infertility." <u>Nat Rev Urol</u> **15**(6):

369-384.

Kuroda, M., J. Sok, L. Webb, H. Baechtold, F. Urano, Y. Yin, P. Chung, D. G. de Rooij, A.

Akhmedov, T. Ashley and D. Ron (2000). "Male sterility and enhanced radiation sensitivity in

TLS(-/-) mice." <u>Embo j</u> **19**(3): 453-462.

Legrand, J. M. D., A. L. Chan, H. M. La, F. J. Rossello, M. L. Änkö, F. V. Fuller-Pace and R. M.

Hobbs (2019). "DDX5 plays essential transcriptional and post-transcriptional roles in the

maintenance and function of spermatogonia." <u>Nat Commun</u> **10**(1): 2278.

Liang, J., Y. Zheng, W. Zeng, L. Chen, S. Yang, P. Du, Y. Wang, X. Yu and X. Zhang (2021).

"Comparison of proteomic profiles from the testicular tissue of males with impaired and normal

spermatogenesis." <u>Syst Biol Reprod Med</u> **67**(2): 127-136.

Means, A. R., J. S. Tash and J. G. Chafouleas (1982). "Physiological implications of the

presence, distribution, and regulation of calmodulin in eukaryotic cells." <u>Physiol Rev</u> **62**(1):

1-39.

Meschede, D. and J. Horst (1997). "The molecular genetics of male infertility." <u>Mol Hum</u>

<u>Reprod</u> **3**(5): 419-430.

Morgan, M., L. Kumar, Y. Li and M. Baptissart (2021). "Post-transcriptional regulation in

spermatogenesis: all RNA pathways lead to healthy sperm." Cell Mol Life Sci **78**(24):

8049-8071.

Moriya, M., K. Fujinaga, M. Yazawa and C. Katagiri (1995). "Immunohistochemical localization

of the calcium/calmodulin-dependent protein phosphatase, calcineurin, in the mouse testis: its

unique accumulation in spermatid nuclei." Cell Tissue Res **281**(2): 273-281.

Nie, D., X. Yang and Z. Yankai (2009). "Molecular cloning and expression profile analysis of a

novel mouse testis-specific expression gene mtIQ1." Mol Biol Rep **36**(5): 1203-1209.

Sano, M., A. S. Ohshima, N. Kawamura, S. Kitajima and A. Mizutani (1987).

"Immunohistochemical study of calmodulin in developing mouse testis." J Exp Zool **241**(1):

51-59.

Sechi, S., A. Frappaolo, A. Karimpour-Ghahnavieh, M. Gottardo, R. Burla, L. Di Francesco, E.

Szafer-Glusman, E. Schininà, M. T. Fuller, I. Saggio, M. G. Riparbelli, G. Callaini and M. G.

Giansanti (2019). "Drosophila Doublefault protein coordinates multiple events during male

meiosis by controlling mRNA translation." Development **146**(22).

Smoake, J. A., S. Y. Song and W. Y. Cheung (1974). "Cyclic 3',5'-nucleotide

phosphodiesterase. Distribution and developmental changes of the enzyme and its protein

activator in mammalian tissues and cells." Biochim Biophys Acta **341**(2): 402-411.

Tian, H. and P. M. Petkov (2021). "Mouse EWSR1 is crucial for spermatid post-meiotic

transcription and spermiogenesis." Development **148**(11).

Traven, E., A. Ogrinc and T. Kunej (2017). "Initiative for standardization of reporting genetics

of male infertility." Syst Biol Reprod Med **63**(1): 58-66.

Valsa, J., K. P. Skandhan, P. S. Khan, K. P. Avni, S. Amith and M. Gondalia (2015). "Calcium

and magnesium in male reproductive system and in its secretion. I. Level in normal human

semen, seminal plasma and spermatozoa." Urologia **82**(3): 174-178.

Wang, X. L., J. M. Li and S. Q. Yuan (2022). "Characterization of the protein expression and

localization of hnRNP family members during murine spermatogenesis." Asian J Androl.

Wen, Y., R. T. Richardson and G. O'Rand M (1999). "Processing of the sperm protein Sp17

during the acrosome reaction and characterization as a calmodulin binding protein." Dev Biol

206(2): 113-122.

Xu, H., J. Guo, W. Wu, Q. Han, Y. Huang, Y. Wang, C. Li, X. Cheng, P. Zhang and Y. Xu

(2022). "Deletion of Hnrnpk Gene Causes Infertility in Male Mice by Disrupting

Spermatogenesis." Cells **11**(8).

YAMAMOTO, N. J. A. h. e. c. (1985). "Immunoelectron microscopic localization of calmodulin

in guinea pig testis and spermatozoa."   **18**(2): 199-211.

Yang, F., Q. Wei, R. S. Adelstein and P. J. Wang (2012). "Non-muscle myosin IIB is essential

for cytokinesis during male meiotic cell divisions." Dev Biol **369**(2): 356-361.

Yin, L. L., J. M. Li, Z. M. Zhou and J. H. Sha (2005). "Identification of a novel testis-specific

gene and its potential roles in testis development/spermatogenesis." Asian J Androl **7**(2):

127-137.

Zhang, P., W. Jiang, N. Luo, W. Zhu and L. Fan (2019). "IQ motif containing D (IQCD), a new

acrosomal protein involved in the acrosome reaction and fertilisation." Reprod Fertil Dev **31**(5):

898-914.

Acknowledgments: We thank the patient and the family members for their support

during this research study.

Competing interests: The authors declare that they have no competing interests.

Author contributions: Y.S. designed and supervised the study experiments. Y.Y.

collected data and conducted the clinical evaluations. T.R., R.Z., and Y.S. performed

experiments and analyzed most of the data. S.C. generated the CRISPR mice. J.G.,

C.J., X.W., G.S. and S.D. performed cell experiments. Y.S. wrote the manuscript with

input from others. All authors revised and approved the article.

Ethics Approval and Consent to Participate: For human participants, this study was

performed in line with the principles of the Declaration of Helsinki. Approval was

granted by Ethics Committee of the Second West China Hospital of Sichuan

University. Informed consent was obtained from all individual participants included in

the study. For the animal experiments, they were performed in accordance with the

recommendation of the Guide for the Care and Use of Laboratory Animals of the

National Institutes of Health and the animal experiments were approved by the

Experimental Animal Management and Ethics Committee of West China Second

University Hospital (IACUC no. 2021(070)).

Funding: This study was funded by the National Key Research and Development

Project (2019YFA0802101)

Data and materials availability: All data needed to evaluate the conclusions in the

paper are present in the paper and/or the Supplementary Materials.

Figure legends

Fig. 1. Identification of a splicing mutation in *IQCH* in a consanguineous family

with male infertility. (A) Pedigree analysis of the consanguineous family with two

infertile males (II-1 and IV-1), with the black arrow pointing to the proband. The

affected males exhibited the homozygous variant in *IQCH* by Sanger sequencing.

Sequence chromatograms are shown below the pedigree. **(B)** (i) The electrophoresis

results of the minigene assay show a decrease in the molecular weight of the RT-PCR

products generated from the Mut-*IQCH* (263 bp) compared with the WT-*IQCH* (381

bp). (ii) Sanger sequencing of the complementary DNA of the splicing mutation

showing the deletion of exon 4 in *IQCH*. (iii) The pattern diagram demonstrating the

splicing effects caused by the *IQCH* mutation. (iv) Western blotting results showed

that the Mut–c*IQCH* plasmids did not express IQCH. NC, negative control. Three

independent experiments were performed. **(C)** Immunofluorescence staining showed

that the expression of IQCH was barely detected in the proband's sperm compared

with the control (blue, DAPI; green, α-Tubulin; red, IQCH; scale bars, 5 μm).

Fig. 2. Abnormal flagellar morphology and defective acrosomes and

mitochondria in the infertile patient. (A and **B)** The Papanicolaou staining (A) and

scanning electron microscopy (SEM) (B) results showed flagellar morphological

abnormalities (scale bars in A, 5 μm; scale bars in B, 2.5 μm). The dotted boxes and

red arrowheads denote the axoneme cracking and exposure. **(C)** Transmission electron

microscopy (TEM) results showing the deformed acrosomes and abnormal

arrangement and diameter mitochondria (scale bars, 500 nm). The red arrowheads

denote the abnormal mitochondria. N, nucleus; M, mitochondria. **(D** and **E)** Defects

of the acrosome and mitochondria were observed in the proband's sperm by PNA (D)

and TFAM (E) staining (blue, DAPI; green, α-Tubulin; red, PNA or TFAM; scale

bars, 5 μm). PNA, peanut agglutinin; TFAM, Transcription Factor A, Mitochondrial.

Fig. 3. The absence of *Iqch* impaired spermatogenesis in mice. (A and **B)** The

Papanicolaou staining (B) and SEM (C) results showing unmasking, bending, or

cracking axoneme in the spermatozoa from the *Iqch* KO mice (n = 3 biologically

independent WT mice and KO mice; scale bars in B, 5 μm; scale bars in C, 2.5 μm).

The red arrowheads point to the axoneme abnormalities. **(C)** TEM revealing dilated

intermembrane spaces of mitochondria and a normal annulus in the testicular and

epididymal spermatozoa from the *Iqch* KO mice (n = 3 biologically independent WT

mice and KO mice; scale bars, 500 nm). The red arrowheads point to the dilated

intermembrane spaces of mitochondria. The green arrowheads point to the normal

annulus. M, mitochondria.

Fig. 4. Poor IVF treatment outcomes from using sperm from the *Iqch* KO mice.

(A) Representative pronucleus embryos, two-cell embryos, and blastocysts obtained

from the WT mice and the *Iqch* KO mice. Significantly reduced fertilization rates,

cleavage rates, and blastocyst formation rates were observed in the *Iqch* KO mice

compared to the WT mice (n = 3 biologically independent WT mice and KO mice;

scale bars, 100 μm; Student's t test; *p < 0.05; error bars, s.e.m). **(B)** The acrosome

reaction rates in the capacitated spermatozoa from the WT mice and *Iqch* KO mice

were determined by Coomassie brilliant blue staining. The acrosome reaction rates

were reduced in the spermatozoa from the *Iqch* KO mice (n = 3 biologically

independent WT mice and KO mice; scale bars, 5 μm; Student's t-test; *p < 0.05;

error bars, s.e.m). The green arrowheads indicate the reacted acrosomes. The red

arrowheads indicate intact acrosomes. (**C** and **D**) PNA (C) and PLCζ (D) staining

showing the abnormal acrosome morphology and aberrant PLCζ location and

expression in the spermatozoa from the *Iqch* KO mice (n = 3 biologically independent

WT mice and KO mice; scale bars, scale bars, 5 μm). The dotted box indicates the

typical pattern of the PLCζ location and expression in the spermatozoa from the WT

mice. PNA, peanut agglutinin; PLCζ, phospholipase C zeta 1.

Fig. 5. IQCH bound and activated male reproduction-related proteins in the

sperm of the mice. **(A)** Bubble plots of the GO analysis showing that the

IQCH-interacting proteins are significantly enriched in spermatogenesis and RNA

processing in three categories: biological process, cellular component, and molecular

function. GO, gene ontology. **(B)** Thirty-three ribosomal proteins interacted with

IQCH in the sperm from the mice. **(C)** A heatmap showing the differential protein

results from the proteomic analysis of the sperm from the WT and *Iqch* KO mice. **(D)**

Bubble plots showing the decreased enrichment of proteins related to the

spermatogenetic process and RNA processing according to the GO analysis of the

849 *Iqch* KO mice compared to the WT mice. **(E)** Venn diagram depicting the 76

850 overlapping proteins between the 1186 downregulated proteins in the sperm from the

851 *Iqch* KO mice and the 288 proteins binding with IQCH. **(F)** Chord diagram showing

852 21 proteins involved in RNA binding, 8 proteins involved in mitochondrial function,

853 and 4 proteins involved in calcium channel activity among the 76 overlapping

854 proteins. HNRPAB showed the greatest reduction of expression in the *Iqch* KO mice.

856 **Fig. 6. IQCH interacted with CaM to regulate the expression of HNRPAB and**

857 **spermatogenesis. (A)** A Co-IP assay of the mouse sperm lysates revealed that the

858 seven proteins most relevant to the phenotype of the *Iqch* KO mice bound with IQCH.

859 **(B)** Western blotting showing the reduced expression of the seven proteins in the

860 sperm from the *Iqch* KO mice compared to the WT mice. **(C)** qPCR analysis of RNA

861 immunoprecipitation (RIP) using HNRPAB antibodies and IgG antibodies on the

862 mouse sperm lysates showed that HNRPAB interacted with several RNAs associated

863 with fertilization and axoneme assembly. The qPCR analysis of RIP performed on the

864 sperm lysates from the *Iqch* KO mice revealed a decrease in the interaction between

865 HNRPAB and the RNA targets compared to the WT mice (Student's t-test; *p < 0.05;

866 error bars, s.e.m). **(D)** The Co-IP assay showed the binding of IQCH and CaM in the

867 WT sperm but not in the *Iqch* KO sperm. **(E)** The Co-IP assay showed that the

868 decreased expression of HNRPAB was due to the reduced binding of IQCH and CaM

869 by the knockout of IQCH or CaM. **(F)** The overexpression of IQCH and the

870 simultaneous knockout of CaM or the overexpression of CaM and the simultaneous

knockout of IQCH in K562 cells further confirmed that the downregulation of

HNRPAB was due to the diminished interaction between IQCH and CaM by western

blotting analysis. **(G)** Analysis of the binding affinity between GFP-IQCH from the

cell lysates (target) and recombinant CaM (ligand) by a microscale thermophoresis

(MST) assay showing the interaction between IQCH and CaM. Their interaction was

destroyed after the deletion of the IQ motif within IQCH. **(H)** The Co-IP assay of the

HEK 293 cells co-transfected with WT-*IQCH* and WT-*CaM* plasmids or

co-transfected with *IQCH* (△IQ) and WT-*CaM* plasmids showed that CaM interacted

with the IQ motif. The downregulation of HNRPAB was due to the disrupted

interaction between IQCH and CaM. Three independent experiments were performed.

Fig.7. Proposed model for the mechanisms of action underlying the involvement

of IQCH in spermatogenesis. IQCH interacts with CaM by IQ motif to regulate the

expression of RNA binding proteins. RNA binding proteins, particularly HNRPAB,

bind and regulate several RNAs that influence the development of acrosome,

mitochondria, and axoneme, thereby playing a critical role in spermatogenesis.

Table 1: Results of the semen analysis and sperm morphology examination of the

patient

Parameter	Patient	Reference
Sperm volume, ml	2.8	≥1.5
Sperm concentration, million/ml	10.0	≥15
Vitality, %	30	≥58
Motility, %	4	≥32
Abnormal morphology (%)	99.5	-

891 **Table 2. Semen analysis using CASA in the mouse model of *Iqch* KO**

	Adult Male Mice		
	WT	**KO**	**P[b] value**
Semen parameters			
Sperm concentration (10^6 /ml) [a]	94.51±10.68	67.75±3.70	0.038
Motility (%)	57.08±2.18	5.49±3.25	<0.001
Progressive motility (%)	57.00±2.05	5.49±3.25	<0.001
Sperm locomotion parameters			
Curvilinear velocity (VCL) (μm/s)	78.29±6.23	9.94±4.03	<0.001
Straight-line velocity (VSL) (μm/s)	36.18±2.09	2.90±1.98	<0.001
Average path velocity (VAP) (μm/s)	45.10±0. 19	4.17±2.49	<0.001
Amplitude of lateral head displacement (ALH) (μm)	0.77±0. 02	0.13±0.03	<0.001
Linearity (LIN)	0.46±0.01	0.27±0.08	0.047
Wobble (WOB, = VAP/VCL)	0.58±0.04	0.40±0.08	0.033
Straightness (STR, = VSL/VAP)	0.80±0.05	0.67±0.07	0.069
Beat-cross frequency (BCF) (Hz)	4.33±0.19	0.65±0.26	<0.001

892 [a] Epididymides and vas deferens, [b] two-tailed Student's t-test

893 n = 3 biologically independent WT mice or KO mice

A

III-2
Heterozygous
mutation

III-3
Heterozygous
mutation

II-1
Homozygous
mutation

IV-1
Homozygous
mutation

M=c.387+1_387+10del

WT allele

B

i

Marker NC pSPL3 WT-*IQCH* Mut-*IQCH*

381bp
263bp

ii

WT-*IQCH*

Exon V1   Exon4 deletion   Exon V2

ACCCAGCAACCTGGAG

Mut-*IQCH*

Exon V1 Exon V2

iii

c.387+1_387+10del

V1   Exon4   V2

WT-*IQCH*   Mut-*IQCH*

V1 Exon4 V2   V1   V2

92bp 118bp 171bp   92bp 171bp

iv

WT-c*IQCH*   Mut-c*IQCH*

IB: IQCH

IB: GAPDH

C

Control   Proband

DAPI/α-Tubulin

DAPI/IQCH

Merge

A

Biological Process | Cellular Component | Molecular Function (dot plots, Log2 Fold enrichment vs GO Terms Description, P value and Protein number legends)

B

Protein accession	Protein names	Gene names	Protein score
Q9D8E6	60S ribosomal protein L4	Rpl4	221
P47911	60S ribosomal protein L6	Rpl6	196
P14148	60S ribosomal protein L7	Rpl7	164
P47963	60S ribosomal protein L13	Rpl13	143
Q9CR57	60S ribosomal protein L14	Rpl14	109
P62908	40S ribosomal protein S3	Rps3	97
Q6ZWN5	40S ribosomal protein S9	Rps9	96
P35980	60S ribosomal protein L18	Rpl18	91
P27659	60S ribosomal protein L3	Rpl3	83
P62264	40S ribosomal protein S14	Rps14	83
P12970	60S ribosomal protein L7a	Rpl7a	73
P14131	40S ribosomal protein S16	Rps16	73
P19253	60S ribosomal protein L13a	Rpl13a	72
P62702	40S ribosomal protein S4, X isoform	Rps4x	66
P62301	40S ribosomal protein S13	Rps13	65
P47915	60S ribosomal protein L29	Rpl29	63
P14115	60S ribosomal protein L27a	Rpl27a	62
P61255	60S ribosomal protein L26	Rpl26	60
P84099	60S ribosomal protein L19	Rpl19	58
P25444	40S ribosomal protein S2	Rps2	53
P47962	60S ribosomal protein L5	Rpl5	53
Q9CPR4	60S ribosomal protein L17	Rpl17	51
P62754	40S ribosomal protein S6	Rps6	48
P14869	60S acidic ribosomal protein P0	Rplp0	47
P62751	60S ribosomal protein L23a	Rpl23a	47
P60867	40S ribosomal protein S20	Rps20	47
O09167	60S ribosomal protein L21	Rpl21	44
Q9CZM2	60S ribosomal protein L15	Rpl15	42
P35979	60S ribosomal protein L12	Rpl12	42
P97351	40S ribosomal protein S3a	Rps3a	36
P62855	40S ribosomal protein S26	Rps26	35
P62082	40S ribosomal protein S7	Rps7	34
P62849	40S ribosomal protein S24	Rps24	32

C

807 Upregulated Proteins   1186 Downregulated Proteins (heatmap, KO vs WT)

D

Biological Process / Cellular Component / Molecular Function dot plots (Log2 Fold enrichment vs GO Terms Description, P value and Protein number legends)

E

Proteomics_gene: 1110   76   LCMS_gene: 212

F

logFC   GO Terms ▢ RNA binding ▢ Mitochondrial function ▢ Calcium channel

IQ motif

IQCH

CaM

HNRPAB

RNA binding proteins

Dnhd1 *Catsper2*
Catsper1 *Catsper3*
Ccdc40 *Ccdc39*
Ccdc65 *Irrc6*

RNAs

Acrosomes

Flagellar axoneme

Acrosome
Acroplaxome
Manchette
Mitochondria
Axoneme