1   RESEARCH ARTICLE

The receptor-like kinase BIR1 inhibits elicitor-induced plasmodesmata callose deposition and PTI gene expression and requires EDS1 and SOBIR1 to cause dose-dependent cell-death in Arabidopsis

7   **Irene Guzmán-Benito[1],** Carmen Robinson[1,2], Chenlei Hua[3], Ana Rocio Sede[4], Laura Elvira-González[4], Isabel Punzón[1], Manfred Heinlein[4], Thorsten Nürnberger[3,5], César Llave[1]*

10   [1]Centro de Investigaciones Biológicas Margarita Salas. Consejo Superior de Investigaciones Científicas, Ramiro de Maeztu 9, 28040 Madrid, Spain; [2]Doctorado en Biotecnología y Recursos Genéticos de Plantas y Microorganismos Asociados, Universidad Politécnica de Madrid, 28040-Madrid, Spain; [3]Department of Plant Biochemistry, Center of Plant Molecular Biology (ZMBP), University of Tübingen, Tübingen, Germany; [4]Institut de Biologie Moléculaire des Plantes, CNRS, Université de Strasbourg, 67000 Strasbourg, France; [5]Department of Biochemistry, University of Johannesburg, Johannesburg, South Africa.

19   Author for correspondence: *César Llave*

20   *E-mail: cesarllave@cib.csic.es*

21   *Phone: +34-91-8373112*

WORD COUNT: 6457
Summary: 197
Introduction: 924
Materials and methods: 356
Results: 3532

Discussion: 1581

Acknowledgments: 64

Number of figures: 7

Number of figures in color: 7

Supporting information

Number of supporting methods: 3

Number of supporting figures: 8

Number of supporting tables: 8

ORCIDS

Irene Guzmán-Benito   0000-0002-9912-8164

Carmen Robinson   0000-0002-7146-0253

Chenlei Hua   0000-0002-6630-9049

Ana Rocio Sede   0000-0002-7868-0638

Laura Elvira-González   0000-0002-2530-8912

Manfred Heinlein   0000-0001-7322-1654

Thorsten Nürnberger   0000-0002-7804-7170

César Llave   0000-0003-3844-4582

Summary

- The receptor-like kinase BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) functions as a negative regulator of cell death and defense in Arabidopsis. Previous studies showed that BIR1 expression is up-regulated during infections with microbes and viruses. However, the biological consequences of BIR1 induction remain unknown.

- Here, we use a dexamethasone (DEX)-inducible expression system in *Arabidopsis thaliana* to investigate the outputs associated with physiological and non-physiological levels of BIR1 expression.

- We show that BIR1 induction at physiological levels significantly interferes with gene expression and plasmodesmata callose deposition triggered by canonical pattern-triggered immunity (PTI) elicitors. Plants that accumulated non-physiological doses of BIR1 displayed morphological defects that concur with transcriptomic changes in multiple plant defense genes. We provide experimental evidence that ENHANCED DISEASE SUSCEPTIBILITY 1 (EDS1) and SUPPRESSOR OF BIR1-1 (SOBIR1) are required for the effector-triggered immunity (ETI)-type cell death phenotypes associated with non-physiological levels of BIR1

- We propose that BIR1 induction may represent a pathogen-triggered mechanism to modulate plant defenses during infection. Our model predicts that when BIR1 regulation is lost, BIR1 integrity is sensed by one or several guarding resistance (R) proteins to initiate an ETI-like response, in which SOBIR1 cooperates with EDS1 to transduce signals downstream of R proteins.

Key words

Arabidopsis, BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1), cell death, ENHANCED DISEASE SUSCEPTIBILITY 1 (EDS1), immune regulation, immune receptor, pattern-triggered immunity, SUPPRESSOR OF BIR1 (SOBIR1)

Introduction

Plants use immunity to defend themselves against invaders such as bacteria, fungi, oomycetes, and viruses (Korner *et al.*, 2013; Nishad *et al.*, 2020). Pattern-triggered immunity (PTI) uses multiple cell-surface pattern recognition receptors (PRR) to recognize pathogen/microbe/damage-associated molecular patterns (PAMP/MAMP/DAMP). Effector-triggered immunity (ETI) involves intracellular receptors that sense pathogen-effector proteins or effector-induced manipulations of host proteins (Zhou & Zhang, 2020; Yu *et al.*, 2021). PRRs are single-transmembrane receptor-like kinases (RLK) or receptor-like proteins (RLP). PRRs form active receptor complexes in a ligand-dependent manner with members of the somatic embryogenesis co-receptor kinases (SERKs), including BRASSINOSTEROID INSENSITIVE1-ASSOCIATED RECEPTOR KINASE 1 (BAK1/SERK3) (Boutrot & Zipfel, 2017; Hohmann *et al.*, 2017; Smakowska-Luzan *et al.*, 2018). BAK1 functions as a positive regulator of PTI (Wu *et al.*, 2020), and has a dual role as positive and negative regulator of cell death during ETI (Belkhadir *et al.*, 2012; Dominguez-Ferreras *et al.*, 2015). RLP-type PRRs lack kinase activity and constitutively associate in a ligand-independent manner with the leucine-rich repeat (LRR)-RLK SUPPRESSOR OF BIR1-1 (SOBIR1) (Gust *et al.*, 2017). Once the microbe is sensed, PRR complexes activate receptor-like cytoplasmic kinases (RLCKs) that initiate downstream signaling cascades (Liang & Zhou, 2018; Rao *et al.*, 2018). Among them, BOTRYTIS INDUCED KINASE1 (BIK1) functions as a positive regulator of immunity when triggered by LRR-RLKs or as a negative regulator when triggered by LRR-RLPs (Bi *et al.*, 2018; Wan *et al.*, 2019).

ETI relies on nucleotide-binding domain LRR intracellular immune receptors (NLRs) encoded by polymorphic disease resistance genes (R). NLRs comprise either proteins with an N-terminal domain similar to the TOLL/INTERLEUKIN RECEPTOR 1 (TIR1) (TIR-NB-LRR, TNL), proteins with a coiled-coil (CC) domain at the N terminus (CC-NB-LRR, CNL) or proteins with a RESISTANCE TO POWDERY MILDEW 8-LIKE DOMAIN (RPW8)-type domain (RNLs) (van Wersch *et al.*, 2020; Wang & Chai, 2020). NLRs function as "sensors" that recognize effectors, or as "helpers" that mediate signaling downstream of sensor NLRs (Jubic *et al.*, 2019). Recent studies demonstrated that PTI and ETI work together to mutually potentiate the immune response (Yuan *et al.*, 2021b; Yuan *et al.*, 2023). NLR-mediated signaling is dependent on multiple PTI-associated components such as BAK1, SOBIR1, or BIK1 in a PTI-independent manner, while TNL genes that are up-regulated during early PTI boost immune responses (Ngou *et al.*, 2021; Tian *et al.*, 2021; Yuan *et al.*, 2021a). ENHANCED

DISEASE SUSCEPTIBILITY (EDS1) and PHYTOALEXIN DEFICIENT 4 (PAD4), which

are essential in TNL-mediated immunity, also play a key role in PTI signaling in response to

different microbial elicitors (Pruitt *et al.*, 2021b; Tian *et al.*, 2021). Furthermore, PTI and ETI

outputs usually converge, albeit with different amplitude and duration (Yuan *et al.*, 2021b).

Early immune responses include apoplastic reactive oxygen species (ROS) burst, cytosolic

calcium (Ca^{+2}) influx and activation of Ca^{+2}-dependent protein kinases (CPK), and activation

of mitogen-activated protein kinases (MAPK). This response is followed by the transcriptional

activation of defense processes that include accumulation of defense phytohormones, regulation

of cell wall and secretome composition (Li *et al.*, 2020).

Plants deploy a variety of mechanisms to avoid the detrimental effects of hyperactivated

defenses (Couto & Zipfel, 2016; Withers & Dong, 2017; Mithoe & Menke, 2018). BAK1-

INTERACTING RECEPTOR-LIKE KINASE (BIR) proteins play a key role in defense

attenuation. The current model postulates that, in the absence of microbes, BIR proteins

constitutively associate with BAK1, maintaining the PRRs in a resting state. During ligand

perception, BAK1 dissociates from BIR and becomes accessible to complex with PRR

receptors (Gao *et al.*, 2009; Halter *et al.*, 2014; Imkampe *et al.*, 2017). The Arabidopsis genome

contains four BIR homologs. Arabidopsis BIR1 negatively regulates cell death signaling

pathways that depend on BAK1, SOBIR1, EDS1 and PAD4, and that likely involve

components of the TNL-mediated ETI response (Gao *et al.*, 2009; Schulze *et al.*, 2022).

Arabidopsis BIR2 and BIR3 have critical roles in the negative regulation of the PRR-mediated

PTI response (Halter *et al.*, 2014; Imkampe *et al.*, 2017). BIR4 has so far not been characterized.

Under normal conditions, BIR1 expression is transcriptionally activated by multiple

microbial and viral pathogens in a salicylic acid (SA)-dependent manner (Guzman-Benito *et*

al., 2019). RNA-directed DNA methylation and post-transcriptional RNA silencing maintain

BIR1 transcripts at physiological levels when BIR1 is actively transcribed (Guzman-Benito *et*

al., 2019). However, transgenic overexpression of BIR1 at non-physiological levels triggers

developmental defects and cell death similar to the autoimmune phenotypes observed in *bir1-*

1 mutants (Gao *et al.*, 2009; Guzman-Benito *et al.*, 2019). Given that perturbations in BIR1

homeostasis have severe growth-defense penalties, effective strategies for BIR1 regulation are

needed to prevent unbearably high levels of BIR1 expression. To date, the effects and

importance of BIR1 induction in plant defense remain largely unknown.

In this work, we conducted a time-course analysis of the phenotypical outputs associated

with physiological and non-physiological levels of BIR1 expression in Arabidopsis. We first

tested if BIR induction interferes with the immune response triggered by canonical PAMPs. We found that physiological induction of BIR1 represses the expression of PTI genes and plasmodesmata (PD) callose deposition in response to several PTI elicitors. Whereas physiological induction of BIR1 has a subtle impact on the plant transcriptome, the morphological and growth defects caused when BIR1 is expressed at non-physiological levels are concomitant with the transcriptional reprogramming of multiple genes encoding components of PTI and ETI pathways. We found that the cell death phenotypes associated with BIR1 overexpression require the TNL-associated EDS1 and/or the RLP-co-receptor SOBIR1, but the underlying mechanism is unknown. Finally, our work demonstrates that a well-balanced expression of BIR1 avoids interference with cellular processes of growth and development, and is critical for plant fitness.

Material and Methods

Plant material and growth condition

Arabidopsis thaliana plants were grown in controlled environmental chambers under long-day conditions (16h day/8h night) at 22°C. Arabidopsis lines used in this study were derived from the Columbia-0 (Col-0) ecotype. Seeds were surface-sterilized and sown on plates containing ½ Murashige and Skoog (MS) medium supplemented with MES (0,5g/L), 1% sucrose and 0,9% Plant agar (Duchefa Biochemie). Seedlings were transferred to soil 7 to 10 days after germination.

Mutants for *bir1-1* and *sobir1-12* were donated by Yuelin Zhang (University of British Columbia, Canada). The Arabidopsis *eds1-2*, *pad4-1* and *eds5-3* alleles were supplied by Jane E. Parker (Max Planck Institute for Plant Breeding Research, Germany). Arabidopsis mutant *bak1-5* was donated by Birgit Kemmerling (Center of Plant Molecular Biology, University of Tübingen, Germany). The *sid2-2* mutant was a gift from Francisco Tenllado (Centro de Investigaciones Biológicas Margarita Salas, CSIC, Spain).

Arabidopsis Col-0 transgenic plants expressing a BIR1-mCherry fusion under a dexamethasone (DEX)-inducible gene expression system (BIR1 L6 and BIR1 L9) have already been described (Guzman-Benito *et al.*, 2019). In this study, the same DEX-inducible expression system containing a mCherry-tagged BIR1 (hereafter BIR1) protein-coding gene was transformed into Arabidopsis mutant backgrounds according to standard floral dipping (Clough & Bent, 1998). Independent homozygous lines were selected in T3 and used for subsequent

174 experiments. Morphological phenotypes after DEX treatments were investigated in two
175 independent transgenic lines of each genotype.

Protein and RNA analyses

177 Protein assays and one-step quantitative RT-PCR (RT-qPCR) using gene-specific primers
178 (Table S1) and CBP20 (At5g44200) as endogenous reference gene are described in Methods
179 S1.

Library preparation for transcriptome sequencing and bioinformatic analyses

A total of 24 samples, including three biological replicates per treatment, were collected for
RNA-Seq. Each sample consisted of RNA pooled from six individual plants. Sequencing
libraries, transcriptome sequencing of paired-end reads using an Illumina PE150 Hiseq platform
and bioinformatic analyses were done at NOVOGENE (HK) COMPANY LIMITED (Wan
Chai, Hong Kong) as described (Methods S2). The raw data was deposited in the Gene
Expression Omnibus under accession number GSE234036.

Analyses of Reactive oxygen species (ROS) and ethylene, plasmodesmata callose deposition and dexamethasone (DEX) treatments.

189 Experimental procedures are described in Methods S3.

Results

BIR1 induction causes dose-dependent transcriptome reprogramming in Arabidopsis

We conducted a RNA-Seq assay on Arabidopsis BIR1 L9 plants, which express a DEX-
inducible BIR1-mCherry transgene (Guzman-Benito *et al.*, 2019), to understand the
downstream biological processes and pathways associated with BIR1 induction, This system
proved to be a valuable tool for studying BIR1 effects in the absence of pathogen elicitors or
effectors. Three-week-old plants were sprayed with a 30 µM DEX solution at 24 h intervals
and rosette leaf samples were harvested for next-generation sequencing. Control plants were
mock-treated with H_2O. Three biological replicates were used as representative samples of each
condition tested in this study: DEX-treated WT (DeWT_), mock-treated WT (MoWT_), DEX-
treated BIR1 L9 (DeL9_) and mock-treated BIR1 L9 (MoL9_) plants at 4 or 12 days of
treatment as indicated **(Table S2)**. After four days of DEX treatment, BIR1 L9 plants developed
normally like WT plants, but they started showing a range of morphological and growth defects
from day 12 onward, including stunting, leaf deformation, premature senescence (yellowing),

and cell-death **(Fig. 1a)**. To avoid extensive damage on the plant transcriptome, DeL9_ samples
were collected before the onset of cell death symptoms. DeL9_ samples accumulated BIR1 at
physiological levels after 4 days of DEX treatment whereas non-physiological levels of BIR1
occurred after long BIR1 induction periods (12 days of DEX treatment) **(Fig. 1a)**. (Guzman-
Benito *et al.*, 2019).

Principal component analysis (PCA) was used to sort RNA-seq-based transcriptomic data
according to gene expression levels. To discard the potential effects of DEX on gene expression,
we first compared DEX-treated WT versus DEX-treated BIR1 L9 plants. PC1 defined a distinct
separation in the PCA score plot between WT (DeWT_12) and BIR1 L9 (DeL9_12) samples
after 12 days of treatment, indicating that the BIR1 expression level was the main source of
variance (36.92%) **(Fig. 1b)**. Samples in the DeL9_4 group, which produced physiological
levels of BIR1, clustered in the proximity of non-transgenic DeWT_4 samples, reflecting their
similar expression profiles. PC2 seems to highlight a shift in gene expression between samples
due to age or developmental stage (22.07%). When DEX-treated BIR1 L9 plants were
compared with mock-treated BIR1 L9 plants, the BIR1 expression level was again the dominant
source of variance (36.85%) underlying PC1, whereas time was the basis for PC2 (19.57%)
separation in the PCA score plot **(Fig. S1a)**. BIR1 abundance [measured in Fragments Per
Kilobase of transcript sequence per Millions of base-pairs sequenced (FPKM)] in replicate
DeL9_12b was substantially lower than in the other two DeL9_12 replicates within the
DeL9_12 group **(Fig. 1a)**. Interestingly, the DeL9_12b sample was located far from the other
DeL9_12 samples in the PCA plot, and was found near the samples within the DeWT_12 or
MoL9_12 clusters **(Figs. 1b, S1a)**. This finding may suggest that moderate BIR1
overexpression produced less transcriptomic alterations than high BIR1 overexpression levels
relative to the controls.

Differentially expressed genes (DEG) from different comparisons are presented in **Table
S3**. Our data indicate that a massive transcriptional reprogramming occurs when BIR1 is
overexpressed in Arabidopsis (12 days of DEX treatment) but not when BIR1 is
transcriptionally induced at physiological levels (4 days of DEX treatment) **(Fig. 1c)** (**Table
S4)**. DEGs using different statistical thresholds are shown in **Figs. S1b,c**. Gene ontology (GO)
analysis showed that genes encoding proteins with oxidoreductase activity, transmembrane
transport and other metabolic processes or genes related to lipid or carbohydrate metabolism
were particularly overrepresented in our whole set of DEGs, including WT and BIR1 L9 plants
(Figs. 1d, S1d, S2d) (Table S5). These alterations presumably reflect the effects of DEX on

the plant transcriptome. However, DEGs with kinase activity were only enriched in BIR1 overexpressing plants, relative to mock-treated BIR1 L9 plants **(Figs. 1d, S1d)** or DEX-treated WT plants **(Fig. S2d),** ruling out an effect caused by DEX. The Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analysis showed the enrichment of genes related to several metabolic processes and genes involved in the biosynthesis of secondary metabolites that were specifically activated after DEX-treatments **(Fig. S3) (Table S6)**. Conversely, DEGs involved in plant hormone signal transduction were significantly enriched only in DEX-treated BIR1 L9, but not in mock-treated BIR1 L9 plants or DEX-treated WT plants. The impact of DEX application on plant morphology and gene expression in WT plants is shown in **Fig. S2** and **Table S4**.

Effects of BIR1 induction on PTI activation in response to bacterial elicitors

We assessed the effects of inducing BIR1 at physiological levels on the early events of PTI response. BIR1 was induced by treating BIR1 transgenic plants (L6 and L9) with a 30 µM DEX solution 4 times at 24 h intervals, whereas control plants were mock-treated with H_2O. The flg22 peptide derived from bacterial flagellin and the nlp20 fragment of NECROSIS AND ETHYLENE-INDUCING PEPTIDE 1 (NEP1)-LIKE proteins (NLP) were used as elicitors. The luminol-based assay showed that ROS production triggered by 1 µM flg22 or nlp20 was comparable between WT, BIR1 L6 and BIR1 L9 plants **(Fig. 2a)**. Likewise, production of the phytohormone ethylene was similarly increased in DEX-treated WT, BIR1 L6 and BIR1 L9 plants in response to flg22 and, to a greater extent, in response to nlp20 at concentrations of 1 µM and 100 nM. These findings indicate that induction of BIR1 had no measurable effects on ROS and ethylene accumulation triggered by canonical elicitors **(Fig. 2b)**. Next, we tested the expression of genes encoding the canonical PTI markers FLG22-INDUCED RECEPTOR-LIKE KINASE 1 (FRK1) and PATHOGENESIS RELATED 1 (PR1) in BIR1 L6 and BIR1 L9 transgenic plants upon 1 µM flg22 elicitation. Whereas all these genes were significantly up-regulated in mock-treated controls, they were no longer activated upon BIR1 induction in DEX-treated BIR1 L6 an BIR1 L9 plants. Both PTI markers were altered by flg22 in WT plants treated with DEX, which discarded a potential repressive effect of DEX on PTI gene expression **(Fig. 2c)**. The induction of BIR1 protein in the BIR1 L6 and BIR1 L9 plants used for ROS and ethylene assays was confirmed by immunocapture followed by western blot, while BIR1 transcripts were monitored in the samples used for RT-qPCR expression assays (**Fig. 2d**).

Collectively, these results indicate that BIR1 induction at physiological levels negligibly

affects ROS production and ethylene accumulation but interferes with downstream events of

the PTI pathway, impacting the transcriptional activation of defense genes.

Induction of BIR1 hinders callose deposition at PD triggered by flg22 or dsRNA

During the early stages of pathogen invasion, conserved PAMPs trigger callose depositions in

the cell-wall and PD (Luna *et al.*, 2011). Deposition of callose at PD regulates symplastic

transport, and limits the ability of the invading pathogen to enter plant tissues (Tang *et al.*, 2017;

Huang *et al.*, 2023). To investigate whether BIR1 induced at physiological levels affects callose

deposition in PD, we used *in vivo* aniline blue staining to quantify PD-associated callose from

digital photographs of BIR1 L9 Arabidopsis plants. BIR1 was induced with a 30 μM DEX

solution 3 times at 24h intervals, whereas control plants were mock-treated with H_2O. We first

investigated callose deposition using 1 μM flg22 as elicitor. Compared to plants treated without

the elicitor (H_2O), flg22 caused a noticeable increase in PD callose intensity BIR1 L9 control

plants (DEX-) **(Figs. 3a,c)**. In contrast, PD-associated callose was no longer induced by flg22

in plants expressing the BIR1 transgene (DEX+) **(Figs. 3a,c)**. BIR1 L9 plants were later treated

with the synthetic dsRNA analog polyinosinic-polycytidylic acid [poly(I:C)] (0.5 μg μl^{-1}) as

dsRNA is known to trigger deposition of callose and PD closure in Arabidopsis (Huang *et al.*,

2023). In the absence of DEX, callose intensity was significantly higher in BIR1 L9 plants

treated with poly(I:C) compared to H_2O-treated plants. In contrast, after DEX treatment,

poly(I:C)-induced callose was significantly repressed by the expression of BIR1 **(Figs. 3b,c)**.

We observed enhanced callose deposition in WT controls and DEX-treated WT plants in

response to flg22 and poly(I:C) relative to H_2O-treated plants, suggesting that DEX does not

affect callose deposition **(Fig. 3d)**. These findings indicate that the induction of BIR1 impairs,

at least partially, PD callose deposition triggered by different PAMP elicitors.

BIR1 overexpression triggers the expression of immunity-related genes

RNA-Seq results were further validated in mock- and DEX-treated BIR1 L9 plants after 12

days of treatment using RT-qPCR. We found that expression of DEGs encoding BIR1, PR1,

RESPIRATORY BURST OXIDASE HOMOLOG PROTEIN D (RBOHD), FRK1 and

CYTOCHROME P450 MONOOXYGENASE 79B2 (CYP79B2) defense markers were all up-

regulated in Arabidopsis plants overexpressing BIR1 compared to mock-treated controls, which

is consistent with their RNA-Seq-based profiles and confirms reproducibility in their expression

patterns **(Fig. 4a)**.

A detailed inspection of the RNA-Seq data revealed that multiple genes involved in the

perception of pathogens and the activation of defense pathways were misregulated when BIR1

was overexpressed [|$\log_2$ (Fold-Change)| ≥ 1 and adjusted p-value ≤ 0.01] **(Table S7, S8)**. Two-

tailed Fisher's exact tests revealed that genes encoding LRR-type RLKs in the Arabidopsis

genome were significantly enriched **(Table S7)**. Cysteine-rich RLKs (CRK), which are

important in the perception of apoplastic ROS and, likely in association with BAK1, in the

activation of PTI and cell death, were overrepresented in the pool of DEGs **(Table S7) (Fig.**

S4) (Chen, 2001)(Yeh *et al.*, 2015; Quezada *et al.*, 2019; Castro *et al.*, 2021). Furthermore, a

significant percentage of RLP coding genes in Arabidopsis were in the list of DEGs **(Tables**

S7) (Fig. S5) (Wang *et al.*, 2008; Lv *et al.*, 2016). Most of LRR-RLK, CRK and RLP transcripts

were up-regulated in response to elevated BIR1, but not in DEX-treated WT plants. BIR1

overexpression boosted the transcription of many other genes involved in PTI and ETI signaling

(Figs. 4b, S6). This includes the RLK co-receptors BAK1 and SOBIR1, RLCKs such as BIK1

and several AvrPphB SUSCEPTIBLE1(PBS1)-LIKE (PBL) kinases (PBL4, PBL12, PBL20)

(Liang & Zhou, 2018; Rao *et al.*, 2018) **(Table S7)**. Active BIK1 positively regulates the Ca^{2+}

influx from the apoplast to the cytosol, which switches on the kinase activity of CPKs (Li *et*

al., 2014; Yip Delormel & Boudsocq, 2019). Collectively, CPKs were not differentially

regulated, but many individual CPKs were regarded as DEGs in plants overexpressing BIR1

(Table S3). For instance, the expression of CPK4-6 and CPK11, which are key players in

transcriptional reprogramming upon elicitor perception, was enhanced only in DEX-treated

BIR1 L9 plants **(Fig. S7a)** (Boudsocq *et al.*, 2010). Upon ligand perception, BIK1 dissociates

from activated receptor complexes and phosphorylates the plasma membrane NADPH oxidase

RBOHD to generate extracellular ROS (Kadota *et al.*, 2014; Li *et al.*, 2014). Interestingly,

RBOHD was induced in DEX-treated BIR1 L9 plants after both 4 and 12 days of treatment, but

not in their corresponding controls **(Figs. 4a,b)**. Genes encoding the lipase-like proteins EDS1,

PAD4, SENESCENCE ASSOCIATED-GENE 101 (SAG101) and ACTIVATED DISEASE

RESISTANCE 1 (ADR1), which are required for TNL-receptor function during ETI, were also

up-regulated (Table S3) **(Fig. 4b)** (Zhou & Zhang, 2020; Ngou *et al.*, 2021; Pruitt *et al.*, 2021b;

Sun *et al.*, 2021; Tian *et al.*, 2021). Accordingly, a large number of DEGs in our assay were

annotated as disease resistance (R) protein-coding genes, of which the TNL class was

particularly enriched in BIR1 overexpressing plants (Swiderski *et al.*, 2009) **(Table S7)**.

Conversely, neither NON-RACE SPECIFIC DISEASE RESISTANCE 1 (NDR1) nor genes encoding CNL intracellular receptors, which require NDR1, were differentially expressed in response to BIR1 overexpression **(Fig. 4b) (Table S3)**.

EDS5, which transduces the signal downstream of SA in Arabidopsis (Nawrath *et al.*, 2002), was up-regulated in DEX-treated BIR1 L9 samples. Interestingly, BIR2 was up-regulated, whereas BIR3 and its partner CONSTITUTIVE SHADE AVOIDANCE 1 (CSA1), which also interacts with BIR1, were unaffected by BIR1 overexpression **(Fig. S6)** (Schulze *et al.*, 2022). Finally, transcripts encoding the plant heterotrimeric G proteins ARABIDOPSIS G PROTEIN α-SUBUNIT1 (GPA1), ARABIDOPSIS G PROTEIN β-SUBUNIT 1 (AGB1), ARABIDOPSIS G PROTEIN γ-SUBUNIT 2 (AGG2), and EXTRA-LARGE G PROTEIN 1 and 2 (XLG1/2) were more abundant in plants overexpressing BIR1 relative to controls after 12 days of DEX treatment (**Fig. S7b**). These proteins positively regulate stomatal defense and PTI signaling, ROS production and BIK1 stability allowing optimal immune activation (Liang *et al.*, 2016; Zhong *et al.*, 2019). Consistent with transcript levels, SOBIR1, PAD4-dependent PR1 and EDS1 proteins were strongly up-regulated by BIR1 overexpression in DEX-treated BIR1 L9 plants, compared to the non-treated controls (**Figs. 4c, 5c**).

In conclusion, in the absence of microbes or microbe-derived elicitors/effectors, the expression of BIR1 at non-physiological levels alters hormone homeostasis, and leads to an inadequate over-activation of multiple immune receptors and co-receptors involved in the perception and signaling of ligands and effectors.

Genetic disruption of SOBIR1, but not BAK1, partially rescues BIR1 overexpression phenotypes

In the following sections we tested if mutations in key components of PTI and ETI signaling alleviated cell death and immune alterations caused by non-physiological expression of BIR1 (Rodriguez *et al.*, 2016; Freh *et al.*, 2022). SOBIR1 and BAK1 are necessary for activating cell death and constitutive defense responses in the *bir1-1* mutant (Gao *et al.*, 2009; Liu *et al.*, 2016). Therefore, we tested if SOBIR1 and BAK1 were also required for the BIR1 phenotypes. A DEX-inducible BIR1-mCherry system was used to generate independent homozygous transgenic lines in the well-characterized *sobir1-12* and *bak1-5* mutant backgrounds (Gao *et al.*, 2009; Schwessinger *et al.*, 2011). Transgenic plants were sprayed with a 30 µM DEX

solution or H₂O (mock-treated controls) for 12 days at 24 h intervals to trigger BIR1 overexpression.

As previously shown, our reference set of DEX-treated BIR1 L9 plants accumulated high levels of BIR1 transcripts, and displayed stunting symptoms, accelerated senescence, and cell death relative to non-treated BIR1 L9 controls or DEX-treated WT plants **(Fig. 5a)** (Guzman-Benito *et al.*, 2019). When *sobir1-12* BIR1 plants were investigated, we observed that the phenotypes after 12 days of DEX treatments, albeit variable between biological replicates, were milder than those in BIR1 L9 plants **(Figs. 5a, S8a,b)**. Interestingly, BIR1 transcript and protein accumulated in *sobir1-12* BIR1 plants like in BIR1 L9 plants after 12 days of DEX application, discarding potential dose effects **(Fig. 5b,c)**. Transcripts of the PTI-markers RBOHD, FRK1 and WRKY DNA-BINDING PROTEIN 29 (WRKY29), were increased, albeit not significantly, in DEX-treated *sobir1-1*2 BIR1 plants relative to non-treated controls **(Figs. 5d, S8c)**. These finding suggest that cell death and PTI gene activation due to BIR1 overexpression were alleviated by the lack of SOBIR1.

bak1-5 BIR1 plants consistently developed strong morphological phenotypes after 12 days of DEX application, that closely resemble those of BIR1 L9 plants **(Figs. 5a, S8a,b)**. RT-qPCR showed that BIR1 transcripts also accumulated in these plants after long DEX treatments, although to a variable degree depending on the transgenic lines and biological replicates. For instance, BIR1 transcript levels in *bak1*-5 BIR1 plants were comparable to those in *sobir1-12* BIR1 plants in **Fig. S8b**. Differences in BIR1 protein accumulation were observed between DEX-treated *bak1-5* BIR1 L4 and L10 lines in this experiment **(Fig. 5c)**. **Figs. 5a** and **b** illustrate data from a different experimental replicate in which BIR1 transcripts in *bak1*-5 BIR1 plants were much less abundant than in BIR1 L9 plants. These interesting observations suggest that the persistent induction of BIR1 over time, rather than an excess of BIR1, was the trigger of the morphological alterations in the *bak1*-5 mutant background. RBOHD, FRK1 and WRKY remained transcriptionally up-regulated in DEX-treated *bak1-5* BIR1 plants although at varying degrees between replicates **(Fig. 5e)**. Collectively, these data indicate that the morphological phenotypes and, perhaps, PTI gene activation when BIR1 is expressed at non-physiological levels are genetically unrelated to BAK1.

Alterations in plant hormone pathways may negatively affect plant growth (Denance *et al.*, 2013). The transcriptional reprogramming of genes involved in plant hormone metabolism and signaling in BIR1-overexpressing plants prompted us to further examine their expression in the mutant backgrounds mentioned above. RT-qPCR revealed that genes encoding salicylic acid

(SA)-dependent PR1, jasmonic acid (JA)-responsive PLANT DEFENSIN 1.2 (PDF1.2), ethylene (ET)- and JA-responsive PR4 as well as genes encoding the cytochrome P450 enzymes CYP71B15 (PAD3) and CYP79B2 involved in camalexin (CA)-biosynthesis and CA-mediated plant defense, were all up-regulated in BIR1-overexpressing plants of *sobir1-12* BIR1 genotypes **(Fig. 5f)**. Intriguingly, the induction of phytohormone-related genes was notably reduced in *bak1-5* BIR1 plants after DEX treatment compared to *sobir1-12* BIR1 or BIR1 L9 plants **(Figs. 4a, 5g)**. This data suggests that the *bak1-5* mutation may, at least partially, compromise hormone defense signaling upon BIR1 overexpression. None of these genes, except PR1, were induced or repressed in DEX-treated WT plants, discarding a potential effect of DEX on their expression **(Table S3)**.

Genetic disruption of EDS1, but not PAD4, partially rescues BIR1 overexpression phenotypes

A previous report showed that stunting, activation of cell death and defense responses in the *bir1-1* mutant were partially dependent on EDS1 and PAD4 (Gao *et al.*, 2009). The EDS1-PAD4 node has a relevant role in PRR signaling and TNL activation, indicating that it mediates both PTI and ETI (Pruitt *et al.*, 2021a). We had observed that TNL-coding transcripts were abundant in BIR1 L9 plants and that EDS1 transcripts and protein were elevated in BIR1 L9, *sobir1-12* BIR1 and *bak1-5* BIR1 plants (**Figs. 5b,c**). Therefore, we wondered if TNL-mediated ETI responses along with, or independently of, PTI could be important for BIR1 overexpression phenotypes. Thus, we tested if the BIR1 phenotypes were affected by EDS1 or PAD4 mutations. The DEX-inducible BIR1-mCherry system was used to generate homozygous transgenic lines from *eds1-2* and *pad4-1* mutants (Parker *et al.*, 1996; Zhou *et al.*, 1998). The *eds1-2* BIR1 plants displayed mostly WT-like morphologies and, occasionally, localized yellowing after 12 days of DEX treatments compared to non-treated plants **(Figs. 6a, S8a)**. RT-qPCR analysis showed that BIR1 accumulated in the *eds1-2* BIR1 plants as much as in the BIR1 L9 plants **(Figs. 6a)**. The expression of FRK1 and RBOHD was significantly enhanced in *eds1-2* BIR1 plants after DEX applications compared to non-treated controls **(Fig. 6b)**. WRKY29 showed a tendency, albeit not significant, to be up-regulated in these plants (Fig. 6b). Our data indicated that the lack of EDS1 attenuates the phenotypic alterations caused by BIR1 overexpression, but does not affect the activation of PTI genes.

Next, we found that *pad4-1* BIR1 plants essentially had BIR1 L9-like phenotypes after 12 days of DEX applications **(Figs. 6a, S8a)**. In these plants, BIR1 accumulated much less than in BIR1 L9 or *eds1-2* BIR1 plants **(Fig. 6a)**. As shown previously for *bak1-5* BIR1 plants, the sustained induction of BIR1, rather than BIR1 overexpression, might be sufficient to cause typical BIR1 phenotypes regardless of the *pad4-1* mutation. Interestingly, the PTI-components FRK1, WRKY29 and RBOHD showed no responsiveness to BIR1 overexpression in *pad4-1* BIR1 plants, which suggests that PTI gene activation in these plants is PAD4-dependent **Figs. 6c, S8c)**. In summary, the genetic elimination of EDS1, but not PAD4, mitigates the morphological phenotypes caused by the induction of BIR1 at non-physiological levels. Likely, symptoms arise from the activation of EDS1-dependent ETI-type cell death in BIR1 overexpressing plants, which is independent of PTI gene activation.

RT-qPCR revealed a significant transcriptional activation of genes encoding markers in the SA, JA, ET and CA pathways in DEX-treated *eds1-2* BIR1 plants versus non-treated controls **(Fig. 6d)**. PDF1.2 and PR4 transcript markers were also up-regulated, but to a lesser extent, in *pad4*-1 BIR1 transgenic plants after 12 days of DEX treatments relative to the non-treated controls, whereas PAD4-dependent PR1 and PAD3 transcripts were not differentially expressed, suggesting that PAD4 was important for PTI gene expression and phytohormone homeostasis **(Fig. 6e)**.

Phenotypes of BIR1 overexpressing lines are unrelated to SA signaling pathways

pad4-1 BIR1 plants were as sensitive to DEX treatments as BIR1 L9 plants, suggesting that morphological phenotypes caused by non-physiological BIR1 levels are independent of SA-mediated signaling. To confirm this prediction, the effects of mutations in SALICYLIC ACID INDUCTION DEFICIENT 2 (SID2) and EDS5, involved in pathogen-induced SA biosynthesis (Huang *et al.*, 2020), were assessed in BIR1 overexpressing plants. DEX-inducible BIR1-mCherry transgenic plants were generated in the *sid2-2* and *eds5-3* (*eds5* is allelic to *sid1*) backgrounds, which contain low levels of SA (Nawrath *et al.*, 2002). We found that *sid2-2* BIR1 plants exhibited from moderate to severe morphological phenotypes after 12 days of DEX treatment. These coincided with moderate levels of BIR1 overexpression relative to those observed in BIR1 L9 plants **(Figs. 7a)**. The transcription levels of PR4, PDF1.2, PAD3 and CYP79B2 were significantly elevated in DEX-treated *sid2-2* BIR1 plants compared to non-treated controls **(Fig. 7b)**. *eds5*-3 BIR1 plants typically displayed strong developmental defects

and necrosis and accumulated BIR1 transcripts at BIR1 L9-like levels after long DEX applications **(Figs. 7a)**. PR4, PAD3 and PDF1.2, but not CYP79B2, were up-regulated in *eds5-3* BIR1 overexpressing plants relative to non-treated controls **(Fig. 7c)**. Interestingly, the canonical SA-marker PR1 was also induced in *eds5-3* BIR1 and, to a much lesser extent, *sid2-2* BIR1 plants. This observation suggests that PR1 induction was only partially dependent on EDS5 and SID2 in these plants (Tsuda *et al.*, 2013) **(Fig. 7b,c)**. Although residual EDS5/SID2-independent SA production in BIR1 overexpressing plants cannot be fully discarded (Lu, 2009), our findings suggest that the ETI-like phenotypes in these plants were not drastically affected by mutations in SA biosynthesis (*sid2-2* and *eds5-3*) and SA signaling (*pad4-1*). This idea is consistent with the observation that the expression of most SA-responsive genes, including PR1, is also controlled by SA-independent mechanisms during EDS1-dependent ETI (Tsuda *et al.*, 2013; Cui *et al.*, 2017).

Discussion

Plants respond to microbes and viruses by enhancing the expression of multiple genes that function as regulators or executors of defense programs (Nishad *et al.*, 2020). Among them, BIR1 was shown to be up-regulated during bacterial and oomycete infections, and also during viral infections in a SA-dependent manner (Gao *et al.*, 2009; Guzman-Benito *et al.*, 2019). The observation that knock-out *bir1-1* mutants have increased resistance to the virulent oomycete *Hyaloperonospora parasitica* Noco2 and to the RNA virus *Tobacco rattle virus* (TRV) suggests an active role of BIR1 in plant immunity (Gao *et al.*, 2009; Guzman-Benito *et al.*, 2019). Since BIR1 is a negative regulator of cell death and defense (Liu *et al.*, 2016), we hypothesized that BIR1 induction might be part of a common strategy to modulate, at least partially, the amplitude of the immune responses during microbial and viral infections. To test this hypothesis, we used a previously described DEX-inducible system for the conditional expression of a mCherry-tagged BIR1 transgene in Arabidopsis (Guzman-Benito *et al.*, 2019). By using different induction times, this system allowed us to investigate the outputs associated with physiological and non-physiological levels of BIR1. Even though adverse effects due to long DEX treatments cannot be fully discarded in our experimental system, specific effects resulting from BIR1 overexpression could be distinguished from other potential combinatorial effects resulting from expressing BIR1 in an activated glucocorticoid-inducible background (Kang *et al.*, 1999). Importantly, BIR1 induction is triggered in the absence of immune elicitors or effectors that could otherwise initiate undesirable localized and systemic defense responses in the plant.

The induction of BIR1 at a level comparable to that observed under natural conditions in bacteria- or virus-infected plants, or plants treated with SA was referred to as physiological in our assay (Gao *et al.*, 2009; Guzman-Benito *et al.*, 2019). At this level, BIR1 had a subtle, if any, impact on both plant transcriptome and plant morphology. In contrast, in the presence of the PTI elicitor flg22, the induction of BIR1 negatively affects the expression of several PTI-marker genes, but has no effects on ROS burst, as previously shown (Liu *et al.*, 2016), or ethylene production. This observation suggests that BIR1 can regulate, directly or indirectly, the activity of components in the PTI signaling pathway that ultimately control immune gene transcription. Furthermore, the physiological induction of BIR1 compromises the callose deposition at PD elicited by either flg22 or the dsRNA-analog poly(I:C). Targeted deposition of callose modulates symplastic connectivity by regulating PD pore size, and therefore constitutes an essential layer in the defense against pathogens (Wang *et al.*, 2021). According to our data, BIR1 induction during pathogen attacks can be interpreted as a strategy to regulate PD trafficking. Although not tested in this study, BIR1 effects on gene expression and callose deposition may render PTI less effective against pathogens. Indeed, PTI gene expression and callose formation are usually blocked/inhibited by many defense-suppressing virulence effectors from different types of microbes (Cui *et al.*, 2009; Wang *et al.*, 2021). Viral-encoded proteins such as the potyviral silencing suppressor helper-component proteinase (HC-Pro) or the movement protein (MP) of several tobamoviruses are capable of suppressing callose deposition (Citovsky *et al.*, 1990; Chowdhury *et al.*, 2020). A recent work shows that tobamovirus MPs promote viral cell-to-cell propagation by blocking dsRNA-induced callose deposition (Huang *et al.*, 2023). It is tempting to speculate that BIR1 induction during the viral infection controls PD callose deposition, and perhaps other antiviral defenses, triggered by virus-derived dsRNA. Further research is needed to test these ideas.

The overexpression of BIR1 at a level as that expected when BIR1 gene regulation is lost was referred to as non-physiological. At this level, BIR1 led to cell death-type morphological phenotypes, and the inappropriate overexpression of plant immune genes (Guzman-Benito *et al.*, 2019). Among them, the up-regulation of a significant number of RLP-coding genes is striking. RLPs have been observed to be predominantly expressed in plants undergoing leaf senescence, supporting the idea that BIR1 overexpression accelerates a senescence program (Wang *et al.*, 2008). In Arabidopsis, RLP1, RLP23, RLP30, RLP32 and RLP42 mediate SOBIR1-dependent immune activation, but the role of other RLPs in the perception of pathogen and defense is unknown (Wang *et al.*, 2008; Zhang *et al.*, 2013; Albert *et al.*, 2019).

The overexpression of genes encoding TNL-resistance proteins, but not CNLs, is another
important feature linked to the non-physiological induction of BIR1. TNL-based immunity is
mediated by two distinct signaling modules, EDS1/SAG101/NRG1 and EDS1/PAD4/ADR1
(Sun *et al.*, 2021). Here, we provide genetic evidence that the BIR1 overexpression
morphological phenotypes are partially dependent on EDS1, but not on PAD4 . We propose
that EDS1, perhaps in concert with SAG101, triggers cell death-like phenotypes upon TNL
activation in BIR1 overexpressing plants **(Fig. 7d)**. How TNL receptors are activated when
BIR1 is strongly expressed is a question to resolve. A current model postulates that loss of BIR1
function leads to the activation of hypothetical guarding R proteins which further feed PAD4-
and SOBIR1-dependent resistance pathways (Gao *et al.*, 2009). Our results are coherent with
BIR1 or a yet unknown BIR1/SERK complex being guarded by one or several EDS1-dependent
TNL proteins. Therefore, induction of BIR1 at non-physiological leves may promote TNL-
mediated resistance pathways that require EDS1 when no effectors or elicitors are present.

In addition to EDS1, we demonstrated that SOBIR1 is necessary for BIR1 overexpression
phenotypes. The role of SOBIR1 in the BIR1 pathway is unclear. Although extensive
investigation is required, SOBIR1 may function as a specific co-regulator of the EDS1-
dependent responses triggered by BIR1 overexpression, as part of the same signaling pathway
(Fig. 7d). This idea is inspired by a recent work showing that the cell-surface RLP23 receptor
uses SOBIR1 to form a supramolecular complex with a EDS1/PAD4/ADR1 node in a ligand-
independent manner to mediate defense signaling cascades that are common to both surface
PRR (PTI) and NLR (ETI) receptors (Pruitt *et al.*, 2021b).

The TNL protein CSA1 senses perturbances of BIR3, BAK1 or BIR3/BAK1 complexes
leading to ETI-type autoimmune responses (Schulze *et al.*, 2022). CSA1 did not show
responsiveness to BIR1 overexpression in our assay, but it has been shown to interact with
BIR1 (Schulze *et al.*, 2022). It will be interesting to see if CSA1 also guards BIR1 integrity,
perhaps in cooperation with SOBIR1 and/or other SERKs **(Fig. 7d)**. Finally, phenotypes linked
to BIR1 overexpression were apparently unaffected by the *bak1-5* mutation, which suggests
that other SERK family members can compensate for the loss of BAK1 in PTI/ETI signaling.

We observed that the upregulation of FRK1, WRKY29 and RBOHD due to BIR1
overexpression was compromised by mutations in SOBIR1 and PAD4, but it was independent
of EDS1. Since EDS1, PAD4 and SOBIR1 are important for both cell surface PRR receptor-
and TNL-mediated signaling and resistance (Pruitt *et al.*, 2021b; Tian *et al.*, 2021), we will
need additional experimentation to establish if changes in PTI gene expression involve PTI-

dependent pathways, TNL activation or other unknown mechanisms yet to be determined. Our study also reveals significant alterations in plant hormone gene expression caused by non-physiological levels of BIR1. Interestingly, genes related to the plant hormones ET, JA, CA, and SA, which play a central role in regulating immune responses, remained up-regulated in *eds1-2* BIR1 and *sobir1-12* BIR1 backgrounds, in which phenotypes were partially rescued. This suggests that i) regulation of hormone homeostasis does not require SOBIR1- and/or EDS1-mediated signaling pathways, and that ii) such hormonal imbalance in BIR1 overexpressing plants has little, if any, influence on the ETI-like phenotypes.

The Arabidopsis heterotrimeric G protein AGB1 (SOBIR2) is a common signaling component of PTI mediated by different RLK-PRRs, and functions in parallel with PAD4 downstream of SOBIR1 to regulate cell death and pathogen resistance in *bir1-1* Arabidopsis mutants (Liu *et al.*, 2013; Liang *et al.*, 2016). Similarly, Gγ-proteins AGG1/AGG2 have been described as positive regulators of cell death and constitutive defense phenotypes in *bir1-1* mutants (Liu *et al.*, 2013). AGB1 and AGG2 were up-regulated in BIR1 overexpressing plants. These findings bring up the idea that AGB1 and AGG1/2 could also function downstream of BIR1 to regulate EDS1-mediated cell death when BIR1 integrity is disrupted. Equally important, the BIR1-responsive GPA1 and XLG, in association with Gβγ modules, also have crucial roles in multiple growth and developmental processes (Wang *et al.*, 2023). Thus, it is conceivable that overactivation of these heterotrimeric G proteins due to expression of BIR1 at non-physiological levels may cause indirect effects on plant development. Further studies are needed to test these plausible hypotheses and their applicability to other plants species. In addition, identification of BIR1 interactors shall increase our understanding of the underlying mechanisms and components of the BIR1-mediated regulation of immunity.

Conclusions

Here, we investigated the implications of BIR1 homeostasis on the immune response in Arabidopsis. BIR1 induction at physiological levels interferes with both immune gene expression and PD callose deposition triggered by canonical PTI elicitors. We posit that BIR1 induction may be a pathogen-driven mechanism to modulate plant defenses during infections, helping to maintain a host-pathogen equilibrium. Expression of BIR1 at non-physiological levels results in severe morphological and growth defects as well as transcriptomic changes related to plant immunity, which suggests a dose-dependent regulatory activity of BIR1. We

show that both EDS1 and SOBIR1 are necessary for the ETI-type cell death phenotypes observed when BIR1 is overexpressed, while PAD4 and BAK1 are not. We hypothesize that BIR1, previously thought to represent a negative regulator of plant immunity, is a guarded protein surveilled by unknown TNLs, with SOBIR1 working alongside EDS1 to transduce signals downstream of the TNLs **(Fig. 7d)**.

Acknowledgements

This work was supported by National Research grants RTI2018-096979-B-I00 and PID2021 127982NB-I00 to CL from Ministerio de Ciencia e Innovación (Spain) and UE-FEDER, and grant iLINKA20415 to CL from CSIC (Spain). Contributions of ARS, LEG and MH were supported by grants ERA-NET SusCrop2 / ANR-21-SUSC-0003-01 and ANR-PRC / ANR-21-CE20-0020-01 to MH. Contributions of CH and TN were supported by DFG grant Nu70/17-1 to TN.

Author contributions

CL and IGB conceived and designed the study. CL, IGB, CR and IP performed expression assays and transgenics. CL and CH performed ROS and ethylene assays. CL, ARS and LEG performed callose assays. CL, IGB, MH and TN analyzed the data. CL wrote the article.

Competing interests

None declared

REFERENCES

Albert I, Zhang L, Bemm H, Nurnberger T. 2019. Structure-Function Analysis of Immune Receptor AtRLP23 with Its Ligand nlp20 and Coreceptors AtSOBIR1 and AtBAK1. *Mol Plant Microbe Interact* **32**(8): 1038-1046.

Belkhadir Y, Jaillais Y, Epple P, Balsemao-Pires E, Dangl JL, Chory J. 2012. Brassinosteroids modulate the efficiency of plant immune responses to microbe-associated molecular patterns. *Proc Natl Acad Sci U S A* **109**(1): 297-302.

Bi G, Zhou Z, Wang W, Li L, Rao S, Wu Y, Zhang X, Menke FLH, Chen S, Zhou JM.
2018. Receptor-Like Cytoplasmic Kinases Directly Link Diverse Pattern Recognition
Receptors to the Activation of Mitogen-Activated Protein Kinase Cascades in Arabidopsis.
Plant Cell **30**(7): 1543-1561.

Boudsocq M, Willmann MR, McCormack M, Lee H, Shan L, He P, Bush J, Cheng SH,
Sheen J. 2010. Differential innate immune signalling via Ca(2+) sensor protein kinases.
Nature **464**(7287): 418-422.

Boutrot F, Zipfel C. 2017. Function, Discovery, and Exploitation of Plant Pattern Recognition
Receptors for Broad-Spectrum Disease Resistance. *Annu Rev Phytopathol* **55**: 257-286.

Castro B, Citterico M, Kimura S, Stevens DM, Wrzaczek M, Coaker G. 2021. Stress-
induced reactive oxygen species compartmentalization, perception and signalling. *Nat*
Plants **7**(4): 403-412.

Chen Z. 2001. A superfamily of proteins with novel cysteine-rich repeats. *Plant Physiol* **126**(2):
473-476.

Chowdhury RN, Lasky D, Karki H, Zhang Z, Goyer A, Halterman D, Rakotondrafara
AM. 2020. HCPro Suppression of Callose Deposition Contributes to Strain-Specific
Resistance Against Potato Virus Y. *Phytopathology* **110**(1): 164-173.

Citovsky V, Knorr D, Schuster G, Zambryski P. 1990. The P30 movement protein of tobacco
mosaic virus is a single-strand nucleic acid binding protein. *Cell* **60**(4): 637-647.

Clough SJ, Bent AF. 1998. Floral dip: a simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. *Plant J* **16**(6): 735-743.

Couto D, Zipfel C. 2016. Regulation of pattern recognition receptor signalling in plants. *Nat*
Rev Immunol **16**(9): 537-552.

Cui H, Gobbato E, Kracher B, Qiu J, Bautor J, Parker JE. 2017. A core function of EDS1
with PAD4 is to protect the salicylic acid defense sector in Arabidopsis immunity. *New*
Phytol **213**(4): 1802-1817.

Cui H, Xiang T, Zhou JM. 2009. Plant immunity: a lesson from pathogenic bacterial effector
proteins. *Cell Microbiol* **11**(10): 1453-1461.

Denance N, Sanchez-Vallet A, Goffner D, Molina A. 2013. Disease resistance or growth: the
role of plant hormones in balancing immune responses and fitness costs. *Front Plant Sci* **4**:
155.

Dominguez-Ferreras A, Kiss-Papp M, Jehle AK, Felix G, Chinchilla D. 2015. An Overdose
of the Arabidopsis Coreceptor BRASSINOSTEROID INSENSITIVE1-ASSOCIATED

RECEPTOR KINASE1 or Its Ectodomain Causes Autoimmunity in a SUPPRESSOR OF BIR1-1-Dependent Manner. *Plant Physiol* **168**(3): 1106-1121.

Freh M, Gao JL, Petersen M, Panstruga R. 2022. Plant autoimmunity-fresh insights into an old phenomenon. *Plant Physiology* **188**(3): 1419-1434.

Gao M, Wang X, Wang D, Xu F, Ding X, Zhang Z, Bi D, Cheng YT, Chen S, Li X, et al. 2009. Regulation of cell death and innate immunity by two receptor-like kinases in Arabidopsis. *Cell Host Microbe* **6**(1): 34-44.

Gust AA, Pruitt R, Nurnberger T. 2017. Sensing Danger: Key to Activating Plant Immunity. *Trends Plant Sci* **22**(9): 779-791.

Guzman-Benito I, Donaire L, Amorim-Silva V, Vallarino JG, Esteban A, Wierzbicki AT, Ruiz-Ferrer V, Llave C. 2019. The immune repressor BIR1 contributes to antiviral defense and undergoes transcriptional and post-transcriptional regulation during viral infections. *New Phytol* **224**(1): 421-438.

Halter T, Imkampe J, Mazzotta S, Wierzba M, Postel S, Bucherl C, Kiefer C, Stahl M, Chinchilla D, Wang X, et al. 2014. The leucine-rich repeat receptor kinase BIR2 is a negative regulator of BAK1 in plant immunity. *Curr Biol* **24**(2): 134-143.

Hohmann U, Lau K, Hothorn M. 2017. The Structural Basis of Ligand Perception and Signal Activation by Receptor Kinases. *Annu Rev Plant Biol* **68**: 109-137.

Huang C, Sede AR, Elvira-González L, Yan Y, Rodriguez M, Mutterer J, Boutant E, Shan L, Heinlein M. 2023. dsRNA-induced immunity targets plasmodesmata and is suppressed by viral movement proteins. *Plant Cell* **35**(10): 3845-3869.

Huang W, Wang Y, Li X, Zhang Y. 2020. Biosynthesis and Regulation of Salicylic Acid and N-Hydroxypipecolic Acid in Plant Immunity. *Mol Plant* **13**(1): 31-41.

Imkampe J, Halter T, Huang S, Schulze S, Mazzotta S, Schmidt N, Manstretta R, Postel S, Wierzba M, Yang Y, et al. 2017. The Arabidopsis Leucine-Rich Repeat Receptor Kinase BIR3 Negatively Regulates BAK1 Receptor Complex Formation and Stabilizes BAK1. *Plant Cell* **29**(9): 2285-2303.

Jubic LM, Saile S, Furzer OJ, El Kasmi F, Dangl JL. 2019. Help wanted: helper NLRs and plant immune responses. *Curr Opin Plant Biol* **50**: 82-94.

Kadota Y, Sklenar J, Derbyshire P, Stransfeld L, Asai S, Ntoukakis V, Jones JD, Shirasu K, Menke F, Jones A, et al. 2014. Direct regulation of the NADPH oxidase RBOHD by the PRR-associated kinase BIK1 during plant immunity. *Mol Cell* **54**(1): 43-55.

Kang HG, Fang YW, Singh KB. 1999. A glucocorticoid-inducible transcription system causes severe growth defects in

and induces defense-related genes. *Plant Journal* **20**(1): 127-133.

Korner CJ, Klauser D, Niehl A, Dominguez-Ferreras A, Chinchilla D, Boller T, Heinlein
M, Hann DR. 2013. The immunity regulator BAK1 contributes to resistance against diverse
RNA viruses. *Mol Plant Microbe Interact* **26**(11): 1271-1280.

Li L, Li M, Yu L, Zhou Z, Liang X, Liu Z, Cai G, Gao L, Zhang X, Wang Y, et al. 2014.
The FLS2-associated kinase BIK1 directly phosphorylates the NADPH oxidase RbohD to
control plant immunity. *Cell Host Microbe* **15**(3): 329-338.

Li P, Lu YJ, Chen H, Day B. 2020. The Lifecycle of the Plant Immune System. *CRC Crit Rev*
Plant Sci **39**(1): 72-100.

Liang X, Ding P, Lian K, Wang J, Ma M, Li L, Li L, Li M, Zhang X, Chen S, et al. 2016.
Arabidopsis heterotrimeric G proteins regulate immunity by directly coupling to the FLS2
receptor. *Elife* **5**: e13568.

Liang X, Zhou JM. 2018. Receptor-Like Cytoplasmic Kinases: Central Players in Plant
Receptor Kinase-Mediated Signaling. *Annu Rev Plant Biol* **69**: 267-299.

Liu J, Ding P, Sun T, Nitta Y, Dong O, Huang X, Yang W, Li X, Botella JR, Zhang Y.
2013. Heterotrimeric G proteins serve as a converging point in plant defense signaling
activated by multiple receptor-like kinases. *Plant Physiol* **161**(4): 2146-2158.

Liu Y, Huang X, Li M, He P, Zhang Y. 2016. Loss-of-function of Arabidopsis receptor-like
kinase BIR1 activates cell death and defense responses mediated by BAK1 and SOBIR1.
New Phytol **212**(3): 637-645.

Lu H. 2009. Dissection of salicylic acid-mediated defense signaling networks. *Plant Signal*
Behav **4**(8): 713-717.

Luna E, Pastor V, Robert J, Flors V, Mauch-Mani B, Ton J. 2011. Callose deposition: a
multifaceted plant defense response. *Mol Plant Microbe Interact* **24**(2): 183-193.

Lv Y, Yang N, Wu J, Liu Z, Pan L, Lv S, Wang G. 2016. New insights into receptor-like
protein functions in Arabidopsis. *Plant Signal Behav* **11**(7): e1197469.

Mithoe SC, Menke FL. 2018. Regulation of pattern recognition receptor signalling by
phosphorylation and ubiquitination. *Curr Opin Plant Biol* **45**(Pt A): 162-170.

Nawrath C, Heck S, Parinthawong N, Metraux JP. 2002. EDS5, an essential component of
salicylic acid-dependent signaling for disease resistance in Arabidopsis, is a member of the
MATE transporter family. *Plant Cell* **14**(1): 275-286.

Ngou BPM, Ahn HK, Ding P, Jones JDG. 2021. Mutual potentiation of plant immunity by
cell-surface and intracellular receptors. *Nature* **592**(7852): 110-115.

Nishad R, Ahmed T, Rahman VJ, Kareem A. 2020. Modulation of Plant Defense System in
Response to Microbial Interactions. *Front Microbiol* **11**: 1298.

Parker JE, Holub EB, Frost LN, Falk A, Gunn ND, Daniels MJ. 1996. Characterization of
eds1, a mutation in Arabidopsis suppressing resistance to Peronospora parasitica specified
by several different RPP genes. *Plant Cell* **8**(11): 2033-2046.

Pruitt RN, Gust AA, Nurnberger T. 2021a. Plant immunity unified. *Nat Plants* **7**(4): 382-
383.

Pruitt RN, Locci F, Wanke F, Zhang L, Saile SC, Joe A, Karelina D, Hua C, Frohlich K,
Wan WL, et al. 2021b. The EDS1-PAD4-ADR1 node mediates Arabidopsis pattern-
triggered immunity. *Nature* **598**(7881): 495-499.

Quezada EH, Garcia GX, Arthikala MK, Melappa G, Lara M, Nanjareddy K. 2019.
Cysteine-Rich Receptor-Like Kinase Gene Family Identification in the Phaseolus Genome
and Comparative Analysis of Their Expression Profiles Specific to Mycorrhizal and
Rhizobial Symbiosis. *Genes (Basel)* **10**(1).

Rao S, Zhou Z, Miao P, Bi G, Hu M, Wu Y, Feng F, Zhang X, Zhou JM. 2018. Roles of
Receptor-Like Cytoplasmic Kinase VII Members in Pattern-Triggered Immune Signaling.
Plant Physiol **177**(4): 1679-1690.

Rodriguez E, El Ghoul H, Mundy J, Petersen M. 2016. Making sense of plant autoimmunity
and 'negative regulators'. *FEBS J* **283**(8): 1385-1391.

Schulze S, Yu L, Hua C, Zhang L, Kolb D, Weber H, Ehinger A, Saile SC, Stahl M, Franz-
Wachtel M, et al. 2022. The Arabidopsis TIR-NBS-LRR protein CSA1 guards BAK1-BIR3
homeostasis and mediates convergence of pattern- and effector-induced immune responses.
Cell Host Microbe **30**(12): 1717-1731 e1716.

Schwessinger B, Roux M, Kadota Y, Ntoukakis V, Sklenar J, Jones A, Zipfel C. 2011.
Phosphorylation-dependent differential regulation of plant growth, cell death, and innate
immunity by the regulatory receptor-like kinase BAK1. *PLoS Genet* **7**(4): e1002046.

Smakowska-Luzan E, Mott GA, Parys K, Stegmann M, Howton TC, Layeghifard M,
Neuhold J, Lehner A, Kong J, Grunwald K, et al. 2018. An extracellular network of
Arabidopsis leucine-rich repeat receptor kinases. *Nature* **553**(7688): 342-346.

Sun X, Lapin D, Feehan JM, Stolze SC, Kramer K, Dongus JA, Rzemieniewski J,
Blanvillain-Baufume S, Harzen A, Bautor J, et al. 2021. Pathogen effector recognition-
dependent association of NRG1 with EDS1 and SAG101 in TNL receptor immunity. *Nat*
Commun **12**(1): 3335.

Swiderski MR, Birker D, Jones JD. 2009. The TIR domain of TIR-NB-LRR resistance

proteins is a signaling domain involved in cell death induction. *Mol Plant Microbe Interact*

22(2): 157-165.

Tang D, Wang G, Zhou JM. 2017. Receptor Kinases in Plant-Pathogen Interactions: More

Than Pattern Recognition. *Plant Cell* **29**(4): 618-637.

Tian H, Wu Z, Chen S, Ao K, Huang W, Yaghmaiean H, Sun T, Xu F, Zhang Y, Wang S,

et al. 2021. Activation of TIR signalling boosts pattern-triggered immunity. *Nature*

598(7881): 500-503.

Tsuda K, Mine A, Bethke G, Igarashi D, Botanga CJ, Tsuda Y, Glazebrook J, Sato M,

Katagiri F. 2013. Dual regulation of gene expression mediated by extended MAPK

activation and salicylic acid contributes to robust innate immunity in Arabidopsis thaliana.

PLoS Genet **9**(12): e1004015.

van Wersch S, Tian L, Hoy R, Li X. 2020. Plant NLRs: The Whistleblowers of Plant

Immunity. *Plant Commun* **1**(1): 100016.

Wan WL, Zhang L, Pruitt R, Zaidem M, Brugman R, Ma X, Krol E, Perraki A, Kilian J,

Grossmann G, et al. 2019. Comparing Arabidopsis receptor kinase and receptor protein-

mediated immune signaling reveals BIK1-dependent differences. *New Phytol* **221**(4): 2080-

2095.

Wang G, Ellendorff U, Kemp B, Mansfield JW, Forsyth A, Mitchell K, Bastas K, Liu CM,

Woods-Tor A, Zipfel C, et al. 2008. A genome-wide functional investigation into the roles

of receptor-like proteins in Arabidopsis. *Plant Physiol* **147**(2): 503-517.

Wang J, Chai J. 2020. Structural Insights into the Plant Immune Receptors PRRs and NLRs.

Plant Physiol **182**(4): 1566-1581.

Wang Y, Li X, Fan B, Zhu C, Chen Z. 2021. Regulation and Function of Defense-Related

Callose Deposition in Plants. *Int J Mol Sci* **22**(5).

Wang Y, Zhang H, Wang P, Zhong H, Liu W, Zhang S, Xiong L, Wu Y, Xia Y. 2023.

Arabidopsis EXTRA-LARGE G PROTEIN 1 (XLG1) functions together with XLG2 and

XLG3 in PAMP-triggered MAPK activation and immunity. *J Integr Plant Biol* **65**(3): 825-

837.

Withers J, Dong X. 2017. Post-translational regulation of plant immunity. *Curr Opin Plant*

Biol **38**: 124-132.

Wu Y, Gao Y, Zhan Y, Kui H, Liu H, Yan L, Kemmerling B, Zhou JM, He K, Li J. 2020.

Loss of the common immune coreceptor BAK1 leads to NLR-dependent cell death. *Proc*

Natl Acad Sci U S A **117**(43): 27044-27053.

Yeh YH, Chang YH, Huang PY, Huang JB, Zimmerli L. 2015. Enhanced Arabidopsis

pattern-triggered immunity by overexpression of cysteine-rich receptor-like kinases. *Front*

Plant Sci **6**: 322.

Yip Delormel T, Boudsocq M. 2019. Properties and functions of calcium-dependent protein

kinases and their relatives in Arabidopsis thaliana. *New Phytol* **224**(2): 585-604.

Yu TY, Sun MK, Liang LK. 2021. Receptors in the Induction of the Plant Innate Immunity.

Mol Plant Microbe Interact **34**(6): 587-601.

Yuan M, Cai B, Xin XF. 2023. Plant immune receptor pathways as a united front against

pathogens. *PLoS Pathog* **19**(2): e1011106.

Yuan M, Jiang Z, Bi G, Nomura K, Liu M, Wang Y, Cai B, Zhou JM, He SY, Xin XF.

2021a. Pattern-recognition receptors are required for NLR-mediated plant immunity. *Nature*

592(7852): 105-109.

Yuan M, Ngou BPM, Ding P, Xin XF. 2021b. PTI-ETI crosstalk: an integrative view of plant

immunity. *Curr Opin Plant Biol* **62**: 102030.

Zhang W, Fraiture M, Kolb D, Loffelhardt B, Desaki Y, Boutrot FF, Tor M, Zipfel C,

Gust AA, Brunner F. 2013. Arabidopsis receptor-like protein30 and receptor-like kinase

suppressor of BIR1-1/EVERSHED mediate innate immunity to necrotrophic fungi. *Plant*

Cell **25**(10): 4227-4241.

Zhong CL, Zhang C, Liu JZ. 2019. Heterotrimeric G protein signaling in plant immunity. *J*

Exp Bot **70**(4): 1109-1118.

Zhou JM, Zhang Y. 2020. Plant Immunity: Danger Perception and Signaling. *Cell* **181**(5):

978-989.

Zhou N, Tootle TL, Tsui F, Klessig DF, Glazebrook J. 1998. PAD4 functions upstream from

salicylic acid to control defense responses in Arabidopsis. *Plant Cell* **10**(6): 1021-1030.

 Figure legends

Fig. 1. Transcriptional reprogramming associated with BAK1-INTERACTING RECEPTOR-
LIKE KINASE 1 (BIR1) induction in Arabidopsis. (a) Morphology of representative BIR1
transgenic Arabidopsis plants (line 9, L9) under the control of a dexamethasone (DEX)-
inducible 35S promoter used for RNA-Seq experiments. Samples were collected after 4 or 12
days of 30 μM DEX treatment. H_2O-treated plants were used as controls. Relative expression
of the BIR1-coding gene based on the number of Fragments Per Kilobase of transcript sequence
per Millions of base-pairs sequenced (FPKM) (top) and Western blot using anti-mCherry
antibody (bottom) in samples used for RNA-Seq. Significant differences between DEX- and
mock-treated samples were analyzed using two-way ANOVA followed by Sidak´s multiple
comparison test; ***, adjusted p-value < 0.001. (b) Principal component analysis (PCA) of
RNA-Seq data. The PCA was performed using normalized RNA-Seq data of differentially
expressed genes (DEGs) between BIR1 L9 and wild-type (WT) plants after 4 and 12 days of
DEX treatment (DeL9_vs DeWT_). Each biological replicate is represented in the score plot.
The variance explained by each component (%) is given in parentheses. The sample
(DeL9_12b) with differential expression within the DEX-treated L9 group is indicated with
black arrows in (a) and (b). (c) Number of up-or down-regulated transcripts, with a |$\log_2$ (Fold-
Change)| ≥1 and adjusted p-value ≤0.01, in each pairwise comparison is indicated. (d)
Scatterplot of selected gene ontology (GO) terms of DEGs. GO terms between WT and BIR1
L9 plants at 4 and 12 days after DEX-treatment (DeL9_ vs DeWT_) are shown. Dot size
represents the count of different genes and the color indicates significance of the term
enrichment. GO terms with adjusted p-value < 0.05 are significantly enriched. The kinase term
(indicated by an arrow) was exclusively assigned to BIR1 L9 plants. WT, wild-type Col-0; L9,
BIR1 L9 transgenic plants; De, DEX-treated plants; Mo, Mock-treated plants; _4, 4 days of
treatment; _12, 12 days of treatment.

Fig. 2. Effect of BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) induction on
immune responses triggered by flg22 and nlp20 treatments. (a) Over time production of reactive
oxygen species (ROS) in dexamethasone (DEX)-treated wild-type (WT) and BIR1 L6 and L9
transgenic plants after elicitation of Arabidopsis leaf disks with 1 μM flg22 (left), 1 μM nlp20
(middle) or H_2O (right). Bars represent means ± SD of relative fluorescence units. (b) Ethylene
accumulation in BIR1 L6 and L9 transgenic plants after treatment with H_2O (mock), flg22 and
nlp20 at 1 μM or 100 nM concentration as indicated. Bars represent average values ± SD.

Within each elicitor/mock treatment, differences between DEX-treated (DEX+) and controls

(DEX-) were analyzed by two-way ANOVA followed by Sidak´s multiple comparison test. (c)

RT-qPCR analysis of the genes encoding the PTI-markers FLG22-INDUCED RECEPTOR-

LIKE KINASE 1 (FRK1) and PATHOGENESIS RELATED 1 (PR1) at 45 min after treatment

with H_2O (mock) or 1 µM flg22, and determined in DEX-treated (DEX+) versus non-treated

(DEX-) samples. (d) Immunoprecipitation of mCherry-BIR1 fusion proteins, indicated with an

arrowhead, from samples of BIR1 L6 and L9 plants used in (a) and (b) (left), and RT-qPCR

analysis of BIR1-coding transcripts in DEX-treated (DEX+) samples of BIR1 L6 and L9 plants

used in (c) (right). In all cases, RT-qPCR values from biologically independent samples were

normalized to the CBP20 internal control, and related to H_2O (mock)-treated plants, set to a

value of 1. Data are means ± SD analyzed by Kruskal-Wallis test; *, adjusted p-value <0.05.

Experiments were repeated three times with similar results.

Fig. 3. BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) induction inhibits flg22-

and poly(I:C)-triggered callose accumulation at plasmodesmata (PD). Callose spots at PD were

visualized upon aniline blue staining of epidermal cells in response to bacterial flg22 (a) or

poly(I:C) (b) in BIR1 L9 transgenic Arabidopsis plants. Plants were previously sprayed with

30 µM dexamethasone (DEX) for three days to elicit BIR1 expression. Mock-treated plants

were used as non BIR1-expressing controls (top panels). Photographs were taken 30 minutes

after treatment with H_2O (control) or 1 µM flg22 or 50 ng/µl of poly(I:C). BIR1-mCherry

fluorescence in DEX-treated BIR1 L9 plants was confirmed within the same images. Inlays

show enlargements of the areas within the dashed boxes. Scale bar, 10 µm. (c) Relative PD

callose content in flg22/poly(I:C)-treated and H_2O (control)-treated leaves from BIR1 L9 plants

previously treated with or without DEX as indicated. d) Callose content at PD in control (DEX

-) and DEX-treated wild-type (WT) plants in response to H_2O and poly (I:C) is shown. In all

cases, black dots represent the mean values of callose intensities in individual PD (>100)

measured in three leaf discs taken from at least three independent biological replicates per each

treatment. Significant differences between samples were analyzed by Kruskal-Wallis test or by

one-way ANOVA followed by Tukey's multiple comparison test; ****, adjusted p-value

<0.0001, ***, <0.001, **, <0.01. Experiments were repeated twice with similar results.

Fig. 4. BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpression leads to the transcriptional activation of pattern-triggered immunity (PTI)-associated co-receptors and effector-triggered immunity (ETI) regulators. (a) Validation of the RNA-Seq results. Expression of markers encoded by selected BIR1-responsive differentially expressed genes (DEGs) after 12 days of dexamethasone (DEX) treatment based on fold-change values ($\log_2$ DeL9_12/DeWT_12 and $\log_2$ DeL9_12/MoL9_12 as indicated) was confirmed by RT-qPCR. Values from biologically independent samples were normalized to the CBP20 internal control, and related to untreated (DEX-) plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test; *, $p < 0.05$. (b) Relative expression of selected genes encoding PTI/ETI-markers based on the number of Fragments Per Kilobase of transcript sequence per Millions of base-pairs sequenced (FPKM) in each of the three independent samples per condition used for RNA-Seq. Actin expression is shown as control. Note that the expression levels of each individual replicate within the DEX-treated BIR1 L9_12 group mirror the abundance of BIR1 transcripts in each replicate. DeL9_12b is indicated by a red arrow. Significant differences between mock-treated and DEX-treated samples were analyzed using two-way ANOVA followed by Sidak´s multiple comparison test; ****, adjusted p-value < 0.0001, *** < 0.001, **< 0.01. WT, wild-type Col-0; L9, BIR1 L9 transgenics; _4d, 4 days of treatment; _12d, 12 days of treatment. (c) Western blot analysis of SUPPRESSOR OF BIR1-1 (SOBIR1) and PATHOGENESIS RELATED 1 (PR1) using specific antibodies in two independent sets of BIR1 L9 samples. Plants were mock-treated (-) or DEX-treated (+) for 12 days. The presence (+) or absence (-) of morphological phenotypes (Ph) in each sample is indicated. Protein loading control by Ponceau S is shown.

Fig. 5. Phenotypes caused by BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpression are partially rescued in *sobir1-12* mutants, but not in a *bak1-5* mutant. Morphology of transgenic *sobir1-12* BIR1 L6 and *bak1*-5 BIR1 L10 mutant plants expressing BIR1 under the control of a dexamethasone (DEX)-inducible 35S promoter. Transgenic BIR1 L9 plants in the wild-type (WT) Col-0 background are shown as a reference. Plants were grown for three weeks on soil and treated with 30 μM dexamethasone (DEX) to induce BIR1 transcription or mock-treated with H_2O. Non-transgenic WT Col-0 plants are shown as controls. Plants were photographed after 12 days of the first DEX/ H_2O application. Images from independent transgenic lines are shown in Fig. S7. RT-qPCR analysis of *BIR1* transcripts in BIR1 L9 plants are shown. b) RT-qPCR analysis of BIR1 and ENHANCED DISEASE

SUSCEPTIBILITY 1 (EDS1) transcripts in *sobir1-12* BIR1 L6 and *bak1-5* BIR1 L10 plants. c) Western blot analysis of BIR1 and EDS1 proteins with specific antibodies in WT BIR1 L9 and two independent lines of *sobir1-12* BIR1 (L6, L8) and *bak1-5* BIR1 (L4, L10). Plants were mock-treated (-) or DEX-treated (+) for 12 days as indicated. The presence (+) or absence (-) of morphological phenotypes (Ph) in each sample is indicated. Protein loading control by Ponceau S is shown. RT-qPCR analysis of genes encoding the PTI-markers RESPIRATORY BURST OXIDASE HOMOLOG PROTEIN D (RBOHD), FLG22-INDUCED RECEPTOR-LIKE KINASE 1 (FRK1) or WRKY DNA-BINDING PROTEIN 29 (WRKY29) in *sobir1-12* BIR1 L6 (d) and *bak1-5* BIR1 L10 (e) plants. RT-qPCR analysis of genes encoding representative markers of the salicylic acid PATHOGENESIS RELATED 1 (PR1), jasmonic acid PLANT DEFENSIN 1.2 (PDF1.2), ethylene PR4 and camalexin [PHYTOALEXIN DEFICIENT 3 (PAD3), CYTOCHROME P450 MONOOXYGENASE 79B2 (CYP79B2)] pathways in *sobir1-12* BIR1 L6 (f) and *bak1-5* BIR1 L10 (g) plants. In all cases, the same set of replicates from each genotype was used to measure all the genes shown in the graphs. Transcript accumulation was determined in mock-treated (DEX-) versus DEX-treated (DEX+) after 12 days of treatment. Values from biologically independent samples were normalized to the CBP20 internal control, and related to mock-treated plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test; *, p <0.05. Experiments were repeated at least twice with similar results.

Fig. 6. Phenotypes caused by BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpression are partially rescued in *eds1-2* mutants, but not in *pad4-1* mutants. (a) Morphology of transgenic *eds1-2* BIR1 L1b and *pad4-1* BIR1 L0 mutant plants expressing BIR1 under the control of a dexamethasone (DEX)-inducible 35S promoter. Plants were grown for three weeks on soil and treated with 30 μM DEX to induce BIR1 transcription or mock-treated with H_2O. Plants were photographed after 12 days of the first DEX/ H_2O application. Images from independent transgenic lines are shown in Fig. S7. RT-qPCR analysis of BIR1-coding transcripts in *eds1-2* BIR1 L1b and *pad4-1* BIR1 L0 mutant lines are shown. RT-qPCR analysis of genes encoding the PTI-markers RESPIRATORY BURST OXIDASE HOMOLOG PROTEIN D (RBOHD) and FLG22-INDUCED RECEPTOR-LIKE KINASE 1 (FRK1) in *eds1-2* BIR1 L1b (b) and *pad4-1* BIR1 L0 (c) plants. RT-qPCR analysis of genes encoding representative markers of the salicylic acid PATHOGENESIS RELATED 1 (PR1), jasmonic acid PLANT DEFENSIN 1.2 (PDF1.2), ethylene PR4 and camalexin [PHYTOALEXIN

DEFICIENT 3 (PAD3), CYTOCHROME P450 MONOOXYGENASE 79B2 (CYP79B2)] pathways in *eds1-2* BIR1 L1b (d) and *pad4-1* BIR1 L0 (e) plants. In all cases, the same set of replicates from each genotype was used to measure all the genes shown in the graphs. Transcript accumulation was determined in mock-treated (DEX-) versus DEX-treated (DEX+) after 12 days of treatment. Values from biologically independent samples were normalized to the CBP20 internal control, and related to mock-treated plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test; *, $p < 0.05$. Experiments were repeated at least twice with similar results.

Fig. 7. Phenotypes caused by BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpression are independent of salicylic acid (SA)-mediated signaling pathways. (a) Morphology of transgenic *sid2-2* BIR1 (L5) and *eds5-3* BIR1 (L10) mutant plants expressing BIR1 under the control of a dexamethasone (DEX)-inducible 35S promoter. Plants were grown for three weeks on soil and treated with 30 μM DEX to induce BIR1 transcription or mock-treated with H_2O. Plants were photographed after 12 days of the first DEX/ H_2O application. Images from independent transgenic lines are shown in Fig. S7a, b. RT-qPCR analysis of genes encoding BIR1 and representative markers of the salicylic acid PATHOGENESIS RELATED 1 (PR1), jasmonic acid PLANT DEFENSIN 1.2 (PDF1.2), ethylene PR4 and camalexin [PHYTOALEXIN DEFICIENT 3 (PAD3), CYTOCHROME P450 MONOOXYGENASE 79B2 (CYP79B2)] pathways in *sid2-2* BIR1 (L5) (b) and *eds5-3* BIR1 (L10) (c) mutant lines. In all cases, the same set of replicates from each genotype was used to measure all the genes shown in the graphs. Transcript accumulation was determined in mock-treated (DEX-) versus DEX-treated (DEX+) after 12 days of treatment. Values from biologically independent samples were normalized to the CBP20 internal control, and related to mock-treated plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test; *, $p < 0.05$. Experiments were repeated at least twice with similar results. (d) Tentative working model for BIR1 homeostasis. SA-mediated transcription of BIR1 is regulated by RNA-directed DNA methylation (RdDM) and post-transcriptional RNA silencing (PTGS) (Guzman-Benito *et al.*, 2019). In these conditions, BIR1 induction interferes with pattern-triggered immunity (PTI) gene expression and plasmodesmata (PD) callose deposition triggered by various elicitors (flg22, nlp20, dsRNA). Loss of BIR1 regulation leads to constitutive BIR1 induction at non-physiological levels that cause SA-independent up-regulation of immune and hormone genes. BIR1 overexpression is sensed by one or several TOLL/INTERLEUKIN RECEPTOR 1 (TIR1)-

977   containing nucleotide-binding domain leucine-rich repeat receptors (TNLs), of which

978   CONSTITUTIVE SHADE AVOIDANCE 1 (CSA1) is a plausible candidate, to promote

979   PAD4-independent effector triggered immunity (ETI)-type cell death that requires

980   ENHANCED DISEASE SUSCEPTIBILITY 1 (EDS1) and SUPPRESSOR OF BIR1-1

981   (SOBIR1).

Supporting information

Additional supporting information may be found in the online version of this article

Supporting methods

Methods S1. RNA and protein analyses.

Methods S2. Gene expression analyses.

Methods S3. Analyses of reactive oxygen species (ROS), ethylene, plasmodesmata callose deposition and dexamethasone (DEX) treatments.

Supporting figures

Fig. S1. Transcriptomic profiling of BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) expressing transgenic lines.

Fig. S2. Transcriptomic profiling of dexamethasone (DEX)-treated wild-type Col-0 plants.

Fig. S3. Enrichment analysis of Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways using RNA-Seq data.

Fig. S4. Differential expression of receptor-like protein (RLP)-coding genes in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants.

Fig. S5. Differential expression of cysteine-rich receptor-like kinase (CRK)-coding genes in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants.

Fig. S6. Differential expression of cell-surface pattern recognition receptors (PRR)-coding genes in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants.

Fig. S7. Differential expression of calcium-dependent protein kinases (CPK)- and HETEROTRIMERIC G PROTEIN-coding genes in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants.

Fig. S8. Morphological and growth defects in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants of different immune mutant genetic backgrounds.

Supporting tables

Table S1. List of primers

Table S2. List of expressed genes.

Table S3. List of differentially expressed genes (DEG. padjust value < 0.05).

Table S4. List of differentially expressed genes (DEG. |Log2fold change > 1| padjust value <
0.01).

Table S5. Enrichment analysis of Gene Ontology (GO) terms.

Table S6. Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis.

Table S7. List of differentially expressed genes (DEG. padjust value < 0.05) within functional
categories.

Table S8. Statistical analysis of enrichment of functional gene groups in plants overexpressing
BIR1 after 12 days of DEX treatment relative to non-expressing controls using Two-tailed
Fisher's exact tests.

1022

1023

Fig. 1. Transcriptional reprogramming associated with BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) induction in Arabidopsis. (a) Morphology of representative BIR1 transgenic Arabidopsis plants (line 9, L9) under the control of a dexamethasone (DEX)-inducible 35S promoter used for RNA-Seq experiments. Samples were collected after 4 or 12 days of 30 μM DEX treatment. H_2O-treated plants were used as controls. Relative expression of the BIR1-coding gene based on the number of Fragments Per Kilobase of transcript sequence per Millions of base-pairs sequenced (FPKM) (top) and Western blot using anti-mCherry antibody (bottom) in samples used for RNA-Seq. Significant differences between DEX- and mock-treated samples were analyzed using two-way ANOVA followed by Sidak´s multiple comparison test; ***, adjusted p-value < 0.001. (b) Principal component analysis (PCA) of RNA-Seq data. The PCA was performed using normalized RNA-Seq data of differentially expressed genes (DEGs) between BIR1 L9 and wild-type (WT) plants after 4 and 12 days of DEX treatment (DeL9_vs DeWT_). Each biological replicate is represented in the score plot. The variance explained by each component (%) is given in parentheses. The sample (DeL9_12b) with differential expression within the DEX-treated L9 group is indicated with black arrows in (a) and (b). (c) Number of up-or down-regulated transcripts, with a |$\log_2$(Fold-Change)| ≥1 and adjusted p-value ≤0.01, in each pairwise comparison is indicated. (d) Scatterplot of selected gene ontology (GO) terms of DEGs. GO terms between WT and BIR1 L9 plants at 4 and 12 days after DEX-treatment (DeL9_ vs DeWT_) are shown. Dot size represents the count of different genes and the color indicates significance of the term enrichment. GO terms with adjusted p-value < 0.05 are significantly enriched. The kinase term (indicated by an arrow) was exclusively assigned to BIR1 L9 plants. WT, wild-type Col-0; L9, BIR1 L9 transgenic plants; De, DEX-treated plants; Mo, Mock-treated plants; _4, 4 days of treatment; _12, 12 days of treatment.

Fig. 2. Effect of BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) induction on immune responses triggered by flg22 and nlp20 treatments. (a) Over time production of reactive oxygen species (ROS) in dexamethasone (DEX)-treated wild-type (WT) and BIR1 L6 and L9 transgenic plants after elicitation of Arabidopsis leaf disks with 1 μM flg22 (left), 1 μM nlp20 (middle) or H_2O (right). Bars represent means ± SD of relative fluorescence units. (b) Ethylene accumulation in BIR1 L6 and L9 transgenic plants after treatment with H_2O (mock), flg22 and nlp20 at 1 μM or 100 nM concentration as indicated. Bars represent average values ± SD. Within each elicitor/mock treatment, differences between DEX-treated (DEX+) and controls (DEX-) were analyzed by two-way ANOVA followed by Sidak´s multiple comparison test. (c) RT-qPCR analysis of the genes encoding the PTI-markers FLG22-INDUCED RECEPTOR-LIKE KINASE 1 (FRK1) and PATHOGENESIS RELATED 1 (PR1) at 45 min after treatment with H_2O (mock) or 1 μM flg22, and determined in DEX-treated (DEX+) versus non-treated (DEX-) samples. (d) Immunoprecipitation of mCherry-BIR1 fusion proteins, indicated with an arrowhead, from samples of BIR1 L6 and L9 plants used in (a) and (b) (left), and RT-qPCR analysis of BIR1-coding transcripts in DEX-treated (DEX+) samples of BIR1 L6 and L9 plants used in (c) (right). In all cases, RT-qPCR values from biologically independent samples were normalized to the CBP20 internal control, and related to H_2O (mock)-treated plants, set to a value of 1. Data are means ± SD analyzed by Kruskal-Wallis test; *, adjusted p-value <0.05. Experiments were repeated three times with similar results.

Fig. 3. BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) induction inhibits flg22- and poly(I:C)-triggered callose accumulation at plasmodesmata (PD). Callose spots at PD were visualized upon aniline blue staining of epidermal cells in response to bacterial flg22 (a) or poly(I:C) (b) in BIR1 L9 transgenic Arabidopsis plants. Plants were previously sprayed with 30 μM dexamethasone (DEX) for three days to elicit BIR1 expression. Mock-treated plants were used as non BIR1-expressing controls (top panels). Photographs were taken 30 minutes after treatment with H_2O (control) or 1 μM flg22 or 50 ng/μl of poly(I:C). BIR1-mCherry fluorescence in DEX-treated BIR1 L9 plants was confirmed within the same images. Inlays show enlargements of the areas within the dashed boxes. Scale bar, 10 μm. (c) Relative PD callose content in flg22/poly(I:C)-treated and H_2O (control)-treated leaves from BIR1 L9 plants previously treated with or without DEX as indicated. d) Callose content at PD in control (DEX -) and DEX-treated wild-type (WT) plants in response to H_2O and poly (I:C) is shown. In all cases, black dots represent the mean values of callose intensities in individual PD (>100) measured in three leaf discs taken from at least three independent biological replicates per each treatment. Significant differences between samples were analyzed by Kruskal-Wallis test or by one-way ANOVA followed by Tukey's multiple comparison test; ****, adjusted *p*-value <0.0001, ***, <0.001, **, <0.01. Experiments were repeated twice with similar results.

Fig. 4. BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpression leads to the transcriptional activation of pattern-triggered immunity (PTI)-associated co-receptors and effector-triggered immunity (ETI) regulators. (a) Validation of the RNA-Seq results. Expression of markers encoded by selected BIR1-responsive differentially expressed genes (DEGs) after 12 days of dexamethasone (DEX) treatment based on fold-change values (log_2 DeL9_12/DeWT_12 and log_2 DeL9_12/MoL9_12 as indicated) was confirmed by RT-qPCR. Values from biologically independent samples were normalized to the CBP20 internal control, and related to untreated (DEX-) plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test; *, p <0.05. (b) Relative expression of selected genes encoding PTI/ETI-markers based on the number of Fragments Per Kilobase of transcript sequence per Millions of base-pairs sequenced (FPKM) in each of the three independent samples per condition used for RNA-Seq. Actin expression is shown as control. Note that the expression levels of each individual replicate within the DEX-treated BIR1 L9_12 group mirror the abundance of BIR1 transcripts in each replicate. DeL9_12b is indicated by a red arrow. Significant differences between mock-treated and DEX-treated samples were analyzed using two-way ANOVA followed by Sidak´s multiple comparison test; ****, adjusted p-value < 0.0001, *** < 0.001, ** < 0.01. WT, wild-type Col-0; L9, BIR1 L9 transgenics; _4d, 4 days of treatment; _12d, 12 days of treatment. (c) Western blot analysis of SUPPRESSOR OF BIR1-1 (SOBIR1) and PATHOGENESIS RELATED 1 (PR1) using specific antibodies in two independent sets of BIR1 L9 samples. Plants were mock-treated (-) or DEX-treated (+) for 12 days. The presence (+) or absence (-) of morphological phenotypes (Ph) in each sample is indicated. Protein loading control by Ponceau S is shown.

Fig. 5. Phenotypes caused by BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpression are partially rescued in *sobir1-12* mutants, but not in a *bak1-5* mutant. Morphology of transgenic *sobir1-12* BIR1 L6 and *bak1-5* BIR1 L10 mutant plants expressing BIR1 under the control of a dexamethasone (DEX)-inducible 35S promoter. Transgenic BIR1 L9 plants in the wild-type (WT) Col-0 background are shown as a reference. Plants were grown for three weeks on soil and treated with 30 µM dexamethasone (DEX) to induce BIR1 transcription or mock-treated with H_2O. Non-transgenic WT Col-0 plants are shown as controls. Plants were photographed after 12 days of the first DEX/ H_2O application. Images from independent transgenic lines are shown in Fig. S7. RT-qPCR analysis of *BIR1* transcripts in BIR1 L9 plants are shown. b) RT-qPCR analysis of BIR1 and ENHANCED DISEASE SUSCEPTIBILITY 1 (EDS1) transcripts in *sobir1-12* BIR1 L6 and *bak1-5* BIR1 L10 plants. c) Western blot analysis of BIR1 and EDS1 proteins with specific antibodies in WT BIR1 L9 and two independent lines of *sobir1-12* BIR1 (L6, L8) and *bak1-5* BIR1 (L4, L10). Plants were mock-treated (-) or DEX-treated (+) for 12 days as indicated. The presence (+) or absence (-) of morphological phenotypes (Ph) in each sample is indicated. Protein loading control by Ponceau S is shown. RT-qPCR analysis of genes encoding the PTI-markers RESPIRATORY BURST OXIDASE HOMOLOG PROTEIN D (RBOHD), FLG22-INDUCED RECEPTOR-LIKE KINASE 1 (FRK1) or WRKY DNA-BINDING PROTEIN 29 (WRKY29) in *sobir1-12* BIR1 L6 (d) and *bak1-5* BIR1 L10 (e) plants. RT-qPCR analysis of genes encoding representative markers of the salicylic acid PATHOGENESIS RELATED 1 (PR1), jasmonic acid PLANT DEFENSIN 1.2 (PDF1.2), ethylene PR4 and camalexin [PHYTOALEXIN DEFICIENT 3 (PAD3)*, CYTOCHROME P450 MONOOXYGENASE 79B2 (CYP79B2)*] pathways in *sobir1-12* BIR1 L6 (f) and *bak1-5* BIR1 L10 (g) plants. In all cases, the same set of replicates from each genotype was used to measure all the genes shown in the graphs. Transcript accumulation was determined in mock-treated (DEX-) versus DEX-treated (DEX+) after 12 days of treatment. Values from biologically independent samples were normalized to the CBP20 internal control, and related to mock-treated plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test; *, *p* <0.05. Experiments were repeated at least twice with similar results.

Fig. 6. Phenotypes caused by BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpression are partially rescued in *eds1-2* mutants, but not in *pad4-1* mutants. (a) Morphology of transgenic *eds1-2* BIR1 L1b and *pad4-1* BIR1 L0 mutant plants expressing BIR1 under the control of a dexamethasone (DEX)-inducible 35S promoter. Plants were grown for three weeks on soil and treated with 30 µM DEX to induce BIR1 transcription or mock-treated with H_2O. Plants were photographed after 12 days of the first DEX/ H_2O application. Images from independent transgenic lines are shown in Fig. S7. RT-qPCR analysis of BIR1-coding transcripts in *eds1-2* BIR1 L1b and *pad4-1* BIR1 L0 mutant lines are shown. RT-qPCR analysis of genes encoding the PTI-markers RESPIRATORY BURST OXIDASE HOMOLOG PROTEIN D (RBOHD) and FLG22-INDUCED RECEPTOR-LIKE KINASE 1 (FRK1) in *eds1-2* BIR1 L1b (b) and *pad4-1* BIR1 L0 (c) plants. RT-qPCR analysis of genes encoding representative markers of the salicylic acid PATHOGENESIS RELATED 1 (PR1), jasmonic acid PLANT DEFENSIN 1.2 (PDF1.2), ethylene PR4 and camalexin [PHYTOALEXIN DEFICIENT 3 (PAD3)*, CYTOCHROME P450 MONOOXYGENASE 79B2 (CYP79B2)] pathways in *eds1-2* BIR1 L1b (d) and *pad4-1* BIR1 L0 (e) plants. In all cases, the same set of replicates from each genotype was used to measure all the genes shown in the graphs. Transcript accumulation was determined in mock-treated (DEX-) versus DEX-treated (DEX+) after 12 days of treatment. Values from biologically independent samples were normalized to the CBP20 internal control, and related to mock-treated plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test; *, p <0.05. Experiments were repeated at least twice with similar results.

Fig. 7. Phenotypes caused by BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpression are independent of salicylic acid (SA)-mediated signaling pathways. (a) Morphology of transgenic *sid2-2* BIR1 (L5) and *eds5-3* BIR1 (L10) mutant plants expressing BIR1 under the control of a dexamethasone (DEX)-inducible 35S promoter. Plants were grown for three weeks on soil and treated with 30 μM DEX to induce BIR1 transcription or mock-treated with H_2O. Plants were photographed after 12 days of the first DEX/ H_2O application. Images from independent transgenic lines are shown in Fig. S7a, b. RT-qPCR analysis of genes encoding BIR1 and representative markers of the salicylic acid PATHOGENESIS RELATED 1 (PR1), jasmonic acid PLANT DEFENSIN 1.2 (PDF1.2), ethylene PR4 and camalexin [PHYTOALEXIN DEFICIENT 3 (PAD3)*, CYTOCHROME P450 MONOOXYGENASE 79B2 (CYP79B2)] pathways in *sid2-2* BIR1 (L5) (b) and *eds5-3* BIR1 (L10) (c) mutant lines. In all cases, the same set of replicates from each genotype was used to measure all the genes shown in the graphs. Transcript accumulation was determined in mock-treated (DEX-) versus DEX-treated (DEX+) after 12 days of treatment. Values from biologically independent samples were normalized to the CBP20 internal control, and related to mock-treated plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test; *, *p* <0.05. Experiments were repeated at least twice with similar results. (d) Tentative working model for BIR1 homeostasis. SA-mediated transcription of BIR1 is regulated by RNA-directed DNA methylation (RdDM) and post-transcriptional RNA silencing (PTGS) (Guzman-Benito *et al.*, 2019). In these conditions, BIR1 induction interferes with pattern-triggered immunity (PTI) gene expression and plasmodesmata (PD) callose deposition triggered by various elicitors (flg22, nlp20, dsRNA). Loss of BIR1 regulation leads to constitutive BIR1 induction at non-physiological levels that cause SA-independent up-regulation of immune and hormone genes. BIR1 overexpression is sensed by one or several TOLL/INTERLEUKIN RECEPTOR 1 (TIR1)-containing nucleotide-binding domain leucine-rich repeat receptors (TNLs), of which CONSTITUTIVE SHADE AVOIDANCE 1 (CSA1) is a plausible candidate, to promote PAD4-independent effector triggered immunity (ETI)-type cell death that requires ENHANCED DISEASE SUSCEPTIBILITY 1 (EDS1) and SUPPRESSOR OF BIR1-1 (SOBIR1).

1 **New Phytologist Supporting Information**
2

Irene Guzmán-Benito[1], Carmen Robinson[1,2], Chenlei Hua[3], Ana Rocio Sede[4], Laura Elvira-

González[4], Isabel Punzón[1], Manfred Heinlein[4], Thorsten Nürnberger[3,5], César Llave[1]*

[1]Centro de Investigaciones Biológicas Margarita Salas. Consejo Superior de Investigaciones

Científicas, Ramiro de Maeztu 9, 28040 Madrid, Spain; [2]Doctorado en Biotecnología y

Recursos Genéticos de Plantas y Microorganismos Asociados, Universidad Politécnica de

Madrid, 28040-Madrid, Spain; [3]Department of Plant Biochemistry, Center of Plant Molecular

Biology (ZMBP), University of Tübingen, Tübingen, Germany; [4]Institut de Biologie

Moléculaire des Plantes, CNRS, Université de Strasbourg, 67000 Strasbourg, France;

[5]Department of Biochemistry, University of Johannesburg, Johannesburg, South Africa.

The following Supporting Information is available for this article:

Methods S1. RNA and protein analyses.

Methods S2. Gene expression analyses.

Methods S3. Reactive oxygen species (ROS), ethylene, plasmodesmata callose deposition and

dexamethasone (DEX) treatments.

Fig. S1. Transcriptomic profiling of BAK1-INTERACTING RECEPTOR-LIKE KINASE 1

(BIR1) expressing transgenic lines.

Fig. S2. Transcriptomic profiling of dexamethasone (DEX)-treated wild-type Col-0 plants.

Fig. S3. Enrichment analysis of Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways

using RNA-Seq data.

Fig. S4. Differential expression of receptor-like protein (RLP)-coding genes in BAK1-
INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants.

Fig. S5. Differential expression of cysteine-rich receptor-like kinase (CRK)-coding genes in
BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants.

Fig. S6. Differential expression of cell-surface pattern recognition receptors (PRR)-coding
genes in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants.

Fig. S7. Differential expression of calcium-dependent protein kinases (CPK)- and
HETEROTRIMERIC G PROTEIN-coding genes in BAK1-INTERACTING RECEPTOR-
LIKE KINASE 1 (BIR1) overexpressing plants.

Fig. S8. Morphological and growth defects in BAK1-INTERACTING RECEPTOR-LIKE
KINASE 1 (BIR1) overexpressing plants of different immune mutant genetic backgrounds.

Table S1. List of primers

Table S2. List of expressed genes.

Table S3. List of differentially expressed genes (DEG. padjust value < 0.05).

Table S4. List of differentially expressed genes (DEG. |Log2fold change > 1| padjust value <
0.01).

Table S5. Enrichment analysis of Gene Ontology (GO) terms.

Table S6. Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis.

Table S7. List of differentially expressed genes (DEG. padjust value < 0.05) within functional
categories.

Table S8. Statistical analysis of enrichment of functional gene groups in plants overexpressing
BIR1 after 12 days of DEX treatment relative to non-expressing controls using Two-tailed
Fisher's exact tests.

55
56

Supporting methods

Methods S1. RNA and protein analyses

RNA analysis. Total RNA was extracted using TRIzol reagent (Invitrogen) and treated with

DNase I (Invitrogen) following manufacturer's instructions to eliminate genomic DNA traces.

One-step quantitative RT-PCR (RT-qPCR) was carried out using Brilliant III Ultra-Fast SYBR

Green QRT-PCR Master Mix (Agilent Technologies) in a Rotor-Gene 6000/Rotor-Gene Q real-

time PCR machine (Corbett/Qiagen) (Fernandez-Calvino *et al.*, 2016). Relative gene

expression was determined using the Delta-delta cycle threshold method and Rotor-Gene 6000

Series Software (Corbett). Transcript levels of the target genes were normalized to the transcript

levels of the housekeeping gene CBP20 (At5g44200) as a reference gene.

Protein analyses. Leaf tissue samples were ground in liquid nitrogen in an Eppendorf tube and

homogenized in extraction buffer (65 mM Tris-HCl, pH 8; 3% sodium dodecyl sulfate [SDS];

1% ß-mercaptoethanol; 10% glycerol). Samples were diluted in Laemmli buffer, incubated for

5 min at 95°C, and loaded onto 10% SDS-PAGE protein gels. After electrophoresis, proteins

were blotted onto an ECL nitrocellulose membrane (Amersham-Pharmacia), and detected using

commercial horseradish peroxide (HRP)-conjugated secondary antibodies and a

chemiluminescent substrate (LiteAblot Plus). Primary and secondary antibodies were diluted

according to the manufacturer's recommendations. The RFP-Trap (Chromotek) was used for

immunoprecipitation of mCherry-tagged BIR1 protein from Arabidopsis leaves following

manufacturer's instruction.

Methods S2. Gene expression analyses

RNA and library preparation for transcriptome sequencing. Total RNA was extracted from each

sample using the RNeasy plant mini kit (Qiagen) followed by DNase treatment according to

manufacturer's instructions. A total of 24 samples, including three biological replicates per

treatment, were collected. Each sample consisted of RNA pooled from six individual plants.

RNA degradation and contamination were monitored on 1% agarose gels, and BIR1 transcript

accumulation was measured by RT-qPCR. The RNA purity and RNA concentration were

checked using a Nanodrop spectrophotometer (ThermoFisher Scientific), while the RNA

integrity was measured using the Agilent Bioanalyzer 2100 system (Agilent Technologies).

Sequencing libraries were generated at NOVOGENE (HK) COMPANY LIMITED (Wan Chai,

Hong Kong) (www.novogene.com) using NEBNext® UltraTM RNA Library Prep Kit for

Illumina® (New England Biolabs) following manufacturer's recommendations and index codes

were added to attribute sequences to each sample. First-strand cDNA from mRNA was

synthesized using random hexamer primer and M-MuLV Reverse Transcriptase (RNase H-).

Second-strand cDNA synthesis was subsequently performed using DNA Polymerase I and

RNase H. NEBNext Adaptors with hairpin loop structure were ligated. In order to select cDNA

fragments of preferentially 150~200 bp in length, the library fragments were purified with the

AMPure XP system (Beckman Coulter). PCR was done with Phusion High-Fidelity DNA

polymerase, Universal PCR primers and Index (X) Primer. PCR products were purified with

the AMPure XP system and library quality was assessed on the Agilent Bioanalyzer 2100

system. Clustering of the index-coded samples was performed on a cBot Cluster Generation

System using the TruSeq PE Cluster Kit v3-cBot-HS (Illumina) according to the manufacturer's

recommendations. Finally, paired-end reads were generated via an Illumina PE150 Hiseq

platform at NOVOGENE (HK) COMPANY LIMITED.

<u>Bioinformatics analysis of RNA-Seq data</u>. Clean reads were obtained by removing adaptor tags

and reads containing poly-N and low-quality reads from raw data. Paired-end clean reads were

mapped to the Arabidopsis TAIR v10 reference genome using HISAT2 software. To quantify

the gene expression levels, HTSeq v0.6.1 was used to calculate the number of mapped reads of

each gene and then normalize the results to the expected number of Fragments Per Kilobase of

transcript sequence per Millions of base-pairs sequenced (FPKM). Differential expression

analysis between groups (three biological replicates per condition) was performed using

DESeq2 R package (1.18.0). The resulting *p-value*s were adjusted using the Benjamini and

Hochberg's approach for controlling the False Discovery Rate (FDR). Genes with an adjusted

p-value ≤0.05 found by DESeq2 were deemed as differentially expressed (DEG). Functional

classification of DEGs including Gene Ontology (GO) and KEGG was performed. GO

enrichment analysis of DEGs was implemented by tGOseq R package, in which gene length

bias was corrected. KOBAS software was used to test the statistical enrichment of DEGs in the

KEGG database GO terms and KEGG pathways with corrected *p-value* ≤0.05 were considered

significantly enriched by DEGs.

Methods S3. Reactive oxygen species (ROS), ethylene, plasmodesmata callose deposition

and dexamethasone (DEX) treatments

ROS and ethylene measurements. The detection of ROS and ethylene was performed as
described (Albert *et al.*, 2015; Wan *et al.*, 2019). Approximately 3-4 week-old BIR1 L9 or WT
Arabidopsis plants were treated with 30 μM dexamethasone (DEX) (H_2O was used as control)
for 4 days at 24 h intervals and leaves were then cut into pieces and left floating on H_2O
overnight before measurement. For ROS assays, leaf pieces were analyzed in a 96-well plate
containing 20 μM L-012 (Wako Pure Chemical Industries Ltd) and 2 μg ml^{-1} peroxidase.
Luminescence was measured both before (background) and over 1 h following flg22 or nlp20
(GenScript Biotech) elicitation or mock treatment using a Mithras LB 940 luminometer
(Berthold Technologies). For ethylene measurement, leaf pieces were incubated in 20 mM MES
buffer, pH 5.6 and the indicated elicitor (GenScript Biotech). Ethylene accumulation was
measured by gas chromatographic analysis (GC-14A) of 1 ml of the air drawn from the closed
tube with a syringe after the indicated incubation time. At least three replicates were performed
for all treatments.

Callose staining. Approximately 3-4 week-old BIR1 L9 or WT Arabidopsis plants were sprayed
with 30 μM DEX (H_2O was used as control) for 3-4 days at 24 h intervals and then leaf disks
were excised and left floating on H_2O. For callose staining, individual leaf disks were soaked
with 0,1% aniline blue solution (in 50 mM potassium phosphate buffer, pH 8.0) containing
either 0.5 μg μl^{-1} poly(I:C) (Sigma) or 1 μM flg22 (GenScript Biotech). Leaf disks were further
evacuated for 1-2 minutes (< 0.8 pa) in a vacuum desiccator. Aniline blue fluorescence was
imaged 30 minutes after poly(I:C)/flg22 or control treatment using a Zeiss LSM 780 confocal
laser scanning microscope with ZEN 2.3 software (Carl Zeiss) and using a 405 nm diode laser
for excitation and filtering the emission at 475-525 nm. mCherry fluorescence was excited at
561 nm filtering the emission at 560-610 nm. 8-bit images were acquired with a Plant-
Apochromat 40x/1.4 Oil DIC M27 objective. Callose fluorescence intensity was quantified with
ImageJ software (http://rsbweb.nih. gov/ij/) using the plug-in calloseQuant (Huang *et al.*,
2022). Callose spots were measured in 3-4 images taken from each leaf disk. If not otherwise
mentioned, three leaf discs per plant from three different plants were evaluated for each
genotype or condition. Regions of interest (ROIs) selected by calloseQuant were verified
visually before measurement. Individual fluorescence intensities that occurred as outliers from
the general distribution of fluorescence intensities (<1%) in the sample were excluded from
analysis.

Dexamethasone (DEX) treatments. DEX (Sigma) was dissolved in 100% ethanol to make a 30
mM DEX stock solution which was stored at –20°C in a light-tight vial. For the time course

experiments monitoring BIR1 expression, Col-0 wild-type (controls) and transgenic BIR1 L9

and L6 plants were grown for 3 weeks in soil and sprayed at 24 h intervals with DEX at 30 µM

as indicated {McNellis, 1998 #2330}. At this concentration, plants display optimal induction

of the transgene {Guzman-Benito, 2019 #2456}. For RNA-seq, samples were harvested after 4

or 12 days of DEX treatments, before the appearance of cell-death-like symptoms. For gene

expression assays in mutant backgrounds, plants were collected by the time symptoms

appeared, which normally occurred after 10-14 days of DEX treatments. After 4 and 12 days

of treatments BIR1 was produced at physiological and non-physiological levels, respectively.

WT plants treated with DEX for 4 days exhibit normal growth and development, whereas a

slight growth retardation was observed in DEX-treated WT plants after 12 days of DEX

treatment **(Fig. 5a)**.

Supporting figures

Fig. S1. Transcriptomic profiling of BAK1-INTERACTING RECEPTOR-LIKE KINASE 1

(BIR1) expressing transgenic lines. (a) Principal component analysis (PCA) of RNA-Seq data.

The PCA was performed using normalized RNA-Seq data of differentially expressed genes

(DEGs) between dexamethasone (DEX)-treated and mock-treated transgenic BIR1 L9 plants

after 4 and 12 days of treatment (DeL9_vs MoL9_). Each biological replicate is represented in

the score plot. The variance explained by each component (%) is given in parentheses. The

sample (DeL9_12b) with differential expression within the DEX-treated L9 group is indicated

with a black arrow. (b) Number of up-or down-regulated transcripts, with a $|\log_2(\text{Fold-Change})|$

≥ 1 and adjusted p-value ≤ 0.05, in each pairwise comparison as indicated. (c) Volcano plots

showing the distribution of DEGs (adjusted p-value ≤ 0.05). The x-axis shows the fold change

in gene expression between samples, and y-axis shows the statistical significance of the

differences. Significantly up- and down-regulated genes are highlighted in red and green,

respectively. Genes that are not differentially regulated are in blue. (d) Scatterplot of selected

gene ontology (GO) terms of DEGs. GO terms between wild-type (WT) and BIR1 L9 plants at

4 and 12 days after DEX-treatment (DeL9_ vs MoL9_) are shown. Dot size represents the count

of different genes and the color indicates significance of the term enrichment. GO terms with

adjusted p-value < 0.05 are significantly enriched. The kinase term (indicated by an arrow) was

exclusively assigned to BIR1 L9 plants. WT, wild-type Col-0; L9, BIR1 L9 transgenic plants;

De, DEX-treated plants; Mo, Mock-treated plants; _4, 4 days of treatment; _12, 12 days of

treatment.

Fig. S2. Transcriptomic profiling of dexamethasone (DEX)-treated wild-type Col-0 plants. (a)

Morphology of representative wild-type (WT) Arabidopsis plants used for RNA-Seq

experiments. Samples were collected after 4 or 12 days of 30 μM DEX treatment and subjected

to RNA sequencing. H_2O-treated plants were used as controls. Note that application of DEX

has a negative effect on plant growth as deduced by the significant reduction of rosette size

after 12 days of DEX treatment relative to the non-treated controls. Box and whisker plot

showing 5-95 percentile was analyzed by Mann-Whitney test; ****, $p < 0.0001$. (b) Volcano

plots showing the distribution of differentially expressed genes (DEGs) (adjusted p-value $\leq$

0.05). The x-axis shows the gene fold change, and y-axis plots the statistical significance (p-

value) of the differences. Significantly up- and down-regulated genes are highlighted in red and

green, respectively. Genes that are not differentially regulated are in blue. (c) Principal

component analysis (PCA) of RNA-Seq data. The PCA was performed using normalized RNA-

Seq data of DEGs between DEX-treated and mock-treated WT plants after 4 and 12 days of

treatment (DeWT_vs MoWT_). Each biological replicate is represented in the score plot. The

variance explained by each component (%) is given in parentheses. (d) Scatterplot of selected

gene ontology (GO) terms of DEGs. GO terms between DEX-treated and mock-treated WT

plants at 4 and 12 days of treatment (DeL9_ vs MoL9_) are shown. Dot size represents the

count of different genes and the color indicates significance of the term enrichment. GO terms

with adjusted p-value < 0.05 are significantly enriched. WT, wild-type Col-0; De, DEX-treated

plants; Mo, Mock-treated plants; _4, 4 days of treatment; _12, 12 days of treatment.

Fig. S3. Enrichment analysis of Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways

using RNA-Seq data. Scatterplot of selected Kyoto Encyclopedia of Genes and Genomes

(KEGG) pathways. Dot size represents the count of different genes and the color indicates

significance of the term enrichment. Terms with adjusted p-value < 0.05 are significantly

enriched. WT, wild-type Col-0; De, DEX-treated plants; Mo, Mock-treated plants; _4, 4 days

of treatment; _12, 12 days of treatment.

Fig. S4. Differential expression of receptor-like protein (RLP)-coding genes in BAK1-
INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants. (a) Relative
expression levels are represented by the number of Fragments Per Kilobase of transcript
sequence per Millions of base-pairs sequenced (FPKM) in each of the three independent
samples per condition used for RNA-Seq. Significant differences between dexamethasone
(DEX)- and mock-treated samples were analyzed using two-way ANOVA followed by Sidak´s
multiple comparison test; ***, adjusted p-value < 0.001, **< 0.01. Note that the expression
levels of each individual replicate within the DEX-treated BIR1 L9_12 group mirror the
abundance of BIR1 transcripts in each replicate. (b) Differential expression based on $\log_2$ fold-
change in the indicated pairwise comparisons. WT, wild-type Col-0; L9, BIR1 L9 transgenics;
De, DEX-treated plants; Mo, mock-treated plants: _4d, 4 days of treatment; _12d, 12 days of
treatment.

Fig. S5. Differential expression of cysteine-rich receptor-like kinase (CRK)-coding genes in
BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants. (a)
Relative expression levels are represented by the number of Fragments Per Kilobase of
transcript sequence per Millions of base-pairs sequenced (FPKM) in each of the three
independent samples per condition used for RNA-Seq. Significant differences between
dexamethasone (DEX)- and mock-treated samples were analyzed using two-way ANOVA
followed by Sidak´s multiple comparison test; ***, adjusted p-value < 0.001, **< 0.01. Note
that the expression levels of each individual replicate within the DEX-treated BIR1 L9_12
group mirror the abundance of BIR1 transcripts in each replicate. (b) Differential expression
based on $\log_2$ fold-change in the indicated pairwise comparisons. WT, wild-type Col-0; L9,
BIR1 L9 transgenics; De, DEX-treated plants; Mo, mock-treated plants: _4d, 4 days of
treatment; _12d, 12 days of treatment.

Fig. S6. Differential expression of cell-surface pattern recognition receptors (PRR)-coding
genes in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants.
Relative expression levels are represented by the number of Fragments Per Kilobase of
transcript sequence per Millions of base-pairs sequenced (FPKM) in each of the three
independent samples per condition used for RNA-Seq. Significant differences between
dexamethasone (DEX)- and mock-treated samples were analyzed using two-way ANOVA

followed by Sidak´s multiple comparison test; ***, adjusted p-value < 0.001, **< 0.01, *< 0.05.

WT, wild-type Col-0; L9, BIR1 L9 transgenics; De, DEX-treated plants; Mo, mock-treated

plants: _4d, 4 days of treatment; _12d, 12 days of treatment.

Fig. S7. Differential expression of calcium-dependent protein kinases (CPK)- and

HETEROTRIMERIC G PROTEIN-coding genes in BAK1-INTERACTING RECEPTOR-

LIKE KINASE 1 (BIR1) overexpressing plants. Relative expression levels are represented by

the number of Fragments Per Kilobase of transcript sequence per Millions of base-pairs

sequenced (FPKM) in each of the three independent samples per condition used for RNA-Seq.

Significant differences between dexamethasone (DEX)- and mock-treated samples were

analyzed using two-way ANOVA followed by Sidak´s multiple comparison test; ***, adjusted

p-value < 0.001, **< 0.01, *< 0.05. WT, wild-type Col-0; L9, BIR1 L9 transgenics; De, DEX-

treated plants; Mo, mock-treated plants: _4d, 4 days of treatment; _12d, 12 days of treatment.

Fig. S8. Morphological and growth defects in BAK1-INTERACTING RECEPTOR-LIKE

KINASE 1 (BIR1) overexpressing plants of different immune mutant genetic backgrounds. (a)

Morphology of transgenic *sobir1-12* BIR1 L8, *bak1-5* BIR1 L4, *eds1-2* BIR1 L2, *pad4-1* BIR1

L12b and *eds5-3* BIR1 L12 mutant plants from independent lines expressing BIR1 under the

control of a dexamethasone (DEX)-inducible 35S promoter. Plants were grown for three weeks

on soil and treated with 30 μM DEX to activate BIR1 transcription or mock-treated with H_2O.

Plants were photographed after 12 days of the first DEX/ H_2O application. (b) Morphology and

BIR1 expression levels of DEX- and mock-treated *sobir1-12* BIR1 (L6, L8) and *bak1-5* BIR1

(L4, L10) plants as observed in different biological replicates of the experiment. qRT-PCR

analysis showed comparable BIR1 transcript levels in both genotypes. Each sample consisted

of RNA pooled from six different plants. (c) RT-qPCR analysis of transcripts encoding PTI-

marker WRKY DNA-BINDING PROTEIN 29 (WRKY29) in *sobir1-12* BIR1 L6 and *pad4-*1

BIR1 L10 mutant backgrounds. In all cases, transcript accumulation was determined in DEX-

treated (DEX+) versus mock-treated (DEX-) after 12 days of treatment. Values from

biologically independent samples were normalized to the CBP20 internal control, and related

to mock-treated plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test;

*, p <0.05. (d) Rosette size (diameter) in wild-type (WT) and BIR1 overexpressing ETI/PTI

mutants after 12 days of DEX treatment relative to the non-treated controls. Box and whisker

plots showing 5-95 percentile were analyzed by Mann-Whitney test; ****, p <0.0001. Note that

rosette size is largely influenced by the effect of DEX on plant development as shown for WT

Col-0 plant in this study.

Supporting tables

Table S1. List of primers

Table S2. List of expressed genes. Values represent the total read counts/ Fragments Per

Kilobase of transcript sequence per Millions of base-pairs sequenced (FRKM) values of each

biological replicate (a,b,c) per condition and their corresponding row mean values. De (DEX-

treated plants), Mo (Mock-treated plants), WT (wild-type plants), L9 (BIR1 L9 transgenic

plants), _4 (4 days of treatment), _12 (12 days of treatment)

Table S3. List of differentially expressed genes (DEG. padjust value < 0.05). Values represent

the mean Fragments Per Kilobase of transcript sequence per Millions of base-pairs sequenced

(FPKM) values from three biological replicates per condition. De (DEX-treated plants), Mo

(Mock-treated plants), WT (wild-type plants), L9 (BIR1 L9 transgenic plants), _4 (4 days of

treatment), _12 (12 days of treatment).

Table S4. List of differentially expressed genes (DEG. |Log2fold change > 1| padjust value <

0.01). Values represent the mean Fragments Per Kilobase of transcript sequence per Millions

of base-pairs sequenced (FPKM) values from three biological replicates per condition. De

(DEX-treated plants), Mo (Mock-treated plants), WT (wild-type plants), L9 (BIR1 L9

transgenic plants), _4 (4 days of treatment), _12 (12 days of treatment).

Table S5. Enrichment analysis of Gene Ontology (GO) terms. CC (cellular component), BP

(biological process) and MF (molecular function). Gene Ratio is the ratio of DEGs to all genes

for this GO term. Bg Ratio is the ratio of all genes concerning this GO term to all genes in

background GO database. Count is the number of differentially expressed genes concerning

this GO term. Adjusted p-value < 0.05 means significant enrichment. De (DEX-treated plants),

Mo (Mock-treated plants), WT (wild-type plants), L9 (BIR1 L9 transgenic plants), _4 (4 days
of treatment), _12 (12 days of treatment).

Table S6. Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis.
Gene Ratio is the ratio of DEGs to all genes for this pathway. Bg Ratio is the ratio of all genes
concerning this KEGG term to all genes in background KEGG database. Count is the number
of differentially expressed genes concerning this term. Adjusted p-value < 0.05 means
significant enrichment. De (DEX-treated plants), Mo (Mock-treated plants), WT (wild-type
plants), L9 (BIR1 L9 transgenic plants), _4 (4 days of treatment), _12 (12 days of treatment).

Table S7. List of differentially expressed genes (DEG. padjust value < 0.05) within functional
categories. Values represent the mean Fragments Per Kilobase of transcript sequence per
Millions of base-pairs sequenced (FPKM) values from three biological replicates per condition.
De (DEX-treated plants), Mo (Mock-treated plants), WT (wild-type plants), L9 (BIR1 L9
transgenic plants), _4 (4 days of treatment), _12 (12 days of treatment).

Table S8. Statistical analysis of enrichment of functional gene groups in plants overexpressing
BIR1 after 12 days of DEX treatment relative to non-expressing controls using Two-tailed
Fisher's exact tests. De (DEX-treated plants), Mo (Mock-treated plants), WT (wild-type plants),
L9 (BIR1 L9 transgenic plants). LRR, leucine-rich repeat; RLK, receptor-like kinase; CRK,
cysteine-rich RLKs; RLP, receptor-like protein; TNL, TIR1-nucleotide-binding domain LRR
immune receptors.

Supporting references

Albert I, Bohm H, Albert M, Feiler CE, Imkampe J, Wallmeroth N, Brancato C,
Raaymakers TM, Oome S, Zhang H, et al. 2015. An RLP23-SOBIR1-BAK1 complex
mediates NLP-triggered immunity. *Nat Plants* **1**: 15140.
Fernandez-Calvino L, Guzman-Benito I, Del Toro FJ, Donaire L, Castro-Sanz AB, Ruiz-
Ferrer V, Llave C. 2016. Activation of senescence-associated Dark-inducible (DIN) genes

during infection contributes to enhanced susceptibility to plant viruses. *Mol Plant Pathol*
17(1): 3-15.

Guzman-Benito I, Donaire L, Amorim-Silva V, Vallarino JG, Esteban A, Wierzbicki AT,
Ruiz-Ferrer V, Llave C. 2019. The immune repressor BIR1 contributes to antiviral defense
and undergoes transcriptional and post-transcriptional regulation during viral infections.
New Phytol **224**(1): 421-438.

Huang C, Mutterer J, Heinlein M. 2022. In Vivo Aniline Blue Staining and Semiautomated
Quantification of Callose Deposition at Plasmodesmata. *Methods Mol Biol* **2457**: 151-165.

McNellis TW, Mudgett MB, Li K, Aoyama T, Horvath D, Chua NH, Staskawicz BJ. 1998.
Glucocorticoid-inducible expression of a bacterial avirulence gene in transgenic Arabidopsis
induces hypersensitive cell death. *Plant J* **14**(2): 247-257.

Wan WL, Zhang L, Pruitt R, Zaidem M, Brugman R, Ma X, Krol E, Perraki A, Kilian J,
Grossmann G, et al. 2019. Comparing Arabidopsis receptor kinase and receptor protein-
mediated immune signaling reveals BIK1-dependent differences. *New Phytol* **221**(4): 2080-
2095.

Fig. S1. Transcriptomic profiling of BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) expressing transgenic lines. (a) Principal component analysis (PCA) of RNA-Seq data. The PCA was performed using normalized RNA-Seq data of differentially expressed genes (DEGs) between dexamethasone (DEX)-treated and mock-treated transgenic BIR1 L9 plants after 4 and 12 days of treatment (DeL9_vs MoL9_). Each biological replicate is represented in the score plot. The variance explained by each component (%) is given in parentheses. The sample (DeL9_12b) with differential expression within the DEX-treated L9 group is indicated with a black arrow. (b) Number of up-or down-regulated transcripts, with a |log$_2$(Fold-Change)| ≥1 and adjusted p-value ≤0.05, in each pairwise comparison as indicated. (c) Volcano plots showing the distribution of DEGs (adjusted p-value ≤ 0.05). The x-axis shows the fold change in gene expression between samples, and y-axis shows the statistical significance of the differences. Significantly up- and down-regulated genes are highlighted in red and green, respectively. Genes that are not differentially regulated are in blue. (d) Scatterplot of selected gene ontology (GO) terms of DEGs. GO terms between wild-type (WT) and BIR1 L9 plants at 4 and 12 days after DEX-treatment (DeL9_ vs MoL9_) are shown. Dot size represents the count of different genes and the color indicates significance of the term enrichment. GO terms with adjusted p-value < 0.05 are significantly enriched. The kinase term (indicated by an arrow) was exclusively assigned to BIR1 L9 plants. WT, wild-type Col-0; L9, BIR1 L9 transgenic plants; De, DEX-treated plants; Mo, Mock-treated plants; _4, 4 days of treatment; _12, 12 days of treatment.

Fig. S2. Transcriptomic profiling of dexamethasone (DEX)-treated wild-type Col-0 plants. (a) Morphology of representative wild-type (WT) Arabidopsis plants used for RNA-Seq experiments. Samples were collected after 4 or 12 days of 30 µM DEX treatment and subjected to RNA sequencing. H_2O-treated plants were used as controls. Note that application of DEX has a negative effect on plant growth as deduced by the significant reduction of rosette size after 12 days of DEX treatment relative to the non-treated controls. Box and whisker plot showing 5-95 percentile was analyzed by Mann-Whitney test; ****, $p < 0.0001$. (b) Volcano plots showing the distribution of differentially expressed genes (DEGs) (adjusted p-value ≤ 0.05). The x-axis shows the gene fold change, and y-axis plots the statistical significance (p-value) of the differences. Significantly up- and down-regulated genes are highlighted in red and green, respectively. Genes that are not differentially regulated are in blue. (c) Principal component analysis (PCA) of RNA-Seq data. The PCA was performed using normalized RNA-Seq data of DEGs between DEX-treated and mock-treated WT plants after 4 and 12 days of treatment (DeWT_vs MoWT_). Each biological replicate is represented in the score plot. The variance explained by each component (%) is given in parentheses. (d) Scatterplot of selected gene ontology (GO) terms of DEGs. GO terms between DEX-treated and mock-treated WT plants at 4 and 12 days of treatment (DeL9_ vs MoL9_) are shown. Dot size represents the count of different genes and the color indicates significance of the term enrichment. GO terms with adjusted p-value < 0.05 are significantly enriched. WT, wild-type Col-0; De, DEX-treated plants; Mo, Mock-treated plants; _4, 4 days of treatment; _12, 12 days of treatment.

Fig. S3. Enrichment analysis of Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways using RNA-Seq data. Scatterplot of selected Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways. Dot size represents the count of different genes and the color indicates significance of the term enrichment. Terms with adjusted p-value < 0.05 are significantly enriched. WT, wild-type Col-0; De, DEX-treated plants; Mo, Mock-treated plants; _4, 4 days of treatment; _12, 12 days of treatment.

Fig. S4. Differential expression of receptor-like protein (RLP)-coding genes in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants. (a) Relative expression levels are represented by the number of Fragments Per Kilobase of transcript sequence per Millions of base-pairs sequenced (FPKM) in each of the three independent samples per condition used for RNA-Seq. Significant differences between dexamethasone (DEX)- and mock-treated samples were analyzed using two-way ANOVA followed by Sidak´s multiple comparison test; ***, adjusted p-value < 0.001, **< 0.01. Note that the expression levels of each individual replicate within the DEX-treated BIR1 L9_12 group mirror the abundance of BIR1 transcripts in each replicate. (b) Differential expression based on log$_2$ fold-change in the indicated pairwise comparisons. WT, wild-type Col-0; L9, BIR1 L9 transgenics; De, DEX-treated plants; Mo, mock-treated plants: _4d, 4 days of treatment; _12d, 12 days of treatment.

Fig. S5. Differential expression of cysteine-rich receptor-like kinase (CRK)-coding genes in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants. (a) Relative expression levels are represented by the number of Fragments Per Kilobase of transcript sequence per Millions of base-pairs sequenced (FPKM) in each of the three independent samples per condition used for RNA-Seq. Significant differences between dexamethasone (DEX)- and mock-treated samples were analyzed using two-way ANOVA followed by Sidak´s multiple comparison test; ***, adjusted p-value < 0.001, **< 0.01. Note that the expression levels of each individual replicate within the DEX-treated BIR1 L9_12 group mirror the abundance of BIR1 transcripts in each replicate. (b) Differential expression based on $\log_2$ fold-change in the indicated pairwise comparisons. WT, wild-type Col-0; L9, BIR1 L9 transgenics; De, DEX-treated plants; Mo, mock-treated plants: _4d, 4 days of treatment; _12d, 12 days of treatment.

Fig. S6. Differential expression of cell-surface pattern recognition receptors (PRR)-coding genes in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants. Relative expression levels are represented by the number of Fragments Per Kilobase of transcript sequence per Millions of base-pairs sequenced (FPKM) in each of the three independent samples per condition used for RNA-Seq. Significant differences between dexamethasone (DEX)- and mock-treated samples were analyzed using two-way ANOVA followed by Sidak´s multiple comparison test; ***, adjusted p-value < 0.001, **< 0.01, *< 0.05. WT, wild-type Col-0; L9, BIR1 L9 transgenics; De, DEX-treated plants; Mo, mock-treated plants: _4d, 4 days of treatment; _12d, 12 days of treatment.

Fig. S7. Differential expression of calcium-dependent protein kinases (CPK)- and HETEROTRIMERIC G PROTEIN-coding genes in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants. Relative expression levels are represented by the number of Fragments Per Kilobase of transcript sequence per Millions of base-pairs sequenced (FPKM) in each of the three independent samples per condition used for RNA-Seq. Significant differences between dexamethasone (DEX)- and mock-treated samples were analyzed using two-way ANOVA followed by Sidak´s multiple comparison test; ***, adjusted p-value < 0.001, **< 0.01, *< 0.05. WT, wild-type Col-0; L9, BIR1 L9 transgenics; De, DEX-treated plants; Mo, mock-treated plants: _4d, 4 days of treatment; _12d, 12 days of treatment.

Fig. S8. Morphological and growth defects in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants of different immune mutant genetic backgrounds. (a) Morphology of transgenic *sobir1-12* BIR1 L8, *bak1-5* BIR1 L4, *eds1-2* BIR1 L2, *pad4-1* BIR1 L12b and *eds5-3* BIR1 L12 mutant plants from independent lines expressing BIR1 under the control of a dexamethasone (DEX)-inducible 35S promoter. Plants were grown for three weeks on soil and treated with 30 μM DEX to activate BIR1 transcription or mock-treated with H_2O. Plants were photographed after 12 days of the first DEX/ H_2O application. (b) Morphology and BIR1 expression levels of DEX- and mock-treated *sobir1-12* BIR1 (L6, L8) and *bak1-5* BIR1 (L4, L10) plants as observed in different biological replicates of the experiment. qRT-PCR analysis showed comparable BIR1 transcript levels in both genotypes. Each sample consisted of RNA pooled from six different plants. (c) RT-qPCR analysis of transcripts encoding PTI-marker WRKY DNA-BINDING PROTEIN 29 (WRKY29) in *sobir1-12* BIR1 L6 and *pad4*-1 BIR1 L10 mutant backgrounds. In all cases, transcript accumulation was determined in DEX-treated (DEX+) versus mock-treated (DEX-) after 12 days of treatment. Values from biologically independent samples were normalized to the CBP20 internal control, and related to mock-treated plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test; *, p <0.05. (d) Rosette size (diameter) in wild-type (WT) and BIR1 overexpressing ETI/PTI mutants after 12 days of DEX treatment relative to the non-treated controls. Box and whisker plots showing 5-95 percentile were analyzed by Mann-Whitney test; ****, p <0.0001. Note that rosette size is largely influenced by the effect of DEX on plant development as shown for WT Col-0 plant in this study.