RESEARCH ARTICLE

The receptor-like kinase BIR1 inhibits elicitor-induced

plasmodesmata callose deposition and PTI gene expression and

requires EDS1 and SOBIR1 to cause dose-dependent cell-death in

Arabidopsis

Irene Guzmán-Benito[1], Carmen Robinson[1], Chenlei Hua[2], Ana Rocio Sede[3], Laura

Elvira-González[3], Isabel Punzón[1], Manfred Heinlein[3], Thorsten Nürnberger[2,4], César

Llave[1]*

[1]Centro de Investigaciones Biológicas Margarita Salas. Consejo Superior de Investigaciones

Científicas, Ramiro de Maeztu 9, 28040 Madrid, Spain; [2]Department of Plant Biochemistry,

Center of Plant Molecular Biology (ZMBP), University of Tübingen, Tübingen, Germany;

[3]Institut de Biologie Moléculaire des Plantes, CNRS, Université de Strasbourg, 67000

Strasbourg, France; [4]Department of Biochemistry, University of Johannesburg, Johannesburg,

South Africa.

Author for correspondence: César Llave

21   E-mail: cesarllave@cib.csic.es

22   Phone: +34-91-8373112

Summary

- The receptor-like kinase BIR1 functions as a negative regulator of cell death and defense in Arabidopsis. Previous studies showed that BIR1 expression is up-regulated during infections with microbes and viruses. However, the biological consequences of BIR1 induction remain unknown.
- Here, we use a DEX-inducible expression system in Arabidopsis to investigate the outputs associated with physiological and non-physiological levels of BIR1 expression.
- We show that BIR1 induction at physiological levels significantly interferes with gene expression and plasmodesmata callose deposition triggered by canonical PTI elicitors. We found that plants that accumulated non-physiological doses of BIR1 displayed morphological defects that concur with transcriptomic changes in multiple plant defense genes. We provide experimental evidence that EDS1 and SOBIR1 are required for the ETI-type cell death phenotypes associated to non-physiological levels of BIR1
- Here we propose that BIR1 induction may represent a pathogen-triggered mechanism to modulate plant defenses during infection. Our model predicts that when BIR1 regulation is lost, BIR1 integrity is sensed by one or several guarding resistance (R) proteins to initiate an ETI-like response, in which SOBIR1 cooperates with EDS1 to transduce signals downstream of R proteins.

Key words

Arabidopsis, immune regulation, immune receptor, cell death, BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1), SUPPRESSOR OF BIR1 (SOBIR1), ENHANCED DISEASE SUSCEPTIBILITY 1 (EDS1)

Introduction

Plants use immunity to defend themselves against invaders such as bacteria, fungi, oomycetes, and viruses (Korner *et al.*, 2013; Macho & Lozano-Duran, 2019; Nishad *et al.*, 2020; Ngou *et al.*, 2022). This system involves multiple cell-surface pattern recognition receptors (PRR) that recognize pathogen/microbe/damage-associated molecular patterns (PAMP/MAMP/DAMP), leading to pattern-triggered immunity (PTI), and intracellular receptors that sense pathogen-effector proteins or effector-induced manipulations of host proteins, leading to effector-triggered immunity (ETI) (Li *et al.*, 2020; Zhou & Zhang, 2020; Lee *et al.*, 2021; Yu *et al.*, 2021). PRRs are single-transmembrane receptor-like kinases (RLK) or receptor-like proteins (RLP), which form active receptor complexes in a ligand-dependent manner with members of the somatic embryogenesis co-receptor kinases (SERKs), including BRASSINOSTEROID INSENSITIVE1-ASSOCIATED RECEPTOR KINASE 1 (BAK1/SERK3) (Boutrot & Zipfel, 2017; Hohmann *et al.*, 2017; Tang *et al.*, 2017; Smakowska-Luzan *et al.*, 2018; Lee *et al.*, 2021). BAK1 functions as a positive regulator of PTI (Liebrand *et al.*, 2014; Wu *et al.*, 2020), and has a dual role as positive and negative regulator of cell death during ETI (Belkhadir *et al.*, 2012; Dominguez-Ferreras *et al.*, 2015). RLP-type PRRs lack kinase activity and constitutively associate in a ligand-independent manner with the leucine-rich repeat (LRR)-RLK adapter SUPPRESSOR OF BIR1-1 (SOBIR1) (Gust *et al.*, 2017). Once the microbe is sensed, PRR receptor complexes activate, via trans-phosphorylation, receptor-like cytoplasmic kinases (RLCKs) that initiate downstream signaling cascades (Tang *et al.*, 2017; Liang & Zhou, 2018; Rao *et al.*, 2018). Among them, BOTRYTIS INDUCED KINASE1 (BIK1) functions either as a positive regulator of immunity triggered by LRR-RLKs such as those induced by the flg22 peptide derived from bacterial flagellin (FLAGELLIN SENSING 2, FLS2), or as a negative regulator of immunity triggered by LRR-RLPs such as those induced by the nlp20 fragment of NECROSIS AND ETHYLENE-INDUCING PEPTIDE 1 (NEP1)-LIKE proteins (NLP, RLP23) (Bi *et al.*, 2018; Wan *et al.*, 2019).

ETI relies on nucleotide-binding domain LRR intracellular immune receptors (NLRs) encoded by polymorphic disease resistance genes (R). NLRs comprise either proteins with an N-terminal domain similar to the TOLL/INTERLEUKIN RECEPTOR 1 (TIR1) (TIR-NB-LRR, TNL), proteins with a coiled-coil (CC) domain at the N terminus (CC-NB-LRR, CNL) or proteins with a RESISTANCE TO POWDERY MILDEW 8-LIKE DOMAIN (RPW8)-type domain (RNLs) (McHale *et al.*, 2006; van Wersch *et al.*, 2020; Wang & Chai, 2020). NLRs function as "sensors" that recognize effectors, or as "helpers" that act downstream of sensor NLRs in ETI

signaling (Jubic *et al.*, 2019). Recent studies demonstrated that PTI and ETI work together to
mutually potentiate the immune response (Yuan *et al.*, 2021b; Yuan *et al.*, 2023). For instance,
NLR-mediated signaling is dependent on multiple PTI-associated components such as BAK1,
SOBIR1, or BIK1 in a PTI-independent manner, while TNL genes that are up-regulated during
early PTI boost immune responses (Ngou *et al.*, 2021; Tian *et al.*, 2021; Yuan *et al.*, 2021a).
ENHANCED DISEASE SUSCEPTIBILITY (EDS1), which is essential in TNL-mediated
immunity, also plays a key role in PTI signaling in response to different microbial elicitors
(Pruitt *et al.*, 2021b; Tian *et al.*, 2021). Furthermore, PTI and ETI outputs usually converge,
albeit with different amplitude and duration (Yuan *et al.*, 2021b). Early responses include
apoplastic reactive oxygen species (ROS) burst, cytosolic calcium (Ca^{+2}) influx and activation
of Ca^{+2}-dependent protein kinases (CPK), and activation of mitogen-activated protein kinases
(MAPK). This response is followed by the transcriptional activation of defense processes that
include accumulation of defense phytohormones, regulation of cell wall and secretome
composition, and cellular motion (Li *et al.*, 2020).

Plants deploy a variety of mechanisms to protect themselves from the detrimental effects of
hyperactivated defenses (Couto & Zipfel, 2016; Withers & Dong, 2017; Mithoe & Menke,
2018). BAK1-INTERACTING RECEPTOR-LIKE KINASE (BIR) proteins play a key role in
defense attenuation. The current model postulates that, in the absence of microbes, BIR proteins
constitutively associate with BAK1, thus maintaining the PRRs in a resting state. During ligand
perception, BAK1 dissociates from BIR and becomes accessible to complex with PRR
receptors (Gao *et al.*, 2009; Halter *et al.*, 2014; Imkampe *et al.*, 2017). The Arabidopsis genome
has four BIR homologs (Halter *et al.*, 2014). BIR1 exerts a tight control of BAK1- and SOBIR1-
mediated cell death signaling, presumably related to PRR-RLP-ligand perception (Albert *et al.*,
2015; Liu *et al.*, 2016). BIR2 and BIR3 have critical roles in the negative regulation of PRR-
mediated PTI (Halter *et al.*, 2014; Imkampe *et al.*, 2017). Unlike BIR2 and BIR3, BIR1
possesses phosphorylation activities (Gao *et al.*, 2009). Interestingly, knocking-out BIR1
triggers resistance pathways that implicate components of TNL-mediated ETI response,
suggesting that loss of BIR1 activates one or several unknown TNL proteins (Gao *et al.*, 2009;
Liu *et al.*, 2016; Schulze *et al.*, 2022).

BIR1 expression is transcriptionally activated in response to multiple microbial and viral
pathogens in a salicylic acid (SA)-dependent manner (Guzman-Benito *et al.*, 2019). To date,
the effects of BIR1 induction on the outcome of plant defenses remain largely unknown.
However, previous work showed that transgenic overexpression of BIR1 triggers

developmental defects and cell death similar to the autoimmune phenotypes observed in *bir1-*
1 mutants (Gao *et al.*, 2009; Guzman-Benito *et al.*, 2019). Although the consequences of BIR1
overexpression have not been fully characterized yet, this finding seems to indicate that
perturbations in BIR1 homeostasis have severe growth-defense penalties. Clearly, effective
strategies for BIR1 regulation are needed to prevent unbearably high levels of BIR1 expression.
In this regard, RNA-directed DNA methylation and post-transcriptional RNA silencing regulate
levels of BIR1 transcripts when BIR1 is actively transcribed (Guzman-Benito *et al.*, 2019).

In this work, we conducted a time-course analysis to investigate the phenotypical outputs
associated with physiological and non-physiological levels of BIR1 expression in Arabidopsis
plants. We show that physiological induction of BIR1 interferes with the expression of PTI
marker genes and plasmodesmata (PD)-associated callose deposition in response to several PTI
elicitors, but not with ethylene accumulation or ROS production. Whereas physiological
induction of BIR1 has a subtle impact on the plant transcriptome, the morphological and growth
defects caused by the sustained expression of BIR1 are concomitant with the transcriptional
reprogramming of multiple genes encoding components of PTI and ETI signaling pathways.
We found that the cell death phenotypes associated with prolonged BIR1 overexpression
require the TNL-associated EDS1 and/or the RLP-co-receptor SOBIR1, but the underlying
mechanism is unknown. Finally, our work demonstrates that a well-balanced expression of
BIR1 avoids interference with cellular processes of growth and development, and is critical for
plant fitness.

Material and methods

Plant material and growth condition

Arabidopsis thaliana plants were grown in controlled environmental chambers under long-day
conditions (16h day/8h night) at 22°C. Arabidopsis lines used in this study were derived from
the Columbia-0 (Col-0) ecotype. Seeds were surface-sterilized and sown on plates containing
½ Murashige and Skoog (MS) medium supplemented with MES (0,5g/L), 1% sucrose and 0,9%
Plant agar (Duchefa Biochemie). Seedlings were transferred to soil seven to ten days after
germination.

Mutants for *bir1-1* and *sobir1-12* were donated by Yuelin Zhang (University of British
Columbia, Canada). The Arabidopsis *eds1-2*, *pad4-1* and *eds5-3* alleles were supplied by Jane
E. Parker (Max Planck Institute for Plant Breeding Research, Germany). Arabidopsis mutant

bak1-5 was donated by Birgit Kemmerling (Center of Plant Molecular Biology, University of
Tübingen, Germany). The *sid2-2* mutant was a gift from Francisco Tenllado (Centro de
Investigaciones Biológicas Margarita Salas, CSIC, Spain).

Arabidopsis Col-0 transgenic plants expressing a BIR1-mCherry fusion under a dexamethasone
(DEX)-inducible gene expression system (BIR1 L6 and BIR1 L9) have already been described
(Guzman-Benito *et al.*, 2019). In this study, the same DEX-inducible expression system
containing a mCherry-tagged BIR1 (herein BIR1) protein-coding gene was transformed into
Arabidopsis mutant backgrounds according to standard floral dipping (Clough & Bent, 1998).
Independent homozygous lines were selected in T3 and used for subsequent experiments.
Unless otherwise indicated, morphological phenotypes after DEX treatments were investigated
in two independent transgenic lines of each genotype.

RNA analysis

Total RNA was extracted using TRIzol reagent (Invitrogen) and treated with DNase I
(Invitrogen) following manufacturer's instructions to eliminate genomic DNA traces. One-step
quantitative RT-PCR (RT-qPCR) was carried out using Brilliant III Ultra-Fast SYBR Green
QRT-PCR Master Mix (Agilent Technologies) in a Rotor-Gene 6000/Rotor-Gene Q real-time
PCR machine (Corbett/Qiagen) (Fernandez-Calvino *et al.*, 2016). Relative gene expression was
determined using the Delta-delta cycle threshold method and Rotor-Gene 6000 Series Software
(Corbett). Transcript levels of the target genes were normalized to the transcript levels of the
housekeeping gene *CBP20* (*At5g44200*) as a reference gene. Primers are listed in Table S1.

Protein analysis

Leaf tissue samples were ground in liquid nitrogen in an Eppendorf tube and homogenized in
extraction buffer (65 mM Tris-HCl, pH 8; 3% sodium dodecyl sulfate [SDS]; 1% ß-
mercaptoethanol; 10% glycerol). Samples were diluted in Laemmli buffer, incubated for 5 min
at 95°C, and loaded onto 10% SDS-PAGE protein gels. After electrophoresis, proteins were
blotted onto an ECL nitrocellulose membrane (Amersham-Pharmacia), and detected using
commercial horseradish peroxide (HRP)-conjugated secondary antibodies and a
chemiluminescent substrate (LiteAblot Plus). Primary and secondary antibodies were diluted
according to the manufacturer's recommendations. The RFP-Trap (Chromotek) was used for
immunoprecipitation of mCherry-tagged BIR1 protein from Arabidopsis leaves following
manufacturer's instruction.

RNA and library preparation for transcriptome sequencing

Total RNA was extracted from each sample using the RNeasy plant mini kit (Qiagen) followed

by DNase treatment according to manufacturer's instructions. A total of 24 samples, including

three biological replicates per treatment, were collected. Each sample consisted of RNA pooled

from six individual plants. RNA degradation and contamination were monitored on 1% agarose

gels, and BIR1 transcript accumulation was measured by RT-qPCR. The RNA purity and RNA

concentration were checked using a Nanodrop spectrophotometer (ThermoFisher Scientific),

while the RNA integrity was measured using the Agilent Bioanalyzer 2100 system (Agilent

Technologies). Sequencing libraries were generated at NOVOGENE (HK) COMPANY

LIMITED (Wan Chai, Hong Kong) (www.novogene.com) using NEBNext® UltraTM RNA

Library Prep Kit for Illumina® (New England Biolabs) following manufacturer's

recommendations and index codes were added to attribute sequences to each sample. First-

strand cDNA from mRNA was synthesized using random hexamer primer and M-MuLV

Reverse Transcriptase (RNase H-). Second-strand cDNA synthesis was subsequently

performed using DNA Polymerase I and RNase H. NEBNext Adaptors with hairpin loop

structure were ligated. In order to select cDNA fragments of preferentially 150~200 bp in

length, the library fragments were purified with the AMPure XP system (Beckman Coulter).

PCR was done with Phusion High-Fidelity DNA polymerase, Universal PCR primers and Index

(X) Primer. PCR products were purified with the AMPure XP system and library quality was

assessed on the Agilent Bioanalyzer 2100 system. Clustering of the index-coded samples was

performed on a cBot Cluster Generation System using the TruSeq PE Cluster Kit v3-cBot-HS

(Illumina) according to the manufacturer's recommendations. Finally, paired-end reads were

generated via an Illumina PE150 Hiseq platform at NOVOGENE (HK) COMPANY LIMITED.

Bioinformatics analysis of RNA-Seq data

Clean reads were obtained by removing adaptor tags and reads containing poly-N and low-

quality reads from raw data. Paired-end clean reads were mapped to the Arabidopsis TAIR v10

reference genome using HISAT2 software. To quantify the gene expression levels, HTSeq

v0.6.1 was used to calculate the number of mapped reads of each gene and then normalize the

results to the expected number of Fragments Per Kilobase of transcript sequence per Millions

of base-pairs sequenced (FPKM). Differential expression analysis between groups (three

biological replicates per condition) was performed using DESeq2 R package (1.18.0). The

resulting *p-value*s were adjusted using the Benjamini and Hochberg's approach for controlling

the False Discovery Rate (FDR). Genes with an adjusted *p-value* ≤0.05 found by DESeq2 were

deemed as differentially expressed (DEG). Functional classification of DEGs including Gene

Ontology (GO) and KEGG was performed. GO enrichment analysis of DEGs was implemented by tGOseq R package, in which gene length bias was corrected. KOBAS software was used to test the statistical enrichment of DEGs in the KEGG database GO terms and KEGG pathways with corrected *p-value* ≤0.05 were considered significantly enriched by DEGs. The raw data was deposited in the Gene Expression Omnibus under accession number GSE234036.

Reactive oxygen species (ROS) and ethylene measurement

The detection of ROS and ethylene was performed as described (Albert *et al.*, 2015; Wan *et al.*, 2019). Approximately 4-wk-old BIR1 L9 or WT Arabidopsis plants were treated with 30 μM DEX (H_2O was used as control) for 4 days at 24 h intervals and leaves were then cut into pieces and left floating on H_2O overnight before measurement. For ROS assays, leaf pieces were analyzed in a 96-well plate containing 20 μM L-012 (Wako Pure Chemical Industries Ltd) and 2 μg ml^{-1} peroxidase. Luminescence was measured both before (background) and over 1 h following flg22 (GenScript Biotech) elicitation or mock treatment using a Mithras LB 940 luminometer (Berthold Technologies). For ethylene measurement, leaf pieces were incubated in 20 mM MES buffer, pH 5.6 and the indicated elicitor (GenScript Biotech). Ethylene accumulation was measured by gas chromatographic analysis (GC-14A) of 1 ml of the air drawn from the closed tube with a syringe after the indicated incubation time. At least three replicates were performed for all treatments.

Callose staining

Callose staining was carried out as described (Huang *et al.*, 2022; Huang *et al.*, 2023). Briefly, approximately 4-wk-old BIR1 L9 or WT Arabidopsis plants were sprayed with 30 μM DEX (H_2O was used as control) for 4 days at 24 h intervals and then leaf disks were excised and left floating on H_2O. For callose staining, individual leaf disks were soaked with 0,1% aniline blue solution (in 50 mM potassium phosphate buffer, pH 8.0) containing either 0.5 μg μl^{-1} poly(I:C) (Sigma) or 1 μM flg22 (GenScript Biotech). Leaf disks were further evacuated for 1-2 minutes (< 0.8 pa) in a vacuum desiccator. Aniline blue fluorescence was imaged 30 minutes after poly(I:C)/flg22 or control treatment using a Zeiss LSM 780 confocal laser scanning microscope with ZEN 2.3 software (Carl Zeiss) and using a 405 nm diode laser for excitation and filtering the emission at 475-525 nm. mCherry fluorescence was excited at 561 nm filtering the emission at 560-610 nm. 8-bit images were acquired with a Plant-Apochromat 40x/1.4 Oil DIC M27 objective. Callose fluorescence intensity was quantified with ImageJ software (http://rsbweb.nih. gov/ij/) using the plug-in calloseQuant (Huang *et al.*, 2022). Callose spots

were measured in 3-4 images taken from each leaf disk. If not otherwise mentioned, three leaf discs per plant from three different plants were evaluated for each genotype or condition. Regions of interest (ROIs) selected by calloseQuant were verified visually before measurement. Individual fluorescence intensities that occurred as outliers from the general distribution of fluorescence intensities (<1%) in the sample were excluded from analysis.

Results

BIR1 induction causes dose-dependent transcriptome reprogramming in Arabidopsis

To understand the downstream biological processes and pathways associated with BIR1 induction, we conducted a global transcriptome RNA-Seq assay in Arabidopsis WT and BIR1 L9 plants that express a DEX-inducible BIR1-mCherry transgene (Guzman-Benito *et al.*, 2019). This system proved to be a valuable tool for studying the phenotypic outcomes caused by different levels of BIR1 expression in the absence of pathogen elicitors or effectors. Three-week-old plants were sprayed with a 30 μM DEX solution at 24 h intervals and rosette leaf samples were harvested for next-generation sequencing. Control plants were mock-treated with H_2O. Three biological replicates were used as representative samples for each condition tested in this study: DEX-treated WT (DeWT_), mock-treated WT (MoWT_), DEX-treated BIR1 L9 (DeL9_) and mock-treated BIR1 L9 (MoL9_) plants at 4 or 12 days of treatment as indicated **(Table S2)**. We found that BIR1 was induced at physiological levels in DeL9_ samples after 4 days of DEX treatment, and plants displayed WT morphologies **(Fig. 1A)**. In contrast, DeL9_ plants accumulated high doses of BIR1 (non-physiological), and exhibited a range of growth defects and premature senescence after long BIR1 induction times (12 days of DEX treatment) **(Fig. 1A)** (Guzman-Benito *et al.*, 2019). Principal component analysis (PCA) was used to sort RNA-seq-based transcriptomic data according to gene expression levels. To discard the potential effects of DEX on gene expression, we first compared DEX-treated WT versus DEX-treated BIR1 L9 plants. PC1 defined a distinct separation in the PCA score plot between WT (DeWT_12) and BIR1 L9 (DeL9_12) samples after 12 days of treatment, indicating that the BIR1 expression level was the main source of variance (36.92%) **(Fig. 1B)**. Samples in the DeL9_4 group, which produced physiological levels of BIR1 after 4 days of treatment, clustered in the proximity of non-transgenic DeWT_4 samples, reflecting their similar expression profiles. PC2 seems to highlight a shift in gene expression between samples due to age or developmental stage (22.07%). When DEX-treated BIR1 L9 plants were compared with

mock-treated BIR1 L9 plants, the BIR1 expression level was again the dominant source of variance (36.85%) underlying PC1, whereas time was the basis for PC2 (19.57%) separation in the PCA score plot **(Fig. S1A).**

We noticed that normalized BIR1 abundance (FPKM) after 12 days of DEX treatment differed between independent BIR1 L9 replicates (DeL9_12), perhaps due to the differences in DEX sensitivity **(Fig. 1A)**. The DeL9_12b sample within the DeL9_12 group was located far from the other DeL9_12 samples in the PCA plot, and positioned near the samples within the DeWT_12 or MoL9_12 clusters **(Figs. 1B, S1A)**. Noticeably, DeL9_12b produced fewer BIR1-derived sequencing reads than the other two samples within the DeL9_12 group **(Fig. 1A)**. Thus, we conclude that BIR1 levels contribute to creating heterogeneous transcriptomes between replicates, although other factors such as the severity of the phenotypes caused by BIR1 overexpression may also yield quantitative differences in their expression profiles.

Differentially expressed genes (DEG) from different comparisons are presented in **Table S3**. A dramatic impact on the transcriptome was registered in plants that overexpress BIR1 (12 days of DEX treatment) (adjusted p-value ≤ 0.05) **(Figs. S1B, S1C)**. Using stricter statistical thresholds ($|\log_2(\text{Fold-Change})| \geq 1$ and adjusted p-value ≤ 0.01), 270 and 2,051 DEGs were detected in BIR1 L9 plants after 4 and 12 days of DEX treatment, respectively, relative to the mock-treated controls (DeL9_ vs MoL9_). Of them, 191 and 1,809 genes were up-regulated and 79 and 242 were down-regulated **(Fig. 1C)** **(Table S4)**. 584 and 4,125 genes were distinctly expressed between BIR1 L9 and WT plants after 4 and 12 days of DEX-treatment (DeL9_ vs DeWT_), respectively, of which 220 and 1,815 genes were up-regulated and 364 and 2,310 were down-regulated **(Fig. 1C)** **(Table S4)**. These findings indicate that a massive transcriptional reprogramming occurs when BIR1 is overexpressed in Arabidopsis but not when BIR1 is transcriptionally induced at physiological levels.

Gene functional enrichment analysis was used to infer biological functions or pathways significantly associated with DEGs. Gene ontology (GO) analysis showed that genes encoding proteins with oxidoreductase activity, transmembrane transport and other metabolic processes or genes related to lipid or carbohydrate metabolism were particularly overrepresented in our whole set of DEGs, including WT and BIR1 L9 plants, after long DEX treatments **(Figs. 1D, S1D, S2D) (Table S5)**. Notably, DEGs with kinase activity were only enriched in BIR1 overexpressing L9 plants, relative to mock-treated BIR1 L9 plants **(Figs. 1D, S1D)**. The kinase activity GO term was not enriched in DEX- or mock-treated WT plants at any point tested, excluding a DEX effect **(Fig. S2D)**. The Kyoto Encyclopedia of Genes and Genomes (KEGG)

pathway analysis showed the enrichment of genes related to several metabolic processes and biosynthesis of secondary metabolites that were specifically regulated after DEX-treatments **(Fig. S3) (Table S6)**. Conversely, DEGs involved in plant hormone signal transduction were significantly enriched only in BIR1 overexpressing BIR1 L9, but not in mock-treated BIR1 L9 plants or WT plants exposed to DEX.

As indicated above, the implemented experimental approach further allowed us to study the impact of DEX application. Macroscopically, WT plant treated with DEX for 12 days exhibited identical morphology as the mock-treated controls but produced significantly smaller rosettes due to the effects of DEX treatments **(Fig. S2A)**. Changes in gene expression due to DEX treatments in WT plants are listed in **Table S4 (Fig. S2B)**. PCA analysis revealed that the duration of the DEX treatment was the dominant source of variance underlying PC1 (36.80%) in the PCA score plot **(Fig. S2C)**. GO and KEGG analyses revealed a significant enrichment of terms related to oxidoreductase activities and metabolic processes in response to DEX treatments **(Figs. S2D, S3)**.

Effects of BIR1 induction on ROS and ethylene production and PTI gene expression in response to bacterial elicitors

We wanted to assess the effects of the physiological induction of BIR1 on early events of the PTI response. BIR1 was induced by treating BIR1 plants with a 30 µM DEX solution 4 times at 24h intervals, whereas control plants were mock-treated with H_2O. The luminol-based assay showed that ROS production triggered by 1 µM flg22 was comparable between WT and BIR1 L6 and L9 plants **(Fig. 2A)**. Then, we tested the production of the phytohormone ethylene in response to flg22 and nlp20 at concentrations of 1 µM and 100 nM. As expected, ethylene levels were significantly increased in WT, BIR1 L6 and L9 plants in response to flg22 and, to a greater extent, nlp20 at both concentrations tested. However, the induction of BIR1 in DEX-treated plants had no measurable consequences on ethylene accumulation compared to the mock-treated controls **(Fig. 2B)**. Next, we tested the expression of the canonical PTI marker genes encoding FLG22-INDUCED RECEPTOR-LIKE KINASE 1 (FRK1) and PATHOGENESIS-RELATED 1 (PR1) in BIR1 L6 and L9 transgenic plants upon 1 µM flg22 elicitation. Whereas all these genes were significantly up-regulated in mock-treated controls (DEX-), BIR1 induction by DEX in BIR1 L6 an L9 plants notably reduced transcript levels (DEX+). Importantly, PTI genes clearly responded to flg22 in WT plants regardless of DEX, which

discards a potential repressive effect of DEX on PTI gene expression (**Fig. 2C**). The induction of BIR1 protein in the BIR1 L6 and L9 plants used for ROS and ethylene assays was confirmed by immunocapture followed by western blot, while BIR1 transcript levels were monitored in the samples used for RT-qPCR expression assays (**Fig. 2D**).

Collectively, these results indicate that BIR1 induction within a physiological threshold has negligible effects on ROS production or ethylene accumulation but interferes with downstream events in the PTI pathway leading to transcriptional activation of defense genes.

Induction of BIR1 hinders callose deposition at PD triggered by flg22 or dsRNA

Callose-containing depositions at the cell-wall and PD triggered by conserved PAMPs occur during the early stages of pathogen invasion (Luna *et al.*, 2011). Deposition of callose at PD regulates symplastic transport, and limits the ability of the invading pathogen to enter plant tissues (De Storme & Geelen, 2014; Xu *et al.*, 2017; Huang *et al.*, 2023). To investigate whether physiological induction of BIR1 could affect PD callose deposition, we used *in vivo* aniline blue staining to quantify PD-associated callose from digital photographs of BIR1 L9 Arabidopsis plants. BIR1 was induced with a 30 µM DEX solution 3 times at 24h intervals, whereas control plants were mock-treated with H_2O. We first investigated callose deposition using 1 µM flg22 as elicitor. Flg22 treatment of control (DEX-) BIR1 L9 plants caused a noticeable increase in PD callose intensity compared to plants treated without elicitor (H_2O) (**Figs. 3A, C**). In contrast, PD-associated callose induced by flg22 remained at H_2O-induced control levels in BIR1-expressing plants (DEX+) (**Figs. 3A, C**), indicating that BIR1 was able to interfere, at least partially, with PD callose deposition triggered by flg22.

To study if BIR1-mediated callose inhibition was a general response, BIR1 L9 plants were treated with the synthetic dsRNA analog polyinosinic-polycytidylic acid [poly(I:C)] (0.5 µg µl^{-1}) since dsRNA is known to trigger deposition of callose and PD closure in Arabidopsis (Huang *et al.*, 2023). In the absence of DEX, callose intensity was significantly higher in BIR1 L9 plants treated with poly(I:C) compared to H_2O-treated plants. In contrast, after DEX treatment, poly(I:C)-induced callose was significantly repressed by the expression of BIR1 (**Figs. 3B, C**). Callose was enhanced in DEX-treated WT plants in response to poly(I:C) relative to H_2O-treated plants, suggesting that DEX does not affect callose deposition in the WT genotype (**Fig. 3C**). These findings indicate that the activation of BIR1 impairs PD callose deposition induced by different PAMP elicitors.

We further explored if callose accumulation elicited by poly(I:C) or flg22 at the PD was SOBIR1-dependent using a *sobir1-12* mutant (Gao *et al.*, 2009; Liu *et al.*, 2016). We observed enhanced callose deposition in WT plants in response to poly(I:C), whereas the median PD callose level detected in *sobir1-12* plants after poly(I:C) treatment was similar to those triggered in H_2O-treated *sobir1-12* plants **(Fig. 3D)**. Surprisingly, PD callose deposition in the *sobir1-12* background was also reduced in response to flg22 (**Fig. 3D**). Callose activation at PD by both elicitors was inhibited in the double *bir1-1 sobir1-1* mutant **(Fig. 3D).** These findings indicate that callose deposition triggered by either dsRNA or bacterial flg22 involved the SOBIR1 co-receptor.

BIR1 overexpression triggers the expression of immunity-related genes

To independently validate the RNA-Seq results, we used RT-qPCR to examine the expression of a subset of DEGs using independent sets (not used for RNA-seq) of mock- and DEX-treated BIR1 L9 plants collected after 12 days of treatment. Our results indicated that expression of BIR1, PR1, RESPIRATORY BURST OXIDASE HOMOLOG PROTEIN D (RBOHD), FRK1 and CYTOCHROME P450 MONOOXYGENASE 79B2 (CYP79B2) defense genes were all significantly up-regulated in Arabidopsis plants overexpressing BIR1 compared to mock-treated controls, which is in good consistency with their RNA-Seq-based expression profiles **(Fig. 4A)**.

A detailed inspection of the RNA-Seq data revealed that multiple genes involved in the perception of pathogens and the activation of downstream defense pathways were misregulated when BIR1 was overexpressed. To increase accuracy, in the following analyses we have considered solely DEGs with a |log$_2$ (Fold-Change)| ≥1 and adjusted *p*-value ≤0.01. Genes encoding LRR-type RLKs in the Arabidopsis genome were significantly enriched in response to high levels of BIR1 [n= 225, two-tailed Fisher's exact test, P = 0.001 (DeL9_12 vs DeWT_12) and P=4.5E-4 (DeL9_12 vs MoL9_12)] **(Table S7)**. Besides, cysteine-rich RLKs (CRK), which contain cysteine-rich repeats in their extracellular domain, were highly overrepresented in the pool of DEGs [n= 42, two-tailed Fisher's exact test, P = 1.06E-6 (DeL9_12 vs DeWT_12) and P=1.86E-9 (DeL9_12 vs MoL9_12)] **(Table S7) (Fig. S4)** (Chen, 2001). CRKs are important in the perception of apoplastic ROS and, likely in association with BAK1, in the activation of PTI, enhanced pathogen resistance and hypersensitive response-associated cell death (Yeh *et al.*, 2015; Quezada *et al.*, 2019; Castro *et al.*, 2021). Furthermore,

we observed a high percentage of RLP coding genes in Arabidopsis, not expected by chance, in the list of DEGs [n=57, two-tailed Fisher's exact test, P = 4.08E-5 (DeL9_12 vs DeWT_12) and P=6.04E-9 (DeL9_12 vs MoL9_12)] **(Table S7) (Fig. S5)** (Wang *et al.*, 2008; Lv *et al.*, 2016). Most of LRR-RLK, CRK and RLP transcripts were up-regulated in response to elevated BIR1, but not in DEX-treated WT plants. RNA-Seq data revealed that BIR1 overexpression boosted the transcription of many other genes involved in PTI and ETI signaling **(Figs. 4B, S6A)**. This includes the RLK co-receptors BAK1 and SOBIR1, RLCKs such as BIK1 and several AvrPphB SUSCEPTIBLE1(PBS1)-LIKE (PBL) (PBL4, PBL12, PBL20) (Liang & Zhou, 2018; Rao *et al.*, 2018) **(Table S7)**. Active BIK1 positively regulates a Ca^{2+} influx from apoplast to cytosol that induces a conformational change in CPKs that switches on their kinase activity (Li *et al.*, 2014; Yip Delormel & Boudsocq, 2019). Collectively, CPKs were not differentially regulated in plants overexpressing BIR1, but many individual CPKs were regarded as DEGs in our assay. For instance, the expression of CPK4-6 and CPK11, which are key players for transcriptional reprogramming upon elicitor perception, was enhanced only in DEX-treated BIR1 L9 plants **(Fig. S6B)** (Boudsocq *et al.*, 2010). Upon ligand perception, BIK1 dissociates from activated receptor complexes and phosphorylates the plasma membrane NADPH oxidase RBOHD to generate extracellular ROS (Kadota *et al.*, 2014; Li *et al.*, 2014; Marcec & Tanaka, 2021). Interestingly, RBOHD was induced in DEX-treated BIR1 L9 plants after both 4 and 12 days of treatment, but not in their corresponding controls **(Figs. 4A, 4B)**. We found that elevated BIR1 levels strongly increased the transcript abundance of genes encoding the lipase-like proteins EDS1, PHYTOALEXIN DEFICIENT 4 (PAD4), SENESCENCE ASSOCIATED-GENE 101 (SAG101) and ACTIVATED DISEASE RESISTANCE 1 (ADR1), which are required for TNL-receptor function during ETI **(Figs. 4B)** (Zhou & Zhang, 2020; Ngou *et al.*, 2021; Pruitt *et al.*, 2021b; Sun *et al.*, 2021; Tian *et al.*, 2021). Accordingly, a large number of DEGs in our assay were annotated as disease resistance (R) protein-coding genes, of which the TNL class was particularly enriched [n=100, two-tailed Fisher's exact test, P = 0.06 (DeL9_12 vs DeWT_12) and P=4.4E-5 (DeL9_12 vs MoL9_12)] (Swiderski *et al.*, 2009) **(Table S7)**. Conversely, neither NON-RACE SPECIFIC DISEASE RESISTANCE 1 (NDR1) nor genes encoding CNL intracellular receptors, which require NDR1, were differentially expressed in response to BIR1 overexpression **(Fig. 4B) (Table S3)**. EDS5, which transduces downstream of SA in Arabidopsis (Nawrath *et al.*, 2002), was highly up-regulated in a BIR1-dependent manner. Interestingly, BIR2 was distinctively up-regulated in DEX-treated BIR1 L9 samples, whereas BIR3 and its partner CONSTITUTIVE SHADE AVOIDANCE 1 (CSA1), which also interacts with BIR1, were unaffected in plants

overexpressing BIR1 **(Fig. S6A)** (Schulze *et al.*, 2022). Consistent with transcript levels, SOBIR1, PAD4-dependent PR1 and EDS1 proteins were strongly up-regulated by BIR1 overexpression in DEX-treated BIR1 L9 plants, compared to the non-treated controls (**Figs. 4C, 5C**).

In conclusion, our data indicated that the overexpression of BIR1 significantly alters hormone homeostasis and up-regulates multiple immune receptors and co-receptors involved in the perception and signaling of ligands and effectors. Defense activation in BIR1 overexpressing plants occurs in the absence of microbes or microbe-derived elicitors/effectors, and correlates with growth phenotypes and cell death that are reminiscent of plants with constitutively activated defenses (Lorrain *et al.*, 2003; Guzman-Benito *et al.*, 2019).

Genetic disruption of SOBIR1, but not BAK1, partially rescues BIR1 overexpression phenotypes

SOBIR1 and BAK1 are necessary for activating cell death and constitutive defense responses in the knock-out *bir1-1* mutant (Gao *et al.*, 2009; Liu *et al.*, 2016). Therefore, we tested if SOBIR1 and BAK1 were also required for the phenotypes associated with non-physiological doses of BIR1. A DEX-inducible BIR1-mCherry system was used to generate independent homozygous transgenic lines in the well-characterized *sobir1-12* and *bak1-5* mutant backgrounds (Gao *et al.*, 2009; Schwessinger *et al.*, 2011). Transgenic plants were sprayed with 30 μM DEX solution or H_2O (mock-treated controls) for 12 days at 24 h intervals to promote BIR1 overexpression.

As previously shown, our reference set of DEX-treated BIR1 L9 plants accumulated high levels of BIR1 transcripts, and displayed a range of stunting symptoms, accelerated senescence (yellowing), and cell death relative to non-treated BIR1 L9 controls or DEX-treated WT plants **(Fig. 5A)** (Guzman-Benito *et al.*, 2019). Nonetheless, growth retardation in DEX-treated plants was partially due to DEX effects as shown for WT plants after 12 days of DEX treatment **(Fig. 5A)**. When *sobir1-12* BIR1 plants were investigated, we observed that the phenotypes after 12 days of DEX treatments, albeit variable between biological replicates, were milder than those in BIR1 L9 plants, and ranged from typical WT appearance to localized yellowing and, occasionally, discrete leaf necrosis **(Figs. 5A, S7A, S7B)**. Interestingly, BIR1 transcript and protein accumulated in *sobir1-12* BIR1 plants like in BIR1 L9 plants after 12 days of DEX application, discarding potential dose effects **(Fig. 5B, C)**. Furthermore, transcripts of the PTI-

markers RBOHD, FRK1 and WRKY DNA-BINDING PROTEIN 29 (WRK29) in DEX-treated *sobir1-1*2 BIR1 plants, although slightly increased, were statistically indistinguishable from non-treated controls **(Figs. 5D, S7C)**, suggesting that the up-regulation of PTI genes due to BIR1 overexpression was affected by the lack of SOBIR1.

bak1-5 BIR1 plants consistently developed strong morphological phenotypes after 12 days of DEX application, that closely resemble those of BIR1 L9 plants **(Figs. 5A, S7A, S7B)**. RT-qPCR showed that BIR1 transcripts also accumulated in these plants after long DEX treatments, although to a variable degree depending on the transgenic lines and biological replicates. For instance, BIR1 transcript levels in *bak1*-5 BIR1 plants were comparable to those in *sobir1-12* BIR1 plants in **Fig. S7B**. Besides, differences in BIR1 protein accumulation were observed between DEX-treated *bak1-5* BIR1 L4 and L10 lines in this experiment **(Fig. 5C)**. Conversely, **Figs. 5A** and **B** illustrate data from a different experimental replicate in which BIR1 transcripts in *bak1*-5 BIR1 plants were much less abundant than in BIR1 L9 plants. These interesting observations suggest that the persistent induction of BIR1 over time, rather than an excess of BIR1, was the trigger of the morphological alterations in the *bak1*-5 mutant background. RBOHD, FRK1 and WRKY remained transcriptionally up-regulated in DEX-treated *bak1-5* BIR1 plants as they did in BIR1 L9 plants **(Figs. 5E)**. Collectively, these data indicate that the morphological phenotypes and the activation of PTI defense genes that occur when BIR1 is continually induced or overexpressed were partially dependent on SOBIR1, but genetically unrelated to BAK1.

Alterations in plant hormone pathways may negatively affect plant growth (Denance *et al.*, 2013). The transcriptional reprogramming of genes involved in plant hormone metabolism and signaling in BIR1-overexpressing plants prompted us to further examine their expression in the mutant backgrounds mentioned above. RT-qPCR revealed that genes encoding salicylic acid (SA)-dependent PR1, jasmonic acid (JA)-responsive PLANT DEFENSIN 1.2 (PDF1.2), ethylene (ET)- and JA-responsive PR4 as well as genes encoding the cytochrome P450 enzymes CYP71B15 (PAD3) and CYP79B2 involved in camalexin (CA)-biosynthesis and CA-mediated plant defense, were all up-regulated in BIR1-overexpressing plants of *sobir1-12* BIR1 genotypes **(Fig. 5F)**. Intriguingly, the induction of these genes was notably reduced in *bak1-5* BIR1 plants after DEX treatment compared to *sobir1-12* BIR1 or BIR1 L9 plants **(Fig. 4A, 5G)**. This data suggests that the *bak1-5* mutation may compromise, at least partially, defense hormone signaling upon BIR1 overexpression. None of these genes, except PR1, were induced

500 or repressed in DEX-treated WT plants, discarding a potential effect of DEX on their expression

501 **(Table S3)**.

Genetic disruption of EDS1, but not PAD4, partially rescues BIR1 overexpression phenotypes

A previous report showed that growth stunting, activation of cell death and defense responses

in the *bir1-1* mutant are partially dependent on EDS1 and PAD4 (Gao *et al.*, 2009). The EDS1-

PAD4 node has a relevant role in PRR signaling and TNL activation, indicating that it mediates

both PTI and ETI (Pruitt *et al.*, 2021a). Given that TNL-coding transcripts were abundant in

BIR1 L9 plants and that EDS1 transcripts and protein were elevated in BIR1 L9, *sobir1-12*

BIR1 and *bak1-5* BIR1 plants **(Figs. 5B, C)**, we wondered if TNL-mediated ETI responses

along with, or independently of, PTI could be important for BIR1 overexpression phenotypes.

Thus, we tested if the phenotypes caused by sustained BIR1 induction were dependent on EDS1

or PAD4. The DEX-inducible BIR1-mCherry system was used to generate homozygous

transgenic lines from *eds1-2* and *pad4-1* mutants (Parker *et al.*, 1996; Zhou *et al.*, 1998). The

eds1-2 BIR1 plants displayed mostly WT-like morphologies and, occasionally, localized

yellowing after 12 days of DEX treatments compared to non-treated plants **(Figs. 6A, S7A)**.

Importantly, RT-qPCR analysis showed that BIR1 accumulated in the *eds1-2* BIR1 plants as

much as in the BIR1 L9 plants **(Figs. 6A)**. The expression of FRK1 and RBOHD was

significantly enhanced in *eds1-2* BIR1 plants after DEX applications compared to non-treated

controls **(Fig. 6B)**. Our data indicate that the lack of EDS1 attenuates the morphological

phenotypes caused by BIR1 overexpression, and pose the possibility that these phenotypic

alterations and the activation of PTI genes are unconnected processes.

Next, we found that *pad4-1* BIR1 plants had essentially BIR1 L9-like phenotypes after 12 days

of DEX applications **(Figs. 6A, S7A)**. In these plants, BIR1 accumulated much less than in

BIR1 L9 or *eds1-2* BIR1 plants **(Fig. 6A)**. As shown for *bak1-5* BIR1 plants, we presume that

sustained induction of BIR1, rather than BIR1 overexpression, was sufficient to cause typical

BIR1 phenotypes regardless of the *pad4-1* mutation. Interestingly, PTI-components FRK1,

WRKY29 and RBOHD showed no responsiveness to BIR1 overexpression in *pad4-1* BIR1

plants, which support the idea that cell death and PTI activation are unrelated in these plants

(Figs. 6C, S7C). In summary, we observed that the genetic elimination of EDS1, but not PAD4,

contributes to mitigating the morphological phenotypes caused by continuous BIR1

induction/overexpression, and suggests that symptoms arise from the activation of EDS1-dependent ETI-type cell death rather than the disruption of PRR-mediated PTI responses.

RT-qPCR revealed a significant transcriptional activation of marker genes in the SA, JA, ET and CA pathways in response to high levels of BIR1 in DEX-treated *eds1-2* BIR1 plants versus non-treated controls **(Fig. 6D)**. On the contrary, the same markers were poorly or no longer up-regulated in *pad4*-1 BIR1 transgenics after 12 days of DEX treatments relative to the non-treated controls, suggesting that PAD4 is important for PTI gene expression and phytohormone homeostasis **(Fig. 6E)**.

Phenotypes of BIR1 overexpressing lines are unrelated to SA signaling pathways

Previously, high SA levels were found as partly responsible for *bir1-1* phenotypes (Gao *et al.*, 2009). However, *pad4-1* BIR1 plants were as sensitive as BIR1 L9 plants to DEX treatments, suggesting that morphological phenotypes caused by non-physiological BIR1 levels are independent of SA-mediated signaling. To confirm this prediction, the effects of mutations in the SA biosynthesis on the growth and defense activation in BIR1 overexpressing plants were assessed. In Arabidopsis, SALICYLIC ACID INDUCTION DEFICIENT 2/ISOCHORISMATE SYNTHASE 1 (SID2/ICS1) and EDS5 are involved in pathogen-induced SA biosynthesis (Huang *et al.*, 2020). Therefore, DEX-inducible BIR1-mCherry transgenics were generated in the *sid2-2* and *eds5-3* genetic backgrounds. We found that *sid2-2* BIR1 plants exhibited from moderate to severe morphological phenotypes after 12 days of DEX treatment that coincided with moderate levels of BIR1 overexpression relative to those observed in BIR1 L9 plants **(Figs. 7A, B)**. *eds5*-3 BIR1 plants typically displayed strong developmental defects and necrosis and accumulated BIR1 transcripts at BIR1 L9 levels after long DEX applications **(Figs. 7A, C)**. The transcription levels of PDF1.2 and CYP79B2, but not PAD3, were high in DEX-treated *sid2*-2 BIR1 plants compared to non-treated controls **(Fig. 7B)**, while PAD3 and PDF1.2, but not CYP79B2 were up-regulated in *eds5-3* BIR1 overexpressing plants relative to non-treated controls **(Fig. 7C)**. Our findings confirm that the phenotypical alterations linked to BIR1 overexpression were not drastically affected by disruptive mutations in SA biosynthesis (*sid2-2* and *eds5-3*) and SA signaling (*pad4-1*).

Discussion

Plants respond to microbes and viruses by enhancing the expression of multiple genes that function as regulators or executors of various defense programs (Nishad *et al.*, 2020). Among them, BIR1 was shown to be up-regulated during bacterial and oomycete infections, and during viral infections in a SA-dependent manner (Gao *et al.*, 2009; Guzman-Benito *et al.*, 2019). The observation that knock-out *bir1-1* mutants have increased resistance to the virulent oomycete *Hyaloperonospora parasitica* Noco2 and to the RNA virus *Tobacco rattle virus* (TRV) suggests an active role in plant immunity (Gao *et al.*, 2009; Guzman-Benito *et al.*, 2019). Since BIR1 is a negative regulator of cell death and defense (Liu *et al.*, 2016; Ma *et al.*, 2017), we hypothesized that BIR1 induction might be part of a common strategy to modulate, at least partially, the amplitude of the immune responses during microbial and viral infections. To test the hypothesis, we used a previously described DEX-inducible system for the conditional expression of a mCherry-tagged BIR1 transgene in Arabidopsis (Guzman-Benito *et al.*, 2019). This system allowed us to investigate the outputs associated with short and long BIR1 induction times. Importantly, BIR1 induction is triggered in the absence of immune elicitors or effectors that could otherwise initiate undesirable localized and systemic defense responses in the plant. Upon short-term DEX treatments, BIR1 accumulates in rosette leaves to levels comparable to those observed under natural conditions in bacteria- or virus-infected plants, or plants treated with SA (Gao *et al.*, 2009; Guzman-Benito *et al.*, 2019). In the absence of pathogens or elicitors, short induction of BIR1 has a subtle, if any, impact on both plant transcriptome and plant morphology. In contrast, in the presence of the PTI elicitor flg22, the physiological induction of BIR1 negatively affects the expression of several PTI-marker genes, whereas it has no effects on ROS burst, as previously shown (Liu *et al.*, 2016), or production of ethylene. This observation suggests that BIR1 can regulate, directly or indirectly, the activity of components in the PTI signaling pathway that ultimately control immune gene transcription. Likewise, BIR1 activation compromises the callose deposition at PD elicited by either flg22 or the dsRNA-analog poly(I:C). PD allow cell-to-cell and long-distance transport of micro- and macromolecules, representing gateways through which plant viruses pass from cell to cell to spread infections (De Storme & Geelen, 2014). Targeted deposition of callose modulates symplastic connectivity by regulating PD pore size, and therefore constitutes an essential layer in the defense against pathogens (Wang *et al.*, 2021). According to our data, induction of BIR1 during pathogen attack can be interpreted as a strategy to regulate PD trafficking. Although not tested in this study, BIR1 effects on gene expression and callose deposition may render PTI less effective against pathogens. Indeed, PTI gene expression and callose formation are usually blocked/inhibited by many defense-suppressing virulence effectors from different types of

microbes (Cui *et al.*, 2009; Wang *et al.*, 2021). Viral-encoded proteins such as the potyviral silencing suppressor helper-component proteinase (HC-Pro) or the movement protein (MP) of several tobamoviruses are capable of suppressing callose deposition (Citovsky *et al.*, 1990; Chowdhury *et al.*, 2020). A recent work shows that tobamovirus MPs promote viral cell-to-cell propagation by blocking dsRNA-induced callose deposition (Huang *et al.*, 2023). It is tempting to speculate that BIR1 induction during the viral infection controls PD callose deposition, and perhaps other antiviral defenses, triggered by viral-derived dsRNA. dsRNA-induced deposition of callose requires PTI signaling components such as SERK1 and BIK1/PBL1 (Huang *et al.*, 2023). Our results show that PD callose deposition elicited by flg22 or poly(I:C) is inhibited in the *sobir1-1* mutant. This observation is puzzling because flg22 signaling is independent of SOBIR1. Further research is needed to understand the role of SOBIR1 in controlling PD callose deposition.

Long DEX treatments of BIR1 transgenic plants led to sustained and/or elevated BIR1 levels (non-physiological) similar to those expected if BIR1 gene regulation were lost. Under these conditions, plants exhibit dramatic changes in gene expression and eventually strong morphological and growth phenotypes, including accelerated senescence (yellowing), stunting, necrosis and cell death (Guzman-Benito *et al.*, 2019). An excess of BIR1 in our transgenic plants also induced the activation of multiple genes involved in different branches of the PTI and ETI signaling responses. Among them, the up-regulation of a significant number of RLP-coding genes is striking. RLPs have been observed predominantly expressed in senescence leaves, supporting the idea that BIR1 overexpression accelerates a senescence program (Wang *et al.*, 2008). In Arabidopsis, RLP1, RLP23, RLP30, RLP32 and RLP42 mediate SOBIR1-dependent immune activation, but the role of other RLPs in the perception of pathogen and defense is unknown (Wang *et al.*, 2008; Zhang *et al.*, 2013; Albert *et al.*, 2019). The overexpression of genes encoding TNL-resistance proteins, but not CNLs, is another important feature linked to high levels of BIR1. Here, we provide genetic evidence that morphological phenotypes due to persistent BIR1 induction are partially dependent on EDS1. TNL-based immunity is mediated by two distinct signaling modules, EDS1/SAG101/NRG1 and EDS1/PAD4/ADR1 (Sun *et al.*, 2021). Since knocking out PAD4 had little influence on BIR1 overexpression phenotypes, we propose that EDS1, perhaps in concert with SAG101, triggers cell death-like phenotypes upon TNL activation in BIR1 overexpressing plants **(Fig. 7D)**. How TNL receptors are activated when BIR1 is strongly expressed is a question to resolve. A current model postulates that loss of BIR1 function leads to the activation of hypothetical guarding R

proteins which further feed PAD4- and SOBIR1-dependent resistance pathways (Gao *et al.*, 2009). Our results are also coherent with BIR1 or a yet unknown BIR1/SERK complex being guarded by one or several EDS1-dependent TNL proteins. Therefore, non-physiological induction of BIR1 may promote TNL-mediated resistance pathways that require EDS1 when no effectors or elicitors are present. In addition to EDS1, we demonstrated that SOBIR1 is necessary for BIR1 overexpression phenotypes. The role of SOBIR1 in the BIR1 pathway is unclear. Although extensive investigation is required, SOBIR1 may function as a specific co-regulator of the EDS1-dependent responses triggered by BIR1 overexpression, as part of the same signaling pathway **(Fig. 7D)**. This idea is inspired by a recent work showing that the cell-surface RLP23 receptor uses SOBIR1 to form a supramolecular complex with a EDS1/PAD4/ADR1 node in a ligand-independent manner to mediate defense signaling cascades that are common to both surface PRR (PTI) and NLR (ETI) receptors (Pruitt *et al.*, 2021b). The TNL protein CSA1 senses perturbances of BIR3, BAK1 or BIR3/BAK1 complexes leading to ETI-type autoimmune responses (Schulze *et al.*, 2022). CSA1 did not show responsiveness to BIR1 overexpression in our assay, but it has been shown to interact with BIR1 (Schulze *et al.*, 2022). It will be interesting to see if CSA1 also guards BIR1 integrity, perhaps in cooperation with SOBIR1 and/or other SERKs **(Fig. 7D)**. Finally, phenotypes linked to BIR1 overexpression were apparently unaffected by the *bak1-5* mutation, which suggests that other SERK family members can compensate for the loss of BAK1 in PTI/ETI signaling.

We observed that the up-regulation of FRK1, WRKY29 and RBOHD due to BIR1 overexpression was compromised by mutations in SOBIR1 and PAD4, but it was independent of EDS1. Since EDS1, PAD4 and SOBIR1 are important for both cell surface PRR receptor- and TNL-mediated signaling and resistance (Pruitt *et al.*, 2021b; Tian *et al.*, 2021), we will need additional experimentation to establish if changes in PTI gene expression involve PTI-dependent pathways, TNL activation or other unknown mechanisms yet to be determined. Our study also revealed significant alterations in plant hormone gene expression. Interestingly, genes related to the plant hormones ET, JA, CA, and SA, which play a central role in regulating immune responses, remained up-regulated in *eds1-2* BIR1 and *sobir1-12* BIR1 backgrounds, in which phenotypes were partially rescued. This suggests that i) regulation of hormone homeostasis does not require SOBIR1- and/or EDS1-mediated signaling pathways, and that ii) such hormonal imbalance has little, if any, influence on the ETI-like phenotypes observed in BIR1 overexpressing plants.

Conclusions

Here, we investigated the implications of BIR1 homeostasis on the immune response in Arabidopsis. BIR1 induction at physiological levels interferes with both immune gene expression and PD callose deposition triggered by canonical PTI elicitors. We posit that BIR1 induction may represent a pathogen-driven mechanism to modulate plant defenses, and that this mechanism helps to maintain a host-pathogen equilibrium during infections. Over time induction of BIR1 typically leads to high levels on BIR1 in the plant tissue, and causes severe morphological and growth defects concomitant with transcriptomic changes affecting plant immunity. These findings indicate a dose-dependency of the BIR1 regulatory activity, in which persistent BIR1 induction/BIR1 overexpression over time results in autoimmunity and fitness penalties. We provide experimental evidence that both EDS1 and SOBIR1, but neither PAD4 nor BAK1, were necessary for the ETI-type cell death phenotypes that occur when BIR1 is overexpressed. We hypothesize that one or several unknown TNLs monitor the integrity of BIR1 or BIR1-co-receptor complexes and that SOBIR1 cooperates with EDS1 to transduce signals downstream of TNL **(Fig. 7D)**.

Acknowledgements

This work was supported by National Research grants RTI2018-096979-B-I00 and PID2021 7982NB-I00 to CL from Ministerio de Ciencia e Innovación (Spain) and UE-FEDER, and grant iLINKA20415 to CL from CSIC (Spain). Contributions of ARS, LEG and MH were supported by grants ERA-NET SusCrop2 / ANR-21-SUSC-0003-01 and ANR-PRC / ANR-21-CE20-0020-01 to MH. Contributions of CH and TN were supported by DFG grant Nu70/17-1 to TN. We thank Rosa Lozano-Durán, Birgit Kemmerling, Laura Medina-Puche and Delphine Pott for critical reading and colleagues who kindly donated seeds and materials.

Author contributions

CL and IGB conceived and designed the study. CL, IGB, CR and IP performed expression assays and transgenics. CL and CH performed ROS and ethylene assays. CL, ARS and LEG performed callose assays. CL, IGB, MH and TN analyzed the data. CL wrote the article.

REFERENCES

Albert I, Bohm H, Albert M, Feiler CE, Imkampe J, Wallmeroth N, Brancato C,
Raaymakers TM, Oome S, Zhang H, et al. 2015. An RLP23-SOBIR1-BAK1 complex
mediates NLP-triggered immunity. *Nat Plants* **1**: 15140.

Albert I, Zhang L, Bemm H, Nurnberger T. 2019. Structure-function analysis of immune
receptor AtRLP23 with its ligand nlp20 and coreceptors AtSOBIR1 and AtBAK1. *Mol*
Plant Microbe Interact **32**(8): 1038-1046.

Belkhadir Y, Jaillais Y, Epple P, Balsemao-Pires E, Dangl JL, Chory J. 2012.
Brassinosteroids modulate the efficiency of plant immune responses to microbe-associated
molecular patterns. *Proc Natl Acad Sci U S A* **109**(1): 297-302.

Bi G, Zhou Z, Wang W, Li L, Rao S, Wu Y, Zhang X, Menke FLH, Chen S, Zhou JM.
2018. Receptor-like cytoplasmic kinases directly link diverse pattern recognition Receptors
to the activation of mitogen-activated protein kinase cascades in Arabidopsis. *Plant Cell*
30(7): 1543-1561.

Boudsocq M, Willmann MR, McCormack M, Lee H, Shan L, He P, Bush J, Cheng SH,
Sheen J. 2010. Differential innate immune signalling via Ca(2+) sensor protein kinases.
Nature **464**(7287): 418-422.

Boutrot F, Zipfel C. 2017. Function, discovery, and exploitation of plant pattern recognition
receptors for broad-spectrum disease resistance. *Annu Rev Phytopathol* **55**: 257-286.

Castro B, Citterico M, Kimura S, Stevens DM, Wrzaczek M, Coaker G. 2021. Stress-
induced reactive oxygen species compartmentalization, perception and signalling. *Nat*
Plants **7**(4): 403-412.

Chen Z. 2001. A superfamily of proteins with novel cysteine-rich repeats. *Plant Physiol* **126**(2):
473-476.

Chowdhury RN, Lasky D, Karki H, Zhang Z, Goyer A, Halterman D, Rakotondrafara
AM. 2020. HCPro suppression of callose deposition contributes to strain-specific
resistance against Potato Virus Y. *Phytopathology* **110**(1): 164-173.

Citovsky V, Knorr D, Schuster G, Zambryski P. 1990. The P30 movement protein of tobacco
mosaic virus is a single-strand nucleic acid binding protein. *Cell* **60**(4): 637-647.

Clough SJ, Bent AF. 1998. Floral dip: a simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. *Plant J* **16**(6): 735-743.

Couto D, Zipfel C. 2016. Regulation of pattern recognition receptor signalling in plants. *Nat*
Rev Immunol **16**(9): 537-552.

Cui H, Xiang T, Zhou JM. 2009. Plant immunity: a lesson from pathogenic bacterial effector
proteins. *Cell Microbiol* **11**(10): 1453-1461.

De Storme N, Geelen D. 2014. Callose homeostasis at plasmodesmata: molecular regulators
and developmental relevance. *Front Plant Sci* **5**: 138.

Denance N, Sanchez-Vallet A, Goffner D, Molina A. 2013. Disease resistance or growth: the
role of plant hormones in balancing immune responses and fitness costs. *Front Plant Sci*
4: 155.

Dominguez-Ferreras A, Kiss-Papp M, Jehle AK, Felix G, Chinchilla D. 2015. An overdose
of the Arabidopsis coreceptor BRASSINOSTEROID INSENSITIVE1-ASSOCIATED
RECEPTOR KINASE1 or its ectodomain causes autoimmunity in a SUPPRESSOR OF
BIR1-1-dependent manner. *Plant Physiol* **168**(3): 1106-1121.

Fernandez-Calvino L, Guzman-Benito I, Del Toro FJ, Donaire L, Castro-Sanz AB, Ruiz-
Ferrer V, Llave C. 2016. Activation of senescence-associated Dark-inducible (DIN) genes
during infection contributes to enhanced susceptibility to plant viruses. *Mol Plant Pathol*
17(1): 3-15.

Gao M, Wang X, Wang D, Xu F, Ding X, Zhang Z, Bi D, Cheng YT, Chen S, Li X, et al.
2009. Regulation of cell death and innate immunity by two receptor-like kinases in
Arabidopsis. *Cell Host Microbe* **6**(1): 34-44.

Gust AA, Pruitt R, Nurnberger T. 2017. Sensing danger: key to activating plant immunity.
Trends Plant Sci **22**(9): 779-791.

Guzman-Benito I, Donaire L, Amorim-Silva V, Vallarino JG, Esteban A, Wierzbicki AT,
Ruiz-Ferrer V, Llave C. 2019. The immune repressor BIR1 contributes to antiviral
defense and undergoes transcriptional and post-transcriptional regulation during viral
infections. *New Phytol* **224**(1): 421-438.

Halter T, Imkampe J, Mazzotta S, Wierzba M, Postel S, Bucherl C, Kiefer C, Stahl M,
Chinchilla D, Wang X, et al. 2014. The leucine-rich repeat receptor kinase BIR2 is a
negative regulator of BAK1 in plant immunity. *Curr Biol* **24**(2): 134-143.

Hohmann U, Lau K, Hothorn M. 2017. The structural basis of ligand perception and signal
activation by receptor kinases. *Annu Rev Plant Biol* **68**: 109-137.

Huang C, Mutterer J, Heinlein M. 2022. In vivo aniline blue staining and semiautomated
quantification of callose deposition at plasmodesmata. *Methods Mol Biol* **2457**: 151-165.

Huang C, Sede AR, Elvira-González L, Yan Y, Rodriguez M, Mutterer J, Boutant E, Shan
L, Heinlein M. 2023. dsRNA-induced immunity targets plasmodesmata and is suppressed
by viral movement proteins. *Plant Cell* https://doi.org/10.1101/2022.11.21.517408.

Huang W, Wang Y, Li X, Zhang Y. 2020. Biosynthesis and regulation of salicylic acid and
N-hydroxypipecolic acid in plant immunity. *Mol Plant* **13**(1): 31-41.

Imkampe J, Halter T, Huang S, Schulze S, Mazzotta S, Schmidt N, Manstretta R, Postel
S, Wierzba M, Yang Y, et al. 2017. The Arabidopsis leucine-rich repeat receptor kinase
BIR3 negatively regulates BAK1 receptor complex formation and stabilizes BAK1. *Plant*
Cell **29**(9): 2285-2303.

Jubic LM, Saile S, Furzer OJ, El Kasmi F, Dangl JL. 2019. Help wanted: helper NLRs and
plant immune responses. *Curr Opin Plant Biol* **50**: 82-94.

Kadota Y, Sklenar J, Derbyshire P, Stransfeld L, Asai S, Ntoukakis V, Jones JD, Shirasu
K, Menke F, Jones A, et al. 2014. Direct regulation of the NADPH oxidase RBOHD by
the PRR-associated kinase BIK1 during plant immunity. *Mol Cell* **54**(1): 43-55.

Korner CJ, Klauser D, Niehl A, Dominguez-Ferreras A, Chinchilla D, Boller T, Heinlein
M, Hann DR. 2013. The immunity regulator BAK1 contributes to resistance against
diverse RNA viruses. *Mol Plant Microbe Interact* **26**(11): 1271-1280.

Lee DH, Lee HS, Belkhadir Y. 2021. Coding of plant immune signals by surface receptors.
Curr Opin Plant Biol **62**: 102044.

Li L, Li M, Yu L, Zhou Z, Liang X, Liu Z, Cai G, Gao L, Zhang X, Wang Y, et al. 2014.
The FLS2-associated kinase BIK1 directly phosphorylates the NADPH oxidase RbohD to
control plant immunity. *Cell Host Microbe* **15**(3): 329-338.

Li P, Lu YJ, Chen H, Day B. 2020. The Lifecycle of the plant immune system. *CRC Crit Rev*
Plant Sci **39**(1): 72-100.

Liang X, Zhou JM. 2018. Receptor-like cytoplasmic kinases: central players in plant receptor
kinase-mediated signaling. *Annu Rev Plant Biol* **69**: 267-299.

Liebrand TW, van den Burg HA, Joosten MH. 2014. Two for all: receptor-associated kinases
SOBIR1 and BAK1. *Trends Plant Sci* **19**(2): 123-132.

Liu Y, Huang X, Li M, He P, Zhang Y. 2016. Loss-of-function of Arabidopsis receptor-like
kinase BIR1 activates cell death and defense responses mediated by BAK1 and SOBIR1.
New Phytol **212**(3): 637-645.

Lorrain S, Vailleau F, Balague C, Roby D. 2003. Lesion mimic mutants: keys for deciphering
cell death and defense pathways in plants? *Trends Plant Sci* **8**(6): 263-271.

Luna E, Pastor V, Robert J, Flors V, Mauch-Mani B, Ton J. 2011. Callose deposition: a
multifaceted plant defense response. *Mol Plant Microbe Interact* **24**(2): 183-193.

Lv Y, Yang N, Wu J, Liu Z, Pan L, Lv S, Wang G. 2016. New insights into receptor-like
protein functions in Arabidopsis. *Plant Signal Behav* **11**(7): e1197469.

Ma C, Liu Y, Bai B, Han Z, Tang J, Zhang H, Yaghmaiean H, Zhang Y, Chai J. 2017.
Structural basis for BIR1-mediated negative regulation of plant immunity. *Cell Res* **27**(12):
1521-1524.

Macho AP, Lozano-Duran R. 2019. Molecular dialogues between viruses and receptor-like
kinases in plants. *Mol Plant Pathol* **20**(9): 1191-1195.

Marcec MJ, Tanaka K. 2021. Crosstalk between calcium and ROS signaling during flg22-
triggered immune response in Arabidopsis leaves. *Plants (Basel)* **11**(1).

McHale L, Tan X, Koehl P, Michelmore RW. 2006. Plant NBS-LRR proteins: adaptable
guards. *Genome Biol* **7**(4): 212.

Mithoe SC, Menke FL. 2018. Regulation of pattern recognition receptor signalling by
phosphorylation and ubiquitination. *Curr Opin Plant Biol* **45**(Pt A): 162-170.

Nawrath C, Heck S, Parinthawong N, Metraux JP. 2002. EDS5, an essential component of
salicylic acid-dependent signaling for disease resistance in Arabidopsis, is a member of the
MATE transporter family. *Plant Cell* **14**(1): 275-286.

Ngou BPM, Ahn HK, Ding P, Jones JDG. 2021. Mutual potentiation of plant immunity by
cell-surface and intracellular receptors. *Nature* **592**(7852): 110-115.

Ngou BPM, Ding P, Jones JDG. 2022. Thirty years of resistance: Zig-zag through the plant
immune system. *Plant Cell* **34**(5): 1447-1478.

Nishad R, Ahmed T, Rahman VJ, Kareem A. 2020. Modulation of plant defense system in
response to microbial interactions. *Front Microbiol* **11**: 1298.

Parker JE, Holub EB, Frost LN, Falk A, Gunn ND, Daniels MJ. 1996. Characterization of
eds1, a mutation in Arabidopsis suppressing resistance to Peronospora parasitica specified
by several different RPP genes. *Plant Cell* **8**(11): 2033-2046.

Pruitt RN, Gust AA, Nurnberger T. 2021a. Plant immunity unified. *Nat Plants* **7**(4): 382-
383.

Pruitt RN, Locci F, Wanke F, Zhang L, Saile SC, Joe A, Karelina D, Hua C, Frohlich K,
Wan WL, et al. 2021b. The EDS1-PAD4-ADR1 node mediates Arabidopsis pattern-
triggered immunity. *Nature* **598**(7881): 495-499.

Quezada EH, Garcia GX, Arthikala MK, Melappa G, Lara M, Nanjareddy K. 2019.
Cysteine-rich receptor-like kinase gene family identification in the phaseolus genome and
comparative analysis of their expression profiles specific to mycorrhizal and rhizobial
symbiosis. *Genes (Basel)* **10**(1).

Rao S, Zhou Z, Miao P, Bi G, Hu M, Wu Y, Feng F, Zhang X, Zhou JM. 2018. Roles of
receptor-like cytoplasmic kinase VII members in pattern-triggered immune signaling.
Plant Physiol **177**(4): 1679-1690.

Schulze S, Yu L, Hua C, Zhang L, Kolb D, Weber H, Ehinger A, Saile SC, Stahl M, Franz-
Wachtel M, et al. 2022. The Arabidopsis TIR-NBS-LRR protein CSA1 guards BAK1-
BIR3 homeostasis and mediates convergence of pattern- and effector-induced immune
responses. *Cell Host Microbe* **30**(12): 1717-1731 e1716.

Schwessinger B, Roux M, Kadota Y, Ntoukakis V, Sklenar J, Jones A, Zipfel C. 2011.
Phosphorylation-dependent differential regulation of plant growth, cell death, and innate
immunity by the regulatory receptor-like kinase BAK1. *PLoS Genet* **7**(4): e1002046.

Smakowska-Luzan E, Mott GA, Parys K, Stegmann M, Howton TC, Layeghifard M,
Neuhold J, Lehner A, Kong J, Grunwald K, et al. 2018. An extracellular network of
Arabidopsis leucine-rich repeat receptor kinases. *Nature* **553**(7688): 342-346.

Sun X, Lapin D, Feehan JM, Stolze SC, Kramer K, Dongus JA, Rzemieniewski J,
Blanvillain-Baufume S, Harzen A, Bautor J, et al. 2021. Pathogen effector recognition-
dependent association of NRG1 with EDS1 and SAG101 in TNL receptor immunity. *Nat*
Commun **12**(1): 3335.

Swiderski MR, Birker D, Jones JD. 2009. The TIR domain of TIR-NB-LRR resistance
proteins is a signaling domain involved in cell death induction. *Mol Plant Microbe Interact*
22(2): 157-165.

Tang D, Wang G, Zhou JM. 2017. Receptor kinases in plant-pathogen interactions: more than
pattern recognition. *Plant Cell* **29**(4): 618-637.

Tian H, Wu Z, Chen S, Ao K, Huang W, Yaghmaiean H, Sun T, Xu F, Zhang Y, Wang S,
et al. 2021. Activation of TIR signalling boosts pattern-triggered immunity. *Nature*
598(7881): 500-503.

van Wersch S, Tian L, Hoy R, Li X. 2020. Plant NLRs: The whistleblowers of plant immunity.
Plant Commun **1**(1): 100016.

Wan WL, Zhang L, Pruitt R, Zaidem M, Brugman R, Ma X, Krol E, Perraki A, Kilian J,
Grossmann G, et al. 2019. Comparing Arabidopsis receptor kinase and receptor protein-
mediated immune signaling reveals BIK1-dependent differences. *New Phytol* **221**(4):
2080-2095.

Wang G, Ellendorff U, Kemp B, Mansfield JW, Forsyth A, Mitchell K, Bastas K, Liu CM,
Woods-Tor A, Zipfel C, et al. 2008. A genome-wide functional investigation into the
roles of receptor-like proteins in Arabidopsis. *Plant Physiol* **147**(2): 503-517.

Wang J, Chai J. 2020. Structural insights into the plant immune receptors PRRs and NLRs.
Plant Physiol **182**(4): 1566-1581.

Wang Y, Li X, Fan B, Zhu C, Chen Z. 2021. Regulation and function of defense-related
callose deposition in plants. *Int J Mol Sci* **22**(5).

Withers J, Dong X. 2017. Post-translational regulation of plant immunity. *Curr Opin Plant*
Biol **38**: 124-132.

Wu Y, Gao Y, Zhan Y, Kui H, Liu H, Yan L, Kemmerling B, Zhou JM, He K, Li J. 2020.
Loss of the common immune coreceptor BAK1 leads to NLR-dependent cell death. *Proc*
Natl Acad Sci U S A **117**(43): 27044-27053.

Xu B, Cheval C, Laohavisit A, Hocking B, Chiasson D, Olsson TSG, Shirasu K, Faulkner
C, Gilliham M. 2017. A calmodulin-like protein regulates plasmodesmal closure during
bacterial immune responses. *New Phytol* **215**(1): 77-84.

Yeh YH, Chang YH, Huang PY, Huang JB, Zimmerli L. 2015. Enhanced Arabidopsis
pattern-triggered immunity by overexpression of cysteine-rich receptor-like kinases. *Front*
Plant Sci **6**: 322.

Yip Delormel T, Boudsocq M. 2019. Properties and functions of calcium-dependent protein
kinases and their relatives in Arabidopsis thaliana. *New Phytol* **224**(2): 585-604.

Yu TY, Sun MK, Liang LK. 2021. Receptors in the Induction of the Plant Innate Immunity.
Mol Plant Microbe Interact **34**(6): 587-601.

Yuan M, Cai B, Xin XF. 2023. Plant immune receptor pathways as a united front against
pathogens. *PLoS Pathog* **19**(2): e1011106.

Yuan M, Jiang Z, Bi G, Nomura K, Liu M, Wang Y, Cai B, Zhou JM, He SY, Xin XF.
2021a. Pattern-recognition receptors are required for NLR-mediated plant immunity.
Nature **592**(7852): 105-109.

Yuan M, Ngou BPM, Ding P, Xin XF. 2021b. PTI-ETI crosstalk: an integrative view of plant
immunity. *Curr Opin Plant Biol* **62**: 102030.

Zhang W, Fraiture M, Kolb D, Loffelhardt B, Desaki Y, Boutrot FF, Tor M, Zipfel C,
Gust AA, Brunner F. 2013. Arabidopsis receptor-like protein30 and receptor-like kinase
suppressor of BIR1-1/EVERSHED mediate innate immunity to necrotrophic fungi. *Plant*
Cell **25**(10): 4227-4241.

Zhou JM, Zhang Y. 2020. Plant immunity: danger perception and signaling. *Cell* **181**(5): 978-
989.

Zhou N, Tootle TL, Tsui F, Klessig DF, Glazebrook J. 1998. PAD4 functions upstream from
salicylic acid to control defense responses in Arabidopsis. *Plant Cell* **10**(6): 1021-1030.

 Figure legends

Figure 1. Transcriptional reprogramming associated to BAK1-INTERACTING
RECEPTOR-LIKE KINASE 1 (BIR1) induction in Arabidopsis.

A) Morphology of representative BIR1 transgenic Arabidopsis plants (line 9, L9) under the
control of a dexamethasone (DEX)-inducible 35S promoter used for RNA-Seq experiments.
Samples were collected after 4 or 12 days of 30 μM DEX treatment. H_2O-treated plants were
used as controls. Relative BIR1 expression based on the number of Fragments Per Kilobase of
transcript sequence per Millions of base-pairs sequenced (FPKM) (top) and Western blot using
anti-mCherry antibody (bottom) in samples used for RNA-Seq. Significant differences between
DEX- and mock-treated samples were analyzed using two-way ANOVA followed by Sidak´s
multiple comparison test; ***, adjusted p-value < 0.001. B) Principal component analysis
(PCA) of RNA-Seq data. The PCA was performed using normalized RNA-Seq data of
differentially expressed genes (DEGs) between BIR1 L9 and wild-type (WT) plants after 4 and
12 days of DEX treatment (DeL9_vs DeWT_). Each biological replicate is represented in the
score plot. The variance explained by each component (%) is given in parentheses. The sample
(DeL9_12) with differential expression within the DEX-treated L9 group is indicated with
black arrows in (a) and (b). C) Number of up-or down-regulated transcripts, with a |$\log_2$ (Fold-
Change)| ≥1 and adjusted p-value ≤0.01, in each pairwise comparison is indicated. D)
Scatterplot of selected gene ontology (GO) terms of DEGs. GO terms between WT and BIR1
L9 plants at 4 and 12 days after DEX-treatment (DeL9_ vs DeWT_) are shown. Dot size
represents the count of different genes and the color indicates significance of the term
enrichment. GO terms with adjusted p-value < 0.05 are significantly enriched. The kinase term
(indicated by an arrow) was exclusively assigned to BIR1 L9 plants. WT, wild-type Col-0; L9,
BIR1 L9 transgenic plants; De, DEX-treated plants; Mo, Mock-treated plants; _4, 4 days of
treatment; _12, 12 days of treatment.

Figure 2. Effect of BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1)
induction on immune responses triggered by flg22 and nlp20 treatments.

A) Over time production of reactive oxygen species (ROS) in dexamethasone (DEX)-treated
wild-type (WT) and BIR1 L6 and L9 transgenic plants after elicitation of Arabidopsis leaf disks
with 1 μM flg22 (left) or H_2O (right). Bars represent means ± SD of relative fluorescence units.
B) Ethylene accumulation in BIR1 L6 and L9 transgenic plants after treatment with H_2O

(mock), flg22 and nlp20 at 1 µM or 100 nM concentration as indicated. Bars represent average

values ± SD. Within each elicitor/mock treatment, differences between DEX-treated (DEX+)

and controls (DEX-) were analyzed by two-way ANOVA followed by Sidak´s multiple

comparison test. C) RT-qPCR analysis of the PTI-marker genes FLG22-INDUCED

RECEPTOR-LIKE KINASE 1 (FRK1) and PATHOGENESIS RELATED 1 (PR1) at 45 min

after treatment with H_2O (mock) or 1 µM flg22, and determined in DEX-treated (DEX+) versus

non-treated (DEX-) samples. D) Immunoprecipitation of mCherry-BIR1 fusion proteins,

indicated with an arrowhead, from samples of BIR1 L6 and L9 plants used in (A) and (B) (left),

and RT-qPCR analysis of BIR1 transcripts in DEX-treated (DEX+) samples of BIR1 L6 and

L9 plants used in (C) (right). In all cases, RT-qPCR values from biologically independent

samples were normalized to the CBP20 internal control, and related to H_2O (mock)-treated

plants, set to a value of 1. Data are means ± SD analyzed by Kruskal-Wallis test; *, adjusted p-

value <0.05. Experiments were repeated three times with similar results.

Figure 3. BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) induction

inhibits flg22- and poly(I:C)-triggered callose accumulation at plasmodesmata (PD).

Callose spots at PD were visualized upon aniline blue staining of epidermal cells in response to

bacterial flg22 (A) or poly(I:C) (B) in BIR1 L9 transgenic Arabidopsis plants. Plants were

previously treated with 30 µM dexamethasone (DEX) to elicit BIR1 expression. Mock-treated

plants were used as non BIR1-expressing controls (top panels). Photographs were taken 30

minutes after treatment with H_2O (control) or 1 µM flg22 or 50 ng/µl of poly(I:C). BIR1-

mCherry fluorescence in DEX-treated BIR1 L9 plants was confirmed within the same images.

Inlays show enlargements of the areas within the dashed boxes. Scale bar, 10 µm. C) Relative

PD callose content in flg22/poly(I:C)-treated and H_2O (control)-treated leaves from BIR1 L9

plants previously treated with or without DEX as indicated. Callose content at PD in DEX-

treated wild-type (WT) plants in response to poly (I:C) is shown. D) flg22/poly(I:C)-induced

callose content at PD in the *sobir1-12* and *bir1-1 sobir1-1* mutants. H_2O-treated controls and

callose deposition at PD from WT Col-0 leaves are shown as controls. In all cases, black dots

represent the mean values of callose intensities in individual PD (>100) measured in three leaf

discs taken from at least three independent biological replicates per each treatment. Significant

differences between samples were analyzed by Kruskal-Wallis test or by one-way ANOVA

followed by Tukey's multiple comparison test; ****, adjusted p-value <0.0001, ***, <0.001,

**, <0.01. Experiments were repeated twice with similar results.

Figure 4. BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpression leads to the transcriptional activation of pattern-triggered immunity (PTI)-associated co-receptors and effector-triggered immunity (ETI) regulators.

A) Validation of the RNA-Seq results. Expression of selected BIR1-responsive differentially expressed genes (DEGs) after 12 days of dexamethasone (DEX) treatment based on fold-change values ($\log_2$ DeL9_12/DeWT_12 and $\log_2$ DeL9_12/MoL9_12 as indicated) was confirmed by RT-qPCR. Values from biologically independent samples were normalized to the CBP20 internal control, and related to untreated (DEX-) plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test; *, p <0.05. B) Relative expression of selected PTI/ETI-related genes based on the number of Fragments Per Kilobase of transcript sequence per Millions of base-pairs sequenced (FPKM) in each of the three independent samples per condition used for RNA-Seq. Actin expression is shown as control. Note that the expression levels of each individual replicate within the DEX-treated BIR1 L9_12 group mirror the abundance of BIR1 transcripts in each replicate. Significant differences between mock-treated and DEX-treated samples were analyzed using two-way ANOVA followed by Sidak´s multiple comparison test; ****, adjusted p-value < 0.0001, *** < 0.001, **< 0.01. WT, wild-type Col-0; L9, BIR1 L9 transgenics; _4d, 4 days of treatment; _12d, 12 days of treatment. C) Western blot analysis of SUPPRESSOR OF BIR1-1 (SOBIR1) and PATHOGENESIS RELATED 1 (PR1) using specific antibodies in two independent sets of BIR1 L9 samples. Plants were mock-treated (-) or DEX-treated (+) for 12 days. The presence (+) or absence (-) of morphological phenotypes (Ph) in each sample is indicated. Protein loading control by Ponceau S is shown.

Figure 5. Phenotypes caused by BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpression are partially rescued in *sobir1-12* mutants, but not in a *bak1-5* mutant.

A) Morphology of transgenic *sobir1-12* BIR1 L6 and *bak1*-5 BIR1 L10 mutant plants expressing BIR1 under the control of a dexamethasone (DEX)-inducible 35S promoter. Transgenic BIR1 L9 plants in the wild-type (WT) Col-0 background are shown as a reference. Plants were grown for three weeks on soil and treated with 30 µM dexamethasone (DEX) to induce BIR1 transcription or mock-treated with H_2O. Non-transgenic WT Col-0 plants are shown as controls. Plants were photographed after 12 days of the first DEX/ H_2O application.

Images from independent transgenic lines are shown in Fig. S7. RT-qPCR analysis of *BIR1* transcripts in BIR1 L9 plants are shown. B) RT-qPCR analysis of BIR1 and ENHANCED DISEASE SUSCEPTIBILITY 1 (EDS1) transcripts in *sobir1-12* BIR1 L6 and *bak1-5* BIR1 L10 plants. C) Western blot analysis of BIR1 and EDS1 proteins with specific antibodies in WT BIR1 L9 and two independent lines of *sobir1-12* BIR1 (L6, L8) and *bak1-5* BIR1 (L4, L10). Plants were mock-treated (-) or DEX-treated (+) for 12 days as indicated. The presence (+) or absence (-) of morphological phenotypes (Ph) in each sample is indicated. Protein loading control by Ponceau S is shown. RT-qPCR analysis of the PTI-marker genes RESPIRATORY BURST OXIDASE HOMOLOG PROTEIN D (RBOHD), FLG22-INDUCED RECEPTOR-LIKE KINASE 1 (FRK1) or WRKY DNA-BINDING PROTEIN 29 (WRKY29) in *sobir1-12* BIR1 L6 (D) and *bak1-5* BIR1 L10 (E) plants. RT-qPCR analysis of representative marker genes of the salicylic acid PATHOGENESIS RELATED 1 (PR1), jasmonic acid PLANT DEFENSIN 1.2 (PDF1.2), ethylene PR4 and camalexin [PHYTOALEXIN DEFICIENT 3 (PAD3)*,* CYTOCHROME P450 MONOOXYGENASE 79B2 (CYP79B2)] pathways in *sobir1-12* BIR1 L6 (F) and *bak1-5* BIR1 L10 (G) plants. In all cases, the same set of replicates from each genotype was used to measure all the genes shown in the graphs. Transcript accumulation was determined in mock-treated (DEX-) versus DEX-treated (DEX+) after 12 days of treatment. Values from biologically independent samples were normalized to the CBP20 internal control, and related to mock-treated plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test; *, $p < 0.05$. Experiments were repeated at least twice with similar results.

Figure 6. Phenotypes caused by BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpression are partially rescued in *eds1-2* mutants, but not in a *pad4-1* mutant.

A) Morphology of transgenic *eds1-2* BIR1 L1b and *pad4-1* BIR1 L0 mutant plants expressing BIR1 under the control of a dexamethasone (DEX)-inducible 35S promoter. Plants were grown for three weeks on soil and treated with 30 μM DEX to induce BIR1 transcription or mock-treated with H_2O. Plants were photographed after 12 days of the first DEX/ H_2O application. Images from independent transgenic lines are shown in Fig. S7. RT-qPCR analysis of BIR1 transcript in *eds1-2* BIR1 L1b and *pad4-1* BIR1 L0 mutant lines are shown. RT-qPCR analysis of the PTI-marker genes RESPIRATORY BURST OXIDASE HOMOLOG PROTEIN D (RBOHD) and FLG22-INDUCED RECEPTOR-LIKE KINASE 1 (FRK1) in *eds1-2* BIR1 L1b (B) and *pad4-1* BIR1 L0 (C) plants. RT-qPCR analysis of representative marker genes of the

salicylic acid PATHOGENESIS RELATED 1 (PR1), jasmonic acid PLANT DEFENSIN 1.2 (PDF1.2), ethylene PR4 and camalexin [PHYTOALEXIN DEFICIENT 3 (PAD3), CYTOCHROME P450 MONOOXYGENASE 79B2 (CYP79B2)] pathways in *eds1-2* BIR1 L1b (D) and *pad4-1* BIR1 L0 (E) plants. In all cases, the same set of replicates from each genotype was used to measure all the genes shown in the graphs. Transcript accumulation was determined in mock-treated (DEX-) versus DEX-treated (DEX+) after 12 days of treatment. Values from biologically independent samples were normalized to the CBP20 internal control, and related to mock-treated plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test; *, p <0.05. Experiments were repeated at least twice with similar results.

Figure 7. Phenotypes caused by BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpression are independent of salicylic acid (SA)-mediated signaling pathways.

A) Morphology of transgenic *sid2-2* BIR1 (L5) and *eds5-3* BIR1 (L10) mutant plants expressing BIR1 under the control of a dexamethasone (DEX)-inducible 35S promoter. Plants were grown for three weeks on soil and treated with 30 µM DEX to induce BIR1 transcription or mock-treated with H_2O. Plants were photographed after 12 days of the first DEX/ H_2O application. Images from independent transgenic lines are shown in Fig. S7a, b. RT-qPCR analysis of BIR1 and representative marker genes of the jasmonic acid PLANT DEFENSIN 1.2 (PDF1.2) and camalexin [PHYTOALEXIN DEFICIENT 3 (PAD3), CYTOCHROME P450 MONOOXYGENASE 79B2 (CYP79B2)] pathways in *sid2-2* BIR1 (L5) (B) and *eds5-3* BIR1 (L10) (C) mutant lines. In all cases, the same set of replicates from each genotype was used to measure all the genes shown in the graphs. Transcript accumulation was determined in mock-treated (DEX-) versus DEX-treated (DEX+) after 12 days of treatment. Values from biologically independent samples were normalized to the *CBP20* internal control, and related to mock-treated plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test; *, p <0.05. Experiments were repeated at least twice with similar results. D) Tentative working model for BIR1 homeostasis. SA-mediated transcription of BIR1 is regulated by RNA-directed DNA methylation (RdDM) and post-transcriptional RNA silencing (PTGS) (Guzman-Benito *et al.*, 2019). In these conditions, BIR1 induction interferes with PTI gene expression and PD callose deposition triggered by various elicitors (flg22, nlp20, dsRNA). Loss of BIR1 regulation leads to constitutive BIR1 induction and ultimately BIR1 overexpression at levels that cause SA-independent up-regulation of immune and hormone genes. BIR1 overexpression is sensed

1058   by one or several TNLs, of which CSA1 is a plausible candidate, to promote PAD4-independent

1059   ETI-type cell death that requires EDS1 and SOBIR1.

1060

1061

 Supporting information

 Figure S1. Transcriptomic profiling of BAK1-INTERACTING RECEPTOR-LIKE
 KINASE 1 (BIR1) expressing transgenic lines.

A) Principal component analysis (PCA) of RNA-Seq data. The PCA was performed using
normalized RNA-Seq data of differentially expressed genes (DEGs) between dexamethasone
(DEX)-treated and mock-treated transgenic BIR1 L9 plants after 4 and 12 days of treatment
(DeL9_vs MoL9_). Each biological replicate is represented in the score plot. The variance
explained by each component (%) is given in parentheses. B) Number of up-or down-regulated
transcripts, with a |log$_2$ (Fold-Change)| $\geq$1 and adjusted p-value $\leq$0.05, in each pairwise
comparison as indicated. C) Volcano plots showing the distribution of DEGs (adjusted p-value
$\leq$ 0.05). The x-axis shows the fold change in gene expression between samples, and y-axis
shows the statistical significance of the differences. Significantly up- and down-regulated genes
are highlighted in red and green, respectively. Genes that are not differentially regulated are in
blue. D) Scatterplot of selected gene ontology (GO) terms of DEGs. GO terms between wild-
type (WT) and BIR1 L9 plants at 4 and 12 days after DEX-treatment (DeL9_ vs MoL9_) are
shown. Dot size represents the count of different genes and the color indicates significance of
the term enrichment. GO terms with adjusted p-value < 0.05 are significantly enriched. The
kinase term (indicated by an arrow) was exclusively assigned to BIR1 L9 plants. WT, wild-
type Col-0; L9, BIR1 L9 transgenic plants; De, DEX-treated plants; Mo, Mock-treated plants;
_4, 4 days of treatment; _12, 12 days of treatment.

1082

 Figure S2. Transcriptomic profiling of dexamethasone (DEX)-treated wild-type Col-0
 plants.

A) Morphology of representative wild-type (WT) Arabidopsis plants used for RNA-Seq
experiments. Samples were collected after 4 or 12 days of 30 µM DEX treatment and subjected
to RNA sequencing. H$_2$O-treated plants were used as controls. Note that application of DEX
has a negative effect on plant growth as deduced by the significant reduction of rosette size
after 12 days of DEX treatment relative to the non-treated controls. Box and whisker plot
showing 5-95 percentile was analyzed by Mann-Whitney test; ****, p <0.0001. B) Volcano
plots showing the distribution of differentially expressed genes (DEGs) (adjusted p-value $\leq$
0.05). The x-axis shows the gene fold change, and y-axis plots the statistical significance (p-
value) of the differences. Significantly up- and down-regulated genes are highlighted in red and

green, respectively. Genes that are not differentially regulated are in blue. C) Principal component analysis (PCA) of RNA-Seq data. The PCA was performed using normalized RNA-Seq data of DEGs between DEX-treated and mock-treated WT plants after 4 and 12 days of treatment (DeWT_vs MoWT_). Each biological replicate is represented in the score plot. The variance explained by each component (%) is given in parentheses. D) Scatterplot of selected gene ontology (GO) terms of DEGs. GO terms between DEX-treated and mock-treated WT plants at 4 and 12 days of treatment (DeL9_ vs MoL9_) are shown. Dot size represents the count of different genes and the color indicates significance of the term enrichment. GO terms with adjusted p-value < 0.05 are significantly enriched. WT, wild-type Col-0; De, DEX-treated plants; Mo, Mock-treated plants; _4, 4 days of treatment; _12, 12 days of treatment.

Figure S3. Enrichment analysis of Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways using RNA-Seq data.

Scatterplot of selected Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways. Dot size represents the count of different genes and the color indicates significance of the term enrichment. Terms with adjusted p-value < 0.05 are significantly enriched. WT, wild-type Col-0; De, DEX-treated plants; Mo, Mock-treated plants; _4, 4 days of treatment; _12, 12 days of treatment.

Figure S4. Differential expression of receptor-like protein (RLP)-coding genes in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants.

A) Relative expression levels of selected RLP-coding genes based on the number of Fragments Per Kilobase of transcript sequence per Millions of base-pairs sequenced (FPKM) in each of the three independent samples per condition used for RNA-Seq. Significant differences between dexamethasone (DEX)- and mock-treated samples were analyzed using two-way ANOVA followed by Sidak´s multiple comparison test; ***, adjusted p-value < 0.001, **< 0.01. Note that the expression levels of each individual replicate within the DEX-treated BIR1 L9_12 group mirror the abundance of BIR1 transcripts in each replicate. B) Differential expression of selected BIR1-responsive RLP-coding genes based on $\log_2$ fold-change in the indicated pairwise comparisons. WT, wild-type Col-0; L9, BIR1 L9 transgenics; De, DEX-treated plants; Mo, mock-treated plants: _4d, 4 days of treatment; _12d, 12 days of treatment.

Figure S5. Differential expression of cysteine-rich receptor-like kinase (CRK)-coding genes in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants.

A) Relative expression levels of selected CRK-coding genes based on the number of Fragments Per Kilobase of transcript sequence per Millions of base-pairs sequenced (FPKM) in each of the three independent samples per condition used for RNA-Seq. Significant differences between dexamethasone (DEX)- and mock-treated samples were analyzed using two-way ANOVA followed by Sidak´s multiple comparison test; ***, adjusted p-value < 0.001, **< 0.01. Note that the expression levels of each individual replicate within the DEX-treated BIR1 L9_12 group mirror the abundance of BIR1 transcripts in each replicate. B) Differential expression of selected BIR1-responsive CRK-coding genes based on $\log_2$ fold-change in the indicated pairwise comparisons. WT, wild-type Col-0; L9, BIR1 L9 transgenics; De, DEX-treated plants; Mo, mock-treated plants: _4d, 4 days of treatment; _12d, 12 days of treatment.

Figure S6. Differential expression of cell-surface pattern recognition receptors (PRR) and calcium-dependent protein kinases (CPK)-encoding genes in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants.

Relative expression levels of selected PRR- (A) and CPK-coding genes (B) based on the number of Fragments Per Kilobase of transcript sequence per Millions of base-pairs sequenced (FPKM) in each of the three independent samples per condition used for RNA-Seq. Significant differences between dexamethasone (DEX)- and mock-treated samples were analyzed using two-way ANOVA followed by Sidak´s multiple comparison test; ***, adjusted p-value < 0.001, **< 0.01, *< 0.05. WT, wild-type Col-0; L9, BIR1 L9 transgenics; De, DEX-treated plants; Mo, mock-treated plants: _4d, 4 days of treatment; _12d, 12 days of treatment.

Figure S7. Morphological and growth defects in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants of different immune mutant genetic backgrounds.

A) Morphology of transgenic *sobir1-12* BIR1 L8, *bak1-5* BIR1 L4, *eds1*-2 BIR1 L2, *pad4-1* BIR1 L12b and *eds5-3* BIR1 L12 mutant plants from independent lines expressing BIR1 under the control of a dexamethasone (DEX)-inducible 35S promoter. Plants were grown for three weeks on soil and treated with 30 µM DEX to activate BIR1 transcription or mock-treated with

H$_2$O. Plants were photographed after 12 days of the first DEX/ H$_2$O application. B) Morphology

and BIR1 expression levels of DEX- and mock-treated *sobir1-12* BIR1 (L6, L8) and *bak1-5*

BIR1 (L4, L10) plants as observed in different biological replicates of the experiment. qRT-

PCR analysis showed comparable BIR1 transcript levels in both genotypes. Each sample

consisted of RNA pooled from six different plants. C) RT-qPCR analysis of the PTI-marker

WRKY DNA-BINDING PROTEIN 29 (WRKY29) gene in *sobir1-12* BIR1 L6 and *pad4*-1

BIR1 L10 mutant backgrounds. In all cases, transcript accumulation was determined in DEX-

treated (DEX+) versus mock-treated (DEX-) after 12 days of treatment. Values from

biologically independent samples were normalized to the CBP20 internal control, and related

to mock-treated plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test;

*, p <0.05. D) Rosette size (diameter) in wild-type (WT) and BIR1 overexpressing ETI/PTI

mutants after 12 days of DEX treatment relative to the non-treated controls. Box and whisker

plots showing 5-95 percentile were analyzed by Mann-Whitney test; ****, p <0.0001. Note that

rosette size is largely influenced by the effect of DEX on plant development as shown for WT

Col-0 plant in this study.

1173

Table S1. List of primers

Table S2. List of expressed genes

Values represent the total read counts/ Fragments Per Kilobase of transcript sequence per

Millions of base-pairs sequenced (FRKM) values of each biological replicate (a,b,c) per

condition and their corresponding row mean values. De (DEX-treated plants), Mo (Mock-

treated plants), WT (wild-type plants), L9 (BIR1 L9 transgenic plants), _4 (4 days of treatment),

_12 (12 days of treatment).

1181

1182   **Table S3. List of differentially expressed genes (DEG. padjust value < 0.05)**

1183   Values represent the mean Fragments Per Kilobase of transcript sequence per Millions of base-

1184   pairs sequenced (FPKM) values from three biological replicates per condition. De (DEX-

1185   treated plants), Mo (Mock-treated plants), WT (wild-type plants), L9 (BIR1 L9 transgenic

1186   plants), _4 (4 days of treatment), _12 (12 days of treatment).

1187

Table S4. List of differentially expressed genes (DEG. |Log2fold change > 1| padjust value < 0.01)

Values represent the mean Fragments Per Kilobase of transcript sequence per Millions of base-pairs sequenced (FPKM) values from three biological replicates per condition. De (DEX-treated plants), Mo (Mock-treated plants), WT (wild-type plants), L9 (BIR1 L9 transgenic plants), _4 (4 days of treatment), _12 (12 days of treatment).

Table S5. Enrichment analysis of Gene Ontology (GO) terms

CC (cellular component), BP (biological process) and MF (molecular function). Gene Ratio is the ratio of DEGs to all genes for this GO term. Bg Ratio is the ratio of all genes concerning this GO term to all genes in background GO database. Count is the number of differentially expressed genes concerning this GO term. Adjusted p-value < 0.05 means significant enrichment. De (DEX-treated plants), Mo (Mock-treated plants), WT (wild-type plants), L9 (BIR1 L9 transgenic plants), _4 (4 days of treatment), _12 (12 days of treatment).

Table S6. Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis

Gene Ratio is the ratio of DEGs to all genes for this pathway. Bg Ratio is the ratio of all genes concerning this KEGG term to all genes in background KEGG database. Count is the number of differentially expressed genes concerning this term. Adjusted p-value < 0.05 means significant enrichment. De (DEX-treated plants), Mo (Mock-treated plants), WT (wild-type plants), L9 (BIR1 L9 transgenic plants), _4 (4 days of treatment), _12 (12 days of treatment).

Table S7. List of differentially expressed genes (DEG. padjust value < 0.05) within functional categories

Values represent the mean Fragments Per Kilobase of transcript sequence per Millions of base-pairs sequenced (FPKM) values from three biological replicates per condition. De (DEX-treated plants), Mo (Mock-treated plants), WT (wild-type plants), L9 (BIR1 L9 transgenic plants), _4 (4 days of treatment), _12 (12 days of treatment).

Figure 1. Transcriptional reprogramming associated to BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) induction in Arabidopsis.
A) Morphology of representative BIR1 transgenic Arabidopsis plants (line 9, L9) under the control of a dexamethasone (DEX)-inducible 35S promoter used for RNA-Seq experiments. Samples were collected after 4 or 12 days of 30 μM DEX treatment. H_2O-treated plants were used as controls. Relative BIR1 expression based on the number of Fragments Per Kilobase of transcript sequence per Millions of base-pairs sequenced (FPKM) (top) and Western blot using anti-mCherry antibody (bottom) in samples used for RNA-Seq. Significant differences between DEX- and mock-treated samples were analyzed using two-way ANOVA followed by Sidak´s multiple comparison test; ***, adjusted p-value < 0.001. B) Principal component analysis (PCA) of RNA-Seq data. The PCA was performed using normalized RNA-Seq data of differentially expressed genes (DEGs) between BIR1 L9 and wild-type (WT) plants after 4 and 12 days of DEX treatment (DeL9_vs DeWT_). Each biological replicate is represented in the score plot. The variance explained by each component (%) is given in parentheses. The sample (DeL9_12) with differential expression within the DEX-treated L9 group is indicated with black arrows in (a) and (b). C) Number of up-or down-regulated transcripts, with a |$\log_2$(Fold-Change)| ≥1 and adjusted p-value ≤0.01, in each pairwise comparison is indicated. D) Scatterplot of selected gene ontology (GO) terms of DEGs. GO terms between WT and BIR1 L9 plants at 4 and 12 days after DEX-treatment (DeL9_ vs DeWT_) are shown. Dot size represents the count of different genes and the color indicates significance of the term enrichment. GO terms with adjusted p-value < 0.05 are significantly enriched. The kinase term (indicated by an arrow) was exclusively assigned to BIR1 L9 plants. WT, wild-type Col-0; L9, BIR1 L9 transgenic plants; De, DEX-treated plants; Mo, Mock-treated plants; _4, 4 days of treatment; _12, 12 days of treatment.

Figure 2. Effect of BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) induction on immune responses triggered by flg22 and nlp20 treatments.

A) Over time production of reactive oxygen species (ROS) in dexamethasone (DEX)-treated wild-type (WT) and BIR1 L6 and L9 transgenic plants after elicitation of Arabidopsis leaf disks with 1 μM flg22 (left) or H_2O (right). Bars represent means ± SD of relative fluorescence units. B) Ethylene accumulation in BIR1 L6 and L9 transgenic plants after treatment with H_2O (mock), flg22 and nlp20 at 1 μM or 100 nM concentration as indicated. Bars represent average values ± SD. Within each elicitor/mock treatment, differences between DEX-treated (DEX+) and controls (DEX-) were analyzed by two-way ANOVA followed by Sidak´s multiple comparison test. C) RT-qPCR analysis of the PTI-marker genes FLG22-INDUCED RECEPTOR-LIKE KINASE 1 (FRK1) and PATHOGENESIS RELATED 1 (PR1) at 45 min after treatment with H_2O (mock) or 1 μM flg22, and determined in DEX-treated (DEX+) versus non-treated (DEX-) samples. D) Immunoprecipitation of mCherry-BIR1 fusion proteins, indicated with an arrowhead, from samples of BIR1 L6 and L9 plants used in (A) and (B) (left), and RT-qPCR analysis of BIR1 transcripts in DEX-treated (DEX+) samples of BIR1 L6 and L9 plants used in (C) (right). In all cases, RT-qPCR values from biologically independent samples were normalized to the CBP20 internal control, and related to H_2O (mock)-treated plants, set to a value of 1. Data are means ± SD analyzed by Kruskal-Wallis test; *, adjusted *p*-value <0.05. Experiments were repeated three times with similar results.

Figure 3. BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) induction inhibits flg22- and poly(I:C)-triggered callose accumulation at plasmodesmata (PD).

Callose spots at PD were visualized upon aniline blue staining of epidermal cells in response to bacterial flg22 (A) or poly(I:C) (B) in BIR1 L9 transgenic Arabidopsis plants. Plants were previously treated with 30 μM dexamethasone (DEX) to elicit BIR1 expression. Mock-treated plants were used as non BIR1-expressing controls (top panels). Photographs were taken 30 minutes after treatment with H_2O (control) or 1 μM flg22 or 50 ng/μl of poly(I:C). BIR1-mCherry fluorescence in DEX-treated BIR1 L9 plants was confirmed within the same images. Inlays show enlargements of the areas within the dashed boxes. Scale bar, 10 μm. C) Relative PD callose content in flg22/poly(I:C)-treated and H_2O (control)-treated leaves from BIR1 L9 plants previously treated with or without DEX as indicated. Callose content at PD in DEX-treated wild-type (WT) plants in response to poly (I:C) is shown. D) flg22/poly(I:C)-induced callose content at PD in the *sobir1-12* and *bir1-1 sobir1-1* mutants. H_2O-treated controls and callose deposition at PD from WT Col-0 leaves are shown as controls. In all cases, black dots represent the mean values of callose intensities in individual PD (>100) measured in three leaf discs taken from at least three independent biological replicates per each treatment. Significant differences between samples were analyzed by Kruskal-Wallis test or by one-way ANOVA followed by Tukey's multiple comparison test; ****, adjusted *p*-value <0.0001, ***, <0.001, **, <0.01. Experiments were repeated twice with similar results.

Figure 4. BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpression leads to the transcriptional activation of pattern-triggered immunity (PTI)-associated co-receptors and effector-triggered immunity (ETI) regulators.

A) Validation of the RNA-Seq results. Expression of selected BIR1-responsive differentially expressed genes (DEGs) after 12 days of dexamethasone (DEX) treatment based on fold-change values ($\log_2$ DeL9_12/DeWT_12 and $\log_2$ DeL9_12/MoL9_12 as indicated) was confirmed by RT-qPCR. Values from biologically independent samples were normalized to the CBP20 internal control, and related to untreated (DEX-) plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test; *, $p < 0.05$. B) Relative expression of selected PTI/ETI-related genes based on the number of Fragments Per Kilobase of transcript sequence per Millions of base-pairs sequenced (FPKM) in each of the three independent samples per condition used for RNA-Seq. Actin expression is shown as control. Note that the expression levels of each individual replicate within the DEX-treated BIR1 L9_12 group mirror the abundance of BIR1 transcripts in each replicate. Significant differences between mock-treated and DEX-treated samples were analyzed using two-way ANOVA followed by Sidak´s multiple comparison test; ****, adjusted p-value < 0.0001, *** < 0.001, **< 0.01. WT, wild-type Col-0; L9, BIR1 L9 transgenics; _4d, 4 days of treatment; _12d, 12 days of treatment. C) Western blot analysis of SUPPRESSOR OF BIR1-1 (SOBIR1) and PATHOGENESIS RELATED 1 (PR1) using specific antibodies in two independent sets of BIR1 L9 samples. Plants were mock-treated (-) or DEX-treated (+) for 12 days. The presence (+) or absence (-) of morphological phenotypes (Ph) in each sample is indicated. Protein loading control by Ponceau S is shown.

Figure 5. Phenotypes caused by BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpression are partially rescued in *sobir1-12* mutants, but not in a *bak1-5* mutant.

A) Morphology of transgenic *sobir1-12* BIR1 L6 and *bak1*-5 BIR1 L10 mutant plants expressing BIR1 under the control of a dexamethasone (DEX)-inducible 35S promoter. Transgenic BIR1 L9 plants in the wild-type (WT) Col-0 background are shown as a reference. Plants were grown for three weeks on soil and treated with 30 μM dexamethasone (DEX) to induce BIR1 transcription or mock-treated with H$_2$O. Non-transgenic WT Col-0 plants are shown as controls. Plants were photographed after 12 days of the first DEX/ H$_2$O application. Images from independent transgenic lines are shown in Fig. S7. RT-qPCR analysis of *BIR1* transcripts in BIR1 L9 plants are shown. B) RT-qPCR analysis of BIR1 and ENHANCED DISEASE SUSCEPTIBILITY 1 (EDS1) transcripts in *sobir1-12* BIR1 L6 and *bak1*-5 BIR1 L10 plants. C) Western blot analysis of BIR1 and EDS1 proteins with specific antibodies in WT BIR1 L9 and two independent lines of *sobir1-12* BIR1 (L6, L8) and *bak1*-5 BIR1 (L4, L10). Plants were mock-treated (-) or DEX-treated (+) for 12 days as indicated. The presence (+) or absence (-) of morphological phenotypes (Ph) in each sample is indicated. Protein loading control by Ponceau S is shown. RT-qPCR analysis of the PTI-marker genes RESPIRATORY BURST OXIDASE HOMOLOG PROTEIN D (RBOHD), FLG22-INDUCED RECEPTOR-LIKE KINASE 1 (FRK1) or WRKY DNA-BINDING PROTEIN 29 (WRKY29) in *sobir1-12* BIR1 L6 (D) and *bak1*-5 BIR1 L10 (E) plants. RT-qPCR analysis of representative marker genes of the salicylic acid PATHOGENESIS RELATED 1 (PR1), jasmonic acid PLANT DEFENSIN 1.2 (PDF1.2), ethylene PR4 and camalexin [PHYTOALEXIN DEFICIENT 3 (PAD3), CYTOCHROME P450 MONOOXYGENASE 79B2 (CYP79B2)] pathways in *sobir1-12* BIR1 L6 (F) and *bak1*-5 BIR1 L10 (G) plants. In all cases, the same set of replicates from each genotype was used to measure all the genes shown in the graphs. Transcript accumulation was determined in mock-treated (DEX-) versus DEX-treated (DEX+) after 12 days of treatment. Values from biologically independent samples were normalized to the CBP20 internal control, and related to mock-treated plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test; *, $p < 0.05$. Experiments were repeated at least twice with similar results.

Figure 6. Phenotypes caused by BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpression are partially rescued in *eds1-2* mutants, but not in a *pad4-1* mutant.

A) Morphology of transgenic *eds1-2* BIR1 L1b and *pad4-1* BIR1 L0 mutant plants expressing BIR1 under the control of a dexamethasone (DEX)-inducible 35S promoter. Plants were grown for three weeks on soil and treated with 30 µM DEX to induce BIR1 transcription or mock-treated with H_2O. Plants were photographed after 12 days of the first DEX/ H_2O application. Images from independent transgenic lines are shown in Fig. S7. RT-qPCR analysis of BIR1 transcript in *eds1-2* BIR1 L1b and *pad4-1* BIR1 L0 mutant lines are shown. RT-qPCR analysis of the PTI-marker genes RESPIRATORY BURST OXIDASE HOMOLOG PROTEIN D (RBOHD) and FLG22-INDUCED RECEPTOR-LIKE KINASE 1 (FRK1) in *eds1-2* BIR1 L1b (B) and *pad4-1* BIR1 L0 (C) plants. RT-qPCR analysis of representative marker genes of the salicylic acid PATHOGENESIS RELATED 1 (PR1), jasmonic acid PLANT DEFENSIN 1.2 (PDF1.2), ethylene PR4 and camalexin [PHYTOALEXIN DEFICIENT 3 (PAD3)*,* CYTOCHROME P450 MONOOXYGENASE 79B2 (CYP79B2)] pathways in *eds1-2* BIR1 L1b (D) and *pad4-1* BIR1 L0 (E) plants. In all cases, the same set of replicates from each genotype was used to measure all the genes shown in the graphs. Transcript accumulation was determined in mock-treated (DEX-) versus DEX-treated (DEX+) after 12 days of treatment. Values from biologically independent samples were normalized to the CBP20 internal control, and related to mock-treated plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test; *, $p < 0.05$. Experiments were repeated at least twice with similar results.

Figure 7. Phenotypes caused by BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpression are independent of salicylic acid (SA)-mediated signaling pathways.

A) Morphology of transgenic *sid2-2* BIR1 (L5) and *eds5-3* BIR1 (L10) mutant plants expressing BIR1 under the control of a dexamethasone (DEX)-inducible 35S promoter. Plants were grown for three weeks on soil and treated with 30 μM DEX to induce BIR1 transcription or mock-treated with H_2O. Plants were photographed after 12 days of the first DEX/ H_2O application. Images from independent transgenic lines are shown in Fig. S7a, b. RT-qPCR analysis of BIR1 and representative marker genes of the jasmonic acid PLANT DEFENSIN 1.2 (PDF1.2) and camalexin [PHYTOALEXIN DEFICIENT 3 (PAD3), CYTOCHROME P450 MONOOXYGENASE 79B2 (CYP79B2)] pathways in *sid2-2* BIR1 (L5) (B) and *eds5-3* BIR1 (L10) (C) mutant lines. In all cases, the same set of replicates from each genotype was used to measure all the genes shown in the graphs. Transcript accumulation was determined in mock-treated (DEX-) versus DEX-treated (DEX+) after 12 days of treatment. Values from biologically independent samples were normalized to the *CBP20* internal control, and related to mock-treated plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test; *, $p < 0.05$. Experiments were repeated at least twice with similar results. D) Tentative working model for BIR1 homeostasis. SA-mediated transcription of BIR1 is regulated by RNA-directed DNA methylation (RdDM) and post-transcriptional RNA silencing (PTGS) (Guzman-Benito *et al.*, 2019). In these conditions, BIR1 induction interferes with PTI gene expression and PD callose deposition triggered by various elicitors (flg22, nlp20, dsRNA). Loss of BIR1 regulation leads to constitutive BIR1 induction and ultimately BIR1 overexpression at levels that cause SA-independent up-regulation of immune and hormone genes. BIR1 overexpression is sensed by one or several TNLs, of which CSA1 is a plausible candidate, to promote PAD4-independent ETI-type cell death that requires EDS1 and SOBIR1.

Figure S1. Transcriptomic profiling of BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) expressing transgenic lines.
A) Principal component analysis (PCA) of RNA-Seq data. The PCA was performed using normalized RNA-Seq data of differentially expressed genes (DEGs) between dexamethasone (DEX)-treated and mock-treated transgenic BIR1 L9 plants after 4 and 12 days of treatment (DeL9_vs MoL9_). Each biological replicate is represented in the score plot. The variance explained by each component (%) is given in parentheses. B) Number of up-or down-regulated transcripts, with a |$\log_2$ (Fold-Change)| ≥1 and adjusted p-value ≤0.05, in each pairwise comparison as indicated. C) Volcano plots showing the distribution of DEGs (adjusted p-value ≤ 0.05). The x-axis shows the fold change in gene expression between samples, and y-axis shows the statistical significance of the differences. Significantly up- and down-regulated genes are highlighted in red and green, respectively. Genes that are not differentially regulated are in blue. D) Scatterplot of selected gene ontology (GO) terms of DEGs. GO terms between wild-type (WT) and BIR1 L9 plants at 4 and 12 days after DEX-treatment (DeL9_ vs MoL9_) are shown. Dot size represents the count of different genes and the color indicates significance of the term enrichment. GO terms with adjusted p-value < 0.05 are significantly enriched. The kinase term (indicated by an arrow) was exclusively assigned to BIR1 L9 plants. WT, wild-type Col-0; L9, BIR1 L9 transgenic plants; De, DEX-treated plants; Mo, Mock-treated plants; _4, 4 days of treatment; _12, 12 days of treatment.

Figure S2. Transcriptomic profiling of dexamethasone (DEX)-treated wild-type Col-0 plants.
A) Morphology of representative wild-type (WT) Arabidopsis plants used for RNA-Seq experiments. Samples were collected after 4 or 12 days of 30 µM DEX treatment and subjected to RNA sequencing. H_2O-treated plants were used as controls. Note that application of DEX has a negative effect on plant growth as deduced by the significant reduction of rosette size after 12 days of DEX treatment relative to the non-treated controls. Box and whisker plot showing 5-95 percentile was analyzed by Mann-Whitney test; ****, $p < 0.0001$. B) Volcano plots showing the distribution of differentially expressed genes (DEGs) (adjusted p-value ≤ 0.05). The x-axis shows the gene fold change, and y-axis plots the statistical significance (p-value) of the differences. Significantly up- and down-regulated genes are highlighted in red and green, respectively. Genes that are not differentially regulated are in blue. C) Principal component analysis (PCA) of RNA-Seq data. The PCA was performed using normalized RNA-Seq data of DEGs between DEX-treated and mock-treated WT plants after 4 and 12 days of treatment (DeWT_vs MoWT_). Each biological replicate is represented in the score plot. The variance explained by each component (%) is given in parentheses. D) Scatterplot of selected gene ontology (GO) terms of DEGs. GO terms between DEX-treated and mock-treated WT plants at 4 and 12 days of treatment (DeL9_ vs MoL9_) are shown. Dot size represents the count of different genes and the color indicates significance of the term enrichment. GO terms with adjusted p-value < 0.05 are significantly enriched. WT, wild-type Col-0; De, DEX-treated plants; Mo, Mock-treated plants; _4, 4 days of treatment; _12, 12 days of treatment.

Figure S3. Enrichment analysis of Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways using RNA-Seq data.
Scatterplot of selected Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways. Dot size represents the count of different genes and the color indicates significance of the term enrichment. Terms with adjusted *p*-value < 0.05 are significantly enriched. WT, wild-type Col-0; De, DEX-treated plants; Mo, Mock-treated plants; _4, 4 days of treatment; _12, 12 days of treatment.

Figure S4. Differential expression of receptor-like protein (RLP)-coding genes in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants.

A) Relative expression levels of selected RLP-coding genes based on the number of Fragments Per Kilobase of transcript sequence per Millions of base-pairs sequenced (FPKM) in each of the three independent samples per condition used for RNA-Seq. Significant differences between dexamethasone (DEX)- and mock-treated samples were analyzed using two-way ANOVA followed by Sidak´s multiple comparison test; ***, adjusted p-value < 0.001, **< 0.01. Note that the expression levels of each individual replicate within the DEX-treated BIR1 L9_12 group mirror the abundance of BIR1 transcripts in each replicate. B) Differential expression of selected BIR1-responsive RLP-coding genes based on $\log_2$ fold-change in the indicated pairwise comparisons. WT, wild-type Col-0; L9, BIR1 L9 transgenics; De, DEX-treated plants; Mo, mock-treated plants: _4d, 4 days of treatment; _12d, 12 days of treatment.

Figure S5. Differential expression of cysteine-rich receptor-like kinase (CRK)-coding genes in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants.

A) Relative expression levels of selected CRK-coding genes based on the number of Fragments Per Kilobase of transcript sequence per Millions of base-pairs sequenced (FPKM) in each of the three independent samples per condition used for RNA-Seq. Significant differences between dexamethasone (DEX)- and mock-treated samples were analyzed using two-way ANOVA followed by Sidak′s multiple comparison test; ***, adjusted p-value < 0.001, **< 0.01. Note that the expression levels of each individual replicate within the DEX-treated BIR1 L9_12 group mirror the abundance of BIR1 transcripts in each replicate. B) Differential expression of selected BIR1-responsive CRK-coding genes based on log$_2$ fold-change in the indicated pairwise comparisons. WT, wild-type Col-0; L9, BIR1 L9 transgenics; De, DEX-treated plants; Mo, mock-treated plants: _4d, 4 days of treatment; _12d, 12 days of treatment.

Figure S6. Differential expression of cell-surface pattern recognition receptors (PRR) and calcium-dependent protein kinases (CPK)-encoding genes in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants.

Relative expression levels of selected PRR- (A) and CPK-coding genes (B) based on the number of Fragments Per Kilobase of transcript sequence per Millions of base-pairs sequenced (FPKM) in each of the three independent samples per condition used for RNA-Seq. Significant differences between dexamethasone (DEX)- and mock-treated samples were analyzed using two-way ANOVA followed by Sidak´s multiple comparison test; ***, adjusted p-value < 0.001, **< 0.01, *< 0.05. WT, wild-type Col-0; L9, BIR1 L9 transgenics; De, DEX-treated plants; Mo, mock-treated plants: _4d, 4 days of treatment; _12d, 12 days of treatment.

Figure S7. Morphological and growth defects in BAK1-INTERACTING RECEPTOR-LIKE KINASE 1 (BIR1) overexpressing plants of different immune mutant genetic backgrounds.

A) Morphology of transgenic *sobir1-12* BIR1 L8, *bak1-5* BIR1 L4, *eds1*-2 BIR1 L2, *pad4-1* BIR1 L12b and *eds5-3* BIR1 L12 mutant plants from independent lines expressing BIR1 under the control of a dexamethasone (DEX)-inducible 35S promoter. Plants were grown for three weeks on soil and treated with 30 µM DEX to activate BIR1 transcription or mock-treated with H_2O. Plants were photographed after 12 days of the first DEX/ H_2O application. B) Morphology and BIR1 expression levels of DEX- and mock-treated *sobir1-12* BIR1 (L6, L8) and *bak1-5* BIR1 (L4, L10) plants as observed in different biological replicates of the experiment. qRT-PCR analysis showed comparable BIR1 transcript levels in both genotypes. Each sample consisted of RNA pooled from six different plants. C) RT-qPCR analysis of the PTI-marker WRKY DNA-BINDING PROTEIN 29 (WRKY29) gene in *sobir1-12* BIR1 L6 and *pad4*-1 BIR1 L10 mutant backgrounds. In all cases, transcript accumulation was determined in DEX-treated (DEX+) versus mock-treated (DEX-) after 12 days of treatment. Values from biologically independent samples were normalized to the CBP20 internal control, and related to mock-treated plants, set to a value of 1. Data are mean ± SD analyzed by Mann-Whitney test; *, p <0.05. D) Rosette size (diameter) in wild-type (WT) and BIR1 overexpressing ETI/PTI mutants after 12 days of DEX treatment relative to the non-treated controls. Box and whisker plots showing 5-95 percentile were analyzed by Mann-Whitney test; ****, p <0.0001. Note that rosette size is largely influenced by the effect of DEX on plant development as shown for WT Col-0 plant in this study.