TITLE:

The haplolethal gene *wupA* of Drosophila exhibits potential as a target

for an X-poisoning gene drive.

Clancy D. Lawler, Natalia Hernandes, Ana Karla Parra Nuñez, Soumitra Bhide,

Simon Baxter and Charles Robin*

School of BioSciences, The University of Melbourne, Victoria, Australia

*Corresponding author:

Charles Robin

Building 184, School of BioSciences, Parkville Campus,

The University of Melbourne, Victoria 3010 Australia

crobin@unimelb.edu.au

Running head: A target for a haplolethal gene drive

Abstract

A synthetic gene drive that targets haplolethal genes on the X-chromosome can skew the sex ratio towards males. Like an 'X-shredder' strategy it does not involve 'homing' and that has advantages including the reduction in gene drive resistance allele formation. We examine this 'X-poisoning' strategy by targeting four of the 11 known X-linked haplolethal/haplosterile genes of *Drosophila melanogaster* with CRISPR/cas9. We find that targeting the *wupA* gene during spermatogenesis skews the sex ratio so fewer than 14% of progeny are daughters. That is unless we cross the mutagenic males to X^XY female flies that bear attached-X chromosomes, which reverses the inheritance of the poisoned X chromosome so that sons inherit it from their mother; in which case only 2% of the progeny are sons. These sex ratio biases suggests that most of the CRISPR/cas9 mutants we induced in the *wupA* gene are haplolethal but some are recessive lethal. We have used simulations to model release strategies of a gene drive where an X-poisoning Y chromosome is introduced to invasive populations of the pest *Drosophila suzukii*. While the reproductive output of carriers of X-poisoning drives is less than that of X-shredding drives due to their post-zygotic mode of action, the threshold dynamics of an X-poisoning drive result in more predictable outcomes than X-shredders that tend to exhibit 'chasing' more often. Like *D. melanogaster* and *D. suzukii*, the *wupA* orthologs in *Anopheles* mosquitos are found on X chromosomes making it a viable target in multiple species.

Introduction

For many decades it has been envisioned that insect populations that vector

diseases of humans and animals or that are pests of food crops could be controlled

using genetic methods (Serebrovsky 1969; Knipling 1955; Curtis 1968; Whitten 1985;

Burt 2003). Indeed, Sterile Insect Techniques have successfully controlled fly species

(Bushland, Lindquist, and Knipling 1955; Hendrichs, Franz, and Rendon 1995; Scott

et al. 2017) and field trials relying on self-limiting transgenes are currently being

deployed and assessed (Yao *et al.* 2022; Waltz 2021). Recently, the prospect of

synthetic gene drives, a distinct class of population manipulation, where transgenes

can be engineered to spread through otherwise natural populations of insect pests has

received much attention (Alphey *et al.* 2020; Hay, Oberhofer, and Guo 2021; Champer,

Buchman, and Akbari 2016). The effectiveness of various gene drive designs, many of

which use CRISPR/cas9 machinery, have been assessed in laboratory populations

(Chen *et al.* 2007; N. Windbichler *et al.* 2007; Akbari *et al.* 2014; Kyrou *et al.* 2018;

Oberhofer, Ivy, and Hay 2019; Webster, Vella, and Scott 2020; Yang *et al.* 2022). Many

of these require 'homing' where double-stranded DNA breaks must be repaired by

homologous recombination (HR) so that the transgenic selfish elements are copied

and thereby spread. A drawback of such homing designs has been that some double

stranded breaks can be repaired using the non-homologous end-joining (NHEJ)

pathway. In these instances not only does homing fail but mutations that are resistant

to subsequent CRISPR/cas9 cleavages can be introduced into the population (Nikolai

Windbichler *et al.* 2011; Gratz *et al.* 2014; Champer *et al.* 2017; 2018; Hammond *et al.*

2017).

One gene drive design that does not require homing is referred to as the X-

shredder (Galizi *et al.* 2014). It targets highly repeated sequences on the X

chromosome during spermatogenesis such that X-bearing sperm are inviable, and the

drive skews the sex ratio towards males. If the X-shredding transgenes could be placed

on the Y chromosome, then the pattern of fathers-only-having-sons will mean the sex

ratio will become more biased over the generations and the population will approach

extinction. Originally, the X-shredder was attempted in lab populations of *Anopheles*

gambiae mosquito using the endonuclease *PpoI*, but subsequent implementations in

A. gambiae mosquitos, *Drosophila melanogaster* and *Ceratitis capitata* fruit flies have

deployed CRISPR/cas9 machinery (Galizi *et al.* 2014; Fasulo *et al.* 2020; Meccariello

et al. 2021).

A related approach, dubbed 'X-poisoning', also targets loci on the X

chromosome; not so the chromosome is physically destroyed but so that mutations are

introduced such that progeny receiving the X chromosome will not be viable (Fasulo *et*

al. 2020). This strategy uses CRISPR/Cas9 to target X-linked haplolethal genes.

Although haplolethal genes are typically thought of as genes where both alleles of a

diploid organism need to be functional, X-linked haplolethal genes can occur in many

dipterans because their dosage compensation mechanism elevates the X

chromosome-encoded output in males (that is in contrast to eutherian mammals where

there is silencing of X loci in females and therefore haplolethals on the X would be

incompatible with maleness) (Rose *et al.* 2016; Cook *et al.* 2012). There may be a few

potential advantages of X-poisoning over X-shredding. Firstly, X-poisoning does not

rely on natural repeats occurring exclusively on the X chromosome and theoretically

this may increase its applicability to some species. Secondly, CRISPR cutting in one

or a few sites on the X chromosome might occur more readily than cutting the X

chromosome at an overwhelming number of places, as is required for X-shredding.

Thirdly, X-shredding has been shown to be ameliorated by the NHEJ pathway thus

potentially limiting its utility (Fasulo *et al.* 2020). Fourthly, as we explore in this

manuscript, it may be that post-zygotic effects of X-poisoning may change the

threshold dynamics of gene drive invasion.

While informative, laboratory cage experiments can fail to incorporate

population dynamics of natural populations. Therefore, computational simulations are

used to estimate the impact of drift, migration, mutation, relative fitness, population

structure and environmental stochasticity upon drive outcome. For example,

seasonality (Eckhoff *et al.* 2017), release range (North, Burt, and Godfray 2013) and

spatial structure have been shown to alter simulation outcomes. While simulations of

panmictic populations predict a suppression drive will either fix, be lost, or reach a

stable equilibrium (Burt 2003; Larner *et al.* 2019; Prowse *et al.* 2017), spatially explicit

simulations show that a suppression gene drive in continuous space may undergo

cyclic patterns of local population elimination followed by recruitment of both wildtype
and gene drive individuals through immigration (Champer *et al.* 2021; Paril and Phillips
2022). This may lead to unstable drive frequencies and cyclic subpopulation
elimination and recolonisation that has been termed a gene drive 'chase' (Champer *et*
al. 2021). This instability, and unpredictability, is likely to be an undesirable feature of
a population control system as it may lead to ambiguity in drive frequencies and long-
term persistence of the gene drive. Gene drives with strong threshold dependent
dynamics may be less likely to undergo a population chase, as the required threshold
to drive through the population is unlikely to be reached by migration alone (Buchman
et al. 2018; Sánchez C. *et al.* 2020). How these dynamics are likely to play out in an
expanding population is currently not well studied.

Here we describe the targeting of haplolethal genes on the X chromosome of
Drosophila melanogaster with CRISPR/cas9. The loci differ from the two targeted in
the X-poisoning study of Fasulo *et al.* (2020) as we examine three other ribosomal
protein encoding genes and the haplolethal gene *Wings up A (wupA)*. In contrast to
the ribosomal protein genes which might be required for the viability of most cell types
including those in spermatogenesis, *wupA*, encodes a troponin that is expressed in
muscles from mid embryogenesis. We therefore sought experimental affirmation that
these genes are indeed haplolethal genes and whether they could be valuable targets
for X-poisoning gene drives.   We then simulate scenarios where X-poisoning
constructs are hypothetically placed on the Y chromosome so that they drive through
populations. As the postzygotic lethality of many haplolethal mutations is likely to
reduce the output of fertile offspring for drive carrying males (Prado, Canal, and Ferrús
1999; Cook *et al.* 2012), we model the drive's dynamics in comparison to the pre-
zygotic X-shredder drive. The simulations model a hypothetical incursion of *Drosophila*
suzukii, an agricultural pest of global significance closely related to *D. melanogaster*
(Mazzi *et al.* 2017; Ørsted and Ørsted 2019; Yeh *et al.* 2020; Knapp, Mazzi, and Finger
2021). Our spatially explicit models of an X-poisoning gene drive and an X-shredder
gene drive were designed to answer three main questions. Firstly, how do these two
population control methods function in a spatially explicit expanding population?
Secondly, under what circumstances, if any, does an X-poisoning drive outperform an
X-shredder drive? Thirdly, how do changes in release strategy impact each of the gene
drives?

Materials and Methods

gRNA design and cloning

A combination of ChopChop v3 (Labun *et al.* 2019) and CRISPR Optimal Target Finder (Gratz *et al.* 2014) was used to identify the gRNA target sites with the highest efficiency scores with a low likelihood of off-target activity. All gRNA sequences were checked against the DGRP reference panel using FlyBase's JBrowse (Larkin *et al.* 2021) to avoid standing variation at the target site that may give natural resistance.

The design protocol of (Port and Bullock 2016) was used to clone gRNA sequences into the pCFD5_w plasmid (Addgene #112645). Three separate plasmids were generated that targeted haplolethal genes on the X chromosome; pRpS19a_RpL35, pRpL36_RpL35, and p*wupA*. pRpS19a_RpL35 encoded four gRNAs, two targeting *RpS19a* (*RpS19*_RNA1: GAAGGATATTGACCAGCACG, *RpS19*_RNA2: AACCGACTCCAGCGGGACTG) and two targeting *RpL35* (*RpL35*_gRNA1: GTGCTCCGAGCTGAGGATCA, *RpL35*_gRNA2: CACAATGTAGACGCGAGCGA). pRpL36_RpL35 encodes four gRNA, two targeting *RpL36* (*RpL36*_gRNA1: GCTGGCTATTGGCCTGAACA, *RpL36*_gRNA2: CATGCGCGACTTGGTCCGCG) and two targeting *RpL35* (as aforementioned). Each of the three sgRNA's used to target *wupA* (*wupA*_gRNA1: ACCAAAAACACAAATCAAAA, *wupA*_gRNA2: TGAGGTGCGCAAGCGCATGG, *wupA*_gRNA3: CGCATCATCGAAGAACGTTG) would affect all 13 alternate transcripts annotated in Flybase. The PAM site for *wupA*_gRNA1 corresponds to the most common start codon and is in the 5'UTR of transcripts using the alternate start codon, the PAM site for *wupA*_gRNA2 corresponds to the alternate start methionine which is a codon used by in all transcripts, and that for *wupA*_gRNA3 is in a coding exon common to all transcripts (Figure 1). All primers used and sgRNA sequences are tabulated in Table S1.

Microinjections

The pRpS19a+RpL35 and p*wupA* were microinjected into the 09w fly line while the pRpL36+RpL35 was injected into the 10w fly line. These two fly lines were obtained from Trent Perry, Bio21 Unimelb who had generated them using the Bloomington stocks #3605 (w[1118]) and #25709 (y[1] v[1] P{y[+t7.7]=nos-phiC31\int.NLS}X;

177 P{y[+t7.7]=CaryP} attP40) and #25710 (P{y[+t7.7]=nos-phiC31\int.NLS}X, y[1] sc[1]

178 v[1] sev[21]; P{y[+t7.7]=CaryP}attP2) to generate second chromosome landing site

179 flies (09W) and third chromosome landing site flies (10W) in a *white* null background.

180 Insertion events were validated by sequencing and single integration lines were

181 maintained.

Assessing sex ratio biasing capabilities of sgRNA constructs

To assess the sex ratio biasing rates of each of the sgRNA expressing

constructs, we first generated male flies that expressed both Cas9 and sgRNAs in their

germ cells. Individual females homozygous for each of the three sgRNA constructs

(RpS19a+RpL35, RpL36+RpL35 and *wupA*) were crossed to males containing Cas9

under the control of the *nanos* promoter (Either NIG CAS-0001; y^2 cho^2 v^1; attP40{nos-

Cas9}/CyO which carries a recessive lethal allele on the second chromosome and

cannot be made homozygous or a line we generated that has {nos-Cas9}attP40 placed

in the 09w second chromosome attP40 landing site(referred to herein as NH-

nosCAS9). The paternal inheritance of cas9 prevented maternal deposition and

somatic activity of the drive (Champer *et al.* 2019). Initially, single males heterozygous

for both cas9 and one of the sgRNA constructs were crossed to three 09W 3–4-day

old virgins and allowed to mate and lay in a vial for seven days before being cleared

and all offspring that emerged counted. Subsequent crosses included single nos-

cas9/Wup-sgRNA males left to mate with single females for four hours, or alternatively

crossed to three-day-old virgin females with the genotype C(1)DX,y^1,f^1/Y (Bloomington

Stock number 4559), which is an 'attached-X' line forces paternal X chromosome

inheritance.

Simulations

All models were generated using SLiM 3.6, an individual-based forward

population genetic simulation program (Haller and Messer 2019) and run on the

Spartan cluster at the University of Melbourne. The simulations were designed to

replicate the biology of *Drosophila suzukii* (Supplementary Table S2; Krüger *et al.*

2019; Lanouette *et al.* 2020) with some parameters taken from *Drosophila*

melanogaster data (Morrow, Stewart, and Rice 2005; Schnakenberg, Siegal, and

Bloch Qazi 2012), and with overlapping generations and an expanding population in a

two-dimensional space. Time was measured in generations with each generation

being the length of time taken for an embryo to reach reproductive maturity. To

simulate the expansion of the invading population, wild-type individuals started within

one generation's movement of the XY coordinates (0,0) within the square two-

dimensional space (Video S1-S8). The maximum per generation movement within the

simulated two-dimensional square varied as a test parameter with values between low,

medium-low, medium-high, and high. The per generation movement varied around a

normal distribution with standard deviation as a percentage of the entire simulated area

– high per-generation movement results in approximately 95% of individuals moving

~0-6% of the entire simulated area and the low per-generation movement speed

resulting in approximately 95% of individuals moving between ~0-4% of the entire

simulated area per-generation along either axis with medium-low and medium-high

sitting equidistant between these values. Both the dynamics of a X-poisoning drive and

a X-shredder drive were tested within the context of these simulations.

Four release strategies, each occurring at least 10 generations after the

population started its invasion into the space, were implemented for each of the two

gene drive systems:

1. The total number of gene drive individuals were released once uniformly over

the entire occupied area.

2. The total number of gene drive individuals were released only at the point where

the invasive population originated.

3. A proportion of the total number of gene drive individuals were released

uniformly over the entire inhabited area every multiple of the release generation.

4. A proportion of the total number of gene drive individuals were released only at

the point where the invasive population originated every multiple of the release

generation.

The number of released individuals varied between 100 and 993. The

generation of the gene drive release varied between generation 10 and 30 with the

multiple release simulations occurring at an interval of between 1 and 20 generation

between generation 10 and generation 110 – all simulations have a 10 generation burn

in period with generations counted after this initial period such that release generation

measures the number of generations after the burn in period. Both the multiple interval-

based release and the single release strategies maintain the same total number of

released individuals – either the total release number is released in a single go, or a

proportion of the total release number is released every generation interval. The X-
shredder gene drive was encoded so that the sex ratio of the offspring was biased
towards males whereas the X-poisoning drive resulted in females (born to a drive
containing father) being made non-viable before they reached sexual maturity –
resulting in post zygotic lethality. The proportion of 100 replicate simulations that led to
population eradication within 110 generations was assessed. For further information
on simulation variables see Supplementary Table S2 and for the simulation code see
files S1 and S2.

Statistical Analyses

All statistical analyses were undertaken in R (R Core team 2023). Sex ratio biasing
in the *Drosophila* X-poisoning gene drive crosses were analysed using an unpaired
two-tail t-test with a Bonferroni correction for multiple comparisons where applicable.
Simulation data was generated using Spartan cluster at the University of Melbourne.
Each combination of variables was independently run 100 times and the average
outcome shown in a heatmap. The R code used to analyse the data is available in
file S3.

Results

260
Four of the haplolethal genes described on the X chromosome of *Drosophila*
*melanogaste*r (Cook *et al.* 2012) were targeted using three multi-sgRNA constructs
and cas9 expressed in the male germline via a *nanos* promoter (Figure 1). The
expectation was that males carrying the cas9 and sgRNA transgenes would have a
reduced number of viable daughters relative to controls yet the same number of sons
(Figure 2a).   All the crosses targeting ribosomal protein encoding genes
(RpS19a+RpL35, RpL36+RpL35) yielded significantly fewer offspring when compared
to the controls ($p<0.0001$, t-test with Bonferroni correction; Figure 2b). The crosses
involving the RpS19a+RpL35 sgRNA construct followed expectations with significantly
fewer daughters than sons ($p<0.0001$, t-test with Bonferroni correction) and no
significant difference in the number of sons compared to the control. In contrast the
RpL36+RpL35 crosses yielded significantly fewer sons *and* daughters compared to

control ($p<0.05$ and $p<0.01$ respectively, t-test with Bonferroni correction) without a

significant sex ratio bias.

When the CRISPR machinery was targeted to *wupA* significant skews in sex ratio

towards maleness was observed ($p<0.0001$, t-test with Bonferroni correction, Fig. 2).

However, we initially found the expression of sgRNAs targeting *wupA* did not result in

a significant reduction in the total number of offspring produced when compared to

controls (Fig. 2b). The highly significant reduction in the number of female offspring

($p<0.0001$, t-test with Bonferroni correction), was offset by the significantly higher

number of male offspring compared to control ($p<0.0001$, t-test with Bonferroni

correction). We hypothesised that the lack of reduction in reproductive output may be

due to a carrying capacity within the vials with an excess of embryos being laid during

the seven day period where remating was allowed. To address this possibility, single

males heterozygous for both X-poisoning drive components (ie. the *wupA* targetting

sgRNAs and nos-cas9) were placed in vials with single females for four hours only. In

these single pair crosses, a significant reduction in the total number of offspring

($p<0.01$, t-test with Bonferroni correction) as well as a highly significant change in sex

ratio was observed ($p<0.0001$, t-test with Bonferroni correction; Figure 2c) and there

was no significant difference in the number of males produced when compared to the

control.

To understand why the sex biasing was not absolute, X-chromosomes that had

survived being passed through the paternal germ line were characterized. If the

targeted loci had been cut and reannealed, indels or single nucleotide mutations may

have been introduced, and they would be revealed by sequence analysis at the sites

targeted by the sgRNA's. To simplify the sequence analysis, we used X^XY females

which reverse the inheritance of the paternal X's so that they are passed to sons.

Consistent with expectations, males expressing *nanos*-Cas9 and sgRNA's targeting

wupA sired significantly more daughters than sons (the sex ratio bias being flipped due

to sons paternally inheriting the X chromosome; $p<0.001$, t-test with Bonferroni

correction). The RpS19a+RpL35 cross showed no significant sex ratio bias but had

significantly fewer male offspring than the control and no significant difference in the

number of females (Figure 2d; $p<0.0001$, t-test with Bonferroni correction). There was

no reduction in the number of offspring produced by males expressing gRNA targeting

RpL36+RpL35.

The X chromosomes of a total of 60 surviving sons (14 for *wupA*, 22 for
RpS19a+RpL35 and 24 for RpL36+RpL35) from 15 independent crosses were
sequenced and only the offspring of one male showed any sign of CRISPR cutting. A
complex indel resulting in a net 9 nucleotide insertion that did not introduce a premature
stop codon was recovered from a cross involving the RpL36+RpL35 sgRNAs (Figure
S1). The fact that this individual survived suggests that gene drive resistance alleles
could arise against RpL35 gRNA1. The data also concurs with the idea that these
genes are indeed lethal genes as no frameshifting indels or nonsense mutations were
recovered. More generally, these data are consistent with the notion that most of the
surviving progeny carrying a paternal X never had their X chromosome cut by the
CRISPR machinery.

The above results suggest that *wupA* could be a good target for an X-poisoning gene
drive. To create such a drive, transgenes encoding cas9 and the sgRNA's targeting
wupA would ideally be placed on the Y chromosome. Placing transgenes on the Y
chromosome has been technically challenging although it has been done (Gamez *et*
al. 2021). Indeed, we examined the one *D. melanogaster* line currently available that
has a cas9 gene on the Y chromosome (BL91386). The cas9 of this line uses a *vasa*
promoter and this expresses outside the germline as well as in it. A cross between
males with this vasa-cas9 Y chromosome with females bearing *wupA* targeting
sgRNA's yielded almost entirely daughters (264 females and two males that were small
and infertile). Presumably the males that would have cas9 and sgRNA expression in
their germline die before they reach adulthood because somatic expression of the
vasa-cas9 with the ubiquitously expressed *wupA* sgRNA's is lethal.

However, as an X-poisoning Y chromosome could in theory be built with a cas9
restricted to the germline, and as the *wupA* gene is found on the X chromosome of
other Drosophila species including the pest species *Drosophila suzukii*, we set out to
model an X-poisoning strategy deployed in a hypothetical invasion of *D. suzukii*.

Simulations of gene drives in an expanding population

Drosophila suzukii is an invasive species that has recently spread throughout
the United States and Europe and it has the potential to spread in Australia if an
incursion establishes (Maino, Schouten, and Umina 2021). We therefore developed
spatially explicit models (in SLiM; Haller and Messer 2019) to explore the dynamics of

an X-poisoning drive and an X-shredder drive in expanding populations. The models

incorporated parameters that attempted to capture key aspects of *D. suzukii* biology

such as seasonal fluctuations in population size (Stockton, Wallingford, and Loeb

2018). The simulated space was a two-dimensional square with the invasive

population expanding from the bottom left corner (videos S1-S8). Populations differing

in movement rates were simulated and gene drive flies were released a varying

number of generations after the incursion and at various release sizes (Figure 3). We

contrasted models that used X-poisoning, in which daughters arising from gene drive

males did not reach reproductive maturity, to X-shredding where fathers only produced

Y-bearing sperm. As the former is post-zygotic and the former pre-zygotic, X-poisoning

drives had half the offspring of X-shredding drives.

We first modelled the scenario where gene drive individuals were released only

once at the incursion point. The X-poisoning drive failed to eradicate the population

under all tested parameters while the X-shredder drive showed increased eradication

likelihood with increased movement of individuals (Figure 3a). In these simulations the

X-shredder is largely unimpacted by 'release size' and 'release generation', although

a mild seasonality effect is observed in the later (see below). As the degree of

movement affects the distance from which a mate can be selected, it impacts geneflow

and metapopulation dynamics; low movement of individuals results in small

populations with uninhabited areas between them. If suppression occurs more quickly

than gene flow between the subpopulations, the X-shredder is unable to suppress the

entire metapopulation. As a result, the likelihood of eradication given a single release

at the incursion point is largely dependent on the degree of movement and the resulting

population geneflow (Figure 3a).

Next, we modelled multiple releases of gene drive individuals at the incursion

point. That strategy slightly increases the likelihood of eradication for the X-shredder

while maintaining the relationship between eradication likelihood and movement

(Figure 3b). Note that the same number of gene drives individuals were released as in

the 'single release', it was just that they were distributed over multiple releases.

We then simulated the release of gene drive individuals across the range of the

invasive species rather than just at the point of origin and found that to be a much more

effective for both drive types in both single and multiple releases (Figures 3c and 3d).

The simulated winter event impacts the likely outcome for both gene drives in these
release strategies, appearing as a striated change in eradication likelihood at release
generation 1, 10 and 20. In the simulated winters that occur every ten generations, no
offspring are produced, and spatial competition is increased by 60%. Releasing the X-
poisoning drive individuals uniformly over the invaded area immediately after incursion
identification (immediately after the ten generation burn in period), or timing it with the
fluctuations in population size, due to the effects, of winter gives the greatest likelihood
of eradication (Figure 3c and d). The X-shredder drive shows similar dynamics during
low movement speeds but is less impacted by release time at high movement speeds
for the multiple release strategy.

X-poisoning appears to be as good as X-shredding when releases occur across
the range and the movement of individuals is low and releases are well timed relative
to the population declines in winter (Figure 3c and 3d). The inability of the X-poisoning
drive to function at high movement speeds, or when released only at the incursion
point, is indicative of a threshold dynamic within this drive. Unlike the X-shredder drive,
the X-poisoning drive is unable to reach the threshold required to drive through the
population by migration alone resulting in the threshold release rates required for
eradication. The X-shredder is seen to have no threshold and the high degree of gene
flow in the simulations with high movement enables the drive to eradicate the
population more effectively.

Empirically informed sex ratio biasing:

In the above simulations the efficiency of the gene drives were maximized; the only
fertile progeny of drive-bearing fathers, are sons. Next, we altered the simulations to
incorporate more realistic gene drive efficiencies. These used the likelihood of female
survival seen in the single mate crosses described in the X-poisoning crosses above
(i.e. proportion of surviving females = 0.13), or the X-shredder drive sex ratio biasing
rates observed by Meccariello *et al.* (2021) in *Ceratitis capitata* X-shredder (biasing
offspring sex ratio towards male by 80%). In general, such changes made X-shredding
as ineffective as X-poisoning given medium-high to high movement speed – the X-
shredder drive still outperforms the X-poisoning drive at low movement speeds.
However little effect was seen in single uniform release scenarios (compare Figure 3c
to Figure 4a).

Gene drive chases:

For many of the simulations, especially for X-shredder gene drive models,
populations were often observed to undergo cyclic fluctuations in population size and
gene drive proportion. This was due to local eradication events, followed by
recolonisation and subsequent reintroduction of the drive through migration – a gene
drive chase. To identify the frequency of a chase occurring, the simulated space was
split into 100 equal cells within the two-dimensional space. A chase was deemed to
occur in a simulation if any cell went twice through the cycle of colonization and loss.
As multiple releases could result in the misclassification, due to drive individuals being
released into a cell rather than migrating into it, only single release strategies were
examined. These simulations revealed that the likelihood of the X-shredder drive
undergoing a gene drive chase was largely independent of release size, with release
generation only having a minimal impact around winter (Figure 4b). As with eradication
likelihood, the likelihood of gene drive chase for the X-shredder drive is most strongly
impacted by the movement speed. The occurrence of a chase in a X-poisoning gene
drive is rare, only occurring at low frequency under uniform release conditions (Figure
4b).

Gene drive persistence

In the case of a pest such as *Drosophila suzukii* the unpredictability of a chase
dynamic is undesirable. As the X-poisoning gene drive has a lower occurrence of gene
drive chase than the X-shredder under all release strategies and tested variables, we
further examined whether X-poisoning provided an advantage over X-shredding by
examining the persistence of gene drives within populations.

We assessed the number of simulated populations that had gene drives
persisting in their last generation. In the simulations of X-poisoning drives there were
none under any release strategy. Whereas the X-shredding simulations exhibited
substantial rates of persistence particularly in scenarios where individuals within the
populations had low to low-medium movement rates (Figure 4c). Due to the threshold
dynamics of the X-poisoning drive, even if it were to undergo a chase it is unlikely to
eradicate a population due to the migration of gene drive individuals alone. Unlike the
X-shredder, which spreads quickly into an expanding population, the X-poisoning drive
expands more slowly than the expanding wild-type population due to the reduction in

reproductive output from the postzygotic lethality of this drive. The reduction in
reproductive output acts as a fitness cost, resulting in the removal of the gene drive if
it is below the threshold required to drive. As a result, the X-poisoning drive either
eradicates the population or is removed from the population before the end of the
observed period.

Discussion
440
For an X-poisoning strategy to elicit an effect in a single generation the targeted
genes need to be haplo-insufficient. Such is the knowledge associated with
Drosophila melanogaster that it is known there are eleven loci on the X chromosome
of *D. melanogaster* that are haplolethal or haplosterile (Cook *et al.* 2012).
Haplolethal genes can only exist on the X chromosome in species where dosage
compensation between the homogametic and heterogametic sex is due to
upregulation - doubling the output of the X chromosome in males, rather than the
inactivation of the X as seen in humans.
449
Eight of the X-linked haplolethals of *D. melanogaster* encode ribosomal proteins,
which contribute to the translational machinery of cells, and an insufficient quantity of
them leads to lethality or to the Minute syndrome where flies are tiny and sterile
(Marygold *et al.* 2007). Previously, Fasulo *et al.* 2020 pioneered the 'X-poisoning'
strategy by targeting Ribosomal protein genes S6 and S5a. They found that when
one particular sgRNA targeting RpS6 was combined with cas9 driven by a sperm-
specific beta tubulin 85D promoter the sex ratio was biased so that greater than 92%
of progeny were males. Our targeting of RpL36, RpL35 and RpL19a did not produce
such extreme biasing. However, Fasulo *et al.* (2020) also showed heterogeneity in
the success of their sgRNA's with different guide RNA's targeting the same genes
skewing to various extents. Therefore, the failure to see skewing in our combination
of RpL36 and RpL35 sgRNA's or the skew to only 80% males in the RpL35 and
RpL19a combination, may well have more to do with the particularities of guide RNA
efficacy than being a property of the genes they target.
464
The gene that showed the greatest skewing in our X-poisoning experiments was the
only non-ribosomal protein haplolethal gene that has been molecularly characterised

on the X chromosome. The Wings-up-A (*wupA*) locus of Drosophila encodes

Troponin I, a protein that inhibits the interaction of myosin and actin in muscles. It is

also thought to have non-muscle functions that relate to the nuclear localization of at

least one of its nine isoforms (Casas-Tintó and Ferrús 2021, Flybase). The gene has

12 exons that are variously combined to produce 13 alternate transcriptional splice

forms. Two of the three sgRNAs used in our study target exons common to all splice

forms, and the third targets a PAM site that is found in the first two codons of

isoforms with the most common start methionine. The latter site was previously

targeted with CRISPR by Casas-Tintó and Ferrús (2021) and the resulting

haplolethality is consistent with our findings.

Curiously, we found that the skew in sex ratio was more extreme when the poisoned

X chromosome was passed to sons (via attached-X females) rather than to

daughters (figure 2d versus figures 2b and 2c). As the fathers in these crosses are

the same, this difference is not due to the generation of X-poisoned alleles. Instead,

heterozygosity in daughters may protect more than the exposed hemizygosity of

sons, which suggests that some of the CRISPR generated alleles may encode

recessive lethals. An alternate explanation is that there is something else about

maleness that makes the CRISPR *wupA* alleles more potent in males than females.

A surprising finding of Fasulo *et al.* (2020) was that some alleles among the

survivors contained frameshift mutations. This led the authors to suggest that the

targeted ribosomal protein genes may not be haplolethal genes after all and that

previously studied alleles may have harboured dominant negative mutations instead.

In our analyses, we did not find such frameshifting alleles among individuals carrying

X-chromosomes that had passed through X-poisoning males. Instead, except for a

single in-frame alteration, the loci showed no signs of being cut at all. While our

analyses did not survey as many individuals as Fasulo *et al.'s* study, it perhaps

avoided issues of sequencing errors, copy number variation or chimerism, which

provide alternate explanations for Fasulo *et al.*'s (2020) viable frameshifts in pooled

ribosomal-protein gene sequences.

WupA has very low expression in the early stages of embryos (first 8 hours) which

contrasts to many ribosomal proteins because it is probably not required for the

viability of all cells. Perhaps this feature explains why the number of sons is not

reduced in the *wupA* cross, whereas they are reduced in the ribosomal-protein

crosses. Such an effect may also extend back to the fitness of Y bearing sperm

produced by male carriers of the *wupA*-poison. They may be as viable as wildtype

sperm in a way that is not true for Rp-poisoning. This makes it an attractive target for

future X-poisoning strategies.

WupA is a highly conserved gene that is found on the X chromosome on numerous

dipteran species, including some important pests. *Drosophila suzukii,* an invasive

pest of soft-skinned fruits, appears to share all 12 exons with *D. melanogaster* and

all the predicted proteins have greater than 98% amino acid identity with *D.*

melanogaster homologs. In fact, two of the three sgRNA's used here in *D.*

melanogaster match *D. suzukii* targets perfectly, and the third has only one

nucleotide different. *WupA* of Anopheles mosquito's has a slightly different gene

structure (eg. Exon 1 and Exon 2 are fused eg. XM_053811545) however it is also

found on the X chromosome in these species suggesting it could be a target of X-

poisoning strategies in the many pest species of this genus.

Our modelling compared X-poisoning to X-shredding drives and suggested that the

latter is often more effective at eradication in an expanding population. The X-

shredder gene drive can expand into the expanding population of insects,

eradicating them as it goes. It does this more effectively than X-poisoning because

its bearers were modelled as having twice the reproductive output when compared to

the X-poisoning (this may be over-conservative as multiple mating's may introduce a

frequency dependent effect that would be less harsh on the X-poisoning). As the

expansion of the wild-type population is constrained by the ability to find a mate and

produce offspring at the invasion front, the X-shredder drive can expand into the

wild-type population at a slightly faster rate than the wild-type population can expand

into the uninhabited space. This results in the expansion of the drive following the

expansion of the population like a wave (e.g. Video S1).

The simulations also emphasize how important the extent of movement of individuals

is to gene drive success. All interactions between individuals are impacted by

movement: competition is based on the number of individuals within a generation's

movement, mating occurs only between individuals within interacting distance. As
movement increases, the interaction distance increases, and gene flow within the
population increases. Individuals are also able to colonise new uninhabited areas
more quickly resulting in the space reaching carrying capacity more quickly. For X-
shredding the greater the movement the more successful the gene drive. However
for X-shredding the effect the effect is quite different; with eradication only successful
when the movement of individuals is low. As the population size increases faster
with greater movement of individuals, yet the number of released individuals is
constrained (it is set to a certain size in our simulations, or to what is logistically
possible in a real scenario), the proportion of released gene drive individuals within
the population decreases and so in the high movement scenarios the released
individuals never cross the necessary threshold required for population eradication.
547
While the likelihood of eradication would be a primary consideration of anyone
deploying a suppression gene drive, many of our simulations showed that the X-
shredder gene drive persisted until the end of the observed period. The persistence
of the gene drive was particularly true when gene drive chasing occurred. Due to the
drive suppressing the population, the X-shredder drive leaves uninhabited space
behind it. If wild-type individuals move behind the expanding wave of the X-shredder
individuals, they are then able to expand in the other direction and thereby escape
the gene drive. This often results in a gene drive chase back in the other direction.
Persistence of gene drives is an undesirable outcome for gene drive population
control as it increases the length of time the drive is present within the population
and that increases the likelihood that the drive will escape beyond the target area,
potentially impacting the species in its native range. X-poisoning on the other hand
has more of a threshold dynamic which limits the chasing dynamic. Thus, there are
circumstances where an X-poisoning may be a better strategy than X-shredding.
562
Our models also illustrate several other features that would need to be considered
before gene drive release. Features such as seasonal declines in population size
had a noticeable effect. Perhaps most significantly our models demonstrated the
importance of gene drive release strategies. They found it was much better to
release the gene drive throughout the invasive species range rather than limiting the
gene drive release to the incursion origin.

Data Availability Statement

570 All data is here or in the supplementary material.

Acknowledgements

572 We'd like to thank Owain Edwards and Max Scott for conversations that helped develop this
573 research and Jackson Champer for suggestions about construct design.

Funding

575 The research was funded by an Australian Research Council Discovery Project Grant
576 (DP190102512).

Conflict of Interests

578 We have no conflicts of interest to declare.

References

Akbari, Omar S., Chun-Hong Chen, John M. Marshall, Haixia Huang, Igor Antoshechkin, and
Bruce A. Hay. 2014. "Novel Synthetic *Medea* Selfish Genetic Elements Drive
Population Replacement in *Drosophila* ; a Theoretical Exploration of *Medea* -
Dependent Population Suppression." *ACS Synthetic Biology* 3 (12): 915–28.
https://doi.org/10.1021/sb300079h.
Alphey, Luke S., Andrea Crisanti, Filippo (Fil) Randazzo, and Omar S. Akbari. 2020.
"Standardizing the Definition of Gene Drive." *Proceedings of the National Academy*
of Sciences 117 (49): 30864–67. https://doi.org/10.1073/pnas.2020417117.
Buchman, Anna, John M. Marshall, Dennis Ostrovski, Ting Yang, and Omar S. Akbari. 2018.
"Synthetically Engineered *Medea* Gene Drive System in the Worldwide Crop Pest
Drosophila suzukii." *Proceedings of the National Academy of Sciences* 115 (18):
4725–30. https://doi.org/10.1073/pnas.1713139115.
Burt, Austin. 2003. "Site-Specific Selfish Genes as Tools for the Control and Genetic
Engineering of Natural Populations." *Proceedings of the Royal Society B: Biological*
Sciences 270 (1518): 921–28. https://doi.org/10.1098/rspb.2002.2319.
Bushland, R. C., A. W. Lindquist, and E. F. Knipling. 1955. "Eradication of Screw-Worms
through Release of Sterilized Males." *Science* 122 (3163): 287–88.
https://doi.org/10.1126/science.122.3163.287.
Casas-Tintó, Sergio, and Alberto Ferrús. 2021. "The Haplolethality Paradox of the WupA
Gene in Drosophila." Edited by R. Scott Hawley. *PLOS Genetics* 17 (3): e1009108.
https://doi.org/10.1371/journal.pgen.1009108.
Champer, Jackson, Anna Buchman, and Omar S. Akbari. 2016. "Cheating Evolution:
Engineering Gene Drives to Manipulate the Fate of Wild Populations." *Nature*
Reviews Genetics 17 (3): 146–59. https://doi.org/10.1038/nrg.2015.34.
Champer, Jackson, Isabel K Kim, Samuel E Champer, Andrew G Clark, and Philipp W Messer.
2021. "Suppression Gene Drive in Continuous Space Can Result in Unstable
Persistence of Both Drive and Wild-Type Alleles." *Molecular Ecology* 30 (December
2020): 1086–1101. https://doi.org/10.1111/mec.15788.
Champer, Jackson, Jingxian Liu, Suh Yeon Oh, Riona Reeves, Anisha Luthra, Nathan Oakes,
Andrew G. Clark, and Philipp W. Messer. 2018. "Reducing Resistance Allele

Formation in CRISPR Gene Drive." *Proceedings of the National Academy of Sciences*
115 (21): 5522–27. https://doi.org/10.1073/pnas.1720354115.
Champer, Jackson, Riona Reeves, Suh Yeon Oh, Chen Liu, Jingxian Liu, Andrew G. Clark, and
Philipp W. Messer. 2017. "Novel CRISPR/Cas9 Gene Drive Constructs Reveal Insights
into Mechanisms of Resistance Allele Formation and Drive Efficiency in Genetically
Diverse Populations." Edited by Harmit S. Malik. *PLOS Genetics* 13 (7): e1006796.
https://doi.org/10.1371/journal.pgen.1006796.
Champer, Jackson, Zhaoxin Wen, Anisha Luthra, Riona Reeves, Joan Chung, Chen Liu, Yoo
Lim Lee, *et al.* 2019. "CRISPR Gene Drive Efficiency and Resistance Rate Is Highly
Heritable with No Common Genetic Loci of Large Effect." *Genetics* 212 (1): 333–41.
https://doi.org/10.1534/genetics.119.302037.
Chen, Chun-Hong, Haixia Huang, Catherine M. Ward, Jessica T. Su, Lorian V. Schaeffer, Ming
Guo, and Bruce A. Hay. 2007. "A Synthetic Maternal-Effect Selfish Genetic Element
Drives Population Replacement in Drosophila." *Science* 316 (5824): 597–600.
https://doi.org/10.1126/science.1138595.
Cook, R Kimberley, Stacey J Christensen, Jennifer A Deal, Rachel A Coburn, Megan E Deal, Jill
M Gresens, Thomas C Kaufman, and Kevin R Cook. 2012. "The Generation of
Chromosomal Deletions to Provide Extensive Coverage and Subdivision of the
Drosophila melanogaster Genome." *Genome Biology* 13 (3).
https://doi.org/10.1186/gb-2012-13-3-r21.
Curtis, C. F. 1968. "A Possible Genetic Method for the Control of Insect Pests, with Special
Reference to Tsetse Flies (Glossina Spp.)." *Bulletin of Entomological Research* 57 (4):
509–23.
Eckhoff, Philip A., Edward A. Wenger, H. Charles J. Godfray, and Austin Burt. 2017. "Impact
of Mosquito Gene Drive on Malaria Elimination in a Computational Model with
Explicit Spatial and Temporal Dynamics." *Proceedings of the National Academy of*
Sciences 114 (2). https://doi.org/10.1073/pnas.1611064114.
Fasulo, Barbara, Angela Meccariello, Maya Morgan, Carl Borufka, Philippos Aris Papathanos,
and Nikolai Windbichler. 2020. "A Fly Model Establishes Distinct Mechanisms for
Synthetic CRISPR/Cas9 Sex Distorters." *PLoS Genetics* 16 (3): 1–22.
https://doi.org/10.1371/journal.pgen.1008647.
Galizi, Roberto, Lindsey A. Doyle, Miriam Menichelli, Federica Bernardini, Anne Deredec,
Austin Burt, Barry L. Stoddard, Nikolai Windbichler, and Andrea Crisanti. 2014. "A
Synthetic Sex Ratio Distortion System for the Control of the Human Malaria
Mosquito." *Nature Communications* 5 (1): 3977.
https://doi.org/10.1038/ncomms4977.
Gamez, Stephanie, Duverney Chaverra-Rodriguez, Anna Buchman, Nikolay P. Kandul, Stelia
C. Mendez-Sanchez, Jared B. Bennett, Héctor M. Sánchez C., *et al.* 2021. "Exploiting a
Y Chromosome-Linked Cas9 for Sex Selection and Gene Drive." *Nature*
Communications 12 (1): 7202. https://doi.org/10.1038/s41467-021-27333-1.
Gratz, Scott J., Fiona P. Ukken, C. Dustin Rubinstein, Gene Thiede, Laura K. Donohue,
Alexander M. Cummings, and Kate M. O'Connor-Giles. 2014. "Highly Specific and
Efficient CRISPR/Cas9-Catalyzed Homology-Directed Repair in Drosophila." *Genetics*
196 (4): 961–71. https://doi.org/10.1534/genetics.113.160713.
Haller, Benjamin C, and Philipp W Messer. 2019. "SLiM 3: Forward Genetic Simulations
Beyond the Wright–Fisher Model." Edited by Ryan Hernandez. *Molecular Biology*
and Evolution 36 (3): 632–37. https://doi.org/10.1093/molbev/msy228.

Hammond, Andrew M., Kyros Kyrou, Marco Bruttini, Ace North, Roberto Galizi, Xenia
Karlsson, Nace Kranjc, *et al.* 2017. "The Creation and Selection of Mutations
Resistant to a Gene Drive over Multiple Generations in the Malaria Mosquito."
Edited by Nick H. Barton. *PLOS Genetics* 13 (10): e1007039.
https://doi.org/10.1371/journal.pgen.1007039.
Hay, Bruce A., Georg Oberhofer, and Ming Guo. 2021. "Engineering the Composition and
Fate of Wild Populations with Gene Drive." *Annual Review of Entomology* 66 (1):
407–34. https://doi.org/10.1146/annurev-ento-020117-043154.
Hendrichs, J., G. Franz, and P. Rendon. 1995. "Increased Effectiveness and Applicability of
the Sterile Insect Technique through Male-Only Releases for Control of
Mediterranean Fruit Flies during Fruiting Seasons." *Journal of Applied Entomology*
119 (1–5): 371–77. https://doi.org/10.1111/j.1439-0418.1995.tb01303.x.
Knapp, Ladina, Dominique Mazzi, and Robert Finger. 2021. "The Economic Impact of
Drosophila Suzukii : Perceived Costs and Revenue Losses of Swiss Cherry, Plum and
Grape Growers." *Pest Management Science* 77 (2): 978–1000.
https://doi.org/10.1002/ps.6110.
Knipling, E.F. 1955. "Possibilities of Insect Control or Eradication through the Use of Sexually
Sterile Males." *Journal of Economic Entomology,* 48 (4): 459–62.
Krüger, Alexandra P., Daniele C. H. Schlesener, Liliane N. Martins, Jutiane Wollmann,
Maríndia Deprá, and Flávio R. M. Garcia. 2019. "Radiation Effects on *Drosophila*
Suzukii (Diptera: Drosophilidae) Reproductive Behaviour." *Journal of Applied*
Entomology 143 (1–2): 88–94. https://doi.org/10.1111/jen.12563.
Kyrou, Kyros, Andrew M. Hammond, Roberto Galizi, Nace Kranjc, Austin Burt, Andrea K.
Beaghton, Tony Nolan, and Andrea Crisanti. 2018. "A CRISPR–Cas9 Gene Drive
Targeting Doublesex Causes Complete Population Suppression in Caged *Anopheles*
gambiae Mosquitoes." *Nature Biotechnology* 36 (11): 1062–66.
https://doi.org/10.1038/nbt.4245.
Labun, Kornel, Tessa G Montague, Maximilian Krause, Yamila N Torres Cleuren, Håkon
Tjeldnes, and Eivind Valen. 2019. "CHOPCHOP v3: Expanding the CRISPR Web
Toolbox beyond Genome Editing." *Nucleic Acids Research* 47 (W1): W171–74.
https://doi.org/10.1093/nar/gkz365.
Lanouette, Geneviève, Jacques Brodeur, François Fournier, Veronique Martel, and
Annabelle Firlej. 2020. "Effect of Irradiation on the Mating Capacity and
Competitiveness of *Drosophila suzukii* (Diptera: Drosophilidae) for the Development
of the Sterile Insect Technique." *The Canadian Entomologist* 152 (4): 563–74.
https://doi.org/10.4039/tce.2020.38.
Larkin, Aoife, Steven J Marygold, Giulia Antonazzo, Helen Attrill, Gilberto dos Santos, Phani V
Garapati, Joshua L Goodman, *et al.* 2021. "FlyBase: Updates to the *Drosophila*
Melanogaster Knowledge Base." *Nucleic Acids Research* 49 (D1): D899–907.
https://doi.org/10.1093/nar/gkaa1026.
Larner, William, Tom Price, Luke Holman, and Nina Wedell. 2019. "An X-Linked Meiotic Drive
Allele Has Strong, Recessive Fitness Costs in Female *Drosophila pseudoobscura*."
Proceedings of the Royal Society B: Biological Sciences 286 (1916): 20192038.
https://doi.org/10.1098/rspb.2019.2038.
Maino, James L., Rafael Schouten, and Paul Umina. 2021. "Predicting the Global Invasion of
Drosophila suzukii to Improve Australian Biosecurity Preparedness." Edited by Bret

Elderd. *Journal of Applied Ecology* 58 (4): 789–800. https://doi.org/10.1111/1365-
2664.13812.
Marygold, Steven J., John Roote, Gunter Reuter, Andrew Lambertsson, Michael Ashburner,
Gillian H. Millburn, Paul M. Harrison, *et al.* 2007. "The Ribosomal Protein Genes and
Minute Loci of *Drosophila melanogaster*." *Genome Biology* 8 (10): R216.
https://doi.org/10.1186/gb-2007-8-10-r216.
Mazzi, Dominique, Esther Bravin, Manuela Meraner, Robert Finger, and Stefan Kuske. 2017.
"Economic Impact of the Introduction and Establishment of *Drosophila suzukii* on
Sweet Cherry Production in Switzerland." *Insects* 8 (1): 18.
https://doi.org/10.3390/insects8010018.
Meccariello, Angela, Flavia Krsticevic, Rita Colonna, Giuseppe Del Corsano, Barbara Fasulo,
Philippos Aris Papathanos, and Nikolai Windbichler. 2021. "Engineered Sex Ratio
Distortion by X-Shredding in the Global Agricultural Pest *Ceratitis capitata*." *BMC*
Biology 19 (1): 78. https://doi.org/10.1186/s12915-021-01010-7.
Morrow, Edward H., Andrew D. Stewart, and William R. Rice. 2005. "Patterns of Sperm
Precedence are not Affected by Female Mating History in *Drosophila melanogaster*."
Evolution 59 (12): 2608. https://doi.org/10.1554/05-238.1.
North, Ace, Austin Burt, and H. Charles J. Godfray. 2013. "Modelling the Spatial Spread of a
Homing Endonuclease Gene in a Mosquito Population." Edited by Yvonne Buckley.
Journal of Applied Ecology 50 (5): 1216–25. https://doi.org/10.1111/1365-
2664.12133.
Oberhofer, Georg, Tobin Ivy, and Bruce A. Hay. 2019. "Cleave and Rescue, a Novel Selfish
Genetic Element and General Strategy for Gene Drive." *Proceedings of the National*
Academy of Sciences 116 (13): 6250–59. https://doi.org/10.1073/pnas.1816928116.
Ørsted, Iben Vejrum, and Michael Ørsted. 2019. "Species Distribution Models of the Spotted
Wing *Drosophila* (*Drosophila suzukii* , Diptera: Drosophilidae) in Its Native and
Invasive Range Reveal an Ecological Niche Shift." Edited by Cristina Garcia. *Journal of*
Applied Ecology 56 (2): 423–35. https://doi.org/10.1111/1365-2664.13285.
Paril, Jeff F., and Ben L. Phillips. 2022. "Slow and Steady Wins the Race: Spatial and
Stochastic Processes and the Failure of Suppression Gene Drives." *Molecular Ecology*
31 (17): 4451–64. https://doi.org/10.1111/mec.16598.
Port, Fillip, and Simon L Bullock. 2016. "Augmenting CRISPR Applications in Drosophila with
tRNA-Flanked SgRNAs." *Nature Methods* 13 (10): 852–54.
https://doi.org/10.1038/nmeth.3972.
Prado, A., I. Canal, and A. Ferrús. 1999. "The Haplolethal Region at the 16F Gene Cluster of
Drosophila melanogaster: Structure and Function." *Genetics* 151 (1): 163–75.
https://doi.org/10.1093/genetics/151.1.163.
Prowse, Thomas A. A., Phillip Cassey, Joshua V. Ross, Chandran Pfitzner, Talia A. Wittmann,
and Paul Thomas. 2017. "Dodging Silver Bullets: Good CRISPR Gene-Drive Design Is
Critical for Eradicating Exotic Vertebrates." *Proceedings of the Royal Society B:*
Biological Sciences 284 (1860): 20170799. https://doi.org/10.1098/rspb.2017.0799.
R Core team. 2023. "R Core Team (2022). R: A Language and Environment for Statistical
Computing. R Foundation for Statistical Computing, Vienna, Austria." https://www.R-
project.org/.
Rose, Graham, Elzbieta Krzywinska, Jan Kim, Loic Revuelta, Luca Ferretti, and Jaroslaw
Krzywinski. 2016. "Dosage Compensation in the African Malaria Mosquito *Anopheles*

gambiae." *Genome Biology and Evolution*, January, evw004.
https://doi.org/10.1093/gbe/evw004.
Sánchez C., Héctor M., Jared B. Bennett, Sean L. Wu, Gordana Rašić, Omar S. Akbari, and
John M. Marshall. 2020. "Modeling Confinement and Reversibility of Threshold-
Dependent Gene Drive Systems in Spatially-Explicit Aedes Aegypti Populations." *BMC*
Biology 18 (1): 50. https://doi.org/10.1186/s12915-020-0759-9.
Schnakenberg, Sandra L., Mark L. Siegal, and Margaret C. Bloch Qazi. 2012. "Oh, the Places
They'll Go: Female Sperm Storage and Sperm Precedence in *Drosophila*
melanogaster." *Spermatogenesis* 2 (3): 224–35.
https://doi.org/10.4161/spmg.21655.
Scott, Maxwell J., Carolina Concha, John B. Welch, Pamela L. Phillips, and Steven R. Skoda.
2017. "Review of Research Advances in the Screwworm Eradication Program over
the Past 25 Years." *Entomologia Experimentalis et Applicata* 164 (3): 226–36.
https://doi.org/10.1111/eea.12607.
Serebrovsky, A.S. 1969. "On the Possibility of a New Method for the Control of Insect Pests."
In *Proceedings of a Panel on Application of the Sterile-Male Technique for the*
Eradication or Control of Harmful Species of Insects, Organised by the Joint FAO/IAEA
Division of Atomic Energy in Food and Agriculture, pp.123-237. Vienna, 27-31 May
1968.
Stockton, Dara, Anna Wallingford, and Gregory Loeb. 2018. "Phenotypic Plasticity Promotes
Overwintering Survival in A Globally Invasive Crop Pest, *Drosophila suzukii.*" *Insects* 9
(3): 105. https://doi.org/10.3390/insects9030105.
Waltz, Emily. 2021. "First Genetically Modified Mosquitoes Released in the United States."
Nature 593 (7858): 175–76. https://doi.org/10.1038/d41586-021-01186-6.
Webster, Sophia H., Michael R. Vella, and Maxwell J. Scott. 2020. "Development and Testing
of a Novel Killer–Rescue Self-Limiting Gene Drive System in *Drosophila*
melanogaster." *Proceedings of the Royal Society B: Biological Sciences* 287 (1925):
20192994. https://doi.org/10.1098/rspb.2019.2994.
Whitten, M.J. 1985. "The Conceptual Basis for Genetic Control." In *Comprehensive Insect*
Physiology Biochemistry and Pharmacology, 12:465–528. Insect Control. Oxford,
New York, Toronto, Sydney, Paris, Frankfurt: Pergamon Press.
Windbichler, N., P. A. Papathanos, F. Catteruccia, H. Ranson, A. Burt, and A. Crisanti. 2007.
"Homing Endonuclease Mediated Gene Targeting in *Anopheles gambiae* Cells and
Embryos." *Nucleic Acids Research* 35 (17): 5922–33.
https://doi.org/10.1093/nar/gkm632.
Windbichler, Nikolai, Miriam Menichelli, Philippos Aris Papathanos, Summer B. Thyme, Hui
Li, Umut Y. Ulge, Blake T. Hovde, *et al.* 2011. "A Synthetic Homing Endonuclease-
Based Gene Drive System in the Human Malaria Mosquito." *Nature* 473 (7346): 212–
15. https://doi.org/10.1038/nature09937.
Yang, Emily, Matthew Metzloff, Anna M Langmüller, Xuejiao Xu, Andrew G Clark, Philipp W
Messer, and Jackson Champer. 2022. "A Homing Suppression Gene Drive with
Multiplexed gRNAs Maintains High Drive Conversion Efficiency and Avoids Functional
Resistance Alleles." Edited by M Ramaswami. *G3 Genes|Genomes|Genetics* 12 (6):
jkac081. https://doi.org/10.1093/g3journal/jkac081.
Yao, Franck Adama, Abdoul-Azize Millogo, Patric Stephane Epopa, Ace North, Florian Noulin,
Koulmaga Dao, Mouhamed Drabo, *et al.* 2022. "Mark-Release-Recapture Experiment
in Burkina Faso Demonstrates Reduced Fitness and Dispersal of Genetically-Modified

Sterile Malaria Mosquitoes." *Nature Communications* 13 (1): 796.
https://doi.org/10.1038/s41467-022-28419-0.
Yeh, D Adeline, Francis A Drummond, Miguel I Gómez, and Xiaoli Fan. 2020. "The Economic
Impacts and Management of Spotted Wing Drosophila (*Drosophila suzukii*): The Case
of Wild Blueberries in Maine." Edited by Cesar Rodriguez-Saona. *Journal of Economic
Entomology* 113 (3): 1262–69. https://doi.org/10.1093/jee/toz360.
803
804

805 ## Figure Legends
806

Figure 1: Targeting Haplolethal genes on the X chromosome. (a) Premeiotic

cells expressing CRISPR/Cas9 (from the autosomes) that effectively target

haplolethal loci on the X-chromosome (orange to red) will result in male only

offspring. If the machinery were put on the Y-chromosome then inheritance of

the elements will be limited to the male germ line and a gene drive would be

created. (b) Eleven haplolethal/haplosteriles were reported by Cook *et al.*

(2012), two (RpS6 and RpS5a) were targeted by Fasulo *et al.* 2020 and the

location of the four loci targeted in this manuscript, and the number of sgRNA

used (rounded blue rectangles) are shown. (c) The WupA gene encodes for

thirteen transcript isoforms. SgRNA1-3 target three conserved regions of the

gene's CDS.

Figure 2: Sex biases with X-poisoning. (a) A skew against daughters (red) is

expected when sgRNA's target haplolethal loci on the X chromosome during

gametogenesis. The number of sons is expected to remain unchanged as they

would arise from sperm carrying the Y chromosome. (b) The observed amount of

sex biasing skew ranged from nothing (RpL36+RpL35), to modest (RpL19a+RpL35),

to promising (WupA). Note that the total number of offspring was also variously

affected. (c) By excluding multiple mating the total number of offspring in the WupA

cross was reduced and there was a strong skew towards males (d) CRISPR/cas9

fathers crossed to Attached X mothers, reverse the inheritance of the sex

chromosomes so that sons inherited the CRISPR modified X chromosome, and

daughters inherited the Y chromosome. (* $P < 0.05$, ** $P < 0.01$, ** $P < 0.001$ **** $P <$

0.0001, unpaired two-tail t-test with a Bonferroni correction for multiple

comparisons).

Figure 3: Simulations of 'X-poisoning' and 'X-shredding' in expanding

populations. The eradication probability of the invasive population is shown in a

black to yellow scale with each coloured element representing 100 simulations. Each

square represents simulations that varied by the number of gene drive individuals

that were released and when they were released relative to the incursion of the

invading species. Four values of movement of individuals are shown (low movement,

medium-low, medium-high, high). Simulations represented in (a) had the gene drive

individuals released at a single time at the incursion point of the invasive species.

Simulations represented in (b) involved multiple smaller releases at the incursion

point. In the simulations of (c) and (d) the gene drive individuals were released

across the occupancy range of the invasive species. Each data point represents the

average outcome of 100 independent simulation runs.

Figure 4: Simulations with realized biasing efficiencies. In contrast to the

analysis of Figure 3 which assumed the poisoning and shredding happened with

absolute efficiency these simulations used the data generated here and in the paper

of Meccariello *et al.* 2021 to inform sex ratio biasing rates. All the simulations shown

here featured a single gene drive release across the present range of the invasive

species. In the simulations of (a) the probability of population eradication is shown

and seasonality can be seen as horizontal striations. In (b) the colour represents the

likelihood that a gene drive 'chase' occur and in (c) the colour represents the

presence of the gene drives at the end of the modelling. Each data point represents

the average outcome of 100 independent simulation runs.

(a)

Cas9
SgRNA
Target gene
Null allele

SPERM:

AFTER FERTILIZATION:

(b)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20

RpL17
RpS6
Fs(1)10A   Hdl   RpL37a
RpS5a
RpS10B

RpL36   RpL35   RpS19a   WupA

= sgRNA
A = RpL36 + RpL35 construct
B = RpS19a + RpL35 construct
C = WupA construct

A

B

C

(c)

WupA gene (11,355 bp)

* SgRNA Targets

Isoforms
A
B
C
D
E
F
G
H
I
J
K
L
M

M A D D
ATGGCTGATGAT
SgRNA1 PAM

M E E A
ATGGAGGAAGCC
SgRNA2 PAM

C G S P
TGCGGCAGTCCC
SgRNA3 PAM

Figure 2

Figure 3

Figure 4