1 TITLE:
2

The haplolethal gene *wupA* of Drosophila exhibits potential as a target

for an X-poisoning gene drive.

Clancy D. Lawler[#], Ana Karla Parra Nuñez[#], Natalia Hernandes, Soumitra Bhide,

Isabelle Lohrey, Simon Baxter and Charles Robin*

School of BioSciences, The University of Melbourne, Victoria, Australia

[#] these authors contributed equally to this work

*Corresponding author:

Charles Robin

Building 184, School of BioSciences, Parkville Campus,

The University of Melbourne, Victoria 3010 Australia

crobin@unimelb.edu.au

Running head: A target for a haplolethal gene drive

Abstract

A synthetic gene drive that targets haplolethal genes on the X-chromosome can skew the sex ratio towards males. Like an 'X-shredder' it does not involve 'homing' and that has advantages including the reduction of gene drive resistance allele formation. We examine this 'X-poisoning' strategy by targeting four of the 11 known X-linked haplolethal/haplosterile genes of *Drosophila melanogaster* with CRISPR/Cas9. We find that targeting the *wupA* gene during spermatogenesis skews the sex ratio so fewer than 14% of progeny are daughters. That is unless we cross the mutagenic males to X^XY female flies that bear attached-X chromosomes, which reverses the inheritance of the poisoned X chromosome so that sons inherit it from their father; in which case only 2% of the progeny are sons. These sex ratio biases suggests that most of the CRISPR/Cas9 mutants we induced in the *wupA* gene are haplolethal but some are recessive lethal. The males generating *wupA* mutants do not suffer from reduced fertility rather the haplolethal mutants arrest development in the late stages of embryogenesis well after fertilized eggs have been laid. This provides a distinct advantage over genetic manipulation strategies involving sterility which can be countered by the remating of females. We also find that *wupA* mutants that destroy the nuclear localization signal of shorter isoforms are not haplolethal as long as the open reading frame remains intact. Like *D. melanogaster wupA* orthologs of *D. suzukii* and *Anopheles* mosquitos are found on X chromosomes making *wupA* a viable X-poisoning target in multiple species.

Introduction

For many decades it has been envisioned that insect populations that vector
diseases or that are pests of crops could be controlled using genetic methods
(Serebrovsky 1969; Knipling 1955; Curtis 1968; Whitten 1985; Burt 2003). Indeed,
Sterile Insect Techniques have successfully controlled fly species (Bushland,
Lindquist, and Knipling 1955; Hendrichs, Franz, and Rendon 1995; Scott *et al.* 2017)
and field trials relying on self-limiting transgenes are currently being deployed and
assessed (Yao *et al.* 2022; Waltz 2021). Recently, the prospect of synthetic gene
drives, a distinct class of population manipulation, where transgenes can be
engineered to spread through otherwise natural populations of insect pests has
received much attention (Alphey *et al.* 2020; Hay, Oberhofer, and Guo 2021;
Champer, Buchman, and Akbari 2016). The effectiveness of various gene drive
designs, many of which use CRISPR/Cas9 machinery, have been assessed in
laboratory populations (Chen *et al.* 2007; N. Windbichler *et al.* 2007; Akbari *et al.*
2014; Kyrou *et al.* 2018; Oberhofer, Ivy, and Hay 2019; Webster, Vella, and Scott
2020; Yang *et al.* 2022). Many of these require 'homing' where double-stranded DNA
breaks must be repaired by homologous recombination (HR) so that the transgenic
selfish elements are copied and thereby spread. A drawback of such homing designs
has been that some double stranded breaks can be repaired using the non-
homologous end-joining (NHEJ) pathway. In these instances not only does homing
fail but mutations that are resistant to subsequent CRISPR/Cas9 cleavages can be
introduced into the population (Nikolai Windbichler *et al.* 2011; Gratz *et al.* 2014;
Champer *et al.* 2017; 2018; Hammond *et al.* 2017).

One genetic control design that does not require homing is referred to as the X-
shredder (Galizi *et al.* 2014). It targets highly repeated sequences on the X
chromosome during spermatogenesis such that X-bearing sperm are inviable, and
the drive skews the sex ratio towards males. If the X-shredding transgenes could be
placed on the Y chromosome, then an inheritance can be established such that
fathers only have sons. Originally, the X-shredder was attempted in lab populations
of *Anopheles gambiae* mosquito using the endonuclease *PpoI*, but subsequent

implementations in *A. gambiae* mosquitos, *Drosophila melanogaster and Ceratitis*
capitata fruit flies have deployed CRISPR/Cas9 machinery (Galizi et al. 2014; Fasulo
et al. 2020; Meccariello et al. 2021; Haber et al. 2023).

A related approach, dubbed 'X-poisoning', also targets loci on the X chromosome;
not so the chromosome is physically destroyed but so that mutations are introduced
such that progeny receiving the X chromosome will not be viable (Burt and Deredec
2018; Fasulo et al. 2020; Haber et al. 2023). This strategy uses CRISPR/Cas9 to
target X-linked haplolethal genes. Although haplolethal genes are typically thought of
as genes where both alleles of a diploid organism need to be functional, X-linked
haplolethal genes can occur in many dipterans because their dosage compensation
mechanism elevates the X chromosome-encoded output in males (that is in contrast
to eutherian mammals where there is silencing of X loci in females and therefore
haplolethals on the X would be incompatible with maleness) (Rose *et al.* 2016; Cook
et al. 2012). There may be a few potential advantages of X-poisoning over X-
shredding. Firstly, X-poisoning does not rely on natural repeats occurring exclusively
on the X chromosome and theoretically this may increase its applicability to some
species. Secondly, CRISPR cutting in one or a few sites on the X chromosome might
occur more readily than cutting the X chromosome at an overwhelming number of
places, as is required for X-shredding. Thirdly, X-shredding has been shown to be
ameliorated by the NHEJ pathway thus potentially limiting its utility (Fasulo *et al.*
2020). Fourthly, it may be that post-zygotic effects of X-poisoning may change the
spread dynamics so it has a higher threshold than X-shredding and that may lead to
more controllable population manipulation.

Here we describe the targeting of haplolethal genes on the X chromosome of
Drosophila melanogaster with CRISPR/Cas9. The loci differ from the two targeted in
the X-poisoning study of Fasulo *et al.* (2020) as we examine three other ribosomal
protein encoding genes and the haplolethal gene *Wings up A (wupA).* In contrast to
the ribosomal protein genes which might be required for the viability of most cell
types including those in spermatogenesis, *wupA*, encodes a troponin that is
expressed in muscles from mid embryogenesis. We therefore sought experimental
affirmation that these genes are indeed haplolethal genes and whether they could be
valuable targets for X-poisoning gene drives.

Materials and Methods

gRNA design and cloning

A combination of ChopChop v3 (Labun *et al.* 2019) and CRISPR Optimal Target Finder (Gratz *et al.* 2014) was used to identify the gRNA target sites with the highest efficiency scores with a low likelihood of off-target activity. All gRNA sequences were checked against the DGRP reference panel using FlyBase's JBrowse (Larkin *et al.* 2021) to avoid standing variation at the target site that may give natural resistance.

The design protocol of Port and Bullock (2016) was used to clone gRNA sequences into the pCFD5_w plasmid (Addgene #112645). Three separate plasmids were generated that targeted haplolethal genes on the X chromosome; pRpS19a_RpL35, pRpL36_RpL35, and p*wupA*. pRpS19a_RpL35 encoded four gRNAs, two targeting *RpS19a* (*RpS19*_RNA1: GAAGGATATTGACCAGCACG, *RpS19*_RNA2: AACCGACTCCAGCGGGACTG) and two targeting *RpL35* (*RpL35*_gRNA1: GTGCTCCGAGCTGAGGATCA, *RpL35*_gRNA2: CACAATGTAGACGCGAGCGA). pRpL36_RpL35 encodes four gRNA, two targeting *RpL36* (*RpL36*_gRNA1: GCTGGCTATTGGCCTGAACA, *RpL36*_gRNA2: CATGCGCGACTTGGTCCGCG) and two targeting *RpL35* (as aforementioned). Each of the three sgRNA's used to target *wupA* (*wupA*_gRNA1: ACCAAAAACACAAATCAAAA, *wupA*_gRNA2: TGAGGTGCGCAAGCGCATGG, *wupA*_gRNA3: CGCATCATCGAAGAACGTTG) would affect all 13 alternate transcripts annotated in Flybase. The PAM site for *wupA*_sgRNA1 corresponds to the most common start codon and is in the 5'UTR of transcripts using the alternate start codon, the PAM site for *wupA*_sgRNA2 corresponds to the alternate *wupA* start methionine which is a codon used by in all transcripts, and that for *wupA*_sgRNA3 is in a coding exon common to all transcripts (Figure 1). All primers used and sgRNA sequences are tabulated in Table S1.

Microinjections

250-500 ng/uL of the pRpS19a+RpL35 and p*wupA* plasmids were microinjected into the 09w fly line while the pRpL36+RpL35 was injected into the 10w fly line. These two fly lines were obtained from Trent Perry, Bio21 Unimelb who had generated them using the Bloomington stocks #24749 and #25709 (y[1] v[1] P{y[+t7.7]=nos-phiC31\int.NLS}X; P{y[+t7.7]=CaryP} attP40) and #25710 (P{y[+t7.7]=nos-

145   phiC31\int.NLS}X, y[1] sc[1] v[1] sev[21]; P{y[+t7.7]=CaryP}attP2) to generate second

146   chromosome landing site flies (09W) and third chromosome landing site flies (10W)

147   in a *white* null background. Insertion events were validated by sequencing and single

148   integration lines were maintained.

Assessing sex ratio biasing capabilities of sgRNA constructs

To assess the sex ratio biasing rates of each of the sgRNA expressing constructs, we

first generated male flies that expressed both Cas9 and sgRNAs in their germ cells.

Individual females homozygous for each of the three sgRNA constructs

(RpS19a+RpL35, RpL36+RpL35 and *wupA*) were crossed to males containing (i)

Cas9 under the control of the *nanos* promoter (Either NIG CAS-0001; y^2 cho^2 v^1;

attP40{nos-Cas9}/CyO which carries a recessive lethal allele on the second

chromosome and cannot be made homozygous) or (ii) a line we generated that has

nos-Cas9 (from Addgene plasmid 62208 courtesy of Simon Bullock which has a

single NLS and a 3'UTR from *nanos*) placed in the 09w second chromosome attP40

landing site (referred to herein as nos-Cas9(II)_NH) or (iii) a *vasa*-cas9 on the Y

chromosome from BL91386 (Gamez et al. 2021). The paternal inheritance of Cas9

prevented maternal deposition and somatic activity of the drive (Champer *et al.* 2019)

and is consistent with the ultimate aim of placing X-poisoning transgenes on the Y

chromosome. Initially, single males heterozygous for Cas9 and one of the sgRNA

constructs were crossed to three 09w 3–4-day old virgins (14 replicates) and allowed

to mate and lay in a vial for seven days before being cleared and all offspring that

emerged counted. Subsequent crosses (14 replicates) included single nos-

Cas9/*wupA*-sgRNA males left to mate with single females for four hours, or

alternatively crossed to 3-day-old virgin females with the genotype C(1)DX,y^1,f^1/Y

(Bloomington Stock number 4559; 7 replicates), which is an 'attached-X' line that

forces paternal X chromosome inheritance.

Analysis of life stage associated with *wupA* lethality

Next we assessed the lifestage in which sex skewing occurred with *wupA* X-

poisoning crosses. Single males with a GFP marked X-chromosome (y[1]

M{RFP[3xP3.PB] GFP[E.3xP3]=vas-int.Dm}Zh-2A w[*]/Y; sg(*wupA*)/nos-Cas9 were

crossed to three wildtype virgin females (DGRP line 897) and allowed to mate for 48

hours. The control cross was the same except there was no Cas9. All flies where

then transferred to a vial with grape juice agar and allowed to lay eggs for 14 hours,

after which the flies were cleared. At the time of clearing, the total number of eggs

laid were counted. Twenty-four hours later the eggs were sexed and the female eggs

(GFP) were separated from the male eggs (non-GFP). We then tracked how many of

the female eggs successfully emerged into larvae, pupae, and adults. We performed

20 replicates for each cross.

Competitive mating assays

We assessed mate choice by placing one *ebony* (dark body colour) homozygous

female in a vial with one *ebony* homozygous male and one test male of the following

genotypes: A) sg(*wupA*)/nos-Cas9, B) sg(*wupA*)/+, C) +/nos-Cas9, D) 09w. The flies

were allowed to mate for 24 hours, after which the parents were cleared and the

progeny was phenotyped for body colour once they reached adulthood. All flies were

2-3 day old virgins at the time of mating.

Sequence characterization of *wupA* survivors

Sons of crosses between X-poisoning males and attached-X females had their DNA

extracted and were characterized with Sanger sequencing using standard protocols.

PCR primers were used to amplify the DNA around all sgRNA targeted sites of a

single gene, these were purified with a spin column and Sanger sequenced by

Macrogen (see Table S1 for sequencing primers).

Two hundred of the surviving female progeny from the cross 09w x sg(*wupA*)/nos-

Cas9 were collected and pooled into batches of ten. Their DNA was extracted

following a standard phenol:chloroform DNA extraction protocol. PCR reactions were

then performed using the primers labelled "*wupA* sg# Illumina sequencing" in Table

S1. Each set of primers amplifies a specific sg(*wupA*) target site. The PCR products

of each female sample were pooled together and purified with Bioneer AccuPrep

PCR/Gel Purification Kit. Samples were sent for MiSeq 300 sequencing with AGRF.

The sequencing results were input into Crispresso2 for allele-specific quantification

of each sg(*wupA*) target site (Clement et al. 2019).

Statistical Analyses

All statistical analyses were undertaken in R (R Core team 2023). Sex ratio biasing in the *Drosophila* X-poisoning gene drive crosses were analysed using an unpaired two-tail t-test with a Bonferroni correction for multiple comparisons where applicable.

Results

X-poisoning targeting ribosomal protein genes

Four of the haplolethal genes described on the X chromosome of *Drosophila melanogaste*r (Cook *et al.* 2012) were targeted using three multi-sgRNA constructs and Cas9 expressed in the male germline via a *nanos* promoter (Figure 1). The expectation was that males carrying the Cas9 and sgRNA transgenes would have a reduced number of viable daughters relative to controls yet the same number of sons (Figure 2a). All the crosses targeting ribosomal protein encoding genes (RpS19a+RpL35, RpL36+RpL35) yielded significantly fewer offspring when compared to the controls ($p < 0.0001$, t-test with Bonferroni correction; Figure 2b; Table 1). The crosses involving the RpS19a+RpL35 sgRNA construct followed expectations with significantly fewer daughters than sons ($p < 0.0001$, t-test with Bonferroni correction) and no significant difference in the number of sons compared to the control. In contrast the RpL36+RpL35 crosses yielded significantly fewer sons *and* daughters compared to control ($p < 0.05$ and $p < 0.01$ respectively, t-test with Bonferroni correction) without a significant sex ratio bias (Table 1).

Wings-up A poisoning skews the sex ratio

When the CRISPR machinery was targeted to *wupA,* significant skews in sex ratio towards maleness was observed ($p < 0.0001$, t-test with Bonferroni correction, Figure 2b). However, we initially found the expression of sgRNAs targeting *wupA* did not result in a significant reduction in the total number of offspring produced when compared to controls (Fig. 2b). The highly significant reduction in the number of female offspring ($p < 0.0001$, t-test with Bonferroni correction), was offset by the significantly higher number of male offspring compared to control ($p < 0.0001$, t-test with Bonferroni correction). We hypothesised that the lack of reduction in reproductive output may be due to a carrying capacity within the vials with an excess of embryos being laid during the seven day period where remating was allowed. To address this possibility, single males heterozygous for both X-poisoning drive components (ie. the *wupA* targeting sgRNAs and nos-Cas9) were placed in vials with single females for four hours only. In these single pair crosses, a significant reduction in the total number of adult offspring ($p < 0.01$, t-test with Bonferroni correction) as well as a highly significant change in sex ratio was observed ($p < 0.0001$, t-test with

Bonferroni correction; Figure 2c) and there was no significant difference in the
number of males produced when compared to the control.

As *wupA* has very low expression in the early stages of embryos (first 8 hours;
modENCODE Temporal Expression Data Flybase) it contrasts to many ribosomal
proteins because it is not required for the viability of all cells and particularly not those
in early development. The gene *wupA* encodes Troponin I, a protein that inhibits the
interaction of myosin and actin in muscles although it is also thought to have non-
muscle functions that relate to the nuclear localization of at least one of its nine
isoforms (Casas-Tintó and Ferrús 2021, Flybase). To explicitly test where the sex-
skewing of *wupA* CRISPR/nos-Cas9 elicited its effect we tracked the fate of X-
chromosomes labelled with GFP. In the crosses we conducted, female embryos
glowed green (after the 3XP3 GFP transgene is turned on), whereas male embryos
and unfertilized eggs did not. Twenty single pair crosses were set up for the
haplolethal cross (males bearing *wupA* sgRNAs and Cas9) and for controls (the
males had *wupA* sgRNAs but lacked Cas9) and these revealed that the number of
female embryos did not differ significantly between the haplolethal cross and the
control cross, but the number of larvae, pupae and adults did (Figure 3). We could
determine that females of the haplolethal cross typically died in late embryogenesis
(dorsal closure was completed and we observed mouthparts and denticle belts
however segmental furrows appeared unevenly spaced which may be attributable to
abnormal muscle contraction; see Figure S1). Only ~13% of GFP eggs became 1st
instar larvae, whereas ~72% of GFP eggs from the control cross were observed to
develop into larvae (Figure 3). Consistent with earlier results the ratio of females
among adults in these experiments was ~11% in the haplolethal cross and ~52% in
the control. We also examined 15 adult females emerging from this cross and all
were fertile.

No fitness cost detected in nos-Cas9 *wupA* poisoning males

Transcriptome datasets (e.g. FlyAtlas2- Anatomy RNA-Seq) indicate that *wupA* is
expressed in the testis and so we also sought to determine whether CRISPR/Cas9 of
wupA during spermatogenesis affected the fertility of the bearer. Males generating
wupA alleles during spermatogenesis could have a phenotype that makes them less
successful in mating, perhaps because of female choice, or they could produce lower

quality sperm. To test for these possibilities we examined the fitness of males in a

competitive assay. Four classes of test males (those with CRISPR/nos-Cas9

targeting *wupA*, and three control lines: one that carried the Cas9 without the sgRNA,

another that had the sgRNA but no Cas9, and a third that had neither CRISPR

component) individually competed for a female homozygous for the recessive ebony

body colour mutation against an ebony male. Eighty-five vials containing two males

and one female were set up (approximately 20 vials for each of the four tester male

genotypes; Table 2) and the number of ebony and wildtype progeny counted. In the

crosses involving males with both the sgRNA's and Cas9 transgenes (cross A, Table

2) 84 of the 112 progeny were wildtype with ten of the former being females. So

these test males were much more competitive than ebony males, even though there

was a massive skew in the sex ratio so that only 12% of the progeny (10/84) were

daughters (Table 2). In contrast, ebony males were more successful than the males

of the three control crosses although only significantly so in cross D (Table 2).

If competition is assessed at the level of vials then 81 of the 85 vials produce either

ebony progeny or wildtype progeny. At this level the males with the full CRISPR/Cas9

machinery were also seen to be more successful than ebony males as they were the

only sires in 16 vials, whereas ebony were the only sires in 5 of the vials (P=0.03).

Again, this was in contrast to the crosses involving the three control genotypes where

the number of ebony either did not differ significantly from wildtype (sgRNA-only

control P=0.82; Cas9-only control P=0.21) or in the case of the background control

line were more fit (P=7.6×10^{-5}). So we found no evidence of a pre-laying fitness

deficit for males that have CRISPR machinery targeting *wupA* during

spermatogenesis in the *nanos* expression domain.

In contrast, we attempted to create *wupA*-poisoning males that would have Cas9

expressed under the *vasa* promoter, but this failed. We crossed males which had

vasa-Cas9 on the Y chromosome and females that were homozygous for the *wupA*

sgRNA construct. This yielded 264 daughters (that did not have Cas-9 because they

did not inherit the Y chromosome) and only two small and infertile males. This is

consistent with earlier reports that *vasa* expresses outside the germline (Gamez et al.

2021), and shows that X-poisoning with *wupA* could have drastic fitness costs if the

CRISPR machinery is active in broader expression domains.

The nos-Cas9 *wupA* sex biasing is not absolute

To understand why the sex biasing was not absolute, X-chromosomes that had survived being passed through the paternal germ line were characterized. If the targeted loci had been cut and reannealed, indels or single nucleotide mutations may have been introduced, and they would be revealed by sequence analysis at the sites targeted by the sgRNA's. Initially, to simplify the sequence analysis, we used $X^\wedge XY$ females (hereafter 'attached-X') which reverse the inheritance of the paternal X's so that they are passed to sons. Consistent with expectations, males expressing *nanos-*Cas9 and sgRNA's targeting *wupA* sired significantly more daughters than sons (the sex ratio bias being flipped due to sons paternally inheriting the X chromosome; $p<0.001$, t-test with Bonferroni correction). The RpS19a+RpL35 cross showed no significant sex ratio bias but had significantly fewer male offspring than the control and no significant difference in the number of females (Figure 2d; $p<0.0001$, t-test with Bonferroni correction). There was no reduction in the number of offspring produced by males expressing gRNA targeting RpL36+RpL35.

The X chromosomes of a total of 60 surviving sons (14 for *wupA*, 22 for RpS19a+RpL35 and 24 for RpL36+RpL35) from 15 independent crosses were Sanger sequenced and only the offspring of one male showed any sign of CRISPR cutting. A complex indel resulting in a net 9 nucleotide insertion that did not introduce a premature stop codon was recovered from a cross involving the RpL36+RpL35 sgRNAs (Figure S2). The fact that this individual survived suggests that gene drive resistance alleles could arise against RpL35 gRNA1. The data also concurs with the idea that these genes are indeed lethal genes as no frameshifting indels or nonsense mutations were recovered. More generally, these data are consistent with the notion that most of the surviving progeny carrying a paternal X never had their X chromosome cut by the CRISPR machinery.

Not all *wupA* mutations are haplolethal

Curiously, we found that the skew in sex ratio was more extreme when the *wupA-*poisoned X chromosome was passed to sons (2% ; via attached-X females) rather than to daughters (14% ; Figure 2d versus Figures 2b and 2c; Table 1). As the fathers in these crosses are the same, this difference is not due to the generation of

X-poisoned alleles. Instead, heterozygosity in daughters may protect against lethality

more than the exposed hemizygosity of sons, which suggests that some of the

CRISPR generated alleles may encode recessive lethals. If true then we reasoned it

might be possible to find classes of mutations in daughters of males that have *wupA*

targeted during spermatogenesis that do not occur in their sons (from crosses with

attached-X). Three PCR amplicons surrounding the three sgRNA sites in *wupA* were

sequenced from twenty pools of ten daughters using Illumina's short read MiSeq

sequencing technology. Strikingly, each of the 20 pools had a high frequency of

deletion alleles (between 5 and 18%) among the reads from the amplicon that

amplified across the site targeted by sgRNA2 (Figure 4). Given that each of the 20

pools was composed of ten daughters, and we saw no such indels among the 14

sons from the attached-X cross, a significant number of the mutations generated by

CRISPR-editing seem to be lethal (purged in sons) but not haplolethal (and so

present in sequences from daughters). Notably all of the high frequency indels

observed had a length divisible by three and therefore did not change the open

reading frame. So it would seem that frameshift alleles are haplolethal but some

indels that maintain the open reading frame are lethal but not haplolethal.

Discussion

For an X-poisoning strategy to elicit an effect in a single generation the targeted

genes need to be haplo-insufficient. Such is the knowledge associated with

Drosophila melanogaster that it is known there are eleven loci on the X chromosome

of *D. melanogaster* that are haplolethal or haplosterile (Cook *et al.* 2012). Haplolethal

genes can only exist on the X chromosome in species such as Diptera where

dosage compensation between the homogametic and heterogametic sex is due to

upregulation - doubling the output of the X chromosome in males, rather than the

inactivation of the X as seen in humans.

Eight of the X-linked haplolethals of *D. melanogaster* encode ribosomal proteins,

which contribute to the translational machinery of cells, and an insufficient quantity of

them leads to lethality or to the Minute syndrome where flies are tiny and sterile

(Marygold *et al.* 2007). Previously, Fasulo *et al.* 2020 pioneered the 'X-poisoning'

strategy by targeting Ribosomal protein genes S6 and S5a. They found that when

one particular sgRNA targeting RpS6 was combined with Cas9 driven by a sperm-
specific beta tubulin 85D promoter the sex ratio was biased so that greater than 92%
of progeny were males. Our targeting of RpL36, RpL35 and RpL19a did not produce
such extreme biasing. However, Fasulo *et al.* (2020) also showed heterogeneity in
the success of their sgRNA's with different guide RNA's targeting the same genes
skewing to various extents. Therefore, the failure to see skewing in our combination
of RpL36 and RpL35 sgRNA's or the skew to only 80% males in the RpL35 and
RpL19a combination, may well have more to do with the particularities of guide RNA
efficacy than being a property of the genes they target, and in our case by subtle
design features such as the position of particular guide RNAs in the array may be
important.
A surprising finding of Fasulo *et al.* (2020) was that some alleles among the
survivors contained frameshift mutations. This led the authors to suggest that the
targeted ribosomal protein genes may not be haplolethal genes after all and that
previously studied alleles may have harboured dominant negative mutations instead.
In our analyses, we did not find such frameshifting alleles among individuals carrying
X-chromosomes that had passed through X-poisoning males. Instead, the loci
showed no signs of being cut at all or exhibited mutations that maintained the frame
(Figure 4b for sgRNA2 of *wupA* and Figure S3 for sgRNA1 and sgRNA3 of *wupA*).
390
The *wupA* locus of Drosophila was the only non-ribosomal protein haplolethal gene
that has been molecularly characterised on the X chromosome and it showed the
greatest skewing in our X-poisoning experiments. It encodes Troponin I, a protein
that inhibits the interaction of myosin and actin in muscles. It is also thought to have
non-muscle functions that relate to the nuclear localization of at least one of its
isoforms (Casas-Tintó and Ferrús 2021, Flybase). The gene has 12 exons that are
variously combined to produce 13 alternate transcriptional splice forms. Two of the
three sgRNAs used in our study target exons common to all splice forms (sgRNA2
and sgRNA3), and the third targets a PAM site that is found in the first two codons of
isoforms with the most common start methionine. The latter site was previously
targeted with CRISPR by Casas-Tintó and Ferrús (2021) and the resulting
haplolethality is consistent with our findings.
403

Our results provide an insight into the distinct functions of *wupA* because indels were

enriched around the sgRNA2 target site which lies on top of the methionine codon

used by *wupA* isoforms that translocate into the nucleus. We infer that such

mutations are recessive lethals that are exposed in the hemizygous condition of

males. However, as heterozygotes in females they produce Troponin I with an intact

actin-binding site and an intact Troponin T interaction site and so they are not

haplolethal. The haplolethality of *wupA* seems to relate to the delicate stoichometry

required for Troponin I and F-actin. Without sufficient Troponin I, calcium is not

required to allow tropomyosin to move and allow actin to interact with myosin. Thus,

lethality arises from the irreversible hyper-contraction of muscles. Such a model is

consistent with observations that mild *wupA* alleles (yielding the wings-up

phenotype) can be rescued by a mild allele of tropomyosin (Naimi et al. 2001).

There is much about *wupA* that makes it an attractive target for future X-poisoning

strategies. As a haplolethal gene not expressed until late embryogenesis, the

progeny of a poisoned cross are unaffected at fertilization and egg-lay. It thus

contrasts with X-poisoning targeting ribosomal proteins which may have reduced

fertility because spermatogenesis may be affected. It also contrast with genetic

systems causing sterility (such as PG-SIT (Kandul et al. 2019)) which can be

counteracted by females mating multiple times. We also did not find males that

poison *wupA* to have reduced fitness, although we acknowledge that our competitive

assays are likely be confounded with genetic background effects (especially eye-

colour attributable to the number of transgenes) but argue that if there is any fitness

cost attributable to *wupA* poisoning using a *nanos*-driven Cas9 it is less than the loss

of a second transgenic copy of *mini-white* and less than that of ebony males.

The *wupA gene* is a highly conserved gene that is found on the X chromosome on

numerous dipteran species, including some important pests (Figure S4). *Drosophila*

suzukii, an invasive pest of soft-skinned fruits, appears to share all 12 exons with *D.*

melanogaster and all the predicted proteins have greater than 98% amino acid

identity with *D. melanogaster* homologs. In fact, two of the three sgRNA's used here

in *D. melanogaster* match *D. suzukii* targets perfectly, and the third has only one

nucleotide different. *wupA* of Anopheles mosquito's has a slightly different gene

structure (eg. Exon 1 and Exon 2 are fused eg. XM_053811545) however it is also

found on the X chromosome in these species suggesting it could be a target of X-

poisoning strategies in the many pest species of this genus. Unfortunately, *wupA*
orthologs are not on the X-chromosome of all economically important dipterans, such
as the tephritids which have Muller element F (chromosome 4 of Drosophila) as an X
chromosome.

Taken together the above results suggest that *wupA* could be a good target for an X-
poisoning gene drive. To create such a drive, transgenes encoding a germline
restricted Cas9 and the sgRNA's targeting *wupA* would ideally be placed on the Y
chromosome. Placing transgenes on the Y chromosome has been technically
challenging although it has been done (Gamez *et al.* 2021). Indeed, we examined the
one *D. melanogaster* line currently available that has a Cas9 transgene on the Y
chromosome however because it uses a *vasa* promoter that expresses outside the
germline males that poisoned the X during spermatogenesis could not be generated.
An important next step in this research program will be to overcome the challenge of
placing germline-limited drive components onto the Y chromosome.

Finally, we note that where an X-poisoning strategy can take the form of a Y-linked
editor (Burt and Deredec 2018) it may not necessarily drive through a population.
While fathers may only have sons, the Y-chromosome bearing the X-poison may not
produce more sons than any other Y-chromosome in a population and so it may not
be considered a gene drive. However, if X-poisoned males produce more sons than
other Y-chromosomes, perhaps as a result of reduced larval competition with sisters
(such as seen in the original *wupA* cross), then a Y-linked editor could indeed be a
gene drive. In this larval-competition scenario, as the frequency of the Y-linked editor
increases, then so too will the strength of the gene drive.

Data Availability Statement
464 All data is here or in the supplementary material.

Acknowledgements

We'd like to thank Owain Edwards, Max Scott and Nina Wedell for conversations that
helped develop this research and Jackson Champer for suggestions about construct design.
We also thank the anonymous reviewers for suggestions that improved the manuscript.

Funding

The research was funded by an Australian Research Council Discovery Project Grant
(DP190102512).

Conflict of Interests

We have no conflicts of interest to declare.

References

Akbari, Omar S., Chun-Hong Chen, John M. Marshall, Haixia Huang, Igor Antoshechkin, and
Bruce A. Hay. 2014. "Novel Synthetic *Medea* Selfish Genetic Elements Drive
Population Replacement in *Drosophila* ; a Theoretical Exploration of *Medea* -
Dependent Population Suppression." *ACS Synthetic Biology* 3 (12): 915–28.
https://doi.org/10.1021/sb300079h.
Alphey, Luke S., Andrea Crisanti, Filippo (Fil) Randazzo, and Omar S. Akbari. 2020.
"Standardizing the Definition of Gene Drive." *Proceedings of the National Academy*
of Sciences 117 (49): 30864–67. https://doi.org/10.1073/pnas.2020417117.
Burt, Austin. 2003. "Site-Specific Selfish Genes as Tools for the Control and Genetic
Engineering of Natural Populations." *Proceedings of the Royal Society B: Biological*
Sciences 270 (1518): 921–28. https://doi.org/10.1098/rspb.2002.2319.
Burt, Austin, and Anne Deredec. 2018. "Self-Limiting Population Genetic Control with Sex-
Linked Genome Editors." *Proceedings of the Royal Society B: Biological Sciences* 285
(1883): 20180776. https://doi.org/10.1098/rspb.2018.0776.
Bushland, R. C., A. W. Lindquist, and E. F. Knipling. 1955. "Eradication of Screw-Worms
through Release of Sterilized Males." *Science* 122 (3163): 287–88.
https://doi.org/10.1126/science.122.3163.287.
Casas-Tintó, Sergio, and Alberto Ferrús. 2021. "The Haplolethality Paradox of the wupA
Gene in Drosophila." Edited by R. Scott Hawley. *PLOS Genetics* 17 (3): e1009108.
https://doi.org/10.1371/journal.pgen.1009108.
Champer, Jackson, Anna Buchman, and Omar S. Akbari. 2016. "Cheating Evolution:
Engineering Gene Drives to Manipulate the Fate of Wild Populations." *Nature*
Reviews Genetics 17 (3): 146–59. https://doi.org/10.1038/nrg.2015.34.
Champer, Jackson, Jingxian Liu, Suh Yeon Oh, Riona Reeves, Anisha Luthra, Nathan Oakes,
Andrew G. Clark, and Philipp W. Messer. 2018. "Reducing Resistance Allele
Formation in CRISPR Gene Drive." *Proceedings of the National Academy of Sciences*
115 (21): 5522–27. https://doi.org/10.1073/pnas.1720354115.
Champer, Jackson, Riona Reeves, Suh Yeon Oh, Chen Liu, Jingxian Liu, Andrew G. Clark, and
Philipp W. Messer. 2017. "Novel CRISPR/Cas9 Gene Drive Constructs Reveal Insights
into Mechanisms of Resistance Allele Formation and Drive Efficiency in Genetically
Diverse Populations." Edited by Harmit S. Malik. *PLOS Genetics* 13 (7): e1006796.
https://doi.org/10.1371/journal.pgen.1006796.
Champer, Jackson, Zhaoxin Wen, Anisha Luthra, Riona Reeves, Joan Chung, Chen Liu, Yoo
Lim Lee, et al. 2019. "CRISPR Gene Drive Efficiency and Resistance Rate Is Highly
Heritable with No Common Genetic Loci of Large Effect." *Genetics* 212 (1): 333–41.
https://doi.org/10.1534/genetics.119.302037.
Chen, Chun-Hong, Haixia Huang, Catherine M. Ward, Jessica T. Su, Lorian V. Schaeffer, Ming
Guo, and Bruce A. Hay. 2007. "A Synthetic Maternal-Effect Selfish Genetic Element

Drives Population Replacement in Drosophila." *Science* 316 (5824): 597–600.
https://doi.org/10.1126/science.1138595.

Clement, Kendell, Holly Rees, Matthew C. Canver, Jason M. Gehrke, Rick Farouni, Jonathan
Y. Hsu, Mitchel A. Cole, et al. 2019. "CRISPResso2 Provides Accurate and Rapid
Genome Editing Sequence Analysis." *Nature Biotechnology* 37 (3): 224–26.
https://doi.org/10.1038/s41587-019-0032-3.

Cook, R Kimberley, Stacey J Christensen, Jennifer A Deal, Rachel A Coburn, Megan E Deal, Jill
M Gresens, Thomas C Kaufman, and Kevin R Cook. 2012. "The Generation of
Chromosomal Deletions to Provide Extensive Coverage and Subdivision of the
Drosophila Melanogaster Genome." *Genome Biology* 13 (3).
https://doi.org/10.1186/gb-2012-13-3-r21.

Curtis, C. F. 1968. "A Possible Genetic Method for the Control of Insect Pests, with Special
Reference to Tsetse Flies (Glossina Spp.)." *Bulletin of Entomological Research* 57 (4):
509–23.

Fasulo, Barbara, Angela Meccariello, Maya Morgan, Carl Borufka, Philippos Aris Papathanos,
and Nikolai Windbichler. 2020. "A Fly Model Establishes Distinct Mechanisms for
Synthetic CRISPR/Cas9 Sex Distorters." *PLoS Genetics* 16 (3): 1–22.
https://doi.org/10.1371/journal.pgen.1008647.

Galizi, Roberto, Lindsey A. Doyle, Miriam Menichelli, Federica Bernardini, Anne Deredec,
Austin Burt, Barry L. Stoddard, Nikolai Windbichler, and Andrea Crisanti. 2014. "A
Synthetic Sex Ratio Distortion System for the Control of the Human Malaria
Mosquito." *Nature Communications* 5 (1): 3977.
https://doi.org/10.1038/ncomms4977.

Gamez, Stephanie, Duverney Chaverra-Rodriguez, Anna Buchman, Nikolay P. Kandul, Stelia
C. Mendez-Sanchez, Jared B. Bennett, Héctor M. Sánchez C., et al. 2021. "Exploiting a
Y Chromosome-Linked Cas9 for Sex Selection and Gene Drive." *Nature
Communications* 12 (1): 7202. https://doi.org/10.1038/s41467-021-27333-1.

Gratz, Scott J., Fiona P. Ukken, C. Dustin Rubinstein, Gene Thiede, Laura K. Donohue,
Alexander M. Cummings, and Kate M. O'Connor-Giles. 2014. "Highly Specific and
Efficient CRISPR/Cas9-Catalyzed Homology-Directed Repair in Drosophila." *Genetics*
196 (4): 961–71. https://doi.org/10.1534/genetics.113.160713.

Haber, Daniella An, Yael Arien, Lee Benjamin Lamdan, Yehonathan Alcalay, Chen Zecharia,
Flavia Krsticevic, Elad Shmuel Yonah, et al. 2023. "Targeting Mosquito X-
Chromosomes Reveals Complex Transmission Dynamics of Sex Ratio Distorting Gene
Drives." Preprint. Synthetic Biology. https://doi.org/10.1101/2023.08.20.554004.

Hammond, Andrew M., Kyros Kyrou, Marco Bruttini, Ace North, Roberto Galizi, Xenia
Karlsson, Nace Kranjc, et al. 2017. "The Creation and Selection of Mutations
Resistant to a Gene Drive over Multiple Generations in the Malaria Mosquito."
Edited by Nick H. Barton. *PLOS Genetics* 13 (10): e1007039.
https://doi.org/10.1371/journal.pgen.1007039.

Hay, Bruce A., Georg Oberhofer, and Ming Guo. 2021. "Engineering the Composition and
Fate of Wild Populations with Gene Drive." *Annual Review of Entomology* 66 (1):
407–34. https://doi.org/10.1146/annurev-ento-020117-043154.

Hendrichs, J., G. Franz, and P. Rendon. 1995. "Increased Effectiveness and Applicability of
the Sterile Insect Technique through Male-Only Releases for Control of
Mediterranean Fruit Flies during Fruiting Seasons." *Journal of Applied Entomology*
119 (1–5): 371–77. https://doi.org/10.1111/j.1439-0418.1995.tb01303.x.

Kandul, Nikolay P., Junru Liu, Hector M. Sanchez C., Sean L. Wu, John M. Marshall, and Omar
563   S. Akbari. 2019. "Transforming Insect Population Control with Precision Guided
564   Sterile Males with Demonstration in Flies." *Nature Communications* 10 (1): 84.
565   https://doi.org/10.1038/s41467-018-07964-7.
566 Knipling, E.F. 1955. "Possibilities of Insect Control or Eradication through the Use of Sexually
567   Sterile Males." *Journal of Economic Entomology,* 48 (4): 459–62.
568 Kyrou, Kyros, Andrew M. Hammond, Roberto Galizi, Nace Kranjc, Austin Burt, Andrea K.
569   Beaghton, Tony Nolan, and Andrea Crisanti. 2018. "A CRISPR–Cas9 Gene Drive
570   Targeting Doublesex Causes Complete Population Suppression in Caged Anopheles
571   Gambiae Mosquitoes." *Nature Biotechnology* 36 (11): 1062–66.
572   https://doi.org/10.1038/nbt.4245.
573 Labun, Kornel, Tessa G Montague, Maximilian Krause, Yamila N Torres Cleuren, Håkon
574   Tjeldnes, and Eivind Valen. 2019. "CHOPCHOP v3: Expanding the CRISPR Web
575   Toolbox beyond Genome Editing." *Nucleic Acids Research* 47 (W1): W171–74.
576   https://doi.org/10.1093/nar/gkz365.
577 Larkin, Aoife, Steven J Marygold, Giulia Antonazzo, Helen Attrill, Gilberto dos Santos, Phani V
578   Garapati, Joshua L Goodman, et al. 2021. "FlyBase: Updates to the *Drosophila*
579   *Melanogaster* Knowledge Base." *Nucleic Acids Research* 49 (D1): D899–907.
580   https://doi.org/10.1093/nar/gkaa1026.
581 Marygold, Steven J., John Roote, Gunter Reuter, Andrew Lambertsson, Michael Ashburner,
582   Gillian H. Millburn, Paul M. Harrison, et al. 2007. "The Ribosomal Protein Genes and
583   Minute Loci of Drosophila Melanogaster." *Genome Biology* 8 (10): R216.
584   https://doi.org/10.1186/gb-2007-8-10-r216.
585 Meccariello, Angela, Flavia Krsticevic, Rita Colonna, Giuseppe Del Corsano, Barbara Fasulo,
586   Philippos Aris Papathanos, and Nikolai Windbichler. 2021. "Engineered Sex Ratio
587   Distortion by X-Shredding in the Global Agricultural Pest Ceratitis Capitata." *BMC*
588   *Biology* 19 (1): 78. https://doi.org/10.1186/s12915-021-01010-7.
589 Naimi, Benyoussef, Andrew Harrison, Mark Cummins, Upendra Nongthomba, Samantha
590   Clark, Inmaculada Canal, Alberto Ferrus, and John C. Sparrow. 2001. "A
591   Tropomyosin-2 Mutation Suppresses a Troponin I Myopathy in *Drosophila*." Edited
592   by John Pringle. *Molecular Biology of the Cell* 12 (5): 1529–39.
593   https://doi.org/10.1091/mbc.12.5.1529.
594 Oberhofer, Georg, Tobin Ivy, and Bruce A. Hay. 2019. "Cleave and Rescue, a Novel Selfish
595   Genetic Element and General Strategy for Gene Drive." *Proceedings of the National*
596   *Academy of Sciences* 116 (13): 6250–59. https://doi.org/10.1073/pnas.1816928116.
597 Port, Fillip, and Simon L Bullock. 2016. "Augmenting CRISPR Applications in Drosophila with
598   tRNA-Flanked sgRNAs." *Nature Methods* 13 (10): 852–54.
599   https://doi.org/10.1038/nmeth.3972.
600 R Core team. 2023. "R Core Team (2022). R: A Language and Environment for Statistical
601   Computing. R Foundation for Statistical Computing, Vienna, Austria." https://www.R-
602   project.org/.
603 Rose, Graham, Elzbieta Krzywinska, Jan Kim, Loic Revuelta, Luca Ferretti, and Jaroslaw
604   Krzywinski. 2016. "Dosage Compensation in the African Malaria Mosquito *Anopheles*
605   *Gambiae*." *Genome Biology and Evolution*, January, evw004.
606   https://doi.org/10.1093/gbe/evw004.
607 Scott, Maxwell J., Carolina Concha, John B. Welch, Pamela L. Phillips, and Steven R. Skoda.
608   2017. "Review of Research Advances in the Screwworm Eradication Program over

the Past 25 Years." *Entomologia Experimentalis et Applicata* 164 (3): 226–36.
https://doi.org/10.1111/eea.12607.
Serebrovsky, A.S. 1969. "On the Possibility of a New Method for the Control of Insect Pests."
In *Proceedings of a Panel on Application of the Sterile-Male Technique for the*
Eradication or Control of Harmful Species of Insects, Organised by the Joint FAO/IAEA
Division of Atomic Energy in Food and Agriculture, pp.123-237. Vienna, 27-31 May
1968.
Waltz, Emily. 2021. "First Genetically Modified Mosquitoes Released in the United States."
Nature 593 (7858): 175–76. https://doi.org/10.1038/d41586-021-01186-6.
Webster, Sophia H., Michael R. Vella, and Maxwell J. Scott. 2020. "Development and Testing
of a Novel Killer–Rescue Self-Limiting Gene Drive System in *Drosophila*
Melanogaster." *Proceedings of the Royal Society B: Biological Sciences* 287 (1925):
20192994. https://doi.org/10.1098/rspb.2019.2994.
Whitten, M.J. 1985. "The Conceptual Basis for Genetic Control." In *Comprehensive Insect*
Physiology Biochemistry and Pharmacology, 12:465–528. Insect Control. Oxford,
New York, Toronto, Sydney, Paris, Frankfurt: Pergamon Press.
Windbichler, N., P. A. Papathanos, F. Catteruccia, H. Ranson, A. Burt, and A. Crisanti. 2007.
"Homing Endonuclease Mediated Gene Targeting in Anopheles Gambiae Cells and
Embryos." *Nucleic Acids Research* 35 (17): 5922–33.
https://doi.org/10.1093/nar/gkm632.
Windbichler, Nikolai, Miriam Menichelli, Philippos Aris Papathanos, Summer B. Thyme, Hui
Li, Umut Y. Ulge, Blake T. Hovde, et al. 2011. "A Synthetic Homing Endonuclease-
Based Gene Drive System in the Human Malaria Mosquito." *Nature* 473 (7346): 212–
15. https://doi.org/10.1038/nature09937.
Yang, Emily, Matthew Metzloff, Anna M Langmüller, Xuejiao Xu, Andrew G Clark, Philipp W
Messer, and Jackson Champer. 2022. "A Homing Suppression Gene Drive with
Multiplexed gRNAs Maintains High Drive Conversion Efficiency and Avoids Functional
Resistance Alleles." Edited by M Ramaswami. *G3 Genes|Genomes|Genetics* 12 (6):
jkac081. https://doi.org/10.1093/g3journal/jkac081.
Yao, Franck Adama, Abdoul-Azize Millogo, Patric Stephane Epopa, Ace North, Florian Noulin,
Koulmaga Dao, Mouhamed Drabo, et al. 2022. "Mark-Release-Recapture Experiment
in Burkina Faso Demonstrates Reduced Fitness and Dispersal of Genetically-Modified
Sterile Malaria Mosquitoes." *Nature Communications* 13 (1): 796.
https://doi.org/10.1038/s41467-022-28419-0.
643
644

Figure Legends

Figure 1: Targeting Haplolethal genes on the X chromosome. (a) Premeiotic cells expressing CRISPR/Cas9 (from the autosomes) that effectively target haplolethal loci on the X-chromosome (blue to red) will result in male only offspring. If the machinery were put on the Y-chromosome then inheritance of the elements will be limited to the male germ line and a gene drive would be created. (b) Eleven haplolethal/haplosteriles were reported by Cook *et al.* (2012), two (RpS6 and RpS5a) were targeted by Fasulo *et al.* 2020 and the location of the four loci targeted in this manuscript, and the number of sgRNA used (rounded blue rectangles) are shown. (c) The *wupA* gene encodes for thirteen transcript isoforms. SgRNA1-3 target three conserved regions of the gene's coding sequence.

Figure 2: Sex biases with X-poisoning. (a) A symbolic representation of what is expected in our crosses. A skew against daughters (red) is expected when sgRNA's target haplolethal loci on the X chromosome during gametogenesis. The number of sons is expected to remain unchanged as they would arise from sperm carrying the Y chromosome. (b) The observed amount of sex biasing skew ranged from nothing (RpL36+RpL35), to modest (RpL19a+RpL35), to promising (wupA). Note that the total number of offspring was also variously affected. (c) By excluding multiple mating the total number of offspring in the *wupA* cross was reduced and there was a strong skew towards males (d) CRISPR/Cas9 fathers crossed to attached-X mothers, reverse the inheritance of the sex chromosomes so that sons inherited the CRISPR modified X chromosome, and daughters inherited the Y chromosome. (* P < 0.05, ** P < 0.01, ** P < 0.001 **** P < 0.0001, unpaired two-tail t-test with a Bonferroni correction for multiple comparisons).

Figure 3: Developmental stage of lethality from *wupA* poisoned fathers (a) The average number of daughters tracked from 20 vials of the haplolethal (*wupA*-poisoned) cross (in red) and 20 vials from a control cross (grey). Error bars represent the standard error of the mean and asterisks represent significance (P<0.05). (b) The

total number of daughters from 20 vials from the haplolethal (red) and control cross

(grey) at each life stage.

Figure 4: Recessive lethal *wupA* mutations revealed in females (a) The three

sites within *wupA* that are targeted by sgRNAs (see figure 1c) vary in the number of

CRISPR mutations uncovered by sequencing 200 daughters of an X-poisoning

cross. Here a Crispresso2 output shows the total number of modified and unmodified

reads from Illumina sequencing at each target site. (b) A nucleotide alignment of the

modified reads of target site 2 showing their frequencies. Amino acid sequences are

shown at the top, the sgRNA sequence is highlighted in grey, the PAM site is

underlined, the red arrow is the cut site, and black blocks represent insertions

(shown more to detail in Figure S3). (c) Isoform G of Tropinin I without any mutations

(blue) and with the largest deletion found in our sequencing results (a 21 bp deletion

shown in red). Protein folding predicted with AlphaFold. The deletion location is

highlighted in grey. No mutations in target site 2 will affect the acting-binding and

troponin-interacting areas.

Table 1: Sex skews in X-poisoning crosses and controls

gene/s targeted with sgRNA	nos-cas9 used	number of fathers used per vial	number of mothers per vial	Number of replicate vials used	Stock origin of mothers	number of sons	number of daughters	proportion of rarer sex	Drive efficiency (%)*
None	CAS-0001	1	3	14	09w	1014	1081	0.48	-3
None	CAS-0001	1	1	14	09w	603	544	0.47	5
None	CAS-0001	1	3	4	C(1)DX, y[1]f[1]/Y	182	151	0.45	9
wupA	CAS-0001	1	3	14	09w	1409	176	0.11	78
wupA	CAS-0001	1	1	14	09w	698	94	0.12	76
wupA	NH-noscas9	3	10	18	09w	1569	257	0.14	72
wupA	CAS-0001	1	3	6	C(1)DX, y[1]f[1]/Y	7	290	0.02	95
wupA	NH-noscas9	3	10	18	C(1)DX, y[1]f[1]/Y	14	780	0.02	96
RpS19a+RpL35	CAS-0001	1	3	14	09w	789	195	0.20	60
RpS19a+RpL35	CAS-0001	1	3	7	C(1)DX, y[1]f[1]/Y	55	143	0.28	44
RpL36+RpL35	CAS-0001	1	3	14	10w	678	717	0.49	-3
RpL36+RpL35	CAS-0001	1	3	6	C(1)DX, y[1]f[1]/Y	234	237	0.50	1

694 *Two times the difference between the expected frequency of 50% and the observed % of rarer sex.
695

696 **Table 2: Competitive mating assays**

Cross	A	B	C	D
Test male genotype	sg(*wupA*)/ nos-Cas9	sg(*wupA*)/+	+/nos-Cas9	+/+
Total vials assayed	21	19	23	22
Vials with only *ebony* offspring	5	9	14	17
Vials with only wild-type offspring	16	9	9	2
Total *ebony* progeny (% female)	28 (64%)	98 (52%)	151 (50%)	213 (51%)
Total wild-type progeny (% female)	84 (12%)	57 (65%)	106 (49%)	76 (47%)

698
699
700

(a)

(b)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20

RpL17   Fs(1)10A   Hdl RpL37a   RpS10B
RpS6   RpS5a

RpL36   RpL35   RpS19a   wupA

= sgRNA

A = RpL36 + RpL35 construct
B = RpS19a + RpL35 construct
C = wupA construct

A

B

C

(c)

wupA gene (11,355 bp)

* SgRNA Targets

Isoforms
A
B
C
D
E
F
G
H
I
K
L
M

M A D D
A T G G C T G A T G A T
SgRNA1 PAM

M E E A
A T G G A G G A A G C C
SgRNA2 PAM

C G S P
T G C G G C A G T C C C
SgRNA3 PAM

Figure 2

(a)

(b)

(a)

(b)

	Change in length	Frequency
	-21	0.044
	-21	0.018
	-18	0.305
	-18	0.008
	-18	0.011
	-18	0.011
	-15	0.130
	-12	0.030
	-12	0.007
	-12	0.009
	-9	0.126
	-9	0.018
	-9	0.007
	-6	0.091
	-9	0.014
	-3	0.103
	+6	0.047
	+3	0.012
	0	0.009

(c)