Metabolic and Neurobehavioral Disturbances Induced by Purine Recycling Deficiency in *Drosophila*

Céline Petitgas[1,4], Laurent Seugnet[2], Amina Dulac[1], Ali Mteyrek[1], Rebecca Fima[1], Marion Strehaiano[1], Joana Dagorret[1], Baya Chérif-Zahar[1], Sandrine Marie[3], Irène Ceballos-Picot[4,5] and Serge Birman[1,*]

[1]*Genes Circuits Rhythms and Neuropathology, Brain Plasticity Unit, CNRS, ESPCI Paris, PSL Research University, 75005 Paris, France*

[2]*Integrated Physiology of the Brain Arousal Systems (WAKING), Lyon Neuroscience Research Centre, INSERM/CNRS/UCBL1, 69675 Bron, France*

[3]*Laboratory of Metabolic Diseases, Cliniques Universitaires Saint-Luc, Université catholique de Louvain, B-1200, Brussels, Belgium*

[4]*Metabolomic and Proteomic Biochemistry Laboratory, Necker-Enfants Malades Hospital, AP-HP, 149 rue de Sèvres, 75015 Paris, France*

[5]*Paris Cité University, 12 rue de l'École de Médecine, 75006 Paris, France*

*Author for correspondence: serge.birman@espci.fr

Abstract

Lesch-Nyhan disease (LND) is a rare genetic disorder induced by deficiency in

hypoxanthine-guanine phosphoribosyltransferase (HGPRT), an enzyme of the purine salvage

pathway. This leads in early age to hyperuricemia and severe neurobehavioral disturbances,

including dystonia, spasticity and compulsive self-injury. To date, no treatment is available for

these neurological symptoms and no animal model recapitulates all the defects observed in LND

patients. Here we studied LND-related mechanisms in the fruit fly *Drosophila melanogaster*. We

confirmed that no HGPRT activity is expressed in this organism, where the only purine-recycling

enzyme is adenine phosphoribosyltransferase (Aprt). This enzyme is also present in humans but

its deficiency does not trigger neurological defects. In contrast, we observed that *Drosophila Aprt*

mutants showed both metabolic and neurobehavioral disturbances, including increased uric acid

levels, locomotor reactivity impairments, sleep alterations, seizure-like behavior, reduced

lifespan, and reduction of adenosine signaling and content. Locomotor defects could be rescued

by neuronal *Aprt* re-expression in mutant context and reproduced by knocking down *Aprt*

selectively in the protocerebral anterior medial (PAM) clusters of dopaminergic neurons, the

mushroom bodies and glia subsets. Ingestion of allopurinol normalized uric acid levels in *Aprt*

mutants but not their neurological defects, as is the case in LND patients, whereas feeding

adenosine or N^6-methyladenosine during development fully rescued the epileptic behavior.

Intriguingly, pan-neuronal expression of an LND-associated mutant form of human HGPRT

(I42T), but not the wild-type enzyme, resulted in early locomotor defects and seizure in flies,

similar to *Aprt* deficiency. Overall, this shows that *Drosophila* can be used as a new model in

different ways to better understand LND and seek a cure for this dramatic disease.

Keywords

Adenine phosphoribosyltransferase (APRT) – Adenosine signaling – *Drosophila melanogaster* – Hypoxanthine-guanine phosphoribosyltransferase (HGPRT) – Lesch-Nyhan disease – Mushroom bodies – PAM dopamine neurons – Purine salvage pathway – Seizure-like behavior – Startle-induced locomotion – Sleep – Uric acid level

Introduction

The purine salvage pathway is an essential component of cellular metabolism that allows the recovery of free purine bases derived from the diet or from the degradation of nucleic acids and nucleotides, thus avoiding the energy cost of *de novo* purine biosynthesis (Nyhan et al. 2014). Energy-intensive tissues, such as cardiac muscle cells and brain neurons, extensively use this pathway to maintain their purine levels (Ipata 2011, Johnson et al. 2019). The two main recycling enzymes involved in the salvage pathway in mammals are hypoxanthine-guanine phosphoribosyltransferase (HGPRT), which converts hypoxanthine and guanine into IMP and GMP, respectively, and adenine phosphoribosyltransferase (APRT), which converts adenine into AMP.

APRT and HGPRT deficiency induce very different disorders in humans. Loss of APRT seems to have only metabolic consequences, leading to the formation of 2,8-dihydroxyadenine crystals in kidney, which can be fatal but is readily prevented by allopurinol treatment (Bollée et al. 2012; Harambat et al. 2012). In contrast, highly inactivating mutations in HGPRT trigger Lesch-Nyhan disease (LND), a rare neurometabolic X-linked recessive disorder with dramatic consequences for child neurodevelopment (Lesch and Nyhan 1964, Seegmiller et al. 1967). The metabolic consequence of HGPRT deficiency is an overproduction of uric acid in the blood

(hyperuricemia) that can lead to gout and tophi, or nephrolithiasis (Sass et al. 1965; Kelley et al.

1967). Affected children also develop severe neurological impairments, such as dystonia,

choreoathetosis, spasticity, and a dramatic compulsive self-injurious behavior (Nyhan 1997;

Jinnah et al. 2006; Torres and Puig 2007; Schretlen et al. 2005; Madeo et al. 2019). They have a

developmental delay from 3 to 6 months after birth and most of them never walk or even sit

without support. Xanthine oxidase inhibitors, such as febuxostat or allopurinol, are automatically

given to patients after diagnosis to decrease their uric acid levels and prevent the formation of

urate crystals in kidney, which can lead to renal failure (Kelley et al. 1967; Torres et al. 2007;

Lahaye et al. 2014). However, no treatment is yet available to abrogate the neurobehavioral

symptoms of LND (Torres et al. 2007; Jinnah et al. 2010; Madeo et al. 2019).

Noticeably, the clinical severity in LND appears to be inversely correlated to the

enzymatic activity of HGPRT (Fu and Jinnah 2012; Fu et al. 2014). The full LND phenotype

which is the most severe corresponds to mutations that leave less than 2% of residual activity,

whereas 2-10% of activity leads to an intermediate phenotype with various neurological

abnormalities but without self-injury, and more than 10% protects from the neurobehavioral

symptoms and only results in hyperuricemia-related disorders. These two last groups are named

Lesch-Nyhan variants (Jinnah et al. 2010).

To date, the causes of the neurobehavioral troubles in LND are still not elucidated and it

remains challenging to understand and cure this dramatic disease (Jinnah et al. 2010, Bell et al.

2016). The most favored hypothesis is a dysfunction of the basal ganglia, and particularly of its

dopaminergic pathways (Baumeister and Frye 1985; Visser 2000; Nyhan 2000; Saito and

Takashima, 2000; Egami et al. 2007). Indeed, analyses of LND patient brains revealed a marked

loss of dopamine (DA) levels (Lloyd et al. 1981; Ernst et al. 1996) and DA transporters (Wong et

al. 1996) specifically in the basal ganglia, but the mechanisms linking HGPRT activity and the

DA system are still under investigation. DA deficits have been reported in HGPRT knockout

rodents, but these animal models do not exhibit the motor or neurobehavioral defects of human

patients (Finger et al. 1988; Dunnett et al. 1989; Jinnah et al. 1993; Jinnah et al. 1994; Meek et al.

2016). Recent studies reported that HGPRT deficiency disrupts the proliferation and migration of

developing midbrain dopamine (mDA) neurons in mouse embryos, arguing for a

neurodevelopmental syndrome (Witteveen et al. 2022). This could result from ATP depletion and

impaired energy metabolism (Bell et al., 2021) or an overactivation of *de novo* purine synthesis

leading to the accumulation of potentially toxic intermediates of this pathway (López, 2008;

López et al., 2020). The reason why dopaminergic neurons would be primarily affected in this

disease is not currently known. Pharmacological models have also been developed by injecting

the dopaminergic neurotoxin 6-hydroxydopamine into neonatally rat brains, which induced a

self-mutilation behavior in response to DA agonist administration in adulthood. However, these

models are of limited utility as they do not reproduce the basic genetic impairment of LND

(Breese et al. 1990; Knapp et al. 2016; Bell et al. 2016).

New animal models are therefore critically needed to study LND pathogenesis and find

efficient therapeutic molecules. The fruit fly *Drosophila melanogaster* presents many advantages

for translational studies and drug discovery (Fernández-Hernández et al. 2016, Perrimon et al.

2016; Papanikolopoulou et al. 2019). It has been widely used to model a variety of human

diseases, including neurodegenerative disorders such as Alzheimer, Huntington or Parkinson

diseases (Jaiswal et al. 2012; McGurk et al 2015; Dung and Thao 2018; Nagoshi 2018), as well

as heritable neurodevelopmental disorders such as Angelman, Rett or Fragile X syndromes

(Gatto and Broadie 2011). Although the importance of this invertebrate model for studying rare

human genetic diseases is now recognized (Oriel and Lasko 2018), a *Drosophila* LND model has

not yet been developed to our knowledge. This is probably due to the fact that no HGPRT

activity has been detected in this organism (Miller and Collins 1973; Becker 1974a; Becker

1974b). However, an ortholog of APRT is expressed in *Drosophila* (Johnson et al. 1987),

encoded by the *Adenine phosphoribosyltransferase* (*Aprt*) gene, which seems to be the only

enzyme of the purine salvage pathway in this insect. It is therefore possible that part of the

functions of human HGPRT, and in particular those essential for nervous system development

and physiology, were endorsed by Aprt in *Drosophila*.

Here, we tested whether purine recycling disruption in *Drosophila* can have similar

consequences as lack of HGPRT activity in humans, by studying the effects of *Aprt* deficiency on

purine metabolism, lifespan and various behaviors, including spontaneous and startle-induced

locomotion, sleep and seizure-like bang-sensitivity. We provide evidence that Aprt prolongs

lifespan and prevents seizure, and is required during development and in adult stage for many

aspects of *Drosophila* life. Its activity seems to be of particular importance in subpopulations of

dopaminergic neurons and glial cells in the brain. To our knowledge, this is the first LND model

that recapitulates both purine recycling disruption and neurobehavioral symptoms. We also find

that the expression of a LND-associated mutant form of HGPRT, but not the wild-type enzyme,

in *Drosophila* neurons, induced neurobehavioral impairments similar to those of *Aprt*-deficient

flies. Such a potential toxic gain-of-function effect of mutated HGPRT has not been

demonstrated in an animal model before.

Results

***Drosophila* lacking Aprt activity have a shortened lifespan**

To assess the importance of purine recycling in brain development and activity in

Drosophila, we analyzed the phenotypes induced by a deficiency in Aprt, which is potentially the

only recycling enzyme of the purine salvage pathway in this organism (Figure S1A and B, *step*

12). The *Aprt⁵* mutant was originally generated by chemical mutagenesis followed by selection of

flies resistant to purine toxicity (Johnson and Friedman, 1981; Johnson and Friedman, 1983).

Enzymatic assays confirmed a strong reduction in Aprt activity in extracts of heterozygous

Aprt⁵/+ mutants and its absence in homozygous and hemizygous (*Aprt⁵/Df(3L)ED4284*) mutants

(Figure S2 and Table S1). Sequencing of the *Aprt⁵* cDNA (Figure S3A) indicated that the *Aprt⁵*

mRNA codes for a protein with several amino acid changes compared to *Drosophila*

melanogaster wild-type Aprt, modifying in particular 3 amino acid residues that have been

conserved in the Aprt sequences from *Drosophila* to humans (Figure S3B). These mutations are

likely to be responsible for the loss of enzymatic activity. The homozygous *Aprt⁵* mutants are

considered viable because they develop and reproduce normally. However, we observed that

these mutants have a significantly reduced longevity, their median lifespan being only 38 days

against 50 days for wild-type flies ($p < 0.001$) (Figure 1A).

Uric acid levels are increased in *Aprt⁵* mutants and normalized by allopurinol

In humans, one of the consequences of HGPRT deficiency on metabolism is the

overproduction of uric acid (Harkness et al. 1988; Fu et al. 2015). We assayed the levels of

purine metabolites by HPLC and found that the level of uric acid was substantially increased by

37.7% on average in *Drosophila Aprt⁵* mutant heads ($p < 0.01$) (Figure 1B). We then tried to

normalize uric acid content in flies by providing allopurinol in the diet, as is usually done for

LND patients. Allopurinol is a hypoxanthine analog and a competitive inhibitor of xanthine

oxidase, an enzyme that catalyzes the oxidation of xanthine into uric acid (Figure S1A and B,

step 19). Remarkably, the administration of 100 µg/ml allopurinol during 5 days decreased uric

acid levels to a normal concentration range in *Aprt⁵* mutant heads (Figure 1B). Therefore, quite

similarly to HGPRT deficit in humans, the lack of Aprt activity in flies increases uric acid levels

and this metabolic disturbance can be prevented by allopurinol.

Figure 1. Aprt deficiency shortens lifespan and induces metabolic and neurobehavioral defects. (**A**) $Aprt^5$ mutant flies have a reduced lifespan compared to wild-type flies (median lifespan: 38 and 50 days, respectively). Three independent experiments were performed on 150 males per genotype with similar results and a representative experiment is shown. ***$p < 0.001$. (**B**) HPLC profiles on head extracts revealed an increase in uric acid levels in $Aprt^5$ mutant flies. Administration of 100 μg/ml allopurinol for 5 days before the test normalized uric acid levels. Mean of 3 independent experiments performed on 40 flies per genotype. *$p < 0.05$, **$p < 0.01$, ns: not significant. (**C-E**) Effect on climbing ability. (**C**) $Aprt^5$ mutants show early defects in the SING paradigm that monitors locomotor reactivity and climbing performance. This deficit was already obvious at 1 d a.E. (days after eclosion) and further decreased up to 8 d a.E., after which it did not change significantly up to 30 d a.E. Mean of 3 independent experiments performed on 50 flies per genotype. **$p < 0.01$, ***$p < 0.001$. (**D**) Downregulating *Aprt* by RNAi in all cells (*da>Aprt*[RNAi]) also led to an early impairment in climbing responses in the SING assay at 10 d a.E., as compared to the driver (*da/+*) and effector (*Aprt*[RNAi]/+) only controls. Mean of 3 independent experiments performed on 50 flies per genotype. ***$p < 0.001$. (**E**) Adult-specific inactivation of *Aprt* (*tub-Gal80*[ts]; *da-Gal4>Aprt*[RNAi]) decreased startle-induced climbing abilities in the SING paradigm, suggesting that the locomotor impairment induced by Aprt deficiency is not caused by a developmental effect. Flies were raised at permissive temperature (18°C) in which Gal80[ts] suppressed Gal4-controlled *Aprt*[RNAi] expression, and were shifted from 18 to 30°C for 3 days before the test (between 7 and 10 d a.E.) to activate transgene expression. Mean of 3 independent experiments performed on 50 flies per genotype. ***$p < 0.001$, ns: not significant.

Aprt deficiency decreases motricity in young flies

LND patients present dramatic motor disorders that prevent them for walking at an early age. To examine whether a deficiency in the purine salvage pathway can induce motor disturbance in young flies, we monitored the performance of *Aprt*-null mutants in startle-induced negative geotaxis (SING), a widely used paradigm to assess climbing performance and locomotor reactivity (Feany and Bender, 2000; Friggi-Grelin et al. 2003; Riemensperger et al. 2013; Sun et al. 2018). Strikingly, $Aprt^5$ mutant flies showed a very early SING defect starting from 1 day after eclosion (d a.E.) (performance index (PI) = 0.73 vs. 0.98 for wild-type flies, $p < 0.001$) that was more pronounced at 8 d a.E. (PI = 0.51 vs. 0.96 for the wild-type flies, $p < 0.001$). The fly locomotor performance did not further decline afterwards until 30 d a.E. (Figure 1C). *Df(3L)ED4284* and *Df(3L)BSC365* are two partially overlapping small genomic deficiencies that covers *Aprt* and several neighbor genes. Hemizygous $Aprt^5$/*Df(3L)ED4284* or

Aprt⁵/Df(3L)BSC365 flies also displayed SING defects at 10 d a.E. (PI = 0.71 and 0.68,

respectively, compared to 0.97 for wild-type flies, $p < 0.01$) (Figure S4). In contrast to its

beneficial effect on uric acid levels, we observed that allopurinol treatment did not improve the

locomotor ability of *Aprt⁵* mutant flies, either administered 5 days before the test or throughout

the development (Figure S5A, B). This is comparable to the lack of effect of this drug against

neurological defects in LND patients.

To confirm the effect of *Aprt* deficiency on the SING behavior, we used an *UAS-Aprt*ᴿᴺᴬⁱ

line that reduced Aprt activity by more than 95% when expressed in all cells with the *da-Gal4*

driver (Table S1). These *Aprt* knocked down flies also showed a strong SING defect at 10 d a.E.

(PI = 0.62 against 0.97 and 0.85 for the driver and *UAS-Aprt*ᴿᴺᴬⁱ alone controls, respectively, $p <$

0.001), like the *Aprt⁵* mutant (Figure 1D). Next, we used *tub-Gal80*ᵗˢ, which inhibits Gal4 activity

at permissive temperature (McGuire et al. 2003), to prevent Aprt knockdown before the adult

stage. *tub-Gal80*ᵗˢ*; da-Gal4>Aprt*ᴿᴺᴬⁱ flies raised at permissive temperature (18°C) did not show

any locomotor impairment (Figure 1E, *black bars*). However, after being transferred for 3 days

(between 7 and 10 d a.E.) at a restrictive temperature (30°C) that inactivates Gal80, the same flies

demonstrated a similar SING defect as *Aprt⁵* mutants (PI = 0.63 compared to 0.94 and 0.83 for

the driver and RNAi alone controls, respectively, $p < 0.001$) (Figure 1E, *yellow bars with dots*).

This demonstrates that *Aprt* inactivation at the adult stage only is sufficient to alter SING

performance in *Drosophila*, strongly suggesting that this genotype does not result from

developmental defects.

Cell specificity of Aprt requirement for startle-induced locomotion

We then tried to identify the neural cells in which Aprt is required to ensure a normal

locomotor reactivity in young flies by expressing *Aprt*ᴿᴺᴬⁱ with selective Gal4 drivers.

Expression in all neurons with *elav-Gal4* led to decreased locomotor performance in the SING

test (PI = 0.68 at 10 d a.E., vs. 0.90 and 0.86 for the driver and RNAi controls, respectively, *p* <

0.05) (Figure 2A), which was comparable to the effects observed with the $Aprt^5$ mutant or after

ubiquitous expression of the RNAi. To confirm the role of neuronal Aprt in locomotor control,

we generated a *UAS-Aprt* line which allowed for a substantial increase in Aprt expression and

activity (Figure S6). We then found that re-expressing *Drosophila* Aprt selectively in neurons in

$Aprt^5$ background partially but significantly rescued the SING phenotype of the null mutant (PI =

0.70 vs. 0.52 for *driver* and *UAS-Aprt* controls in $Aprt^5$ background, *p* < 0.05) (Figure 2B).

Furthermore, *Aprt* knockdown in all glial cells with *repo-Gal4*, or in sub-populations of

glial cells that express the glutamate transporter Eaat1 with *Eaat1-Gal4*, which includes

astrocyte-like glia, cortex glia and some subperineurial glia (Rival et al. 2004; Mazaud et al.

2019), also led to a lower SING performance of 10-day-old flies (PI = 0.72 for *repo-Gal4* vs.

0.91 for both controls, *p* < 0.05, and PI = 0.56 for *Eaat1-Gal4* vs. 0.77, *p* < 0.05, and 0.88, *p* <

0.01, for the driver and RNAi controls, respectively (Figure 2C, D). In contrast, *MZ0709-Gal4*

(Doherty et al. 2009) and *NP6520-Gal4* (Awasaki et al. 2008) that selectively target the

ensheathing glia, did not induce any significant locomotor defects when used to express the *Aprt*

RNAi (Figure S7). Noticeably, re-expressing Aprt with *Eaat1-Gal4* in the $Aprt^5$ background did

not rescue the SING phenotype (Figure 2E), in contrast to the positive effect of neuronal re-

expression (Figure 2B). Overall this suggests that Aprt is required both in neurons and glia

subsets to ensure a normal SING performance in young flies, and that neuronal but not glial *Aprt*

re-expression is sufficient to ensure a partially functional locomotor behavior.

Figure 2. *Aprt* knockdown in neurons or glial cells disrupts startle-induced locomotion in *Drosophila*. (**A**) *Aprt*RNAi expression in all neurons with *elav-Gal4* decreased SING performance in 10-day-old flies. (**B**) Pan-neuronal expression of *Drosophila Aprt* with the UAS-Gal4 system partially rescued the locomotor response of *Aprt*[5] mutants. (**C-D**) Downregulation of *Aprt* expression in all glia with *repo-Gal4* (**C**) or in glial cell that express the glutamate transporter Eaat1 with *Eaat1-Gal4* (**D**) also altered SING performances. (**E**) *Aprt* re-expression in glial cells with *Eaat1-Gal4* driver did not rescue the climbing response of *Aprt*[5] mutants. Results of 3 or 4 independent experiments performed on 50 flies per genotype at 10 d a. E. *$p < 0.05$, **$p < 0.01$, ns: not significant.

To identify the neuronal subpopulations in which Aprt is required to ensure proper

locomotor responses in young flies, we first tested dopaminergic drivers since we previously

showed that DA neurons play an important role in the control of the SING behavior

(Riemensperger et al. 2011; Riemensperger et al. 2013; Sun et al. 2018). *Aprt* knockdown

targeted to these neurons with *TH-Gal4* did not induce significant impairments (Figure 3A). This

driver expresses in all brain DA neurons except a large part of the protocerebral anterior medial

(PAM) cluster (Friggi-Grelin et al. 2003; Mao and Davis 2009; Pech et al. 2013). In contrast,

downregulating *Aprt* with the *TH-Gal4, R58E02-Gal4* double driver, which labels all DA

neurons including the PAM cluster (Sun et al. 2018), induced a quite similar locomotor defect as

did pan-neuronal *Aprt* knockdown (PI = 0.72 vs. 0.96 and 0.93 for the driver and RNAi controls,

respectively, $p < 0.001$) (Figure 3B). Besides, downregulating *Aprt* in a majority of the

serotoninergic neurons with *TRH-Gal4* (Cassar et al. 2015) did not induce a SING defect (Figure

3C).

These results suggest that some DA neurons in the PAM cluster are specifically involved

in the locomotor impairments induced by *Aprt* deficiency. It has been previously shown in our

laboratory that inhibiting DA synthesis in a subset of 15 PAM DA neurons cluster was able to

alter markedly SING performance in aging flies (Riemensperger et al. 2013). We and others also

reported that the degeneration or loss of PAM DA neurons is involved in the SING defects

observed in several *Drosophila* models of Parkinson disease (Riemensperger et al. 2013; Bou Dib

et al. 2014; Tas et al. 2018; Pütz et al. 2021). Here we found that expressing *Aprt*[RNAi] in the PAM

cluster with *R58E02-Gal4* reproduced the same motor disturbance as pan-neuronal expression (PI

= 0.74 vs. 0.96, $p < 0.001$, and 0.85, $p < 0.05$, for the driver and RNAi controls, respectively)

(Figure 3D), and this result was confirmed by using two other PAM drivers (*NP6510-Gal4* –

expressing only in 15 neurons - *and R76F05-Gal4*) (Figure 3E, F). This strongly suggests that

purine recycling deficiency compromises the correct functioning of these neurons, leading to a

defective startle-induced locomotion.

Because PAM DA neurons innervate the horizontal lobes of the mushroom bodies (Liu et

al., 2012; Riemensperger et al. 2013), and because this structure has been shown to be involved

in the control of startle-induced climbing (Riemensperger et al. 2013; Bou Dib et al. 2014; Sun et

Figure 3. *Aprt* downregulation in DA neurons of the PAM cluster and in mushroom body neurons impairs startle-induced locomotion. (**A**) RNAi-mediated *Aprt* inactivation in brain DA neurons except a large part of the PAM cluster with the *TH-Gal4* driver did not lead to locomotor defects in the SING assay. (**B**) In contrast, *Aprt* knockdown in all dopaminergic neurons including the PAM cluster with the *TH-Gal4, R58E02-Gal4* double-driver led to a decrease in SING performance. (**C**) *Aprt* downregulation in serotoninergic neurons with *TRH-Gal4* did not alter startle-induced climbing response of the flies. (**D-F**) *Aprt* knockdown selectively in DA neurons of the PAM cluster using either *R58E02-Gal4* (**D**), *NP6510-Gal4* (**E**) or *R76F05-Gal4* (**F**) significantly decreased climbing performance. (**G-H**) *Aprt* knockdown in all the mushroom body intrinsic neurons with *238Y-Gal4* (**G**) or *VT30559-Gal4* (**H**) also led to a decrease in SING performance. Results of 3 or 4 independent experiments performed on 50 flies per genotype at 10 d a. E. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ns: not significant.

al. 2018), we also inactivated *Aprt* in MB neurons with *238Y-Gal4* that strongly expresses in all the MB cells (Aso et al. 2009). Interestingly, this driver did induce a locomotor reactivity impairment (PI = 0.70 vs. 0.89 for both controls, $p < 0.01$) (Figure 3G), and the same result was observed with *VT30559-Gal4*, which is a very specific driver for all the MB intrinsic neurons (Plaçais et al. 2017) (Figure 3H). Overall, these results show that normal expression of the SING behavior depends on *Aprt* expression in the PAM and MB neurons in *Drosophila*.

Sleep disturbances induced by Aprt deficiency

Both the mushroom body and subpopulations of PAM DA neurons are known to be regulators of sleep in *Drosophila* (Nall et al. 2016; Artiushin and Sehgal 2017). The fact that *Aprt* deficiency in some of these cells impaired locomotor regulation prompted us to monitor the spontaneous locomotion and the sleep pattern of *Aprt* mutants. Compared to controls, *Aprt*-deficient flies did not have an altered circadian activity profile (Figure S8), nor any difference in total spontaneous locomotor activity during the day, as quantified by the number of infra-red beam cuts (*events*) per 30 min, but they showed a 26.2% increase in total activity during the night ($p < 0.001$) (Figure 4A). As usual, a sleep bout was defined as 5 min or more of fly immobility (Huber et al. 2004). We found that *Aprt*[5] mutants slept significantly less than wild-type flies and

that it was the case both during day and night (Figure 4B, C). These mutants indeed showed a reduced walking speed (Figure 4D) and a smaller average sleep bout duration (Figure 4E), indicating a difficulty to maintain sleep. The reduced speed does not seem to be caused by a decreased energetic metabolism, since we could not detect different ATP levels in head and thorax of $Aprt^5$ mutants compared to wild-type flies (Figure S9).

Figure 4. *Aprt*-deficient flies sleep less and walk slower than wild-type flies. (**A**) Quantification of total spontaneous locomotor activity during day and night over 5 LD cycles. *Aprt*[5] mutants show no difference in spontaneous locomotion with wild-type flies during the day but a higher activity at night. 3 independent experiments were performed on 32 flies per genotype and mean ± SEM was plotted. ***$p < 0.001$, ns: not significant. (**B**) Sleep pattern during a typical 24h LD cycle showing that the total amount of sleep is smaller during day and night in *Aprt*[5] mutants compared to wild-type flies. (**C**) Quantification of day (ZT1-12), night (ZT13-24) and total sleep in *Aprt*[5] mutants. ZT is for zeitgeber. (**D**) Locomotion speed during waking is reduced in *Aprt*[5] mutants. (**E**) The average sleep bout duration is also decreased, indicating that *Aprt*[5] mutants have a difficulty to maintain sleep. (**F**) Sleep pattern of *elav>Aprt*[RNAi] flies, showing that knockdown of Aprt in all neurons led to sleep reduction during the night, similarly to the mutant condition, and an even more profound sleep defect during the day. (**G**) Quantification of total amount of sleep when *Aprt* was downregulated in all neurons (*elav-Gal4*), all glial cells (*repo-Gal4*) and both neurons and glial cells (*elav-Gal4*; *repo-Gal4*). Except for glia only, the resulting effect was a significant sleep reduction. For sleep and locomotion speed, means ± SEM were plotted. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ns: not significant.

Interestingly, RNAi-mediated downregulation of Aprt selectively in neurons (*elav>Aprt*[RNAi] flies) also significantly decreased sleep duration during day and night (Figure 4F), whereas glial-only expression (*repo>Aprt*[RNAi]) had no effect (Figure S10A). Expressing the RNAi in both neurons and glia (*elav; repo>Aprt*[RNAi]) had similar effects as in neurons alone (Figure S10B). Quantification of total sleep in these experiments, and the total amount of day and night sleep, are shown in Figure 4G and Figure S10C, D, respectively. Overall, these results suggest that *Aprt* expression is selectively needed in neurons for a normal sleep pattern in *Drosophila*.

Brain DA synthesis and levels are increased in *Aprt*-deficient flies

Since we found that induced locomotion and sleep, two behaviors controlled by DA neurons in *Drosophila*, were altered in *Aprt*-deficient flies, we logically expected brain DA levels to be reduced in these mutants, as is the case in the basal ganglia of LND patients. To check for that, we performed comparative immunostaining for tyrosine hydroxylase (TH), the specific enzyme of DA synthesis (Friggi-Grelin et al. 2003; Riemensperger et al. 2011), on dissected

adult brains from wild-type and $Aprt^5$ mutant flies. Surprisingly, the global TH protein level

appeared not to be decreased, but relatively increased by 17.5% in the $Aprt^5$ mutant brain ($p <$

0.01), in particular around the mushroom bodies, a structure that receives dense dopaminergic

projections (Figure 5A, B). Moreover, DA immunostaining carried out on whole-mount dissected

brains revealed a similarly increased level of this neuromodulator in $Aprt^5$ flies, by 17.0% on

average in the entire brain ($p < 0.01$), but not specifically in the mushroom bodies or another part

of the brain (Figure 5C, D). This indicates that disruption of the purine salvage pathway does not

impede DA synthesis and levels in the *Drosophila* brain, which are instead slightly increased,

perhaps as a compensatory mechanism. The behavior defects suggest that the lack of Aprt

activity could interfere with correct functioning of the DA system in an indirect way. We

therefore searched for another neurotransmitter system that could be affected by *Aprt* deficiency.

Figure 5. *Aprt* deficiency increases DA synthesis and content in the *Drosophila* brain. (**A**) Representative confocal projections of TH-immunostained whole-mount adult brains from wild-type flies and *Aprt⁵* mutants. MB: mushroom body. Scale bars: 100 μm. (**B**) Quantification of TH immunofluorescence intensity normalized to the controls in the entire brain. 4 to 6 brains were dissected per experiment and genotype, and 6 independent experiments were performed. **$p < 0.01$. (**C**) Representative confocal projections of DA immunostaining in whole-mount adult brains of wild type and *Aprt⁵* mutants. Scale bars: 100 μm. (**D**) Quantification of DA immunofluorescence intensity over the entire brain showed a slight increase in DA content in *Aprt⁵* mutants compared to wild-type controls. 6 brains were dissected per experiment and genotype, and 3 independent experiments were performed. **$p < 0.01$.

Interactions between Aprt and the adenosinergic system

Aprt catalyzes the conversion of adenine into AMP, and AMP breakdown by the enzyme 5'-nucleotidase produces adenosine, primarily in the extracellular space. Adenosine then acts as a widespread neuromodulator in the nervous system by binding to adenosine receptors. We therefore suspected that adenosine level could be reduced in the absence of Aprt activity, leading to alterations in adenosinergic neurotransmission. Indeed, we found a significant decrease in adenosine level either in whole flies (by 61.0% on average, $p < 0.01$) or in heads (by 48.0%, $p < 0.05$) in $Aprt^5$ mutants (Figure 6A). We then examined the consequences of this reduction on molecular components of the adenosinergic system, namely G protein-coupled adenosine receptors and equilibrative nucleoside transporters (ENTs), which carry out nucleobase and nucleoside uptake of into cells. Only one seven-transmembrane-domain adenosine receptor, AdoR, is present in *Drosophila*, that is very similar to mammalian adenosine receptors (Dolezelova et al. 2007; Brody and Cravchik, 2000), and three putative equilibrative nucleoside transporters (Ent1-3) have been identified (Sankar et al. 2002) but only one, Ent2, showed nucleoside transport activity (Machado et al. 2007). Interestingly, Ent2 has been shown to be enriched in the mushroom bodies and its transcript level to be highly elevated in *AdoR* mutant flies (Knight et al. 2010). In $Aprt^5$ background, we also observed a strong increase in *Ent2* mRNAs (2.3-fold higher than wild-type flies, $p < 0.001$), but no noticeable effect on *AdoR* transcript level (Figure 6B). The increased expression of *Ent2* could correspond to a compensatory response to adenosine shortage in $Aprt^5$ mutants.

Figure 6. Relations between Aprt and molecular components of adenosinergic signaling. (**A-B**) Impacts of the lack of Aprt activity on the adenosinergic system. (**A**) Adenosine level was measured in whole flies or heads of $Aprt^5$ flies by UPLC. Compared to wild-type flies, adenosine level was significantly reduced in the mutants. Results of 3 independent experiments performed with 5 males per genotype in triplicates, and 2 independent experiments with 30 heads per genotype in triplicates. $*p < 0.05$, $**p < 0.01$. (**B**) $Aprt^5$ mutation did not affect $AdoR$ expression but induced a marked increase in adenosine transporter $Ent2$ mRNA abundance. 3-6 different RNA extractions were performed on 20-30 male heads. Results of 3 to 6 independent experiments. $***p < 0.001$, ns: not significant. (**C**) Null $AdoR^{KGex}$ mutants showed decreased $Aprt$ expression (*left panel*) and a stronger decrease in Aprt activity (*right panel*). 4 independent RNA extractions were carried out on 20-30 male heads and 4 independent real-time PCR determinations were done per RNA sample. For Aprt activity, 3 independent determinations were performed on 20 whole flies per genotype. $***p < 0.001$.

The AdoR receptor is highly expressed in adult fly heads and its ectopic overexpression leads to early larval lethality (Dolezelova et al. 2007). In contrast, a null mutant of this receptor, $AdoR^{KGex}$, in which the entire coding sequence is deleted, is fully viable (Wu et al. 2009). This enabled us to examine the consequences of a complete lack of AdoR on purine recycling in adult

flies. We found that *Aprt* transcripts were decreased by 29.5% on average ($p < 0.001$) in

AdoR[KGex] mutant heads (Figure 6C, *left panel*), while Aprt activity was even more decreased by

78.4% ($p < 0.001$) compared to wild-type flies (Figure 6C, *right panel*). This effect likely results

from the much increased level of adenosine uptake in the *AdoR* mutants, which would

downregulate *Aprt* activity by a feedback mechanism. Overall, these data indicate that

extracellular adenosine levels must be strongly decreased in Aprt mutant flies, both from the

general reduction in adenosine levels and the increased expression of the Ent2 transporter.

Although AdoR expression is not affected, AdoR signaling is therefore expected to be

significantly reduced in *Aprt* mutants.

***Aprt* mutants show epilepsy-like seizure behavior**

A number of *Drosophila* mutants with disrupted metabolism or neural function show

increased susceptibility to seizure and paralysis following strong mechanical or

electrophysiological stimulation (Fergestad et al. 2006, Parker et al. 2011, Kroll et al. 2015).

These mutants are called bang-sensitive (BS) and commonly used as models of epileptic seizure

(Song and Tanouye 2008). Here we checked whether Aprt deficiency could trigger a BS

phenotype. We observed that aged *Aprt* mutants (at 30 d a.E.) recovered slowly after a strong

mechanical stimulation applied by vortexing the vial for 10 s at high speed. These flies took on

average 17.3 s to recover and get back on their legs, compared to 2.5 s for wild-type flies of the

same age ($p < 0.01$) (Figure 7A and Movie S1). Some of the mutant flies appeared more deeply

paralyzed as they did not spontaneously recover unless the vial was stirred, so their recovery time

could not be scored. Hemizygous flies of the same age containing the *Aprt*[5] mutation over two

partially overlapping genomic deficiencies covering *Aprt* also showed a BS behavior, with an

average longer recovery time of 28.9 s and 33.2 s, respectively ($p < 0.001$) (Figure 7B).

We then downregulated *Aprt* by RNAi in all cells with the *da-Gal4* driver to check if this

could also induce BS behavior. As shown in Figure 7C, *da>Aprt*[RNAi] flies at 30 d a.E. indeed

displayed seizure after mechanical shock, quite similar to that of the *Aprt*[5] mutants (22.7 s

recovery time on average compared to 1.55 s and 0.72 s for the driver and RNAi controls,

respectively, $p < 0.05$). In contrast, inactivating *Aprt* selectively in neurons *(elav-Gal4)*, glial

cells *(repo-Gal4)* or muscles *(24B-Gal4)* did not induce a BS phenotype (Figure S11). This

suggests that the BS phenotype requires *Aprt* knockdown in other cells or in several of these cell

types. Finally, in contrast to the SING defect, we observed that adult-specific *Aprt* knockdown in

all cells with *da-Gal4* for 3 days did not trigger a BS behavior (Figure 7D), indicating that the

bang sensitivity requires a longer downregulation of the gene or might be the consequence of a

developmental defect.

Administration of Adenosine or *N*[6]-methyladenosine to *Aprt*-deficient flies prevents seizure

Drosophila disease models are advantageously tractable for drug screening *in vivo*

(Fernández-Hernández et al. 2016, Perrimon et al. 2016). We thus administered several

compounds related to purine metabolism to *Aprt*[5] flies to check if they can rescue

neurobehavioral impairments (locomotor defects and seizure). Feeding allopurinol at the same

concentration used for uric acid normalization (100 µg/ml, Figure 1B), either in adults 5 days

before the test or throughout all developmental stages, did not alter the BS phenotype (Figure

S12), as was the case for the SING assay (Figure S5). Similarly in humans, it has been shown

that the daily intake of allopurinol, even from infancy, does not mitigate the neurobehavioral

impairments in LND patients (Marks et al. 1968; Torres et al. 2007; Jinnah et al. 2010; Madeo et

al. 2019).

Figure 7. *Aprt* deficiency triggers a seizure-like phenotype. (**A**) At 30 d a.E., *Aprt⁵* mutants need a much longer time than wild-type flies to recover from a strong mechanical shock, showing a bang-sensitive (BS) paralysis comparable to tonic-clonic seizure. Results of 3 independent experiments performed on 50 flies per genotype. **$p < 0.01$. (**B**) At 30 d a. E., hemizygous *Aprt⁵* mutants also showed a marked BS phenotype. Results of 2-4 independent experiments performed on 50 flies per genotype. ***$p < 0.001$. (**C**) RNAi-mediated downregulation of *Aprt* in all cells (*da>Aprt*^RNAi) also led to BS phenotype in adults at 30 d a.E., but not with the driver and effector controls. Results of 3 independent experiments performed on 50 flies per genotype. *$p < 0.05$. (**D**) *Aprt* knockdown by RNAi during the adult stage for 3 days before the test was not sufficient to induce bang-sensitivity, suggesting that this phenotype could be caused by a developmental defect or a longer downregulation of *Aprt*. Results of 2 independent experiments performed on 50

483 flies per genotype. ns: not significant. (**E, F**) The BS phenotype of 30-day-old $Aprt^5$ mutants was
484 rescued by feeding either 500 μM adenosine (ado) (**E**) or 500 μM N^6-methyladenosine (m6A) (**F**)
485 during all developmental stages plus 5 days before the test. Results of 4 or 6 independent
486 experiments performed on 50 flies per genotype. ***$p < 0.001$, ns: not significant.
487

Then, we tried to supplement *Aprt* mutants with various purine compounds, including

adenine, hypoxanthine, adenosine and N^6-methyladenosine (m6A), at 100 or 500 μM, either in

adult stage 5 days before the test or throughout larval development plus 5 days before the test.

None of these drugs was able to rescue the SING defect (data not shown). In contrast and

interestingly, administration of 500 μM adenosine or m6A during development rescued the BS

phenotype of $Aprt^5$ mutants (Figure 7E-F). This further indicates that different mechanisms

underpin SING alteration and BS behavior in *Aprt* mutants and provide another evidence that the

bang sensitivity may be caused by a developmental defect. The results also suggest that the lower

adenosine levels of *Aprt* mutant flies could be at the origin of their bang sensitivity.

Neuronal expression of mutant HGPRT induces early locomotor defects and seizure

behavior

In concordance with reports from about 50 years ago (Miller and Collins 1973; Becker

1974a; 1974b), we could confirm that no HGPRT enzymatic activity can be detected in wild-type

Drosophila extracts (Table S2). Furthermore, searching the *Drosophila* genome with BLAST did

not retrieve any *Drosophila* homologue of *HPRT1*, the human gene encoding HGPRT. In order to

potentially develop another *Drosophila* model mimicking LND conditions, we generated new

transgenic UAS lines to express in living flies either the human wild-type HGPRT (HGPRT-WT)

or a pathogenic LND-associated mutant form of this protein (HGPRT I42T), both isoforms being

inserted at the same genomic docking site. These lines were validated by showing that they are

transcribed at similar levels (Figure 8A, B). Enzymatic assays on adult extracts of flies

expressing the wild-type form HGPRT-WT in all cells showed significant HGPRT enzyme

activity, while an 80.5% lower activity was detected in *Drosophila* expressing the mutant form

HGPRT-I42T (Table S2).

Figure 8. Expression of a pathogenic mutant isoform of human HGPRT induces neurobehavioral
defects in flies. (**A-B**) Ubiquitous expression of human *HPRT1* with *da-Gal4*. (**A**) Amplification
of human *HPRT1* transcripts detected by RT-PCR in head extracts of *da>HPRT1-WT* and
da>HPRT1-I42T flies. A band with lower intensity was also detected in the effector controls
(*UAS-HPRT1-WT/+ and UAS-HPRT1-I42T/+*), and not in the driver control (*da/+*), which
indicates a small amount of driver-independent transgene expression. (**B**) Quantification of the
previous experiment. (**C, D**) Expression of the LND-associated I42T isoform in all neurons
(*elav>HPRT1-I42T*), but not of the wild-type form (*elav>HPRT1-WT*), induced an early
locomotor SING defect at 15 d a.E. (**C**) and a prominent bang-sensitivity phenotype at 30 d a.E.
(**D**), as compared to the driver (*elav/+*) and effector (*UAS-HPRT1 I42T/+*) only controls. Results
of 3 independent experiments performed on 50 flies per genotype. **$p < 0.01$, ***$p < 0.001$, ns:
not significant.

We next analyzed the consequences of human HGPRT expression on the SING and bang

sensitivity behaviors. Interestingly, the pan-neuronal expression of mutant I42T isoform

specifically induced a significant early locomotor defect at 15 d a.E. (PI = 0.71 vs. 0.92 and 0.90

for the driver and effector controls, respectively, $p < 0.01$) (Figure 8C) and a relatively small but

robust bang-sensitivity behavior at 30 d a.E. (2.3 s average recovery time vs. 0.64 s and 0.31 s for

the driver and effector only controls, $p < 0.001$) (Figure 8D). These defective phenotypes could

not be seen when HGPRT-WT was expressed. Therefore, and remarkably, whereas wild-type

HGPRT expression appears to be innocuous in *Drosophila*, we observed that the neuronal

expression of a pathogenic LND-associated isoform triggered neurobehavioral impairments quite

similar to those of *Aprt*-deficient flies.

Discussion

Over the past 35 years or so, several animal models of LND have been developed in

rodents based on HGPRT mutation in order to better understand the cause of the disease and test

potential therapeutic treatments (Finger et al. 1988; Dunnett et al. 1989; Jinnah et al. 1993; Engle

et al 1996; Meek et al. 2016; Witteveen et al. 2022). However, none of these models recapitulated

the full LND syndrome and, particularly, motor or neurobehavioral symptoms as a consequence

of HGPRT deficiency. To date, the causes of the neurobehavioral impairments in LND are not

yet clearly elucidated and the disease is still incurable (Fu et al. 2014; Bell et al. 2016; López et

al., 2020; Bell et al., 2021; Witteveen et al. 2022). Here we have tried to develop new models of

this disease in *Drosophila*, a useful organism to conduct genetic and pharmacological studies. An

apparent drawback of this project was that no homolog of the mammalian gene encoding HGPRT

is conserved in the fly genome, which may explain why no LND models have been developed to

date in this organism. To try to overcome that, we used two different strategies. First, because

Aprt seems to be the only enzyme of the purine salvage pathway in *Drosophila*, we hypothesized

that it could have analogous functions, both for purine homeostasis and behavior control, as

HGPRT in humans. Indeed, we show here that *Aprt* deficiency induces both metabolic and

neurobehavioral disturbances in *Drosophila*, similar to the loss of HGPRT, but not APRT,

activity in humans. Second, we have expressed the wild-type human HGPRT and a LND-

associated mutant form of this protein in *Drosophila* neurons to analyze their respective effects in

living flies. Here we provide evidence that the fruit fly can be used to study the consequences of

defective purine recycling pathway and HGPRT mutation in the nervous system, and perform

pharmacological screens to find new therapy for LND.

***Aprt* deficient flies replicate lifespan and metabolic defects caused by HGPRT deficiency**

Flies that carry a homozygous null-mutation in *Aprt* develop normally and live until the

adult stage (Johnson and Friedman, 1983). Furthermore, a previous study reported that

heterozygous *Aprt*/+ flies have an extended lifespan (Stenesen et al. 2013). We observed that

homozygous $Aprt^5$ mutants have in contrast a significantly reduced longevity compared to wild-

type flies. It can be noted that the lack of HGPRT activity in LND also causes reduced lifespan

expectancy, which is generally less than 40 years of age for properly cared patients. Stenesen et

al. (2013) have demonstrated that dietary supplementation with adenine, the Aprt substrate,

prevented the longevity extension conferred either by dietary restriction or heterozygous

mutations of AMP biosynthetic enzymes. This suggests that longevity depends on an accurate

regulation of adenine levels. It is possible that adenine could accumulate to toxic levels during

aging in homozygous *Aprt* mutants, explaining their shorter lifespan. Alternatively, we could

speculate that AMP, the Aprt product, will be decreased in the absence of a functional Aprt, and

as a consequence, AMP-activated protein kinase (AMPK), an enzyme that protects cells from

stresses inducing ATP depletion, would be less activated. Multiple publications explored the role

of AMPK in lifespan regulation (Sinnett and Brenman 2016) and it has been demonstrated that

downregulating AMPK by RNAi in *Drosophila* adult fat body or muscles reduces lifespan

(Stenesen et al. 2013), as well as its ubiquitous inactivation under starvation (Johnson et al.

2010).

In humans, HGPRT deficiency induces an accumulation of hypoxanthine and guanine,

resulting from the lack of recycling and an increased *de novo* purine synthesis (Harkness et al.

1988, Fu et al. 2015; Ceballos-Picot et al. 2015) leading to an overproduction of uric acid which

increases the risk for nephrolithiasis, renal failure and gouty arthritis if not properly treated. In

insects, the end-product of purine catabolism is not uric acid, as in humans, but allantoin.

However, urate oxidase, the enzyme which converts uric acid into allantoin, is expressed

selectively in the Malpighi tubules, an excretory organ that forms the pre-urine in flies (Wallrath

et al. 1990). Hence, an accumulation of uric acid could occur in the hemolymph if the purine

salvage pathway is impaired. Here, we have indeed observed an increase in uric acid level in

Drosophila Aprt mutant heads. Interestingly, we were able to reduce uric acid to normal levels in

Aprt mutant flies by feeding them allopurinol, a xanthine oxidase inhibitor that is used to

decrease uric acid levels and protect against renal failure in LND patients.

Aprt is required in dopamine and mushroom body neurons for young fly motricity

We observed that *Aprt*-null adult flies have a strikingly reduced performance in the SING

test, which monitors locomotor reactivity to startle and climbing ability. This phenotype was

already obvious at an early age, starting from 1 d a.E. The performance continued to decrease

until 8 d a.E. and then appeared to stabilize up to 30 d a.E. Remarkably, this phenotype is clearly

different from the defects usually found in α-synuclein-induced *Drosophila* Parkinson disease

models, in which the locomotion starts declining at around 25 d a.E. and then decreases steadily

with aging (Feany and Bender 2000, Riemensperger et al. 2013, Rahmani et al. 2022). Such a

locomotor defect affecting newly eclosed *Aprt*-deficient flies could be reminiscent of the early

onset of motor symptoms in LND patients, which most often appear from 3 to 6 months of age

(Jinnah et al. 2006). Interestingly, we were also able to induce an impairment in startle-induced

locomotion by knocking down *Aprt* during 3 days only in adult flies, which suggests that the

motricity phenotype does not necessarily require a developmental flaw. Although *Aprt* mutants

walked slower than wild-type flies, they were no less active and their ATP levels were not

different than in controls, excluding that reduced SING performance resulted from a major failure

in energetic metabolism.

Since the complete absence of *Aprt* induced a strong and early SING defect, we tried to

identify in which cells *Aprt* is required to ensure a normal locomotor behavior. Downregulating

Aprt in all neurons by RNAi reproduced the locomotor defect of the *Aprt*[5] mutant, and *Aprt*

mutant locomotion could be partially rescued by neuronal Aprt re-expression. The fact that

rescue was not complete was possibly due to a dominant negative effect of the *Aprt*[5] mutation on

Aprt activity, as suggested by enzymatic assays on extracts of heterozygous mutant flies (Figure

S2). We then considered that DA neurons could be involved in this motor phenotype. Previous

studies from our and other laboratories have indeed demonstrated the role of DA neurons in the

control of SING in *Drosophila* (Feany and Bender 2000; Friggi-Grelin et al. 2003;

Riemensperger et al. 2011; Vaccaro et al. 2017; Sun et al. 2018), and that the degeneration or loss

of PAM DA neurons underpin the SING defects observed in different *Drosophila* Parkinson

disease models (Riemensperger et al. 2013; Bou Dib et al. 2014; Tas et al. 2018; Pütz et al.

2021). Here, we showed that knocking down *Aprt* in all DA neurons or only in the PAM DA

neurons was sufficient to induce early SING defects. Moreover, inactivation of Aprt with *TH-*

Gal4 that labels all DA neurons except for a large part of the PAM cluster did not affect the

SING, suggesting that Aprt expression in subsets of PAM neurons is critical for this locomotor

behavior.

Here we found that inactivating *Aprt* in all MB neurons also induced a lower performance

in the SING assay. This important brain structure receives connections from DA neurons,

including the PAM, and is enriched in DA receptors (Waddell 2010). We have recently shown

that activation of MB-afferent DA neurons decreased the SING response, an effect that requires

signalization by the DA receptor dDA1/Dop1R1 in MB neurons (Sun et al. 2018). We also

observed a SING defect after knocking down *Aprt* either in all glial cells, or more selectively in

glial subpopulations expressing the glutamate transporter Eaat1, but not in the ensheathing glia.

We reported before that some astrocyte-like glial cells that express *Eaat1* extend processes

forming a thick mesh-like network around and inside the entire MB neuropil (Sinakevitch et al.

2010). So it could be hypothesized that these MB-associated astrocytes also require Aprt

expression to ensure a full SING behavior. However, re-expressing *Aprt* with *Eaat1-Gal4* did not

lead to SING rescue in Aprt mutant background, in contrast to the positive effect observed with

elav-Gal4. This suggests that the presence of Aprt in neurons can somehow compensate for *Aprt*

deficiency in glia, but the reverse is not true. In conclusion, proper startle-induced locomotion in

young flies depends on Aprt activity in PAM DA and MB neurons, and in *EAAT1*-expressing

astrocytes.

Neuronal Aprt regulates spontaneous activity and sleep

Because sleep is regulated by the mushroom body as well as DA neurons in flies,

(Artiushin and Sehgal 2017), we have also monitored the spontaneous locomotor activity and

sleep pattern of *Aprt^5* mutants. Their activity profile appeared normal, with unaltered morning

and evening anticipation, indicating that the circadian rhythms are maintained in light-dark

conditions in the absence of a functional purine recycling pathway (Figure S8). This experiment

also highlighted that *Aprt*-deficient flies are hyperactive during the night, which suggests that

they sleep less than wild-type flies. This could be confirmed by measuring their sleep pattern on

video recordings. *Aprt* mutants show a reduced walking speed, sleep less during both day and

night, and have a difficulty to maintain sleep. Downregulating *Aprt* selectively in neurons

reproduced the sleep defect, whereas doing it in glial cells only had no effect. We have not

attempted here to identify further the neuronal cells involved in this phenotype. Like for the

SING behavior defect, it could involve the PAM DA neurons, as subpopulations of this cluster

have been shown to regulate sleep in *Drosophila* (Nall et al. 2016). Therefore, the lack of purine

recycling markedly disrupts sleep in *Aprt*-deficient flies, making them more active at night. This

is strikingly comparable to young LND patients who have a much disturbed sleep-time during the

night (Mizuno et al. 1979).

Lack of Aprt reduces adenosine signaling leading to DA neuron overactivation

Here we found that disruption of the purine salvage pathway induced by *Aprt* deficiency

in *Drosophila* did not decrease the level of DA in brain, although the mutant flies showed

reduction in induced locomotion and sleep. This prompted us to study another neuromodulator,

adenosine, whose level was likely to be altered in *Aprt* mutants, since this nucleoside is indirectly

a product of Aprt enzymatic activity. The purine nucleoside adenosine is one of the building

blocks of RNA and the precursor of essential molecules such as ATP and cAMP, but it is also the

endogenous ligand of adenosine receptors that modulate a wide range of physiological functions.

In the brain, adenosine regulates many motor and cognitive processes, such as the sleep-wake

cycle, anxiety, depression, epilepsy, memory and drug addiction (Soliman et al. 2018). In normal

metabolic conditions, adenosine is present at low concentrations in the extracellular space and its

level is highly regulated, either taken up by cells and incorporated into ATP stores or deaminated

by adenosine deaminase into inosine. In mammals, several nucleoside transporters mediate the

uptake of adenosine and other nucleosides into cells, named equilibrative and concentrative

nucleoside transporters (ENTs and CNTs), respectively (Gray et al. 2004; Boswell-Casteel and

Hays 2017; Pastor-Anglada and Pérez-Torras 2018).

As expected, we observed a marked reduction in adenosine levels in *Aprt* mutant flies.

While we did not observe any alteration in the transcript levels of *AdoR*, the gene coding for the

only adenosine receptor in flies, transcript levels of the known adenosine transporter, Ent2, were

increased more than 2-fold in *Aprt* mutants. Interestingly, one paper reported a similar strong

increase in *Ent2* transporter expression in *AdoR* mutant flies (Knight et al. 2010). This suggests

that AdoR signaling is less activated in Aprt-deficient flies compared to controls, and the

decrease in adenosine we observed in *Aprt* mutants also supports this hypothesis. The fact that

AdoR expression was not affected in these conditions suggests that a role of this receptor is to

inform the cells about the external adenosine concentration and accordingly regulate Ent2

expression level. We also observed that the lack of AdoR decreased Aprt expression and activity

in *Drosophila*, probably as a consequence of the increase in Ent2 expression, which would lead

to higher adenosine influx and then Aprt downregulation by a feedback mechanism.

Adenosine and dopamine receptors are known to interact closely in mammals (Franco et

al. 2000; Kim and Palmiter 2008), whereas the function of AdoR has not been extensively

studied in *Drosophila*, in particular with regard to its potential interactions with the DA system.

However, a previous study performed in fly larvae showed that an increase in astrocytic Ca^{2+}

signaling can silence DA neuron activity through AdoR stimulation on DA neurons, by a

mechanism potentially involving the breakdown of released astrocytic ATP into adenosine (Ma et

al. 2016). Conversely, the AdoR antagonist caffeine was shown to increase DA neuron signaling

in the PAM cluster, although it is not certain if caffeine acts through AdoR or another receptor to

activate these neurons (Nall et al. 2016). This suggests that a reduction in AdoR signaling, as

seems to be the case in Aprt-deficient flies, will lead to DA neuron overactivation. We have

previously shown that thermogenetic or optogenetic activation of the DA neurons strongly

reduces fly performance in the SING test (Sun et al. 2018). As described here, this is precisely

the locomotor phenotype of young Aprt-deficient flies. Furthermore, it has been shown in

Drosophila that nocturnal hyperactivity is caused by an increase in dopamine signaling (Kumar

et al. 2012; Lee et al. 2013). This is fully in accordance with our present observation that Aprt-

deficient flies sleep less and are more active during the night. Therefore, both the locomotor and

sleep defects induced by the lack of Aprt activity can be explained by an overactivation of DA

neurons that would be a direct consequence of reduced adenosinergic signaling.

At first view, these observations do not seem to be in favor of the idea that the Aprt

deficient fly could represent a useful model to study LND, since a constant feature of patients

with this disease is a marked reduction in DA level in the basal ganglia. However, it has been

previously proposed that adenosine could be a mediator of some of the neurological

consequences of LND pathogenesis (Visser et al., 2000). Indeed, some studies showed that

adenosine transport is decreased in peripheral blood lymphocytes of LND patients (Torres et al.

2004), as well as A_{2A} adenosine receptor mRNA and protein levels (García et al. 2009, García et

al. 2011). In a murine HPRT knockout model of LND, expression of the adenosine A_1 receptor

was found to be strongly increased and that of A_{2A} slightly decreased, while the expression of

A_{2B} was not affected (Bertelli et al. 2006). Chronic administration of high doses of caffeine,

which also acts as an antagonist of adenosine receptors in the mammalian brain, caused a self-

injurious behavior in rats (Miñana et al. 1984; Jinnah et al. 1990). Moreover, central injection of

an adenosine agonist is sufficient to prevent self-mutilation induced by dopamine agonist

administration in neonatally 6-OHDA lesioned rats (Criswell et al. 1988). These striking

observations encouraged us to study further the potential involvement of adenosine signaling in

the neurobehavioral effects induced by *Aprt* deficiency in *Drosophila*.

Adenosine or N6-methyladenosine supplementation rescues the epilepsy behavior of *Aprt*

mutants

LND is characterized by several behavioral troubles including dystonia, spasticity and

choreoathetosis. This mainly corresponds to uncontrolled, sustained or repetitive muscular

contractions, and stiffness or involuntary twitching and writhing that can affect the posture,

walking ability and normal movements. Some patients can also show an epileptic disorder

(Jinnah et al. 2006; Madeo et al. 2019). In flies, the bang-sensitivity test is often used to model

epileptic seizures (Song et al. 2008; Parker et al. 2011). Here, we have found that 30-day-old *Aprt*

mutant flies show a typical BS phenotype after a strong mechanical shock. Previous works have

shown that bang sensitivity is often linked to neuronal dysfunction in flies (Parker et al. 2011;

Kroll and Tanouye 2013; Kroll et al. 2015; Saras and Tanouye 2016). The fact that we could not

reproduce the BS phenotype by knocking down *Aprt* in specific cells such as neurons, glia, or

muscles, indicates that Aprt has to be inactivated in several or all cell types to induce this defect.

This may suggest that the crucial component missing in the absence of Aprt activity is diffusible

and can be provided to neurons from the extracellular space or the hemolymph.

Interestingly, inactivating *Aprt* by RNAi in all cells during development reproduced the

same seizure-like behavior, whereas inactivation of *Aprt* for 3 days only in adult flies failed to

induce bang sensitivity, in contrast to the SING phenotype. These results indicate that the

absence of Aprt during development somehow alters a mechanism that controls bang sensitivity,

but only has a repercussion with age. The role of Aprt being to recycle adenine into AMP, it was

conceivable that *Drosophila Aprt* mutants lack nucleotides in the brain during development,

possibly inducing neuronal defects that could trigger later the BS behavior. We have tried,

therefore, to feed the mutants with a diet supplemented with various compounds involved in

purine metabolism, such as allopurinol, adenine, hypoxanthine, adenosine or its analog N6-

methyladenosine (m^6A), either throughout larval development or only in adult stage. Only

adenosine and m^6A, ingested during development, were able to prevent the BS behavior. This

suggests that the absence of Aprt induces a lack of adenosine in the developing nervous system,

as we demonstrated, which has a profound impact on the formation or function of neural circuits

controlling the BS behavior in adults. Adenosine is methylated at the nitrogen-6 position to form

m^6A in RNA molecules by a large methyltransferase complex (Batista 2017). Once released

during RNA turnover or degradation, m^6A cannot be incorporated back into nucleic acids and so

it is excreted in urine (Schram 1998). The rescuing effect we observed with m^6A on the BS

phenotype indicates thereby that both this compound and adenosine rather act as adenosine

receptor agonists or allosteric regulators, than as nucleotide precursors, to prevent the nervous

system disturbance induced by Aprt deficiency.

Adenosine has been shown to strongly inhibit brain electrical activity, and its role as an

endogenous anticonvulsant and seizure terminator is well-established in humans (Boison 2005;

Masino et al. 2014; Weltha et al. 2019). Accordingly, deficiencies in the adenosine-based

neuromodulatory system can contribute to epileptogenesis. For instance, increased expression of

astroglial adenosine kinase (ADK), which converts adenosine into AMP, leads to a reduction in

brain adenosine level that plays a major role in epileptogenesis (Weltha et al. 2019). Hence,

therapeutic adenosine increase is a rational approach for seizure control. One explanation for how

adenosine release is responsible for seizure arrest implies the attenuation of the GABA$_A$ receptor-

mediated depolarization (Ilie et al. 2012). Overall, such data could help explain why our

observation that feeding adenosine or its derivative m^{6}A rescued the seizure-like phenotype of

Aprt mutant flies, and further suggests that adenosinergic signaling has similar functions in the

fly and mammalian brains. In addition, we did observe that adenosine level was decreased in *Aprt*

mutant flies, perhaps because of an elevated ADK activity to compensate the absence of AMP

production by Aprt. Although the reason why adenosine administration only in adult stage could

not rescue the BS behavior remains to be elucidated, it is consistent with the fact that

downregulating Aprt for three days in aged adult flies was not sufficient to give rise to the BS

phenotype.

We and others recently identified m^{6}A, and other compounds sharing an adenosine

moiety, as able to rescue the viability of LND fibroblasts and neural stem cells derived from

induced pluripotent stem cells (iPSCs) of LND patients, when cultured in the presence of

azaserine, a potent blocker of *de novo* purine synthesis (Petitgas and Ceballos-Picot, unpublished

results; Ruillier et al. 2020). Like for flies again, allopurinol was not capable of rescuing the cell

viability in this *in vitro* model. The similarity of these results provides further evidence

suggesting that the *Aprt*-deficient *Drosophila* could be used as a model to study LND.

Expression of mutant HGPRT triggers locomotor and seizure phenotypes similar to *Aprt*

deficiency

Why HGPRT deficit causes such dramatic neurobehavioral troubles in LND patients still

remains a crucial question. To date, cellular (Smith & Friedman 2000; Torres et al. 2004;

Ceballos-Picot et al. 2009; Cristini et al. 2010; Guibinga et al. 2012; Sutcliffe et al. 2021) and

rodent (Finger et al. 1988; Dunnett et al. 1989; Jinnah et al. 1993; Meek et al. 2016; Witteveen et

al. 2022) models only focused on the consequences of HGPRT deficiency to phenocopy the

disease. Such an approach was justified by the fact that the lower the residual activity of mutant

HGPRT, the more severe the symptoms are in patients (Fu and Jinnah 2012. Fu et al. 2014).

However, it would be conceivable that part of the symptoms observed in the disease result from

compensatory physiological mechanisms or a deleterious gain-of-function conferred to the

HGPRT protein by the pathogenic mutations. Here we observed that neuronal expression of

mutant human HGPRT-I42T, which expresses a low enzymatic activity, but not the wild-type

protein, induced early locomotor defects in young flies and bang-sensitivity in older flies, very

similarly to the defects induced by Aprt deficiency. This suggests a potential neurotoxicity of the

pathological mutant form of HGPRT, which could be related to disturbances in purine

metabolism or other signaling pathways. The human mutant form might not be properly degraded

and accumulate as aggregates, potentially exerting neuronal toxicity. Such an approach opens

interesting perspectives to better understand the endogenous function of HGPRT and its

pathogenic forms. Indeed, the identification of a potential inherent neurotoxicity of defective

forms of human HGPRT is a new element, which could be explored in further work in the fly and

also in rodent models.

A new model of LND in an invertebrate organism

LND, a rare X-linked metabolic disorder due to mutations of the *HPRT1* gene, has

dramatic neurological consequences for affected children. To date, no treatment is available to

abrogate these troubles, and no fully satisfactory *in vivo* models exist to progress in the

understanding and cure of this disease. HGPRT knockout rodents do not show comparable motor

and behavioral troubles, which makes these models problematic for testing new therapeutic

treatments. In *Drosophila*, no HGPRT homologue is expressed, indicating that Aprt is the only

enzyme of the purine salvage pathway in this organism. Here we find that *Aprt* mutant flies show

symptoms partly comparable to the lack of HGPRT in humans, including increase in uric acid

levels, reduced longevity, and various neurobehavioral defects such as early locomotor decline,

sleep disorders and epilepsy behavior. This animal model therefore recapitulates both salvage

pathway disruption and motor symptoms, as observed in LND patients. Moreover, these results

highlighted that *Aprt* deficiency in *Drosophila* has more similarities with HGPRT than APRT

deficiency in humans. *Aprt* mutant flies also show a disruption of adenosine signaling and we

found that adenosine itself or a derivative compound can relieve the epileptic symptoms.

Moreover, we observed that neuronal expression of a mutant form of human HGPRT that causes

LND also triggers abnormalities in fly locomotion and seizure-like behavior, which has not been

documented to date in other models. The use of *Drosophila* to study LND should therefore prove

valuable to better understand the link between purine recycling deficiency and brain functioning

and carry out drug screening in a living organism, paving the way towards new improvements in

the cure of this dramatic disease.

Materials and Methods

***Drosophila* culture and strains**.

Flies were raised at 25°C on standard cornmeal-yeast-agar medium supplemented with

methylhydroxybenzoate as an antifungal under a 12h:12h light-dark (LD) cycle. The *Drosophila*

mutant lines were obtained either from the Bloomington *Drosophila* Stock Center (BDSC), the

Vienna *Drosophila* Resource Center (VDRC) or our own collection: *Aprt*[5] (Johnson and

Friedman, 1983) (BDSC #6882), *Df(3L)ED4284* (BDSC #8056), *Df(3L)BSC365* (BDSC

#24389), *UAS-Aprt*[RNAi] (VDRC #106836), *AdoR*[KG03964ex] (Wu et al. 2009) here named *AdoR*[KGex]

(BDSC #30868), and the following Gal4 drivers: *238Y-Gal4*, *24B-Gal4*, *da-Gal4*, *EAAT1-Gal4*

(Rival et al. 2004), *elav-Gal4* (*elav*[C155]), *MZ0709-Gal4*, *NP6510-Gal4*, *NP6520-Gal4*, *R58E02-*

Gal4 (Liu et al. 2012), *R76F05-Gal4*, *repo-Gal4*, *TH-Gal4* (Friggi-Grelin et al. 2003), *TRH-Gal4*

(Cassar et al. 2015), *tub-Gal80*[ts] and *VT30559-Gal4*. The Canton-S line was used as wild-type

control. The simplified driver>effector convention was used to indicate genotypes in figures, e.g.

elav>Aprt[RNAi] for *elav-Gal4*; *UAS-Aprt*[RNAi]. In some experiments, to restrict RNAi-mediated *Aprt*

inactivation at the adult stage, we have used the TARGET system (McGuire et al. 2003). Flies

were raised at 18°C (permissive temperature) where Gal4 transcriptional activity was inhibited by

tub-Gal80[ts], and shifted to 30°C (restrictive temperature) for 3 days before the test to enable

Gal4-driven *Aprt*[RNAi] expression.

***APRT and HPRT* transgenic lines**

To generate a *UAS-Aprt* strain, a 549-bp *Aprt* insert containing the coding sequence was

PCR amplified from the ORF clone BS15201 (*Drosophila* Genomics Resource Center,

Bloomington, IN, USA) using primers with added restriction sites (Table S3). After digestion with

EcoRI and XbaI, the *Aprt* cDNA was subcloned into the *pUASTattB* vector (Bischof et al. 2007)

using the In-Fusion HD Cloning Kit (Takara Bio, Kyoto, Japan) according to the manufacturer's

instructions, and the insertion verified by sequencing (Eurofins Genomics, Ebersberg, Germany).

The construction was sent to BestGene (Chino Hills, CA, USA) for *Drosophila* germline

transformation into the attP14 docking site on the 2d chromosome. The *UAS-Aprt* line yielding the

strongest expression was selected and used in the experiments.

A clone containing the human wild-type *HPRT1* cDNA was kindly provided to us by Prof.

Hyder A. Jinnah (Emory University, GA, USA). We constructed the *HPRT1-I42T* cDNA from this

clone by site-directed mutagenesis using the Quick Change II Site-Directed Mutagenesis Kit

(Agilent Technologies, Santa Clara, CA, USA). Primers sequences used to create the mutation

are shown in Table S3. The cDNA obtained was verified by sequencing. Then, the 657-bp

HPRT1-WT and *HPRT1-I42T* inserts were PCR amplified using primers with added restriction sites

and *Drosophila* translation start consensus sequence (Table S3). They were subcloned into

pUASTattB and verified by sequencing. The transgenes were sent to BestGene for *Drosophila*

transformation and inserted into the attP40 docking site on the 2d chromosome.

Sequencing of *Aprt*[5]

For sequencing of the *Aprt*[5] cDNA, total RNA was isolated by standard procedure from 20-30

heads of homozygous *Aprt*[5] flies and reverse transcribed using oligo d(T) primers (PrimeScript RT

Reagent Kit, Takara Bio, Kyoto, Japan). At least 750 ng of the first strand cDNA was amplified by

PCR in 20 µl of reaction mixture using PrimeStar Max DNA polymerase (Takara Bio, Kyoto

Japan) with a Techne Prime Thermal Cycler apparatus (Bibby Scientific, Burlington, NJ, USA).

The program cycles included 10 s denaturation at 98°C, 10 s annealing at 55°C and 10 s elongation

at 72°C, repeated 35 times. 1µL of the PCR product was amplified again with the same program, in

30 µl of reaction mixture. After elution on 1% agarose gel, DNA were purified using the

Wizard® SV Gel and PCR Clean-Up System protocol (Promega, Madison, Wisconsin, USA)

according to the manufacturer's instructions. Finally, 7.5 µL of purified DNA were sent with 2.5

µL of primers (sense and antisense in separate tubes) for sequencing (Eurofins Genomics,

Ebersberg, Germany).

Lifespan assay

Longevity study was performed as previously described (Riemensperger et al. 2011).

Drosophila adult males were collected within 24h of emergence and maintained on standard

medium at 25°C under a 12:12h LD cycle. They were transferred into fresh bottles every 2-3

days and the number or surviving flies was scored. 50 flies per bottle in triplicate were tested for

each genotype and the experiment was done 3 times.

Startle-induced negative geotaxis (SING)

SING assays were performed as previously described (Rival et al. 2004; Riemensperger et

al. 2011; Sun et al. 2018). For each genotype, 50 adult males divided into 5 groups of 10 flies

were placed in a vertical column (25-cm long, 1.5-cm diameter) with a conic bottom end and left

for about 30 min for habituation. Then, columns were tested individually by gently tapping them

down (startle), to which flies normally responded by climbing up. After 1 min, flies having

reached at least once the top of the column (above 22 cm) and flies that never left the bottom

(below 4 cm) were counted. Each fly group was assayed three times at 15 min intervals. The

performance index (PI) for each column is defined as $\frac{1}{2}[1 + (n_{top}-n_{bot})/n_{tot}]$, where n_{tot} is the total

number of flies in the column, n_{top} the number of flies at the top and n_{bot} the number of flies at the

bottom after 1 min. SING was performed at 1, 8, 10 and 30 days after adult eclosion (d a.E.).

Spontaneous locomotion and sleep monitoring

Spontaneous locomotor activity was recorded as previously described (Vaccaro et al.

2017), using *Drosophila* activity infra-red beam monitors (DAM, TriKinetics Inc, Waltham, MA

USA) placed in incubators at 25°C equipped with standard white light. 8-day-old male flies were

maintained individually for 5-6 days under 12:12h LD cycle in 5 x 65 mm glass tubes containing

5% sucrose, 1.5% agar medium. Data analysis was performed with the FaasX software (Klarsfeld

et al. 2003). Histograms represent the distribution of the activity through 24h in 30-min bins,

averaged for 32 flies per genotype over 4-5 cycles.

For sleep monitoring, 2-4-day-old virgin female flies were transferred individually into 5

x 65 mm glass tubes containing standard food and their movements were recorded for up to 5

days using DAM infra-red beam monitors (TriKinetics Inc, Waltham, MA USA) or the

Drosophila ARousal Tracking (DART) video system (Faville et al. 2015), in a 12h:12h LD cycle

with 50-60% humidity. Control and experimental flies were recorded simultaneously. Each

experiment included at least 14 flies for each condition and was repeated 2-3 times with

independent groups of flies. Fly sleep, defined as periods of immobility lasting more than 5 min

(Huber et al. 2004), was computed with a Microsoft Excel macro for the infra-red beam data and

with the DART software for the video data. Distribution and homogeneity as well as statistical

group comparisons were tested using Microsoft Excel plugin software StatEL. The *p* value shown

is the highest obtained among *post-hoc* comparisons and means ± SEM were plotted.

Immunohistochemistry

Brains of 8 to 10-day-old adult flies were dissected in ice-cold Ca^{2+}-free *Drosophila*

Ringer's solution and processed for whole-mount anti-TH or anti-DA immunostaining as

previously described (Riemensperger et al. 2011, Cichewicz et al. 2017). The primary antibodies

used were mouse monoclonal anti-TH (1:1000, cat. n° 22941, ImmunoStar, Hudson, WI, USA)

and mouse monoclonal anti-DA (1:100, cat. n° AM001, GemacBio, Saint-Jean-d'Illac, France).

The secondary antibody was goat anti-mouse conjugated to Alexa Fluor 488 (1:1000,

ThermoFisher Scientific, Waltham, MA, USA). Brains were mounted with antifade reagent,

either Prolong Gold (ThermoFisher Scientific, Waltham, MA, USA) or, alternatively, 65% 2,2'-

thiodiethanol (Sigma-Aldrich, St. Louis, MI, USA) (Cichewicz et al. 2017) for DA staining.

Images were acquired on a Nikon A1R confocal microscope with identical laser, filter and gain

settings for all samples in each experiment. Immunofluorescence intensity levels were quantified

using the Fiji software. Experiments were repeated independently at least 3 times on 4 to 6 brains

per genotype.

Bang-sensitivity test

Bang-sensitivity assays were performed as previously described (Howlett et al. 2013). 30-

day-old males were divided into 5 groups of 10 flies under CO_2 and allowed to recover overnight.

The following day, each group was placed in a vial without food and after 20 min of habituation,

the vials were stimulated vigorously with a vortex mixer for 10 s at 2,500 rpm. The recovery time

was measured for each fly, from the end of the stimulation until they reached a normal standing

position. Results are the mean of the recovery time for at least 50 flies per genotype.

Drug administration

Allopurinol (A8003, Sigma Aldrich, St. Louis, MI, USA) was diluted in standard medium

at 100 μg/ml and flies were placed for 5 days on this medium before metabolite extraction.

Adenine, adenosine and hypoxanthine (A2786, A9251 and H9377, Sigma Aldrich, St. Louis, MI,

USA) or N^6-methyladenosine (m^6A) (QB-1055, Combi-Blocks, San Diego, CA, USA) were

diluted in fly food medium at 500 μM. Parents were allowed to lay eggs on this medium in order

to have exposition to the drug throughout all larval development of the progeny. Adults were

collected and placed in normal medium until 5 days before the test, when they were placed again

in food supplemented with adenosine or m^6A at the same concentrations.

Uric acid quantification

For purine metabolite extraction, 40 heads from 8-day-old flies were ground in 80% ethanol, heated for 3 min at 80°C and evaporated in a Speedvac apparatus. Dried residues were resuspended in MilliQ water and total protein content of each homogenate was measured with the Pierce BCA Protein Assay Kit (ThermoFisher Scientific, Waltham, MA, USA). 20 µl of each sample was injected into a 25 cm x 4.6 mm C18 Nucleosil column with 5-µm particles (Interchim, Montluçon, France). Purine metabolites were detected by an Agilent 1290 Infinity HPLC system coupled with a diode array detector as recommended by the ERDNIM advisory document. 7 wavelengths were used for detection (230, 240, 250, 260, 266, 270 and 280 nm) in order to have the spectrum of each compound for identification in addition to the retention time. The mobile phases contained 0.05 M monopotassium phosphate pH 5 and 65% (v/v) acetonitrile. The flux was fixed at 1 ml/min. For analysis, the maximum height of the compound was normalized to protein content and compared to the control genotype.

ATP assay

ATP level was measured by bioluminescence using the ATP Determination Kit (A22066, ThermoFisher Scientific, Waltham, MA, USA) on a TriStar 2 Spectrum LB942S microplate reader (Berthold Technologies, Bad Wildbad, Germany), according to the manufacturer's instructions. Samples were prepared as described previously (Fergestad et al. 2006). Briefly, 30 heads or 5 thoraces from 8-day-old flies were homogenized in 200 µL of 6 M guanidine-HCl to inhibit ATPases, boiled directly 5 min at 95°C and then centrifuged for 5 min at 13,000 g and 4°C. Total protein content of each supernatant was measured with the BCA Protein Assay Kit.

Each supernatant was then diluted at 1:500 in TE buffer pH 8 and 10 µl were placed in a 96-well

white-bottom plate. The luminescent reaction solution (containing D-luciferine, recombinant

firefly luciferase, dithiothreitol and reaction buffer) was added to each well with an injector and

high-gain 1 sec exposure glow reads were obtained at 15 min after reaction initiation. Results

were compared to a standard curve generated with known ATP concentrations and final values

were normalized to the protein content.

Enzyme activity assay

HGPRT and APRT activities were assessed in *Drosophila* by adapting methods

previously established for human cells (Cartier and Hamet 1968, Ceballos-Picot et al. 2009). 20

whole male flies were homogenized in 250 µl of 110 mM Tris, 10 mM $MgCl_2$ pH 7.4 (Tris-

$MgCl_2$ buffer), immediately frozen and kept at least one night at -80°C before the assay. After 5

min of centrifugation at 13,000 g, total protein content of each supernatant was measured with the

BCA Protein Assay Kit to normalize activity level. Kinetics of [^{14}C]-hypoxanthine conversion to

IMP and [^{14}C]-adenine conversion to AMP were assessed for HGPRT and Aprt assays,

respectively. Compositions of reaction mediums were: 25 µl of a radioactive solution made with

38 µl of [^{14}C]-hypoxanthine (25 µCi/ml) diluted in 1 ml of 1.2 mM cold hypoxanthine, for

HGPRT assay, or 25 µl of a radioactive solution made with 75 µl of [^{14}C]-adenine (50 µCi/ml)

diluted in 1 ml of 1.2 mM cold adenine for Aprt assay, and a volume of fly extract equivalent to

200 µg protein diluted in Tris-$MgCl_2$ buffer to 150 µl. Reactions were monitored at 37°C and

started by adding 25 µl of the co-factor 5-phosphoribosyl-1-pyrophosphate (PRPP) at 10 mM.

After 6, 12, 24 and 36 min, 40 µL of each pool was placed in a tube containing either 25 µl of

HIE (3 mM hypoxanthine, 6 mM IMP, 200 mM EDTA) or AAE (3 mM adenine, 6 mM AMP,

200 mM EDTA) solutions and incubated for 3 min at 95°C to stop the reaction. The different

radioactive compounds were separated by paper chromatography on Whatman 3MM strips using

28% NH$_4$OH, 50 mM EDTA as solvent for about 1h30. Then, the substrates and products were

visualized under a UV lamp at 254 nm and placed separately in vials in 2 ml of Scintran (VWR,

Radnor, PA, USA). The radioactivity in disintegrations per minute (dpm) was measured in a

Packard Tri-Carb 1600 TR liquid scintillation analyzer (PerkinElmer, Waltham, MA, USA). The

percentage of substrate transformation as a function of time was converted in nmol/min/mg

protein and finally normalized to wild-type control values.

1005

Reverse transcription-PCR

Total RNA was isolated by standard procedure from 20-30 heads of 8-day-old males

collected on ice and lysed in 600 µl QIAzol Reagent (Qiagen, Venlo, Netherlands). 1 µg of total

RNA was treated by DNase (DNase I, RNase-free, ThermoFisher Scientific, Waltham, MA,

USA) according to the manufacturer's instructions. 5 µL of treated RNA was reverse transcribed

using oligo d(T) primers (PrimeScript RT Reagent Kit – Takara Bio, Kyoto Japan). Then, at least

750 ng of the first strand cDNA was amplified in 20 µl of reaction mixture using PrimeStar Max

DNA polymerase (Takara Bio, Kyoto Japan) with a Techne Prime Thermal Cycler apparatus

(Bibby Scientific, Burlington, NJ, USA). The program cycles included 10 s denaturation at 98°C,

10 s annealing at 55°C and 10 s elongation at 72°C, repeated 30 times. PCR product levels were

measured after electrophoresis by densitometry analysis with the Fiji software. Data were

normalized to amplification level of the ribosomal protein *rp49/RpL32* transcripts as internal

control.   Sequences   of   the   primers   used   were: sense   *HPRT*,   5'-

GAGATACAAAATGGCGACCCGCAGCCCT;   antisense   *HPRT*,   5'-

GCTCGGATCCTTATCATTACTAGGCTTTG (amplicon 686bp); sense *rp49*, 5'-

GACGCTTCAAGGGACAGTATC; antisense *rp49*, 5'- AAACGCGGTTCTGCATGAG

(amplicon 144bp).

1023

Quantitative reverse transcription PCR

For RT-qPCR, total RNA was isolated by standard procedure from 20-30 heads of 8-day-

old males collected on ice and lysed in 600 µl QIAzol Lysis Reagent (Qiagen, Venlo,

Netherlands). 1 µg total RNA was reverse transcribed using oligo d(T) primers (PrimeScript RT

Reagent Kit, Takara Bio, Kyoto, Japan). Approximately 40 ng of the first strand cDNA was

amplified in 10 µl of reaction mixture using the LightCycler 480 SYBR Green I Master reaction

mix (Roche Applied Science, Mannheim, Germany) and a LightCycler 480 Instrument (Roche

Applied Science, Mannheim, Germany). The program cycles included a 10-min preincubation

step at 95°C, 40 cycles of amplification (10 s denaturation at 95°C, 10 s annealing at 55°C, 20 s

elongation at 72°C), followed by a melting curves analysis for PCR product identification. Data

were normalized to amplification level of the ribosomal protein *rp49/RpL32* transcripts as

internal control. The genes analyzed and primer sequences used for qPCR are indicated in Table

S4.

1037

Adenosine assay

Adenosine was determined by Ultra Performance Liquid Chromatography (UPLC). 5

whole flies or 30 heads were homogenized in 120 µl 0.9% (w/v) NaCl using a Minilys apparatus

(Bertin Instruments, Montigny-le-Bretonneux, France) and frozen at -80°C. After unfreezing and

5 min of microcentrifugation, 20 µl of 10% perchloric acid were added to 70 µl of the

supernatant and the mixture was left for 5 min on ice. After a new centrifugation, 20 µl of a

neutralization solution (made by mixing 3 volumes of 3 M K_2CO_3 with 1 volume of 6.4 mM

NaOH containing 0.4 mg/ml bromothymol blue) were added and the mixture was centrifuged

again before injection (5 µl). Samples were analyzed with a UV diode-array detector on an

Acquity UPLC HSS T3 column (1,8 µm, 2.1 x 150mm) (Waters Corporation, Milford, MA,

USA). The mobile phases consisted in Buffer A (30 mM ammonium acetate, pH 4.0 with

1:10,000 heptafluorobutyric acid (HPFA)) and Buffer B (acetonitrile with 1:10,000 HPFA), using

a flow rate of 0.3 ml/min. Chromatographic conditions were: 3.5 min 100% Buffer A, 16.5 min

up to 6.3% Buffer B, 2 min up to 100% Buffer B, and 1 min 100% Buffer B. The gradient was

then returned over 5 min to 100% Buffer A, restoring the initial conditions. Results were

normalized to protein levels for each sample.

1054

Statistics

Statistical significance was determined with the Prism 6 software (GraphPad Software, La

Jolla, CA, USA). Survival curves for longevity experiments were analyzed using the log-rank

test. The Student's *t*-test was used to compare two genotypes or conditions, and one-way or two-

way ANOVA with Tukey's or Sidak's *post-hoc* multiple comparison tests for three or more

conditions. Results are presented as mean ± SEM. Probability values in all figures: *$p < 0.05$, **p

< 0.01, ***$p < 0.001$.

1062

Acknowledgements

We are grateful to Pr. Hyder A. Jinnah for HPRT cDNA gift and Dr Thomas

Riemensperger for helpful discussion. This work was supported by fundings from PSL Research

University and ESPCI Paris to SB and Institut de France to ICP. CP was recipient of PhD

fellowships from Association Lesch-Nyhan Action (LNA), Association Malaury and Labex

MemoLife.

1069

Authors contributions

1071   Conceptualization, C.P., L.S., S.M., I.C.P. and S.B.; investigation, C.P., L.S., A.D., R.F.,

1072   M.S., J.D. and S.M.; formal analysis, C.P., L.S., A.D., R.F., M.S., J.D. and S.M.; methodology,

1073   C.P., L.S., A.M., B.C.Z and S.M.; validation, C.P., I.C.P and S.B.; writing, original draft, C.P.,

1074   I.C.P. and S.B.; writing, review & editing, C.P., I.C.P and S.B.; funding acquisition and

1075   supervision, I.C.P and S.B.; project administration, S.B.

References

Artiushin. G., & Sehgal, A. (2017). The *Drosophila* circuitry of sleep-wake regulation. *Current Opinion in Neurobiology*, *44*, 243-250. https://doi.org/10.1016/j.conb.2017.03.004

Aso, Y., Grübel, K., Busch, S., Friedrich, A. B., Siwanowicz, I., & Tanimoto, H. (2009). The mushroom body of adult *Drosophila* characterized by Gal4 drivers. *Journal of Neurogenetics*, *23*(1-2), 156-172. https://doi.org/10.1080/01677060802471718

Awasaki, T., Lai, S.-L., Ito, K., & Lee, T. (2008). Organization and postembryonic development of glial cells in the adult central brain of *Drosophila*. *The Journal of Neuroscience*, *28*(51), 13742-13753. https://doi.org/10.1523/JNEUROSCI.4844-08.2008

Batista, P. J. (2017). The RNA modification N6-methyladenosine and its implications in human disease. *Genomics, Proteomics & Bioinformatics*, *15*(3), 154-163. https://doi.org/10.1016/j.gpb.2017.03.002

Baumeister, A. A., & Frye, G. D. (1985). The biochemical basis of the behavioral disorder in the Lesch-Nyhan syndrome. *Neuroscience & Biobehavioral Reviews*, *9*(2), 169-178. https://doi.org/10.1016/0149-7634(85)90043-0

Becker, J.-L. (1974a). Purine metabolism pathways in *Drosophila* cells growth "*in vitro*": phosphoribosyl transferase activities. *Biochimie*, *56*(5), 779-781. https://doi.org/10.1016/S0300-9084(74)80051-9

Becker, J.-L. (1974b). Purine metabolism in *Drosophila melanogaster* cells in culture *in vitro*: Purine interconversion [*in French*]. *Biochimie*, *56*(9), 1249-1253. https://doi.org/10.1016/S0300-9084(74)80018-0

Bell, S., McCarty, V., Peng, H., Jefri, M., Hettige, N., Antonyan, L., Crapper, L., O'Leary, L. A., Zhang, X., Zhang, Y., Wu, H., Sutcliffe, D., Kolobova, I., Rosenberger, T. A., Moquin, L., Gratton, A., Popic, J., Gantois, I., Stumpf, P. S., Schuppert, A. A., Mechawar, N., Sonenberg, N., Tremblay, M. L., Jinnah, H. A., & Ernst, C. (2021). Lesch-Nyhan disease causes impaired energy metabolism and reduced developmental potential in midbrain dopaminergic cells. *Stem Cell Reports*, *16*(7):1749-1762. http://doi.org/10.1016/j.stemcr.2021.06.003

Bell, S., Kolobova, I., Crapper, L., & Ernst, C. (2016). Lesch-Nyhan syndrome: models, theories, and therapies. *Molecular Syndromology*, *7*(6), 302-311. https://doi.org/10.1159/000449296

Bertelli, M., Cecchin, S., Lapucci, C., Jacomelli, G., Jinnah, H.A., Pandolfo, M., & Micheli, V. (2006). Study of the adenosinergic system in the brain of HPRT knockout mouse (Lesch-Nyhan disease). *Clinica Chimica Acta; International Journal of Clinical Chemistry*, *373*(1-2): 104-7. https://doi.org/10.1016/j.cca.2006.05.013.

Birman, S. (2005). Arousal mechanisms: speedy flies don't sleep at night. *Current Biology*, *15*(13), R511-R513. https://doi.org/10.1016/j.cub.2005.06.032

Bischof, J., Maeda, R. K., Hediger, M., Karch, F., & Basler, K. (2007). An optimized transgenesis system for *Drosophila* using germ-line-specific phiC31 integrases. *Proceedings of the National Academy of Sciences*, *104*(9), 3312-3317. https://doi.org/10.1073/pnas.0611511104

Boison, D. (2005). Adenosine and epilepsy: from therapeutic rationale to new therapeutic strategies. *The Neuroscientist*, *11*(1), 25-36. https://doi.org/10.1177/1073858404269112

Bollée, G., Harambat, J., Bensman, A., Knebelmann, B., Daudon, M., & Ceballos-Picot, I. (2012). Adenine phosphoribosyltransferase deficiency. *Clinical Journal of the American Society of Nephrology*, *7*(9), 1521-1527. https://doi.org/10.2215/CJN.02320312

Booth, W. T., Schlachter, C. R., Pote, S., Ussin, N., Mank, N. J., Klapper, V., Offermann, L. R., Tang, C., Hurlburt, B. K., & Chruszcz, M. (2018). Impact of an N-terminal polyhistidine tag on protein thermal stability. *ACS Omega*, *3*(1), 760-768. https://doi.org/10.1021/acsomega.7b01598

Boswell-Casteel, R.C., & Hays, F.A. (2017). Equilibrative nucleoside transporters-a review. *Nucleosides, Nucleotides and Nucleic Acids*, *36*(1): 7-30. https://doi.org/10.1080/15257770.2016.1210805.

Bou Dib, P., Gnägi, B., Daly, F., Sabado, V., Tas, D., Glauser, D. A., Meister, P., & Nagoshi, E. (2014). A conserved role for p48 homologs in protecting dopaminergic neurons from

oxidative stress. *PLoS Genetics*, *10*(10), e1004718. https://doi.org/10.1371/journal.pgen.1004718

Breese, G. R., Criswell, H. E., Duncan, G. E., & Mueller, R. A. (1990). A dopamine deficiency model of Lesch-Nyhan disease—the neonatal-6-OHDA-lesioned rat. *Brain Research Bulletin*, *25*(3), 477–484. https://doi.org/10.1016/0361-9230(90)90240-Z

Brody,T., & Cravchik, A. (2000). *Drosophila melanogaster* G protein-coupled receptors. *Journal of Cell Biology*, *150*(2), F83–F88. 8. https://doi.org/10.1083/jcb.150.2.F83

Cartier, P., & Hamet, M. (1968). Determination of the purine phosphoribosyl transferase activity of human red cells [*in French*]. *Clinica Chimica Acta; International Journal of Clinical Chemistry*, *20*(2), 205-214. https://doi.org/10.1016/0009-8981(68)90152-6

Cassar, M., Issa, A.-R., Riemensperger, T., Petitgas, C., Rival, T., Coulom, H., Iche-Torres, M., Han, K.-A., & Birman, S. (2015). A dopamine receptor contributes to paraquat-induced neurotoxicity in *Drosophila*. *Human Molecular Genetics*, *24*(1), 197-212. https://doi.org/10.1093/hmg/ddu430

Ceballos-Picot, I., Mockel, L., Potier, M.-C., Dauphinot, L., Shirley, T. L., Torero-Ibad, R., Fuchs, J., & Jinnah, H. A. (2009). Hypoxanthine-guanine phosphoribosyl transferase regulates early developmental programming of dopamine neurons: implications for Lesch-Nyhan disease pathogenesis. *Human Molecular Genetics*, *18*(13), 2317-2327. https://doi.org/10.1093/hmg/ddp164

Ceballos-Picot,I, Le Dantec A., Brassier A., Jaïs J-P., Ledroit M., Cahu J., Ea H-K., Daignan-Fornier B., Pinson B. New biomarkers for early diagnosis of Lesch-Nyhan disease revealed by metabolic analysis on a large cohort of patients. Orphanet J Rare Dis. 2015 Jan 23; 10:7. https://doi.org/10.1186/s13023-014-0219-0.

Cichewicz, K., Garren, E.J., Adiele, C., Aso, Y., Wang, Z., Wu, M., Birman, S., Rubin, G.M., & Hirsh, J. (2017). A new brain dopamine-deficient *Drosophila* and its pharmacological and genetic rescue. *Genes, Brain and Behavior*, *16*(3), 394-403. https://doi.org/10.1111/gbb.12353.

Cristini, S., Navone, S., Canzi, L., Acerbi, F., Ciusani, E., Hladnik, U., de Gemmis, P., Alessandri, G., Colombo, A., Parati, E., & Invernici, G. (2010). Human neural stem cells: A model system for the study of Lesch-Nyhan disease neurological aspects. *Human Molecular Genetics*, *19*(10), 1939-1950. https://doi.org/10.1093/hmg/ddq072

Criswell, H., Mueller, R.A., & Breese, G.R. (1988). Assessment of purine-dopamine interactions in 6-hydroxydopamine-lesioned rats: evidence for pre- and postsynaptic influences by adenosine. *Journal of Pharmacology and Experimental Therapeutics*, *244*(2): 493-500.

Doherty, J., Logan, M. A., Tasdemir, O. E., & Freeman, M. R. (2009). Ensheathing glia function as phagocytes in the adult *Drosophila* brain. *The Journal of Neuroscience*, *29*(15), 4768-4781. https://doi.org/10.1523/JNEUROSCI.5951-08.2009

Dolezelova, E., Nothacker, H.-P., Civelli, O., Bryant, P.J., & Zurovec, M. (2007). A *Drosophila* adenosine receptor activates cAMP and calcium signaling. *Insect Biochemistry and Molecular Biology*, *37*(4): 318-29. https://doi.org/10.1016/j.ibmb.2006.12.003.

Dung, V. M., & Thao, D. T. P. (2018). Parkinson's disease model. *Advances in Experimental Medicine and Biology*, *1076*, 41-61. https://doi.org/10.1007/978-981-13-0529-0_4

Dunnett, S. B., Sirinathsinghji, D. J. S., Heavens, R., Rogers, D. C., & Kuehn, M. R. (1989). Monoamine deficiency in a transgenic (*Hprt*⁻) mouse model of Lesch-Nyhan syndrome. *Brain Research*, *501*(2), 401-406. https://doi.org/10.1016/0006-8993(89)90659-8

Egami, K., Yitta, S., Kasim, S., Lewers, J. C., Roberts, R. C., Lehar, M., & Jinnah, H. A. (2007). Basal ganglia dopamine loss due to defect in purine recycling. *Neurobiology of Disease*, *26*(2), 396-407. https://doi.org/10.1016/j.nbd.2007.01.010

Engle, S. J., Womer, D. E., Davies, P. M., Boivin, G., Sahota, A., Simmonds, H. A., Stambrook, P. J., & Tischfield, J. A. (1996). HPRT-APRT-deficient mice are not a model for Lesch-Nyhan syndrome. *Human Molecular Genetics*, *5*(10), 1607–1610. https://doi.org/10.1093/hmg/5.10.1607

Ernst, M., Zametkin, A. J., Matochik, J. A., Pascualvaca, D., Jons, P. H., Hardy, K., Hankerson, J. G., Doudet, D. J., & Cohen, R. M. (1996). Presynaptic dopaminergic deficits in Lesch–Nyhan disease. *New England Journal of Medicine*, *334*(24), 1568–1572. https://doi.org/10.1056/NEJM199606133342403

Faville, R., Kottler, B., Goodhill, G. J., Shaw, P. J., & van Swinderen, B. (2015). How deeply does your mutant sleep? Probing arousal to better understand sleep defects in *Drosophila*. *Scientific Reports*, *5*, 8454. https://doi.org/10.1038/srep08454

Feany, M. B., & Bender, W. W. (2000). A *Drosophila* model of Parkinson's disease. *Nature*, *404*(6776), 394-398. https://doi.org/10.1038/35006074

Fergestad, T., Bostwick, B., & Ganetzky, B. (2006). Metabolic disruption in *Drosophila* bang-sensitive seizure mutants. *Genetics*, *173*(3), 1357-1364. https://doi.org/10.1534/genetics.106.057463

Fernández-Hernández, I., Scheenaard, E., Pollarolo, G., & Gonzalez, C. (2016). The translational relevance of *Drosophila* in drug discovery. *EMBO Reports*, *17*(4), 471-472. https://doi.org/10.15252/embr.201642080

Finger, S., Heavens, R. P., Sirinathsinghji, D. J., Kuehn, M. R., & Dunnett, S. B. (1988). Behavioral and neurochemical evaluation of a transgenic mouse model of Lesch-Nyhan syndrome. *Journal of the Neurological Sciences*, *86*(2-3), 203-213. https://doi.org/10.1016/0022-510x(88)90099-8

Franco, R., Ferré, S., Agnati, L., Torvinen, M., Ginés, S., Hillion, J., Casadó, V., Lledó, P.-M., Zoli, M., Lluis, C., et al. (2000). Evidence for adenosine/dopamine receptor interactions: indications for heteromerization. *Neuropsychopharmacology*, *23*: S50–S59. https://doi.org/10.1016/S0893-133X(00)00144-5.

Friggi-Grelin, F., Coulom, H., Meller, M., Gomez, D., Hirsh, J., & Birman, S. (2003). Targeted gene expression in *Drosophila* dopaminergic cells using regulatory sequences from tyrosine hydroxylase. *Journal of Neurobiology*, *54*(4), 618-627. https://doi.org/10.1002/neu.10185

Fu, R., Ceballos-Picot, I., Torres, R. J., Larovere, L. E., Yamada, Y., Nguyen, K. V., Hegde, M., Visser, J. E., Schretlen, D. J., Nyhan, W. L., Puig, J. G., O'Neill, P. J., Jinnah, H. A., & for the Lesch-Nyhan Disease International Study Group. (2014). Genotype-phenotype correlations in neurogenetics: Lesch-Nyhan disease as a model disorder. *Brain*, *137*(5), 1282-1303. https://doi.org/10.1093/brain/awt202

Fu, R., & Jinnah, H. A. (2012). Genotype-phenotype correlations in Lesch-Nyhan disease: moving beyong the gene. *Journal of Biological Chemistry*, *287*(5), 2997-3008. https://doi.org/10.1074/jbc.M111.317701

Fu, Rong, Sutcliffe, D., Zhao, H., Huang, X., Schretlen, D. J., Benkovic, S., & Jinnah, H. A. (2015). Clinical severity in Lesch–Nyhan disease: the role of residual enzyme and compensatory pathways. *Molecular Genetics and Metabolism*, *114*(1), 55-61. https://doi.org/10.1016/j.ymgme.2014.11.001

García, M.G., Puig, J.G., & Torres, R.J. (2009). Abnormal adenosine and dopamine receptor expression in lymphocytes of Lesch–Nyhan patients. *Brain Behavior and Immunity*, *23*(8): 1125-31. https://doi.org/10.1016/j.bbi.2009.07.006.

García, M.G., Puig, J.G., & Torres, R.J. (2012). Adenosine, Dopamine and Serotonin Receptors Imbalance in Lymphocytes of Lesch-Nyhan Patients. *Journal of Inherited Metabolic Disease*, *35*(6): 1129-35. https://doi.org/10.1007/s10545-012-9470-5.

Gatto, C. L., & Broadie, K. (2011). *Drosophila* modeling of heritable neurodevelopmental disorders. *Current Opinion in Neurobiology*, *21*(6), 834-841. https://doi.org/10.1016/j.conb.2011.04.009

Gray, J.H., Owen, R.P., & Giacomini, K.M. (2004). The concentrative nucleoside transporter family, SLC28. *Pflugers Archiv European Journal of Physiology*, *447*(5): 728-34. https://doi.org/10.1007/s00424-003-1107-y.

Guibinga, G.-H., Hrustanovic, G., Bouic, K., Jinnah, H. A., & Friedmann, T. (2012). MicroRNA-mediated dysregulation of neural developmental genes in HPRT deficiency: clues for Lesch-Nyhan disease? *Human Molecular Genetics*, *21*(3), 609-622. https://doi.org/10.1093/hmg/ddr495

Harambat, J., Bollée, G., Daudon, M., Ceballos-Picot, I., & Bensman, A. (2012). Adenine phosphoribosyltransferase deficiency in children. *Pediatric Nephrology*, *27*(4), 571-579. https://doi.org/10.1007/s00467-011-2037-0

Hardie, S. L., & Hirsh, J. (2006). An improved method for the separation and detection of biogenic amines in adult *Drosophila* brain extracts by high performance liquid chromatography. *Journal of Neuroscience Methods*, *153*(2), 243-249. https://doi.org/10.1016/j.jneumeth.2005.11.001

Harkness, R. A., McCreanor, G. M., & Watts, R. W. E. (1988). Lesch-Nyhan syndrome and its pathogenesis: purine concentrations in plasma and urine with metabolite profiles in CSF. *Journal of Inherited Metabolic Disease*, *11*(3), 239-252. https://doi.org/10.1007/BF01800365

Howlett, I. C., Rusan, Z. M., Parker, L., & Tanouye, M. A. (2013). *Drosophila* as a model for intractable epilepsy: *Gilgamesh* suppresses seizures in *para*[bss1] heterozygote flies. *G3; Genes|Genomes|Genetics*, *3*(8), 1399-1407. https://doi.org/10.1534/g3.113.006130

Huber R, Hill SL, Holladay C, Biesiadecki M, Tononi G, & Cirelli C. (2004). Sleep homeostasis in *Drosophila melanogaster*. *Sleep*, *15*;27(4):628-39. https://doi.org/10.1093/sleep/27.4.628

Ilie, A., Raimondo, J. V., & Akerman, C. J. (2012). Adenosine release during seizures attenuates $GABA_A$ receptor-mediated depolarization. *The Journal of Neuroscience*, *32*(15), 5321-5332. https://doi.org/10.1523/JNEUROSCI.5412-11.2012

Ipata, P. L. (2011). Origin, utilization, and recycling of nucleosides in the central nervous system. *Advances in Physiology Education*, *35*(4), 342-346. https://doi.org/10.1152/advan.00068.2011

Issa, A.-R., Sun, J., Petitgas, C., Mesquita, A., Dulac, A., Robin, M., Mollereau, B., Jenny, A., Chérif-Zahar, B., & Birman, S. (2018). The lysosomal membrane protein LAMP2A promotes autophagic flux and prevents SNCA-induced Parkinson disease-like symptoms in the *Drosophila* brain. *Autophagy*, *14*(11), 1898-1910. https://doi.org/10.1080/15548627.2018.1491489

Jaiswal, M., Sandoval, H., Zhang, K., Bayat, V., & Bellen, H. J. (2012). Probing mechanisms that underlie human neurodegenerative diseases in *Drosophila*. *Annual Review of Genetics*, *46*, 371-396. https://doi.org/10.1146/annurev-genet-110711-155456

Jinnah, H. A., Ceballos-Picot, I., Torres, R. J., Visser, J. E., Schretlen, D. J., Verdu, A., Larovere, L. E., Chen, C.-J., Cossu, A., Wu, C.-H., Sampat, R., Chang, S.-J., de Kremer, R. D., Nyhan, W., Harris, J. C., Reich, S. G., Puig, J. G., & for the Lesch-Nyhan Disease International Study Group. (2010). Attenuated variants of Lesch-Nyhan disease. *Brain*, *133*(3), 671-689. https://doi.org/10.1093/brain/awq013

Jinnah, H. A., Page, T., & Friedmann, T. (1993). Brain purines in a genetic mouse model of Lesch-Nyhan disease. *Journal of Neurochemistry*, *60*(6), 2036-2045. https://doi.org/10.1111/j.1471-4159.1993.tb03488.x

Jinnah, H. A., Visser, J. E., Harris, J. C., Verdu, A., Larovere, L., Ceballos-Picot, I., Gonzalez-Alegre, P., Neychev, V., Torres, R. J., Dulac, O., Desguerre, I., Schretlen, D. J., Robey, K. L., Barabas, G., Bloem, B. R., Nyhan, W., De Kremer, R., Eddey, G. E., Puig, J. G., Reich S. G.; Lesch-Nyhan Disease International Study Group. (2006). Delineation of the motor disorder of Lesch-Nyhan disease. *Brain*, *129*(Pt 5), 1201-1217. https://doi.org/10.1093/brain/awl056

Jinnah, H. A., Wojcik, B. E., Hunt, M., Narang, N., Lee, K. Y., Goldstein, M., Wamsley, J. K., Langlais, P. J., & Friedmann, T. (1994). Dopamine deficiency in a genetic mouse model of Lesch-Nyhan disease. *The Journal of Neuroscience*, *14*(3), 1164-1175. https://doi.org/10.1523/JNEUROSCI.14-03-01164.1994

Jinnah, H.A., Gage, F.H., & Friedmann, T. (1990). Animal models of Lesch-Nyhan syndrome. *Brain Research Bulletin*, *25*(3): 467-75. https://doi.org/10.1016/0361-9230(90)90239-V.

Johnson, D. H., Edström, J.-E., Burnett, J. B., & Friedman, T. B. (1987). Cloning of a *Drosophila melanogaster* adenine phosphoribosyltransferase structural gene and deduced amino acid sequence of the enzyme. *Gene*, *59*(1), 77-86. https://doi.org/10.1016/0378-1119(87)90268-X

Johnson, D. H., & Friedman, T. B. (1981). Purine resistant mutants of *Drosophila* are adenine phosphoribosyltransferase deficient. *Science*, *212*(4498), 1035-1036. https://doi.org/10.1126/science.212.4498.1035

Johnson, D. H., & Friedman, T. B. (1983). Purine-resistant *Drosophila melanogaster* result from mutations in the adenine phosphoribosyltransferase structural gene. *Proceedings of the National Academy of Sciences*, *80*(10), 2990-2994. https://doi.org/10.1073/pnas.80.10.2990

Johnson, E. C., Kazgan, N., Bretz, C. A., Forsberg, L. J., Hector, C. E., Worthen, R. J., Onyenwoke, R., & Brenman, J. E. (2010). Altered metabolism and persistent starvation behaviors caused by reduced AMPK function in *Drosophila*. *PLoS One*, *5*(9), e12799. https://doi.org/10.1371/journal.pone.0012799

Johnson, T. A., Jinnah, H. A., & Kamatani, N. (2019). Shortage of cellular ATP as a cause of diseases and strategies to enhance ATP. *Frontiers in Pharmacology*, *In press*. https://doi.org/10.3389/fphar.2019.00098

Kelley, W. N., Rosenbloom, F. M., Henderson, J. F., & Seegmiller, J. E. (1967). A specific enzyme defect in gout associated with overproduction of uric acid. *Proceedings of the National Academy of Sciences of the United States of America*, *57*(6), 1735-1739. https://doi.org/10.1073/pnas.57.6.1735

Kim, D.S., & Palmiter, R.D. (2008). Interaction of dopamine and adenosine receptor function in behavior: studies with dopamine-deficient mice. *Frontiers in Bioscience*, *13*: 2311-18. https://doi.org/10.2741/2845.

Klarsfeld, A., Leloup, J.-C., & Rouyer, F. (2003). Circadian rhythms of locomotor activity in *Drosophila*. *Behavioural Processes*, *64*(2), 161-175. https://doi.org/10.1016/S0376-6357(03)00133-5

Knapp, D. J., & Breese, G. R. (2016). The use of perinatal 6-hydroxydopamine to produce a rodent model of Lesch-Nyhan disease. *Current Topics in Behavioral Neurosciences*, *29*, 265-277. https://doi.org/10.1007/7854_2016_444

Knight, D., Harvey, P.J., Iliadi, K.G., Klose, M.K., Iliadi, N., Dolezelova, E., Charlton, M.P., Zurovec, M., & Boulianne, G.L. (2010). Equilibrative nucleoside transporter 2 regulates associative learning and synaptic function in *Drosophila*. *Journal of Neuroscience*, *30*(14): 5047-57. https://doi.org/10.1523/JNEUROSCI.6241-09.2010.

Kroll, J. R., Wong, K. G., Siddiqui, F. M., & Tanouye, M. A. (2015). Disruption of endocytosis with the dynamin mutant shibire[ts1] suppresses seizures in *Drosophila*. *Genetics*, *201*(3), 1087-1102. https://doi.org/10.1534/genetics.115.177600

Kroll, Jason R., & Tanouye, M. A. (2013). Rescue of *easily shocked* mutant seizure sensitivity in *Drosophila* adults: ethanolamine kinase and seizure susceptibility. *Journal of Comparative Neurology*, *521*(15), 3500-3507. https://doi.org/10.1002/cne.23364

Kumar, S., Chen, D., & Sehgal, A. (2012). Dopamine acts through cryptochrome to promote acute arousal in *Drosophila*. *Genes & Development*, *26*(11), 1224-1234. https://doi.org/10.1101/gad.186338.111

Lahaye, C., Augé, F., Soubrier, M., & Ceballos-Picot, I. (2014). New mutation affecting hypoxanthine phosphoribosyltransferase responsible for severe tophaceous gout. *The Journal of Rheumatology*, *41*(6), 1252-1254. https://doi.org/10.3899/jrheum.131168

Lee, G., Kikuno, K., Bahn, J.-H., Kim, K.-M., & Park, J. H. (2013). Dopamine D2 receptor as a cellular component controlling nocturnal hyperactivities in *Drosophila melanogaster*. *Chronobiology International*, *30*(4), 443-459. https://doi.org/10.3109/07420528.2012.741169

Lesch, M., & Nyhan, W. L. (1964). A familial disorder of uric acid metabolism and central nervous system function. *The American Journal of Medicine*, *36*(4), 561-570. https://doi.org/10.1016/0002-9343(64)90104-4

Liu, C., Plaçais, P.-Y., Yamagata, N., Pfeiffer, B. D., Aso, Y., Friedrich, A. B., Siwanowicz, I., Rubin, G. M., Preat, T., & Tanimoto, H. (2012). A subset of dopamine neurons signals reward for odour memory in *Drosophila*. *Nature*, *488*(7412), 512-516. https://doi.org/10.1038/nature11304

Lloyd, K. G., Hornykiewicz, O., Davidson, L., Shannak, K., Farley, I., Goldstein, M., Shibuya, M., Kelley, W. N., & Fox, I. H. (1981). Biochemical evidence of dysfunction of brain neurotransmitters in the Lesch-Nyhan syndrome. *New England Journal of Medicine*, *305*(19), 1106-1111. https://doi.org/10.1056/NEJM198111053051902

López, J. M. Is ZMP the toxic metabolite in Lesch-Nyhan disease? (2008). *Med Hypotheses*, *71*(5):657-63. http://doi.org/10.1016/j.mehy.2008.06.033

López, J. M., Outtrim, E. L., Fu, R., Sutcliffe, D. J., Torres, R. J., & Jinnah, H. A. (2020). Physiological levels of folic acid reveal purine alterations in Lesch-Nyhan disease. *Proc Natl Acad Sci U S A*, *117*(22):12071-12079. http://doi.org/10.1073/pnas.2003475117

Machado, J., Abdulla, P., Hanna, W.J.B., Hilliker, A.J., & Coe, I.R. (2007). Genomic analysis of nucleoside transporters in *Diptera* and functional characterization of *DmENT2*, a *Drosophila* equilibrative nucleoside transporter. *Physiological Genomics*, *28*(3): 337-47. https://doi.org/10.1152/physiolgenomics.00087.2006.

Madeo, A., Di Rocco, M., Brassier, A., Bahi-Buisson, N., De Lonlay, P., & Ceballos-Picot, I. (2019). Clinical, biochemical and genetic characteristics of a cohort of 101 French and Italian patients with HPRT deficiency. *Molecular Genetics and Metabolism*, *127*(2), 147-157. https://doi.org/10.1016/j.ymgme.2019.06.001

Marks, J. F., Baum, J., Keele, D. K., Kay, J. L., & MacFarlen, A. (1968). Lesch-Nyhan syndrome treated from the early neonatal period. *Pediatrics*, *42*(2), 357-359.

Masino, S. A., Kawamura, M., & Ruskin, D. N. (2014). Adenosine receptors and epilepsy: current evidence and future potential. *International Review of Neurobiology*, *119*, 233-255. https://doi.org/10.1016/B978-0-12-801022-8.00011-8

Mazaud, D., Kottler, B., Gonçalves-Pimentel, C., Proelss, S., Tüchler, N., Deneubourg, C., Yuasa, Y., Diebold, C., Jungbluth, H., Lai, E. C., Hirth, F., Giangrande, A., Fanto, M. (2019). Transcriptional regulation of the glutamate/GABA/glutamine cycle in adult glia controls motor activity and seizures in *Drosophila*. *Journal of Neuroscience*, *39*(27):5269-5283. https://doi.org/10.1523/JNEUROSCI.1833-18.2019

McGuire, S. E., Le, P. T., Osborn, A. J., Matsumoto, K., & Davis, R. L. (2003). Spatiotemporal rescue of memory dysfunction in *Drosophila*. *Science (New York, N.Y.)*, *302*(5651), 1765-1768. https://doi.org/10.1126/science.1089035

McGurk, L., Berson, A., & Bonini, N. M. (2015). *Drosophila* as an *in vivo* model for human neurodegenerative disease. *Genetics*, *201*(2), 377-402. https://doi.org/10.1534/genetics.115.179457

Meek, S., Thomson, A. J., Sutherland, L., Sharp, M. G. F., Thomson, J., Bishop, V., Meddle, S. L., Gloaguen, Y., Weidt, S., Singh-Dolt, K., Buehr, M., Brown, H. K., Gill, A. C., & Burdon, T. (2016). Reduced levels of dopamine and altered metabolism in brains of HPRT knock-out rats: a new rodent model of Lesch-Nyhan Disease. *Scientific Reports*, *6*, 25592. https://doi.org/10.1038/srep25592

Miller, S., & Collins, J.M. (1973). Metabolic purine pathways in the developing ovary of the housefly, *Musca domestica*. *Comparative Biochemistry and Physiology*, *44B*(4): 1153-63. https://doi.org/10.1016/0305-0491(73)90267-8.

Miñana, M.D., Portolés, M., Jordá, A., & Grisolía, S. (1984). Lesch-Nyhan syndrome, caffeine model: increase of purine and pyrimidine enzymes in rat brain. *Journal of Neurochemistry*, *43*(6): 1556-60. https://doi.org/10.1111/j.1471-4159.1984.tb06078.x.

Mizuno, T., Ohta, R., Kodama, K., Kitazumi, E., Minejima, N., Takeishi, M., & Segawa, M. (1979). Self-mutilation and sleep stage in the Lesch-Nyhan syndrome. *Brain & Development*, *1*(2):121-5. https://doi.org/10.1016/S0387-7604(79)80020-0

Mundell, S., & Kelly, E. (2011). Adenosine receptor desensitization and trafficking. *Biochimica et Biophysica Acta - Biomembranes*, *1808*(5): 1319-28. https://doi.org/10.1016/j.bbamem.2010.06.007.

Nagoshi, E. (2018). *Drosophila* models of sporadic Parkinson's disease. *International Journal of Molecular Sciences*, *19*(11), 3343. https://doi.org/10.3390/ijms19113343

Nall, A. H., Shakhmantsir, I., Cichewicz, K., Birman, S., Hirsh, J., & Sehgal, A. (2016). Caffeine promotes wakefulness via dopamine signaling in *Drosophila*. *Scientific Reports*, *6*, 20938. https://doi.org/10.1038/srep20938

Nyhan, W. L. (1997). The recognition of Lesch-Nyhan syndrome as an inborn error of purine metabolism. *Journal of Inherited Metabolic Disease*, *20*(2), 171-178. https://doi.org/10.1023/A:1005348504512

Nyhan, W L. (2000). Dopamine function in Lesch-Nyhan disease. *Environmental Health Perspectives*, *108*(Suppl 3), 409-411. https://doi.org/10.1289/ehp.00108s3409

Nyhan, William L. (2014). Nucleotide synthesis via salvage pathway. In John Wiley & Sons Ltd (Éd.), *ELS*. John Wiley & Sons, Ltd. https://doi.org/10.1002/9780470015902.a0001399.pub3

Oriel, C., & Lasko, P. (2018). Recent developments in using *Drosophila* as a model for human genetic disease. *International Journal of Molecular Sciences*, *19*(7), 2041. https://doi.org/10.3390/ijms19072041

Papanikolopoulou, K., Mudher, A., & Skoulakis, E. (2019). An assessment of the translational relevance of *Drosophila* in drug discovery. *Expert Opinion on Drug Discovery*, *14*(3), 303-313. https://doi.org/10.1080/17460441.2019.1569624

Parker, Louise, Howlett, I. C., Rusan, Z. M., & Tanouye, M. A. (2011). Seizure and epilepsy: studies of seizure disorders in *Drosophila*. *International Review of Neurobiology*, *99*, 1-21. https://doi.org/10.1016/B978-0-12-387003-2.00001-X

Pastor-Anglada, M., & Pérez-Torras, S. (2018). Emerging roles of nucleoside transporters. *Frontiers in Pharmacology*, *9*: 606. https://doi.org/10.3389/fphar.2018.00606.

Pech, U., Pooryasin, A., Birman, S., & Fiala, A. (2013). Localization of the contacts between Kenyon cells and aminergic neurons in the *Drosophila melanogaster* brain using splitGFP reconstitution: splitGFP in *Drosophila* mushroom bodies. *Journal of Comparative Neurology*, *521*(17), 3992-4026. https://doi.org/10.1002/cne.23388

Perrimon, N., Bonini, N. M., & Dhillon, P. (2016). Fruit flies on the front line: the translational impact of *Drosophila*. *Disease Models & Mechanisms*, *9*(3), 229-231. https://doi.org/10.1242/dmm.024810

Plaçais, P.-Y., de Tredern, E., Scheunemann, L., Trannoy, S., Goguel, V., Han, K.-A., Isabel, G., & Preat, T. (2017). Upregulated energy metabolism in the *Drosophila* mushroom body is the trigger for long-term memory. *Nature Communications*, *8*, 15510. https://doi.org/10.1038/ncomms15510

Pütz, S. M., Kram, J., Rauh, E., Kaiser, S., Toews, R., Lueningschroer-Wang, Y., Rieger, D., & Raabe, T. (2021) Loss of p21-activated kinase Mbt/PAK4 causes Parkinson-like phenotypes in *Drosophila*. *Disease Models and Mechanisms*, *14*(6):dmm047811. https://doi.org/10.1242/dmm.047811

Rahmani, Z., Surabhi, S., Rojo-Cortés, F., Dulac, A., Jenny, A. & Birman S. (2022). *Lamp1* deficiency enhances sensitivity to α-synuclein and oxidative stress in *Drosophila* models of Parkinson disease. *International Journal of Molecular Sciences*, *23*(21):13078. https://doi.org/10.3390/ijms232113078

Reynolds, E. R. (2018). Shortened lifespan and other age-related defects in bang sensitive mutants of *Drosophila melanogaster*. *G3 (Bethesda, Md.)*, *8*(12), 3953-3960. https://doi.org/10.1534/g3.118.200610

Riemensperger, T., Isabel, G., Coulom, H., Neuser, K., Seugnet, L., Kume, K., Iche-Torres, M., Cassar, M., Strauss, R., Preat, T., Hirsh, J., & Birman, S. (2011). Behavioral consequences of dopamine deficiency in the *Drosophila* central nervous system. *Proceedings of the National Academy of Sciences*, *108*(2), 834-839. https://doi.org/10.1073/pnas.1010930108

Riemensperger, Thomas, Issa, A.-R., Pech, U., Coulom, H., Nguyễn, M.-V., Cassar, M., Jacquet, M., Fiala, A., & Birman, S. (2013). A single dopamine pathway underlies progressive locomotor deficits in a *Drosophila* model of Parkinson disease. *Cell Reports*, *5*(4), 952-960. https://doi.org/10.1016/j.celrep.2013.10.032

Rival, T., Soustelle, L., Strambi, C., Besson, M.-T., Iché, M., & Birman, S. (2004). Decreasing glutamate buffering capacity triggers oxidative stress and neuropil degeneration in the *Drosophila* brain. *Current Biology*, *14*(7), 599-605. https://doi.org/10.1016/j.cub.2004.03.039

Ruillier, V., Tournois, J., Boissart, C., Lasbareilles, M., Mahé, G., Chatrousse, L., Cailleret, M., Peschanski, M. & Benchoua, A. (2020). Rescuing compounds for Lesch-Nyhan disease identified using stem cell-based phenotypic screening. *JCI Insight*, *5*(4):e132094. https://doi.org/10.1172/jci.insight.132094

Sankar, N., Machado, J., Abdulla, P., Hilliker, A.J., & Coe, I.R. (2002). Comparative genomic analysis of equilibrative nucleoside transporters suggests conserved protein structure despite limited sequence identity. *Nucleic Acids Research*, *30*(20): 4339-50. https://doi.org/10.1093/nar/gkf564.

Saito, Y., & Takashima, S. (2000). Neurotransmitter changes in the pathophysiology of Lesch–Nyhan syndrome. *Brain and Development*, *22*, S122-S131. https://doi.org/10.1016/S0387-7604(00)00143-1

Saras, A., & Tanouye, M. A. (2016). Seizure suppression by high temperature via cAMP modulation in *Drosophila*. *G3; Genes|Genomes|Genetics*, *6*(10), 3381-3387. https://doi.org/10.1534/g3.116.034629

Sass, J. K., Itabashi, H. H., & Dexter, R. A. (1965). Juvenile gout with brain involvement. *Archives of Neurology*, *13*(6), 639-655. https://doi.org/10.1001/archneur.1965.00470060075008

Schram, K. H. (1998). Urinary nucleosides. *Mass Spectrometry Reviews*, *17*(3), 131-251. https://doi.org/10.1002/(SICI)1098-2787(1998)17:3<131::AID-MAS1>3.0.CO;2-O

Schretlen, D. J., Ward, J., Meyer, S. M., Yun, J., Puig, J. G., Nyhan, W. L., Jinnah, H. A., & Harris, J. C. (2005). Behavioral aspects of Lesch-Nyhan disease and its variants. *Developmental Medicine and Child Neurology*, *47*(10), 673-677. https://doi.org/10.1017/S0012162205001374

Seegmiller, J. E., Rosenbloom, F. M., & Kelley, W. N. (1967). Enzyme defect associated with a sex-linked human neurological disorder and excessive purine synthesis. *Science (New York, N.Y.)*, *155*(3770), 1682-1684. https://doi.org/10.1126/science.155.3770.1682

Sinakevitch, I., Grau, Y., Strausfeld, N. J., & Birman, S. (2010). Dynamics of glutamatergic signaling in the mushroom body of young adult Drosophila. *Neural Development*, *5*(1), 10. https://doi.org/10.1186/1749-8104-5-10

Sinnett, S. E., & Brenman, J. E. (2016). The role of AMPK in *Drosophila melanogaster*. In M. D. Cordero & B. Viollet (Éds.), *AMP-activated Protein Kinase* (Vol. 107, p. 389-401). Springer International Publishing. https://doi.org/10.1007/978-3-319-43589-3_16

Smith, D. W., & Friedmann, T. (2000). Characterization of the dopamine defect in primary cultures of dopaminergic neurons from hypoxanthine phosphoribosyltransferase knockout mice. *Molecular Therapy*, *1*(5), 486-491. https://doi.org/10.1006/mthe.2000.0057

Soliman, A.M., Fathalla, A.M., & Moustafa, A.A. (2018). Adenosine role in brain functions: pathophysiological influence on Parkinson's disease and other brain disorders. *Pharmacological Reports*, *70*(4): 661-67. https://doi.org/10.1016/j.pharep.2018.02.003.

Song, J., & Tanouye, M. (2008). From bench to drug: human seizure modeling using *Drosophila*. *Progress in Neurobiology*, *84*(2), 182-191. https://doi.org/10.1016/j.pneurobio.2007.10.006

Stenesen, D., Suh, J. M., Seo, J., Yu, K., Lee, K.-S., Kim, J.-S., Min, K.-J., & Graff, J. M. (2013). Dietary adenine controls adult lifespan via adenosine nucleotide biosynthesis and AMPK, and regulates the longevity benefit of caloric restriction. *Cell metabolism*, *17*(1), 101-112. https://doi.org/10.1016/j.cmet.2012.12.006

Sun, J., Xu, A. Q., Giraud, J., Poppinga, H., Riemensperger, T., Fiala, A., & Birman, S. (2018). Neural control of startle-induced locomotion by the mushroom bodies and associated neurons in *Drosophila*. *Frontiers in Systems Neuroscience*, *12*, 6. https://doi.org/10.3389/fnsys.2018.00006

Sutcliffe, D. J., Dinasarapu, A. R., Visser, J. E., Hoed, J. D., Seifar, F., Joshi, P., Ceballos-Picot, I., Sardar, T., Hess, E. J., Sun, Y. V., Wen, Z., Zwick, M. E., & Jinnah, H. A. (2021). Induced pluripotent stem cells from subjects with Lesch-Nyhan disease. *Scientific Reports*, *11*(1):8523. https://doi.org/10.1038/s41598-021-87955-9

Tas, D., Stickley, L., Miozzo, F., Koch, R., Loncle, N., Sabado, V., Gnägi, B., & Nagoshi, E. (2018) Parallel roles of transcription factors dFOXO and FER2 in the development and maintenance of dopaminergic neurons. *PLoS Genetics*, *14*(3):e1007271. https://doi.org/10.1371/journal.pgen.1007271

Torres, R. J., Deantonio, I., Prior, C., & Puig, J. G. (2004). Adenosine transport in peripheral blood lymphocytes from Lesch-Nyhan patients. *Biochemical Journal*, *377*(Pt 3), 733-739. https://doi.org/10.1042/BJ20031035

Torres, R. J., Prior, C., & Puig, J. G. (2007). Efficacy and safety of allopurinol in patients with hypoxanthine-guanine phosphoribosyltransferase deficiency. *Metabolism*, *56*(9), 1179-1186. https://doi.org/10.1016/j.metabol.2007.04.013

Torres, R. J., & Puig, J. G. (2007). Hypoxanthine-guanine phosphoribosyltransferase (HPRT) deficiency: Lesch-Nyhan syndrome. *Orphanet Journal of Rare Diseases*, *2*(1), 48. https://doi.org/10.1186/1750-1172-2-48

Vaccaro, A., Issa, A.-R., Seugnet, L., Birman, S., & Klarsfeld, A. (2017). *Drosophila* clock is required in brain pacemaker neurons to prevent premature locomotor aging independently of its circadian function. *PLOS Genetics*, *13*(1), e1006507. https://doi.org/10.1371/journal.pgen.1006507

Visser, J. E., Bär, P. R., & Jinnah, H. A. (2000). Lesch–Nyhan disease and the basal ganglia. *Brain Research Reviews*, *32*(2–3), 449-475. https://doi.org/10.1016/S0165-0173(99)00094-6

Waddell, S. (2010). Dopamine reveals neural circuit mechanisms of fly memory. *Trends in Neurosciences*, *33*(10), 457-464. https://doi.org/10.1016/j.tins.2010.07.001

Wallrath, L. L., Burnett, J. B., & Friedman, T. B. (1990). Molecular characterization of the *Drosophila melanogaster* urate oxidase gene, an ecdysone-repressible gene expressed only in the malpighian tubules. *Molecular and Cellular Biology*, *10*(10), 5114-5127. https://doi.org/10.1128/MCB.10.10.5114

Weltha, L., Reemmer, J., & Boison, D. (2019). The role of adenosine in epilepsy. *Brain Research Bulletin*, *151*, 46-54. https://doi.org/10.1016/j.brainresbull.2018.11.008

Witteveen, J. S., Loopstok, S. R., Ballesteros, L. L., Boonstra, A., van Bakel, N. H. M., van Boekel, W. H. P., Martens, G. J. M., Visser, J. E., & Kolk, S. M. (2022). HGprt deficiency disrupts dopaminergic circuit development in a genetic mouse model of Lesch-Nyhan disease. *Cell Molecular Life Sciences*, *79*(6):341. https://doi.org/10.1007/s00018-022-04326-x.

Wong, D. F., Harris, J. C., Naidu, S., Yokoi, F., Marenco, S., Dannals, R. F., Ravert, H. T., Yaster, M., Evans, A., Rousset, O., Bryan, R. N., Gjedde, A., Kuhar, M. J., & Breese, G. R. (1996). Dopamine transporters are markedly reduced in Lesch-Nyhan disease *in vivo*. *Proceedings of the National Academy of Sciences of the United States of America*, *93*(11), 5539-5543. https://doi.org/10.1073/pnas.93.11.5539

Wu, M.N., Ho, K., Crocker, A., Yue, Z., Koh, K., & Sehgal, A. (2009). The effects of caffeine on sleep in *Drosophila* require PKA activity, but not the adenosine receptor. *The Journal of Neuroscience*, *29*(35): 11029-37. https://doi.org/10.1523/JNEUROSCI.1653-09.2009.