1 **Field phenomics reveals genetic variation for transpiration response to vapor**
2 **pressure deficit in sorghum**
3

4 Rubí Raymundo[1]*, Xu Wang[2,3], Terry Felderhoff[4], Sarah Sexton-Bowser[4], Jesse Poland[2,5],
5 Alexander E. Lipka[6], Geoffrey P. Morris[1]*
6
[1] Department of Soil and Crop Science, Colorado State University, Fort Collins, CO 80523, USA
[2] Department of Plant Pathology, Kansas State University, Manhattan, KS 66506, USA
[3] Department of Agricultural and Biological Engineering, University of Florida, IFAS Gulf Coast
Research and Education Center, Wimauma, FL 33598, USA
[4] Department of Agronomy, Kansas State University, Manhattan, KS 66506, USA
[5] Center for Desert Agriculture, King Abdullah University of Science and Technology, Thuwal
23955-6900, Kingdom of Saudi Arabia
[6] Department of Crop Sciences , University of Illinois , Urbana, IL 61801, USA
15
16 * Corresponding authors: r.raymundo@colostate.edu and Geoff.Morris@colostate.edu
17

18 Prepared for submission to *Journal of Experimental Botany*

ABSTRACT

Drought adaptation for water-limited environments relies on traits that optimize plant water budgets. Limited transpiration (LT) reduces water demand under high vapor pressure deficit (VPD) (i.e., dry air condition), conserving water for efficient use during the reproductive stage. Although studies in controlled environments report genetic variation for LT, confirming its replicability in field conditions is critical for developing water-resilient crops. Here we test the existence of genetic variation for LT in sorghum in field trials and whether canopy temperature (T_C) is a surrogate method to discriminate this trait. We phenotyped transpiration response to VPD (TR-VPD) via stomatal conductance (g_s), canopy temperature (T_C) from fixed IRT sensors (T_{Cirt}), and unoccupied aerial system thermal imagery (T_{Cimg}) in 11 genotypes. Replicability among phenomic approaches for three genotypes revealed genetic variability for TR-VPD. Genotypes BTx2752 and SC979 carry the LT trait, while genotype DKS54-00 has the non-LT trait. T_C can determine differences in TR-VPD. However, the broad sense heritability (H^2) and correlations suggest that canopy architecture and stand count hampers T_{Cirt} and T_{Cimg} measurement. Unexpectedly, observations of g_s and VPD showed non-linear patterns for genotypes with LT and non-LT traits. Our findings provide further insights into the genetics of plant water dynamics.

Keywords: Stress tolerance, crop adaptation, plant breeding, limited transpiration, stomatal conductance, canopy temperature, sorghum.

INTRODUCTION

Water scarcity is the main threat to agriculture in semi-arid regions. The increasing risk of crop failure in such areas is exacerbated by recurrent droughts, which vary over space and time (Yuan et al., 2019). Thereby ensuring demand for food and fiber in a world facing climate variability and climate change is essential. Adapting with development of climate-resilient crops is a promising strategy to cope with water scarcity (Lorite et al., 2018). However, the nature of drought adaptation traits is complex (Blum, 2011; Cooper and Messina, 2022) since transpiration is driven by many environmental factors. Hence, dissecting these traits requires understanding plant response to each variable (Figure 1A). Depending on the soil water status, plants sense water stress in leaves or roots, causing partial stomatal closure (Sampaio Filho et al., 2018). Water stress in leaves is triggered by vapor pressure deficit (VPD) (Grantz, 1990), while water stress in root tissue is stimulated by soil water deficit (Turner et al., 1985).

Limiting transpiration (LT) during periods of high VPD would cause partial stomatal closure and redistribute plant use of water within the growing season (Sinclair et al., 2005). Simulation studies revealed that unlike the non-limited transpiration (non-LT) trait, the LT trait reduces water uptake during the vegetative stage, saving soil moisture for grain filling (Figure 1B and 1C). Studies in controlled environments reported genetic variability via breakpoint (LT) and linear (non-LT) transpiration response to VPD (TR-VPD). Under the hypothesis that breakpoint thresholds are heritable, breeding programs could develop commercial hybrids and allocate them to geographies that match the hybrid's breakpoint TR-VPD to increase water productivity. However, studies were pseudo-replicated, and some studies reported the lack of reproducibility between controlled environments and field trials varied for different crops (Gilbert et al., 2011; Guiguitant et al., 2017; Shekoofa et al., 2014). Thus, replicability of these findings requires rigorous field testing since TR-VPD phenotypic characterization to-date is limited to controlled environments, and most genotypes were not tested in independent field trials (Fletcher et al., 2007; Gilbert et al., 2011; Guiguitant et al., 2017; Schoppach and Sadok, 2013; Shekoofa et al., 2014; Yang et al., 2012).

Phenomic approaches, such as whole plant transpiration and gas exchange, to discriminate TR-VPD are expensive, time-consuming, and limited for few genotypes (Kirkham, 2014). To accelerate selection, breeding programs require large-scale high-throughput plant phenotyping (HTPP) methods, which can effectively replicate the phenotype identified via traditional methods. Canopy temperature (T_C) is a proxy to estimate plant water stress (Belko et al., 2013; Jackson et al., 1981; Jones et al., 2009) due to its relationship with stomatal response (Deery et al., 2019; Leinonen and Jones, 2004), which is grounded in the theory of energy balance (Jackson et al., 1981). Studies under well-watered conditions suggest that warmer T_C and low transpiration can save soil moisture (Pinter et al., 1990). Further, it was hypothesized that genotypes with warmer T_C could yield better in drought environments (Pinter et al., 1990). While some studies suggest T_C as a reliable approach to dissecting water use traits (Anderegg et al., 2021; Belko et al., 2013; Mutava et al., 2011) other studies indicate that T_C can be an artifact of canopy architecture or micro-environmental variation (Prashar and Jones, 2014).

Sorghum is grown and adapted to water limitations. Despite this advantage, its full
potential still needs to be explored to identify traits contributing to water productivity (Borrell et
al., 2014; Vadez et al., 2014). This study aims to validate and identify the TR-VPD in field trials
in the target environment of the sorghum-producing region of the United States, predominantly
overlaying the semiarid Great Plains. We hypothesize that sorghum has a genetic variation for
TR-VPD under field conditions. Under this hypothesis, our results will be comparable to
previous studies. First, we expect low stomatal conductance (g_s) and high T_C for genotypes with
the LT trait and the opposite for genotypes with the non-LT trait (Figure 1D). Second, we expect
observations to fit breakpoint and linear TR-VPD for genotypes with LT and non-LT traits,
respectively. Our second hypothesis is that T_C is a surrogate method to discriminate TR-VPD.
Under this hypothesis, first we expect a significant negative correlation between g_s and T_C.
Second, no significant correlation between T_C and canopy architecture traits.

MATERIALS AND METHODS

Design and management of field experiments
This study tested the TR-VPD of various genotypes in Ashland Bottoms (AB), Kansas and
Greeley (GR), Colorado (Figure 2A). Based on the PRISM dataset
(https://prism.oregonstate.edu/), the maximum VPD in summer in AB is 2.5 and 3.5 in GR
respectively, corresponding to subhumid and semiarid climates (Figure 2B and 2C). Field trials
were planted in the summer seasons of 2019, 2020, and 2021 in AB and GR. The experiment
included genotypes with putative TR-VPD and commercial hybrids (Table 1). In 2019 and 2020,
the field trials comprised 10 genotypes and only 6 in 2021 (Table 1). Fewer genotypes were
planted in 2021 because the expected TR-VPD (Figure 1D) in AB19 and AB20 matched only for
genotypes DKS54-00, Tx430, BTx2752 and SC979.

Experiments were planted under a randomized complete block with four replications in
2019 and 2020. In 2021, experiments comprised two water treatments (irrigated and rainfed)
planted in a split-plot design. In these experiments, the rainfed treatment tested genotypes'
performance under drought conditions. Experiments were irrigated to isolate transpiration driven
by atmospheric VPD and avoid confounding effects due to low soil moisture (Turner et al., 1985;
Zhang et al., 2018). In AB19 and AB20, experiments were irrigated daily from full canopy cover
to flowering time using drip lines. In GR21, the experiment was irrigated every week from
planting to hard dough stage. Irrigations at this site aimed to fill soil moisture up to 100%.

Atmospheric and soil variables
In each experiment, a weather station (ATMOS 41, METER Group, Inc., USA) recorded
temperature, relative humidity, solar radiation, precipitation, wind speed, and VPD at 15-minute
intervals. In 2019, soil moisture sensors (TEROS 10, METER Group, Inc., USA) were installed
at two depths (15 cm and 30 cm) in a plot of genotype 84G62. In 2021 three soil moisture
sensors were installed on three experimental units at 30 cm. In Colorado (GR21), the seasonal
precipitation was 107 mm; in Kansas, it ranged from 338 (AB21) to 520 mm (AB19). Over three
years of assessment and across all locations, the maximum VPD ranged from 1.1 (AB) to 4.2 kPa
(GR), and soil moisture at 30 cm ranged from 0.20 m^3 m^{-3} to 0.31 m^3 m^{-3}. Experiments with the

lowest and highest VPD were AB19 and GR21, respectively (Figure 2B and 2C). General
characteristics of the field site, crop management, and environmental conditions for each season
are listed in Table S2.

Stomatal conductance phenotyping

A steady-state SC-1 leaf porometer measured (METER Group, Inc. USA) the water diffusion in
the stomatal cavity (Kirkham, 2014). Abaxial stomatal conductance (g_s) was taken on the third
fully expanded sunlit leaf on three representative plants per plot during the vegetative stage.
Information was taken between 12:00 to 16:00 hrs. Evaluations were done without cloud cover
to ensure observed transpiration was due to atmospheric VPD and not poor light quality. In
GR21, stomatal conductance was evaluated 24 hours after irrigation. Environmental conditions
(Table S2) for each evaluation were pulled from the ATMOS 41 weather station.

Canopy temperature phenotyping

Canopy temperature (T_C) was recorded via proximal sensing by infrared thermometer (IRT)
sensors fixed on the ground and remote sensing by thermal cameras carried by an unoccupied
aerial vehicle (UAV). A network of fixed IRT sensors wirelessly transmitted T_C measurements to
a data logger (Dynamax Inc.). T_C collected by IRT sensors (T_{Cirt}) was recorded in three
replications in AB19 and two replications in AB20 and GR21. Each IRT sensor has a 20-degree
field of view and covers a circle area on the target with a diameter-to-sensor distance ratio of 1:3
and a 0.5 °C accuracy over a temperature range of 0 °C to 50 °C. In this study, IRT sensors were
installed at around 1.5 m above the crop canopy, hence covering circle areas on the canopy with
a $0.5m^2$ diameter. Measurements of T_{Cirt} were analyzed from full canopy cover to pre-flowering
time to avoid any confounding effect of soil or pollen/flower temperature. For quality control,
records of T_{Cirt} (minutes) were aggregated at an hourly time step, and the time series was plotted
for visual inspection as a quality control.
A UAV high-throughput plant phenotyping (HTPP) system, including a quadcopter
(Matrice 100, DJI, Shenzhen, China) and thermal camera (VUE Pro R, FLIR, USA), were
integrated to collect thermal images for T_C extraction in AB19, AB21, and GR21 field trials.
Raw thermal images were collected under different heat conditions within the same day (i.e.,
morning, noon, and afternoon) to monitor T_C variation between genotypes (Table S2). Aerial
image overlap rate between two geospatially adjacent images was set to 80% both sequentially
and laterally to ensure optimal orthomosaic photo stitching quality. All data collection flights
were operated at 35m above ground level at 3.5 m s^{-1}. After taking off, the quadcopter will hover
at a waypoint outside the field trials for at least 2 min before real image collection, allowing the
thermal camera to self-calibrate under relatively stable ambient air condition. Multiple ground
temperature reference panels (Figure 3B) for thermal imaging calibration were placed beside the
field trials at least 20 min. before the real data collection. Ground reference temperature values
were measured and recorded in a data logger (CR3000, Campbell Scientific, USA). A semi-
automated image processing pipeline was used to generate field orthomosaic photos and trait
data extraction (Wang et al., 2020). Plot-level T_{Cimg} is the median of all T_C values extracted

within each manually-generated plot boundary. The two T_C methods were compared using the
RMSE (Wallach et al., 2014).

In-season field field phenotyping
After seedling emergence, crop establishment was scored via plant density (Figure S6). In AB19,
the plant density ranged from 19 to 25 plants m^{-2}. Even though seeding rate emergence was
similar in AB20 and GR21 (Table S2), plant density differed in both experiments. For instance,
in AB20 and GR21, plant density ranged from 24 to 35 and 13 to 19 plants m^{-2}, respectively. In
all experiments, commercial hybrids had the highest plant density; while the Tx430 genotype, an
inbred line, had the lowest plant density (Figure S6).
Flowering time was evaluated each year and determined when 50% of the two central
rows were flowering at 50%. In AB19 and AB20, one plant from each experimental unit was
harvested around flowering to estimate leaf area. In 2019, the leaf area per plant was calculated
using LI-3100C (LI-COR, Inc, USA). In AB20, the leaf area and size of each leaf across the
canopy profile were estimated using photos processed with the ImageJ program (Rueden et al.,
2017). In AB20, photographs of a plant on each experimental unit were taken before flowering.
These photographs were printed to measure the leaf angle with a protractor. The angle of the
adaxial leaf relative to the stalk was measured in the middle of the canopy. In AB19, AB21, and
GR21, the total biomass of a single plant was harvested at physiological maturity to estimate
grain weight and harvest index (HI). Similarly, in AB19 and GR21, plant height, panicle length,
and panicle exertion were measured on three plants per experimental unit.

Statistical analysis
Differences between putative traits (non-LT vs. LT) and among sorghum genotypes are expected
to occur under high VPD. For this reason, we analyzed records of g_s and T_{Cirt} between 12:00 to
16:00 hours (periods with high solar irradiation). Information on T_{Cirt} was recorded from June 23
to August 10 in AB19 and August 11, 15, 18, and 19 in AB20. Each year the variance was
analyzed with a mixed model in two stages: first, to test the size effect of the putative trait (eq.
1), and second the size effect of each genotype (eq. 2). The study conducted an additional
analysis for genotypes evaluated over all experiments (eq. 3). The models were specified as
follows:

$$Y_{ijklm} = \mu + T_i + G_j + H_k + D_l + r_m + \varepsilon_{ijklm} \quad \text{(Eq. 1)}$$

where Y_{ijklm} is the response variable in the m^{th} block in the l^{th} day in the k^{th} hour in the j^{th}
genotype with the i^{th} trait, μ is the grand mean, T_i is the fixed main effect of the i^{th} trait, G_j is the
random main effect of the j^{th} genotype, H_k is the random effect of k^{th} hour, and r_m is the random
effect of the m^{th} replicate.

$$Y_{jklm} = \mu + G_j + H_k + D_l + r_m + \varepsilon_{jklm} \quad \text{(Eq. 2)}$$

$$Y_{ijklm} = \mu + G_i + E_j + H_k + D_l + r_m + \varepsilon_{ijklm} \quad \text{(Eq. 3)}$$

where Y_{jklm} is the response variable in the m^{th} replicate in the l^{th} day in the k^{th} hour in the j^{th}
genotype, Y_{ijklm} is the response variable in the m^{th} block in the l^{th} day in the k^{th} hour in the j^{th}
environment with the i^{th} genotype, G_i is the fixed main of the i^{th} genotype, E_j is the fixed effect of
the j^{th} environment, H_k is the random effect of k^{th} hour, and r_m is the random effect of the m^{th}
replicate. A pairwise comparison (Sidak test) was performed when the F value fell below $\alpha =$
0.05 significance threshold.

Broad sense heritability (H^2) was estimated for g_s, T_{Cirt}, and T_{Cimg}. H^2 was estimated with
the lmer library (Bates et al., 2005) by fitting a random effects model which estimated variance
components for each random factor on each experiment (Eq. 4) and across experiments (Eq. 5):

$$Y_{iklm} = \mu + G_i + H_k + D_l + r_m + \varepsilon_{ijkm} \qquad \text{(Eq. 4)}$$

202
$$Y_{ijklm} = \mu + G_i + E_j + (GE)_{ij} + H_k + D_l + r_m + \varepsilon_{ijklm} \qquad \text{(Eq. 5)}$$

where Y_{iklm} is the response variable in the m^{th} replicate in the l^{th} day in the k^{th} hour with the i^{th}
genotype, Y_{ijklm} is the response variable in the m^{th} replicate in the l^{th} day in the k^{th} hour in the j^{th}
environment with the i^{th} genotype, G_i is the random effect of the i^{th} genotype, E_j is the random
effect of the j_{th} environment, $(GE)_{ij}$ is the random two way interaction of i^{th} genotype j^{th}
environment, H_k is the random effect of k^{th} hour, and r_m is the random effect of the m^{th} replicate.
The H^2 for each experiment and across experiments was calculated with Eq 6 and 7, respectively.
The variance for environmental effects H and D were disregarded in the analysis since they do
not contribute to selecting the target phenotype.

$$H^2 = \frac{\bar{\sigma}_G^2}{\bar{\sigma}_G^2 + \frac{\bar{\sigma}_\varepsilon^2}{n_r}} \qquad \text{(Eq. 6)}$$

$$H^2 = \frac{\bar{\sigma}_G^2}{\bar{\sigma}_G^2 + \frac{\bar{\sigma}_{G:E}^2}{n_E} + \frac{\bar{\sigma}_\varepsilon^2}{n_E \times n_r}} \qquad \text{(Eq.7)}$$

213 where: σ_G^2, $\sigma_{G:E}^2$, and σ_ϵ^2 represent variance components of G, $G\times E$, and error, respectively.
214 While, n_E is the number of environments, and n_r is the number of replicates.

215 Information on g_s, T_{Cirt}, and canopy–air difference was paired with corresponding VPD.
216 Next, g_s was analyzed via a linear (Eq. 8), segmented (Eq. 9 and Eq.10), and asymptotic (non-
217 linear, Eq.11) regression analysis.
218

$$Y = \beta_0 + \beta_1 VPD + \varepsilon \qquad \text{(Eq. 8)}$$

$$\text{If } VPD <= BP: \qquad Y = \beta_0 + \beta_1 VPD + \varepsilon \qquad \text{(Eq. 9)}$$

$$\text{If } VPD > BP: \qquad Y = \beta_0 + \beta_1 BP + (\beta_1 + \beta_2)(VPD - BP) + \varepsilon \qquad \text{(Eq. 10)}$$

$\qquad Y = a + (a - b)e^{-cVPD} \qquad\qquad\qquad\qquad\qquad\qquad \text{(Eq. 11)}$

where Y represents g_s, BP is the VPD breakpoint, and β_0 is the intercept, β_1 is the slope, and β_2 is
the second slope, a is the plateau (maximum attainable value) for g_s, b is the initial g_s when VPD
is zero, c is the inflexion point or the relative rate of increase for VPD when g_s increases. The
reason for testing different regressions was to identify the best model that fitted the observed
data. Linear regressions were performed using lm library (Bates et al., 2015), the breakpoint
regression was analyzed with the library segmented (Muggeo, 2016), non-linear regressions were
analyzed with libraries nlme, drc, and aomisc (Ritz et al., 2015). The dependence of g_s on VPD
was tested via 1000 permutations for linear and breakpoint regressions. To determine this
dependence for each genotype, the distribution of the regression coefficient for these
permutations was plotted and contrasted against the observed coefficients, β_1 for linear
regression; β_1, β_2, and BP for segmented regressions. The likelihood-ratio test (library lmtest),
and residual standard error (rse) indicated the accuracy of the best regression model
(Archontoulis and Miguez, 2015; Kuznetsova et al., 2017).

$\qquad$ The effect of canopy architecture traits (i.e., leaf area and leaf size) on transpiration VPD
response were visualized using a principal component analysis (PCA) and a correlation analysis.
The PCA was estimated with the R prcomp function ("R Core Team," 2017) and conducted only
for AB20 since most canopy architecture traits were evaluated in this experiment. In this
analysis, the leaf area represents the maximum leaf area per plant, which occurs around
flowering time. The leaf size corresponds to the largest leaf across the canopy profile (Figure
S3). The leaf angle is at the attachment between the adaxial leaf and stalk in the central part of
the canopy. Plant density represents the number of plants scored after seedling emergence
(Figure S6). A correlation analysis was conducted for AB19 and GR21. Grain weight and harvest
index for experiments in AB19, AB21, and GR21 were analyzed to characterize agronomic traits
of potential donors of putative LT trait.

RESULTS

Classification of genotypes for their TR-VPD
To test if VPD during the experiment's growing seasons represented a typical VPD range, we
compared the average daily maximum VPD over 22 years (2000-2021) against the daily
maximum VPD recorded on each experiment (Figure 2B and 2C). Results indicate that growing
seasons AB19 and AB20 underrepresented average VPD in Kansas; while the growing season in
GR21 represented a regular season. In AB19 and AB20, the VPD ranged between 0.2 to 1.9 kPa
during the evaluation period (canopy closure to pre-flowering time, Figures 2B). Otherwise in
GR21 the VPD ranged between 2.5 to 4.2 kPa.

253   To determine the existence of genetic variability for TR-VPD in field conditions, we
254 tested i) the putative non-LT versus LT classification and ii) the genotype classification in each
255 experiment (Table 1). To confirm this classification, significance was expected for the fixed
256 effect and differences among mean groups in post hoc analysis. Specifically, higher g_s for the
257 putative non-LT trait and higher T_{Cirt} and T_{Cimg} for the LT trait (Figure 1D). Results in all
258 experiments indicated no significance for the trait effect for g_s, T_{Cirt}, and T_{Cimg} (Figure 4, upper
259 panels). However, as expected, the putative non-LT group exhibited slightly high g_s (0.06 mol m$^-$
260 ^{2} s^{-1}) and lower T_{Cirt} (-0.1 C), as indicated in Table S3.
261   The analysis of variance for each experiment where genotype was the fixed effect
262 indicated differences ($p < 0.001$) for g_s, T_{Cirt}, and T_{Cimg} (Figure 4). In AB19 (Figure 4A, lower
263 panel), as expected, genotypes with putative LT (BTx2752, 0.62 mol m^{-2} s^{-1}) and non-LT
264 (DKS54-00, 0.77 mol m^{-2} s^{-1}) differed significantly. In this experiment, a genotype classified as
265 LT (SC803) trait unexpectedly exhibited higher g_s. In AB20 (Figure 4B, lower panel), the mean
266 comparison indicated three groups that differed in g_s. The first group comprised a genotype with
267 the LT trait SC979, which had the lowest g_s (0.73 mol m^{-2} s^{-1}). The second group included
268 genotypes with LT (Macia, BTx2752, and BTx2363). The third group had genotypes with non-
269 LT (DKS54-00, Tx430) and LT (Tx7000, BTx642, and SC803) traits, which had the highest g_s.
270 In GR21 (Figure 4C, lower panel), the genotype Tx430 with the non-LT trait unexpectedly
271 exhibited the lowest g_s (Tx430, 0.72 mol m^{-2} s^{-1}). As expected, the genotype DKS54-00 with the
272 non-LT trait had the highest (0.90 mol m^{-2} s^{-1}) g_s. Despite genotypes with the LT trait, BTx2752,
273 and SC979, presenting lower g_s than DKS54-00, they were not significantly different from this
274 genotype.
275   In AB19 (Figure 4D, lower panel), phenotypes identified via T_{Cirt} confirmed that six of
276 nine genotypes matched the expectation (Table 1). Genotypes with the LT trait SC979 (29.9 °C),
277 BTx2752 (29.9 °C), Macia (29.9 °C), and SC803 (30.0 °C) had higher T_{Cirt} and differed from
278 genotypes with the non-LT trait DKS54-00 (29.5 °C) and Tx430 (29.6 °C). In AB20 (Figure 4E,
279 lower panel) T_{Cirt} confirmed the classification for Tx7000 (LT), SC979 (LT), BTx642 (LT),
280 BTx2752 (LT), and Tx430 (non-LT). But, it contradicted the classification for genotypes SC803
281 (LT), Macia (LT), BTx623 (LT), and DKS54-00 (non-LT) (Figure 1D). Genotypes with the LT
282 (BTx2752, 29.0 °C) and non-LT (Tx430, 28.1 °C) traits exhibited the highest and lowest T_{Cirt},
283 respectively. In GR21, all genotypes matched the expectation. The lowest and highest canopy
284 temperature corresponded to genotypes ADV2275 and BTx2752, respectively. In all
285 experiments, T_{Cirt} was replicable only for genotypes with the putative LT trait BTx2752 and
286 SC979.
287   As expected, in AB21, T_{Cimg} for genotypes with the non-putative-LT trait (DKS54-00,
288 DKS28-05, and SC1345) was lower overall than for genotypes with the putative LT trait (SC803,
289 Macia, BTx623, BTx2752). Discrepant results were obtained for genotype SC979, for which
290 high T_{Cimg} was expected but had low T_{Cimg}, and for genotype Tx430 for which low T_{Cimg} was
291 expected but exhibited high T_{Cimg}. In AB21 (Figure 4H, lower panel) and GR21 (Figure 4I, lower
292 panel), genotypes SC979 and BTx2752, classified as genotypes with the LT trait, exhibited

significantly higher TCimg than genotype DKS54-00, classified as a genotype with the non-LT

trait. The only genotype that contradicted its non-LT classification in both experiments was

Tx430, which had the highest TCimg.

Over three years of field experiments g_s and T_{Cirt} significantly differed ($p < 0.001$), and

the covariate effect of VPD on g_s was non-significant (Table S5). The pairwise comparison for g_s

revealed that genotypes DKS54-00 (0.81 mol m^{-2} s^{-1}), BTx2752 (0.78 mol m^{-2} s^{-1}), and SC979

(0.72 mol m^{-2} s^{-1}) belonged to different groups (Table S5). Results of g_s for Tx430 (non-LT)

were statistically similar to genotypes BTx2752 (LT) and SC979 (LT) (Table S5). The pairwise

comparison for T_{Cirt} showed two groups. Genotype DKS54-00 with the non-LT trait had lower

T_{Cirt} (28.2 °C) than genotype BTx2752 with the LT trait (28.6 °C). The analysis of T_{Cirt} for four

genotypes over three years matched with the TR-VPD classification (Table S5).

To test that different phenomic approaches capture the biology of TR-VPD, we estimated

the broad sense of heritability (H^2) and expected similar H^2 for g_s, T_{Cirt}, and T_{Cimg}. However, H^2

differed among phenomic approaches in each experiment (Figure 5). In AB19, the highest and

lowest H^2 corresponded for g_s (0.5) and T_{Cimg} (0.06). Estimates of H^2 Across all experiments for

each method indicate a greater H^2 for g_s (0.4), followed by T_{Cimg} (0.3) and T_{Cirt} (0.2).

Genotypic variation for stomatal response to VPD

To test that transpiration is driven by VPD and not due to soil water deficit, we compared

stomatal response in rainfed and irrigated treatments. This comparison was conducted on a day

when the VPD ranged from 3.2 to 4.0 kPa in GR21 (Figure 6A). In this comparison, we expected

a significant difference ($p < 0.05$) between treatments, and observations confirmed this

expectation. On average, g_s under rainfed treatments was 50% lower than in irrigated treatments.

In this VPD range, the regression slope for g_s in the irrigated treatment was close to zero (0.006

mol m^{-2} s^{-1} kPa); in the rainfed treatment, g_s declined (-0.21 mol m^{-2} s^{-1} kPa) as VPD increased.

Next, we conducted a regression analysis for each genotype. The stomatal response to VPD was

analyzed via linear, segmented, and non-linear models. In all experiments, g_s ranged between 0.3

to 1.2 mol m^{-2} s^{-1} and VPD between 1 to 4.5 kPa (Figure 6). The mean g_s conductance of 0.79,

0.86, 0.82, and 0.82 mol m^{-2} s^{-1} corresponded to experiments AB19, AB20, AB21, and GR21,

respectively. Unexpectedly, high VPD did not increase the mean g_s for genotypes with the

putative non-LT trait (DKS54-00 and Tx430).

Linear regression for genotypes 84G62 (NN), DKS54-00 (non-LT), BTx2752 (LT), and

SC979 (LT) indicated a positive slope but an unexpected negative one for genotype Tx430 (non-

LT) (Figure 6B, Table 3). Segmented (breakpoint) regression for all genotypes suggested a

restriction in g_s between 1.3 to 2 kPa (Figure 6C). Genotype Tx430, which has a putative non-LT

trait, had the lowest breakpoint and the most negative second slope. Breakpoint patterns were

similar for commercial hybrids 84G62 (NN) and DKS54-00 (putative non-LT) indicating a

negative value for the second slope. Genotypes with the putative LT trait SC979 and BTx2752

exhibited similar patterns, revealing positive values for the first and second slope (Figure 6C).

Non-linear regressions revealed an asymptotic curve fitting the observed patterns for all

genotypes 84G62, DKS54-00, BTx2752, and SC979, except Tx430. A non-linear regression

revealed that differences on transpiration under high VPD are represented via the maximum
plateau and the inflection point. Parameters for each regression and genotypes are indicated on
Table 3.
To test that the g_s is dependent on VPD in linear regression, we compared the observed
versus permuted slopes (Figure S3). For genotypes with the non-LT trait DKS54-00 and Tx430,
we expected the observed slope to be positive and significantly higher than permuted slopes. For
genotypes with the LT trait BTx2752 and SC979, which restrict gas exchange at high VPD, we
expected the observed slope to be non-significantly different from the permuted slopes.
However, observations contradicted our expectation, indicating that g_s and VPD are independent
for DKS54-00 ($\beta_1 = 0$, $p = 0.1$), while g_s depends on VPD, positive slope, for genotypes
BTx2752 and SC979 ($\beta_1 = 0$, $p < 0.05$). Similarly, a permutation analysis for a breakpoint
regression revealed that the regressed variable (g_s) is independent of the regressor (VPD, Figure
S4). To determine the goodness of fit for different regression models, we used the likelihood
ratio test. In this analysis, we expected significance for linear regression for genotypes with the
non-LT, whereas significance for a breakpoint regression for genotypes with the LT trait.
Unexpectedly, breakpoint and non-linear regressions fitted the observed data ($p < 0.001$) for a
genotype with non-LT trait DKS54-00 (Figure 6). Otherwise, for a genotype with non-LT trait
Tx430, the linear and breakpoint regression revealed no significant difference ($p > 0.05$).
Similarly, all regressions were not significantly different for genotypes with LT trait BTx2752
and SC979 ($p > 0.05$, Figure 6).

Canopy temperature (T_C) as surrogate method to discriminate TR-VPD
To test if T_C is a surrogate method of g_s, and not an artifact of canopy architecture trais, a PCA
was conducted for AB20 and a correlation analysis for AB19 and GR21. In AB20, PC1, PC2 and
PC3 represented 38.9, 23.9 and 22.1% of the total variability in the data (Figure 7). PCA1 had a
moderately positive loading for leaf area (0.4) and leaf size (0.4) and a negative loading for plant
density (-0.4) and T_{Cirt} (-0.4). PCA2 had a positive and negative loading with leaf size (0.5) and
g_s (-0.6), respectively. PCA3 is dominated by negative loading of the number of leaves (-0.7) and
positive loading for T_{Cirt} (0.4). The biplot for PC1 versus PC2 and PC2 versus PC3 revealed that
T_{Cirt} and g_s are in distinct quadrants suggesting a moderate negative correlation (Figure 7SA) for
these phenotypes. Similarly, in GR21, the correlation between g_s versus T_{Cimg} was negative and
significant ($r = -0.9$, $p < 0.05$). A biplot PC2 versus PC3 showed a substantial correlation
between T_{Cirt} and the number of leaves. These results indicate that T_{Cirt} is a good proxy of g_s, but
canopy architecture can affect T_C. Otherwise, biplots suggested a strong correlation among
canopy architecture traits. For instance, leaf area negatively correlated with the number of leaves
(PC1 versus PC2) and leaf angle (PC2 versus PC3).
To test if TR-VPD phenotype is an artifact of stand count, we conducted correlations
between g_s, T_{Cirt}, T_{Cimg}, and plant density (Figure S7). In AB20, the correlation was close to zero
for plant density and T_{Cirt}, while a negative correlation (-0.5) between plant density and g_s was
non-significant ($p > 0.05$). Significant negative correlations between plant density and T_{Cimg}
(Figure S7B and S7C) were obtained in AB19 ($r = -0.8$, $p < 0.01$) and GR21 ($r = 0.9$, $p < 0.05$).

373 Likewise, in GR21 (Figure S7C), the correlation between plant density and g_s was significant (r
374 $= 0.9$, $p < 0.01$). Overall, results suggest that stand count is the main trait that can hamper the
375 effectiveness of any phenomic approach when dissecting TR-VPD.

376 **DISCUSSION**

377 Testing the existence of genetic variability for the putative LT trait via different phenomic
378 approaches is pivotal when deciding to include this trait in a breeding pipeline. This is the first
379 study that phenotyped the TR-VPD in field trials in sorghum production regions of the United
380 States. Our findings i) help to better understand the challenges when testing the replicability of
381 the trait via different phenomic approaches, ii) discuss the best representation of genetic diversity
382 by comparing different regression models, iii) review the underlying physiological mechanism of
383 the putative LT trait and iv) outline future steps for crop breeding when using the LT to deliver
384 water efficient sorghum hybrids.

385 *A phenotype can be similar or vary among phenomic approaches*
386 Our study confirmed the TR-VPD classification via three phenomic approaches for three out of
387 eleven genotypes (Figure 4). Results for genotypes SC979 (LT), BTx2752 (LT), and DKS54-00
388 (non-LT) corresponded with prior references (Gholipoor et al., 2010; Shekoofa et al., 2014). A
389 PCA (Figure 7) and a negative correlation between g_s and T_C, although non-significant, to some
390 extent, suggested T_C as a surrogate phenotype to discriminate differences in transpiration (Figure
391 7, Figure S7). The effectiveness of T_C in determining variability for LT transpiration has been
392 demonstrated by comparing T_C with whole-plant transpiration in controlled environments (Belko
393 et al., 2013).

394   For other genotypes, the g_s phenotype contradicted prior classification (Figure 4). For
395 instance, unexpected high g_s for genotypes with the putative LT trait SC803, Tx7000, and
396 BTx642 (Table 1) suggest they had the non-LT trait. Discrepant results were also reported for
397 lentils where a genotype with the putative LT in the greenhouse displayed a non-LT trait in field
398 trials (Guiguitant et al., 2017). These observations imply that growth chamber experiments do
399 not fully represent field conditions. Phenotyping for TR-VPD in growth chambers is limited to
400 early growth stages and environmental conditions are more erratic in field settings. Indeed, prior
401 studies suggested that early growth stages in field conditions (Shekoofa et al., 2014) and
402 temperatures beyond 30 °C (Riar et al., 2015) suppress the expression of the LT trait.

403   The phenotype identified with each phenomic approach varied for most genotypes
404 (Figure 4). These differences can be attributed to the sample size utilized in each method (Figure
405 4), variability in canopy architecture, and stand count. The interaction of canopy architechture
406 traits with wind speed influences convective heat transfer and T_C (Gates, 1980; Leigh et al.,
407 2017; Melcher et al., 1994). Canopy architecture traits can be more important determinants of T_C
408 than physiological traits (Still et al., 2021; Woods et al., 2018) since canopy architechture traits
409 and wind speed influence convective heat transfer and T_C. Genotypes with an acute leaf angle
410 likely disrupted air movements through the canopy, leading to high T_{Cirt}, as observed in genotype
411 BTx642 (Figure 7A). Soil exposure in plots of genotypes with fewer leaves would have reflected
412 solar energy, contributing to increasing air temperature and T_C (Figure 7B). This would explain

the unexpectedly high T_{Cirt} in AB19 for genotypes DKS28-00 and SC1345 with the putative non-
LT trait and low leaf area (Table S6).
Microenvironmental conditions created by low plant density and high tillering can
override the phenotypic expression of a target trait (Jones, 2007). The phenotype identified via g_s
for genotype Tx430 shifted from non-LT (Figure 4A, 4B) to LT (Figure 4C) when the plant
density was lower than 13 pl m^{-2} (Figure S8A). Although the experiment was well-irrigated, at
low plant density, it is possible that soil evaporation surpassed plant transpiration. High tillering
(Figure S8B) likely exacerbated water demand, promoting water competition among culms
(Borrell et al., 2014), leading to low g_s and high T_{Cimg} (Figures 4H and 4I).
T_{Cimg}, an HTPP approach, can accelerate breeding selection by bringing higher precision
(Deery et al., 2019), if such an approach can capture the nature of the target trait. Nevertheless,
H^2 for T_{Cimg} was close to zero and lower than H^2 for g_s in AB19. While the H^2 for g_s (0.16-0.55)
in our study aligned with an investigation for cotton (0.16 to 0.44) (Percy et al., 1996), the H^2 for
T_{Cimg} ranged from 0.05 to 05, which are lower than the values reported for wheat (0.3 to 0.8)
(Anderegg et al., 2021; Deery et al., 2019). Different H^2 for T_{Cimg} (Figure 5) among experiments
indicates that this approach is prone to microenvironmental conditions likely caused by the
variability of canopy architecture traits (Figure 7 and S7A). Additionally, stand count can
override the expression of the TR-VPD (Figure S7B and S7B). Overall, using T_{Cimg} to screen
large breeding populations would require considering the number of leaves and stand count as
covariate effects.

Non-linear function models better represent genetic variability for TR-VPD
Genetic variability identified via different phenomic approaches in this study provides candidate
donor lines to breed sorghum yields for water deficit conditions (Figure 4). Our study has shown
that genetic variability occurs under low and high (<2 kPa) VPD (Figure 6D). Genetic variability
for TR-VPD has been proposed as differences in slopes and restriction (breakpoint) in gas
exchange (Gholipoor et al., 2010; Shekoofa et al., 2014). In our study, observations indicated
that no genotypes with the non-LT trait increased g_s (Figure S3). Evidence of whether observed
data fit a linear or segmented regression was never shown or discussed in prior studies. For
instance, observed data for outdoor pots suggested asymptotic patterns for chickpeas with LT
and non-LT traits, but the study reported a breakpoint and LT and linear response (Zaman-Allah
et al., 2011).
In our research, stomatal response to VPD fit different regressions (Figure 6), indicating
the breakpoint or asymptotic as the best model for a genotype with non-LT trait. Although
disparities between a breakpoint (Figure 6C) and an asymptotic model (Figure 6D) seem to be
negligible, independent data support that at higher VPD, g_s will reach its maximum value and
remain constant (Figure 6A). Our findings align with studies where transpiration under well-
watered conditions and high VPD remained constant. A wheat study reported breakpoints
response for 100% of commercial varieties tested in growth chambers (Schoppach et al., 2017).
For woody and herbaceous plants, transpiration under controlled environments reached a
maximum plateau at a VPD of 2.5 kPa (Turner et al., 1984). An asymptotic function better

portrays the stomatal response to VPD, as this non-linear model has a biological meaning and is
utilized to describe photosynthesis, light intensity, and light interception (Archontoulis and
Miguez, 2015). The correct mathematical representation of TR-VPD has a significant
implication when modeling the effect of the LT at crop system scales. Simulation studies using
segmented and nonlinear functions are most likely overestimating the impact of the LT trait on
harvested yield.

The underlying mechanism of TR-VPD
The underlying physiological mechanism of TR-VPD remains enigmatic. The LT trait has been
associated with aquaporin inhibition (Maurel et al., 2016) and low hydraulic conductivity
(Choudhary et al., 2013b) that causes partial stomatal closure. In arabidopsis and angiosperms,
stomatal response to VPD is controlled by hydropassive and hydroactive processes (Merilo et al.,
2018; Wang et al., 2001). Nevertheless, studies in soybean and peanut claim that LT is
modulated via a hydropassive mechanism (Sinclair et al., 2017). While the hydropassive
response involves changes in the turgor cell, the hydroactive mechanism is modulated via
metabolic signaling (Merilo et al., 2018; Wang et al., 2001). Studies reported leaf-derived ABA
when plants are exposed to high VPD (Hu et al., 2016; Jalakas et al., 2021), although findings
indicate that *Open Stomata 1* (*OST1*) kinase rather than ABA controls stomatal response to
humidity (Merilo et al., 2018). Otherwise, differences in leaf hydraulic conductivity reported for
genotypes SC1205 (dwarf, non-LT) and SC15 (tall, putative LT) (Choudhary et al., 2013b;
Ocheltree et al., 2013) can result from contrasting hydraulic anatomy (Du et al., 2020) and plant
architecture. Indeed, under well-water conditions, a study reported differences in hydraulic
anatomy (xylem diameter) for a short (ICSSH58, 1.5 m) and tall (ICSV25280, 1.7 m) sorghum
(Guha et al., 2018).
476   Discrepancies in TR-VPD between non-LT and LT traits are hypothetically due to
stomatal openness (Sinclair et al., 2017) with expected open stomata for the genotype with the
non-LT trait and partial stomatal closure for a genotype with the LT trait (Figure 8A).
Determining whether partial stomatal closure is controlled by hydropassive or hydroactive
mechanisms requires testing the expression of metabolic signals (Figure 8B). Otherwise,
asymptotic patterns of TR-VPD (Figure 6D) suggest that the stomata remain open at high VPD
for genotypes with LT and non-LT traits (Figure 8C). Then, differences in gas exchange are
hypothetically due to variations in hydraulic anatomy traits such as variability in stomatal density
(Bheemanahalli et al., 2021), xylem diameter, or boundary layer thickness (Figure 8D). From a
breeding perspective, a hydroactive mechanism would allow identifying known genes such as
OST1 or *slow anion channel 1* (*SLAC1*), which are ABA-dependent or -independent (Merilo et
al., 2018). Meanwhile, the hydropassive mechanism implies that the aquaporin response is
independent of any signaling pathway and unknown genes. Testing these hypotheses in
identified lines (DKS54-00, BTX2752, and SC979) can elucidate mechanisms underlying the
TR-VPD. Further quantifying the carbon assimilation trade-offs of the LT trait in near-isogenic
lines would indicate the impact of this trait on drought-prone regions. This knowledge can guide
breeding programs to consider different options when designing a breeding pipeline.

Future steps
The results of this study represent TR-VPD when soil water is not limited in the soil profile.
However, drought-prone regions are subjected to simultaneous high VPD and soil water deficit.
Results under a rainfed treatment and soil water deficit in a semiarid environment revealed a
decline in g_s as VPD increases (Figure 6A), suggesting that soil water deficit and high VPD lead
to stomatal closure to avoid dehydration and physiological damage (Oren et al., 1999). So far,
the confounding effect of VPD and soil moisture has been ignored in studies of LT. Then it
would be worthwhile to determine whether the effect of VPD on stomatal closure prevails over
the impact of soil water deficit on stomatal closure or vice versa. Genotypes with putative LT
traits identified in this study correspond to full-season backgrounds. A comparison of yield
components under well-watered and rainfed treatments in a semiarid environment (GR21)
demonstrated that early flowering time overrode the effect of the putative LT trait, either in
irrigated or rainfed treatments (Figure S10). In silico simulations showed that the LT trait can
increase yields by more than 5% in western regions of the sorghum belt where medium and
short-season hybrids are planted. Then it would be needed to identify and introgress the LT trait
in short and medium-season backgrounds. Developing a mapping population and near-isogenic
lines with donors SC979, BTx2752 would allow identifying QTLs via T_{Cimg}; nevertheless,
confounding effects detailed in this study need to be considered when using this approach.

ACKNOWLEDGMENT

This study was supported by funding from the Foundation for Food and Agriculture Research - Seeding Solution "CA18-SS-0000000094 – Bridging the Genome-to-Phenome Breeding Gap for Water-Efficient Crop Yields (G2P Bridge)" to G.P.M. and S.S.B.; the Kansas Department of Agriculture "Collaborative Sorghum Investment Program Water Optimized Sorghum for Kansas" to G.P.M; and the Kansas Grain Sorghum Commission.

AUTHOR CONTRIBUTIONS

G.P.M., S.S.B., T.F., and R.R. contributed to the conception and design of the work. T.F. and R.R. collected field experimental data. X.W was in charge of the UAS imaging. R.R. and X.W. conducted data analysis, interpretation, and drafting of the article. G.P.M., S.S.B., T.F., R.R., X.W., J.P., A.E.L. contributed to the final manuscript.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

REFERENCES

Anderegg, J., Aasen, H., Perich, G., Roth, L., Walter, A., Hund, A., 2021. Temporal trends in canopy temperature and greenness are potential indicators of late-season drought avoidance and functional stay-green in wheat. Field Crops Research 274, 108311. https://doi.org/10.1016/j.fcr.2021.108311

Archontoulis, S.V., Miguez, F.E., 2015. Nonlinear Regression Models and Applications in Agricultural Research. Agronomy Journal 107, 786–798. https://doi.org/10.2134/agronj2012.0506

Bates, D., Maechler, M., Bolker, B.M., Walker, S.C., 2015. Fitting Linear Mixed-Effects Models Using lme4. J. Stat. Softw. 67, 1–48. https://doi.org/10.18637/jss.v067.i01

Belko, N., Zaman-Allah, M., Diop, N.N., Cisse, N., Zombre, G., Ehlers, J.D., Vadez, V., 2013. Restriction of transpiration rate under high vapour pressure deficit and non-limiting water conditions is important for terminal drought tolerance in cowpea. Plant Biol (Stuttg) 15, 304–316. https://doi.org/10.1111/j.1438-8677.2012.00642.x

Bheemanahalli, R., Wang, C., Bashir, E., Chiluwal, A., Pokharel, M., Perumal, R., Moghimi, N., Ostmeyer, T., Caragea, D., Jagadish, S.V.K., 2021. Classical phenotyping and deep learning concur on genetic control of stomatal density and area in sorghum. Plant Physiology 186, 1562–1579. https://doi.org/10.1093/plphys/kiab174

Blum, A., 2011. Drought Resistance and Its Improvement, Plant Breeding for Water-Limited Environments. Springer-Verlag Berlin, Berlin. https://doi.org/10.1007/978-1-4419-7491-4_3

Borrell, A.K., Oosterom, E.J. van, Mullet, J.E., George-Jaeggli, B., Jordan, D.R., Klein, P.E., Hammer, G.L., 2014. Stay-green alleles individually enhance grain yield in sorghum under drought by modifying canopy development and water uptake patterns. New Phytologist 203, 817–830. https://doi.org/10.1111/nph.12869

Choudhary, S., Mutava, R.N., Shekoofa, A., Sinclair, T.R., Prasad, P.V.V., 2013a. Is the stay-green trait in sorghum a result of transpiration sensitivity to either soil drying or vapor pressure deficit? Crop Science 53, 2129. https://doi.org/10.2135/cropsci2013.01.0043

Choudhary, S., Sinclair, T.R., Prasad, P.V.V., 2013b. Hydraulic conductance of intact plants of two contrasting sorghum lines, SC15 and SC1205. FUNCTIONAL PLANT BIOLOGY 40, 730–738. https://doi.org/10.1071/FP12338

Cooper, M., Messina, C.D., 2022. Breeding crops for drought-affected environments and improved climate resilience. The Plant Cell koac321. https://doi.org/10.1093/plcell/koac321

Deery, D.M., Rebetzke, G.J., Jimenez-Berni, J.A., Bovill, W.D., James, R.A., Condon, A.G., Furbank, R.T., Chapman, S.C., Fischer, R.A., 2019. Evaluation of the Phenotypic Repeatability of Canopy Temperature in Wheat Using Continuous-Terrestrial and Airborne Measurements. Frontiers in Plant Science 10.

Du, Q., Jiao, X., Song, X., Zhang, J., Bai, P., Ding, J., Li, J., 2020. The Response of Water Dynamics to Long-Term High Vapor Pressure Deficit Is Mediated by Anatomical Adaptations in Plants. Frontiers in Plant Science 11.

Fletcher, A.L., Sinclair, T.R., Allen, L.H., 2007. Transpiration responses to vapor pressure deficit in well watered 'slow-wilting' and commercial soybean. Environmental and Experimental Botany 61, 145–151. https://doi.org/10.1016/j.envexpbot.2007.05.004

Gates, D.M., 1980. Energy Budgets of Plants, in: Gates, D.M. (Ed.), Biophysical Ecology, Springer Advanced Texts in Life Sciences. Springer, New York, NY, pp. 345–381.

573 https://doi.org/10.1007/978-1-4612-6024-0_11

574 Gholipoor, M., Prasad, P.V.V., Mutava, R.N., Sinclair, T.R., 2010. Genetic variability of
575  transpiration response to vapor pressure deficit among sorghum genotypes. Field Crops
576  Research 119, 85–90. https://doi.org/10.1016/j.fcr.2010.06.018

577 Gilbert, M.E., Holbrook, N.M., Zwieniecki, M.A., Sadok, W., Sinclair, T.R., 2011. Field
578  confirmation of genetic variation in soybean transpiration response to vapor pressure
579  deficit and photosynthetic compensation. Field Crops Research 124, 85–92.
580  https://doi.org/10.1016/j.fcr.2011.06.011

581 Grantz, D.A., 1990. Plant response to atmospheric humidity. Plant, Cell & Environment 13, 667–
582  679. https://doi.org/10.1111/j.1365-3040.1990.tb01082.x

583 Guha, A., Chhajed, S.S., Choudhary, S., Sunny, R., Jansen, S., Barua, D., 2018. Hydraulic
584  anatomy affects genotypic variation in plant water use and shows differential organ
585  specific plasticity to drought in Sorghum bicolor. Environmental and Experimental
586  Botany 156, 25–37. https://doi.org/10.1016/j.envexpbot.2018.08.025

587 Guiguitant, J., Marrou, H., Vadez, V., Gupta, P., Kumar, S., Soltani, A., Sinclair, T.R., Ghanem,
588  M.E., 2017. Relevance of limited-transpiration trait for lentil (Lens culinaris Medik.) in
589  South Asia. Field Crops Research 209, 96–107. https://doi.org/10.1016/j.fcr.2017.04.013

590 Hu, B., Cao, J., Ge, K., Li, L., 2016. The site of water stress governs the pattern of ABA
591  synthesis and transport in peanut. Sci Rep 6, 32143. https://doi.org/10.1038/srep32143

592 Jackson, R.D., Idso, S.B., Reginato, R.J., Pinter Jr., P.J., 1981. Canopy temperature as a crop
593  water stress indicator. Water Resources Research 17, 1133–1138.
594  https://doi.org/10.1029/WR017i004p01133

595 Jalakas, P., Takahashi, Y., Waadt, R., Schroeder, J.I., Merilo, E., 2021. Molecular mechanisms
596  of stomatal closure in response to rising vapour pressure deficit. New Phytologist 232,
597  468–475. https://doi.org/10.1111/nph.17592

598 Jones, H.G., 2007. Monitoring plant and soil water status: established and novel methods
599  revisited and their relevance to studies of drought tolerance. Journal of Experimental
600  Botany 58, 119–130. https://doi.org/10.1093/jxb/erl118

601 Jones, H.G., Serraj, R., Loveys, B.R., Xiong, L., Wheaton, A., Price, A.H., 2009. Thermal
602  infrared imaging of crop canopies for the remote diagnosis and quantification of plant
603  responses to water stress in the field. Funct Plant Biol 36, 978–989.
604  https://doi.org/10.1071/FP09123

605 Kirkham, M.B., 2014. Chapter 24 - Stomatal Anatomy and Stomatal Resistance, in: Kirkham,
606  M.B. (Ed.), Principles of Soil and Plant Water Relations (Second Edition). Academic
607  Press, Boston, pp. 431–451. https://doi.org/10.1016/B978-0-12-420022-7.00024-0

608 Kuznetsova, A., Brockhoff, P.B., Christensen, R.H.B., 2017. lmerTest Package: Tests in Linear
609  Mixed Effects Models. Journal of Statistical Software 82, 1–26.
610  https://doi.org/10.18637/jss.v082.i13

611 Leigh, A., Sevanto, S., Close, J. d., Nicotra, A. b., 2017. The influence of leaf size and shape on
612  leaf thermal dynamics: does theory hold up under natural conditions? Plant, Cell &
613  Environment 40, 237–248. https://doi.org/10.1111/pce.12857

614 Leinonen, I., Jones, H.G., 2004. Combining thermal and visible imagery for estimating canopy
615  temperature and identifying plant stress. Journal of Experimental Botany 55, 1423–1431.

616 Lorite, I.J., Ruiz-Ramos, M., Gabaldón-Leal, C., Cruz-Blanco, M., Porras, R., Santos, C., 2018.
617  Chapter 1 - Water Management and Climate Change in Semiarid Environments, in:
618  García Tejero, I.F., Durán Zuazo, V.H. (Eds.), Water Scarcity and Sustainable

Agriculture in Semiarid Environment. Academic Press, pp. 3–40.
https://doi.org/10.1016/B978-0-12-813164-0.00001-6

Maurel, C., Verdoucq, L., Rodrigues, O., 2016. Aquaporins and plant transpiration. Plant, Cell &
Environment 39, 2580–2587. https://doi.org/10.1111/pce.12814

Melcher, P.J., Goldstein, G., Meinzer, F.C., Minyard, B., Giambelluca, T.W., Loope, L.L., 1994.
Determinants of thermal balance in the Hawaiian giant rosette plant, Argyroxiphium
sandwicense. Oecologia 98, 412–418. https://doi.org/10.1007/BF00324231

Merilo, E., Yarmolinsky, D., Jalakas, P., Parik, H., Tulva, I., Rasulov, B., Kilk, K., Kollist, H.,
2018. Stomatal VPD Response: There Is More to the Story Than ABA. Plant Physiology
176, 851–864. https://doi.org/10.1104/pp.17.00912

Muggeo, V.M.R., 2016. Testing with a nuisance parameter present only under the alternative: a
score-based approach with application to segmented modelling. Journal of Statistical
Computation and Simulation 86, 3059–3067.
https://doi.org/10.1080/00949655.2016.1149855

Mutava, R.N., Prasad, P.V.V., Tuinstra, M.R., Kofoid, K.D., Yu, J., 2011. Characterization of
sorghum genotypes for traits related to drought tolerance. Field Crops Research 123, 10–
18. https://doi.org/10.1016/j.fcr.2011.04.006

Ocheltree, T.W., Nippert, J.B., Kirkham, M.B., Prasad, P.V.V., 2013. Partitioning hydraulic
resistance in Sorghum bicolor leaves reveals unique correlations with stomatal
conductance during drought. Funct Plant Biol 41, 25–36.
https://doi.org/10.1071/FP12316

Oren, R., Sperry, J.S., Katul, G.G., Pataki, D.E., Ewers, B.E., Phillips, N., Schäfer, K.V.R., 1999.
Survey and synthesis of intra- and interspecific variation in stomatal sensitivity to vapour
pressure deficit. Plant, Cell & Environment 22, 1515–1526.
https://doi.org/10.1046/j.1365-3040.1999.00513.x

Percy, R.G., Lu, Z., Radin, J.W., Turcotte, E.L., Zeiger, E., 1996. Inheritance of stomatal
conductance in cotton (Gossypium barbadense). Physiologia Plantarum 96, 389–394.
https://doi.org/10.1111/j.1399-3054.1996.tb00449.x

Pinter, P.J., Zipoli, G., Reginato, R.J., Jackson, R.D., Idso, S.B., Hohman, J.P., 1990. Canopy
temperature as an indicator of differential water use and yield performance among wheat
cultivars. Agricultural Water Management 18, 35–48. https://doi.org/10.1016/0378-
3774(90)90034-V

Prashar, A., Jones, H.G., 2014. Infra-Red Thermography as a High-Throughput Tool for Field
Phenotyping. Agronomy 4, 397–417. https://doi.org/10.3390/agronomy4030397

R Core Team, 2017. . R: A language and environment for statistical computing. Foundation for
Statistical Computing.

Riar, M.K., Sinclair, T.R., Prasad, P.V.V., 2015. Persistence of limited-transpiration-rate trait in
sorghum at high temperature. Environmental and Experimental Botany 115, 58–62.
https://doi.org/10.1016/j.envexpbot.2015.02.007

Ritz, C., Baty, F., Streibig, J.C., Gerhard, D., 2015. Dose-Response Analysis Using R. PLOS
ONE 10, e0146021. https://doi.org/10.1371/journal.pone.0146021

Rueden, C.T., Schindelin, J., Hiner, M.C., DeZonia, B.E., Walter, A.E., Arena, E.T., Eliceiri,
K.W., 2017. ImageJ2: ImageJ for the next generation of scientific image data. BMC
Bioinformatics 18, 529. https://doi.org/10.1186/s12859-017-1934-z

Sampaio Filho, I. de J., Jardine, K.J., de Oliveira, R.C.A., Gimenez, B.O., Cobello, L.O., Piva,
L.R. de O., Candido, L.A., Higuchi, N., Chambers, J.Q., 2018. Below versus above

Ground Plant Sources of Abscisic Acid (ABA) at the Heart of Tropical Forest Response
to Warming. Int J Mol Sci 19, 2023. https://doi.org/10.3390/ijms19072023
Schoppach, R., Fleury, D., Sinclair, T.R., Sadok, W., 2017. Transpiration Sensitivity to
Evaporative Demand Across 120 Years of Breeding of Australian Wheat Cultivars.
Journal of Agronomy and Crop Science 203, 219–226. https://doi.org/10.1111/jac.12193
Schoppach, R.M., Sadok, W., 2013. Transpiration sensitivities to evaporative demand and leaf
areas vary with night and day warming regimes among wheat genotypes. Funct Plant Biol
40, 708–718. https://doi.org/10.1071/FP13028
Shekoofa, A., Balota, M., Sinclair, T.R., 2014. Limited-transpiration trait evaluated in growth
chamber and field for sorghum genotypes. Environmental and Experimental Botany 99,
175–179. https://doi.org/10.1016/j.envexpbot.2013.11.018
Sinclair, T.R., Devi, J., Shekoofa, A., Choudhary, S., Sadok, W., Vadez, V., Riar, M., Rufty, T.,
2017. Limited-transpiration response to high vapor pressure deficit in crop species. Plant
Science 260, 109–118. https://doi.org/10.1016/j.plantsci.2017.04.007
Sinclair, T.R., Hammer, G.L., Oosterom, E.J. van, 2005. Potential yield and water-use efficiency
benefits in sorghum from limited maximum transpiration rate. Functional Plant Biol. 32,
945–952. https://doi.org/10.1071/FP05047
Still, C.J., Rastogi, B., Page, G.F.M., Griffith, D.M., Sibley, A., Schulze, M., Hawkins, L., Pau,
S., Detto, M., Helliker, B.R., 2021. Imaging canopy temperature: shedding (thermal) light
on ecosystem processes. New Phytologist 230, 1746–1753.
https://doi.org/10.1111/nph.17321
Turner, N.C., Schulze, E.-D., Gollan, T., 1985. The Responses of Stomata and Leaf Gas
Exchange to Vapour Pressure Deficits and Soil Water Content. II. In the Mesophytic
Herbaceous Species Helianthus annuus. Oecologia 65, 348–355.
Turner, N.C., Schulze, E.-D., Gollan, T., 1984. The responses of stomata and leaf gas exchange
to vapour pressure deficits and soil water content. Oecologia 63, 338–342.
https://doi.org/10.1007/BF00390662
Vadez, V., Kholova, J., Medina, S., Kakkera, A., Anderberg, H., 2014. Transpiration efficiency:
new insights into an old story. Journal of Experimental Botany 65, 6141–6153.
https://doi.org/10.1093/jxb/eru040
Wang, G.X., Zhang, J., Liao, J.-X., Wang, J.-L., 2001. Hydropassive evidence and effective
factors in stomatal oscillations of Glycyrrhiza inflata under desert conditions. Plant
Science 160, 1007–1013. https://doi.org/10.1016/S0168-9452(01)00344-2
Wang, X., Silva, P., Bello, N.M., Singh, D., Evers, B., Mondal, S., Espinosa, F.P., Singh, R.P.,
Poland, J., 2020. Improved Accuracy of High-Throughput Phenotyping From Unmanned
Aerial Systems by Extracting Traits Directly From Orthorectified Images. Frontiers in
Plant Science 11.
Woods, H.A., Saudreau, M., Pincebourde, S., 2018. Structure is more important than physiology
for estimating intracanopy distributions of leaf temperatures. Ecol Evol 8, 5206–5218.
https://doi.org/10.1002/ece3.4046
Yang, Z., Sinclair, T.R., Zhu, M., Messina, C.D., Cooper, M., Hammer, G.L., 2012. Temperature
effect on transpiration response of maize plants to vapour pressure deficit. Environmental
and Experimental Botany 78, 157–162. https://doi.org/10.1016/j.envexpbot.2011.12.034
Yuan, W., Zheng, Y., Piao, S., Ciais, P., Lombardozzi, D., Wang, Y., Ryu, Y., Chen, G., Dong,
W., Hu, Z., Jain, A.K., Jiang, C., Kato, E., Li, S., Lienert, S., Liu, S., Nabel, J.E.M.S.,
Qin, Z., Quine, T., Sitch, S., Smith, W.K., Wang, F., Wu, C., Xiao, Z., Yang, S., 2019.

Increased atmospheric vapor pressure deficit reduces global vegetation growth. Sci Adv
5, eaax1396. https://doi.org/10.1126/sciadv.aax1396
Zaman-Allah, M., Jenkinson, D.M., Vadez, V., 2011. Chickpea genotypes contrasting for seed
yield under terminal drought stress in the field differ for traits related to the control of
water use. Functional Plant Biol. 38, 270–281. https://doi.org/10.1071/FP10244
Zhang, H., Li, Y., Jian-Kang, Z., 2018. Developing naturally stress-resistant crops for a
sustainable agriculture. Nature Plants; London 4, 989–996.
http://dx.doi.org/10.1038/s41477-018-0309-4
719

FIGURES AND TABLES

Table 1. Summary of hypotheses on transpiration response to VPD classification for sorghum genotypes reported by previous studies. Stomatal conductance (g_s) and canopy temperature (T_{Cirt}) are proxies of transpiration response to VPD evaluated in field experiments from 2019 to 2021 in Kansas and Colorado (see Table S1). Information on canopy temperature was not collected in AB21.

Genotype	Maturity	Inferred VPD response			g_s (years)			T_{Cirt} (years)		
		Growth chambers		Field	19	20	21	19	20	21
		31 °C	37 °C							
SC803	Med	LT [a]	non-LT [d]	LT[e]	+	+		+	+	
SC979[1]	Med	LT [a]	non-LT [d]	-		+	+	+	+	+
BTx2752[2]	Late	LT [a]	non-LT [d]	LT	+	+	+	+	+	+
BTx623	Late	LT [a,b]	non-LT [d]	-		+		+	+	
BTx642	Late	LT [a]	non-LT [d]	LT [e]		+			+	
Macia[3]	Late	LT [a]	LT [d]	-		+		+	+	
Tx7000	Med	LT [b]	-	-		+			+	
SC1345[1,3]	Early	non-LT [a]	-	-				+		
Tx430[3]	Late	non-LT [a]	-	-		+	+	+	+	+
DKS28-05[4]	*Early*	*non-LT [a]*	-	-				+		
DKS54-004[4]	*Late*	*non-LT [a]*	-	*non-LT [e]*	+	+	+	+	+	+
84G62[4]	*Late*	*NN*	*NN*	*NN*	+	+	+	+	+	+
ADVG2275	*Med*	*NN*	*NN*	*NN*		+			+	

NN not known (Commercial checks)

- not available.

+ genotypes evaluated in each experiment.

Germplasm characteristics: [1] Has a Tx430 recombinant inbred line (RIL) family, [2] Elite genotype (G2P Bridge female tester), [3] Nested association mapping parent (NAM) Parent, and [4] Commercial hybrids.

References: [a] (Gholipoor et al., 2010), [b] (Choudhary et al., 2013a), [d] (Riar et al., 2015), and [e] (Shekoofa et al., 2014).

735 **Table 2. Regression parameters for linear, breakpoint and non-linear stomatal**
736 **conductance (g_s) response to VPD for five genotypes.** Information represents data collected in
737 AB19, AB20, AB21, and GR21. 84G62:176 observations, DKS54-00: 173 observations, Tx430:
738 133 observations, Btx2752: 175 observations, and Sc979: 128 observations. RSE: residual
739 standard error.
740

Genotype *Linear*	intercept mol m^{-2} s^{-1}	slope mol m^{-2} s^{-1} kPa^{-1}			R^2	RSE
84G62 (NN)	0.70 ± 0.04	0.05 ± 0.01			0.03	0.23
DKS54-00 (non-LT)	0.80 ± 0.04	0.02 ± 0.02			0.00	0.27
Tx430 (non-LT)	0.98 ± 0.06	-0.08 ± 0.02			0.00	0.24
BTx2752 (LT)	0.64 ± 0.04	0.05 ± 0.02			0.04	0.24
SC979 (LT)	0.80 ± 0.04	0.02 ± 0.02			0.00	0.25
Breakpoint	intercept mol m^{-2} s^{-1}	slope1 mol m^{-2} s^{-1} kPa^{-1}	slope2 mol m^{-2} s^{-1} kPa^{-1}	BP kPa	R^2	RSE
84G62 (NN)	0.37 ± 0.07	0.26 ± 0.04	-0.06 ± 0.04	2.08 ± 0.18	0.16	0.21
DKS54-00 (non-LT)	0.49 ± 0.08	0.20 ± 0.05	-0.07 ± 0.03	2.10 ± 0.20	0.09	0.22
TX430 (non-LT)	-0.057 ± 1.42	0.76 ± 1.21	-0.12 ± 0.29	1.29 ± 0.20	0.11	0.24
BTx2752 (LT)	0.47 ± 0.04	0.16 ± 0.09	0.00 ± 0.02	1.99 ± 0.45	0.06	0.21
SC979 (LT)	0.46 ± 0.16	0.13 ± 0.09	0.01 ± 0.04	2.07 ± 0.56	0.01	0.21
Asymptotic	plateau mol m^{-2} s^{-1}	inflexion point mol m^{-2} s^{-1} kPa-1	start			RSE
84G62 (NN)	0.89 ± 0.02	1.47 ± 0.40	-0.00 ± 0.36			0.22
DKS54-00 (non-LT)	0.89 ± 0.02	1.96 ± 0.68	-0.09 ± 0.60			0.23
Tx430 (non-LT)	-	-	-			-
BTx2752 (LT)	0.81 ± 0.03	1.52 ± 0.81	0.03 ± 0.62			0.21
SC979 (LT)	0.48 ± 0.06	0.69 ± 1.86	0.48 ± 0.62			0.24

742
Figure 1. Drivers of plant transpiration, the hypothetical effect for transpiration response to VPD traits during a growing season, and predictions of field phenotyping for transpiration response to VPD traits. A) Transpiration responses to light intensity represented with an asymptotic function (Q), transpiration response to soil moisture represented with an linear function (SW), hypothesis for the transpiration response to VPD for a putative non-LT (linear) and LT trait (breakpoint). Light intensity regulates stomatal response at night (☾), from

sunrise to noon (⛅), or during any overcast period or day (⛅). LT, as a response to light
saturation occurs around noon (☀). In this period, transpiration is modulated by SW and VPD in
water-limited and well-watered conditions, respectively. B and C) Hypothesis of the effect of the
putative LT and non-LT trait in water dynamics during the growing season. Leaves of a genotype
with a putative non-LT trait are insensitive to high VPD (low humidity) and keep the stomata
open. Leaves of a genotype with putative LT trait perceive a high VPD (low humidity), causing
partial stomatal closure. Blue arrows indicate the effect of the LT on soil moisture during the
growing season. D) Phenomic approaches to discriminate TR-VPD under well-watered
conditions phenotyped via stomatal conductance (g_s) and canopy temperature (T_{Cirt} and T_{Cimg}).
758

760
Figure 2. Study system to dissect transpiration response to VPD in sorghum germplasm. A)
Spatial variability of annual precipitation and maximum VPD in Kansas and Colorado.
Precipitation and VPDe deficit information for sorghum-producing areas were acquired from the
PRISM Climate Group (https://prism.oregonstate.edu/). B and C) The annual trajectory of daily
maximum VPD from 2000 to 2021 and daily maximum VPD for field experiments in Ashland
Bottoms (AB19, AB20) and Greeley (GR21). Maximum daily VPD at each site were acquired
from the PRISM Climate Group (https://prism.oregonstate.edu/). Maximum VPD for each
experiment was recorded with the ATMOS41 weather station. Information of AB21 not included
in Figure 2B.
770
771

772
Figure 3. Approach for comparison of sources for canopy temperature (T_C) in AB19. A)
Multiple ground temperature reference panels, including stainless steel boards painted in white,
gray, and black, flat square containers filled with water and mixture of ice and water in AB19. B
and C) Thermal image reflecting estimated T_C in AB19. D) IRT sensors in GR21. Each sensor
was located on top of the canopy at a 1.5-meter distance. E) Trajectory of T_C obtained with IRT
sensors (black symbol) and UAS-based imaging (magenta symbol) in 16 sorghum plots in
Ashland Bottoms on July 26, 2019 (AB19). Black symbols represent T_{Cirt} every minute during
24 hours, and magenta symbols represent T_{Cimg} every two seconds during a flight time of 12
minutes. UAS-based imaging was collected in motion at 35 meters above ground level. F)
Comparison of T_C obtained with IRT sensors and UAS-based imaging in 16 sorghum plots at
10:00, 12:00, and 14:00 hours in Ashland Bottoms on July 26, 2019 (AB19).

Figure 4. Field Phenomic approaches to discriminate variability in transpiration response to VPD . A, B, and C) Variability for stomatal conductance (g_s) D, E, and F) Canopy temperature from IRT sensors (T_{Cirt}) in hours with high VPD (12:00 to 16:00 hours). H, I and J) Canopy temperature from thermal imagery (T_{Cimg}) Colors represent the reported putative TR-VPD (Table 1): not known (NN, gray), non-limited transpiration (non-LT, blue) and limited transpiration (LT, magenta). Error lines indicate the standard error. Letters indicate significant differences ($\alpha < 0.05$) of all pairwise comparisons using the Sidak test. g_s

represents individual leaves in the sunlit canopy, T_{Cirt} represents the 0.5 m^2 diameter of the sunlit canopy, while T_{Cimg} represents the median T_C at plot level (3 m^2)

Figure 5. The broad sense of heritability (H^2) for different phenomic approaches to discriminate variability in transpiration response to VPD. Stomatal conductance (g_s), canopy temperature from IRT sensors (T_{Cirt}), and canopy temperature from thermal imagery (T_{Cimg}).

Genotype (trait)	*p*-value		
	linear	breakpoint	Non-linear
84G62 (NN)	ns	***	***
DKS54-00 (non-LT)	ns	***	**
Tx430 (non-LT)	ns	ns	-
BTx2752 (LT)	ns	.	ns
SC979 (LT)	ns	ns	ns

Figure 6. Stomatal response to VPD for sorghum germplasm evaluated over three years of
field experiment. A) Comparison of stomatal conductance (g_s) to VPD under irrigated and
rainfed treatments in GR21. The information comprises six genotypes evaluated for their
stomatal response to VPD in GR21. B, C and D) variability of stomatal response to VPD for
sorghum germplasm represented via linear, breakpoint and non-linear regression models. Each
line was fitted with observation indicated in Figure S2.

Figure 7. Principal component analysis (PCA) for genotypes characterized for their TR-
VPD and canopy architecture traits in AB20. P.density: plant density (plants m^{-2}); Leaf.angle:
leaf angle (°); Leaf.area: leaf area per plant (leaf area plant^{-1}); Leaf.size : maximum leaf size
(cm^{2}); N.leaf: maximum number of leaves. Details on each canopy architecture trait is provided
in Table S4 and Figures S3, S5, and S6.

814
Figure 8. Hypotheses on mechanisms underlying variation in transpiration VPD response.
A) Water demand varies due to stomatal response, where the genotype with putative non-LT
maintains open stomata, and the genotype with LT trait partially closes the stomata. B) Both
hydroactive and hydropassive mechanisms can cause stomatal closure, but hormonal signaling is
involved in the hydroactive mechanism. C) Genotypes with non-LT (pink line) and LT (blue)
maintain stomata open, and water demand varies due to differences in hydraulic anatomy. D)
Expected hydraulic anatomic traits for genotypes with putative non-LT and LT traits. The
hydropassive and hydroactive mechanism for the LT trait was adapted from Merilo et al. (2018).
823