Maternal choline supplementation lessens the behavioral dysfunction produced by developmental manganese exposure in a rodent model of ADHD

Authors
Shanna L. Howard[1], Stephane A. Beaudin[1], Barbara J. Strupp[2]*, Donald R. Smith[1]*

[1]Department of Microbiology and Environmental Toxicology, University of California, Santa Cruz, CA, USA
[2]Division of Nutritional Sciences and Department of Psychology, Cornell University, Ithaca, NY, USA
*Co-corresponding authors

Key terms
Manganese, choline, maternal nutrition, dietary supplements, attention, emotion regulation, Attention Deficit Hyperactivity Disorder, motor skills, infant development, environmental exposure

Highlights
1. Developmental Mn exposure causes lasting dysfunction consistent with ADHD symptomology.
2. Maternal choline supplementation (MCS) protects against Mn-induced deficits in attention and behavioral reactivity.
3. MCS in control animals produces lasting benefits to offspring in learning, attention, and error reactivity.
4. These data support efforts to increase choline intake during pregnancy, particularly for individuals at risk of neurotoxicant exposure.

Abstract

Studies in children have reported associations between elevated manganese (Mn) exposure and ADHD-related symptoms of inattention, impulsivity/hyperactivity, and psychomotor impairment. Maternal choline supplementation (MCS) during pregnancy/lactation may hold promise as a protective strategy because it has been shown to lessen cognitive dysfunction caused by numerous early insults. Our objectives were to determine whether (1) developmental Mn exposure alters behavioral reactivity/emotion regulation, in addition to impairing learning, attention, impulse control, and sensorimotor function, and (2) MCS protects against these Mn-induced impairments. Pregnant Long-Evans rats were given standard diet, or a diet supplemented with additional choline throughout gestation and lactation (G3 - PND 21). Male offspring were exposed orally to 0 or 50 mg Mn/kg/day over PND 1–21. In adulthood, animals were tested in a series of learning, attention, impulse control, and sensorimotor tasks. Mn exposure caused lasting dysfunction in attention, reactivity to errors and reward omission, learning, and sensorimotor function, recapitulating the constellation of symptoms seen in ADHD children. MCS lessened Mn-induced attentional dysfunction and partially normalized reactivity to committing an error or not receiving an expected reward but provided no protection against Mn-induced learning or sensorimotor dysfunction. In the absence of Mn exposure, MCS produces lasting offspring benefits in learning, attention, and reactivity to errors. To conclude, developmental Mn exposure produces a constellation of deficits consistent with ADHD symptomology, and MCS offered some protection against the adverse Mn effects, adding to the evidence that maternal choline supplementation is neuroprotective for offspring and improves offspring cognitive functioning.

1. Introduction

Attention Deficit Hyperactivity Disorder (ADHD) is the most prevalent neurodevelopmental disorder among children, affecting ~5-7% of youths up through age 18 years, and ~2–3% of adults[1]. ADHD is characterized by symptoms of inattention, impulsivity/hyperactivity, emotion dysregulation, and in some cases psychomotor deficits[2–4]. Notably, emotion dysregulation has been shown to have a greater impact than inattention and hyperactivity on overall well-being and self-esteem, including peer relationships, academic performance and occupational attainment[5,6]. While the cause(s) of ADHD and related symptoms remains poorly understood, there is an emerging body of epidemiological evidence showing that developmental environmental insults, such as exposure to elevated manganese (Mn) levels, is associated with increased risk of ADHD and related symptoms in children and adolescents[7,8,17,9–16]. Although these epidemiological studies have played a key role in raising concerns about elevated Mn exposure, they are unable to establish causality, in part because of challenges in accurately measuring Mn exposure, and quantifying and controlling for potential confounding factors.

Our prior rodent model studies have shown that developmental Mn exposure can cause lasting ADHD-like impairments in attention, impulse control, and sensorimotor function. In addition, these deficits are associated with hypofunctioning of the catecholaminergic system in the prefrontal cortex and striatum - brain regions that in part mediate attention, impulse control, emotion regulation, and sensorimotor function[18–25]. While our rat model of developmental Mn exposure recapitulates many of the features of ADHD, it remains unclear whether Mn exposure also causes lasting disruption of behavioral reactivity/emotion regulation. This is a significant knowledge gap, given the central role that emotion dysregulation plays in ADHD symptomology and quality of life for children with ADHD[3–5], and the fact that Mn exposure is a risk factor for ADHD diagnosis[7].

While therapies such as methylphenidate (Ritalin) have proven to be effective for children diagnosed with ADHD, there are no established treatments available for children exposed to high levels of Mn during development. A number of nutritional interventions, including maternal supplementation with choline, folic acid, genistein, and L-methionine have been shown to mitigate the effects of a variety of environmental insults[26,27]. In particular, maternal choline supplementation (MCS) has been shown to lessen cognitive

dysfunction caused by prenatal alcohol exposure[28,29], maternal stress[30], and Down syndrome[31-33] in
animal models, to normalize some gene expression and epigenetic molecular alterations caused by
perinatal alcohol exposure[34,35], and to mitigate some of the detrimental effects of prenatal alcohol
exposure in children of alcohol-using mothers[36,37]. In addition, MCS has been shown to improve cognitive
function of typically developing children[38-40] and animal models[41-49]. These benefits of MCS may reflect,
in part, the growing evidence that the amount of choline most commonly consumed by pregnant women
– and pregnant laboratory animals – is insufficient to meet the increased demand for choline during fetal
and neonatal development[50-52]. Physiological demands for choline increase markedly during pregnancy
due to choline's numerous roles in fetal development, including as a precursor biomolecule for
acetylcholine, phosphatidylcholine, sphingomyelin, and betaine[38,53,54]. However, over 90% of pregnant
women consume less than the Adequate Intake (AI) for choline[40,50,51,55] – an amount which itself appears
to be insufficient to meet the demands of pregnancy[38-40].

The present study used our rodent model of early childhood environmental Mn exposure to test two
hypotheses: (1) That developmental Mn exposure causes lasting alterations in behavioral
reactivity/emotion regulation (assessed via the reaction to committing an error or not receiving an
expected reward) in addition to the deficits in attention, impulse control, learning, and sensorimotor
function – all areas of dysfunction seen in children with ADHD; and (2) that supplementing the maternal
diet with additional choline during pregnancy and lactation will offer protection against the adverse effects
of early postnatal Mn exposure on attention, impulse control, learning, sensorimotor function, and
behavioral reactivity (if shown). Collectively, these findings will add to our understanding of the
constellation of effects produced by developmental Mn exposure as a risk factor for ADHD, and they will
provide evidence to evaluate the efficacy of MCS to protect against Mn-induced dysfunctions.

2. Materials and Methods

2.1 Subjects
Sixty-four (64) male Long-Evans rats were used for neurobehavioral assessment. Additional littermates
were used for analyses of tissue Mn, choline, and choline metabolites. Subjects were born in-house from
16 nulliparous timed-pregnant dams (Charles River; gestational age 3; 8-9 weeks old). Twelve to 24
hours after parturition (designated PND 1, birth = PND 0), litters were weighed, sexed, and culled to eight
pups per litter such that each litter was composed of as many males as possible (2-8) and the remainder
females. Each treatment group contained n = 16 animals, and no more than 1-2 males per litter were
assigned to a particular treatment group. At PND 22, all pups were weaned and pair-housed with an
animal of the same treatment group and maintained on a reversed 12:12 hour light/dark cycle. Animals
were housed in polycarbonate cages at a constant temperature of 21 ± 2°C. All aspects of behavioral
testing and feeding were carried out during the active (dark) phase of the animals' diurnal cycle. All
animal care and treatments were approved by the institutional IACUC (protocol Smitd1803) and adhered
to National Institutes of Health guidelines set forth in the Guide for the Care and Use of Laboratory
Animals[56].

2.2 Maternal choline supplementation
Pregnant dams were randomly selected to receive one of two concentrations of choline chloride in their
diet (Dyets AIN-76A Purified Rodent Diet), both of which were provided *ad libitum* during gestation and
lactation, from G3 until PND 21. The control purified diet (unsupplemented) contained 1.1 g choline/kg,
whereas the supplemented purified diet contained 5.0 g choline/kg; both are reported by the
manufacturer to contain 54 mg Mn/kg. Upon weaning on PND 21, all pups were provided Irradiated
Teklad Global Soy Protein-Free Extruded Rodent Diet 2920, reported by the manufacturer to contain 80
mg Mn/kg and 1.2 g choline/kg. See Supplemental Figure S1 for a timeline of maternal choline
supplementation, Mn dosing, and behavioral testing.

2.3 Manganese exposure

Neonate animals were orally exposed to 0 or 50 mg Mn/kg/day in a vehicle solution (5% w/v stevia in
Milli-Q water) from PND 1–21, as in our prior studies[18–24]. Briefly, Mn was delivered once per day directly
into the mouth of each pup (~15-25 µL/dose) using a micropipette fitted with a flexible polyethylene
pipette tip. This Mn exposure regimen is relevant to children exposed to elevated Mn via drinking water,
diet, or both, as pre-weaning exposure to 50 mg Mn/kg/day produces relative increases in Mn intake that
approximates the relative increases reported in infants and young children exposed to Mn-contaminated
water or soy-based formulas (or both)[20,22].
139
2.4 Testing apparatus
Sixteen identical automated 5-Choice Serial Reaction Time Task (5-CSRTT) testing chambers fitted with
odor delivery systems (#MED-NP5L-OLF, Med Associates, Inc., St. Albans, VT) were used to assess
specific cognitive processes, including learning, focused and selective attention, inhibitory control, and
behavioral reactivity, as described previously[20,21,57]. Briefly, each testing chamber contained a curved
wall equipped with five 2.5 × 2.5 cm response ports fitted with light-emitting diodes that served as the
visual cue, an infrared beam to register nose pokes, and pneumatic inlet and vacuum outlet ports to
introduce and remove air-based odor distractors. Opposite the response wall was the food magazine wall
that contained a 45 mg food pellet reward dispensing port fitted with an infrared beam to register nose
pokes; each unit also contained a small house light. The entire testing chamber was enclosed in a sound
attenuating cubicle. Each animal was assigned to one of the 16 testing chambers, and each chamber
was balanced for the four treatment conditions.
152
2.5 Behavioral testing
Animals were mildly food restricted starting on PND 45 in preparation for behavioral testing, with water
available *ad libitum* throughout behavioral assessment. Animals were provided food for 2 hours each day
immediately after behavioral testing, so that their body weights were maintained at ~90–95% of free-
feeding weights. All rats were weighed and tested 6 days/week throughout training and testing.
Behavioral assessment occurred during the active (dark) period of the diurnal cycle at the same time
each day and in the same chamber for each individual rat. A daily test session consisted of 150 trials or
50 min, whichever came first. Each trial sequence was initiated by a nose-poke in the food magazine
port, which was followed by a 3 sec turnaround time to allow the animal to reorient from the food
magazine wall to the response wall; trial onset began after the 3 sec turnaround time. All behavioral
testing was conducted by individuals blind to the treatment condition of the subjects.
164
Behavioral testing began on ~ PND 50. Briefly, testing began with food magazine and nose-poke training
for 11 days, followed by a 5-choice visual discrimination learning task in which the visual cue was
presented immediately after trial onset (i.e., no pre-cue delay), and stayed illuminated for 15 sec (for
more detail, see Beaudin et al.[20]). Animals stayed on this task until they reached the criterion of 80%
correct trials on 2 of 3 consecutive testing days. Following this task, animals were administered a second
visual discrimination task with a fixed 1 sec cue duration and no pre-cue delay for 5 days. This training
was followed by a series of attention and behavioral reactivity tasks as described below (see
Supplemental Figure S1 for the timeline of testing).
173
5-CSRTT assessment is conducted via a series of progressively more difficult tasks, described in greater
detail below, that challenge the ability of animals to sustain attention, inhibit premature responses, and
ignore olfactory distractors while maintaining focus on the predictive visual cue. Recorded response
types for all 5-CSRTT tests included the following: 1) premature responses (responses made after trial
onset but before presentation of the visual cue; assessed in the first focused attention, second focused
attention, and selective attention tasks); 2) correct responses (responses made to the correct port
following presentation of the visual cue); 3) incorrect responses (responses made to the incorrect port
following presentation of the visual cue); and 4) omissions (failure to respond within 15 sec after visual
cue presentation). Premature and incorrect responses and omission errors were not rewarded and were
immediately followed by a 5 sec time-out, in which the house light was turned off for 5 sec. In addition,
the latency (sec) for correct and incorrect responses was recorded (response latency), as was the

latency (sec) to retrieve the food pellet reward following a correct response (collection latency).
Perseverative responses, defined as additional nose pokes into the response port after a correct
response, were also recorded. The calculated response outcomes were: (1) %Correct, calculated as
number of correct responses/(correct + incorrect + premature + omissions) × 100; (2) %Incorrect,
calculated as above but with incorrect responses in the numerator; (3) %Premature, calculated as above
but with premature responses in the numerator; and (4) %Omissions, calculated as above but with
omissions in the numerator. Additionally, we assessed attentional accuracy via a %Accurate outcome,
calculated as number of correct responses/(correct + incorrect) × 100. The number of perseverative
responses per correct response was calculated simply as #perseverative responses/#correct responses.
194

2.6 Focused attention tasks

Focused attention can be defined as the ability to maintain attentional focus on a specific task or stimulus
(e.g., a visual cue). Two successive focused attention tasks (FA1 and FA2) were administered over
~PND 85-103, and PND 104-121, respectively, following completion of the visual learning discrimination
task. Both focused attention tasks assessed the ability of animals to detect and respond to a visual cue
presented unpredictably in time and location (one of the five response ports). The first focused attention
task used variable pre-cue delays of 0, 1, 2, or 3 sec and a fixed visual cue duration of 0.7 sec and was
administered for 15 daily sessions. The second focused attention task included variable pre-cue delays
of 0, 3, 4, or 5 sec and variable visual cue durations of 0.5 or 0.7 sec and was administered for 15
sessions.
205

2.7 Selective attention task with olfactory distractors

Selective attention can be defined as the ability to maintain a behavioral or cognitive set in the face of
distracting or competing stimuli. The selective attention task (SAT) with olfactory distractors was
administered across 12 sessions, from PND 129-140. In this task, the pre-cue delay varied between 3
and 4 sec, with the two delays balanced across the trials within each test daily session, and the cue
duration was 0.5 sec. On one third of the trials in each session, an olfactory distractor was presented 1 or
2 sec after trial onset (i.e., 1–3 sec before the visual cue, depending on the pre-cue delay). The delay
between odor distractor presentation and visual cue presentation (1-3 sec) is referred to as the odor-to-
cue interval (OtoC) and was included in statistical analysis. The nine different olfactory distractors were
made from liquid odorants (McCormick & Company, Inc.) diluted in propylene glycol, and delivered into
an incorrect response port as scented air. The SAT was preceded by a baseline task (three sessions),
administered from PND 126-128. This task was identical to the SAT, except it did not include the
presentation of olfactory distractors.
219

2.8 Assessment of behavioral reactivity and compulsiveness in the reward omission task

Behavioral reactivity/emotion regulation was assessed by measuring the animal's performance on trials
following an error or not receiving an expected reward, areas where children with ADHD have been
shown to differ from typically developing children[3,4]. Reactivity to committing an error was assessed in
two tasks: (1) The second focused attention task, and (2) a reward omission task. In this latter task, a 0.7
sec visual cue was presented immediately at trial onset, and the food pellet reward was randomly
omitted on three out of 10 correct trials. It is notable that the consequences of committing an error versus
a reward omission are very different for the animal. Upon committing an error (in all tasks), the rat
receives a 5 sec 'time-out' during which the house light turns off and the animal sits in darkness, and no
pellet reward is provided. In contrast, on a reward omission trial the house light remains on and all other
trial conditions/cues remain unchanged from a normal rewarded correct trail. Attentional performance on
the reward omission task was assessed as a function of whether the preceding trial was either (1) correct
(and rewarded), (2) incorrect (including omission errors), or (3) correct but food reward omitted. We also
assessed perseverative responses in the reward omission task as a measure of compulsivity, consistent
with prior studies[58,59]. Perseverative responses are additional nose-pokes into the correct port following a
correct response. Animals were tested in the reward omission task for 12 sessions from PND 175-187.
236

2.9 Montoya staircase task

The Montoya staircase test was used to assess sensorimotor function, as described previously by us[19].
Performance on the staircase test is a sensitive measure of skilled forelimb function directly relevant to
pediatric Mn studies. This test taps essential sensorimotor functions required to successfully reach and
grasp small food pellets from the descending steps of the staircase, including advancing the limb over
the food, opening the digits in preparation for grasping, grasping and manipulating the food, and
withdrawing the paw to place the food in the mouth[60]. Optimal performance on this test requires several
sensorimotor skills that are similarly assessed in human studies, including forelimb movements and digit
flexion for manipulating objects. Thus, the staircase test measures aspects of fine psychomotor skills that
are known to be disrupted in children with a history of developmental Mn exposure[8,61].
247
248 Eight identical Plexiglas staircase devices were used, modeled after the original design of Montoya et
249 al.[62]. Each device contained a platform and a left-and-right side staircase. The staircases contained 6
250 descending steps on each side, and each step was baited with three 45 mg food pellets for a total of 36
251 pellets available at the start of a session. To determine the step of origin and final location of each pellet,
252 pellets were color coded with a unique color per step level[63].
253
The staircase study was conducted over 1 month, beginning on PND 85. Each staircase session
occurred immediately following operant testing every day. Staircase training occurred for the first 11
days. During training, rats retrieved the food pellets from the descending steps of the staircases until all
subjects reached asymptotic performance of total pellets retrieved. The next 19 days comprised the
testing phase of the task. Throughout training and testing, each subject was given one 10-min trial per
day, 6 days per week. After each trial, the rat was removed, and the final distribution of colored pellets
was quantified per step and side. The order and time of testing were balanced by treatment and
remained the same each day for each rat.
262
Skilled forelimb performance was evaluated step-by-step for (1) the number of pellets taken (i.e., pellets
that were grasped and removed from their step of origin), (2) the number of pellets eaten (pellets that
were taken and consumed), (3) the corresponding percent of success (the ratio of the number of pellets
eaten/pellets taken x 100), (4) the number of pellets misplaced (all the pellets that were grasped but
ended up on a different step level than their step of origin), and (5) the number of pellets lost (pellets that
ended up on the floor of the apparatus, out of reach of the animal).
269
2.10 Tissue Mn and choline/choline metabolite levels
For blood and brain Mn levels, tissue samples were collected from weaned pups on PND 24, and from
the behaviorally tested animals on PND 202. Briefly, animals were euthanized via CO_2 asphyxiation, and
whole blood was immediately collected from the left ventricle of the surgically-exposed heart and stored
in EDTA Vacutainers at − 20 °C for analyses. Whole brain was immediately removed, and the hind-brain
region collected and stored at −80 °C for Mn concentration determinations. Blood and brain Mn
concentrations were determined by inductively coupled plasma–mass spectrometry (Thermo Element
XR), as previously described[18,20]. The analytical detection limit for Mn was 0.01 ng/mL.
278
Choline/choline metabolite levels were measured in plasma, brain, and/or liver tissues from dams at
weaning (plasma and liver), and/or from the same PND 24 animals (liver and hindbrain) used for Mn
analyses described above. Briefly, concentrations of free choline, betaine, dimethylglycine, methionine,
and trimethylamine N-oxide were measured by liquid chromatography tandem mass spectrometry
according to the method of Holm et al.[64], while phosphocholine, glycerophosphocholine,
phosphatidylcholine, acetylcholine, and lysophosphatidylcholine were measured by liquid
chromatography/mass spectroscopy according to the method of Koc et al.[65] with modifications[66].
286
2.11 Statistical methods
The behavioral data were modeled using structured covariance mixed models. Fixed effects included in
the models were the between-subject factors MCS (two levels corresponding to the unsupplemented and
MCS groups), and Mn exposure treatment (two levels corresponding to the 0 and 50 mg/kg/d groups),

and the within-subject factors pre-cue delay, cue duration, session block, distraction condition, and/or
previous trial outcome, depending on the outcome analyzed. In all models, rat was included as a random
effect to account for correlations within observations from the same animal. Statistical tests used the
Kenward-Roger correction to improve estimation of the variance-covariance of the fixed effects
parameters. Plots of residuals by experimental condition were used to examine the assumption of
homogeneity. Random effects with high variance in the residuals across the levels of the within-subject
factor (e.g., distraction condition) were added to the model to achieve homogeneity if necessary. The
distribution of each random effect was inspected for approximate normality and presence of influential
outliers. The significance level was set at $p \leq 0.05$, and p-values between 0.05 and 0.10 were considered
to be trending towards significance; trending effects are presented if they aid in clarifying the nature of
the Mn and/or MCS effects. Significant main effects or interaction effects were followed by single-degree
of freedom contrasts to better understand the nature of the interactions, using the Student's t-test for
pairwise comparisons of least squared means. Analyses were conducted using SAS (version 9.4) for
Windows. For tissue Mn and choline/metabolite data, analyses were performed using one-way (for
dams) or two-way (for offspring) analysis of variance and Tukey's post hoc test for pairwise comparisons,
using JMP (version 16.0; SAS Institute, Inc.).

3. Results

For the 5-CSRTT measures, we focus primarily on outcomes of %Premature, %Correct, and %Accurate
responses because our prior studies have shown that these measures best differentiated control vs Mn
animals[20,21] and did so again in this study. Note that in general, we graphically depict the data for all four
treatment groups (defined by diet and Mn exposure), regardless of whether the interaction of Mn and
MCS was significant, due to our *a priori* hypotheses relating to the magnitude of the Mn effect under
each of the two diet conditions.

3.1 Visual learning

Visual learning in the 5-choice task is not altered by Mn or MCS
Visual learning (i.e., learning to make a nosepoke into the illuminated port) is foundational to the
subsequent, more challenging 5-CSRTT tasks. Here, there was no effect of early postnatal Mn exposure
or MCS on visual learning. Specifically, there was no main effect of Mn or MCS on trials to criterion or
days to criterion, and no Mn x MCS interaction for these two outcomes (p's >0.295). Moreover, there also
was no effect of Mn or MCS on the percentage of animals per treatment group failing to reach the
learning criterion. The lack of treatment group differences in learning the task criterion indicates that
visual acuity and motivation did not differ between groups and therefore cannot account for group
differences seen in subsequent 5-CSRTT tasks.

3.2 Impulsivity

Developmental Mn exposure did not affect impulse control, but did impair the animals in learning
to wait for the visual cue, and this effect is not rescued by MCS
332 We evaluated %Premature responses as a measure of inhibitory control in the two focused attention
333 tasks and the selective attention task.

In the first focused attention task, Mn exposure impairs learning to withhold premature responses, and
MCS does not alter this effect.
In the first focused attention task, there is a significant Mn x session block interaction [$F_{(2, 435.8)} =$
11.07, p<0.0001] for %Premature responses. Whereas both control and Mn groups commit a relatively
high percentage of premature responses at the start of the task (~ 50%, since this was the first task in
which animals faced longer pre-cue delays), the Mn animals improve less rapidly than controls. As a
result, the Mn animals commit significantly more premature responses than controls in session block 3
(p=0.0084; Figure 1a1). There is also a significant MCS x session block interaction [$F_{(2, 435.8)} = 17.06,$
p<0.0001], but none of the contrasts achieved significance (this two-way interaction is not plotted, but all

four groups are presented in figure 1A.) The three-way interaction of MCS, Mn and block was not
significant; p = 0.69).
346
In the second focused attention task, neither Mn nor MCS significantly alter premature responses.
In the subsequent second focused attention task, neither main effect of Mn (p = 0.75) nor MCS (p = 0.39)
was significant. However, there is a Mn x session block interaction [F(2, 933.7) = 5.53, p = 0.0041],
although there are no significant contrasts between control and Mn animals in any session block (p's
>0.40; Figure 1b1) (see also Figure 1B, which depicts differences between Mn and controls under each
of the two diet conditions; the Mn x MCS x block interaction p = 0.40). There is also a MCS x pre-cue
delay interaction [F(2, 932.8) = 3.18, p = 0.0422], though again, none of the specific contrasts between
unsupplemented and MCS animals reached significance at any pre-cue delay (p's > 0.21, Figure 1c1).
Differences between Mn and controls under each of the two diet conditions is presented in Figure 1C
(the three-way interaction of MCS, Mn and pre-cue delay was not significant; p = 0.24).
357
In the selective attention task, MCS lessens the difference in premature responses between treatment
groups in early session blocks.
In the selective attention task with olfactory distractors, there is a Mn x MCS x session block interaction
for %Premature responses [F(3, 1665) = 4.44, p = 0.0041], although none of the specific contrasts
between treatment groups or diet conditions reached significance for any session block (p's > 0.12,
Figure 1D). Nonetheless, the significant three-way interaction appears to reflect the fact that the
difference in premature responses between Mn and control groups in the first two session blocks is
greater for unsupplemented animals than for the MCS animals, suggesting that MCS lessened the
impulsivity caused by developmental Mn exposure.
367
368 **3.3 Attention**
369
370 **Mn impairs focused attention and MCS lessens this effect**
371 As described earlier, the two focused attention tasks were used to assess whether Mn exposure alters
372 the ability of animals to maintain attentional focus, and whether the Mn impairment was lessened by
373 MCS.
374
In the first focused attention task, Mn exposure impairs attention, and MCS improves learning of the
control group.
In the first focused attention task, there is a significant Mn x session block interaction for %Correct
responses [F(2, 595.6) = 4.12, p = 0.0167]. This reflects that performance in the Mn animals improves
more slowly than control animals across session blocks, such that the Mn animals have significantly
fewer %Correct responses than the controls in session blocks two and three (p=0.007 and 0.002,
respectively, Figure 2a1) (see Figure 2A, which depicts differences between Mn and controls under each
of the two diet conditions; the three-way interaction p = 0.22).
383
Consistent with the %Correct outcome noted above, there is a significant Mn x session block interaction
for %Accurate responses [F(2, 601.3) = 3.33, p = 0.0365; not shown], and a trending three-way
interaction of Mn x MCS x session block for %Accurate responses [F(2, 601.3) = 2.58, p = 0.0769]
(Figure 2B). This three-way interaction reflects the fact that, while all groups improve in attentional
accuracy across session blocks (p's<0.0014), the difference in %Accuracy of Mn and control animals
was greater in the MCS condition than the unsupplemented condition. This appears to be because the
controls benefitted more from MCS than the Mn animals, in terms of improvement in %Accuracy across
session blocks. As a result, under MCS, a Mn deficit in %Accuracy emerges under session blocks two
and three (p's = 0.018 and 0.049, respectively; Figure 2B).
393
394 *In the second focused attention task, MCS improves performance of both the Mn and control animals,*
395 *but in different functional domains.*

In the second focused attention task there is a significant Mn x MCS x pre-cue delay interaction for %Correct responses [$F_{(3, 1277)} = 7.32$, $p < 0.0001$] (Figure 2C). For animals on the unsupplemented diet, the Mn-exposed rats have significantly lower %Correct responses than controls at the 3 sec delay ($p = 0.0477$), and this Mn deficit is also trending at the 4 and 5 sec delays (p's = 0.0518 and 0.0699, respectively). However, in the MCS animals, the Mn deficit at the 3 sec delay is only trending ($p = 0.0762$), and the deficit is no longer present under the 4 and 5 sec pre-cue delays (p's = 0.27 and 0.69, respectively). Thus, the Mn x MCS x pre-cue delay interaction reflects the fact that the differences between control and Mn animals at the longer delays is smaller under MCS than under the unsupplemented diet. Additionally, control animals that received MCS have significantly more correct responses at the 0 sec delay compared to unsupplemented controls ($p = 0.0113$), reflecting a benefit of MCS to control animals for trials when the cue was presented immediately at trial onset, with no pre-cue delay.

We also assessed attentional accuracy (as %Accurate responses) for the second focused attention task, and again observed a significant Mn x MCS x pre-cue delay interaction [$F_{(3, 1271)} = 3.95$, $p = 0.0081$] (Figure 2D), reflecting a pattern in attentional performance similar to what was seen with the %Correct outcome described above. Under the unsupplemented diet, there is a trending Mn deficit at the 3 and 4 sec pre-cue delays (p's = 0.058 and 0.067, respectively). In contrast, under MCS, this Mn deficit is still present at the 3 sec delay ($p = 0.043$), but not at the 4 sec pre-cue delay ($p = 0.23$). While there was no significant Mn deficit at the 5 sec pre-cue delay under the unsupplemented diet, MCS also appears to reduce the difference between control and Mn animals at this longer delay ($p = 0.11$ under the unsupplemented diet and 0.53 under MCS; Figure 2D).

While there was a significant benefit of MCS to controls at the 0 sec pre-cue delay for %Correct responses ($p = 0.0113$), for %Accurate this benefit did not reach significance ($p = 0.129$). There was, however, a trending Mn deficit that emerged at the 0 sec delay under MCS ($p = 0.09$, Figure 2D).

Collectively, this evidence suggests that the lasting attentional deficits caused by developmental Mn exposure persist into the second focused attention task. Additionally, MCS improves performance of the controls but not the Mn animals for trials with a 0 sec pre-cue delay, whereas it reduces the difference between Mn and control animals at the longer (more challenging) delays. The MCS benefit derived by the controls at the 0 sec delay appears to be due to a lessened reaction to committing an error on the prior trial (discussed below; Figure 4), whereas the benefit derived by the Mn animals at the longer delays appears to reflect improved attention.

MCS normalizes the performance of the Mn rats on distraction trials in the selective attention task
To determine the effects of Mn exposure and MCS on selective attention, we assessed animal performance on a selective attention task with olfactory distractors randomly presented either 1, 2, or 3 sec prior to the onset of the visual cue (referred to as the odor-to-cue interval, 'OtoC').

Mn causes attentional dysfunction and MCS reduces the difference between Mn and control animals on trials with olfactory distractors.
For %Correct responses in the selective attention task, there was a significant Mn x OtoC interaction [$F_{(3, 239.2)} = 2.94$, $p = 0.0339$], where the Mn animals were more impaired relative to controls for trials without a distractor ($p = 0.0096$; Figure 3a1). Although the Mn x MCS x OtoC interaction was not significant ($p = 0.269$) (Figure 3A), the pattern described by the Mn x OtoC two-way interaction seems to be driven by the fact that MCS diminished the differences between the Mn and control group for distraction trials (Figure 3A); this pattern is not seen in the unsupplemented groups. In light of this, we conducted follow-up analyses where the three trial conditions with olfactory distractors (i.e., OtoC of -1, -2, -3 sec) were collapsed into a single "distraction" trial condition, thus creating only two distraction conditions (distractor and no-distractor trials; Figure 3B). In this follow-up analysis, there was a trending interaction of Mn x MCS x distractor [$F_{(1, 52.16)} = 3.29$, $p = 0.0754$]. This trend reflects that, for non-distraction trials, the Mn animals perform more poorly than controls under both diets, but for distraction

449 trials the Mn deficit emerges only for unsupplemented animals (Figure 3B); for the MCS condition, the
450 Mn and controls perform similarly for trials with distractors.
451
For %Accurate responses, there was a significant Mn x OtoC interaction [$F(3, 235.7) = 3.14$, $p = 0.026$],
where the Mn deficit was significant for trials without a distractor and trending for trials with the -3 sec
distractor condition (p's = 0.0084 and 0.098; Figure 3c1); the groups were not different for trials with a -1
or -2 distractor. The Mn x MCS x OtoC three-way interaction was not significant [$F(3, 235.8) = 0.8$, $p = 0.495$];
0.495]; however, there is a similar pattern of group differences as noted above for %Correct, where MCS
reduces the difference in %Accuracy between control and Mn animals under the distractor conditions
(Figure 3C). However, neither the full (Figure 3C) nor distractor-simplified (not shown) %Accurate models
revealed a significant Mn x MCS x odor interaction (p's > 0.31). Collectively, these findings indicate that
MCS lessens the difference between control and Mn animals for trials with olfactory distractors.
461
462 **3.4 Behavioral reactivity**
463
Mn alters reactivity to errors and reward omission, effects which are partially normalized by MCS
In the second focused attention task, Mn animals born to dams on the unsupplemented diet do not
exhibit behavioral reactivity to an error, and MCS improves performance in control animals following an
error.
In the second focused attention task, control animals that received MCS had significantly more correct
responses than their unsupplemented counterparts at the 0 sec pre-cue delay (reported above in Figure
2C). We hypothesized that this might be due to MCS reducing the disruptive effect of committing an error
on the prior trial, as trials with a 0 sec pre-cue delay are closest temporally to the prior trial (i.e., if the
outcome of the prior trial was disruptive to the animal, it would be most apparent for trials for which the
cue was presented immediately after trial onset). Therefore, we assessed %Correct responses on trials
with a 0 sec pre-cue delay, as a function of the outcome of the prior trial (correct or error).
475
The analysis of %Correct for trials with a 0 sec pre-cue delay revealed a significant Mn x MCS x previous
trial outcome interaction [$F(1, 273.2) = 6.13$, $p = 0.0139$]. This interaction reflects that the reaction to
committing an error varied as a function of both Mn exposure and maternal diet. Specifically, under the
unsupplemented diet, only the controls exhibited fewer correct responses on trials following an error
(relative to after a correct response); the Mn animals showed no decline in performance due to making
an error on the prior trial ($p < 0.0001$ for control and $p = 0.79$ for Mn animals comparing error vs. correct;
Figure 4). In contrast, under the MCS condition, both control and Mn groups showed a trending reduction
in %Correct for trials following an error (relative to after a correct response; $p = 0.074$ for control and 0.17
for Mn), and the magnitude of the disruptive effect of making an error was similar in the two groups
(Figure 4). Moreover, in the control animals, MCS (vs. the unsupplemented condition) significantly
increased correct responses after an error, but not following a correct response (p's=0.05 and 0.50,
respectively), while in the Mn animals, MCS did not improve performance following either a correct
response ($p = 0.34$) or error ($p = 0.73$), compared to unsupplemented animals (Figure 4).
489
490 In contrast to performance on trials with a 0 sec pre-cue delay, there was no Mn x MCS x previous trial
491 outcome interaction for performance on trials with a 3 sec pre-cue delay [$F(1, 273.8) = 1.72$, $p = 0.19$]
492 (not shown). The absence of this three-way interaction for trials with a longer pre-cue delay supports the
493 inference that the group differences seen at the 0 sec pre-cue delay (described above) reflects reactivity
494 to the outcome of the prior trial.
495
Developmental Mn exposure and MCS both alter reactivity to committing an error and reward
omission in the reward omission task
In the reward omission task, we introduced a novel trial condition in which the food pellet reward is
unexpectedly omitted on a subset (30%) of correct response trials and assessed how this affects
performance on the subsequent trial. Both errors and reward omission have been shown to be disruptive
to performance in our prior studies with similar operant tasks[33,67], and both are often altered in children

with ADHD compared to typically developed children[3,4,68–70]. Specifically, we assessed performance on
each trial as a function of whether the preceding trial response was: (1) correct with a food pellet reward,
(2) an error (incorrect, no food pellet reward + 5 sec time out), or (3) correct but with the food pellet
reward omitted (RO).

In the unsupplemented animals, Mn lessens reactivity to an error but increases reactivity to the omission
of an expected reward
For %Correct responses, there was a significant three-way interaction of Mn x MCS x previous trial
outcome [$F_{(2, 410.9)} = 4.13$, $p = 0.0168$] (Figure 5A). This reflects that for the unsupplemented controls,
committing an error on the prior trial caused a reduction in %Correct on the subsequent trial (versus a
correct response on the prior trial; $p < 0.0001$), whereas the unexpected omission of the food reward on
the prior (correct) trial had no effect on performance ($p = 0.17$). A markedly different pattern was seen in
the unsupplemented Mn animals; for these animals, committing an error on the prior trial did not affect
performance compared to trials following a correct response ($p = 0.152$) (Figure 5A). In contrast,
performance of the unsupplemented Mn animals was significantly decreased on trials following the
omission of the expected food reward relative to after a correct response ($p = 0.0032$). These findings
demonstrate that the unsupplemented animals exposed to Mn early in life respond differently than
unexposed controls (no Mn) to both committing an error and not receiving an expected reward.

MCS normalizes behavioral reactivity to an omitted reward in the Mn animals
Notably, a different pattern of performance was seen for the MCS animals. Specifically, for the Mn-
exposed animals, MCS normalized performance on trials following a reward omission, such that
performance was the same as for trials following a correct response – a pattern matching that of the
unsupplemented controls (no Mn) (Figure 5A). However, MCS did not alter how the Mn animals perform
following an error (vs a correct response), compared to their unsupplemented Mn-exposed counterparts
(Figure 5A). Finally, under the MCS diet, performance on trials following a correct response was
significantly worse in the Mn animals than controls ($p = 0.0189$), a deficit that was not present in the
unsupplemented groups; this appears to be because the controls benefited from MCS with improved
performance on trials following a correct response, but the Mn animals did not (Figure 5A).

For unexposed controls, MCS lessens the disruptive effect of committing an error
For the unexposed controls, there was a trend for MCS to lessen the disruptive effect of committing an
error on the prior trial; i.e., to improve performance on trials that followed an error (relative to post-error
trials for the unsupplemented controls) ($p = 0.076$, Figure 5A). However, for the MCS controls, %Correct
responses on trials following an error was still lower than on trials that followed a correct response ($p <$
0.0001). Furthermore, for the MCS controls, performance on trials following a reward omission was lower
than that on trials following a correct response ($p = 0.0015$), an effect that was not seen in the
unsupplemented control animals. However, it is notable that performance on trials following a reward
omission did not differ between the unsupplemented and supplemented controls; thus, the relative drop
in performance by the MCS controls on trials following a reward omission (versus performance on trials
after a correct response) was due to an *increase* in performance on trials following a correct response,
not a decline in performance following reward omission.

Mn animals exhibit less post-error slowing than controls, and MCS normalizes this effect.
For response latency, there was a three-way interaction of Mn x MCS x previous trial outcome [$F_{(2,}$
$_{121.4)} = 5.11$, $p = 0.0074$] (Figure 5B). This reflects that for animals born to dams on the
unsupplemented diet, both control and Mn groups exhibit post-error slowing (i.e., increased response
latency following an error vs. a correct response; $p < 0.0001$); however, for the unsupplemented groups,
the post-error slowing was significantly less pronounced in Mn animals than controls ($p = 0.0046$), and
the magnitude of the difference in response latency following a correct response compared to an error is
smaller in the Mn versus control animals (Figure 5B). Maternal choline supplementation normalizes post-
error slowing in the Mn animals, such that the response latency following an error is the same as in
controls, and the magnitude of the slowing (i.e., increase in response latency following an error versus a

correct response) is comparable in the Mn and control animals. Finally, trials following a reward omission have a comparable response latency to trials following a correct response across treatment groups, except in control animals that received MCS; these animals exhibit a slight but statistically significant increase in response latency following a reward omission compared to a correct response (p = 0.044) (Figure 5B).

Mn increases perseverative responding following a correct response in the unsupplemented animals, but this abnormality is not seen in the Mn animals born to supplemented dams.
The analysis of perseverative responses per correct response revealed a three-way interaction of Mn x MCS x previous trial outcome [F(2, 111.2) = 3.54, p = 0.0322] (Figure 5C). All animals make more perseverative responses on trials following a correct response than on trials following an error or reward omission (p's < 0.0001), and they make the fewest perseverative responses on trials following a reward omission (compared to trials following either a correct response or error; p's < 0.0001). For groups in the unsupplemented diet condition, the Mn animals tend to make more perseverative responses following a correct trial than controls (p = 0.0663). However, for the MCS groups, Mn animals trended to have fewer perseverative responses than controls following a correct response (p = 0.053), and there is a significant decrease in perseverative responses following a correct response in the MCS+Mn animals compared to their unsupplemented counterparts (p = 0.0019) (Figure 5C).

Mn increases food pellet collection latency; MCS reduces pellet collection latency in Mn animals and increases collection latency in controls, mirroring the effect seen for perseverative responses.
For food pellet collection latency (i.e., the time to collect the food pellet reward following a correct response), there was a Mn x MCS x previous trial outcome interaction [F(2, 410.3) = 6.69, p = 0.0014] (Figure 5D). The pattern of effects seen for pellet collection latency was similar to that seen for perseverative responses, as expected, because committing more perseverative responses delays collecting the pellet reward. In general, pellet collection latency is longest on trials following a correct response, and shortest for trials following a correct response with an omitted reward. Unsupplemented Mn animals have a significantly longer collection latency following a correct response compared to controls (p = 0.0265), while under MCS, Mn animals have a significant decrease in collection latency compared to unsupplemented Mn animals following a correct response (p = 0.0005). In addition, Mn animals that received MCS have significantly shorter collection latency than controls following a correct response (p = 0.0006) (Figure 5D). This effect is enhanced by the fact that MCS also increases collection latency in control animals following a correct response, compared to unsupplemented animals (p = 0.039).

3.5 Sensorimotor function

Early life Mn exposure causes lasting impairment in sensorimotor function that is not altered by MCS.
To determine whether MCS can mitigate the lasting sensorimotor impairments caused by developmental Mn exposure, animals were assessed using the Montoya staircase test of skilled forelimb performance. Mn exposure caused lasting impairment in sensorimotor function, as demonstrated by a significant Mn x step level interaction for the number of pellets taken [F(5, 229.3) = 10.66, p < 0.0001], number of pellets eaten [F(5, 232.6) = 4.57, p = 0.0005], %Success [F(5, 231.4) = 3.26, p = 0.0073], and the number of pellets misplaced [F(5, 356.7) = 5.76, p < 0.0001]. Specifically, there was a significant Mn deficit on steps 4 and 5 of the staircase, where the unsupplemented Mn animals ate significantly fewer pellets and misplaced more pellets than their unsupplemented control counterparts (pellets eaten, p's = 0.0017 and 0.0002 for steps 4 and 5 respectively; pellets misplaced, p's = 0.0005 and 0.002) (Figure 6a1 and 6b1, respectively). MCS did not affect this Mn-induced impairment in reaching and grasping, based on findings that neither the main effect of MCS (p's >0.16), nor any interactions involving MCS and Mn were significant for any of the outcomes (e.g., Mn x MCS, P's >0.32; Mn x MCS x step level, p's >0.72); see figure panels 6A and 6B for the Mn x MCS x step plots for pellets eaten and misplaced).

3.6 Choline and choline metabolite levels

MCS measurably altered choline and choline metabolite levels in dams and offspring.
To confirm that oral MCS over gestation and lactation altered levels of choline and/or its metabolites in dams and their offspring as expected, and whether oral Mn exposure altered choline/metabolite status in the offspring of MCS-treated dams, we measured levels of choline and selected metabolites in dam and offspring tissue (dam plasma and liver upon completion of MCS at PND 21, offspring liver and brain at PND 24). As expected, MCS produces measurable changes in choline and some of its metabolites in dam plasma and liver, and PND 24 pup liver and brain (see Supplemental Table S2 for choline and metabolite measurements in dams, and Supplemental Table S3 for measurements in pups).

3.7 Blood and brain Mn levels

Oral Mn dosing increased blood and brain Mn levels in weanlings, and returned to control levels in adulthood.
Oral Mn exposure over PND 1-21 increased Mn levels in both blood and brain tissue in PND 24 weanlings. Specifically, Mn exposure significantly increases blood Mn levels ~eight-fold, from ~30 to 250 ng/mL in control vs. Mn exposed groups [$F_{(1, 45)}$ = 398.4, $p < 0.0001$] (Supplemental Figure S4A). There was also a significant interaction of Mn x MCS on blood Mn levels [$F_{(1, 42)}$ = 8.37, $p = 0.0059$]; however, Tukey's contrasts revealed no significant difference between blood Mn levels in unsupplemented versus MCS animals exposed to Mn.

Mn exposure also causes an approximately three-fold increase in brain Mn levels in PND 24 offspring, from approximately 3.8 to 11.8 µg/g in control versus Mn-dosed animals, as demonstrated by a significant main effect of Mn [$F_{(1, 36)}$ = 129.7, $p < 0.0001$] (Supplemental Figure S4B). There is also a trending main effect of MCS on brain Mn levels in offspring [$F_{(1, 36)}$ = 3.96, $p = 0.0543$], as well as a trending Mn x MCS interaction [$F_{(1, 36)}$ = 3.13, $p = 0.085$]. Tukey's pairwise comparisons revealed that MCS animals treated with Mn had significantly higher brain Mn levels than their unsupplemented counterparts.

Finally, while blood and brain Mn levels in the PND 200 behaviorally-tested animals return to baseline levels, there remained a statistically significant but biologically unremarkable effect of MCS [$F_{(1, 27)}$ = 7.87, $p = 0.0092$] to reduce blood Mn levels compared to the unsupplemented group, and a similar trending effect of MCS to reduce brain Mn levels in the Mn-exposed group [$F_{(1, 20)}$ = 3.19, $p = 0.089$] (Supplemental Figure S4D). However, Tukey's pairwise comparisons revealed no significant difference between any groups.

4. Discussion

ADHD, which is characterized by impairments in attention, impulse control, and emotion regulation, is the most prevalent neurobehavioral disorder in children, yet the causes of ADHD remain poorly understood[1]. Studies in children and adolescents have revealed associations between developmental Mn exposure and inattention, impulsivity/hyperactivity, emotion dysregulation, and sensorimotor dysfunction[7,8,17,9–16]. However, there are currently no recognized interventions, other than exposure prevention, to mitigate the neurotoxic effects of elevated developmental Mn exposure. Here, using a rodent model of environmental Mn exposure, we demonstrate that developmental Mn exposure causes lasting dysfunction in attention, behavioral reactivity, learning, and sensorimotor function, and that increasing maternal choline intake during pregnancy was effective in protecting against some of these Mn-induced impairments, particularly in the domains of attentional function and behavioral reactivity. MCS also was beneficial for the unexposed animals in the domains of learning, attention, and behavioral reactivity. Collectively, these findings further establish developmental Mn exposure as a significant risk factor for environmentally-induced ADHD and related symptoms. They also demonstrate that MCS exerts lifelong cognitive benefits to offspring, including providing some protection against the neurotoxic impacts of elevated Mn exposure.

661 Given recent evidence that 90% of pregnant women consume less choline than the recommended
662 Adequate Intake (AI), these data provide additional support for efforts to increase choline intake during
663 pregnancy and lactation, particularly for women at risk of environmental exposure to Mn. These findings
664 and inferences are discussed in detail below.
665
4.1 Developmental Mn exposure causes lasting dysfunction in attention, learning, behavioral
reactivity, and sensorimotor function, recapitulating the pattern of symptoms reported in children
with ADHD.
Our findings demonstrate that developmental Mn exposure causes lasting dysfunction in attention,
impulse control, sensorimotor function, and behavioral reactivity (i.e., emotion regulation) – a
constellation of impairments commonly seen in children diagnosed with ADHD[2–4]. For example, across
the three attention tasks, the Mn-exposed animals are particularly impaired in the later session blocks of
testing (FA 1 task), and/or under trial conditions with the longest pre-cue delays (FA 2 task). The former
finding reflects that the Mn animals improve in attentional accuracy more slowly over testing than
controls, while the latter likely reflects deficits in sustained/focused attention at the most challenging
(longer) pre-cue delays. Similarly, the Mn exposed animals appear to be more impulsive than controls,
based on higher rates of premature responses across all three attention tasks (Figure 1). We should
note, however, that the pattern of %Premature responses across session blocks within each attention
task for the offspring of unsupplemented dams suggests that the higher rates of premature responses in
the Mn animals in some session blocks may be a result of Mn animals learning to withhold premature
responses more slowly than controls, rather than inherently greater impulsivity per se. Finally,
developmental Mn exposure also causes lasting impairments in sensorimotor function in the Montoya
staircase test – impairment that is most pronounced at the steps that are the most challenging to reach,
based on significant reductions in the number of pellets taken, pellets eaten, %Success, and increased
number of pellets misplaced (Figure 6). Collectively, these findings are fully consistent with our multiple
prior studies[18–24], and they further implicate elevated developmental Mn exposure as a significant risk
factor for environmentally-based ADHD and related symptoms[7,8,17,9–16].
688
Notably, we also report new findings showing that developmental Mn exposure leads to lasting
dysregulation in behavioral reactivity, an essential functional domain comparable to emotion regulation in
children[3,4]. Here, we evaluated behavioral reactivity by assessing how attentional performance was
affected by either (a) committing an error on the prior trial, or (b) not receiving an expected food pellet
reward. As expected from prior studies [33,67,71–74], we found that control animals perform significantly more
poorly on trials following an error (relative to after a correct response), and that they slow their response
time on trials following an error (Figure 4, 5A, B), a phenomenon called "post-error slowing" [33,67,71–77].
This pattern of changes likely reflects the fact that processing an error diverts attentional resources,
leading to both impaired performance and longer response latencies on post-error trials[78]. In contrast, for
the Mn animals, committing an error on the prior trial did not affect their %Correct responses, and had
only a small effect on their correct response latency (Figure 4, 5A, B). These findings indicate that the
Mn-exposed animals may be less attentive to performance errors (i.e., negative feedback), and have
diminished capacity to learn from trial history and adjust their behavioral strategy accordingly, suggesting
impairment in performance monitoring and reactive control[75]. The pattern seen in the Mn exposed
animals on trials following an error – less performance disruption and less post-error slowing than seen
in controls – mirrors findings showing that ADHD children are less responsive to committing errors than
typically developing children[75], and exhibit diminished post-error slowing[79,80]. Consistent with these
findings, neuroimaging results have revealed that children with ADHD have significantly weaker error-
related activation in the salience network than typically developing children[81], suggesting that ADHD
children may be less sensitive to detecting the saliency of their errors.
709
710 Interestingly, in contrast to their reduced sensitivity to committing an error, the unsupplemented Mn
711 exposed animals exhibit a heightened reaction to the omission of an expected food reward on the prior
712 trial, relative to that seen in controls (Figure 5A, B). The unsupplemented Mn animals exhibit a significant
713 drop in %Correct performance on trials following a reward omission, compared to their performance after

a correct response (Figure 5A, B), a pattern not seen in controls. This finding suggests that the Mn animals are more sensitive to reward (and its omission) than controls, a pattern which is again reminiscent of findings with ADHD children. Specifically, studies show that ADHD children often prefer smaller, more immediate rewards over larger, long-term rewards, are more sensitive than typically-developing children to the positive effects of a reward[68,82]; and are more affected by the loss of an anticipated reward[70]. This pattern thus shows clear parallels with our current findings that the Mn rats exhibit greater drops in performance on trials following a reward omission than do controls.

Consistent with these findings, the Mn-exposed animals commit a higher incidence of perseverative responses on trials following a correct response than do controls. Specifically, while animals across all treatment groups commit more perseverative responses on trials following a correct rewarded response trial than on trials following an error or omitted reward trial, this increase is more pronounced for the Mn animals than controls (Figure 5C), a pattern also seen for food pellet reward collection latency (Figure 5D). These findings provide additional support for the inference that the Mn exposed animals are more sensitive to reward/reinforcement than controls, a pattern also seen in ADHD children (noted above). The findings also support that the Mn animals are more compulsive than controls, a suggestion that is again consistent with the suite of symptoms reported in ~10 – 25% of ADHD children that are co-morbid for both ADHD and Obsessive Compulsive Disorder (OCD)[83–87].

Collectively, our findings demonstrate a constellation of behavioral/cognitive impairments in developmentally Mn-exposed animals that are also seen in Mn exposed children and those with ADHD more generally; namely, impairments in attention, impulse control, and sensorimotor function, as well as a reduced response to committing an error and a heightened sensitivity to reward/reward omission[3,4,68,69]. These latter *new* findings relating to Mn-induced dysfunction in behavioral reactivity/emotion regulation and error monitoring are highly relevant clinically, since the ability to maintain emotional and cognitive functioning in the face of unexpected events, as well as to learn from trial history and performance and adjust behavioral strategy accordingly, provides benefits in a variety of situations[81,88,89] . These findings are also clinically relevant in light of evidence showing that emotion dysregulation occurs in ~25-45% of children and ~30-70% of adults with ADHD, and that emotional problems have a greater negative impact on well-being and quality of life than other ADHD symptoms such as hyperactivity and inattention[3–5]. Thus, our rat model of environmental Mn exposure, used here and in prior studies[18–24], exhibits the hallmark areas of dysfunction seen in the ADHD phenotype[2].

4.2 MCS offers some protection against the adverse effects of early developmental Mn exposure.
Supplementing the maternal diet with additional choline during pregnancy and lactation offers protection against some of the effects of developmental Mn exposure, although the degree of benefit varied by cognitive domain. Two areas where MCS significantly benefited the Mn animals are in the realms of behavioral reactivity/emotion regulation and attention. In the realm of behavioral reactivity, the pattern of findings indicates that MCS offers protection against the effects of Mn exposure on reward sensitivity and reaction to errors. Specifically, animals born to dams on the choline-supplemented diet and then exposed to Mn in the early postnatal period did not exhibit a drop in performance on trials following a reward omission (similar to the unexposed unsupplemented controls), and very different from the Mn animals born to dams on the unsupplemented diet (Figure 5A). In addition, MCS partially normalized the reaction to committing an error for the Mn animals; this was seen for %Correct in the second focused attention task where the drop in performance after an error was similar for the MCS control and Mn animals. In addition, in the reward omission task, MCS normalizes correct response latency on trials following an error for the Mn animals; that is, the Mn+MCS animals exhibit the same degree of post-error slowing to that seen in the control+MCS animals (Figure 5B). Finally, MCS reduced perseverative responding of the Mn animals in the reward omission task on trials after a correct (rewarded) response trial to levels below the control+MCS animals (Figure 5C), and this effect of MCS was accompanied by a reduction in food pellet reward collection latency (Figure 5D). Together, these findings support that MCS normalized compulsivity and reward omission sensitivity in the Mn animals, and partially normalized error reactivity.

Maternal choline supplementation also offered protection against the detrimental effects of Mn exposure
on attentional function. This normalization of attentional function is seen in the second focused attention
task, where MCS lessens the impairment of the Mn animals at the longer and more challenging pre-cue
delays (Figure 2C and D). Similarly, in the selective attention task, MCS normalizes the performance of
the Mn animals on distraction trials, such that they no longer differ from controls (Figure 3). In contrast to
the protection provided by MCS for the adverse effects of Mn in the realms of behavioral reactivity and
attention, no protection was provided with respect to the impairments seen in learning and sensorimotor
function. In the first focused attention task, the Mn animals learned to wait for the visual cue more slowly
than controls, and MCS did not alter this Mn effect (Figure 2A, B). Further, the grasping and reaching
impairments of the Mn animals born to choline-supplemented dams was as pronounced as that seen in
the unsupplemented animals (Figure 6).

Collectively, these findings demonstrate that supplementing the maternal diet with additional choline
during pregnancy and lactation partially, but not completely, normalizes Mn deficits in the domains of
behavioral reactivity/emotion regulation, compulsivity, and attention, but not learning or sensorimotor
function. These neuroprotective effects are consistent with other studies that have shown that MCS
lessens the adverse cognitive effects of various developmental insults, including models of various
neurological disorders (e.g., Down syndrome. Alzheimer's disease, autism), toxicant exposure (e.g.,
alcohol), and other environmental influences (e.g., prenatal stress)[28,29,94–99,30–33,90–93]. For example, MCS
has been shown to lessen the functional deficits produced by developmental alcohol exposure in both
rodent models [28,35,96,98–100], and humans[36,101]. And of particular interest to the present study, MCS has
been shown to lessen the attentional dysfunction, as well as to normalize behavioral reactivity to
committing an error or not receiving an expected reward in a mouse model of Down syndrome[33].
Collectively, these studies provide additional support for increasing choline intake during pregnancy,
particularly for women at risk of toxicant exposure.

4.3 MCS benefits cognitive function in offspring not exposed to Mn
In addition to the benefits provided to the Mn exposed animals, supplementing the maternal diet with
additional choline was also beneficial for the control animals in the realms of behavioral reactivity,
attention, and learning. Specifically, for control animals, MCS lessens the disruptive effect of committing
an error on the prior trial (Figure 4, 5A; discussed above in section 4.2), suggesting an improved ability to
regulate the behavioral disruption/emotional reaction to committing an error. For attentional function, the
MCS benefit in control animals is especially evident at the 0 sec pre-cue delay in the second focused
attention task (Figure 2C), an effect which may also partly reflect this reduction in error reactivity because
trials with a 0 sec pre-cue delay are closest temporally to the prior trial, some of which were error trials.
In addition, trending effects for improved attentional performance were seen for the rate of improvement
across blocks in the first focused attention task (Figure 2A, B), as well as for performance in the reward
omission task on trials following a correct response (Figure 5A).

Altogether, these findings showing cognitive benefits of MCS in control animals are consistent with the
MCS benefits reported in prior studies in humans and animal models[38–41,43,46,47,49]. For example, a
number of studies by Meck, Williams and colleagues demonstrate that MCS improves attention and
memory in rodents[41–44]. Additionally, studies in humans have demonstrated that increased choline intake
during pregnancy produces lasting benefits for offspring attention, memory, and executive functioning [38–
40]. Notably, in one controlled feeding study that followed a cohort of children until 7 years of age, random
assignment of mothers to a higher choline intake (vs the AI) during the third trimester produced faster
processing speed during infancy[39], as well as superior memory, sustained attention, and executive
functioning at 7 years of age[38].

4.4 Putative neurobiological mechanisms
Our rodent model of developmental Mn exposure recapitulates many of the adverse behavioral
symptoms reported in ADHD children. While the neurobiological mechanisms underlying ADHD and
related symptoms are not well understood, studies suggest that the combination of multiple

environmental and biological risk factors lead to hypofunctioning of the catecholaminergic system within
the fronto-cortico-striatal loop and other neural circuits underlying executive attention and emotion
regulation[102–106]. Similarly, studies from our lab and others have demonstrated that developmental Mn
exposure causes hypofunctioning of the catecholaminergic system in the prefrontal cortex and
striatum[22–24], including reducing stimulated dopamine and norepinephrine release[24,25], and altered levels
of dopaminergic and noradrenergic system proteins[22–24]. The catecholaminergic system in the prefrontal
cortex/anterior cingulate cortex has been well-established as critical to executive functioning, including
attention, emotion regulation, error monitoring, and behavioral inhibition[107–113], and thus Mn-induced
alterations to this system may well underlie the ADHD-like behavioral phenotype of the Mn exposed
animals. This inference is further supported by studies from our group showing that oral methylphenidate
(a catecholamine reuptake inhibitor) ameliorates the Mn behavioral deficits in attention and sensorimotor
function[19,21,114]. Notably, our group has also demonstrated that developmental Mn exposure causes
lasting alterations in DNA methylation and expression of genes related to DNA methylation, neuronal
development and catecholaminergic neuronal systems in our rat model[115], and other studies have also
shown that Mn exposure can lead to altered DNA methylation[116–119]. Related to this, it is known that
choline serves as, among other things, a methyl donor in key biochemical reactions, and supplementing
the maternal diet with additional choline has been demonstrated to alter DNA methylation in both
humans and animal models[35,120–126]. Additionally, MCS has been shown to normalize alterations in DNA
methylation caused by fetal alcohol exposure[34,35]. Therefore, it is plausible that the benefits of MCS to
partially normalize Mn-induced behavioral reactivity and attention deficits in the present study may be
due in part to normalization of Mn-induced epigenetic alterations, though this suggestion is in need of
investigation.
842
4.5 Conclusion
The present study provides new evidence that developmental Mn exposure produces lasting
dysregulation of behavioral reactivity/emotion regulation, in addition to its detrimental effects on attention,
impulse control, and sensorimotor function. These findings largely recapitulate the constellation of
symptoms reported in Mn-exposed children, and separately in ADHD children, thus further supporting
concerns over developmental Mn exposure as a significant risk factor for ADHD and related symptoms.
In addition, the present study demonstrates that supplementing the maternal diet with additional choline
offers some protection to the adverse effects of early developmental Mn exposure, as well as provides
lasting cognitive benefits for the unexposed offspring. In light of the markedly increased physiological
demand for choline during pregnancy and the low choline intakes of most pregnant women, these results
provide additional support for efforts to increase choline intake during pregnancy, via diet and/or adding
choline to standard prenatal vitamin regimens.

855
Figure 1. Effects of Mn and MCS on premature responses (impulsivity) in the focused attention 1 task
(panel A), the focused attention 2 task (panels B and C), and the selective attention task (panel D), as a
function of session block or pre-cue delay (sec). Sub-panels a1, b1, and c1 depict significant two-way
interactions of Mn x session block or MCS x pre-cue delay, respectively. Mn exposure impaired the rate
of learning to wait for the cue. MCS did not alter premature responses generally, although there is a
suggestion that MCS lessened group differences in the selective attention task. * indicates p ≤ 0.05
versus controls. Data are lsmeans ± SEM (n=14-16/group).

Figure 2. Effects of Mn and MCS on %Correct and %Accurate responding in the two focused attention tasks. Panels A and B depict performance in the focused attention 1 task as a function of session blocks; sub-panel a1 depicts the significant two-way interaction of Mn x session block. Panels C and D depict performance in the focused attention task 2, as a function of the pre-cue delay (sec). Mn animals exhibit an impairment in focused attention, and MCS affects attention in both groups, but in different ways. * and & indicate $p \leq 0.05$ and $p \leq 0.10$ versus controls, respectively; # indicates $p \leq 0.05$ versus unsupplemented animals. Data are lsmeans ± SEM (n=14-16/group).

871
Figure 3: Effects of Mn and MCS on performance in the selective attention task. Panel A depicts
%Correct for the analysis in which all four odor distraction conditions were included separately, whereas
Panel B depicts %Correct responses for the analysis in which the four odor distraction conditions were
collapsed into distractor (odor) vs no distractor (no odor) conditions. Panel C depicts %Accurate
responses for all four distraction conditions. Sub-panels a1 and c1 depict significant two-way interactions
of Mn x odor distractor condition. Mn causes a deficit in attention, but not selective attention specifically,
and MCS reduces the difference between Mn and control animals under distraction conditions. *
indicates p ≤ 0.05 versus controls. Data are lsmeans ± SEM (n=14-16/group).

880
Figure 4: Effects of Mn and MCS on performance in the current trial as a function of the prior trial
outcome (i.e., correct or incorrect) in the second focused attention task for trials with a 0 sec pre-cue
delay. For animals born to dams on the unsupplemented diet (left side), Mn blunts reactivity to a prior
error. Maternal choline supplementation (right) normalizes error reactivity of the Mn animals. Prior trial
outcomes are Corr (correct response) or Error (incorrect, premature, or omission error). % and + indicate
$p \leq 0.05$ and $p \leq 0.10$ versus following a correct response, respectively; # indicates $p \leq 0.05$ versus
unsupplemented animals. Data are lsmeans ± SEM (n=14-16/group).

888
Figure 5: Effects of Mn and MCS on performance in the current trial as a function of the prior trial
outcome in the reward omission task. Panels A, B, C and D depict, respectively, %Correct, response
latency, perseverative responses, and food pellet collection latency. For the unsupplemented animals,
Mn blunts error reactivity, but heightens reactivity to an omitted reward. Maternal choline
supplementation (MCS) partially normalizes reactivity to both errors and reward omissions of the Mn
animals. Prior trial outcomes are Corr (correct response), Error (incorrect or omission error), or RO
(correct with reward omitted). % indicates p ≤ 0.05 versus following a correct response; * and & indicate
p ≤ 0.05 and p ≤ 0.10 versus controls, respectively; # and ^ indicates p ≤ 0.05 and p ≤ 0.10 versus
unsupplemented animals, respectively. Data are lsmeans ± SEM (n=14-16/group).

898
899 **Figure 6:** Effects of Mn exposure and MCS on sensorimotor function on the Montoya staircase task.
900 Panel A depicts the number of pellets eaten, whereas Panel B depicts number of pellets misplaced, both
901 as a function of step level. The insert panels show performance collapsed across the two diet conditions.
902 Mn exposure causes a lasting deficit in fine motor function that is not rescued by MCS. * indicates p ≤
903 0.05 versus controls. Data are lsmeans ± SEM (n=14-16/group).

Funding Statement
This study was supported by NIEHS grant #R01ES028369.

Conflicts of interest (COI)
The authors declare no actual competing or potential financial interests.

Acknowledgments
The authors would like to thank Bahtia Bilow, Charlyn Agbayani, David Woodfin, Eric Hsu, Gladys Chiang, Jonathan Do, Jordan Altshul, Katie Sharifi, Levon Shahnazarian, Naomi Schlipp, Neha Chenna, Ramtin Poustinchi, Roshana Basto, Sarah Hanson, Sasha (Aleksandra) Nikolaeva, Shannon Twardy, Sofia Bayliss Grijalva, Tyler Waite, Vanessa Howland, Victoria Deng, and Wendy Valdez Rodriguez for their invaluable assistance in behavioral testing. We also thank Tom Jursa, Dr. Marie Caudill, and Olga Malysheva for analytical assistance, Dr. Lynn Johnson for assistance with statistical analyses, and Dr. Nicholas Santiago for assistance in the study.

Author contributions
D.R.S., S.A.B., B.J.S., and S.L.H. conceived and designed research. S.L.H., S.A.B., and D.R.S. performed experiments. S.L.H., S.A.B., B.J.S., and D.R.S. analyzed and interpreted results of experiments. S.L.H. prepared figures. S.L.H., S.A.B., B.J.S., and D.R.S. drafted manuscript. S.L.H., S.A.B., B.J.S. and D.R.S. edited and revised manuscript.

References
1. Faraone, S. V. *et al.* The World Federation of ADHD International Consensus Statement: 208 Evidence-based conclusions about the disorder. *Neuroscience and Biobehavioral Reviews* **128**, 789–818 (2021).
2. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, 5th Edition.* (American Psychiatric Association, 2013).
3. Shaw, P., Stringaris, A., Nigg, J. & Leibenluft, E. Emotion dysregulation in attention deficit hyperactivity disorder. *American Journal of Psychiatry* **171**, 276–293 (2014).
4. Faraone, S. V. *et al.* Practitioner Review: Emotional dysregulation in attention-deficit/hyperactivity disorder – implications for clinical recognition and intervention. *Journal of Child Psychology and Psychiatry and Allied Disciplines* **60**, 133–150 (2019).
5. Riley, A. W. *et al.* Factors related to Health-Related Quality of Life (HRQoL) among children with ADHD in Europe at entry into treatment. *Eur. Child Adolesc. Psychiatry* **15**, i38–i45 (2006).
6. Wehmeier, P. M., Schacht, A. & Barkley, R. A. Social and Emotional Impairment in Children and Adolescents with ADHD and the Impact on Quality of Life. *J. Adolesc. Heal.* **46**, 209–217 (2010).
7. Schullehner, J. *et al.* Exposure to Manganese in Drinking Water during Childhood and Association with Attention-Deficit Hyperactivity Disorder: A Nationwide Cohort Study. *Environ. Health Perspect.* **128**, 1–10 (2020).
8. Lucchini, R. G. *et al.* Tremor, olfactory and motor changes in Italian adolescents exposed to historical ferro-manganese emission. *Neurotoxicology* **33**, 687–96 (2012).
9. Liu, W. *et al.* Biomarkers of environmental manganese exposure and associations with childhood neurodevelopment: A systematic review and meta-analysis. *Environ. Heal. A Glob. Access Sci. Source* **19**, 1–22 (2020).
10. Broberg, K. *et al.* Manganese transporter genetics and sex modify the association between environmental manganese exposure and neurobehavioral outcomes in children. *Environ. Int.* **130**, 104908 (2019).
11. Bauer, J. A. *et al.* Manganese in teeth and neurobehavior: Sex-specific windows of susceptibility. *Environ. Int.* **108**, 299–308 (2017).
12. Bauer, J. A. *et al.* Critical windows of susceptibility in the association between manganese and neurocognition in Italian adolescents living near ferro-manganese industry. *Neurotoxicology* **87**, 51 (2021).
13. Mora, A. M. *et al.* Prenatal and postnatal manganese teeth levels and neurodevelopment at 7, 9,

957   and 10.5 years in the CHAMACOS cohort. *Environ. Int.* **84**, 39 (2015).

958   14.   Haynes, E. N. *et al.* Manganese Exposure and Neurocognitive Outcomes in Rural School-Age
959   Children: The Communities Actively Researching Exposure Study (Ohio, USA). *Environ. Health*
960   *Perspect.* **123**, 1066 (2015).

961   15.   Oulhote, Y. *et al.* Neurobehavioral function in school-age children exposed to manganese in
962   drinking water. *Environ. Health Perspect.* **122**, 1343–50 (2014).

963   16.   Hong, S. B. *et al.* Blood manganese levels in relation to comorbid behavioral and emotional
964   problems in children with attention-deficit/hyperactivity disorder. *Psychiatry Res.* **220**, 418–425
965   (2014).

966   17.   Shin, D. W. *et al.* Association of Hair Manganese Level with Symptoms in Attention-
967   Deficit/Hyperactivity Disorder. *Psychiatry Investig.* **12**, 66 (2015).

968   18.   Beaudin, S. A., Nisam, S. & Smith, D. R. Early life versus lifelong oral manganese exposure
969   differently impairs skilled forelimb performance in adult rats. *Neurotoxicol. Teratol.* **38**, 36–45
970   (2013).

971   19.   Beaudin, S. A. *et al.* Oral methylphenidate alleviates the fine motor dysfunction caused by chronic
972   postnatal manganese exposure in adult rats. *Toxicol. Sci.* **144**, 318–327 (2015).

973   20.   Beaudin, S. A., Strupp, B. J., Strawderman, M. & Smith, D. R. Early postnatal manganese
974   exposure causes lasting impairment of selective and focused attention and arousal regulation in
975   adult rats. *Environ. Health Perspect.* **125**, 230–237 (2017).

976   21.   Beaudin, S. A. *et al.* Methylphenidate alleviates manganese-induced impulsivity but not
977   distractibility. *Neurotoxicol. Teratol.* **61**, 17–28 (2017).

978   22.   Kern, C. H., Stanwood, G. D. & Smith, D. R. Preweaning manganese exposure causes
979   hyperactivity, disinhibition, and spatial learning and memory deficits associated with altered
980   dopamine receptor and transporter levels. *Synapse* **64**, 363–78 (2010).

981   23.   Kern, C. H. & Smith, D. R. Preweaning Mn exposure leads to prolonged astrocyte activation and
982   lasting effects on the dopaminergic system in adult male rats. *Synapse* **65**, 532–544 (2011).

983   24.   Conley, T. E. *et al.* Early postnatal manganese exposure causes arousal dysregulation and lasting
984   hypofunctioning of the prefrontal cortex catecholaminergic systems. *J. Neurochem.* **153**, 631–649
985   (2019).

986   25.   Lasley, S. M. *et al.* Early Postnatal Manganese Exposure Reduces Rat Cortical and Striatal
987   Biogenic Amine Activity in Adulthood. *Toxicol. Sci.* **173**, 144 (2020).

988   26.   Dolinoy, D. C., Huang, D. & Jirtle, R. L. Maternal nutrient supplementation counteracts bisphenol
989   A-induced DNA hypomethylation in early development. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 13056–
990   61 (2007).

991   27.   Wright, K. N. *et al.* Methyl supplementation attenuates cocaine-seeking behaviors and cocaine-
992   induced c-Fos activation in a DNA methylation-dependent manner. *J. Neurosci.* **35**, 8948–58
993   (2015).

994   28.   Thomas, J. D., Idrus, N. M., Monk, B. R. & Dominguez, H. D. Prenatal choline supplementation
995   mitigates behavioral alterations associated with prenatal alcohol exposure in rats. *Birth Defects*
996   *Res. A. Clin. Mol. Teratol.* **88**, 827–37 (2010).

997   29.   Thomas, J. D., La Fiette, M. H., Quinn, V. R. . & Riley, E. P. Neonatal choline supplementation
998   ameliorates the effects of prenatal alcohol exposure on a discrimination learning task in rats.
999   *Neurotoxicol. Teratol.* **22**, 703–711 (2000).

30.   Schulz, K. M. *et al.* Dietary choline supplementation to dams during pregnancy and lactation
mitigates the effects of in utero stress exposure on adult anxiety-related behaviors. *Behav. Brain*
Res. **268**, 104–10 (2014).

31.   Ash, J. A. *et al.* Maternal choline supplementation improves spatial mapping and increases basal
forebrain cholinergic neuron number and size in aged Ts65Dn mice. *Neurobiol. Dis.* **70**, 32–42
(2014).

32.   Velazquez, R. *et al.* Maternal choline supplementation improves spatial learning and adult
hippocampal neurogenesis in the Ts65Dn mouse model of Down syndrome. *Neurobiol. Dis.* **58**,
92–101 (2013).

33.   Moon, J. *et al.* Perinatal Choline Supplementation Improves Cognitive Functionning and Emotional

Regulation in the Ts65Dn Mouse Model of Down Syndrome. *Behav. …* **124**, 346–361 (2010).

34. Bekdash, R. A., Zhang, C. & Sarkar, D. K. Gestational choline supplementation normalized fetal
alcohol-induced alterations in histone modifications, DNA methylation, and proopiomelanocortin
(POMC) gene expression in β-endorphin-producing POMC neurons of the hypothalamus. *Alcohol.*
Clin. Exp. Res. **37**, 1133–1142 (2013).

35. Otero, N. K. H., Thomas, J. D., Saski, C. A., Xia, X. & Kelly, S. J. Choline supplementation and
DNA methylation in the hippocampus and prefrontal cortex of rats exposed to alcohol during
development. *Alcohol. Clin. Exp. Res.* **36**, 1701–9 (2012).

36. Jacobson, S. W. *et al.* Efficacy of maternal choline supplementation during pregnancy in mitigating
adverse effects of prenatal alcohol exposure on growth and cognitive function: A randomized,
double-blind, placebo-controlled clinical trial. *Alcohol. Clin. Exp. Res.* **42**, 1327–1341 (2018).

37. Warton, F. L. *et al.* Maternal choline supplementation mitigates alcohol exposure effects on
neonatal brain volumes. *Alcohol. Clin. Exp. Res.* **45**, 1762 (2021).

38. Bahnfleth, C. L., Strupp, B. J., Caudill, M. A. & Canfield, R. L. Prenatal choline supplementation
improves child sustained attention: A 7-year follow-up of a randomized controlled feeding trial.
FASEB J. **36**, (2022).

39. Caudill, M. A., Strupp, B. J., Muscalu, L., Nevins, J. E. H. & Canfield, R. L. Maternal choline
supplementation during the third trimester of pregnancy improves infant information processing
speed: a randomized, double-blind, controlled feeding study. *FASEB J.* **32**, 2172–2180 (2017).

40. Boeke, C. E. *et al.* Choline intake during pregnancy and child cognition at age 7 years. *Am. J.*
Epidemiol. **177**, 1338–1347 (2013).

41. Meck, W. H. & Williams, C. L. Simultaneous temporal processing is sensitive to prenatal choline
availability in mature and aged rats. *Neuroreport* **8**, 3045–51 (1997).

42. Meck, W. H. & Williams, C. L. Characterization of the facilitative effects of perinatal choline
supplementation on timing and temporal memory. *Neuroreport* **8**, 2831–5 (1997).

43. Meck, W. H. & Williams, C. L. Metabolic imprinting of choline by its availability during gestation:
implications for memory and attentional processing across the lifespan. *Neurosci. Biobehav. Rev.*
27, 385–399 (2003).

44. Meck, W. H., Williams, C. L., Cermak, J. M. & Blusztajn, J. K. Developmental periods of choline
sensitivity provide an ontogenetic mechanism for regulating memory capacity and age-related
dementia. *Front. Integr. Neurosci.* **1**, (2008).

45. Mellott, T. J., Williams, C. L., Meck, W. H. & Blusztajn, J. K. Prenatal choline supplementation
advances hippocampal development and enhances MAPK and CREB activation. *FASEB J.* **18**,
545–547 (2004).

46. Buhusi, C. V., Lamoureux, J. A. & Meck, W. H. Prenatal choline supplementation increases
sensitivity to contextual processing of temporal information. *Brain Res.* **1237**, 204 (2008).

47. Cheng, R.-K., MacDonald, C. J., Williams, C. L. & Meck, W. H. Prenatal choline supplementation
alters the timing, emotion, and memory performance (TEMP) of adult male and female rats as
indexed by differential reinforcement of low-rate schedule behavior. *Learn. Mem.* **15**, 153–62
(2008).

48. Wong-Goodrich, S. J. E. *et al.* Spatial memory and hippocampal plasticity are differentially
sensitive to the availability of choline in adulthood as a function of choline supply in utero. *Brain*
Res. **1237**, 153–66 (2008).

49. Mohler, E. G., Meck, W. H. & Williams, C. L. Sustained Attention in Adult Mice is Modulated by
Prenatal Choline Availability. *Int. J. Comp. Psychol.* **14**, 136–150 (2001).

50. Wallace, T. C. & Fulgoni, V. L. Assessment of Total Choline Intakes in the United States. *J. Am.*
Coll. Nutr. (2016). doi:10.1080/07315724.2015.1080127

51. Wallace, T. C. & Fulgoni, V. L. Usual Choline Intakes Are Associated with Egg and Protein Food
Consumption in the United States. *Nutrients* **9**, (2017).

52. Wallace, T. C. *et al.* The underconsumed and underappreciated essential nutrient. *Nutr. Today* **53**,
240–253 (2018).

53. Abreu-Villaça, Y., Filgueiras, C. C. & Manhães, A. C. Developmental aspects of the cholinergic
system. *Behav. Brain Res.* **221**, 367–378 (2011).

54.   Korsmo, H. W., Jiang, X. & Caudill, M. A. Choline: Exploring the growing science on its benefits for
1064   moms and babies. *Nutrients* **11**, (2019).
1065   55.   Institute of Medicine (US) Food and Nutrition Board. *Dietary Reference Intakes for Thiamin,*
1066   *Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline.*
1067   (National Academies Press , 1998).
1068   56.   National Research Council. *Guide for the Care and Use of Laboratory Animals, Eighth Edition.*
1069   (The National Academic Press, 2011).
1070   57.   Stangle, D. E. *et al.* Succimer Chelation Improves Learning, Attention, and Arousal Regulation in
1071   Lead-Exposed Rats but Produces Lasting Cognitive Impairment in the Absence of Lead Exposure.
1072   *Environ. Health Perspect.* **115**, 201 (2007).
1073   58.   Asinof, S. K. & Paine, T. A. The 5-Choice Serial Reaction Time Task: A Task of Attention and
1074   Impulse Control for Rodents. *J. Vis. Exp.* 51574 (2014). doi:10.3791/51574
1075   59.   Paterson, N. E., Wetzler, C., Hackett, A. & Hanania, T. Impulsive action and impulsive choice are
1076   mediated by distinct neuropharmacological substrates in rat. *Int. J. Neuropsychopharmacol.* **15**,
1077   1473–1487 (2012).
1078   60.   Whishaw, I. Q., Woodward, N. C., Miklyaeva, E. & Pellis, S. M. Analysis of limb use by control rats
1079   and unilateral DA-depleted rats in the Montoya staircase test: movements, impairments and
1080   compensatory strategies. *Behav. Brain Res.* **89**, 167–177 (1997).
1081   61.   Takser, L., Mergler, D., Hellier, G., Sahuquillo, J. & Huel, G. Manganese, Monoamine Metabolite
1082   Levels at Birth, and Child Psychomotor Development. *Neurotoxicology* **24**, 667–674 (2003).
1083   62.   Montoya, C. P., Campbell-Hope, L. J., Pemberton, K. D. & Dunnett, S. B. The "staircase test": a
1084   measure of independent forelimb reaching and grasping abilities in rats. *J. Neurosci. Methods* **36**,
1085   219–228 (1991).
1086   63.   Kloth, V., Klein, A., Loettrich, D. & Nikkhah, G. Colour-coded pellets increase the sensitivity of the
1087   staircase test to differentiate skilled forelimb performances of control and 6-hydroxydopamine
1088   lesioned rats. *Brain Res. Bull.* **70**, 68–80 (2006).
1089   64.   Holm, P. I., Ueland, P. M., Kvalheim, G. & Lien, E. A. Determination of Choline, Betaine, and
1090   Dimethylglycine in Plasma by a High-Throughput Method Based on Normal-Phase
1091   Chromatography–Tandem Mass Spectrometry. *Clin. Chem.* **49**, 286–294 (2003).
1092   65.   Koc, H., Mar, M. H., Ranasinghe, A., Swenberg, J. A. & Zeisel, S. H. Quantitation of choline and
1093   its metabolites in tissues and foods by liquid chromatography/electrospray ionization-isotope
1094   dilution mass spectrometry. *Anal. Chem.* **74**, 4734–4740 (2002).
1095   66.   Ruiz, J., Antequera, T., Andres, A. I., Petron, M. J. & Muriel, E. Improvement of a solid phase
1096   extraction method for analysis of lipid fractions in muscle foods. *Anal. Chim. Acta* **1**–**2**, 201–205
1097   (2004).
1098   67.   Beaudin, S. A., Stangle, D. E., Smith, D. R., Levitsky, D. A. & Strupp, B. J. Succimer chelation
1099   normalizes reactivity to reward omission and errors in lead-exposed rats. *Neurotoxicol. Teratol.*
1100   **29**, 188–202 (2007).
1101   68.   Luman, M., Oosterlaan, J. & Sergeant, J. A. The impact of reinforcement contingencies on AD/HD:
1102   A review and theoretical appraisal. *Clin. Psychol. Rev.* **25**, 183–213 (2005).
1103   69.   Shiels, K. & Hawk, L. W. Self-regulation in ADHD: The role of error processing. *Clinical*
1104   *Psychology Review* **30**, 951–961 (2010).
1105   70.   Douglas, V. I. & Parry, P. A. Effects of reward and nonreward on frustration and attention in
1106   attention deficit disorder. *J. Abnorm. Child Psychol.* **22**, 281–302 (1994).
1107   71.   Gendle, M. H. *et al.* Impaired sustained attention and altered reactivity to errors in an animal
1108   model of prenatal cocaine exposure. *Dev. Brain Res.* **147**, 85–96 (2003).
1109   72.   Gendle, M. H. *et al.* Enduring Effects of Prenatal Cocaine Exposure on Selective Attention and
1110   Reactivity to Errors: Evidence from an Animal Model. *Behav. Neurosci.* **118**, 290–297 (2004).
1111   73.   Morgan, R. E. *et al.* Early lead exposure produces lasting changes in sustained attention,
1112   response initiation, and reactivity to errors. *Neurotoxicol. Teratol.* **23**, 519–531 (2001).
1113   74.   Rushworth, M. F. S., Hadland, K. A., Gaffan, D. & Passingham, R. E. The effect of cingulate
1114   cortex lesions on task switching and working memory. *J. Cogn. Neurosci.* **15**, 338–353 (2003).
75.   Cai, W. *et al.* Both reactive and proactive control are deficient in children with ADHD and

predictive of clinical symptoms. *Transl. Psychiatry* **13**, (2023).

76. Rabbitt, P. M. Errors and error correction in choice-response tasks. *J. Exp. Psychol.* **71**, 264–272
(1966).

77. Danielmeier, C. & Ullsperger, M. Post-error adjustments. *Front. Psychol.* **2**, (2011).

78. Notebaert, W. *et al.* Post-error slowing: An orienting account. *Cognition* **111**, 275–279

79. Balogh, L. & Czobor, P. Post-Error Slowing in Patients With ADHD: A Meta-Analysis. *J. Atten.*
Disord. **20**, 1004–1016 (2016).

80. Shiels, K., Tamm, L. & Epstein, J. N. Deficient post-error slowing in children with ADHD is limited
to the inattentive subtype. *J. Int. Neuropsychol. Soc.* **18**, 612–617 (2012).

81. Cai, W., Griffiths, K., Korgaonkar, M. S., Williams, L. M. & Menon, V. Inhibition-related modulation
of salience and frontoparietal networks predicts cognitive control ability and inattention symptoms
in children with ADHD. *Mol. Psychiatry 2019 268* **26**, 4016–4025 (2021).

82. Marx, I., Höpcke, C., Berger, C., Wandschneider, R. & Herpertz, S. C. The Impact of Financial
Reward Contingencies on Cognitive Function Profiles in Adult ADHD. *PLoS One* **8**, (2013).

83. Brem, S., Grünblatt, E., Drechsler, R., Riederer, P. & Walitza, S. The neurobiological link between
OCD and ADHD. *Atten. Defic. Hyperact. Disord.* **6**, 175 (2014).

84. Sheppard, B. *et al.* ADHD prevalence and association with hoarding behaviors in childhood-onset
OCD. *Depress. Anxiety* **27**, 667–674 (2010).

85. Masi, G. *et al.* Comorbidity of obsessive-compulsive disorder and attention-deficit/hyperactivity
disorder in referred children and adolescents. *Compr. Psychiatry* **47**, 42–47 (2006).

86. Masi, G. *et al.* A Naturalistic Exploratory Study of the Impact of Demographic, Phenotypic and
Comorbid Features in Pediatric Obsessive-Compulsive Disorder. *Psychopathology* **43**, 69–78
(2010).

87. Walitza, S. *et al.* Children and adolescents with obsessive-compulsive disorder and comorbid
attention-deficit/hyperactivity disorder: Preliminary results of a prospective follow-up study. *J.*
Neural Transm. **115**, 187–190 (2008).

88. Russo, S. J., Murrough, J. W., Han, M. H., Charney, D. S. & Nestler, E. J. Neurobiology of
resilience. *Nat. Neurosci. 2012 1511* **15**, 1475–1484 (2012).

89. Schutte, N. S., Schuettpelz, E. & Malouff, J. M. Emotional Intelligence and Task Performance.
Imagin. Cogn. Pers. **20**, 347–354 (2001).

90. Powers, B. E. *et al.* Maternal choline supplementation in a mouse model of Down syndrome:
Effects on attention and nucleus basalis/substantia innominata neuron morphology in adult
offspring. *Neuroscience* **340**, 501–514 (2017).

91. Strupp, B. J. *et al. Maternal choline supplementation: A potential prenatal treatment for Down*
syndrome and Alzheimer's disease. *Curr Alzheimer Res* **10962**, (Nathan Kline Institute, 2016).

92. Langley, E. A., Krykbaeva, M., Blusztajn, J. K. & Mellott, T. J. High maternal choline consumption
during pregnancy and nursing alleviates deficits in social interaction and improves anxiety-like
behaviors in the BTBR T + Itpr3tf/J mouse model of autism. *Behav. Brain Res.* **278**, 210–220
(2015).

93. Holmes, G. L. *et al.* Seizure-induced memory impairment is reduced by choline supplementation
before or after status epilepticus. *Epilepsy Res.* **48**, 3–13 (2002).

94. Velazquez, R. *et al.* Maternal choline supplementation ameliorates Alzheimer's disease pathology
by reducing brain homocysteine levels across multiple generations. *Mol. Psychiatry* 1 (2019).
doi:10.1038/s41380-018-0322-z

95. Thomas, J. D., Biane, J. S., O'Bryan, K. A., O'Neill, T. M. & Dominguez, H. D. Choline
supplementation following third-trimester-equivalent alcohol exposure attenuates behavioral
alterations in rats. *Behav. Neurosci.* **121**, 120–130 (2007).

96. Thomas, J. Perinatal choline supplementation attenuates behavioral alterations associated with
neonatal alcohol exposure in rats. *Neurotoxicol. Teratol.* **26**, 35–45 (2004).

97. Alldred, M. J., Pidikiti, H., Heguy, A., Roussos, P. & Ginsberg, S. D. Basal forebrain cholinergic
neurons are vulnerable in a mouse model of Down syndrome and their molecular fingerprint is
rescued by maternal choline supplementation. *FASEB J.* **37**, (2023).

98. Idrus, N. M., Breit, K. R. & Thomas, J. D. Dietary choline levels modify the effects of prenatal

alcohol exposure in rats. *Neurotoxicol. Teratol.* **59**, 43–52 (2017).

99.   Baker, J. A., Breit, K. R., Bodnar, T. S., Weinberg, J. & Thomas, J. D. Choline Supplementation
Modifies the Effects of Developmental Alcohol Exposure on Immune Responses in Adult Rats.
Nutrients **14**, (2022).

100.   Thomas, J. D. Choline Supplementation as an Intervention for Fetal Alcohol Spectrum Disorders:
A commentary. *Alcohol. Clin. Exp. Res.* **45**, 2465 (2021).

101.   Coles, C. D. *et al.* Dose and Timing of Prenatal Alcohol Exposure and Maternal Nutritional
Supplements: Developmental Effects on 6-Month-Old Infants. doi:10.1007/s10995-015-1779-x

102.   Faraone, S. V. *et al.* Attention-deficit/hyperactivity disorder. *Nat. Rev. Dis. Prim. 2015 11* **1**, 1–23
(2015).

103.   Hinshaw, S. P. Attention Deficit Hyperactivity Disorder (ADHD): Controversy, Developmental
Mechanisms, and Multiple Levels of Analysis. *https://doi.org/10.1146/annurev-clinpsy-050817-*
084917 **14**, 291–316 (2018).

104.   Smith, D. R. & Strupp, B. J. Animal Models of Childhood Exposure to Lead or Manganese:
Evidence for Impaired Attention, Impulse Control, and Affect Regulation and Assessment of
Potential Therapies. *Neurother. 2023 201* **20**, 3–21 (2023).

105.   Arnsten, A. F. T. The Emerging Neurobiology of Attention Deficit Hyperactivity Disorder: The Key
Role of the Prefrontal Association Cortex. *J Pediatr* **154**, 43 (2009).

106.   Posner, J. *et al.* Dissociable attentional and affective circuits in medication-naïve children with
attention-deficit/hyperactivity disorder. *Psychiatry Res.* **213**, 24 (2013).

107.   Arnsten, A. F. T. & Li, B. M. Neurobiology of executive functions: Catecholamine influences on
prefrontal cortical functions. *Biological Psychiatry* **57**, 1377–1384 (2005).

108.   Arnsten, A. F. T. Catecholamine Influences on Dorsolateral Prefrontal Cortical Networks. *Biol.*
Psychiatry **69**, e89–e99 (2011).

109.   Egner, T. Surprise! A unifying model of dorsal anterior cingulate function? *Nat. Neurosci.* **14**,
1219–1220 (2011).

110.   Elston, T. W., Kalhan, S. & Bilkey, D. K. Conflict and adaptation signals in the anterior cingulate
cortex and ventral tegmental area. *Sci. Rep.* **8**, (2018).

111.   Elston, T. W., Croy, E. & Bilkey, D. K. Communication between the Anterior Cingulate Cortex and
Ventral Tegmental Area during a Cost-Benefit Reversal Task. *Cell Rep.* **26**, 2353-2361.e3 (2019).

112.   Holroyd, C. B. & Coles, M. G. H. The neural basis of human error processing: Reinforcement
learning, dopamine, and the error-related negativity. *Psychol. Rev.* **109**, 679–709 (2002).

113.   Jocham, G. & Ullsperger, M. Neuropharmacology of performance monitoring. *Neuroscience and*
Biobehavioral Reviews **33**, 48–60 (2009).

114.   Beaudin, S. A., Howard, S., Santiago, N., Strupp, B. J. & Smith, D. R. Methylphenidate alleviates
attentional and sensorimotor dysfunction induced by developmental manganese exposure:
Assessment of catecholaminergic receptor involvement. doi:10.1101/2023.06.27.546786

115.   Santiago, N. A. *et al.* Developmental Manganese Exposure Causes Lasting Attention Deficits
Accompanied by Dysregulation of mTOR Signaling and Catecholaminergic Gene Expression in
Brain Prefrontal Cortex. *bioRxiv* 2023.07.16.549215 (2023). doi:10.1101/2023.07.16.549215

116.   Tarale, P. *et al.* Global DNA methylation profiling of manganese-exposed human neuroblastoma
SH-SY5Y cells reveals epigenetic alterations in Parkinson's disease-associated genes. *Arch.*
Toxicol. **91**, 2629–2641 (2017).

117.   Guo, Z. *et al.* Manganese chloride induces histone acetylation changes in neuronal cells: Its role in
manganese-induced damage. *Neurotoxicology* **65**, 255–263 (2018).

118.   Wang, L. *et al.* Aberration in Epigenetic Gene Regulation in Hippocampal Neurogenesis by
Developmental Exposure to Manganese Chloride in Mice. *Toxicol. Sci.* (2013).
doi:10.1093/toxsci/kft183

119.   Maccani, J. Z. J. *et al.* DNA Methylation Changes in the Placenta Are Associated With Fetal
Manganese Exposure . *Reprod. Toxicol.* **57**, 43–49 (2015).

120.   M. Kelley, C. *et al.* Effects of Maternal Choline Supplementation on the Septohippocampal
Cholinergic System in the Ts65Dn Mouse Model of Down Syndrome. *Curr. Alzheimer Res.* (2016).

121.   Kwan, S. T. C. *et al.* Maternal choline supplementation during normal murine pregnancy alters the

placental epigenome: Results of an exploratory study. *Nutrients* **10**, 417 (2018).
122. Medici, V. *et al.* Maternal choline modifies fetal liver copper, gene expression, DNA methylation,
and neonatal growth in the tx-j mouse model of Wilson disease. *Epigenetics* **9**, 286–296 (2014).
123. Kovacheva, V. P. *et al.* Gestational choline deficiency causes global and Igf2 gene DNA
hypermethylation by up-regulation of Dnmt1 expression. *J. Biol. Chem.* **282**, 31777–88 (2007).
124. Waterland, R. A. *et al.* Maternal methyl supplements increase offspring DNA methylation at Axin
fused. *genesis* **44**, 401–406 (2006).
125. Niculescu, M. D., Craciunescu, C. N. & Zeisel, S. H. Dietary choline deficiency alters global and
gene-specific DNA methylation in the developing hippocampus of mouse fetal brains. *FASEB J.*
20, 43–9 (2006).
126. Jiang, X. *et al.* Maternal choline intake alters the epigenetic state of fetal cortisol-regulating genes
in humans. *FASEB J.* **26**, 3563–3574 (2012).
1234