Maternal choline supplementation: A potential therapy for developmental Manganese exposure?

Authors

Shanna L. Howard[1], Stephane A. Beaudin[1], Barbara J. Strupp[2]*, Donald R. Smith[1]*

[1]Department of Microbiology and Environmental Toxicology, University of California, Santa Cruz, CA, USA
[2]Division of Nutritional Sciences and Department of Psychology, Cornell University, Ithaca, NY, USA
*Co-corresponding authors

Key terms

Manganese, maternal choline supplementation, attention deficit, sensorimotor, neurodevelopment

Highlights

1. The MCS intervention was partially, but not completely, effective in protecting against Mn-induced deficits; the degree of benefit varied with the various functional domains. Supplementing the maternal diet with choline during pregnancy and lactation provides some benefits to the Mn-exposed animals, including lessening differences between Mn and control animals in attentional function. MCS also partially normalizes the reaction of Mn exposed animals to committing an error or not receiving an expected reward.

2. This study has uncovered the effect of developmental Mn exposure to cause lasting abnormalities in behavioral reactivity to committing an error or not receiving an expected reward. Additionally, we have replicated the effects seen in our prior animal model studies in which Mn causes deficits in attention, learning, and sensorimotor function. The Mn deficits reported here parallel many of the behavioral deficits seen in children exposed to high levels of Mn during development, establishing developmental Mn exposure as an environmental risk factor for ADHD symptoms more broadly.

3. In the absence of Mn exposure, maternal choline supplementation produced lasting offspring benefits in the area of attentional function, recapitulating benefits of MCS demonstrated in other studies.

4. These findings underscore the lasting health risks to children posed by developmental Mn exposure and provide new evidence that MCS lessens some areas of Mn-induced dysfunction.

5. In light of evidence that 90% of pregnant women consume less choline than the current Adequate Intake (AI) levels, these data provide additional support for efforts to add choline to standard prenatal vitamin regimens.

Abstract

Background: Studies in children and adolescents have associated early developmental manganese (Mn) exposure with inattention, impulsivity, hyperactivity, and fine motor deficits, and our rodent studies of early life Mn exposure have recapitulated many of these outcomes, demonstrating causality. There are currently no recognized therapies/interventions other than exposure prevention to mitigate the neurotoxic effects of developmental Mn exposure. One possible preventive treatment is to supplement the maternal diet with additional choline during pregnancy. Maternal choline supplementation (MCS) has been shown to improve offspring cognitive function in humans and animal models, and lessens dysfunction caused by various developmental insults.

Objectives: Determine whether MCS during pregnancy and lactation protects against Mn-induced impairments in attention, impulse control, learning, behavioral reactivity, and sensorimotor function.

Methods: Starting at gestational day 3 (G3), pregnant dams were given standard diet or diet with additional choline (4x the level in standard diets) throughout gestation and lactation until offspring were weaned on PND 21. Pups were exposed orally to 0 or 50 mg Mn/kg/day during early postnatal life (PND 1–21). In adulthood, animals were tested in the five-choice serial reaction time task and the Montoya staircase task, to assess impulsivity, focused and selective attention, behavioral reactivity to errors or omission of an expected reward, and sensorimotor function.

Results: MCS intervention was partially effective in protecting against the Mn-induced deficits, with the degree of benefit varying with the specific domain of function. Specifically, MCS lessens differences between Mn and control animals in attentional function and reactivity to committing an error or not receiving an expected reward. MCS does not protect against Mn-induced sensorimotor dysfunction. Finally, in the absence of Mn exposure, MCS produces lasting benefits in attentional function and reactivity to errors.

Conclusions: MCS was partially effective in alleviating Mn-induced deficits – MCS normalized Mn exposed animals in the domains of attentional function and behavioral reactivity. These findings have implications for understanding the molecular mechanisms by which both MCS and Mn produce lasting cognitive changes, and provide further evidence that MCS benefits offspring. These findings, taken with evidence from other studies demonstrating the benefit of MCS to offspring and the fact that ~90% of women consume less than the adequate intake (AI) of choline, inform the recommendation that MCS should be considered for pregnant women.

1. Introduction

Attention and impulse control dysfunction, including ADHD, is the most prevalent neurodevelopmental syndrome among children, affecting ~5 - 7% of youths up through age 18 years, and ~2 – 3% of adults[1]. There is an emerging body of epidemiological evidence showing that environmental manganese (Mn) exposure during neurodevelopment is associated with increased risk of ADHD and/or attention/impulsivity/hyperactivity symptoms in children and adolescents[2,3,12,4–11]. However, while these epidemiologic studies have played key roles in highlighting associations between elevated Mn exposure and increased risk of ADHD-related symptoms, they are unable to establish causality, in part because of challenges identifying and/or controlling for confounding factors, and in accurately assessing exposure.

Our prior rodent model studies have shown that developmental Mn exposure can cause lasting ADHD-like deficits in attention, impulse control, and sensorimotor function, and that these deficits are associated with a hypofunctioning catecholaminergic system in the prefrontal cortex and striatum, the brain regions

80 mediating attentional, impulse control, and sensorimotor function[13–20]. Notably, gene expression of
81 several of the catecholaminergic system proteins altered by developmental Mn exposure are known to
82 be regulated in part by epigenetic mechanisms[21–25], and thus Mn-induced epigenetic alterations may
83 contribute to the lifelong alterations we see in these protein levels due to developmental Mn exposure.
84
While therapies such as methylphenidate (Ritalin) exist for children diagnosed with ADHD, there are no
established treatments available for children exposed to high levels of Mn during development. A number
of nutritional interventions, including maternal dietary supplementation with choline, folic acid, genistein,
and L-methionine, have been shown to mitigate the effects of a variety of environmental insults[26,27]. In
particular, maternal choline supplementation (MCS) has been shown in animal models to lessen
cognitive dysfunction caused by prenatal alcohol exposure[28–30], maternal stress[31], and Down
syndrome[32–34], and to normalize some gene expression and epigenetic molecular alterations caused by
perinatal alcohol exposure[35,36]. In addition, MCS has been shown to improve cognitive function of
typically developing human offspring[37–39] and animal models[40–48]. Consistent with this, current
recommendations for nutritionally sufficient levels of maternal choline intake may be inadequate,
particularly over periods of increased nutritional demand such as during fetal and neonatal development
(pregnancy and lactation)[49–51]. Notably, despite the fact that the adequate intake (AI) level of choline is
likely insufficient, over 90% of women do not meet this intake level of choline during pregnancy[39,49,50,52].
Thus, while numerous studies show that offspring may derive cognitive benefits from MCS, it is unknown
whether MCS provides protection against the neurocognitive deficits caused by developmental Mn
exposure, or whether elevated Mn exposure mitigates some of the cognitive benefits of MCS.
101
102 Here, we used a rodent model of elevated childhood Mn exposure to determine whether MCS protects
103 against the attention, impulse control, behavioral reactivity, and sensorimotor deficits caused by
104 developmental Mn exposure.
105
106   **2. Materials and methods**
107
Subjects
Sixty-four (64) male Long-Evans rats were used for neurobehavioral assessment. Additional littermates
were used for tissue Mn analysis and choline/choline metabolite analysis. Subjects were born in-house
from 16 nulliparous timed-pregnant dams (Charles River; gestational age 3; 8-9 weeks old). Twelve to 24
hours after parturition (designated PND 1, birth = PND 0), litters were weighed, sexed, and culled to eight
pups per litter such that each litter was composed of as many males as possible (2-8) and the remainder
females. Each treatment group contained n = 16 animals, and no more than 1-2 males per litter were
assigned to a particular treatment group. Animals were housed in polycarbonate cages at a constant
temperature of 21 ± 2°C. At PND 22, all pups were weaned and pair-housed with an animal of the same
treatment group and maintained on a reversed 12:12 hour light/dark cycle. All aspects of behavioral
testing and feeding were carried out during the active (dark) phase of the animals' diurnal cycle. All
animal care and treatments were approved by the institutional IACUC (protocol Smitd1803) and adhered
to National Institutes of Health guidelines set forth in the Guide for the Care and Use of Laboratory
Animals.
122
Maternal choline supplementation
Pregnant dams were randomly selected to receive one of two concentrations of choline chloride in their
diet (Dyets AIN-76A Purified Rodent Diet), both of which were provided ad libitum during gestation and

lactation, from G3 until PND 21. The unsupplemented purified diet food contained 1.1 g choline/kg, while
the supplemented purified diet food contained 5.0 g choline/kg; both are reported by the manufacturer to
contain 54 mg Mn/kg. Upon weaning on PND 21, all pups were provided Irradiated Teklad Global Soy
Protein-Free Extruded Rodent Diet 2920, reported by the manufacturer to contain 80 mg Mn/kg and 1200
mg choline/kg. See Figure 1 for a timeline of maternal choline supplementation, Mn dosing, and
behavioral testing.
132
Manganese exposure
Neonate animals were orally exposed to 0 or 50 mg Mn/kg/day in a vehicle solution (5% w/v stevia in
Milli-Q water) from PND 1–21, as in our prior studies[13–19]. Briefly, Mn was delivered once per day directly
into the mouth of each pup (~15-25 µL/dose) using a micropipette fitted with a flexible polyethylene
pipette tip. This Mn exposure regimen is relevant to children exposed to elevated Mn via drinking water,
diet, or both, as pre-weaning exposure to 50 mg Mn/kg/day produces relative increases in Mn intake that
approximates the increases reported in infants and young children exposed to Mn-contaminated water or
soy- based formulas (or both)[15,17].
141
Testing apparatus
Sixteen identical automated 5-Choice Serial Reaction Time Task (5-CSRTT) testing chambers fitted with
odor delivery systems (#MED-NP5L-OLF, Med Associates, Inc., St. Albans, VT) were used to assess
specific cognitive processes, including focused and selective attention, inhibitory control, and behavioral
reactivity, as described previously[15,16,53]. Briefly, each testing chamber contained a curved wall equipped
with five 2.5 × 2.5 cm response ports fitted with light-emitting diodes that served as the visual cue, an
infrared beam to register nose pokes, and pneumatic inlet and vacuum outlet ports to introduce and
remove air-based odor distractors. Opposite the response wall was the food magazine wall that
contained a 45 mg food pellet reward dispensing port fitted with an infrared beam to register nose pokes;
each unit also contained a small house light. The entire testing chamber was enclosed in a sound
attenuating cubicle. Each animal was assigned to one of the 16 testing chambers, and each chamber
was balanced for the four treatment conditions.
154
Behavioral testing
Animals were mildly food restricted starting on PND 45 in preparation for behavioral testing, with water
available ad libitum throughout behavioral assessment. Animals were provided food for 2 hours each day
immediately after behavioral testing, so that their body weights were maintained at ~90–95% of free-
feeding weights. All rats were weighed and tested 6 days/week throughout training and testing.
Behavioral assessment occurred during the active (dark) period of the diurnal cycle at the same time
each day and in the same chamber for each individual rat. A daily test session consisted of 150 trials or
50 min, whichever came first. Each trial sequence was initiated by a nose-poke in the food magazine
port, and followed by a 3 sec turnaround time to allow the animal to reorient from the food magazine wall
to the response wall; trial onset began after the 3 sec turnaround time. All behavioral testing was
conducted by individuals blind to the treatment condition of the subjects.
166
Behavioral testing began on ~ PND 50, as described elsewhere[15]. Briefly, testing began with food
magazine and nose-poke training for 11 days, followed by a 5-choice visual discrimination learning task
with a fixed 15 sec cue duration and no pre-cue delay. Animals stayed on this task until they reached the
criteria of 80% Correct trials on 2 of 3 consecutive testing days. Following this task, animals were moved
to a second visual discrimination task with a fixed 1 sec cue duration and no pre-cue delay for 5 days.

This training was followed by a series of attention and behavioral reactivity tasks as described below
(see Figure 1 for the timeline of testing).
174
5-CSRTT assessment is conducted via a series of progressively more difficult tasks, described in greater
detail below, that challenge the ability of animals to sustain attention, inhibit premature responses, and
ignore olfactory distractions in favor of the salient visual cue. Recorded response types for all 5-CSRTT
tests included the following: 1) premature responses (responses made after trial onset but before
presentation of the visual cue; in the first focused attention, second focused attention, and selective
attention tasks); 2) correct response (responses made to the correct port following presentation of the
visual cue); 3) incorrect response (responses made to the incorrect port following presentation of the
visual cue); and 4) omissions (failure to respond within the 15 sec response interval following the visual
cue). Premature and incorrect responses and omission errors were not rewarded and were immediately
followed by a 5 sec time-out, in which the house light was turned off for 5 sec. In addition, the latency
(sec) for correct and incorrect responses was recorded, as was the latency to retrieve the food pellet
reward following a correct response. Perseverative responses, defined as additional nose pokes into the
correct port after a correct response, were also recorded. The calculated response outcomes were %
Correct, calculated as number of correct responses/(correct + incorrect + premature + omissions) × 100;
% Incorrect, calculated as above but with incorrect responses in the numerator; % Premature, calculated
as number of premature responses/(correct + incorrect + premature + omissions) × 100; and %
Omissions, calculated as number of omissions/(correct + incorrect + premature + omissions) × 100.
Additionally, because %Correct includes all trial response outcomes in the denominator, all error types
(incorrect, premature, or omission) may contribute to differences between treatment groups. Therefore,
we also assessed attentional accuracy via the %Accurate outcome, which does not include premature or
omitted responses, and is calculated as number of correct responses/(correct + incorrect) × 100.
196
Focused attention tasks
Focused attention can be defined as the ability to maintain attentional focus on a specific task or stimulus
(e.g., a visual cue). Two successive focused attention tasks (FA1 and FA2) were administered over
~PND 85-103, and PND 104-121, respectively, following completion of the visual learning discrimination
task. Both of these tasks assessed the ability of animals to detect and respond to a visual cue presented
unpredictably in time and location (one of the five response ports). The first focused attention task used
variable pre-cue delays of 0, 1, 2, or 3 sec and a fixed visual cue duration of 0.7 sec and was
administered for 15 sessions. The second focused attention task included variable pre-cue delays of 0, 3,
4, or 5 sec and variable visual cue durations of 0.5 or 0.7 sec, and was administered for 15 sessions.
206
Selective attention task with olfactory distractors
Selective attention can be defined as the ability to maintain a behavioral or cognitive set in the face of
distracting or competing stimuli. The two selective attention tasks administered were the selective
attention baseline (SAB) task and the selective attention (SA) task with olfactory distractors, from PND
126-128 and 129-140, respectively. Animals were tested in the selective attention baseline task for 3
daily test sessions. In this baseline task, the pre-cue delay varied between 3 and 4 sec, with the two
delays balanced across the trials within each test daily session, and the cue duration was 0.5 sec. This
task was followed by 12 sessions of the selective attention task, which was identical to the baseline task
except that on one third of the trials in each session, an olfactory distractor was presented 1 or 2 sec
after trial onset (i.e., 1–3 sec before the visual cue). This delay between odor presentation and cue
illumination (1-3 sec) was referred to as the odor-to-cue interval (OtoC) and was included in statistical

analysis. The nine different olfactory distractors were made from liquid odorants (McCormick & Company, Inc.) diluted in propylene glycol, and delivered as scented air.

Assessment of behavioral reactivity and the reward omission task
Behavioral reactivity was assessed the second focused attention task, and again in the reward omission task. In the second focused attention task, performance (e.g., %Correct) on the current trial was assessed as a function of the outcome of the prior trail (correct or incorrect). Similarly, the reward omission task was included as an assessment of behavioral reactivity, but with the addition of an omitted reward on a subset of trials. Specifically, the food pellet reward was randomly omitted on three out of ten correct trials, and attentional performance on the current trial was assessed as a function of whether the preceding trial outcome was either (1) correct, (2) incorrect (including omission errors), or (3) correct but food reward withheld. This task used a 0.7 sec cue duration, and there was no pre-cue delay present. Animals were tested in the reward omission task for 12 sessions from PND 175-187.

Montoya Staircase Test
Montoya staircase test was used to assess sensorimotor function, as described in our prior studies[14]. In the Montoya Staircase test, the rat reaches down on the left or the right side of the central platform of the apparatus to grasp and retrieve food pellets from the descending steps of the staircase. Briefly, eight identical Plexiglas staircase devices were used, modeled after the original design of Montoya et al. (1991)[54]. Each device contained a platform and a left-and-right side staircase. The staircases contained 6 descending steps on each side, and each step was baited with three 45 mg food pellets for a total of 36 pellets available at the start of a session. To determine the step of origin and final location of each pellet, pellets were color coded with a unique color per step level[55].

The staircase study was conducted over 1 month, beginning on PND 85. Each staircase session occurred immediately following operant testing every day. Staircase training occurred for the first 11 days. During training, rats retrieved the food pellets from the descending steps of the staircases until all subjects reached asymptotic performance of total pellets retrieved. The next 19 days comprised testing phase of the task. Throughout training and testing, each subject was given one 10-min trial per day, 6 days per week. After each trial, the rat was removed, and the final distribution of colored pellets was quantified per step and side. The order and time of testing were balanced by treatment and remained the same each day for each rat.

Skilled forelimb performance was evaluated step-by-step for (1) the number of pellets taken (i.e., pellets that were grasped and removed from their step of origin), (2) the number of pellets eaten (pellets that were taken and consumed), (3) the corresponding percent of success (the ratio of the number of pellets eaten/pellets taken x 100), (4) the number of pellets misplaced (all the pellets that were grasped but ended up on a different step level than their step of origin), and (5) the number of pellets lost (pellets that ended up on the floor of the apparatus, out of reach to the animal).

Tissue Mn and choline/choline metabolite levels
For blood and brain Mn levels, tissue samples were collected from weaned pups on PND 24, and from the behaviorally-tested animals on PND 202. Briefly, animals were euthanized via CO_2 asphyxiation, and whole blood was immediately collected from the left ventricle of the surgically-exposed heart and stored in EDTA Vacutainers at − 20 °C for analyses. Whole brain was immediately removed, and the hind-brain region collected and stored at −80 °C for Mn concentration determinations. Blood and brain Mn

264 concentrations were determined by inductively coupled plasma–mass spectrometry (Thermo Element
265 XR), as previously described[13,15]. The analytical detection limit for Mn was 0.01 ng/mL.
266
Choline/choline metabolite levels were measured in plasma, brain, and/or liver tissues from dams at
weaning (plasma and liver), and/or from the same PND 24 animals (liver and hindbrain) used for Mn
analyses described above. Briefly, concentrations of free choline, betaine, dimethylglycine, methionine,
and trimethylamine N-oxide were measured by liquid chromatography tandem mass spectrometry
according to the method of Holm et al (2003)[56], while phosphocholine, glycerophosphocholine,
phosphatidylcholine, acetylcholine, and lysophosphatidylcholine were measured by liquid
chromatography/mass spectroscopy according to the method of Koc et al. (2002)[57] with modifications[58].
274

Statistical methods

The behavioral data were modeled using structured covariance mixed models. Fixed effects included in
the model were the between-subject factors MCS (two levels corresponding to the unsupplemented and
MCS groups), and Mn exposure treatment (two levels corresponding to the 0 and 50 mg/kg/d groups),
and the within-subject factors pre-cue delay, cue duration, session block, distraction condition, and/or
previous trial outcome, depending on the outcome analyzed. In all models, rat was included as a random
effect to account for correlations within observations from the same animal. Statistical tests used the
Kenward-Roger correction to improve estimation of the variance-covariance of the fixed effects
parameters. Plots of residuals by experimental condition were used to examine the assumption of
homogeneity. Random effects with high variance in the residuals across the levels of the factor (e.g.,
distraction condition) were added to the model to achieve homogeneity if necessary. The distribution of
each random effect was inspected for approximate normality and presence of influential outliers. The
significance level was set at $p \leq 0.05$, and p-values between 0.05 and 0.10 were considered to be
trending towards significance. Trending effects are presented if they aid in clarifying the nature of the Mn
and/or MCS effects. Significant main effects or interaction effects were followed by single-degree of
freedom contrasts to better understand the nature of the interactions, using the Student's t-test for
pairwise comparisons of least squared means. Analyses were conducted using SAS (version 9.4) for
Windows. For tissue Mn and choline/metabolite data, analyses were performed using one-way (for
dams) or two-way (for offspring) analysis of variance and Tukey's post hoc test for pairwise comparisons,
using JMP (version 16.0; SAS Institute, Inc.).
295

3. Results

297
For The 5-CSRTT measures, we focus primarily on outcomes of %Premature, %Correct, and %Accurate
responses because these measures have historically best differentiated control vs Mn animals, and did
so as well in this study. Results for each outcome are plotted in the way that best illustrates the Mn and
MCS effects. When there is a significant two-way but not three-way statistical interaction, we plotted both
the two-way (shown in an insert panel) and the three-way interaction to show the performance of each of
the four treatment groups under specific conditions due to our *a priori* hypotheses relating to the Mn
effect under both diet conditions.
305

3.1 Visual learning

307
Visual learning in the 5-choice task is not altered by Mn or MCS

Visual learning (i.e., learning to make a nosepoke in the illuminated port) is foundational to preparing the
animals for subsequent, more challenging 5-CSRTT tasks. Animals were trained on a visual
discrimination learning task over PND 58-79, and there was no effect of early postnatal Mn exposure or
MCS on visual learning. Specifically, there was no main effect of Mn or MCS on trials to criterion or days
to criterion, and no Mn x MCS interaction for these two outcomes (P's > 0.295). Moreover, there also
was no effect of Mn or MCS on the number of animals per treatment group to reach the learning criterion.
The lack of treatment group differences in learning the task criterion indicates that visual acuity and
motivation did not differ between groups and therefore cannot account for group differences seen in
subsequent 5-CSRTT tasks.
318
319 **3.2 Impulsivity**
320
321 ***Developmental Mn exposure did not affect impulse control, but did impair the animals in learning***
322 ***to wait for the visual cue, and this effect is not rescued by MCS***
323 We evaluated premature responses, a measure of impulse control in the 5-CSRTT, as a percent of total
324 responses in the two successive focused attention tasks and the selective attention task (SA).
325
In the first focused attention task, Mn animals are impaired in learning to withhold premature responses,
and MCS does not alter this effect.
In the focused attention 1 task, there is a significant Mn x session block interaction [$F_{(2, 435.8)}$ = 11.07,
p<0.0001] for %Premature responses. While both control and Mn groups commit a relatively high level of
premature responses at the start of the task (~ 50%), the Mn animals improve less rapidly than controls.
As a result, the Mn animals commit significantly more premature responses than controls in session
block 3 (p=0.0084; Figure 2a1). There is also a significant MCS x session block interaction [$F_{(2, 435.8)}$ =
17.06, p<0.0001], and while the MCS and unsupplemented groups don't statistically differ from each
other in any block, the interaction likely appears because MCS animals have slightly fewer premature
responses in block 1 but more premature responses in block 3. Refer to Figure 2A, which depicts
differences between Mn and controls under each of the two diet conditions (three-way interaction p =
0.69).
338
339 *In the second focused attention task, Mn and MCS don't interact to alter premature responses.*
340 In the subsequent focused attention task, there is a Mn x session block interaction [$F_{(2, 933.7)}$ = 5.53, p
341 = 0.0041], although there are no significant contrasts between control and Mn animals in any session
342 block (P's > 0.40; Figure 2b1). Refer to Figure 2B, which depicts differences between Mn and controls
343 under each of the two diet conditions (three-way interaction p = 0.40).
344
There is also a MCS x pre-cue delay interaction [$F_{(2, 932.8)}$ = 3.18, p = 0.0422], though again, none of
the specific contrasts between unsupplemented and MCS animals reached significance at any pre-cue
delay (P's > 0.21, Figure 2c1). However, MCS animals had non-significantly lower premature responses
across pre-cue delays compared to unsupplemented animals, with the apparent difference being
greatest at the shorter (and less challenging) 3 sec pre-cue delay. Refer to Figure 2C, which depicts
differences between Mn and controls under each of the two diet conditions (three-way interaction p =
0.24).
352
353 *In the selective attention task, MCS lessens the difference in premature responses between treatment*
354 *groups in early session blocks.*

In the selective attention task with olfactory distractors, there is a Mn x MCS x session block interaction for %Premature responses [F(3, 1665) = 4.44, p = 0.0041], although none of the specific contrasts between treatment groups or diet conditions reached significance for any session block (P's > 0.12, Figure 2D). Here, the significant three-way interaction arises because the difference in premature responses between Mn and control groups in the first two session blocks is greater for unsupplemented animals than for the MCS animals.

3.3 Attention

Mn impairs focused attention, and while MCS lessens the difference between control and Mn animals on focused attention, the nature of the MCS effect differs in control vs. Mn animals
As described earlier, the two focused attention tasks were used to assess whether Mn exposure or MCS alters the ability of animals to maintain attentional focus to detect the visual cue. We evaluated %Correct and %Accurate responses, which assess attentional function in the 5-CSRTT.

In the first focused attention task, developmental Mn impairs attention, and MCS alters the rate of performance improvement differently in control compared to Mn animals.
In the focused attention 1 task, there is a significant Mn x session block interaction for %Correct responses [F(2, 595.6) = 4.12, p = 0.0167]. This reflects that performance in the Mn animals improves more slowly than control animals across session blocks, such that the Mn animals have significantly fewer %Correct responses than the controls during blocks 2 and 3 (p=0.007 and 0.002, respectively, figure 3a1). Refer to Figure 3A, which depicts differences between Mn and controls under each of the two diet conditions (three-way interaction p = 0.22).

Consistent with the %Correct outcome noted above, there is a significant Mn x session block interaction for %Accurate responses [F(2, 601.3) = 3.33, p = 0.0365; not shown], and a trending three-way interaction of Mn x MCS x session block for %Accurate responses [F(2, 601.3) = 2.58, p = 0.0769] (Figure 3B). For %Accurate responses, the Mn deficit emerges under session blocks 2 and 3 (p's = 0.018 and 0.049 respectively under MCS; Figure 3B). This three-way interaction reflects the fact that the difference in performance of the Mn and controls was greater in the MCS condition than the unsupplemented condition; this appears to be because the controls benefitted more from the MCS than the Mn rats, in terms of learning rate.

In the second focused attention task, MCS improves performance of both the Mn and control animals, but in different functional domains.
In the focused attention 2 task, there is a significant Mn x MCS x pre-cue delay interaction for %Correct responses [F(3, 1277) = 7.32, p < 0.0001] (Figure 3C). For animals on the unsupplemented diet, the Mn-exposed rats have significantly fewer %Correct responses than controls at the 3 sec delay (p = 0.0477), and this Mn deficit is also trending at the 4 and 5 sec delays (p's = 0.0518 and 0.0699, respectively). However, in the MCS animals, the Mn deficit at the 3 sec delay is only trending (p = 0.0762), and the deficit is no longer present under the 4 and 5 sec pre-cue delays (p's = 0.27 and 0.69, respectively). Thus, the Mn x MCS x pre-cue delay interaction reflects the fact that the differences between control and Mn animals at the longer delays is smaller under MCS than under the unsupplemented diet. Additionally, control animals that received MCS have significantly more correct responses at the 0 sec delay compared to unsupplemented controls (p = 0.0113), reflecting a benefit of MCS to control animals for trials when the cue was presented immediately at trial onset, with no pre-cue delay.

401
We also assessed attentional accuracy for the second focused attention task, and again observed a
significant Mn x MCS x pre-cue delay interaction for %Accuracy [$F_{(3, 1271)}$ = 3.95, p = 0.0081] (Figure
3D), reflecting a pattern in attentional performance similar to what was seen in the %Correct outcome
described above. Under the unsupplemented diet, there is a trending Mn deficit at the 3 and 4 sec pre-
cue delays (p's = 0.058 and 0.067, respectively). In contrast, under MCS, this Mn deficit is still present at
the 3 sec delay (0.043), but not at the 4 sec pre-cue delay (p = 0.23). Although there was no significant
Mn deficit at the 5 sec pre-cue delay under the unsupplemented diet, MCS also appears to reduce the
difference between control and Mn animals at this longer delay (p = 0.11 under the unsupplemented diet
and 0.53 under MCS; Figure 3D).
411
412 While there was a significant benefit of MCS to controls at the 0 sec pre-cue delay for %Correct
413 responses (p = 0.0113), for %Accurate this benefit did not reach significance (p = 0.129). There was,
414 however, a trending Mn deficit that emerged at the 0 sec delay under MCS (p = 0.09, Figure 3D).
415
Collectively, this evidence suggests that the lasting attentional deficits caused by developmental Mn
exposure persist into the second focused attention task. Additionally, MCS improves performance of the
controls but not the Mn animals for trials with a 0 sec pre-cue delay, whereas it reduces the difference
between Mn and control animals at the longer delays. The MCS benefit derived by the controls at the 0
sec delay appears to be due to a lessened reaction to committing an error on the prior trial (discussed
below; Figure 5), whereas the benefit derived by the Mn animals at the longer delays appears to reflect
improved attention.
423
424 ***MCS reduces the Mn effect on distraction trials in the selective attention task***
425 To determine the effects of Mn exposure and MCS on selective attention, we assessed animal
426 performance on a selective attention task with olfactory distractors randomly presented either 1, 2, or 3
427 sec prior to the onset of the visual cue (referred to as the odor-to-cue interval, 'OtoC').
428
Mn causes a deficit in attention, but not selective attention specifically, and MCS reduces the difference
between Mn and control animals on trials with olfactory distractors.
For %Correct responses in the selective attention task, there was a significant Mn x OtoC interaction for
%Correct [$F_{(3, 239.2)}$ = 2.94, p = 0.0339], where the Mn animals were more impaired relative to controls
for trials without a distractor (p = 0.0096; Figure 4a1), than for distraction trials. The Mn x MCS x OtoC
interaction for %Correct responses was not significant (p = 0.269) (Figure 4A). However, qualitatively
based on a plot of this interaction, it appears that under the MCS diet, Mn and control animals have
comparable performance under the three distraction conditions (Figure 4A). In light of this, we conducted
follow-up analyses where the three trial conditions with olfactory distractors (i.e., -1, -2, -3 OtoC) were
collapsed into a single "distraction" trial condition, thus creating only two distraction conditions (distractor
and no-distractor trials; Figure 4B). In this follow-up analysis, there was a trending interaction of Mn x
MCS x distractor [$F_{(1, 52.16)}$ = 3.29, p = 0.0754]. This trend reflects that for nondistraction trials there is
a Mn deficit under both diets, but for distraction trials there is only a Mn deficit for unsupplemented
animals (Figure 4B).
443
For %Accurate responses, there was a significant Mn x OtoC interaction [$F_{(3, 235.7)}$ = 3.14, p = 0.026],
where the Mn deficit was significant for trials without a distractor and trending for trials in the -3 sec
distractor condition (p's = 0.0084 and 0.098; Figure 4c1); the groups were not different for trials with a -1

or -2 distractor. The three way interaction (Mn x MCS x OtoC) was not significant [F(3, 235.8) = 0.8, p =
0.495]; however, we see a similar pattern of group differences as noted above for %Correct, where MCS
reduces the difference in %Accuracy between control and Mn animals under the distractor conditions
(Fig 4C). However, neither the original (Figure 4C) nor collapsed (not shown) %Accurate models
revealed a significant Mn x MCS x odor interaction (p's > 0.31).
452
453   Collectively, this evidence demonstrates that MCS lessens the difference between control and Mn
454   animals for trials with olfactory distractors.
455
456   **3.4 Behavioral reactivity**
457
458   **Mn alters behavioral reactivity compared to controls, and MCS differentially affects control vs Mn**
459   **animals in this domain**
460   To determine if developmental Mn exposure and/or MCS alter reactivity to committing an error or not
461   receiving an expected reward, we analyzed performance outcomes on the current trial based on the
462   outcome of the preceding trial for the second focused attention and reward omission tasks.
463
For the second focused attention task, control animals that received MCS had significantly more
%Correct responses than their unsupplemented counterparts at the 0 sec pre-cue delay (reported above
in Figure 3C). We hypothesized that this might be due to MCS reducing the behavioral reaction to
making an error on the prior trial, as trials without a pre-cue delay are closest temporally to the prior trial.
If the outcome of the prior trial was disruptive to the animal, it would be most apparent for trials for which
the cue was presented immediately at trial onset. Therefore, we assessed %Correct responses on
current trials with a 0 sec pre-cue delay, as a function of the outcome of the prior trial.
471
472   The animal's reaction to committing an error on the prior trial was also assessed in a subsequent Reward
473   Omission task designed to assess behavioral reactivity to two different types of potentially disruptive
474   events: (1) making an error on the previous trial (accompanied by a timeout and no reward), or (2)
475   making a correct response but not receiving an expected reward (see Methods). The results of both
476   tasks are described below.
477
In the second focused attention task trials with a 0 sec delay, MCS improves performance in control
animals following an error, and Mn animals do not exhibit behavioral reactivity to an error under
unsupplemented or MCS diet.
In the second focused attention task, analysis revealed a significant Mn x MCS x previous trial outcome
interaction for trials with a 0 sec pre-cue delay [F(1, 273.2) = 6.13, p = 0.0139]. This interaction reflects
that the reaction to committing an error varied as a function of both Manganese exposure and prenatal
maternal diet. Specifically, under the unsupplemented diet, only the controls exhibited a lower percent
correct for trials following an error (relative to after a correct response); the Mn animals showed no
decline in performance due to making an error on the prior trial (p < 0.0001 for control and 0.79 for Mn
animals comparing error vs. correct; Figure 5). In contrast, for animals born to dams given the choline
supplemented diet, both groups showed a reduction in percent correct for trials following an error
(relative to after a correct response; p = 0.074 for control and 0.17 for Mn), and the magnitude of this
disruptive effect was similar in the two groups.
491

492 In the control animals, MCS significantly increased %Correct responses after an error, but not following a
493 correct response (p's=0.05 and 0.50, respectively) (Figure 5). In Mn animals, MCS did not improve
494 performance following either a correct response (p = 0.34) or error (p = 0.73) compared to
495 unsupplemented animals.
496
497 In contrast to trials with a 0 sec pre-cue delay, trials with a 3 sec pre-cue delay did not have a significant
498 interaction of Mn x MCS x previous trial outcome [F(1, 273.8) = 1.72, p = 0.19] (not shown). Therefore,
499 the effect of MCS to improve performance of the control group animals and mitigate behavioral reactivity
500 on trials following an error is most pronounced on trials that are closest in time to the prior trials (i.e.,
501 trails with a 0 sec pre-cue delay).
502
MCS and developmental Mn exposure differently alter behavioral reactivity in response to an
error or omitted reward on the prior trail in a reward omission task
In the reward omission task, we introduced a novel trial condition in which the food pellet reward is
unexpectedly omitted on a subset of correct response trials, and assessed how this affects performance
on the subsequent trial. This was done because we wanted to explore the effect of another type of
behaviorally disruptive stimuli (i.e., reward omission) in contrast to the disruptive effect of an incorrect
response. Specifically, we assessed performance on the current trial depending on whether the
preceding trial response was either (1) correct with a food pellet delivery, (2) an error (incorrect, no food
pellet delivery + 5 sec time out), or (3) correct but with food pellet reward omitted (RO; occurred pseudo-
randomly on three out of each ten consecutive correct trials).
513
Under the unsupplemented diet, Mn eliminates the reactivity to an error that is seen in control animals,
but causes reactivity to the omission of an expected reward. MCS eliminates the deficit following an
omitted reward in Mn animals and improves performance in control animals following an error.
In the reward omission task, there was a Mn x MCS x previous trial outcome interaction for %Correct
[F(2, 410.6) = 4.51, p = 0.0116] (Figure 6). This interaction reflects that both prenatal diet and Mn
exposure affects reactivity to these two conditions. In the unsupplemented controls, committing an error
on the prior trial (versus a correct response) caused a reduction in %Correct responses on the
subsequent (current) trial (p < 0.0001), whereas the unexpected omission of the food reward on the prior
correct trial had no effect on the performance on the subsequent trial (p = 0.20). In contrast, a different
pattern of post–error performance was seen in the unsupplemented Mn animals. Notably, for these
animals, committing an error on the prior trial did not affect their performance on the subsequent
(current) trial (p = 0.152), compared to trials following a correct response (Figure 6). In contrast,
performance of the unsupplemented Mn animals was significantly decreased on trials following the
omission of the expected food reward, compared to a correct response (p = 0.0031). These findings
demonstrate that, for offspring of dams fed an unsupplemented diet, animals exposed to Mn early in life
respond differently than controls (no Mn) to both committing an error and not receiving an expected
reward.
531
A different pattern was seen for animals born to dams on a choline supplemented diet. In MCS control
animals, there was a trend for MCS to lessen the disruptive effect of committing an error on the prior trial;
i.e., improve performance on trials that followed an error (relative to the unsupplemented controls) (p =
0.071, Figure 6). However, within the MCS controls, %Correct responses on trials following an error was
still lower than on trials that followed a correct response (p < 0.0001). Furthermore, in the MCS controls,
the unexpected omission of a food reward on the prior trial also impaired performance compared to

following a correct response (p = 0.0013), an effect that was not seen in the unsupplemented animals. In
addition, for the Mn animals, MCS normalized performance on trials following a reward omission (Figure
6).
541
542 Finally, under the MCS diet, performance on trials following a correct response was significantly worse in
543 the Mn animals than controls (p = 0.0176), a deficit that was not present in the unsupplemented groups
544 due to the fact that the controls benefited from MCS for trials following a correct response but the Mn
545 animals did not.
546
547 **3.5 Sensorimotor function**
548
Early life Mn exposure causes a lasting deficit in sensorimotor function that is not altered by
MCS.
To determine whether MCS can mitigate the lasting sensorimotor deficits caused by developmental Mn
exposure, animals were assessed using the Montoya staircase test of skilled forelimb performance. As
expected, Mn exposure caused lasting deficits in sensorimotor function, as demonstrated by a significant
Mn x step level interaction for number of pellets taken [$F_{(5, 229.3)}$ = 10.66, p < 0.0001], number of
pellets eaten [$F_{(5, 232.6)}$ = 4.57, p = 0.0005], %Success [$F_{(5, 231.4)}$ = 3.26, p = 0.0073], and number of
pellets misplaced [$F_{(5, 356.7)}$ = 5.76, p < 0.0001]. Specifically, there was a significant Mn deficit on steps
4 and 5 of the staircase, where the unsupplemented Mn animals ate significantly fewer pellets and
misplaced more pellets compared to unsupplemented control counterparts (pellets eaten, p's = 0.0017
and 0.0002 for steps 4 and 5 respectively; pellets misplaced, p's = 0.0005 and 0.002) (Figure 7a1 and
7b1, respectively). However, MCS did not alter sensorimotor function in either the controls or the Mn
groups, based on findings that the main effect of MCS (P's >0.16), or any interactions involving MCS and
Mn were not significant for any of the outcomes (e.g., Mn x MCS, P's >0.32; Mn x MCS x step level, P's
>0.72); see figure panels 7A and 7B for the Mn x MCS x step plots for pellets eaten and misplaced).
564
565 **3.6 Choline and choline metabolite levels**
566
MCS measurably altered choline and choline metabolite levels in dams and offspring.
To confirm that oral MCS over gestation and lactation altered levels of choline and/or its metabolites in
dams and their offspring as expected, and whether oral Mn exposure altered choline/metabolite status in
the offspring of MCS-treated dams, we measured levels of choline and selected metabolites in dam and
offspring tissue (dam plasma and liver upon completion of MCS at PND 21, offspring liver and brain at
PND 24). As expected, MCS produces measurable changes in choline and some of its metabolites in
dam plasma and liver, and PND 24 pup liver and brain. See supplemental table 1 for choline and
metabolite measurements in dams, and supplemental table 2 for measurements in pups.
575
576 **3.7 Blood and brain Mn levels**
577
Oral Mn dosing increased blood and brain Mn levels in weanlings, and returned to control levels
in adulthood.
Mn exposure from postnatal days (PND) 1-21 increased levels of Mn in both blood and brain tissue in
PND 24 weanlings. Specifically, oral Mn significantly increases blood Mn levels approximately 8-fold,
from ~30 to 250 ng/mL in control vs. Mn exposed groups [$F_{(1, 45)}$ = 398.4, p < 0.0001] (Figure 8A).
There was also a significant interaction of Mn x MCS on blood Mn levels [$F_{(1, 42)}$ = 8.37, p = 0.0059];

however, Tukey's contrasts revealed no significant difference between blood Mn levels in
unsupplemented vs. MCS animals that received Mn.
586
Mn dosing also causes an approximately 3-fold increase in brain Mn levels in PND 24 offspring, from
approximately 3.8 to 11.8 µg/g in control vs. Mn-dosed animals, as demonstrated by a significant main
effect of Mn [$F(1, 36) = 129.7$, $p < 0.0001$] (Figure 8B). There is also a trending main effect of MCS on
brain Mn levels in offspring [$F(1, 36) = 3.96$, $p = 0.0543$], as well as a trending Mn x MCS interaction [$F(1,$
$36) = 3.13$, $p = 0.085$]. Tukey's pairwise comparisons revealed that, of animals that received Mn, MCS
animals had significantly higher levels of brain Mn than unsupplemented animals.
593
594 Finally, blood and brain Mn return to baseline levels in PND 200 behaviorally tested animals. However,
595 there is a statistically significant but biologically unremarkable effect of MCS [$F(1, 27) = 7.87$, $p = 0.0092$]
596 to reduce blood Mn levels (Figure 8C). In the brain tissue of PND 200 animals, there was a trending
597 effect of MCS to reduce Mn levels [$F(1, 20) = 3.19$, $p = 0.089$]; however, Tukey's pairwise comparisons
598 revealed no significant difference between any groups (Figure 8D).
599
600 **4. Discussion summary**
601
Studies in children and adolescents have associated early developmental Mn exposure with inattention,
impulsivity, hyperactivity, and sensorimotor dysfunction. There are currently no recognized
therapies/interventions, other than exposure prevention, to mitigate the neurotoxic effects of elevated
developmental Mn exposure. In this study, we used a rodent model of developmental Mn exposure to
determine whether MCS protects against the lasting Mn-induced deficits in attention, behavioral
reactivity, learning, impulsivity, and sensorimotor function. Our results demonstrate that developmental
Mn exposure causes lasting deficits in attention, behavioral reactivity, learning, and sensorimotor
function, and that the MCS intervention was partially effective in protecting against Mn-induced deficits,
particularly in the domains of attentional function and behavioral reactivity. Collectively, these findings
demonstrate that MCS may hold promise as nutritional intervention to protect against the neurotoxic
impacts of elevated Mn exposure, as has been seen previously for other early insults. Given that recent
evidence suggests that 90% of pregnant women consume less choline than the current recommended
Adequate Intake (AI) levels, these data provide additional support for efforts to add choline to standard
prenatal vitamin regimens. These findings are discussed in more detail below.
616
4.1 Developmental Mn exposure causes lasting abnormalities in behavioral reactivity, and replicates the
effects seen in our studies in which Mn causes deficits in attention, learning, and sensorimotor function.
In this study, Mn exposure caused deficits in attentional function (Figs 3 and 4), learning to withhold
impulsive responses (Fig 2), and sensorimotor function (Fig 7) in ways that are largely consistent with
our prior studies[14–16,59]. In addition, we established that elevated exposure to Mn during development
leads to lasting abnormalities deficits in behavioral reactivity (Figs 5 and 6), including how animals
respond to an incorrect response, as well as the omission of an expected reward.
624
625 To understand the MCS effects on behavioral reactivity in control and Mn exposed rats, it is important to
626 first describe the effects of Mn in the absence of MCS. While control animals exhibit fewer correct
627 responses on trials following an error than on trials following a correct response, as previously observed
628 [34,53,60–63], Mn blunts this 'normal' reactivity to an error; Mn animals perform the same on trials that follow
629 an error vs. a correct response in both the second focused attention and reward omission tasks (Fig 5,

6). However, in contrast to the lack of effect on performance for trials that follow an error, Mn exposed animals exhibit more errors on trials that follow the unexpected omission of a reward than on trials following a correct response, whereas control animals exhibit no performance deficit under this condition (relative to trials following a correct response) (Fig 6).

4.2 MCS provides some benefits to the Mn-exposed animals by lessening the difference between control and Mn animals on behavioral reactivity and attention.

MCS in some instances normalized the behavioral reactivity of the Mn-exposed animals. Specifically, the Mn animals in the MCS condition respond like controls to the unexpected omission of reward; i.e. they do not show a decrement in performance relative to trials after a correct response. Note that control animals also derive a benefit from MCS, which improves performance in trials following an error compared to the unsupplemented diet (Fig 5, 6); there is still a drop in correct responses after an error (compared to after a correct response), but it is less pronounced. These findings support that developmental Mn exposure produces lasting disruptions in arousal and behavioral reactivity – not in ways that simply blunt or augment behavioral reactivity, but rather modify interpretation/neural processing of committing an error or not receiving an expected reward.

In addition to normalizing some aspects of behavioral reactivity in Mn animals, MCS provides benefits in the domain of attentional function by lessening differences between Mn and control animals on the key measures of attention (%Correct and %Accurate). This is seen in the focused attention 2 task, particularly at the longer and more challenging pre-cue delays (Figure 3C and D), and in the selective attention task, where MCS reduces the difference between Mn and control animals on distraction trials (Figure 4).

4.3 Mn exposure mitigates some of the benefits of MCS.

It is noteworthy that developmental Mn exposure diminishes some of the cognitive benefits of MCS seen in control animals, particularly in learning rate in the first focused attention task. MCS facilitates learning the rules of this new task to a greater extent for the controls than the Mn rats. The improvement in %Accurate responses between block 1 and 2, under MCS, was greater in controls than Mn animals, and thus a Mn deficit emerges under blocks 2 and 3 (significant difference between control and Mn). This is interpreted as a learning benefit provided by MCS to control animals, but this benefit is not seen in the Mn animals.

4.4 MCS benefits control animal cognitive function.

Maternal choline supplementation in the absence of Mn exposure provided a number of cognitive benefits. For example, in control animals, supplementing the maternal diet with additional choline during gestation and lactation lessens the disruptive effect of committing an error (Fig 5, 6; discussed above in section 4.2) and improves attentional function (Fig 3C). The benefit to attention is statistically significant at the 0 sec delay in the FA2 task (Fig 3C), which may also be due in part to the benefit to behavioral reactivity described above. In addition, there is a pattern effects that did not quite reach statistical significance, but further provide evidence of a benefits of MCS on attention. This included the MCS benefit to control attentional function in FA1(3A/B) task, as well as increasing correct responses in the RO task on trials following a correct response (Fig 6). These benefits show that in the absence of environmental exposure, MCS provides cognitive benefits to offspring, [i]which recapitulate benefits seen in prior studies in humans and animal models[37–40,42,45,46,48].

676
4.5 Conclusion
In conclusion, these results demonstrate that MCS is beneficial to both the control and Mn exposed
animals. MCS normalizes, to some degree, the Mn deficits in the areas of attention and behavioral
reactivity. These findings also underscore the lasting neurobehavioral health risks to children posed by
developmental Mn exposure. Finally, in light of the low choline intakes of most pregnant women, these
results provide additional support for adding choline to standard prenatal vitamin regimens.

Figures

Figure 1. Study timeline with task parameters, and study design. (A) Study timeline, including MCS and Mn dosing, as well as tasks with parameters. X-axis is days of age (PND). (B) 2x2 study design, ending with 16 pups per treatment group that were used for behavioral testing.

690
691
Figure 2. Mn animals are impaired in learning to withhold premature responses, and MCS lessens
some differences between control and Mn treatment groups. Premature responses (%) in the first
focused attention task as a function of Mn treatment, maternal diet, and session block (A), and a function
of Mn treatment and session block (a1). Premature responses (%) in the focused attention 2 task as a
function of Mn treatment, maternal diet, and session block (B), and a function of Mn treatment and
session block (b1). Premature responses (%) in the focused attention 2 task as a function of Mn
treatment, maternal diet, and pre-cue delay (sec) (C), and a function of MCS treatment and pre-cue
delay (d1). Premature responses (%) in the SA task as a function of Mn treatment, maternal diet, and
session block (D). The order of testing was the focused attention 1 task followed by the focused attention
2 task and then the selective attention task. * indicates p ≤ 0.05 versus controls. Data are lsmeans ±
SEM (n=14-16/group).

703
704
Figure 3. Mn animals exhibit an impairment in focused attention, and MCS differentially affects
control vs. Mn animal performance. Focused attention 1 task %Correct (A, a1) and %Accurate (B)
responses as a function of Mn treatment, maternal diet, and session block. Focused attention 2 task
%Correct (C) and %Accurate (D) responses as a function of Mn treatment, maternal diet, and pre-cue
delay in sec. * and & indicate $p \leq 0.05$ and $p \leq 0.10$ versus controls, respectively; # indicates $p \leq 0.05$
versus unsupplemented animals. Data are lsmeans ± SEM (n=14-16/group).

711
712
Figure 4: Mn causes a deficit in attention, but not selective attention specifically, and MCS
reduces the difference between Mn and control animals under odor conditions. SA task %Correct
responses as a function of Mn treatment, maternal diet, and distraction condition (no distractor, odor
distractor 1, 2, or 3 sec prior to the cue illuminating) (A), Mn treatment and distraction condition (a1), or
Mn treatment, maternal diet, and odor presence (no distractor, odor distractor present) (B). %Accurate
responses as a function of Mn treatment, maternal diet, and distraction condition (C), and Mn treatment
and distraction condition (c1). * indicates p ≤ 0.05 versus controls. Data are lsmeans ± SEM (n=14-
16/group).

721
722
Figure 5: Under the unsupplemented diet, Mn blunts error reactivity. MCS improves performance in control animals following an error. FA 2 task %Correct responses at trial conditions with a 0 sec PCD, as a function of Mn treatment, maternal diet, and previous trial outcome. Corr = correct, Error = incorrect, premature, or omission error. % and + indicate $p \leq 0.05$ and $p \leq 0.10$ versus following a correct response, respectively; # indicates $p \leq 0.05$ versus unsupplemented animals. Data are lsmeans ± SEM (n=14-16/group).

Figure 6: Mn blunts error reactivity, but causes reactivity to an omitted reward, which is normalized by MCS. RO task (no reward on one-third of the correct response trials) %Correct responses as a function of Mn treatment, maternal diet, and previous trial outcome. Corr = correct, Error = incorrect or omission error, RO = reward omitted. % indicates $p \leq 0.05$ versus following a correct response; * indicates $p \leq 0.05$ versus controls; ^ indicates $p \leq 0.10$ versus unsupplemented animals. Data are lsmeans ± SEM (n=14-16/group).

Figure 7: Mn exposure causes a lasting deficit in fine motor function that is not rescued by MCS.
Number of pellets eaten by control and Mn-exposed groups that received unsupplemented or
supplemented maternal choline, as a function of step level (A), number of pellets eaten as a function of
Mn and step level (a1). Number of pellets misplaced by control and Mn-exposed groups that received
unsupplemented or supplemented maternal choline, as a function of step level (B), number of pellets
misplaced as a function of Mn and step level (b1). * indicates p ≤ 0.05 versus controls. Data are lsmeans
± SEM (n=14-16/group).

Figure 8: **Oral Mn dosing increased blood and brain Mn concentrations in weanling littermates of the behaviorally tested animals (PND 24), and returned to baseline in the behaviorally tested animals at sacrifice (PND 200)**. PND 24 animal blood (n = 11-14) (A) and brain (n = 9-11) (B) measurements show that Mn was increased in Mn-exposed animals. PND 200 animal blood (n = 7-8) (C) and brain (n = 6) (D) measurements show that Mn levels return to baseline. Graphs show arithmetic means with standard error. PND 24 blood and brain statistics were performed on log-transformed data; PND 200 statistics were performed on non-transformed data. Within an age group and tissue, treatment groups with different capital letter superscripts are statistically different from one another (p < 0.05), based on Tukey's *post hoc* test.

Funding Statement
This study was supported by NIEHS grant #R01ES028369.

Conflicts of interest (COI)
The authors declare no actual competing or potential financial interests.

Acknowledgments
The authors would like to thank B. Bilow, C. Agbayani, D. Woodfin, E. Hsu, G. Chiang, J. Do, J. Altshul, K. Sharifi, L. Shahnazarian, N. Schlipp, N. Chenna, R. Poustinchi, R. Basto, S. Hanson, S. Nikolaeva, S. Twardy, S. Bayliss Grijalva, T. Waite, V. Howland, V. Deng, and W. Valdez Rodriguez for their invaluable assistance in behavioral testing. We also thank T. Jursa, M. Caudill, and O. Malysheva for analytical assistance, L. Johnson for the assistance with statistical analyses, and N. Santiago for assistance in the study.

References

1. Faraone, S. V. *et al.* The World Federation of ADHD International Consensus Statement: 208 Evidence-based conclusions about the disorder. *Neuroscience and Biobehavioral Reviews* **128**, 789–818 (2021).
2. Schullehner, J. *et al.* Exposure to manganese in drinking water during childhood and association with attention-deficit hyperactivity disorder: A nationwide cohort study. *Environ. Health Perspect.* **128**, 1–10 (2020).
3. Lucchini, R. G. *et al.* Tremor, olfactory and motor changes in Italian adolescents exposed to historical ferro-manganese emission. *Neurotoxicology* **33**, 687–96 (2012).
4. Liu, W. *et al.* Biomarkers of environmental manganese exposure and associations with childhood neurodevelopment: A systematic review and meta-analysis. *Environ. Heal. A Glob. Access Sci. Source* **19**, 1–22 (2020).
5. Broberg, K. *et al.* Manganese transporter genetics and sex modify the association between environmental manganese exposure and neurobehavioral outcomes in children. *Environ. Int.* **130**, 104908 (2019).
6. Bauer, J. A. *et al.* Manganese in teeth and neurobehavior: Sex-specific windows of susceptibility. *Environ. Int.* **108**, 299–308 (2017).
7. Bauer, J. A. *et al.* Critical windows of susceptibility in the association between manganese and neurocognition in Italian adolescents living near ferro-manganese industry. *Neurotoxicology* **87**, 51 (2021).
8. Mora, A. M. *et al.* Prenatal and postnatal manganese teeth levels and neurodevelopment at 7, 9, and 10.5 years in the CHAMACOS cohort. *Environ. Int.* **84**, 39 (2015).
9. Haynes, E. N. *et al.* Manganese Exposure and Neurocognitive Outcomes in Rural School-Age Children: The Communities Actively Researching Exposure Study (Ohio, USA). *Environ. Health Perspect.* **123**, 1066 (2015).
10. Oulhote, Y. *et al.* Neurobehavioral function in school-age children exposed to manganese in drinking water. *Environ. Health Perspect.* **122**, 1343–50 (2014).
11. Hong, S. B. *et al.* Blood manganese levels in relation to comorbid behavioral and emotional problems in children with attention-deficit/hyperactivity disorder. *Psychiatry Res.* **220**, 418–425 (2014).
12. Shin, D. W. *et al.* Association of Hair Manganese Level with Symptoms in Attention-Deficit/Hyperactivity Disorder. *Psychiatry Investig.* **12**, 66 (2015).
13. Beaudin, S. A., Nisam, S. & Smith, D. R. Early life versus lifelong oral manganese exposure differently impairs skilled forelimb performance in adult rats. *Neurotoxicol. Teratol.* **38**, 36–45 (2013).
14. Beaudin, S. A. *et al.* Oral methylphenidate alleviates the fine motor dysfunction caused by chronic postnatal manganese exposure in adult rats. *Toxicol. Sci.* **144**, 318–327 (2015).

15. Beaudin, S. A., Strupp, B. J., Strawderman, M. & Smith, D. R. Early postnatal manganese
exposure causes lasting impairment of selective and focused attention and arousal regulation in
adult rats. *Environ. Health Perspect.* **125**, 230–237 (2017).
16. Beaudin, S. A. *et al.* Methylphenidate alleviates manganese-induced impulsivity but not
distractibility. *Neurotoxicol. Teratol.* **61**, 17–28 (2017).
17. Kern, C. H., Stanwood, G. D. & Smith, D. R. Preweaning manganese exposure causes
hyperactivity, disinhibition, and spatial learning and memory deficits associated with altered
dopamine receptor and transporter levels. *Synapse* **64**, 363–78 (2010).
18. Kern, C. H. & Smith, D. R. Preweaning Mn exposure leads to prolonged astrocyte activation and
lasting effects on the dopaminergic system in adult male rats. *Synapse* **65**, 532–544 (2011).
19. Conley, T. E. *et al.* Early postnatal manganese exposure causes arousal dysregulation and lasting
hypofunctioning of the prefrontal cortex catecholaminergic systems. *J. Neurochem.* **153**, 631–649
(2019).
20. Lasley, S. M. *et al.* Early Postnatal Manganese Exposure Reduces Rat Cortical and Striatal
Biogenic Amine Activity in Adulthood. *Toxicol. Sci.* **173**, 144 (2019).
21. Shumay, E., Fowler, J. S. & Volkow, N. D. Genomic Features of the Human Dopamine
Transporter Gene and Its Potential Epigenetic States: Implications for Phenotypic Diversity. *PLoS*
One **5**, e11067 (2010).
22. Day, J. J. *et al.* DNA methylation regulates associative reward learning. *Nat. Neurosci.* **16**, 1445
(2013).
23. Schreiner, F. *et al.* Association of COMT genotypes with S-COMT promoter methylation in growth-
discordant monozygotic twins and healthy adults. *BMC Med. Genet.* **12**, 1–12 (2011).
24. Green, A. L. *et al.* Epigenetic Regulation of Dopamine Transporter mRNA Expression in Human
Neuroblastoma Cells. *Neurochem. Res.* **40**, 1372–8 (2015).
25. Muench, C., Wiers, C. E., Cortes, C. R., Momenan, R. & Lohoff, F. W. Dopamine Transporter
Gene Methylation is Associated with Nucleus Accumbens Activation during Reward Processing in
Healthy but not Alcohol-Dependent Individuals. *Alcohol. Clin. Exp. Res.* **42**, 21 (2018).
26. Dolinoy, D. C., Huang, D. & Jirtle, R. L. Maternal nutrient supplementation counteracts bisphenol
A-induced DNA hypomethylation in early development. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 13056–
61 (2007).
27. Wright, K. N. *et al.* Methyl supplementation attenuates cocaine-seeking behaviors and cocaine-
induced c-Fos activation in a DNA methylation-dependent manner. *J. Neurosci.* **35**, 8948–58
(2015).
28. Thomas, J. D., Idrus, N. M., Monk, B. R. & Dominguez, H. D. Prenatal choline supplementation
mitigates behavioral alterations associated with prenatal alcohol exposure in rats. *Birth Defects*
Res. A. Clin. Mol. Teratol. **88**, 827–37 (2010).
29. Thomas, J. D., Biane, J. S., O'Bryan, K. A., O'Neill, T. M. & Dominguez, H. D. Choline
supplementation following third-trimester-equivalent alcohol exposure attenuates behavioral
alterations in rats. *Behav. Neurosci.* **121**, 120–130 (2007).
30. Thomas, J. D., La Fiette, M. H., Quinn, V. R. . & Riley, E. P. Neonatal choline supplementation
ameliorates the effects of prenatal alcohol exposure on a discrimination learning task in rats.
Neurotoxicol. Teratol. **22**, 703–711 (2000).
31. Schulz, K. M. *et al.* Dietary choline supplementation to dams during pregnancy and lactation
mitigates the effects of in utero stress exposure on adult anxiety-related behaviors. *Behav. Brain*
Res. **268**, 104–10 (2014).
32. Ash, J. A. *et al.* Maternal choline supplementation improves spatial mapping and increases basal
forebrain cholinergic neuron number and size in aged Ts65Dn mice. *Neurobiol. Dis.* **70**, 32–42
(2014).
33. Velazquez, R. *et al.* Maternal choline supplementation improves spatial learning and adult
hippocampal neurogenesis in the Ts65Dn mouse model of Down syndrome. *Neurobiol. Dis.* **58**,
92–101 (2013).
34. Moon, J. *et al.* Perinatal Choline Supplementation Improves Cognitive Functionning and Emotional
Regulation in the Ts65Dn Mouse Model of Down Syndrome. *Behav. …* **124**, 346–361 (2010).

35. Bekdash, R. A., Zhang, C. & Sarkar, D. K. Gestational choline supplementation normalized fetal
alcohol-induced alterations in histone modifications, DNA methylation, and proopiomelanocortin
(POMC) gene expression in β-endorphin-producing POMC neurons of the hypothalamus. *Alcohol.*
Clin. Exp. Res. **37**, 1133–1142 (2013).
36. Otero, N. K. H., Thomas, J. D., Saski, C. A., Xia, X. & Kelly, S. J. Choline Supplementation and
DNA Methylation in the Hippocampus and Prefrontal Cortex of Rats Exposed to Alcohol During
Development. *Alcohol. Clin. Exp. Res.* **36**, 1701–1709 (2012).
37. Bahnfleth, C. L., Strupp, B. J., Caudill, M. A. & Canfield, R. L. Prenatal choline supplementation
improves child sustained attention: A 7-year follow-up of a randomized controlled feeding trial.
FASEB J. **36**, (2022).
38. Caudill, M. A., Strupp, B. J., Muscalu, L., Nevins, J. E. H. & Canfield, R. L. Maternal choline
supplementation during the third trimester of pregnancy improves infant information processing
speed: a randomized, double-blind, controlled feeding study. *FASEB J.* **32**, 2172–2180 (2017).
39. Boeke, C. E. *et al.* Choline intake during pregnancy and child cognition at age 7 years. *Am. J.*
Epidemiol. **177**, 1338–1347 (2013).
40. Meck, W. H. & Williams, C. L. Simultaneous temporal processing is sensitive to prenatal choline
availability in mature and aged rats. *Neuroreport* **8**, 3045–51 (1997).
41. Meck, W. H. & Williams, C. L. Characterization of the facilitative effects of perinatal choline
supplementation on timing and temporal memory. *Neuroreport* **8**, 2831–5 (1997).
42. Meck, W. H. & Williams, C. L. Metabolic imprinting of choline by its availability during gestation:
implications for memory and attentional processing across the lifespan. *Neurosci. Biobehav. Rev.*
27, 385–399 (2003).
43. Meck, W. H., Williams, C. L., Cermak, J. M. & Blusztajn, J. K. Developmental periods of choline
sensitivity provide an ontogenetic mechanism for regulating memory capacity and age-related
dementia. *Front. Integr. Neurosci.* **1**, (2008).
44. Mellott, T. J., Williams, C. L., Meck, W. H. & Blusztajn, J. K. Prenatal choline supplementation
advances hippocampal development and enhances MAPK and CREB activation. *FASEB J.* **18**,
545–547 (2004).
45. Buhusi, C. V., Lamoureux, J. A. & Meck, W. H. Prenatal choline supplementation increases
sensitivity to contextual processing of temporal information. *Brain Res.* **1237**, 204 (2008).
46. Cheng, R. K., MacDonald, C. J., Williams, C. L. & Meck, W. H. Prenatal choline supplementation
alters the timing, emotion, and memory performance (TEMP) of adult male and female rats as
indexed by differential reinforcement of low-rate schedule behavior. *Learn. Mem.* **15**, 153–162
(2008).
47. Wong-Goodrich, S. J. E. *et al.* Spatial memory and hippocampal plasticity are differentially
sensitive to the availability of choline in adulthood as a function of choline supply in utero. *Brain*
Res. **1237**, 153–66 (2008).
48. Mohler, E. G., Meck, W. H. & Williams, C. L. Sustained Attention in Adult Mice is Modulated by
Prenatal Choline Availability. *Int. J. Comp. Psychol.* **14**, 136–150 (2001).
49. Wallace, T. C. & Fulgoni, V. L. Assessment of Total Choline Intakes in the United States. *J. Am.*
Coll. Nutr. (2016). doi:10.1080/07315724.2015.1080127
50. Wallace, T. C. & Fulgoni, V. L. Usual Choline Intakes Are Associated with Egg and Protein Food
Consumption in the United States. *Nutrients* **9**, (2017).
51. Wallace, T. C. *et al.* The underconsumed and underappreciated essential nutrient. *Nutr. Today* **53**,
240–253 (2018).
52. Institute of Medicine (US) Food and Nutrition Board. *Dietary Reference Intakes for Thiamin,*
Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline.
(National Academies Press , 1998).
53. Stangle, D. E. *et al.* Succimer Chelation Improves Learning, Attention, and Arousal Regulation in
Lead-Exposed Rats but Produces Lasting Cognitive Impairment in the Absence of Lead Exposure.
Environ. Health Perspect. **115**, 201 (2007).
54. Montoya, C. P., Campbell-Hope, L. J., Pemberton, K. D. & Dunnett, S. B. The "staircase test": a
measure of independent forelimb reaching and grasping abilities in rats. *J. Neurosci. Methods* **36**,

219–228 (1991).

55.   Kloth, V., Klein, A., Loettrich, D. & Nikkhah, G. Colour-coded pellets increase the sensitivity of the
staircase test to differentiate skilled forelimb performances of control and 6-hydroxydopamine
lesioned rats. *Brain Res. Bull.* **70**, 68–80 (2006).

56.   Holm, P. I., Ueland, P. M., Kvalheim, G. & Lien, E. A. Determination of Choline, Betaine, and
Dimethylglycine in Plasma by a High-Throughput Method Based on Normal-Phase
Chromatography–Tandem Mass Spectrometry. *Clin. Chem.* **49**, 286–294 (2003).

57.   Koc, H., Mar, M. H., Ranasinghe, A., Swenberg, J. A. & Zeisel, S. H. Quantitation of choline and
its metabolites in tissues and foods by liquid chromatography/electrospray ionization-isotope
dilution mass spectrometry. *Anal. Chem.* **74**, 4734–4740 (2002).

58.   Ruiz, J., Antequera, T., Andres, A. I., Petron, M. J. & Muriel, E. Improvement of a solid phase
extraction method for analysis of lipid fractions in muscle foods. *Anal. Chim. Acta* **1**–**2**, 201–205
(2004).

59.   Beaudin, S. A., Nisam, S. & Smith, D. R. Early life versus lifelong oral manganese exposure
differently impairs skilled forelimb performance in adult rats. *Neurotoxicol. Teratol.* **38**, 36–45
(2013).

60.   Gendle, M. H. *et al.* Impaired sustained attention and altered reactivity to errors in an animal
model of prenatal cocaine exposure. *Dev. Brain Res.* **147**, 85–96 (2003).

61.   Gendle, M. H. *et al.* Enduring Effects of Prenatal Cocaine Exposure on Selective Attention and
Reactivity to Errors: Evidence from an Animal Model. *Behav. Neurosci.* **118**, 290–297 (2004).

62.   Morgan, R. E. *et al.* Early lead exposure produces lasting changes in sustained attention,
response initiation, and reactivity to errors. *Neurotoxicol. Teratol.* **23**, 519–531 (2001).

63.   Rushworth, M. F. S., Hadland, K. A., Gaffan, D. & Passingham, R. E. The effect of cingulate
cortex lesions on task switching and working memory. *J. Cogn. Neurosci.* **15**, 338–353 (2003).