The rules of sex and strain: deciphering the chemical language of inbred and wild mouse conspecific scents

Maximilian Nagel[1,6], Marco Niestroj[1], Rohini Bansal[2], David Fleck[1], Angelika Lampert[3,4], Romana Stopkova[5], Pavel Stopka[5,*], Yoram Ben-Shaul[2,*], Marc Spehr[1,4,*,‡]

[1] Department of Chemosensation, Institute for Biology II, RWTH Aachen University, 52074 Aachen, Germany

[2] Department of Medical Neurobiology, Institute for Medical Research Israel Canada, Faculty of Medicine, The Hebrew University of Jerusalem, Jerusalem, Israel

[3] Institute of Neurophysiology, Uniklinik RWTH Aachen University, 52074 Aachen, Germany

[4] Research Training Group 2416 *MultiSenses – MultiScales*, RWTH Aachen University, 52074 Aachen, Germany

[5] BIOCEV group, Department of Zoology, Faculty of Science, Charles University, Prague, Czech Republic

[6] present address: Sensory Cells and Circuits Section, National Center for Complementary and Integrative Health, Bethesda, MD, 20892, USA

[*] correspondence should be addressed to: Pavel Stopka (pavel.stopka@natur.cuni.cz), Yoram Ben-Shaul (yoramb@ekmd.huji.ac.il), Marc Spehr (m.spehr@sensorik.rwth-aachen.de)

[‡] lead contact

Summary

In most mammals, conspecific chemosensory communication relies on semiochemical release
within complex bodily secretions and subsequent stimulus detection by the vomeronasal organ
(VNO). Urine, a rich source of ethologically relevant chemosignals, conveys detailed information
about sex, social hierarchy, health and reproductive state, which becomes accessible to a
conspecific via vomeronasal sampling. So far, however, numerous aspects of social
chemosignaling along the vomeronasal pathway remain unclear. Moreover, since virtually all
research on vomeronasal physiology is based on secretions derived from inbred laboratory mice,
it remains uncertain whether such stimuli provide a true representation of potentially more relevant
cues found in the wild. Here, we combine a robust low-noise VNO activity assay with comparative
molecular profiling of sex- and strain-specific mouse urine samples from two inbred laboratory
strains as well as from wild mice. With comprehensive molecular portraits of these secretions,
VNO activity analysis now enables us to (*i*) assess whether and, if so, how much sex- / strain-
selective "raw" chemical information in urine is accessible via vomeronasal sampling; (*ii*) identify
which chemicals exhibit sufficient discriminatory power to signal an animal's sex, strain, or both;
(*iii*) determine the extent to which wild mouse secretions are unique; and (*iv*) analyze whether
vomeronasal response profiles differ between strains. We report both sex- and, in particular,

strain-selective VNO representations of chemical information. Within the urinary 'secretome', both
volatile compounds and proteins exhibit sufficient discriminative power to provide sex- and strain-
specific molecular fingerprints. While total protein amount is substantially enriched in male urine,
females secrete a larger variety at overall comparatively low concentrations. Surprisingly, the
molecular spectrum of wild mouse urine does not dramatically exceed that of inbred strains.
Finally, vomeronasal response profiles differ between C57BL/6 and BALB/c animals, with
particularly disparate representations of female semiochemicals.

Introduction

In rodents and most other mammals, the accessory olfactory system controls conspecific chemical
communication during social interactions [1–4]. Ethologically meaningful chemostimuli that trigger
stereotypic social and sexual behaviors are predominantly detected by the vomeronasal organ
(VNO), a tubular sensory structure at the anterior base of the nasal septum. In a crescent-shaped
medial neuroepithelium, the VNO harbors approximately 100,000 to 200,000 vomeronasal
sensory neurons (VSNs [5]), each extending a single unbranched apical dendrite that terminates in
a paddle-shaped swelling [4]. At the dendritic tips, microvilli are immersed in mucus that fills a central
luminal canal, which extends via the vomeronasal duct into the nasal cavity. During social
investigatory behavior, which in mice primarily involves periods of intense licking and sniffing of
both facial and anogenital regions [6], semiochemicals are sucked into the VNO lumen. Upon
binding to cognate vomeronasal receptors (VRs), the chemical signal is transduced into electrical
VSN activity and, ultimately, neuronal discharge.

Behaviourally relevant natural chemostimuli are, typically, complex blends of compounds [7] in
various bodily secretions [8]. By far, the most widely studied secretion in animal chemosensory
research is urine (see [4] and references therein), which is a rich source of semiochemicals that
serves a well-established function in social communication. While we still lack a comprehensive
molecular description of this broadband vomeronasal stimulus, previous work has identified
several putative semiochemicals in mouse urine, which cover many structural groups and feature
dimensions [7,9–14]. So far, virtually all research on vomeronasal physiology is based on urinary
stimuli derived from inbred laboratory mice (but see [15]). This facilitates standardization across
studies, but it remains unclear whether secretions collected from inbred mice accurately represent
the potentially more ethologically relevant stimuli found in the wild. In fact, while wild *Mus musculus*
populations exhibit several-fold higher levels of genetic variation than those in human populations,
commonly studied inbred strains of laboratory mice derive from a limited set of founders.
Therefore, such strains contain only a small subset of the genetic variation that is present in nature

[16]. Classical inbred strains of house mice are genetic mosaics of the three main wild subspecies,

M. m. domesticus, *musculus*, and *castaneus*, which started to diverge ~350–500 thousand years

ago. However, the genomes of prominent inbred strains, such as C57BL/6 or BALB/c, are

predominantly derived from *M. m. domesticus* [17]. The mitochondrial genomes of both laboratory

strains are identical, implying a common descent along the maternal line. Given their overlapping

geographical distribution, wild house mice subspecies have also undergone secondary contact

and hybridization [18], which has diversified their genetic landscape.

Laboratory inbreeding, which has been ongoing for more than 100 years for C57BL/6 and BALB/c

strains [16], may have affected the chemical composition of bodily secretions and, consequently,

their information content. Indeed, both within and between strains, laboratory mice lack variation

in major urinary protein (MUP) patterns typically found among wild mice [19]. This inbred

homogeneity could have important implications for research investigating, for example, social

recognition or mate choice: if there are marked qualitative and/or quantitative differences in

chemical composition of secretions and, accordingly, in the neuronal representations of wild as

compared to inbred derived stimuli, one might question the conclusions based on a large body of

work using inbred secretions [15].

On the recipients' end, the level of chemosensory information available from con-/heterospecific

secretions is determined by individual VR expression profiles. Across strains, VR repertoires vary

as a function of both genetics and experience [20–23]. Therefore, it is unclear whether receptor arrays

in laboratory strains, despite hundreds of generations of inbreeding and domestication

("microevolution") in a laboratory environment, have retained selectivity for more ethologically

relevant wild-derived stimuli. Moreover, it remains uncertain whether VR tuning profiles enable

VSNs to capture key ethological features from molecular concentration differences between sex-

and strain-specific secretions [24,25]. Together, four key biological questions arise from these lines

of reasoning: (*i*) Do VSN response profiles reflect the global molecular content of urine (i.e., are

these neurons sensitive to many/most compounds) or, by contrast, is the VNO a highly selective

molecular detector (responding to just a few select molecules)? (*ii*) Are VSN response profiles

strain-specific? (*iii*) Which semiochemicals provide information about sex and/or strain? (*iv*) Is

there something unique about wild mouse secretions and/or the VSN response profiles they

trigger?

Here, to address these unresolved issues, we combine a robust VSN activity assay with

comparative molecular profiling of sex- and strain-specific mouse urine from two inbred laboratory

strains and wild mice. Our study provides comprehensive molecular portraits of these secretions.

We report that large fractions of generic urine compounds are shared among both male and

female mice of all genetic backgrounds. We further show that the urinary 'secretome' in wild mice

does not differ dramatically from that found in laboratory strains. Surprisingly, while male urine
contains much higher protein concentrations, females secrete a larger variety of proteins
(including MUPs). For proteins common to all strains, concentrations are relatively low in C57BL/6,
moderate in BALB/c, and high in wild animals. However, despite this concentration bias, there is
no overrepresentation of wild-selective proteins. Notably, both the volatile organic compound
(VOC) and protein profile of urine, each provides sufficient information to decode a given sex/strain
combination (with protein content exhibiting stronger discriminative power). Moreover, we identify
a rich lipocalin repertoire in urine, which alone could allow chemosensory discrimination of
sex/strain combinations.

A key strength of this study is the use of the exact same stimuli as previously employed to
investigate sensory representations in the accessory olfactory bulb (AOB) [15], the first central
processing stage along the accessory olfactory system pathway. Our previous work demonstrated
that AOB representations of ethologically relevant urine stimuli are similar for male mice from two
different inbred strains (C57BL/6 and BALB/c), despite potential differences in VR repertoires. In
addition, we found that wild mouse stimuli elicit responses that, although not identical, are
nevertheless qualitatively similar to those from commonly used inbred strains [15]. VSN activity
analysis now enables us to ask whether these features also manifest in the VNO and, thus, to
assess (*i*) whether the information inherent in a sex- and strain-specific urinary secretome is
accessible via vomeronasal sampling; and (*ii*) if, on the population level, VSN sensory
representations differ between strains. Comparing responses from male C57BL/6 and BALB/c
mice (as done previously on AOB level [15]), our data demonstrate that recipient strain identity is
reflected by their VSN activity patterns. Moreover, when exposed to stimuli from different strains,
large VSN fractions (often >50 %) respond to only one stimulus, suggesting a substantial degree
of selective sampling. It is striking that such selective VSN responses to wild stimuli are rather
rare. Population response patterns among non-selective VSNs are largely unaffected by
concentration differences among common urine components, indicating either that concentration
carries relatively little information, or that common but relevant semiochemicals are secreted at
similar, maybe saturating concentrations. Finally, at VSN level, representation of sensory features
differs between C57BL/6 and BALB/c animals, with particularly disparate representation of female
semiochemicals. Together, our study reveals selective and strain-dependent representations of
urine chemical content, a surprising scarceness of selective responses to wild stimuli, as well as
remarkably rich and sex-/strain-specific molecular profiles that likely preserve most of the
biologically relevant information.

Results

In mice, the VNO is the predominant sensory structure mediating conspecific chemical
communication. In this study, by comparing samples derived from two inbred strains (BALB/c and
C57BL/6) as well as from wild mice, we pursue four main questions: (***i***) Which chemical
components in mouse urine – a major source of semiochemicals – distinguish sex and/or strain?
(***ii***) To what degree has inbreeding affected the chemical composition of urine (i.e., how unique
are wild mouse secretions)? (***iii***) How much of the sex- and strain-specific chemical information in
urine is accessible to a conspecific via vomeronasal sampling (i.e., how selective are VSN
response profiles)? (***iv***) Upon exposure to the same stimuli, do VSN response patterns differ
between strains?

A "low noise" assay to capture VSN population activity

To record VSN signal fingerprints in response to naturalistic stimuli, it is essential to establish a
robust population activity assay that reliably captures the raw information content inherent in bodily
secretions. To this end, we analyzed single-cell Ca^{2+} transients among large VSN populations in
acute coronal VNO sections (Figure 1a,b). Throughout this study, we routinely compared pairs of
pooled stimuli following a standard experimental paradigm (Figure 1c), in which sample pairs
differed either in the donors' sex (male *versus* female) or in their genetic background (BALB/c,
C57BL/6, or wild). We repeated brief alternating stimulus presentations twice at inter-stimulus
intervals that ensured recovery from VSN adaptation [35]. Experiments concluded with a brief
exposure to elevated extracellular K^+ (**S₃**) to depolarize neurons and test for integrity of each
neuron's spike generation machinery. VSNs were categorized as either specialists (selective
response to one stimulus) or generalists (responsive to both stimuli). Overall, we recorded urine-
dependent Ca^{2+} signals from a total of 16,715 VSNs, of which 61.4% displayed generalist profiles,
whereas 38.6% were categorized as specialists (Figure 1c,d). As a measure of intrinsic signal
variability, we calculated reliability indices that quantify the similarity (or lack thereof) of two
successive responses to the same stimulus (with small values reflecting high reliability; see
Materials and Methods). We analyzed both Ca^{2+} signal amplitudes (Figure 1d) and integrals
(Figure S1a) as measures of response magnitude. For both generalist and specialist VSNs,
response reliability indices are normally distributed around zero. While the distribution of reliability
indices derived from response integral measurements is somewhat broad (Figure S1a), indices
based on signal amplitudes proved relatively homogeneous (Figure 1d). Accordingly, we use
average response amplitudes as indicators of signal strength throughout this study.

In a first set of control experiments, we asked how VSNs from male C57BL/6 mice respond when
challenged with pooled urine samples from two groups of animals of the same sex and inbred

strain (Figure S1b,c). Since the chemical composition of both stimuli should be similar, we
expected that the vast majority of urine-sensitive VSNs will display generalist response profiles.
That was indeed the case. Less than 2% of VSNs showed stimulus selectivity (Figure S1b,c).
Moreover, response preference indices (reflecting a bias toward either of the paired stimuli) are
normally (and narrowly) distributed around zero (Figure S1b,c). Both observations confirm a low
level of biological and experimental noise in this setting. Thus, our assay is well-suited to detect
and compare VSN sensory responses upon pairwise stimulation.
8

Figure 1: A population activity assay captures sex-dependent stimuli representations. (*a*) Anatomical location of the rodent VNO. Schematic depicting a sagittal section through a mouse head with an overlay of a VNO image. The vomeronasal duct that opens into the anterior nasal cavity is also highlighted. (***b***) Overview (left) and zoomed-in (middle) differential interference contrast micrograph showing an acute coronal VNO section from an adult C57BL/6 mouse. Confocal fluorescence image (right; dashed rectangle in middle) depicting a depolarization (elevated K⁺) dependent cytosolic Ca²⁺ increase in VSN somata after bulk loading with Cal-520 AM. (*c*) Original traces

showing changes in cytosolic Ca²⁺ concentration over time in two representative VSN somata. VNO slices were briefly
challenged with two mixtures of diluted mouse urine (1:100; 10s; top yellow bars/droplets). Repeated stimulation in
alternating sequence at 180 s inter-stimulus intervals [35] was followed by membrane depolarization upon exposure to
elevated extracellular K⁺ (50 mM; 10 s). According to individual response type, VSNs were categorized as 'generalists'
(top) or 'specialists' (bottom). Signal amplitudes in response to the same stimulus allowed calculation of a reliability
index (RI) as a measure of signal robustness. (*d*) Amplitude reliability index histograms of all generalist VSNs (gray; n
= 10,258) and all specialist VSNs (red; n = 6,457) recorded in this study. Note that for both response types, indices are
normally distributed with a narrow central peak around zero. (*e*) Quantification of results obtained from recordings in
VSNs from male C57BL/6 mice challenged with male *versus* female C57BL/6 urine stimuli. Pie chart (top) illustrates the
proportions of generalist (19%) and specialist neurons (1% and 4%, respectively) among all K⁺-sensitive VSNs (n =

1,999). Bar graph (middle, left) breaks down the summed total of urine-sensitive neurons by categories and compares their distribution with the proportions of VOCs (yellow background; n = 405) and proteins (red background; n = 601) found either in both male and female urine (grey bars) or exclusively in samples from one sex (male, dark blue; female, purple). Box-and-whisker plots (bottom, left) illustrating generalists-to-specialists ratios over individual experiments (n = 19). Boxes represent the first-to-third quartiles. Whiskers represent the 10th and 90th percentiles, respectively. Outliers (1.5 IQR; red x) are plotted individually. The central red band represents the population median (P$_{0.5}$). Results are shown in relation to box-and-whisker plots that outline chemical content data obtained from paired comparisons (n = 10 individuals per group). (***f***) Molecular composition of male *versus* female urine from BALB/c and wild mice. Bar graphs (top) display proportions of VOCs (yellow background; BALB/c, n = 514; wild, n = 462) and proteins (red background; BALB/c, n = 407; wild, n = 526) found either in both male and female urine (grey bars) or exclusively in samples from one sex (male, dark blue; female, purple). Box-and-whisker plots (bottom) quantify category ratios for individual paired experiments (n = 10 individuals per group). (***g***) Quantification (Bradford assay) of protein/peptide content in urine samples from C57BL/6, BALB/c, and wild mice, respectively (n = 10 each). Note the substantially increased protein content in male samples. (***h***) Response index histogram (top) obtained from generalist C57BL/6 VSNs that responded to both male and female same-strain urine (data corresponding to (***e***)). Fitted Gaussian curve (dashed line) centers close to zero (peak = -0.03) and shows a relatively narrow width (σ = 0.13). By contrast, concentration index histograms (bottom), calculated for VOCs (yellow) and proteins (red) found in both male and female urine samples, are heterogeneous and not normally distributed. Asterisks (*) indicate statistical significance, *p* < 0.05; Wilcoxon singed ranked test (for VSN functional data), Mann-Whitney U test (for molecular profiling), and unpaired t-test (for protein content comparison shown in (***g***)).

Sex-specific stimuli elicit distinct VSN sensory representations

We next investigated how sex differences are reflected in VSN response profiles. When challenged with pooled male *versus* female urine from C57BL/6 animals, the fraction of specialist neurons more than doubled to 5%. Notably, female urine recruited more specialist neurons than male urine (Figure 1e). We then asked whether this pattern is correlated with the chemical compositions of male and female urine. In-depth molecular analysis of urine content via GCxGC-MS as well as nLC-MS/MS identified a total of 1006 molecules (detected in ≥3 out of 10 male or female samples, respectively), of which approximately 40% are low molecular weight VOCs, while 60% are proteins. Roughly half of the molecules in either group are found in both male and female urine (Figure 1e). Unexpectedly, while we hardly identify any male-specific proteins in C57BL/6 urine, we find a large fraction of female-specific proteins. We asked whether this phenomenon is (*i*) a distinct feature of C57BL/6 mice, (*ii*) common among inbred laboratory strains, or (*iii*) also observed in wild animals. Therefore, we included urine samples from both BALB/c and wild mice in extended molecular profiling. In both groups, we observed similarly increased levels of female-specific proteins (Figure 1f). The total amount of protein, however, is substantially enriched in male urine (Figure 1g). Our data thus suggest that, while females secrete a larger variety of proteins, overall concentrations are comparatively low.

Finally, we asked whether common compounds (i.e., molecules identified in both male and female urine) show concentration disparities between sexes and, if so, how such differences are reflected in VSN response profiles. When VSNs were challenged with male *versus* female C57BL/6 urine, 19% of all neurons responded to both stimuli (Figure 1e). Here, again, response preference indices are narrowly distributed and center around zero (Figure 1h). By contrast, concentration

indices that can reflect potential disparities are distributed more broadly and non-normally (Figure
1h). This apparent incongruence between nonpreferential generalist sensitivity and relatively large
chemical concentration differences indicates that VSN responses (at least in C57BL/6 males) do
not trivially reflect the information theoretically available via concentration differences between
male and female same-strain stimuli. One explanation for this is that generalist ligands are not
well-represented by the global observed concentration disparities. Alternatively, it may be that
even low concentrations fully activate a given VSN, and thus concentration differences are not
reflected by response strengths. Notably, a broad and non-normal distribution of concentration
indices among molecular components of male *versus* female urine is also found in samples from
both BALB/c and wild mice (Figure S1d).

Comprehensive chemical characterization of urine content identifies distinctive molecular
signatures of sex and strain

Which differences in chemical composition (i.e., which molecules) characterize sex-/strain-specific
secretions and may thus provide information about sex and strain to a recipient? Comprehensive
molecular profiling of urine content provides a unique opportunity to address this question and
identify enriched compounds (i.e., present in ≥6 of 10 individual samples) with the potential
capacity to signal an animal's sex, strain, or both. Adopting this more conservative criterion, we
detected 208 abundant VOCs and 264 abundant proteins. Almost half of all VOCs (47.6%) and
about one-third of all proteins (31.4%) are found in all six sex/strain combinations tested and are,
thus, considered generic mouse urine components (Figure 2a,b). Moreover, a total of 33
compounds (10 VOCs, 23 proteins) were identified across all strains, but in a sex-specific fashion.
Similarly, 22 molecules (10 VOCs, 12 proteins) revealed strain selectivity, independent of sex.
Notably, a large fraction of compounds, i.e., 34.1% of all VOCs (71 / 208) and 22.3% of proteins
(59 / 264), were detected exclusively in one of the six sex/strain combinations. While 32.9% of all
proteins (87 / 264) could not be categorized as either generic or specific (for either sex, or strain,
or a unique sex/strain combination), only 18 of the 208 VOCs (8.7%) did not fall into either
category. Among the 82 proteins that showed sex specificity, either across strains (23 proteins) or
as part of a unique sex/strain combination (59 proteins), the vast majority (89.0%) was found in
female samples. For VOCs, however, the opposite picture emerged. Here, 52 of 81 sex-specific
compounds (64.2%) were selectively detected in male samples. Together, chemical profiling
revealed (*i*) that large fractions of urine content are shared among both laboratory and wild mice;
(*ii*) that roughly one-third of urinary VOCs and one-fourth of proteins are exclusively found in a
given sex/strain combination; and (*iii*) that male and female mice might have adopted different
chemical secretion strategies to signal their sex.

Figure 2: **Chemical profiling of urine content identifies unique sex- and strain-specific molecular fingerprints.** (*a*&*b*) Matrix layout for all intersections of VOCs (*a*) and proteins (*b*) among the six sex/strain combinations, sorted by size. Colored circles in the matrix indicate combinations that are part of the intersection. Bars above the matrix columns represent the number of compounds in each intersection. Empty intersections have been removed to save space. Horizontal bar charts (bottom, left) depict the number of VOCs (*a*) and proteins (*b*) detected in each urine set. Proteomics data are available via ProteomeXchange with identifier PXD042324. (*c-h*) Sparse Partial Least-Squares Discriminant Analysis (sPLS-DA) score plots depicting the first two sPLS-DA components, which explain 7-10% (1st

component) and 5-8% (2nd component) of VOC data variance (*c-e*) as well as 21-24% (1st component) and 11-12% (2nd component) of protein data variance (*f-h*), respectively. Ellipses represent 95% confidence intervals. Plots demonstrate sample clustering according to the urine donors' sex (*c*&*f*), genetic background (*d*&*g*), or sex/strain combination (*e*&*h*). Each data point represents a sample from an individual animal (n = 60; 10 samples per sex/strain combination), with sample type colored according to symbol legend (bottom).

If information coding along the accessory olfactory pathway would strictly follow a 'labeled-line' logic [42], absence or presence of a given molecule could be adequate to signal sex or strain. Several recent lines of evidence, however, suggest a combinatorial coding strategy that also involves some level of circuit plasticity [4,21,43]. Therefore, we next asked if the total VOC or protein content of a given urine sample preserves sufficient predictive / discriminative information to classify samples according to sex, strain, or a specific sex/strain combination. We used sparse Partial Least-Squares Discriminant Analysis (sPLS-DA) [44], a chemometrics machine learning technique, to reduce data dimensionality and optimize sample separation [45]. When plotted on two-dimensional coordinates that represent the most discriminative variables (Figure 2c-h), both VOCs and proteins provide sufficient information to cluster stimuli according to sex (Figure 2c,f), strain

(Figure 2d,g), or a combination of both variables (Figure 2e,h). We then calculated Variable
Importance in Projection (VIP) scores as measures of a particular variable's informative power,
which is correlated to the variance explained by the model [44,45]. Generally, we find that protein
content exhibits stronger discriminative power (21-24% & 11-12% of explained variance) than
VOC content (7-10% & 5-8% of explained variance).

We next aimed to identify the most relevant variables (i.e., molecules) for sex or strain
classification. Training Random Forest classifiers [40], we obtained feature significance scores, the
"Gini importance" [40], that provide relative relevance rankings of the individual variables.
Supplementary Figure 2 lists the 20 most informative VOCs and proteins that discriminate sex and
strain, respectively (Figure S2a,b), along with the abundance of the corresponding top 5 molecules
(VOCs or proteins) for each of the six sex/strain combinations (Figure S2c-f). Notably, major
urinary protein 20 (Mup20 / darcin) [46], fatty acid-binding protein 5 (Fabp5) [47], and N-
acetylgalactosamine-6-sulfatase (Galns) [48] exhibit substantial power to discriminate sex across
strains (Figure S2b,f). While Mup20 and Galns are considerably more abundant in male urine,
Fabp5 appears to be specific for female samples. Strain discrimination, on the other hand, is
optimal with protein-tyrosine kinase 2-beta (Ptk2b) [49], RNase T2 (Rnaset2) [50], prosaposin (Psap)
[51], lymphocyte antigen 6A-2/6E-1 (Ly6a) [52], and superoxide dismutase 1 (Sod1) [53] (Figure S2b,d).
Specifically, Ptk2b is detected exclusively in C57BL/6 mice. Ly6a is absent in BALB/c animals,
whereas Rnaset2 is largely missing in both C57BL/6 and wild mice. With proteins exhibiting
stronger discriminative power than VOCs (Figure 2c-h), future studies will have to focus on these
proteins to identify potential functions as vomeronasal chemosignals.

Overall, we have detected similar amounts of proteins / VOCs across strains. However, individual
variability in protein and VOC content was significantly higher in wild mice than in both laboratory
strains (Figure S3a). We confirmed secretion of previously reported putative semiochemicals
[10,11,24,43,54], including both VOCs (Figure S3b) and proteins (Figure S3c) in both wild and laboratory
mice. Notably, however, all Mups (including Mup20) and most such VOCs were found in samples
from either sex, albeit at male biased concentrations for Mup3, Mup17, and Mup20 (Figure S3c).
The only compounds showing male-specific secretion are 2-sec-butyl-4,5-dihydrothiazole (SBT)
and farnesenes (Figure S3b), which have previously been implicated as facilitators of female
mouse puberty acceleration [11].

Notably, we find a rich repertoire of 27 lipocalins in mouse urine. When based exclusively on
lipocalin content, hierarchical clustering groups individual samples into a set of clusters that, with
very few exceptions (5 out of 60), correspond to the six sex/strain combinations (Figure S4). This
finding, therefore, demonstrates the power of urinary lipocalins for potential chemosensory
discrimination of sex and/or strain.

Increased VSN selectivity upon exposure to strain-specific stimuli

Next, we challenged neurons with stimulus pairs from two same-sex / different-strain combinations and asked whether VSN response profiles reflect the molecular fingerprint of corresponding urine samples (e.g., male C57BL/6 *versus* male BALB/c (Figure 3a)). We analyzed a total of 17,416 K$^+$-sensitive VSNs (Figure 3a-f). Again, we distinguished between specialist VSNs that responded exclusively to one stimulus, and generalists, which responded to both stimuli. Along the same lines, we categorized either strain-specific or broadly detected VOCs and proteins. Several conclusions emerge from these classifications: (*i*) with one exception (i.e., upon exposure to male C57BL/6 *versus* wild stimuli (Figure 3b)), roughly half of all urine-sensitive VSNs are generalists – a result consistent with our finding that generally ~50% of urine molecules are shared among compared strains. Accordingly, the fraction of strain-selective (specialist) responses is considerably larger than observed for sex-specific responses (Figure 1e); (*ii*) in female urine, BALB/c-specific proteins are substantially underrepresented, a fact not reflected by VSN response

profiles (Figure 3d,f); (*iii*) surprisingly, the amount of strain-specific molecules in wild mouse urine does not vastly exceed that in inbred strains; and (*iv*) accordingly, selective VSN responses to wild stimuli are by no means more common (Figure 3b,c,e,f).

Figure 3: Selective VSN response profiles upon exposure to strain-specific signatures. Quantitative comparison between VSN responses to paired stimuli and their respective chemical signatures. Neurons of male C57BL/6 mice were challenged with male (***a-c***) or female (***d-f***) urine, respectively. Response profiles are compared upon exposure to C57BL/6 *versus* BALB/c urine (***a&d***), C57BL/6 *versus* wild stimuli (***b&e***), and BALB/c *versus* wild urine (***c&f***), respectively. Pie charts (top) illustrate the proportions of generalist (light gray) and specialist neurons (dark gray, white, and purple, respectively) among all K$^+$-sensitive VSNs (***a***, n = 1855; ***b***, n = 4116; ***c***, n = 2462; ***d***, n = 2376; ***e***, n = 3230; ***f***, n = 3377). Bar graphs

(left) break down the urine-sensitive neurons by categories and compare their distribution with the proportions of VOCs
(middle; yellow background; ***a***, n = 450; ***b***, n = 448; ***c***, n = 479; ***d***, n = 480; ***e***, n = 405; ***f***, n = 492) and proteins (right;
red background; ***a***, n = 317; ***b***, n = 330; ***c***, n = 334; ***d***, n = 657; ***e***, n = 715; ***f***, n = 584) found either in both urine types
(gray bars) or exclusively in samples from one group (color code as in pie charts). Box-and-whisker plots (bottom)
illustrate generalist-to-specialist VSN category ratios over individual experiments (***a***, n = 6; ***b***, n = 10; ***c***, n = 7; ***d***, n = 9;
e, n = 10; ***f***, n = 12). Boxes represent the first-to-third quartiles. Whiskers represent the 10th and 90th percentiles,
respectively. Outliers (1.5 IQR; red x) are plotted individually. The central red band represents the population median
($P_{0.5}$). Results are shown in relation to box-and-whisker plots that outline chemical content data obtained from paired
comparisons (n = 10 individuals per group).

Pronounced strain-dependent concentration imbalances between common urinary compounds are not reflected by generalist VSNs

When comparing generalist VSN responses to male *versus* female C57BL/6 urine (Figure 1e,h)
we noted that the narrow normally distributed stimulus preference index histograms did not match
the broader and heterogeneous distributions of concentration disparities between sexes (Figure
1h; S1b-d). For sex-dependent cues, as mentioned above, this could indicate that semiochemical
concentration differences carry only limited information. Next, we therefore asked whether strain-
specific concentration differences between urine samples exist and, if so, whether such
differences can convey information about strain. In total, we recorded strain-independent
generalist responses from 3,366 VSNs in male C57BL/6 mice (Figure 4). In all but one
experimental condition (i.e., when comparing male stimuli from BALB/c *versus* wild mice)
response preference indices were normally distributed and well fit by relatively narrow Gaussian
curves that centered around zero. By contrast, chemical analysis revealed that many compounds
identified in urine from both strains differ substantially in concentration. For proteins, in particular,
strong concentration disparities exist in all strain combinations analyzed. In both male and female
samples, concentrations of common proteins are relatively low in C57BL/6, moderate in BALB/c,
and high in wild animals. As shown in Figures 2b and 3, this concentration bias towards wild
proteins does not translate into any dramatic overrepresentation of proteins selectively found in
wild male and/or female urine. In fact, the massively skewed distributions of protein concentration
indices are not reflected by generalist VSN profiles. The latter better match VOC concentration
distributions, which generally display broad, yet Gaussian shapes. We conclude that VSN
population response strength is hardly affected by strain-dependent concentration differences
among common urinary proteins. Thus, it appears somewhat unlikely that individual VSN activity
provides fine-tuned information about distinct semiochemical concentrations. Alternatively,
generalist VSNs might sample information from only a subset of compounds which, in fact, are
secreted at roughly similar concentrations.

Vomeronasal representation of female semiochemicals differs between two inbred strains

So far, our approach was restricted to VSN signals recorded from male C57BL/6 mice. An

important question, of course, is whether the response profiles we observed are themselves

recipient strain-dependent. Thus, we next aimed to assess the extent to which our findings

generalize to VSN populations from another laboratory animal strain. We therefore repeated all

six pairwise same-sex / different-strain stimulation experiments, using acute VNO slices from male

BALB/c mice (Figures 5 & S5). Categorization as generalist or specialist VSNs revealed that

proportions differed significantly between VSNs from BALB/c *versus* C57BL/6 mice, albeit at

varying degrees. Similar to C57BL/6 neurons, selectivity of BALB/c neurons to wild-derived stimuli

was rather rare. In fact, (*i*) we recorded hardly any specialist responses upon exposure to urine

from wild females, and (*ii*) we found comparatively few generalist signals when wild-derived female

urine was among the paired stimuli (Figure 5e,f). This striking insensitivity is not observed in VSNs

from C57BL/6 mice, suggesting substantial differences in VR expression between the two inbred

laboratory strains.

Figure 4: Strain-dependent concentration imbalances exert relatively mild effects on VSN population response homogeneity. Comparison of male C57BL/6 generalist VSN response preferences, upon exposure to paired urine stimuli from different strains, with strain-dependent concentration (im)balances among VOCs and proteins, respectively. (*a*&*b*) Response index histograms (top rows) depict distributions of generalist data outlined in Figure 3 (gray bars). With one exception (*a* (right), male BALB/c *versus* male wild), histograms are well fitted by single Gaussian curves (dashed lines) that each center relatively close to zero (*a* (left), peak = 0.08, σ = 0.18; *a* (middle), peak = -0.11, σ = 0.21; *a* (right), 1st peak = -0.12, 1st σ = 0.14; 2nd peak = 0.21, 2nd σ = 0.09; *b* (left), peak = 0.07, σ = 0.12; *b* (middle), peak = 0.03, σ = 0.23; *b* (right), peak = 0.05, σ = 0.18). Concentration index histograms (middle & bottom rows), calculated for VOCs (yellow) and proteins (red) found in both tested urine samples, are more heterogeneous. Notably, while most VOC concentration index histograms are also fitted by single, albeit broader Gaussian curves (*a* (left), peak = -0.07, σ = 0.37; *a* (middle), peak = 0.02, σ = 0.45; *a*

1   (right), 1st peak = 0.07, 1st σ = 0.32; 2nd peak = 0.64, 2nd σ = 0.06; **b** (left), peak = 0.15, σ = 0.29; **b** (middle), peak =
2   0.21, σ = 0.35; **b** (right), peak = -0.01, σ = 0.42), protein concentration imbalances are not normally distributed.
3

Next, we examined whether generalist VSN response profiles differ between male BALB/c and
C57BL/6 animals (Figure S5). In total, we recorded generalist responses from 1,741 BALB/c
neurons. When plotting response preference indices, we noticed less homogeneous distributions
than previously observed in C57BL/6 mice. Five of the six histograms showed multiple peaks and
thus could not be fitted by a single Gaussian, whereas the only histogram adhering to a normal
distribution was comparatively broad (Figure S5a). As observed in C57BL/6 neurons, the skewed
distributions of protein concentration indices were not reflected by BALB/c generalist VSN profiles.
Comparison of generalist VSN response histograms between BALB/c and C57BL/6 mice (Figure
S5c,d) revealed strong and consistent differences upon exposure to female stimuli (Figure S5d).
While single Gaussians around zero characterize C57BL/6 generalist distributions upon exposure
to female stimuli (Figure S5d), several prominent peaks emerged when fitting histograms derived
from BALB/c VSNs. Notably, for some generalist BALB/c neurons, wild-derived female stimuli are
less potent than their inbred strain counterparts. This finding either indicates reduced
concentrations of the corresponding molecules in wild urine (rendering most proteins unlikely
candidates (Figure S5a,b)), or suggests some yet to be determined form of cooperativity upon
receptor-ligand interaction. Together, for both generalist and specialist VSNs, vomeronasal
representation of female semiochemicals differs considerably between the two inbred mouse
strains.
22

In the present study, we have established a robust VSN activity assay that allows pairwise
comparison of neural selectivity and response strength upon exposure to chemically defined
natural stimuli. In-depth chemical analysis of sex- and strain-specific individual urine samples
revealed that (*i*) large fractions of urine content are shared among mice of all sex / strain
combinations; that (*ii*) the amount of molecules selectively found in wild mouse urine does not
dramatically exceed the urinary secretome of inbred strains; that (*iii*) across strains, female-
specific proteins vastly outnumber the male-specific variety, while (*iv*) overall protein concentration
is substantially enriched in male urine; that (*v*) concentrations of common proteins are relatively
low in C57BL/6, moderate in BALB/c, and high in wild animals; that (*vi*) both secreted VOC and
protein profiles provide sufficient information to distinguish sex, strain, or both; and that (*vii*) the
rich urinary lipocalin repertoire alone might allow chemosensory discrimination of sex and/or
strain.

Figure 5: Vomeronasal representation of female semiochemicals differs between inbred strains. Comparison of VSN response profiles between male BALB/c and C57BL/6 mice. Pie charts (top) illustrate the proportions of generalist (light gray) and specialist BALB/c neurons (dark gray, white, and purple, respectively) among all K+-sensitive VSNs (**a**, n = 1244; **b**, n = 2063; **c**, n = 1903; **d**, n = 1770; **e**, n = 1934; **f**, n = 1818). VSNs were challenged with male (**a-c**) or female (**d-f**) urine, respectively. Response profiles are compared upon exposure to C57BL/6 *versus* BALB/c urine (**a**&**d**), C57BL/6 *versus* wild stimuli (**b**&**e**), and BALB/c *versus* wild urine (**c**&**f**), respectively. Bar graphs break down the urine-sensitive neurons by categories and compare distributions among BALB/c neurons (left) to responses recorded from C57BL/6 VSNs (right; gray background). Box-and-whisker plots (bottom) illustrate generalist-to-specialist VSN category ratios over individual experiments (**a**, n = 8; **b**, n = 10; **c**, n = 9; **d**, n = 10; **e**, n = 9; **f**, n = 10). Boxes represent the first-to-third quartiles. Whiskers represent the 10th and 90th percentiles, respectively. Outliers (1.5 IQR; red x) are plotted individually. The central red band represents the population median ($P_{0.5}$). Asterisks (∗) indicate statistical significance, *p* < 0.05, Wilcoxon singed ranked test.

When asking how much of this chemical information is accessible to inbred male mice via vomeronasal sampling we observe that (*i*) VSN population response profiles do not reflect the global molecular content of urine, suggesting that the VNO functions as a rather selective molecular detector; that (*ii*) selective VSN responses to wild stimuli are by no means more common (in fact, selectivity to wild-derived stimuli is rather rare); that (*iii*) VSN generalist signal strength is unlikely to encode semiochemical concentrations across the entire range of compounds; that (*iv*) male BALB/c neurons display striking insensitivity when challenged with urine from wild females; and, thus, that (*v*) vomeronasal representation of female semiochemicals differs considerably between inbred strains.

Discussion

Urine is the primary source of social chemosignals among mice (and, in fact, many other

mammals) and contains both 'fixed' (i.e., genomic) information about strain, sex, individual identity,

genetic histocompatibility and background, as well as 'variable' (i.e., metabolic) information on

current social, reproductive and health status [55]. The ability to glean ethologically meaningful

information from chemosensory sampling of urine (or any conspecific bodily secretion) depends

on (*i*) the specific (semio)chemical composition of a urine sample, and (*ii*) the sensory apparatus

used for sampling. For most mammals, the VNO is the key chemosensory structure involved in

detecting conspecific chemical cues [4]. The virtually universal use of inbred laboratory mice in

research aimed at understanding VNO physiology – as both the donors of stimuli and the

experimental subjects employed in these studies – could have resulted in misconceptions and

biased notions about vomeronasal signaling and, thus, conspecific chemical communication. If

that were the case, one might question the relevance and ethological validity of conclusions drawn

from a large body of work using inbred secretions [15]. Here, we addressed this issue from both a

chemical ecology and a physiological perspective. In-depth comparative molecular profiling of

urine from two classical laboratory strains as well as wild mice reveals several shared features,

but also qualitative and quantitative differences in composition. Sex- and strain-specific chemical

profiles give rise to unique VSN activity patterns. Furthermore, we observe substantial differences

in vomeronasal representations of stimuli between C57BL/6 and BALB/c sensory neuron

populations.

For analytical purposes, we separate the urinary 'volatilome' and proteome. In chemosensory

research, this distinction has often been conceptualized as general (i.e., airborne) odors, which

activate the main olfactory system, *versus* vomeronasal stimuli [56]. This notion, however, is

misleading since organic compounds with low molecular weight and high vapor pressure (i.e.,

VOCs) in bodily secretions do not instantly evaporate, of course. Rather, they are readily

accessible for vomeronasal sampling upon direct contact during investigatory behavior. For both

urinary VOCs and proteins, large fractions are shared among both male and female mice of either

genetic background (Figure 2a & b). Such compounds could be considered generic (mouse) urine

components and might not even serve any chemosensory signaling functions. Notably, both the

urinary volatilome and proteome on their own, each entail sufficient information to discern an

individual's sex and strain, with protein content exhibiting stronger discriminative power. The

protein that is most informative for discriminating between sexes is, perhaps not surprisingly,

Mup20 (darcin) [46]. This well-described "maleness signal" had previously been reported to elicit

innate attraction and generate a conditioned place preference in females [54,57], whereas, in males,

Mup20 promotes aggression [43]. Chemosensory roles of the second and third best protein

determinants of sex discrimination are basically unexplored. Fabp5 [47] and Galns [48] are

substantially enriched in female and male urine, respectively. Fatty acid-binding proteins, including
Fabp5, are evolutionary conserved intracellular lipid chaperones that coordinate cellular lipid
trafficking and signaling and are thus linked to metabolic and inflammatory pathways [47]. N-
acetylgalactosamine-6-sulfatase (Galns) on the other hand is a lysosomal hydrolase. Five urinary
proteins – Ptk2b [49], Rnaset2 [50], Psap [51], Ly6a [52], and Sod1 [53] – display pronounced strain-
dependent differences in concentration. None of these has previously been attributed a
chemosensory function. Challenging the mouse VNO with purified recombinant protein(s) will help
elucidate whether such functions exist.

Proteomic profiling revealed three additional, rather unexpected findings: First, while male urine
contains much higher protein concentrations, females of a given strain secrete a larger variety of
proteins (including MUPs). In fact, we do not find a single protein that is exclusively detected in
males across strains. By contrast, 23 urinary proteins, while present in all three strains, are found
only in females. In line with these observations, only one additional Mup (i.e., Mup21) made the
list of the 20 most informative proteins that discriminate sex. In general, our data provide little
evidence for a sparse molecular code of (fe)maleness. Rather, the concept of 'signature mixtures'
[7,58,59], which emphasizes a combinatorial ratio code instead of mere presence/absence
phenomena, gains traction. Second, we identify a surprisingly rich lipocalin repertoire in urine,
which alone could allow chemosensory discrimination of sex/strain combinations. Within a total of
27 lipocalins, individual patterns allow hierarchical clustering into sex-/strain-specific groups
(Figure S4). These data, thus, support the notion that the lipocalin 'code', if relevant, is
combinatorial. Third, regarding its molecular spectrum, the urinary 'secretome' in wild mice does
not differ dramatically from the repertoire found in laboratory strains. To paraphrase this generally
reassuring conclusion: while inbreeding could have dramatically modified the nature of chemical
secretions and, consequently, their perception by other mice, inbred chemostimuli are largely
representative of (potentially) ethologically more relevant wild chemosignals. Notably, however,
while mean VOC and protein concentrations show similar distributions across sex/strain
combinations, individual variability is strongly increased in wild mouse urine (Figure S3a).
Accordingly, this finding confirms previous reports of increased individual variation in wild mice
[19,60]. Another inherent factor that could account for differences in urine secretions among each of
the groups, and particularly for comparisons between inbred and wild stimuli, is the microbiome
[15,61]. Yet, we stress that all urine donors were housed in the same facility and fed with the same
diet [15].

Some limitations of our study need to be acknowledged. Regarding stimuli, the six secretion sets
we used do not cover the entire coding capacity of the accessory olfactory system [62]. Moreover,
for VSN response profiling, stimulus samples were pooled across ten individuals in each of the six

sex/strain categories. While pooling stimuli reduces individual variability across samples (e.g.,

regarding fluctuating physiological states), relevant stimulus aspects could be masked. Thus,

given the increased chemical variability we observed among individual wild urine samples, pooling

might obscure distinctive molecular features of wild mouse secretions. The same holds true for

estrus cycle-dependent female stimuli. Because we did not monitor the estrus stage of female

urine donors, pooled mixes likely contain samples across the entire cycle. This is relevant as VSN

responses may be affected by a donor's cycle stage [15,34,63]. Another limitation stems from the use

of male inbred mice as experimental subjects. The rationale behind this experimental strategy is

to allow for comparisons with our previous study on AOB response profiles [15], which used the

exact same settings (see below). Nonetheless, future efforts will have to reveal (*i*) whether our

main findings apply to female recipients; and (*ii*) if our observations generalize to other inbred,

outbred, or wild mouse strains. As we recently declared [15], the latter considerations underscore

the importance of recording from wild recipients. While this endeavor presents significant practical

challenges, it remains an important goal for future studies.

A surprising, but overall reassuring observation is that responses to wild and inbred stimuli are

qualitatively similar. Strikingly, and somewhat counterintuitively, selective VSN responses to wild

stimuli are rather rare. This does not result from a generally reduced compound content in wild

urine as molecular profiling revealed comparably rich chemical portfolios in wild and inbred

samples alike. Rather, we speculate that inbreeding over hundreds of generations in laboratory

settings [16] has resulted in "microevolutionary" pressure to maintain sensitivity to signals from

same- or similar-strain individuals. Indeed, compared to the C57BL/6 reference genome, genetic

variability within VR repertoires is massively increased among wild-derived mice (particularly of

the *M. m. musculus* subspecies) [23]. Among the approximately 200 orthologous receptor genes

compared, BALB/c genes display 184 non-synonymous and just one private (i.e., unique to a

given strain) single-nucleotide polymorphism (SNP). By contrast, VR genes of the wild-derived *M.*

m. musculus PWK/PhJ strain show 789 non-synonymous SNPs and 508 private SNPs [23].

Initially, we asked whether VSN response profiles reflect the global molecular content of urine or,

by contrast, if the VNO serves as a rather selective semiochemical detector. Our findings support

a high level of selectivity. When challenged with same-sex / different-strain stimuli, large VSN

fractions selectively respond to just one stimulus. Intriguingly, this fraction of strain-selective

specialists is considerably larger than observed for sex-specific responses. In line with the notion

of highly selective vomeronasal sampling is our observation that the concentration differences

between compounds shared among strains, which are often substantial, are not reflected by

similarly pronounced differences in response strength among generalist VSNs. There are several,

not necessarily mutually exclusive explanations for this finding: First, concentration could simply

not be a read-out parameter for VSNs, which would support previous ideas of concentration-invariant VSN activity [24]. Second, the concentrations in freshly released urine could just exceed the dynamic tuning range of VSNs since, particularly for VOCs, natural signals (e.g., in scent marks) must be accessible to a recipient for a prolonged amount of time (sometimes days). A similar rationale could explain the increased protein concentrations in male urine, since male mice use scent marking to establish and maintain their territories and urinary lipocalins serve as long-lasting reservoirs of VOCs [64]. Third, generalist VSNs might sample information only from a select subset of urinary compounds, which, given their role as biologically relevant chemosignals, might be released at tightly controlled (and thus similar) concentrations.

While, compared to wild-derived mice, the genetic differences in VR repertoires between C57BL/6 and BALB/c animals appear rather modest [23] (see above), vomeronasal representation of female semiochemicals differs considerably between both inbred strains. We conclude that, even in closely related inbred mice, strain-to-strain VR variation must be prominent, an idea supported by various reports of differences in genetic VR makeup across strains [65–67]. With monoallelic VR expression and 184 described non-synonymous SNPs between C57BL/6 and BALB/c receptor genes, it is likely that even individuals of the same strain express functionally different arrays of VSNs. Adding another layer of complexity, state- and experience-dependent changes in VSN sensitivity have recently been described [21,68].

By adopting the same experimental design (i.e., using the same sets of stimuli in male C57BL/6 *versus* BALB/c mice) in both this study on VSN response profiles and our previous analysis of sensory representations in the AOB [15], we provide a unique comparative perspective on signal transformation along the initial processing nodes of the accessory olfactory pathway. We observe several differences in representations of ethologically relevant urine stimuli between the VNO and AOB. Notably, while stimulus representations across the two inbred recipient strains were very similar in AOB recordings [15], we here observe clear differences in VSN activity, particular upon exposure to female stimuli. Moreover, a substantial fraction of AOB neurons were selective to wild rather than inbred stimuli, whereas relatively few VSNs showed such selectivity. Consistent with the elaborate wiring patterns within the AOB, these observations imply the presence of non-trivial transformations between VSN and AOB representations. Understanding the exact nature, purpose, and neuronal substrates of these transformations remains an important topic for future studies.

Acknowledgements

This work was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research
Foundation) – 368482240/GRK2416 (MS); 378028035 (YBS & MS) – by the Volkswagen
Foundation (I/83533, MS), and the German-Israeli Foundation for Scientific Research and
Development (1-1193-153.13/2012, YBS & MS). We thank Corinna H. Engelhardt, Stefanie Kurth,
and Jessica von Bongartz (RWTH-Aachen University) for excellent technical assistance. We are
grateful to Pavel Talacko and Petr Zacek for Proteomic Core Facility support (BIOCEV, Charles
University).

Author contributions

Conceptualization, M.Na., P.S., Y.B.-S., and M.S.; Formal Analysis, M.Na. and P.S.; Methodology,
M.Na., D.F., R.S., P.S., and M.S.; Investigation, M.Na., M.Ni., R.B., and R.S.; Writing – Original
Draft, M.Na. and M.S.; Writing – Review & Editing, M.Na., R.B., D.F., A.L., P.S., Y.B.-S., and M.S.;
Funding Acquisition, Y.B.-S. and M.S.; Project Administration, P.S., Y.B.-S., and M.S.; Resources,
M.Na., P.S., and M.S.; Supervision, P.S., Y.B.-S., and M.S.; Visualization, M.Na., P.S., A.L., and
M.S.

Declaration of Interests

The authors declare no competing interests.

Methods

Animals.

All animal procedures were approved by local authorities at RWTH Aachen University, were
performed in accordance with local Animal Care and Use Committees' regulations, and in
compliance with European Union legislation (Directive 2010/63/EU) and recommendations by the
Federation of European Laboratory Animal Science Associations. C57BL/6 and BALB/c mice
(Charles River Laboratories, Sulzfeld, Germany) were housed in groups of both sexes [room
temperature (RT); 12:12 h light-dark cycle; food and water available *ad libitum*]. All Ca^{2+} imaging
experiments used slices from young male adults.

Urine collection from two strains of inbred mice (C57BL/6NCrl and BALB/cAnNCr) as well as first-
generation offspring of wild mice was performed at Charles University (Prague, Czech Republic)
according to the institute's ethical committee guidelines. Inbred mice were purchased from
pathogen-free facilities of the Institute of Molecular Genetics (Czech Academy of Sciences in
Prague). Wild *M. m. musculus* mice were caught in house shelters and agricultural buildings near

Prague (Czechia), transferred to the local animal facility at Charles University, and bred for one
generation. All mice that served as urine donors were fed on the same diet. Food and water for
all strains were provided *ad libitum* under stable conditions (13:11 h light-dark cycle; 23°C).

Chemicals and solutions.

The following solutions (S_1–S_6) were used:

(S_1) 4-(2-Hydroxyethyl)piperazine-1-ethanesulfonic acid (HEPES) buffered extracellular solution
containing (in mM) 145 NaCl, 5 KCl, 1 $CaCl_2$, 1 $MgCl_2$, 10 HEPES, pH 7.3 (adjusted with NaOH),
300 mOsm (adjusted with glucose).

(S_2) Oxygenated (95% O_2, 5% CO_2) extracellular solution containing (in mM) 125 NaCl, 25
$NaHCO_3$, 5 KCl, 1 $MgSO_4$, 1 $CaCl_2$, 5 N,N-bis(2-hydroxyethyl)-2-aminoethanesulfonic acid (BES);
pH = 7.3; 300 mOsm (adjusted with glucose).

(S_3) Elevated extracellular K^+ solution containing (in mM) 100 NaCl, 50 KCl, 1 $CaCl_2$, 1 $MgSO_4$,
10 HEPES; pH = 7.3 (adjusted with NaOH); 300 mOsm (adjusted with glucose).

If not stated otherwise, chemicals were purchased from Sigma (Schnelldorf, Germany). Solutions
and stimuli were applied from air pressure-driven reservoirs via an eight-in-one multibarrel
"perfusion pencil" (Science Products, Hofheim, Germany). Changes in focal superfusion [26] were
software-controlled and synchronized with data acquisition by transistor–transistor logic input to
12 V DC solenoid valves using a TIB 14S digital output trigger interface (HEKA Elektronik,
Lambrecht/Pfalz, Germany).

Stimuli

For all three donor types (C57BL/6NCrl, BALB/cAnNCr, and wild mice), we collected fresh urine
by gentle bladder massage from 10 adult male and female individuals, respectively (resulting in a
total of 60 individual samples). To minimize concentration differences that might result from
sample-to-sample volume (i.e., dilution) variability, we collected and pooled four to six samples
from each individual over several days until each of the 60 animals had provided a total urine
volume of >500 μl. Next, we measured general protein content for each sample (Bradford assay).
Aliquots of 10 μl were subjected to comprehensive two-dimensional gas chromatography–mass
spectrometry (GCxGC-MS) and nanoliquid chromatography–tandem mass spectrometry (nLC-
MS/MS; see below).

The remaining samples were divided into ready-to-use aliquots and stored at −86°C. Prior to
experiments, aliquots were thawed and diluted 1:100 in S_1 [27]. For each of the six strain-/sex-
specific stimulus combinations, we created pools from all 10 individuals to minimize individual-to-
individual variability. For both inbred and wild female mice, estrus stage was not determined.
However, urine collection over several days and pooling across 10 individuals in each stimulus
set is designed to reduce variability. Notably, the urine samples employed in this study are the
same stimuli used previously to compare sensory representations of inbred and wild stimuli in the
AOB of male C57BL/6 and BALB/c mice [15].

Comprehensive two-dimensional gas chromatography-mass spectrometry (GCxGC-MS)

Urine VOCs were sampled with Headspace Solid Phase Micro Extraction (HS SPME) on fiber

(DVB/CAR/PDMS grey; Supelco, USA) after 5 min incubation at 55 °C. Next, VOCs were analyzed

using two-dimensional comprehensive gas chromatography with mass detection (Pegasus 4D,

LECO Europe B.V., Geleen, The Netherlands) with a combination of mid-polar and non-polar

separation columns [primary column: SLB-IL60 (30 m x 0.25 mm, SigmaAldrich, USA); secondary

column Rxi-5sil MS (1.4 m x 0.25 mm, Restek, Australia)]. Parameters were set as follows: inlet

temperature 270 °C, splitless injection mode, constant He flow 1 ml min^{-1}, modulation time 4 s (hot

pulse 0.6 s), modulation temperature offset with respect to secondary oven 15 °C. Temperature

program for primary oven: 50 °C (1 min), increase to 320 °C (10 °C min^{-1}), 320 °C (3 min). +5 °C

temperature offset on secondary column. Transfer line temperature was held at 250 °C. Mass

detector was equipped with an electron ionization source and time-of-flight analyzer enabling unit

mass resolution (scanned mass range was 30 – 500 *m/z*). The ion source chamber was held at

250 °C. ChromaTOF® v4.5 software (LECO Europe B.V.) was employed for instrument control

and data processing. Selected compounds were identified by mass spectra library matching (NIST

MS 2.2, USA). Compounds identified only in blanks (or that were highly abundant in blanks) were

removed from analysis (e.g., silanes, siloxanes, propylphosphines etc.).

Protein digestion & nano-scale liquid chromatographic tandem mass spectrometry (nLC-MS/MS)

Urine proteins were precipitated with cold acetone and centrifuged (14,000 x *g*; 10 min; 0 °C),

followed by re-suspension of dried pellets in digestion buffer (1% SDC, 100 mM TEAB; pH 8.5).

Next, protein concentration in each lysate was determined (BCA assay kit; Fisher Scientific). We

used tris(2-carboxyethyl)phosphine (TCEP; 5 mM; 60 °C; 60 min) as reducing agent and S-methyl

methanethiosulfonate (MMTS; 10 mM; 10 min; RT) to block free cysteines. After trypsin digestion

(1 µg per sample; 37 °C; overnight), peptides were desalted on a Michrom C18 column. We used

reverse-phase nanocolumns (EASY-Spray column, 50 cm x 75 µm ID, PepMap C18, 2 µm

particles, 100 Å pore size) for high resolution peptide separation. Eluting peptide cations were

converted to gas-phase ions by electrospray ionization and analyzed on a Thermo Orbitrap Fusion

(Q-OT-qIT; Thermo Fisher, Waltham, MA, USA) as previously described [28,29]. LC-MS data were

pre-processed with MaxQuant software (version 1.5.3.8) [30]. The false discovery rate (FDR) was

set to 1% for both proteins and peptides. We specified a minimum peptide length of seven amino

acids. The Andromeda search engine was used for MS/MS spectra search against Uniprot *Mus*

musculus database (downloaded June 2015), containing 44,900 entries. From this database, all

MUP and OBP sequences were removed and replaced by complete lists of MUPs (Ensembl) and

OBPs [31], respectively. We also added some sequences from TrEMBL that were missing in Uniprot

(e.g., KLKs, BPIs, SPINKs, SCGB/ABPs, and LCNs). Enzyme specificity was set as C-terminal to

Arg and Lys, also allowing cleavage at proline bonds [32] and a maximum of two missed cleavages.

Quantifications were performed using label-free algorithms [33] with a combination of unique and

razor peptides.

VNO slice preparation

For confocal Ca^{2+} imaging, acute coronal VNO slices were prepared as previously described [34,35].
Briefly, C57BL/6 and BALB/c mice were euthanized by brief exposure to a CO_2 atmosphere,
cervical dislocation and decapitation. The lower jaw and palate were rapidly removed. The VNO
was dissected, embedded in 5% low-gelling temperature agarose (VWR International, Erlangen,
Germany), placed in ice-cold oxygenated **S₂**, and coronal slices (150 µm) were cut on a VT1000S
vibrating microtome (Leica Biosystems, Nussloch, Germany). Slices were transferred to a
submerged, chilled, and oxygenated storage chamber with circulating **S₂** until use.

Ca^{2+} imaging

In vitro imaging of VSN activity in acute coronal VNO slices was performed as described [35]. Briefly,
for bulk loading, slices were incubated (90 min; 5°C) in circulating **S₂** with the Ca^{2+}-sensitive dye
CAL520/AM (4.5 µM; Biomol, Hamburg, Germany) and 0.05 % Pluronic® F-127 (20 % solution in
DMSO; Thermo Fisher Scientific, Schwerte, Germany). After washing (5x, **S₂**), slices were
transferred to a recording chamber (Luigs & Neumann, Ratingen, Germany) mounted on an
upright fixed-stage scanning confocal microscope (TCS SP5 DM6000CFS, Leica Microsystems)
equipped with a 20x / 1.0 NA water immersion objective (HCX APO L, Leica Microsystems), and
infrared-optimized differential interference contrast optics. Slices were continuously superfused
with oxygenated **S₂** (~5 ml / min; gravity flow). CAL520 was excited at 488 nm (multi-line argon
laser; <25% laser power) and fluorescence was detected within a 500–600 nm spectral band.
Changes in cytosolic Ca^{2+} were monitored over time at 1.0 Hz frame rate (1024 x 512 pixels; 400
Hz bidirectional scanning frequency) using LAS AF software (Leica Microsystems).

Experimental Design and Statistical Analysis.

Ca^{2+} imaging in VNO slices – All data were obtained from independent experiments performed in
≥3 sessions using ≥3 different animals. Individual numbers of cells / experiments (n) are denoted
in figures and/or captions. Data were analyzed offline using Leica LAS AF 2.4 (Leica
Microsystems), ImageJ 1.51n (Wayne Rasband, National Institutes of Health, USA), MATLAB
R2017b (MathWorks, Natick, MA), and Excel (Microsoft, Seattle, WA) software. If not stated
otherwise, results are presented as box-and-whisker plots, where boxes represent the first-to-third
quartiles and whiskers represent the 10^{th} and 90^{th} percentiles, respectively. Outliers (1.5 IQR) are
plotted individually. The central band represents the population median ($P_{0.5}$). Statistical analyses
were performed using unpaired t-tests, Wilcoxon signed ranked tests, Mann-Whitney U tests, and
two-sample Kolmogorov-Smirnov tests (as dictated by data distribution and experimental design).
Tests and corresponding *p*-values that report statistical significance (≤0.05) are individually
specified in figure captions.

For individual cell analysis, regions-of-interest (ROIs) were defined to outline essentially all
depolarization-sensitive (**S₃**) somata per field of view, based on DIC imaging of cell morphology
at rest. After movement correction (StackReg/Rigid Body transformation plugin[36] in ImageJ) of
time-lapse image stacks, changes in relative fluorescence intensity were calculated as ΔF/F and
measured in arbitrary units. Neurons were classified as 'responsive' when showing stimulus-

dependent Ca^{2+} elevations in somata according to the following three criteria[35,37]: (*i*) exposure to
high extracellular K^+ concentrations (50 mM; **S₃**) induced a robust Ca^{2+} transient; (*ii*) for at least
one exposure to diluted urine, a transient increase in fluorescence intensity was observed during
the stimulation period; and (*iii*) the signal peak intensity exceeded the average prestimulation
baseline intensity plus two standard deviations for a continuous period of at least 3 s ($I_{resp} > I_{baseline}$
$+ 2 \times SD(I_{baseline})$). All responses were normalized to positive controls (i.e., responses evoked by
elevated extracellular K^+). Responsive neurons were categorized as: (*i*) neurons only sensitive to
depolarization (K^+); (*ii*) *specialist* neurons that selectively responded to one of two presented urine
stimuli (and K^+); and (*iii*) *generalist* neurons that responded to both urine stimuli (and K^+). Raw
data (i.e., original intensity *versus* time traces) from each responsive cell were visually inspected
to control for potential unspecific signals (e.g., caused by spontaneous activity). Based on peak
amplitudes recorded from each urine-sensitive neuron (both generalists and specialists), we
calculated a cell-specific *preference index* (PI):

$$PI = \frac{mean\left(\frac{\Delta F}{F}\right)_{stim1} - mean\left(\frac{\Delta F}{F}\right)_{stim2}}{mean\left(\frac{\Delta F}{F}\right)_{stim1} + mean\left(\frac{\Delta F}{F}\right)_{stim2}}$$

where mean ($^{\Delta F}/_F$) is the average signal amplitude evoked by consecutive exposure to the same
stimulus (i.e., either stim1 or stim2). As a measure of how reliably the same stimulus evokes a
response upon consecutive exposures, we additionally calculated a cell-specific *Reliability Index*
(RI), which is based either on response amplitudes or their integrals (i.e., area under curve (AUC)):

$$RI = \frac{response\left(\frac{\Delta F}{F} \, or \, AUC\right)_{1st\,stim} - response\left(\frac{\Delta F}{F} \, or \, AUC\right)_{2nd\,stim}}{response\left(\frac{\Delta F}{F} \, or \, AUC\right)_{1st\,stim} + response\left(\frac{\Delta F}{F} \, or \, AUC\right)_{2nd\,stim}}$$

where consecutive responses of similar strength (i.e., high reliability) are reflected by a small RI
value.

Proteomics and metabolomics – Pairwise comparison between different urine samples allowed
comparative analysis of physiological activity (VSN Ca^{2+} signals) *versus* urine composition
(molecular content). Chemical content was categorized as follows: (*i*) a specific compound is
considered "present" in urine from a given sex/strain combination (i.e., male or female BALB/c,
C57BL/6, or wild, respectively) if it is identified in ≥3 out of 10 individual samples; (*ii*) a compound
is thus considered "absent" if it is found only in ≤2 individual samples. Accordingly, compounds
that are detected as "present" in urine samples from both sex/strain combinations being compared
are designated as *generic* in binary comparisons. By contrast, a compound that is "present" in
only one of the two sex/strain combinations is designated as *specific*. To identify compounds that
are *enriched* in a given sex/strain combination (Figure 2a,b) we raised the criterion for a "present"
call to identification in ≥6 out of 10 individual samples.
To quantify concentration differences between samples we calculated *Concentration Indices* (CIs)
for generic compounds, both VOCs (*GCxGC-MS*) and peptides/proteins (*nLC-MS/MS*):

$$CI = \frac{mean \, conc. \, \left(X_{group1}\right) - mean \, conc. \, \left(X_{group2}\right)}{mean \, conc. \, \left(X_{group1}\right) + mean \, conc. \, \left(X_{group2}\right)}$$

where $mean\ conc.$ is the average concentration of a given compound *X* among all ten samples
within a group (i.e., specific sex/strain combination).
3
Proteomic and metabolomic analysis

Because both GCxGC-MS and nLC-MS/MS data have similar (negative binomial) distributions,
we processed them using the same procedure. First, we performed Sparse Partial Least Squares
Analysis - sPLS-DA[38] to detect potential sources of variation in quantile normalized datasets. Next,
for pairwise comparisons, data reduction eliminated all table entries (rows) if a given metabolite
or protein was detected only in ≤3 individuals in both groups. However, if a compound was found
in samples from ≥4 individuals in a group (e.g., in males or females), we included the full table
entry (row). Accordingly, a chemical is considered "unique" / group-specific if found in ≥4 individual
samples from one group, but is missing in all 10 samples of the other group. For calculation of
sexual dimorphism, we used the Power Law Global Error Model - PLGEM[39] to detect differentially
expressed / abundant proteins and VOCs. To detect the importance of significant molecules in
discriminating either sex or strain, we used a machine learning technique – Random Forest for
Classification [40] (implemented in R software) – and inferred feature importance scores (Gini
importance [40]) that provide relative rankings of individual variable relevance. nLC-MS/MS
proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE [41]
partner repository with the dataset identifier PXD042324. Metabolomics data have been deposited
to the EMBL-EBI MetaboLights database with the identifier MTBLS7439.
21
22
23   ***Supplemental information***
24   *Supplementary Figures S1 - S5*

References

1. Dulac, C. & Torello, A. T. Molecular detection of pheromone signals in mammals: from genes to behaviour. *Nat. Rev. Neurosci.* **4**, 551–562 (2003).

2. Brennan, P. A. & Zufall, F. Pheromonal communication in vertebrates. *Nature* **444**, 308–315 (2006).

3. Tirindelli, R., Dibattista, M., Pifferi, S. & Menini, A. From Pheromones to Behavior. *Physiol. Rev.* **89**, 921–956 (2009).

4. Mohrhardt, J., Nagel, M., Fleck, D., Ben-Shaul, Y. & Spehr, M. Signal detection and coding in the accessory olfactory system. *Chem. Senses* **43**, 667–695 (2018).

5. Wilson, K. C. P. & Raisman, G. Age-related changes in the neurosensory epithelium of the mouse vomeronasal organ: Extended period of post-natal growth in size and evidence for rapid cell turnover in the adult. *Brain Res.* **185**, 103–113 (1980).

6. Luo, M., Fee, M. S. & Katz, L. C. Encoding Pheromonal Signals in the Accessory Olfactory Bulb of Behaving Mice. *Science* **299**, 1196–1201 (2003).

7. Wyatt, T. D. Pheromones. *Curr. Biol.* **27**, R739–R743 (2017).

8. Albone, E. S. *Mammalian semiochemistry : the investigation of chemical signals between mammals*. (John Wiley & Sons, Inc., 1984).

9. Nodari, F. *et al.* Sulfated steroids as natural ligands of mouse pheromone-sensing neurons. *J. Neurosci.* **28**, 6407–18 (2008).

10. Hurst, J. L. *et al.* Individual recognition in mice mediated by major urinary proteins. *Nature* **414**, 631–634 (2001).

11. Novotny, M. V. Pheromones, binding proteins and receptor responses in rodents. *Biochem. Soc. Trans.* **31**, 117–122 (2003).

12. Röck, F., Mueller, S., Weimar, U., Rammensee, H.-G. & Overath, P. Comparative analysis of volatile constituents from mice and their urine. *J. Chem. Ecol.* **32**, 1333–46 (2006).

13. Sturm, T. *et al.* Mouse urinary peptides provide a molecular basis for genotype discrimination by nasal sensory neurons. *Nat. Commun.* **4**, 1616 (2013).

14. Overath, P., Sturm, T. & Rammensee, H.-G. Of volatiles and peptides: in search for MHC-dependent olfactory signals in social communication. *Cell. Mol. Life Sci.* (2014) doi:10.1007/s00018-014-1559-6.

15. Bansal, R. *et al.* Do all mice smell the same? Chemosensory cues from inbred and wild mouse strains elicit stereotypic sensory representations in the accessory olfactory bulb. *BMC Biol.* **19**, 133 (2021).

16. Phifer-Rixey, M. & Nachman, M. W. Insights into mammalian biology from the wild house mouse Mus musculus. *Elife* **4**, 1–13 (2015).

17. Wade, C. M. & Daly, M. J. Genetic variation in laboratory mice. *Nat. Genet.* **37**, 1175–1180 (2005).

18. Duvaux, L., Belkhir, K., Boulesteix, M. & Boursot, P. Isolation and gene flow: inferring the speciation history of European house mice. *Mol. Ecol.* **20**, 5248–5264 (2011).

19. Cheetham, S. A., Smith, A. L., Armstrong, S. D., Beynon, R. J. & Hurst, J. L. Limited variation in the major urinary proteins of laboratory mice. *Physiol. Behav.* **96**, 253–261 (2009).

20. Duyck, K., DuTell, V., Ma, L., Paulson, A. & Yu, C. R. Pronounced strain-specific chemosensory receptor gene expression in the mouse vomeronasal organ. *BMC Genomics* **18**, 965 (2017).

21. Xu, P. S., Lee, D. & Holy, T. E. Experience-Dependent Plasticity Drives Individual Differences in Pheromone-Sensing Neurons. *Neuron* **91**, 878–92 (2016).

22. Ibarra-Soria, X. *et al.* Variation in olfactory neuron repertoires is genetically controlled and environmentally modulated. *Elife* **6**, 1–29 (2017).

23. Wynn, E. H., Sánchez-Andrade, G., Carss, K. J. & Logan, D. W. Genomic variation in the vomeronasal receptor gene repertoires of inbred mice. *BMC Genomics* **13**, 415 (2012).

24. Leinders-Zufall, T. *et al.* Ultrasensitive pheromone detection by mammalian vomeronasal neurons. *Nature* **405**, 792–796 (2000).

25. He, J. *et al.* Distinct signals conveyed by pheromone concentrations to the mouse vomeronasal organ. *J. Neurosci.* **30**, 7473–7483 (2010).

26. Veitinger, S. *et al.* Purinergic signalling mobilizes mitochondrial Ca^{2+} in mouse Sertoli cells. *J. Physiol.* **589**, 5033–55 (2011).

27. Hagendorf, S., Fluegge, D., Engelhardt, C. H. & Spehr, M. Homeostatic control of sensory output in basal vomeronasal neurons: activity-dependent expression of ether-à-go-go-related gene potassium channels. *J. Neurosci.* **29**, 206–21 (2009).

28. Černá, M., Kuntová, B., Talacko, P., Stopková, R. & Stopka, P. Differential regulation of vaginal lipocalins (OBP, MUP) during the estrous cycle of the house mouse. *Sci. Rep.* **7**, 1–10 (2017).

29. Stopkova, R., Klempt, P., Kuntova, B. & Stopka, P. On the tear proteome of the house mouse (Mus musculus musculus) in relation to chemical signalling. *PeerJ* **2017**, (2017).

30. Voukali, E., Veetil, N. K., Němec, P., Stopka, P. & Vinkler, M. Comparison of plasma and cerebrospinal fluid proteomes identifies gene products guiding adult neurogenesis and neural differentiation in birds. *Sci. Rep.* **11**, 1–16 (2021).

31. Stopková, R., Otčenášková, T., Matějková, T., Kuntová, B. & Stopka, P. Biological Roles of Lipocalins in Chemical Communication, Reproduction, and Regulation of Microbiota. *Front. Physiol.* **12**, (2021).

32. Rodriguez, J., Gupta, N., Smith, R. D. & Pevzner, P. A. Does Trypsin Cut Before Proline? *J. Proteome Res.* **7**, 300–305 (2007).

33. Cox, J. *et al.* Accurate Proteome-wide Label-free Quantification by Delayed Normalization and Maximal Peptide Ratio Extraction, Termed MaxLFQ. *Mol. Cell. Proteomics* **13**, 2513–2526 (2014).

34. Cichy, A. *et al.* Extracellular pH Regulates Excitability of Vomeronasal Sensory Neurons. *J. Neurosci.* **35**, 4025–4039 (2015).

35. Wong, W. M. *et al.* Sensory Adaptation to Chemical Cues by Vomeronasal Sensory Neurons. *Eneuro* **5**, ENEURO.0223-18.2018 (2018).

36. Thevenaz, P., Ruttimann, U. E. & Unser, M. A pyramid approach to subpixel registration based on intensity. *IEEE Trans. Image Process.* **7**, 27–41 (1998).

37. Fluegge, D. *et al.* Mitochondrial Ca2+ mobilization is a key element in olfactory signaling. *Nat. Neurosci.* **15**, 754–762 (2012).

38. Rohart, F., Gautier, B., Singh, A. & Lê Cao, K.-A. mixOmics: An R package for 'omics feature selection and multiple data integration. *PLOS Comput. Biol.* **13**, 1–19 (2017).

39. Pavelka, N. *et al.* A power law global error model for the identification of differentially expressed genes in microarray data. *BMC Bioinformatics* **5**, 203 (2004).

40. Breiman, L. Random Forests. *Mach. Learn. 2001 451* **45**, 5–32 (2001).

41. Perez-Riverol, Y. *et al.* The PRIDE database resources in 2022: a hub for mass spectrometry-based proteomics evidences. *Nucleic Acids Res.* **50**, D543–D552 (2022).

42. Ishii, K. *et al.* A Labeled-Line Neural Circuit for Pheromone-Mediated Sexual Behaviors in Mice. *Neuron* **1**, 1–15 (2017).

43. Kaur, A. W. *et al.* Murine Pheromone Proteins Constitute a Context-Dependent Combinatorial Code Governing Multiple Social Behaviors. *Cell* **157**, 676–688 (2014).

44. Lê Cao, K. A., Boitard, S. & Besse, P. Sparse PLS discriminant analysis: Biologically relevant feature

selection and graphical displays for multiclass problems. *BMC Bioinformatics* **12**, 0–16 (2011).

45. Gromski, P. S. *et al.* A tutorial review: Metabolomics and partial least squares-discriminant analysis – a marriage of convenience or a shotgun wedding. *Anal. Chim. Acta* **879**, 10–23 (2015).

46. Roberts, S. A. *et al.* Darcin: a male pheromone that stimulates female memory and sexual attraction to an individual male's odour. *BMC Biol.* **8**, 75 (2010).

47. Furuhashi, M. & Hotamisligil, G. S. Fatty acid-binding proteins: role in metabolic diseases and potential as drug targets. *Nat. Rev. Drug Discov.* **7**, 489–503 (2008).

48. Hori, T. *et al.* Mucopolysaccharidosis type IVA: common double deletion in the N-acetylgalactosamine-6-sulfatase gene (GALNS). *Genomics* **26**, 535–542 (1995).

49. Lev, S. *et al.* Protein tyrosine kinase PYK2 involved in Ca2+-induced regulation of ion channel and MAP kinase functions. *Nature* **376**, 737–745 (1995).

50. Thorn, A. *et al.* Structure and activity of the only human RNase T2. *Nucleic Acids Res.* **40**, 8733–8742 (2012).

51. O'Brien, J. S. & Kishimoto, Y. Saposin proteins: structure, function, and role in human lysosomal storage disorders. *FASEB J.* **5**, 301–308 (1991).

52. van de Rijn, M., Heimfeld, S., Spangrude, G. J. & Weissman, I. L. Mouse hematopoietic stem-cell antigen Sca-1 is a member of the Ly-6 antigen family. *Proc. Natl. Acad. Sci.* **86**, 4634–4638 (1989).

53. McCord, J. M. & Fridovich, I. Superoxide Dismutase. *J. Biol. Chem.* **244**, 6049–6055 (1969).

54. Roberts, S. A., Davidson, A. J., McLean, L., Beynon, R. J. & Hurst, J. L. Pheromonal Induction of Spatial Learning in Mice. *Science (80-.).* **338**, 1462–1465 (2012).

55. Hurst, J. L. Scent marking and social communication. in *Animal Communication Networks* 219–244 (Cambridge University Press, 2005). doi:10.1017/CBO9780511610363.014.

56. Mucignat, C., Redaelli, M. & Caretta, A. One nose one brain: contribution of the main and accessory olfactory system to chemosensation. *Front. Neuroanat.* **6**, 1–9 (2012).

57. Demir, E. *et al.* The pheromone darcin drives a circuit for innate and reinforced behaviours. *Nature* **578**, 137–141 (2020).

58. Kahan, A. & Ben-Shaul, Y. Extracting Behaviorally Relevant Traits from Natural Stimuli: Benefits of Combinatorial Representations at the Accessory Olfactory Bulb. *PLOS Comput. Biol.* **12**, e1004798 (2016).

59. Ben-Shaul, Y. Extracting Social Information from Chemosensory Cues: Consideration of Several Scenarios and Their Functional Implications. *Front. Neurosci.* **9**, 1–15 (2015).

60. Beynon, R. J. *et al.* Polymorphism in major urinary proteins: Molecular heterogeneity in a wild mouse population. *J. Chem. Ecol.* **28**, 1429–1446 (2002).

61. Moudra, A. *et al.* Phenotypic and Clonal Stability of Antigen-Inexperienced Memory-like T Cells across the Genetic Background, Hygienic Status, and Aging. *J. Immunol.* ji2001028 (2021) doi:10.4049/jimmunol.2001028.

62. Isogai, Y. *et al.* Molecular organization of vomeronasal chemoreception. *Nature* **478**, 241–5 (2011).

63. Uchida, N., Poo, C. & Haddad, R. Coding and Transformations in the Olfactory System. *Annu. Rev. Neurosci.* 363–385 (2013) doi:10.1146/annurev-neuro-071013-013941.

64. Hurst, J. L., Robertson, D. H. L., Tolladay, U. & Beynon, R. J. Proteins in urine scent marks of male house mice extend the longevity of olfactory signals. *Anim. Behav.* **55**, 1289–1297 (1998).

65. Miller, C. H., Campbell, P. & Sheehan, M. J. Distinct evolutionary trajectories of V1R clades across mouse species. *BMC Evol. Biol.* **20**, 99 (2020).

66. Park, S. H., Podlaha, O., Grus, W. E. & Zhang, J. The microevolution of V1r vomeronasal receptor

    genes in mice. *Genome Biol. Evol.* **3**, 401–12 (2011).

67. Lilue, J. *et al.* Sixteen diverse laboratory mouse reference genomes define strain-specific haplotypes and novel functional loci. *Nat. Genet.* **50**, 1574–1583 (2018).

68. Dey, S. *et al.* Cyclic Regulation of Sensory Perception by a Female Hormone Alters Behavior. *Cell* **161**, 1334–1344 (2015).

Supplementary Figures

Supplementary Figure S1: (*a*) Area under curve response reliability index histograms of all generalist VSNs (gray; n = 10,258) and all specialist VSNs (red; n = 6,457) recorded from in this study. Note that for both response types, indices are normally distributed. While histograms peak around zero, distributions are much broader as compared to amplitude-based reliability index histograms (*Fig. 1d*). (*b* & *c*) Quantification of VSN responses recorded from male C57BL/6 mice challenged with pooled urine stimuli from two different male C57BL/6 mouse cohorts (*b*) or two different female C57BL/6 mouse cohorts (*c*). Pie charts (top) illustrate the proportions of generalist (gray; 33 % (*b*) and 16 % (*c*), respectively) and specialist neurons (≤1%; dark *versus* light blue (*b*) and dark *versus* light red (*c*), respectively) among all K⁺-sensitive VSNs (n = 1867 (*b*) and n = 1946 (*c*), respectively). Box-and-whisker plots (bottom, left) break down the urine-sensitive neurons by categories and illustrate the generalists-to-specialists ratios over individual experiments (n = 21). Boxes represent the first-to-third quartiles. Whiskers represent the 10th and 90th percentiles, respectively. Outliers (1.5 IQR; red x) are plotted individually. The central red band represents the population median ($P_{0.5}$). Response index histograms (bottom, left) depict signal amplitude strength distributions from generalist VSNs that responded to both paired stimuli (i.e., cohort / group 1 (g1) and cohort / group 2 (g2)). Multi-peak fitting with Gaussian curves (dashed lines) resulted either in a narrow single Gaussian that centers close to zero (*b*; peak = -0.06, σ = 0.16) or two additive Gaussian curves that best represented the calculated index distribution (*c*; peak 1 = -0.01, peak 2 = 0.25). Asterisk (*) indicates statistical significance, *p* < 0.05; Wilcoxon singed ranked test. (*d*) Concentration index histograms that outline individual compound concentration (im)balances of VOCs (yellow; top) and proteins (red; bottom) that are found in both male and female urine samples of either BALB/c (left) or wild (right) mice. Histograms are heterogeneous and data are not normally distributed.

Supplementary Figure S2: (*a*&*b*) After training Random Forest classifiers [40], lists of "Gini importance" [40] scores rank urine VOCs (*a*; yellow) and proteins (*b*; red) that are likely most informative to decode sex and strain, respectively. (*c* - *f*) Box-and-whisker plots quantify concentrations of the respective five VOCs (*c*&*e*) and proteins (*d*&*f*) that are ranked most important / informative (*a*&*b*). Data are plotted for each of the six sex/strain combinations (10 individual samples each). Red (female) and blue (male) boxes represent sex; strain as indicated. Boxes represent the first-to-third quartiles. Whiskers represent the 10th and 90th percentiles, respectively. Outliers (1.5 IQR; black dots) are plotted individually. Central bands represent the population median ($P_{0.5}$).

Supplementary Figure S3: (*a*) Box-and-whisker plots illustrating mean protein and VOC content (left) as well as individual compound concentration variance (right) for the six sex/strain combinations (10 individual samples each). Asterisks (∗) indicate statistical significance, *p* < 0.05, Kruskal-Wallis test. (*b*&*c*) Box-and-whisker plots quantify concentrations of previously reported VOC (*b*) and Mup (*c*) semiochemicals, respectively. Data are plotted for each of the six sex/strain combinations (10 individual samples each). Red (female) and blue (male) boxes represent sex; strain as indicated. Boxes depict the first-to-third quartiles. Whiskers represent the 10th and 90th percentiles, respectively. Outliers (1.5 IQR; black dots) are plotted individually. Central bands represent the population median (P$_{0.5}$).

Supplementary Figure S4: Hierarchical clustering of mouse urine lipocalin content reveals an unexpectedly diverse repertoire of 27 lipocalins. The horizontal tree clusters lipocalins according to proteomic homology based on similarities in protein abundances. The vertical tree clusters the 60 individual urine samples (10 samples for each of the 6 sex/strain combinations) according to patterns of lipocalin abundance (pseudocolors).

Supplementary Figure S5: (***a***&***b***) Comparison of male BALB/c generalist VSN response preferences, upon exposure to paired male (***a***) and female (***b***) urine stimuli from different strains, with strain-dependent concentration (im)balances among VOCs and proteins, respectively. Response index histograms (top rows) depict distributions of generalist data outlined in Figure 5 (gray bars). With one exception (***a*** (middle), male C57BL/6 *versus* male wild), histograms are not well fitted by single Gaussian curves. Rather, histogram shape is best represented by additive multi-peak fitting (dashed lines) with two (***a*** (right), ***b*** (left)), three (***a*** (left), ***b*** (right)), or even four (***b*** (middle)) Gaussians. Individual Gaussian curves center at -0.37, 0.07, and 0.3 (***a***; left), -0.06 (***a***; middle), -0.39 and -0.02 (***a***; right), -0.32 and 0.04 (***b***; left), -0.37, -0.18, 0.01, and 0.37 (***b***; middle), -0.38, -0.1, and 0.11 (***b***; right), respectively. For comparison, transparent and overlapping concentration index histograms (bottom rows) for VOCs (yellow) and proteins (red) are also shown (data correspond to results shown in Figure 4). (***c***&***d***) Comparative overlay of generalist VSN response index histograms (top rows) and corresponding fits (bottom rows) calculated for recordings from male BALB/c (white transparent bars) *versus* male C57BL/6 (black bars) animals that were stimulated with male (***c***) and female (***d***) urine, respectively. Vertical lines represent mean values. Asterisks (∗) indicate statistical significance between BALB/c and C57BL/6 response index distributions, *p* < 0.05, two-sample Kolmogorov-Smirnov test.