The rules of sex and strain: deciphering the chemical language of inbred and wild mouse conspecific scents

Maximilian Nagel[1,6], Marco Niestroj[1], Rohini Bansal[2], David Fleck[1], Angelika Lampert[3,4], Romana Stopkova[5], Pavel Stopka[5,*], Yoram Ben-Shaul[2,*], Marc Spehr[1,4,*,‡]

[1] Department of Chemosensation, Institute for Biology II, RWTH Aachen University, 52074 Aachen, Germany

[2] Department of Medical Neurobiology, Institute for Medical Research Israel Canada, Faculty of Medicine, The Hebrew University of Jerusalem, Jerusalem, Israel

[3] Institute of Neurophysiology, Uniklinik RWTH Aachen University, 52074 Aachen, Germany

[4] Research Training Group 2416 *MultiSenses – MultiScales*, RWTH Aachen University, 52074 Aachen, Germany

[5] BIOCEV group, Department of Zoology, Faculty of Science, Charles University, Prague, Czech Republic

[6] present address: Sensory Cells and Circuits Section, National Center for Complementary and Integrative Health, Bethesda, MD, 20892, USA

[*] correspondence should be addressed to: Pavel Stopka (pavel.stopka@natur.cuni.cz), Yoram Ben-Shaul (yoramb@ekmd.huji.ac.il), Marc Spehr (m.spehr@sensorik.rwth-aachen.de)

[‡] lead contact

1 Summary

In most mammals, conspecific chemosensory communication relies on semiochemical release
within complex bodily secretions and subsequent stimulus detection by the vomeronasal organ
(VNO). Urine, a rich source of ethologically relevant chemosignals, conveys detailed information
about sex, social hierarchy, health and reproductive state, which becomes accessible to a
conspecific via vomeronasal sampling. So far, however, numerous aspects of social
chemosignaling along the vomeronasal pathway remain unclear. Moreover, since virtually all
research on vomeronasal physiology is based on secretions derived from inbred laboratory mice,
it remains uncertain whether such stimuli provide a true representation of potentially more relevant
cues found in the wild. Here, we combine a robust low-noise VNO activity assay with comparative
molecular profiling of sex- and strain-specific mouse urine samples from two inbred laboratory
strains as well as from wild mice. With comprehensive molecular portraits of these secretions,
VNO activity analysis now enables us to (*i*) assess whether and, if so, how much sex- / strain-
selective "raw" chemical information in urine is accessible via vomeronasal sampling; (*ii*) identify
which chemicals exhibit sufficient discriminatory power to signal an animal's sex, strain, or both;
(*iii*) determine the extent to which wild mouse secretions are unique; and (*iv*) analyze whether
vomeronasal response profiles differ between strains. We report both sex- and, in particular,

strain-selective VNO representations of chemical information. Within the urinary 'secretome', both
volatile compounds and proteins exhibit sufficient discriminative power to provide sex- and strain-
specific molecular fingerprints. While total protein amount is substantially enriched in male urine,
females secrete a larger variety at overall comparatively low concentrations. Surprisingly, the
molecular spectrum of wild mouse urine does not dramatically exceed that of inbred strains.
Finally, vomeronasal response profiles differ between C57BL/6 and BALB/c animals, with
particularly disparate representations of female semiochemicals.

Introduction

In rodents and most other mammals, the accessory olfactory system controls conspecific chemical
communication during social interactions [1–4]. Ethologically meaningful chemostimuli that trigger
stereotypic social and sexual behaviors are predominantly detected by the vomeronasal organ
(VNO), a tubular sensory structure at the anterior base of the nasal septum. In a crescent-shaped
medial neuroepithelium, the VNO harbors approximately 100,000 to 200,000 vomeronasal
sensory neurons (VSNs [5]), each extending a single unbranched apical dendrite that terminates in
a paddle-shaped swelling [4]. At the dendritic tips, microvilli are immersed in mucus that fills a central
luminal canal, which extends via the vomeronasal duct into the nasal cavity. During social
investigatory behavior, which in mice primarily involves periods of intense licking and sniffing of
both facial and anogenital regions [6], semiochemicals are sucked into the VNO lumen. Upon
binding to cognate vomeronasal receptors (VRs), the chemical signal is transduced into electrical
VSN activity and, ultimately, neuronal discharge.

Behaviourally relevant natural chemostimuli are, typically, complex blends of compounds [7] in
various bodily secretions [8]. By far, the most widely studied secretion in animal chemosensory
research is urine (see [4] and references therein), which is a rich source of semiochemicals that
serves a well-established function in social communication. While we still lack a comprehensive
molecular description of this broadband vomeronasal stimulus, previous work has identified
several putative semiochemicals in mouse urine and other bodily secretions, which activate VSNs
and cover many structural groups and feature dimensions [7,9,18–21,10–17]. Prominent molecularly
identified VSN stimuli include various sulfated steroids [9,21–26], which could reflect the dynamic
endocrine state of an individual.

So far, virtually all research on vomeronasal physiology is based on urinary stimuli derived from
inbred laboratory mice (but see [27]). This facilitates standardization across studies but, while VSN

sex- and strain-selectivity has previously been explored [21,28], it remains unclear whether secretions
collected from inbred mice accurately represent the potentially more ethologically relevant stimuli
found in the wild. In fact, while wild *Mus musculus* populations exhibit several-fold higher levels of
genetic variation than those in human populations, commonly studied inbred strains of laboratory
mice derive from a limited set of founders. Therefore, such strains contain only a small subset of
the genetic variation that is present in nature [29]. Classical inbred strains of house mice are genetic
mosaics of the three main wild subspecies, *M. m. domesticus*, *musculus*, and *castaneus*, which
started to diverge ~350–500 thousand years ago. However, the genomes of prominent inbred
strains, such as C57BL/6 or BALB/c, are predominantly derived from *M. m. domesticus* [30]. The
mitochondrial genomes of both laboratory strains are identical, implying a common descent along
the maternal line. Given their overlapping geographical distribution, wild house mice subspecies
have also undergone secondary contact and hybridization [31], which has diversified their genetic
landscape.

Laboratory inbreeding, which has been ongoing for more than 100 years for C57BL/6 and BALB/c
strains [29], may have affected the chemical composition of bodily secretions and, consequently,
their information content. Indeed, both within and between strains, laboratory mice lack variation
in major urinary protein (MUP) patterns typically found among wild mice [32]. This inbred
homogeneity could have important implications for research investigating, for example, social
recognition or mate choice: if there are marked qualitative and/or quantitative differences in
chemical composition of secretions and, accordingly, in the neuronal representations of wild as
compared to inbred derived stimuli, one might question the conclusions based on a large body of
work using inbred secretions [27].

On the recipients' end, the level of chemosensory information available from con-/heterospecific
secretions is determined by individual VR expression profiles. Across strains, VR repertoires vary
as a function of both genetics and experience [33–36]. Therefore, it is unclear whether receptor arrays
in laboratory strains, despite hundreds of generations of inbreeding and domestication
("microevolution") in a laboratory environment, have retained selectivity for more ethologically
relevant wild-derived stimuli. Moreover, it remains uncertain whether VR tuning profiles enable
VSNs to capture key ethological features from molecular concentration differences between sex-
and strain-specific secretions [15,37]. Together, four key biological questions arise from these lines
of reasoning: (*i*) Do VSN response profiles reflect the global molecular content of urine (i.e., are
these neurons sensitive to many/most compounds) or, by contrast, is the VNO a highly selective
molecular detector (responding to just a few select molecules)? (*ii*) Are VSN response profiles
strain-specific? (*iii*) Which semiochemicals provide information about sex and/or strain? (*iv*) Is
there something unique about wild mouse secretions and/or the VSN response profiles they
trigger?

Here, to address these unresolved issues, we combine a robust VSN activity assay with
comparative molecular profiling of sex- and strain-specific mouse urine from two inbred laboratory
strains and wild mice. Our study provides molecular portraits of these secretions. We report that
large fractions of generic urine compounds are shared among both male and female mice of all
genetic backgrounds. We further show that the urinary 'secretome' in wild mice does not differ
dramatically from that found in laboratory strains. Surprisingly, while male urine contains much
higher protein concentrations, females secrete a larger variety of proteins (including MUPs). For
proteins common to all strains, concentrations are relatively low in C57BL/6, moderate in BALB/c,
and high in wild animals. However, despite this concentration bias, there is no overrepresentation
of wild-selective proteins. Notably, both the volatile organic compound (VOC) and protein profile
of urine, each provides sufficient information to decode a given sex/strain combination (with
protein content exhibiting stronger discriminative power). Moreover, we identify a rich lipocalin
repertoire in urine, which alone could allow chemosensory discrimination of sex/strain
combinations.

A key strength of this study is the use of the exact same stimuli as previously employed to
investigate sensory representations in the accessory olfactory bulb (AOB) [27], the first central
processing stage along the accessory olfactory system pathway. Our previous work demonstrated
that AOB representations of ethologically relevant urine stimuli are similar for male mice from two
different inbred strains (C57BL/6 and BALB/c), despite potential differences in VR repertoires. In
addition, we found that wild mouse stimuli elicit responses that, although not identical, are
nevertheless qualitatively similar to those from commonly used inbred strains [27]. VSN activity
analysis now enables us to ask whether these features also manifest in the VNO and, thus, to
assess (*i*) whether the information inherent in a sex- and strain-specific urinary secretome is
accessible via vomeronasal sampling; and (*ii*) if, on the population level, VSN sensory
representations differ between strains. Comparing responses from male C57BL/6 and BALB/c
mice (as done previously on AOB level [27]), our data demonstrate that recipient strain identity is
reflected by their VSN activity patterns. Moreover, when exposed to stimuli from different strains,
large VSN fractions (often >50 %) respond to only one stimulus, suggesting a substantial degree
of selective sampling. Together, our study reveals selective and strain-dependent representations
of urine chemical content, a surprising scarceness of selective responses to wild stimuli, as well
as remarkably rich and sex-/strain-specific molecular profiles that likely preserve most of the
biologically relevant information.

Results

In mice, the VNO is the predominant sensory structure mediating conspecific chemical
communication. In this study, by comparing samples derived from two inbred strains (BALB/c and
C57BL/6) as well as from wild mice, we pursue four main questions: (***i***) Which chemical
components in mouse urine – a major source of semiochemicals – distinguish sex and/or strain?
(***ii***) To what degree has inbreeding affected the chemical composition of urine (i.e., how unique
are wild mouse secretions)? (***iii***) How much of the sex- and strain-specific chemical information in
urine is accessible to a conspecific via vomeronasal sampling (i.e., how selective are VSN
response profiles)? (***iv***) Upon exposure to the same stimuli, do VSN response patterns differ
between strains?

A "low noise" assay to capture VSN population activity

To record VSN signal fingerprints in response to naturalistic stimuli, it is essential to establish a
robust population activity assay that reliably captures the raw information content inherent in bodily
secretions. To this end, we analyzed single-cell Ca^{2+} transients among large VSN populations in
acute coronal VNO sections (Figure 1a,b). Our experimental design followed a standard pairwise
comparison paradigm (Figure 1c). Pairs of pooled stimuli differed either in the donors' sex (male
versus female) or in their genetic background (BALB/c, C57BL/6, or wild). We repeated brief (10
s) alternating stimulus presentations twice at 180 s inter-stimulus intervals that ensured recovery
from VSN adaptation [38]. Experiments concluded with a brief exposure to elevated extracellular K^+
(**S₃**) to depolarize neurons and test for integrity of each neuron's spike generation machinery.
Dependent on their individual urine response profiles, VSNs were categorized as either specialists
(selective response to one stimulus) or generalists (responsive to both stimuli). Overall, we
recorded >43,000 K^+-sensitive neurons, of which a total of 16,715 VSNs (38.4%) responded to
urine stimulation. Of these urine-sensitive neurons, 61.4% displayed generalist profiles, whereas
38.6% were categorized as specialists (Figure 1c,d). As a measure of intrinsic signal variability,
we calculated reliability indices that quantify the similarity (or lack thereof) of two successive
responses to the same stimulus (with small values reflecting high reliability; see Materials and
Methods). We analyzed both Ca^{2+} signal amplitudes (Figure 1d) and integrals (Figure 1 – figure
supplement 1a) as measures of response magnitude. For both generalist and specialist VSNs,
response reliability indices are normally distributed around zero. While the distribution of reliability
indices derived from response integral measurements is somewhat broad (Figure 1 – figure
supplement 1a), indices based on signal amplitudes proved relatively homogeneous (Figure 1d).
Accordingly, we use average response amplitudes as indicators of signal strength throughout this
study.

In a first set of control experiments, we asked how VSNs from male C57BL/6 mice respond when
challenged with urine samples from two groups of animals of the same sex and inbred strain
(Figure 1 – figure supplement 1b,c). Since the chemical composition of both stimuli should be
similar, we expected that the vast majority of urine-sensitive VSNs will display generalist response
profiles. That was indeed the case. Less than 2% of VSNs showed stimulus selectivity (Figure 1
– figure supplement 1b,c). Moreover, response response indices (reflecting a bias toward either
of the paired stimuli) are normally (and narrowly) distributed around zero (Figure 1 – figure
supplement 1b,c). Both observations confirm a low level of biological and experimental noise in
this setting. Thus, our assay is well-suited to detect and compare VSN sensory responses upon
pairwise stimulation.

Figure 1: **A population activity assay captures sex-dependent stimuli representations.** (***a***) Anatomical location of the rodent VNO. Schematic depicting a sagittal section through a mouse head with an overlay of a VNO image. The vomeronasal duct that opens into the anterior nasal cavity is also highlighted. (***b***) Overview (left) and zoomed-in (middle) differential interference contrast micrograph showing an acute coronal VNO section from an adult C57BL/6 mouse. Confocal fluorescence image (right; dashed rectangle in middle) depicting a depolarization (elevated K$^+$) dependent cytosolic Ca^{2+} increase in VSN somata after bulk loading with Cal-520 AM. (***c***) Original traces showing changes in cytosolic Ca^{2+} concentration over time in two representative VSN somata. VNO slices were briefly challenged with two

mixtures of diluted mouse urine (1:100; 10s; top yellow bars/droplets). Repeated stimulation in alternating sequence at
180 s inter-stimulus intervals [38] was followed by membrane depolarization upon exposure to elevated extracellular K$^+$
(50 mM; 10 s). According to individual response type, VSNs were categorized as 'generalists' (top) or 'specialists'
(bottom). Signal amplitudes in response to the same stimulus allowed calculation of a reliability index (RI) as a measure
of signal robustness. (***d***) Amplitude reliability index histograms of all generalist VSNs (gray; n = 10,258) and all specialist
VSNs (red; n = 6,457) recorded in this study. Note that for both response types, indices are normally distributed with a
narrow central peak around zero. (***e***) Quantification of results obtained from recordings in VSNs from male C57BL/6
mice challenged with male *versus* female C57BL/6 urine stimuli. Pie chart (top) illustrates the proportions of generalist
(19%) and specialist neurons (1% and 4%, respectively) among all K$^+$-sensitive VSNs (n = 1,999). Bar graph (middle,

left) breaks down the summed total of urine-sensitive neurons by categories and compares their distribution with the
proportions of VOCs (yellow background; n = 405) and proteins (red background; n = 601) found either in both male
and female urine (grey bars) or exclusively in samples from one sex (male, dark blue; female, purple). Box-and-whisker
plots (bottom, left) illustrating generalists-to-specialists ratios over individual experiments (n = 19). Boxes represent the
first-to-third quartiles. Whiskers represent the 10th and 90th percentiles, respectively. Outliers (1.5 IQR; red x) are plotted
individually. The central red band represents the population median ($P_{0.5}$). Results are shown in relation to box-and-
whisker plots that outline chemical content data obtained from paired comparisons (n = 10 individuals per group). (***f***)
Molecular composition of male *versus* female urine from BALB/c and wild mice. Bar graphs (top) display proportions of
VOCs (yellow background; BALB/c, n = 514; wild, n = 462) and proteins (red background; BALB/c, n = 407; wild, n =
526) found either in both male and female urine (grey bars) or exclusively in samples from one sex (male, dark blue;
female, purple). Box-and-whisker plots (bottom) quantify category ratios for individual paired experiments (n = 10
individuals per group). (***g***) Quantification (Bradford assay) of protein/peptide content in urine samples from C57BL/6,
BALB/c, and wild mice, respectively (n = 10 each). Note the substantially increased protein content in male samples.
(***h***) Response index histogram (top) obtained from generalist C57BL/6 VSNs that responded to both male and female
same-strain urine (data corresponding to (***e***)). Fitted Gaussian curve (dashed line) centers close to zero (peak = -0.03)
and shows a relatively narrow width (σ = 0.13). By contrast, concentration index histograms (bottom), calculated for
VOCs (yellow) and proteins (red) found in both male and female urine samples, are heterogeneous and not normally
distributed. Asterisks (*) indicate statistical significance, $p < 0.05$; Wilcoxon signed-rank test (for VSN functional data),
Mann-Whitney U test (for molecular profiling), and unpaired t-test (for protein content comparison shown in (***g***)).
20

Sex-specific stimuli elicit distinct VSN sensory representations

We next investigated how sex differences are reflected in VSN response profiles. When

challenged with male *versus* female urine from C57BL/6 animals, the fraction of specialist neurons

more than doubled to 5% of all K^+-sensitive neurons. Notably, female urine recruited more

specialist neurons than male urine (Figure 1e). We then asked whether this pattern is correlated

with the chemical compositions of male and female urine. In-depth molecular analysis of urine

content via GCxGC-MS as well as nLC-MS/MS identified a total of 1006 molecules (detected in

≥3 out of 10 male or female samples, respectively), of which approximately 40% are low molecular

weight VOCs, while 60% are proteins. Roughly half of the molecules in either group are found in

both male and female urine (Figure 1e). Unexpectedly, while we hardly identify any male-specific

proteins in C57BL/6 urine, we find a large fraction of female-specific proteins. We asked whether

this phenomenon is (*i*) a distinct feature of C57BL/6 mice, (*ii*) common among inbred laboratory

strains, or (*iii*) also observed in wild animals. Therefore, we included urine samples from both

BALB/c and wild mice in extended molecular profiling. In both groups, we observed similarly

increased levels of female-specific proteins (Figure 1f). The total amount of protein, however, is

substantially enriched in male urine (Figure 1g). Our data thus suggest that, while females secrete

a larger variety of proteins, overall concentrations are comparatively low.

Finally, we asked whether common compounds (i.e., molecules identified in both male and female

urine) show concentration disparities between sexes and, if so, how such differences are reflected

in VSN response profiles. When VSNs were challenged with male *versus* female C57BL/6 urine,

19% of all K^+-sensitive neurons responded to both stimuli (Figure 1e). Here, again, response

preference indices are narrowly distributed and center around zero (Figure 1h). By contrast,

analogously calculated concentration indices (see Materials and Methods) that can reflect

potential disparities are distributed more broadly and non-normally (Figure 1h). This apparent

incongruence between nonpreferential generalist sensitivity and relatively large chemical

concentration differences indicates that VSN responses (at least in C57BL/6 males) do not simply

reflect the information theoretically available via concentration differences between male and

female same-strain stimuli. One explanation for this is that common ligands are not well-

represented by the observed general concentration disparities. Alternatively, it may be that even

low concentrations fully activate a given VSN, and thus concentration differences are not reflected

by response strengths. Notably, a broad and non-normal distribution of concentration indices

among molecular components of male *versus* female urine is also found in samples from both

BALB/c and wild mice (Figure 1 – figure supplement 1d).

Chemical characterization of urine content identifies distinctive molecular signatures of

sex and strain

Which differences in chemical composition (i.e., which molecules) characterize sex-/strain-specific

secretions and may thus provide information about sex and strain to a recipient? Molecular

profiling of urine content provides a unique opportunity to address this question and identify

enriched compounds (i.e., present in ≥6 of 10 individual samples) with the potential capacity to

signal an animal's sex, strain, or both. Adopting this more conservative criterion, we detected 208

enriched VOCs and 264 enriched proteins. Almost half of all VOCs (47.6%) and about one-third

of all proteins (31.4%) are found in all six sex/strain combinations tested and are, thus, considered

generic mouse urine components (Figure 2a,b). Moreover, a total of 33 compounds (10 VOCs, 23

proteins) were identified across all strains, but in a sex-specific fashion. Similarly, 22 molecules

(10 VOCs, 12 proteins) revealed strain selectivity, independent of sex. Notably, a large fraction of

compounds, i.e., 34.1% of all VOCs (71 / 208) and 22.3% of proteins (59 / 264), were detected

exclusively in one of the six sex/strain combinations. Only 18 of the 208 VOCs (8.7%) could not

be categorized as either generic or specific (for either sex, or strain, or a unique sex/strain

combination). Among the 82 proteins that showed sex specificity, either across strains (23

proteins) or as part of a unique sex/strain combination (59 proteins), the vast majority (89.0%) was

found in female samples. For VOCs, however, the opposite picture emerged. Here, 52 of 81 sex-

specific compounds (64.2%) were selectively detected in male samples. Together, chemical

profiling revealed (*i*) that large fractions of urine content are shared among both laboratory and

wild mice; (*ii*) that roughly one-third of urinary VOCs and one-fourth of proteins are exclusively

found in a given sex/strain combination; and (*iii*) that male and female mice might have adopted

different chemical secretion strategies to signal their sex.

Figure 2: **Chemical profiling of urine content identifies unique sex- and strain-specific molecular fingerprints.** (*a*&*b*) Matrix layout for all intersections of VOCs (*a*) and proteins (*b*) among the six sex/strain combinations, sorted by size. Colored circles in the matrix indicate combinations that are part of the intersection. Bars above the matrix columns represent the number of compounds in each intersection. Empty intersections have been removed to save space. Horizontal bar charts (bottom, left) depict the number of VOCs (*a*) and proteins (*b*) detected in each urine set. Proteomics data are available via ProteomeXchange with identifier PXD042324. (*c-h*) Sparse Partial Least-Squares Discriminant Analysis (sPLS-DA) score plots depicting the first two sPLS-DA components, which explain 7-10% (1st component) and 5-8% (2nd component) of VOC data variance (*c-e*) as well as

21-24% (1st component) and 11-12% (2nd component) of protein data variance (*f-h*), respectively. Ellipses represent 95% confidence intervals. Plots demonstrate sample clustering according to the urine donors' sex (*c*&*f*), genetic background (*d*&*g*), or sex/strain combination (*e*&*h*). Each data point represents a sample from an individual animal (n = 60; 10 samples per sex/strain combination), with sample type colored according to symbol legend (bottom).

If information coding along the accessory olfactory pathway would strictly follow a 'labeled-line' logic [39], absence or presence of a given molecule could be adequate to signal sex or strain. Several recent lines of evidence, however, suggest a combinatorial coding strategy that also involves some level of circuit plasticity [4,34,40]. Therefore, we next asked if the total VOC or protein content of a given urine sample preserves sufficient predictive / discriminative information to classify samples according to sex, strain, or a specific sex/strain combination. We used sparse Partial Least-Squares Discriminant Analysis (sPLS-DA) [41], a chemometrics machine learning technique, to reduce data dimensionality and optimize sample separation [42]. When plotted on two-dimensional coordinates that represent the most discriminative variables (Figure 2c-h), both VOCs (Table 1) and proteins provide sufficient information to cluster stimuli according to sex (Figure 2c,f), strain (Figure 2d,g), or a combination of both variables (Figure 2e,h). We then calculated

Variable Importance in Projection (VIP) scores as measures of a particular variable's informative
power, which is correlated to the variance explained by the model [41,42]. Generally, we find that
protein content exhibits stronger discriminative power (21-24% & 11-12% of explained variance)
than VOC content (7-10% & 5-8% of explained variance).

We next aimed to identify the most relevant variables (i.e., molecules) for sex or strain
classification. Training Random Forest classifiers [43], we obtained feature significance scores, the
"Gini importance" [43], that provide relative relevance rankings of the individual variables.
Supplementary Figure 2 lists the 20 most informative VOCs (Table 2) and proteins that
discriminate sex and strain, respectively (Figure 2 – figure supplement 1a,b), along with the
abundance of the corresponding top 5 molecules (VOCs or proteins) for each of the six sex/strain
combinations (Figure 2 – figure supplement 1c-f). Notably, major urinary protein 20 (Mup20 /
darcin) [44], fatty acid-binding protein 5 (Fabp5) [45], and N-acetylgalactosamine-6-sulfatase (Galns)
[46] exhibit substantial power to discriminate sex across strains (Figure 2 – figure supplement 1b,f).
While Mup20 and Galns are considerably more abundant in male urine, Fabp5 appears to be
specific for female samples. Strain discrimination, on the other hand, is optimal with protein-
tyrosine kinase 2-beta (Ptk2b) [47], RNase T2 (Rnaset2) [48], prosaposin (Psap) [49], lymphocyte
antigen 6A-2/6E-1 (Ly6a) [50], and superoxide dismutase 1 (Sod1) [51] (Figure 2 – figure supplement
1b,d). Specifically, Ptk2b is detected exclusively in C57BL/6 mice. Ly6a is absent in BALB/c
animals, whereas Rnaset2 is largely missing in both C57BL/6 and wild mice. With proteins
exhibiting stronger discriminative power than VOCs (Figure 2c-h), future studies will have to focus
on these proteins to identify potential functions as vomeronasal chemosignals.

Overall, we have detected similar amounts of proteins / VOCs across strains. However, individual
variability in protein and VOC content was significantly higher in wild mice than in both laboratory
strains (Figure 2 – figure supplement 2a). We confirmed secretion of previously reported putative
semiochemicals [10,11,15,40,52], including both VOCs (Figure 2 – figure supplement 2b) and proteins
(Figure 2 – figure supplement 2c) in both wild and laboratory mice. Notably, however, all Mups
(including Mup20) and most such VOCs were found in samples from either sex, albeit at male-
biased concentrations for Mup3, Mup17, and Mup20 (Figure 2 – figure supplement 2c). The only
compounds showing male-specific secretion are 2-sec-butyl-4,5-dihydrothiazole (SBT) and
farnesenes (Figure 2 – figure supplement 2b), which have previously been implicated as
facilitators of female mouse puberty acceleration [11].

Notably, we find a rich repertoire of 27 lipocalins in mouse urine. When based exclusively on
lipocalin content, hierarchical clustering groups individual samples into a set of clusters that, with
very few exceptions (5 out of 60), correspond to the six sex/strain combinations (Figure 2 – figure
supplement 3). This finding, therefore, demonstrates the power of urinary lipocalins for potential
chemosensory discrimination of sex and/or strain.

VSNs are more selective for strain than for sex

Next, we challenged neurons with stimulus pairs from two same-sex / different-strain combinations

and asked whether VSN response profiles reflect the molecular fingerprint of corresponding urine

samples (e.g., male C57BL/6 *versus* male BALB/c (Figure 3a)). We analyzed a total of 17,416 K$^+$-

sensitive VSNs (Figure 3a-f). Again, we distinguished between specialist VSNs that responded

exclusively to one stimulus, and generalists, which responded to both stimuli. Along the same

lines, we categorized either strain-specific or broadly detected VOCs and proteins. Several

conclusions emerge from these classifications: (*i*) with one exception (i.e., upon stimulation with

male C57BL/6 *versus* wild stimuli (Figure 3b)), roughly half of all urine-sensitive VSNs are

generalists – a result consistent with our finding that generally ~50% of urine molecules are shared

among compared strains. Accordingly, the fraction of strain-selective (specialist) responses is

considerably larger than observed for sex-specific responses (Figure 1e); (*ii*) in female urine,

BALB/c-specific proteins are substantially underrepresented, a fact not reflected by VSN response

profiles (Figure 3d,f); (*iii*) surprisingly, the amount of strain-specific molecules in wild mouse urine

does not vastly exceed that in inbred strains; and (*iv*) accordingly, selective VSN responses to

wild stimuli are by no means more common (Figure 3b,c,e,f).

Figure 3: **Selective VSN response profiles upon exposure to strain-specific signatures.** Quantitative comparison between VSN responses to paired stimuli and their respective chemical signatures. Neurons of male C57BL/6 mice were challenged with male (*a-c*) or female (*d-f*) urine, respectively. Response profiles are compared upon stimulation with C57BL/6 *versus* BALB/c urine (*a*&*d*), C57BL/6 *versus* wild stimuli (*b*&*e*), and BALB/c *versus* wild urine (*c*&*f*), respectively. Pie charts (top) illustrate the proportions of generalist (light gray) and specialist neurons (dark gray, white, and purple, respectively) among all K$^+$-sensitive VSNs (*a*, n = 1855; *b*, n = 4116; *c*, n = 2462; *d*, n = 2376; *e*, n = 3230; *f*, n = 3377). Bar graphs (left) break down the urine-sensitive neurons by categories and compare their distribution with the proportions of VOCs

(middle; yellow background; *a*, n = 450; *b*, n = 448; *c*, n = 479; *d*, n = 480; *e*, n = 405; *f*, n = 492) and proteins (right;
red background; *a*, n = 317; *b*, n = 330; *c*, n = 334; *d*, n = 657; *e*, n = 715; *f*, n = 584) found either in both urine types

(gray bars) or exclusively in samples from one group (color code as in pie charts). Box-and-whisker plots (bottom)
illustrate generalist-to-specialist VSN category ratios over individual experiments (*a*, n = 6; *b*, n = 10; *c*, n = 7; *d*, n = 9;
e, n = 10; *f*, n = 12). Boxes represent the first-to-third quartiles. Whiskers represent the 10th and 90th percentiles,
respectively. Outliers (1.5 IQR; red x) are plotted individually. The central red band represents the population median
($P_{0.5}$). Results are shown in relation to box-and-whisker plots that outline chemical content data obtained from paired
comparisons (n = 10 individuals per group). Asterisks (∗) indicate statistical significance, *p* < 0.05, Wilcoxon signed-
rank test.

Pronounced strain-dependent concentration imbalances between common urinary
compounds are not reflected by generalist VSNs

When comparing generalist VSN responses to male *versus* female C57BL/6 urine (Figure 1e,h)

we noted that the narrow normally distributed stimulus response index histograms did not match

the broader and heterogeneous distributions of concentration disparities between sexes (Figure

1h; S1b-d). For sex-dependent cues, as mentioned above, this could indicate that semiochemical

concentration differences carry only limited information. Next, we therefore asked whether strain-

specific concentration differences between urine samples exist and, if so, whether such

differences can convey information about strain. In total, we recorded strain-independent

generalist responses from 3,366 VSNs in male C57BL/6 mice (Figure 4). In all but one

experimental condition (i.e., when comparing male stimuli from BALB/c *versus* wild mice)

response preference indices were normally distributed and well fit by relatively narrow Gaussian

curves that centered around zero. By contrast, chemical analysis revealed that many compounds

identified in urine from both strains differ substantially in concentration. For proteins, in particular,

strong concentration disparities exist in all strain combinations analyzed. In both male and female

samples, concentrations of common proteins are relatively low in C57BL/6, moderate in BALB/c,

and high in wild animals. As shown in Figures 2b and 3, this concentration bias towards wild

proteins does not translate into any dramatic overrepresentation of proteins selectively found in

wild male and/or female urine. In fact, the massively skewed distributions of protein concentration

indices are not reflected by generalist VSN profiles. The latter better match VOC concentration

distributions, which generally display broad, yet Gaussian shapes. We conclude that VSN

population response strength might not be so strongly affected by strain-dependent concentration

differences among common urinary proteins. In that case, it would appear somewhat unlikely that

individual VSN activity provides fine-tuned information about distinct semiochemical

concentrations. Alternatively, as some (or even many) of the identified proteins could not serve as

vomeronasal ligands at all, generalist VSNs might sample information from only a subset of

compounds which, in fact, are secreted at roughly similar concentrations.

Vomeronasal representation of female semiochemicals differs between two inbred strains

So far, our approach was restricted to VSN signals recorded from male C57BL/6 mice. An

important question, of course, is whether the response profiles we observed are themselves

recipient strain-dependent. Thus, we next aimed to assess the extent to which our findings

generalize to VSN populations from another laboratory animal strain. We therefore repeated all

six pairwise same-sex / different-strain stimulation experiments, using acute VNO slices from male

BALB/c mice (Figures 5 & S5). Categorization as generalist or specialist VSNs revealed that

proportions differed significantly between VSNs from BALB/c *versus* C57BL/6 mice, albeit at

varying degrees. Similar to C57BL/6 neurons, selectivity of BALB/c neurons to wild-derived stimuli

was rather rare. In fact, (*i*) compared to an average specialist rate of 11.2% ± 6.6% (mean ± SD)

calculated over all 13 binary stimulus pairs (n = 26 specialist types), we recorded hardly any

specialist responses upon stimulation with urine from wild females (2% and 3%, respectively), and

(*ii*) we found comparatively few generalist signals when wild-derived female urine was among the

paired stimuli (Figure 5e,f). This striking insensitivity is not observed in VSNs from C57BL/6 mice,

suggesting substantial differences in VR expression between the two inbred laboratory strains.

Notably, we can rule out that the observed differences result from differential sampling along the

VNO's anterior-to-posterior axis since, for each experiment, we routinely sample and average

across slices cut along the organ's entire anterior-to-posterior length (see Materials and Methods).

Figure 4: **Strain-dependent concentration imbalances exert relatively mild effects on VSN population response homogeneity.** Comparison of male C57BL/6 generalist VSN response preferences, upon stimulation with paired urine stimuli from different strains, with strain-dependent concentration imbalances among VOCs and proteins, respectively. (*a*&*b*) Response index histograms (top rows) depict distributions of generalist data outlined in Figure 3 (gray bars). With one exception (*a* (right), male BALB/c *versus* male wild), histograms are well fitted by single Gaussian curves (dashed lines) that each center relatively close to zero (*a* (left), peak = 0.08, σ = 0.18; *a* (middle), peak = -0.11, σ = 0.21; *a* (right), 1st peak = -0.12, 1st σ = 0.14; 2nd peak = 0.21, 2nd σ = 0.09; *b* (left), peak = 0.07, σ = 0.12; *b* (middle), peak = 0.03, σ = 0.23; *b* (right), peak = 0.05, σ = 0.18). Concentration index histograms (middle & bottom rows), calculated for VOCs (yellow) and proteins (red) found in both tested urine samples, are more heterogeneous. Notably, while most VOC concentration index histograms are also fitted by single, albeit broader Gaussian curves (*a* (left), peak = -0.07, σ = 0.37; *a* (middle), peak = 0.02, σ = 0.45; *a* (right), 1st peak = 0.07, 1st σ = 0.32; 2nd peak = 0.64, 2nd σ = 0.06; *b* (left), peak = 0.15, σ = 0.29; *b* (middle), peak = 0.21, σ = 0.35; *b* (right), peak = -0.01, σ = 0.42), protein concentration imbalances are not normally distributed.

Next, we examined whether generalist VSN response profiles differ between male BALB/c and
C57BL/6 animals (Figure 5 – figure supplement 1). In total, we recorded generalist responses from
1,741 BALB/c neurons. When plotting response preference indices, we noticed less
homogeneous distributions than previously observed in C57BL/6 mice. Five of the six histograms
showed multiple peaks and thus could not be fitted by a single Gaussian, whereas the only
histogram adhering to a normal distribution was comparatively broad (Figure 5 – figure supplement
1a). As observed in C57BL/6 neurons, the skewed distributions of protein concentration indices
were not reflected by BALB/c generalist VSN profiles. Comparison of generalist VSN response
histograms between BALB/c and C57BL/6 mice (Figure 5 – figure supplement 1c,d) revealed
strong and consistent differences upon stimulation with female stimuli (Figure 5 – figure
supplement 1d). While single Gaussians around zero characterize C57BL/6 generalist
distributions upon stimulation with female stimuli (Figure 5 – figure supplement 1d), several
prominent peaks emerged when fitting histograms derived from BALB/c VSNs. Notably, for some
generalist BALB/c neurons, wild-derived female stimuli are less potent than their inbred strain
counterparts. This finding either indicates reduced concentrations of the corresponding molecules
in wild urine (rendering most proteins unlikely candidates (Figure 5 – figure supplement 1a,b)), or
suggests some yet to be determined form of cooperativity upon receptor-ligand interaction.
Together, for both generalist and specialist VSNs, vomeronasal representation of female
semiochemicals differs considerably between the two inbred mouse strains.

In the present study, we have established a robust VSN activity assay that allows pairwise
comparison of neural selectivity and response strength upon stimulation with chemically defined
natural stimuli. In-depth chemical analysis of sex- and strain-specific individual urine samples
revealed that (*i*) large fractions of urine content are shared among mice of all sex / strain
combinations; that (*ii*) the amount of molecules selectively found in wild mouse urine does not
dramatically exceed the urinary secretome of inbred strains; that (*iii*) across strains, female-
specific proteins vastly outnumber the male-specific variety, while (*iv*) overall protein concentration
is substantially enriched in male urine; that (*v*) concentrations of common proteins are relatively
low in C57BL/6, moderate in BALB/c, and high in wild animals; that (*vi*) both secreted VOC and
protein profiles provide sufficient information to distinguish sex, strain, or both; and that (*vii*) the
rich urinary lipocalin repertoire alone might allow chemosensory discrimination of sex and/or
strain.

Figure 5: **Vomeronasal representation of female semiochemicals differs between inbred strains.** Comparison of VSN response profiles between male BALB/c and C57BL/6 mice. Pie charts (top) illustrate the proportions of generalist (light gray) and specialist BALB/c neurons (dark gray, white, and purple, respectively) among all K+-sensitive VSNs (*a*, n = 1244; *b*, n = 2063; *c*, n = 1903; *d*, n = 1770; *e*, n = 1934; *f*, n = 1818). VSNs were challenged with male (*a-c*) or female (*d-f*) urine, respectively. Response profiles are compared upon exposure to C57BL/6 *versus* BALB/c urine (*a&d*), C57BL/6 *versus* wild stimuli (*b&e*), and BALB/c *versus* wild urine (*c&f*), respectively. Bar graphs break down the urine-sensitive neurons by categories and compare distributions among BALB/c neurons (left) to responses recorded from C57BL/6 VSNs (right; gray background). Box-and-whisker plots (bottom) illustrate generalist-to-specialist VSN category ratios over individual experiments (*a*, n = 8; *b*, n = 10; *c*, n = 9; *d*, n = 10; *e*, n = 9; *f*, n = 10). Boxes represent the first-to-third quartiles. Whiskers represent the 10^{th} and 90^{th} percentiles, respectively. Outliers (1.5 IQR; red x) are plotted individually. The central red band represents the population median ($P_{0.5}$). Asterisks (∗) indicate statistical significance, $p < 0.05$, Wilcoxon signed-rank test.

When asking how much of this chemical information is accessible to inbred male mice via vomeronasal sampling we observe that (*i*) VSN population response profiles do not reflect the global molecular content of urine, suggesting that the VNO functions as a rather selective molecular detector; that (*ii*) selective VSN responses to wild stimuli are by no means more common (in fact, selectivity to wild-derived stimuli is rather rare); that (*iii*) VSN generalist signal strength is unlikely to encode semiochemical concentrations across the entire range of compounds; that (*iv*) male BALB/c neurons display striking insensitivity when challenged with urine from wild females; and, thus, that (*v*) vomeronasal representation of female semiochemicals differs considerably between inbred strains.

Discussion

Urine is the primary source of social chemosignals among mice (and, in fact, many other
mammals) and contains both 'fixed' (i.e., genomic) information about strain, sex, individual identity,
genetic histocompatibility and background, as well as 'variable' (i.e., metabolic) information on
current social, reproductive and health status [53]. The ability to glean ethologically meaningful
information from chemosensory sampling of urine (or any conspecific bodily secretion) depends
on (*i*) the specific (semio)chemical composition of a urine sample, and (*ii*) the sensory apparatus
used for sampling. For most mammals, the VNO is the key chemosensory structure involved in
detecting conspecific chemical cues [4]. The virtually universal use of inbred laboratory mice in
research aimed at understanding VNO physiology – as both the donors of stimuli and the
experimental subjects employed in these studies – could have resulted in misconceptions and
biased notions about vomeronasal signaling and, thus, conspecific chemical communication. If
that were the case, one might question the relevance and ethological validity of conclusions drawn
from a large body of work using inbred secretions [27]. Here, we addressed this issue from both a
chemical ecology and a physiological perspective. In-depth comparative molecular profiling of
urine from two classical laboratory strains as well as wild mice reveals several shared features,
but also qualitative and quantitative differences in composition. Furthermore, we observe
substantial differences in vomeronasal representations of stimuli between C57BL/6 and BALB/c
sensory neuron populations.

For analytical purposes, we separate the urinary 'volatilome' and proteome. In chemosensory
research, this distinction has often been conceptualized as general (i.e., airborne) odors, which
activate the main olfactory system, *versus* vomeronasal stimuli [54]. This notion, however, is
misleading since organic compounds with low molecular weight and high vapor pressure (i.e.,
VOCs) in bodily secretions do not instantly evaporate, of course. Rather, they are readily
accessible for vomeronasal sampling upon direct contact during investigatory behavior. Notably,
our chemical profiling approach omits (sulfated) steroids other non-volatile small organic
molecules, which have previously been identified in mouse urine as VSN stimuli [9]. Caution should
thus be exerted to not attempt to fully explain VSN response specificity based on VOC and protein
content alone.

For both urinary VOCs and proteins, large fractions are shared among both male and female mice
of either genetic background (Figure 2a & b). Such compounds could be considered generic
(mouse) urine components and might not even serve any chemosensory signaling functions.
Notably, both the urinary volatilome and proteome on their own, each entail sufficient information
to discern an individual's sex and strain, with protein content exhibiting stronger discriminative
power. The protein that is most informative for discriminating between sexes is, perhaps not

surprisingly, Mup20 (darcin) [44]. This well-described "maleness signal" had previously been

reported to elicit innate attraction and generate a conditioned place preference in females [52,55],

whereas, in males, Mup20 promotes aggression [40]. Chemosensory roles of the second and third

best protein determinants of sex discrimination are basically unexplored. Fabp5 [45] and Galns [46]

are substantially enriched in female and male urine, respectively. Fatty acid-binding proteins,

including Fabp5, are evolutionary conserved intracellular lipid chaperones that coordinate cellular

lipid trafficking and signaling and are thus linked to metabolic and inflammatory pathways [45]. N-

acetylgalactosamine-6-sulfatase (Galns) on the other hand is a lysosomal hydrolase. Five urinary

proteins – Ptk2b [47], Rnaset2 [48], Psap [49], Ly6a [50], and Sod1 [51] – display pronounced strain-

dependent differences in concentration. None of these has previously been attributed a

chemosensory function. Challenging the mouse VNO with purified recombinant protein(s) will help

elucidate whether such functions exist.

Proteomic profiling revealed three additional, rather unexpected findings: First, while male urine

contains much higher protein concentrations, females of a given strain secrete a larger variety of

proteins (including MUPs). In fact, we do not find a single protein that is exclusively detected in

males across strains. By contrast, 23 urinary proteins, while present in all three strains, are found

only in females. In line with these observations, only one additional Mup (i.e., Mup21) made the

list of the 20 most informative proteins that discriminate sex. In general, our data provide little

evidence for a sparse molecular code of (fe)maleness. Rather, the concept of 'signature mixtures'

[7,56,57], which emphasizes a combinatorial ratio code instead of mere presence/absence

phenomena, gains traction. Second, we identify a surprisingly rich lipocalin repertoire in urine,

which alone could allow chemosensory discrimination of sex/strain combinations. Within a total of

27 lipocalins, individual patterns allow hierarchical clustering into sex-/strain-specific groups

(Figure 2 – figure supplement 3). These data, thus, support the notion that the lipocalin 'code', if

relevant, is combinatorial. Third, regarding its molecular spectrum, the urinary 'secretome' in wild

mice does not differ dramatically from the repertoire found in laboratory strains. To paraphrase

this generally reassuring conclusion: while inbreeding could have dramatically modified the nature

of chemical secretions and, consequently, their perception by other mice, inbred chemostimuli are

largely representative of (potentially) ethologically more relevant wild chemosignals. Notably,

however, while mean VOC and protein concentrations show similar distributions across sex/strain

combinations, individual variability is strongly increased in wild mouse urine (Figure 2 – figure

supplement 2a). Accordingly, this finding confirms previous reports of increased individual

variation in wild mice [32,58]. Another inherent factor that could account for differences in urine

secretions among each of the groups, and particularly for comparisons between inbred and wild

stimuli, is the microbiome [27,59]. Yet, we stress that all urine donors were housed in the same facility

and fed with the same diet [27].

Some limitations of our study need to be acknowledged. Regarding stimuli, the six secretion sets
we used do not cover the entire coding capacity of the accessory olfactory system [26]. Moreover,
for VSN response profiling, stimulus samples were pooled across ten individuals in each of the six
sex/strain categories. While pooling stimuli reduces individual variability across samples (e.g.,
regarding fluctuating physiological states), relevant stimulus aspects could be masked. Thus,
given the increased chemical variability we observed among individual wild urine samples, pooling
might obscure distinctive molecular features of wild mouse secretions. The same holds true for
estrus cycle-dependent female stimuli. Because we did not monitor the estrus stage of female
urine donors, mixes likely contain samples across the entire cycle. This is relevant as VSN
responses may be affected by a donor's cycle stage [27,60,61]. Another limitation stems from the use
of male inbred mice as experimental subjects. The rationale behind this experimental strategy is
to allow for comparisons with our previous study on AOB response profiles [27], which used the
exact same settings (see below). Nonetheless, future efforts will have to reveal (*i*) whether our
main findings apply to female recipients; and (*ii*) if our observations generalize to other inbred,
outbred, or wild mouse strains. As we recently declared [27], the latter considerations underscore
the importance of recording from wild recipients. While this endeavor presents significant practical
challenges, it remains an important goal for future studies.

A surprising, but overall reassuring observation is that responses to wild and inbred stimuli are
qualitatively similar. Strikingly, and somewhat counterintuitively, selective VSN responses to wild
stimuli are rather rare. This does not result from a generally reduced compound content in wild
urine as molecular profiling revealed comparably rich chemical portfolios in wild and inbred
samples alike. Rather, we speculate that inbreeding over hundreds of generations in laboratory
settings [29] has resulted in "microevolutionary" pressure to maintain sensitivity to signals from
same- or similar-strain individuals. Indeed, compared to the C57BL/6 reference genome, genetic
variability within VR repertoires is massively increased among wild-derived mice (particularly of
the *M. m. musculus* subspecies) [36]. Among the approximately 200 orthologous receptor genes
compared, BALB/c genes display 184 non-synonymous and just one private (i.e., unique to a
given strain) single-nucleotide polymorphism (SNP). By contrast, VR genes of the wild-derived *M.*
m. musculus PWK/PhJ strain show 789 non-synonymous SNPs and 508 private SNPs [36].

Initially, we asked whether VSN response profiles reflect the global molecular content of urine or,
by contrast, if the VNO serves as a rather selective semiochemical detector. Our findings support
a high level of selectivity. When challenged with same-sex / different-strain stimuli, large VSN
fractions selectively respond to just one stimulus. Intriguingly, this fraction of strain-selective
specialists is considerably larger than observed for sex-specific responses. In line with the notion
of highly selective vomeronasal sampling is our observation that the concentration differences

between compounds shared among strains, which are often substantial, are not reflected by
similarly pronounced differences in response strength among generalist VSNs. There are several,
not necessarily mutually exclusive explanations for this finding: First, concentration could simply
not be a read-out parameter for VSNs, which would support previous ideas of concentration-
invariant VSN activity [15]. Second, the concentrations in freshly released urine could just exceed
the dynamic tuning range of VSNs since, particularly for VOCs, natural signals (e.g., in scent
marks) must be accessible to a recipient for a prolonged amount of time (sometimes days). A
similar rationale could explain the increased protein concentrations in male urine, since male mice
use scent marking to establish and maintain their territories and urinary lipocalins serve as long-
lasting reservoirs of VOCs [62]. Third, generalist VSNs might sample information only from a select
subset of urinary compounds, which, given their role as biologically relevant chemosignals, might
be released at tightly controlled (and thus similar) concentrations. In fact, in the most extreme
scenario, several compounds that do display substantial strain- and/or sex-specific differences in
concentration might not act as chemosignals at all. Forth, to some extent, different response
profiles could be attributed to non-volatile small organic molecules such as steroids [9], which were
beyond the focus of our chemical analysis.

While, compared to wild-derived mice, the genetic differences in VR repertoires between C57BL/6
and BALB/c animals appear rather modest [36] (see above), vomeronasal representation of female
semiochemicals differs considerably between both inbred strains. We conclude that, even in
closely related inbred mice, strain-to-strain VR variation must be prominent, an idea supported by
various reports of differences in genetic VR makeup across strains [63–65]. With monoallelic VR
expression and 184 described non-synonymous SNPs between C57BL/6 and BALB/c receptor
genes, it is likely that even individuals of the same strain express functionally different arrays of
VSNs. Adding another layer of complexity, state- and experience-dependent changes in VSN
sensitivity have recently been described [34,66].

By adopting the same experimental design (i.e., using the same sets of stimuli in male C57BL/6
versus BALB/c mice) in both this study on VSN response profiles and our previous analysis of
sensory representations in the AOB [27], we provide a unique comparative perspective on signal
transformation along the initial processing nodes of the accessory olfactory pathway. We observe
several differences in representations of ethologically relevant urine stimuli between the VNO and
AOB. Notably, while stimulus representations across the two inbred recipient strains were very
similar in AOB recordings [27], we here observe clear differences in VSN activity, particular upon
stimulation with female stimuli. Moreover, a substantial fraction of AOB neurons were selective to
wild rather than inbred stimuli, whereas relatively few VSNs showed such selectivity. Consistent
with the elaborate wiring patterns within the AOB, these observations imply the presence of non-

trivial transformations between VSN and AOB representations. Understanding the exact nature,

purpose, and neuronal substrates of these transformations remains an important topic for future

studies.

Acknowledgements

This work was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research

Foundation) – 368482240/GRK2416 (MS); 378028035 (YBS & MS) – by the Volkswagen

Foundation (I/83533, MS), and the German-Israeli Foundation for Scientific Research and

Development (1-1193-153.13/2012, YBS & MS). We thank Corinna H. Engelhardt, Stefanie Kurth,

and Jessica von Bongartz (RWTH-Aachen University) for excellent technical assistance. We are

grateful to Pavel Talacko and Petr Zacek for Proteomic Core Facility support (BIOCEV, Charles

University).

Author contributions

Conceptualization, M.Na., P.S., Y.B.-S., and M.S.; Formal Analysis, M.Na. and P.S.; Methodology,

M.Na., D.F., R.S., P.S., and M.S.; Investigation, M.Na., M.Ni., R.B., and R.S.; Writing – Original

Draft, M.Na. and M.S.; Writing – Review & Editing, M.Na., R.B., D.F., A.L., P.S., Y.B.-S., and M.S.;

Funding Acquisition, Y.B.-S. and M.S.; Project Administration, P.S., Y.B.-S., and M.S.; Resources,

M.Na., P.S., and M.S.; Supervision, P.S., Y.B.-S., and M.S.; Visualization, M.Na., P.S., A.L., and

M.S.

Declaration of Interests

The authors declare no competing interests.

Methods

Animals.

All animal procedures were approved by local authorities at RWTH Aachen University, were

performed in accordance with local Animal Care and Use Committees' regulations, and in

compliance with European Union legislation (Directive 2010/63/EU) and recommendations by the

Federation of European Laboratory Animal Science Associations. C57BL/6 and BALB/c mice
(Charles River Laboratories, Sulzfeld, Germany) were housed in groups of both sexes [room
temperature (RT); 12:12 h light-dark cycle; food and water available *ad libitum*]. All Ca^{2+} imaging
experiments used slices from young male adults.

Urine collection from two strains of inbred mice (C57BL/6NCrl and BALB/cAnNCr) as well as first-
generation offspring of wild mice was performed at Charles University (Prague, Czech Republic)
according to the institute's ethical committee guidelines. Inbred mice were purchased from
pathogen-free facilities of the Institute of Molecular Genetics (Czech Academy of Sciences in
Prague). Wild *M. m. musculus* mice were caught in house shelters and agricultural buildings near
Prague (Czechia), transferred to the local animal facility at Charles University, and bred for one
generation. 20 male (M) and 27 female (F) wild mice were caught at six different sites in the
broader Prague area (i.e., Bohnice (50.13415N, 14.41421E; 2M+4F), Dolni Brezany (49.96321N,
14.4585E; 3M+4F), Hodkovice (49.97227N, 14.48039E; 5M+6F), Písnice (49.98988N,
14.46625E; 3M+6F), Lhota (49.95369N, 14.43087E; 1M+2F), and Zalepy (49.9532N, 14.40829E;
6M+5F). 18 of the 27 wild females were caught pregnant. The remaining 9 females were mated
with males caught at the same site and produced offspring within a month. All mice that served as
urine donors were fed on the same diet. Food and water for all strains were provided *ad libitum*
under stable conditions (13:11 h light-dark cycle; 23°C).

Chemicals and solutions.

The following solutions (S_1–S_6) were used:

(S_1) 4-(2-Hydroxyethyl)piperazine-1-ethanesulfonic acid (HEPES) buffered extracellular solution
containing (in mM) 145 NaCl, 5 KCl, 1 $CaCl_2$, 1 $MgCl_2$, 10 HEPES, pH 7.3 (adjusted with NaOH),
300 mOsm (adjusted with glucose).

(S_2) Oxygenated (95% O_2, 5% CO_2) extracellular solution containing (in mM) 125 NaCl, 25
$NaHCO_3$, 5 KCl, 1 $MgSO_4$, 1 $CaCl_2$, 5 N,N-bis(2-hydroxyethyl)-2-aminoethanesulfonic acid (BES);
pH = 7.3; 300 mOsm (adjusted with glucose).

(S_3) Elevated extracellular K^+ solution containing (in mM) 100 NaCl, 50 KCl, 1 $CaCl_2$, 1 $MgSO_4$,
10 HEPES; pH = 7.3 (adjusted with NaOH); 300 mOsm (adjusted with glucose).

If not stated otherwise, chemicals were purchased from Sigma (Schnelldorf, Germany). Solutions
and stimuli were applied from air pressure-driven reservoirs via an eight-in-one multibarrel
"perfusion pencil" (Science Products, Hofheim, Germany). Changes in focal superfusion [67] were
software-controlled and synchronized with data acquisition by transistor–transistor logic input to
12 V DC solenoid valves using a TIB 14S digital output trigger interface (HEKA Elektronik,
Lambrecht/Pfalz, Germany).

Stimuli

For all three donor types (C57BL/6NCrl, BALB/cAnNCr, and wild mice), we collected fresh urine
by gentle bladder massage from 10 adult male and female individuals, respectively (resulting in a
total of 60 individual samples), starting at an age of 10 weeks. When selecting wild individuals
(first-generation offspring) for urine collection, we ensured that all six capture sites (see above)

were represented and that age-matched animals displayed similar weight (~17g). To minimize
concentration differences that might result from sample-to-sample volume (i.e., dilution) variability,
we collected and pooled four to six samples from each individual over several days until each of
the 60 animals had provided a total urine volume of >500 µl. Next, we measured general protein
content for each sample (Bradford assay). Aliquots of 10 µl were subjected to two-dimensional
gas chromatography–mass spectrometry (GCxGC-MS) and nanoliquid chromatography–tandem
mass spectrometry (nLC-MS/MS; see below).

The remaining samples were divided into ready-to-use aliquots and stored at −86°C. Prior to
experiments, aliquots were thawed and diluted 1:100 in S_1 [68]. For each of the six strain-/sex-
specific stimulus combinations, we created pools from all 10 individuals to minimize individual-to-
individual variability. For both inbred and wild female mice, estrus stage was not determined.
However, urine collection over several days and pooling across 10 individuals in each stimulus
set is designed to reduce variability. Notably, the urine samples employed in this study are the
same stimuli used previously to compare sensory representations of inbred and wild stimuli in the
AOB of male C57BL/6 and BALB/c mice [27].

Two-dimensional gas chromatography-mass spectrometry (GCxGC-MS)

Urine VOCs were sampled with Headspace Solid Phase Micro Extraction (HS SPME) on fiber
(DVB/CAR/PDMS grey; Supelco, USA) after 5 min incubation at 55 °C. Next, VOCs were analyzed
using two-dimensional gas chromatography with mass detection (Pegasus 4D, LECO Europe
B.V., Geleen, The Netherlands) with a combination of mid-polar and non-polar separation columns
[primary column: SLB-IL60 (30 m x 0.25 mm, SigmaAldrich, USA); secondary column Rxi-5sil MS
(1.4 m x 0.25 mm, Restek, Australia)]. Parameters were set as follows: inlet temperature 270 °C,
splitless injection mode, constant He flow 1 ml min^{-1}, modulation time 4 s (hot pulse 0.6 s),
modulation temperature offset with respect to secondary oven 15 °C. Temperature program for
primary oven: 50 °C (1 min), increase to 320 °C (10 °C min^{-1}), 320 °C (3 min). +5 °C temperature
offset on secondary column. Transfer line temperature was held at 250 °C. Mass detector was
equipped with an electron ionization source and time-of-flight analyzer enabling unit mass
resolution (scanned mass range was 30 – 500 *m/z*). The ion source chamber was held at 250 °C.
ChromaTOF® v4.5 software (LECO Europe B.V.) was employed for instrument control and data
processing. Selected compounds were identified by mass spectra library matching (NIST MS 2.2,
USA). Compounds identified only in blanks (or that were highly abundant in blanks) were removed
from analysis (e.g., silanes, siloxanes, propylphosphines etc.).

Protein digestion & nano-scale liquid chromatographic tandem mass spectrometry (nLC-MS/MS)

Urine proteins were precipitated with cold acetone and centrifuged (14,000 x *g*; 10 min; 0 °C),
followed by re-suspension of dried pellets in digestion buffer (1% SDC, 100 mM TEAB; pH 8.5).
Next, protein concentration in each lysate was determined (BCA assay kit; Fisher Scientific). We
used tris(2-carboxyethyl)phosphine (TCEP; 5 mM; 60 °C; 60 min) as reducing agent and S-methyl
methanethiosulfonate (MMTS; 10 mM; 10 min; RT) to block free cysteines. After trypsin digestion
(1 µg per sample; 37 °C; overnight), peptides were desalted on a Michrom C18 column. We used

reverse-phase nanocolumns (EASY-Spray column, 50 cm x 75 µm ID, PepMap C18, 2 µm
particles, 100 Å pore size) for high resolution peptide separation. Eluting peptide cations were
converted to gas-phase ions by electrospray ionization and analyzed on a Thermo Orbitrap Fusion
(Q-OT-qIT; Thermo Fisher, Waltham, MA, USA) as previously described [69,70]. LC-MS data were
pre-processed with MaxQuant software (version 1.5.3.8) [71]. The false discovery rate (FDR) was
set to 1% for both proteins and peptides. We specified a minimum peptide length of seven amino
acids. The Andromeda search engine was used for MS/MS spectra search against Uniprot *Mus
musculus* database (downloaded June 2015), containing 44,900 entries. From this database, all
MUP and OBP sequences were removed and replaced by complete lists of MUPs (Ensembl) and
OBPs [72], respectively. We also added some sequences from TrEMBL that were missing in Uniprot
(e.g., KLKs, BPIs, SPINKs, SCGB/ABPs, and LCNs). Enzyme specificity was set as C-terminal to
Arg and Lys, also allowing cleavage at proline bonds [73] and a maximum of two missed cleavages.
Quantifications were performed using label-free algorithms [74] with a combination of unique and
razor peptides.

VNO slice preparation

For confocal Ca^{2+} imaging, acute coronal VNO slices were prepared as previously described [38,61].
Briefly, C57BL/6 and BALB/c mice were euthanized by brief exposure to a CO_2 atmosphere,
cervical dislocation and decapitation. The lower jaw and palate were rapidly removed. The VNO
was dissected, embedded in 5% low-gelling temperature agarose (VWR International, Erlangen,
Germany), placed in ice-cold oxygenated **S₂**, and coronal slices (150 µm) were cut on a VT1000S
vibrating microtome (Leica Biosystems, Nussloch, Germany). For each experiment, we routinely
prepared VNO slices along the organ's entire anterior-to-posterior axis (not including the most
anterior tip, where the VNO lumen tapers into the vomeronasal duct, and the most posterior part,
the lumen ''twists'' toward the ventral aspect and its volume decreases [75]. This usually yields ~7
slices per individual experiment / session. Therefore, we routinely sample and average across the
entire VNO anterior-to-posterior axis for each experiment. Slices were transferred to a submerged,
chilled, and oxygenated storage chamber with circulating **S₂** until use.

Ca^{2+} *imaging*

In vitro imaging of VSN activity in acute coronal VNO slices was performed as described [38]. Briefly,
for bulk loading, slices were incubated (90 min; 5°C) in circulating **S₂** with the Ca^{2+}-sensitive dye
CAL520/AM (4.5 µM; Biomol, Hamburg, Germany) and 0.05 % Pluronic® F-127 (20 % solution in
DMSO; Thermo Fisher Scientific, Schwerte, Germany). After washing (5x, **S₂**), slices were
transferred to a recording chamber (Luigs & Neumann, Ratingen, Germany) mounted on an
upright fixed-stage scanning confocal microscope (TCS SP5 DM6000CFS, Leica Microsystems)
equipped with a 20x / 1.0 NA water immersion objective (HCX APO L, Leica Microsystems), and
infrared-optimized differential interference contrast optics. Slices were continuously superfused
with oxygenated **S₂** (~5 ml / min; gravity flow). CAL520 was excited at 488 nm (multi-line argon
laser; <25% laser power) and fluorescence was detected within a 500–600 nm spectral band.
Changes in cytosolic Ca^{2+} were monitored over time at 1.0 Hz frame rate (1024 x 512 pixels; 400
Hz bidirectional scanning frequency) using LAS AF software (Leica Microsystems).

Experimental Design and Statistical Analysis.

Ca^{2+} imaging in VNO slices – All data were obtained from independent experiments performed in ≥3 sessions using ≥3 different animals. Individual numbers of cells / experiments (n) are denoted in figures and/or captions. Since, for each experiment, we prepare slices along the VNO's anterior-to-posterior axis (see above), we routinely sample and average across the entire VNO anterior-to-posterior axis for each experiment. Data were analyzed offline using Leica LAS AF 2.4 (Leica Microsystems), ImageJ 1.51n (Wayne Rasband, National Institutes of Health, USA), MATLAB R2017b (MathWorks, Natick, MA), and Excel (Microsoft, Seattle, WA) software. If not stated otherwise, results are presented as box-and-whisker plots, where boxes represent the first-to-third quartiles and whiskers represent the 10th and 90th percentiles, respectively. Outliers (1.5 IQR) are plotted individually. The central band represents the population median ($P_{0.5}$). Statistical analyses were performed using unpaired t-tests, Wilcoxon signed-rank tests, Mann-Whitney U tests, and two-sample Kolmogorov-Smirnov tests (as dictated by data distribution and experimental design). Tests and corresponding *p*-values that report statistical significance (≤0.05) are individually specified in figure captions.

For individual cell analysis, regions-of-interest (ROIs) were defined to outline essentially all depolarization-sensitive (**S_3**) somata per field of view, based on DIC imaging of cell morphology at rest. After movement correction (StackReg/Rigid Body transformation plugin [76] in ImageJ) of time-lapse image stacks, changes in relative fluorescence intensity were calculated as ΔF/F and measured in arbitrary units. Neurons were classified as 'responsive' when showing stimulus-dependent Ca^{2+} elevations in somata according to the following three criteria [38,77]: (*i*) exposure to high extracellular K^+ concentrations (50 mM; **S_3**) induced a robust Ca^{2+} transient; (*ii*) for at least one stimulation with diluted urine, a transient increase in fluorescence intensity was observed during the stimulation period; and (*iii*) the signal peak intensity exceeded the average prestimulation baseline intensity plus two standard deviations for a continuous period of at least 3 s ($I_{resp} > I_{baseline} + 2 \times SD(I_{baseline})$). All responses were normalized to positive controls (i.e., responses evoked by elevated extracellular K^+). Responsive neurons were categorized as: (*i*) neurons only sensitive to depolarization (K^+); (*ii*) *specialist* neurons that selectively responded to one of two presented urine stimuli (and K^+); and (*iii*) *generalist* neurons that responded to both urine stimuli (and K^+). Raw data (i.e., original intensity *versus* time traces) from each responsive cell were visually inspected to control for potential unspecific signals (e.g., caused by spontaneous activity). Based on peak amplitudes recorded from each urine-sensitive neuron (both generalists and specialists), we calculated a cell-specific *response index* (RI):

$$RI = \frac{mean\left(\frac{\Delta F}{F}\right)_{stim1} - mean\left(\frac{\Delta F}{F}\right)_{stim2}}{mean\left(\frac{\Delta F}{F}\right)_{stim1} + mean\left(\frac{\Delta F}{F}\right)_{stim2}}$$

where mean ($^{\Delta F}/_F$) is the average signal amplitude evoked by consecutive stimulation with the same stimulus (i.e., either stim1 or stim2). As a measure of how reliably the same stimulus evokes a response upon consecutive exposures, we additionally calculated a cell-specific *Reliability Index*

(ReI), which is based either on response amplitudes or their integrals (i.e., area under curve
(AUC)):

$$\text{ReI} = \frac{\text{response}\left(\frac{\Delta F}{F} \text{ or } AUC\right)_{1st\ stim} - \text{response}\left(\frac{\Delta F}{F} \text{ or } AUC\right)_{2nd\ stim}}{\text{response}\left(\frac{\Delta F}{F} \text{ or } AUC\right)_{1st\ stim} + \text{response}\left(\frac{\Delta F}{F} \text{ or } AUC\right)_{2nd\ stim}}$$

where consecutive responses of similar strength (i.e., high reliability) are reflected by a small ReI
value.

Proteomics and metabolomics – Pairwise comparison between different urine samples allowed
comparative analysis of physiological activity (VSN Ca^{2+} signals) *versus* urine composition
(molecular content). Chemical content was categorized as follows: (*i*) a specific compound is
considered "present" in urine from a given sex/strain combination (i.e., male or female BALB/c,
C57BL/6, or wild, respectively) if it is identified in ≥3 out of 10 individual samples; (*ii*) a compound
is thus considered "absent" if it is found only in ≤2 individual samples. Accordingly, compounds
that are detected as "present" in urine samples from both sex/strain combinations being compared
are designated as *generic* in binary comparisons. By contrast, a compound that is "present" in
only one of the two sex/strain combinations is designated as *specific*. To identify compounds that
are *enriched* in a given sex/strain combination (Figure 2a,b) we raised the criterion for a "present"
call to identification in ≥6 out of 10 individual samples.

To quantify concentration differences between samples we calculated *Concentration Indices* (CIs)
for generic compounds, both VOCs (*GCxGC-MS*) and peptides/proteins (*nLC-MS/MS*):

$$\text{CI} = \frac{mean\ conc.\ (X_{group1}) - mean\ conc.\ (X_{group2})}{mean\ conc.\ (X_{group1}) + mean\ conc.\ (X_{group2})}$$

where $mean\ conc.$ is the average concentration of a given compound *X* among all ten samples
within a group (i.e., specific sex/strain combination).

Proteomic and metabolomic analysis

Because both GCxGC-MS and nLC-MS/MS data have similar (negative binomial) distributions,
we processed them using the same procedure. First, we performed Sparse Partial Least Squares
Analysis - sPLS-DA[78] to detect potential sources of variation in quantile normalized datasets. Next,
for pairwise comparisons, data reduction eliminated all table entries (rows) if a given metabolite
or protein was detected only in ≤3 individuals in both groups. However, if a compound was found
in samples from ≥4 individuals in a group (e.g., in males or females), we included the full table
entry (row). Accordingly, a chemical is considered "unique" / group-specific if found in ≥4 individual
samples from one group, but is missing in all 10 samples of the other group. For calculation of
sexual dimorphism, we used the Power Law Global Error Model - PLGEM[79] to detect differentially
expressed / abundant proteins and VOCs. To detect the importance of significant molecules in
discriminating either sex or strain, we used a machine learning technique – Random Forest for
Classification [43] (implemented in R software) – and inferred feature importance scores (Gini
importance [43]) that provide relative rankings of individual variable relevance. nLC-MS/MS
proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE [80]

partner repository with the dataset identifier PXD042324. Metabolomics data have been deposited
to the EMBL-EBI MetaboLights database with the identifier MTBLS7439.
3
4
Supplemental information
Figure Supplements
• *Figure 1 – figure supplement 1*
• *Figure 2 – figure supplement 1*
• *Figure 2 – figure supplement 2*
• *Figure 2– figure supplement 3*
• *Figure 5 – figure supplement 1*
12
13   ***Tables***
14   • *Table 1 – relating to Figure 2*
15   • *Table 2 – relating to Figure 2 – figure supplement 1*

References

1. Dulac, C. & Torello, A. T. Molecular detection of pheromone signals in mammals: from genes to behaviour. *Nat. Rev. Neurosci.* 4, 551–562 (2003).

2. Brennan, P. A. & Zufall, F. Pheromonal communication in vertebrates. *Nature* 444, 308–315 (2006).

3. Tirindelli, R., Dibattista, M., Pifferi, S. & Menini, A. From Pheromones to Behavior. *Physiol. Rev.* 89, 921–956 (2009).

4. Mohrhardt, J., Nagel, M., Fleck, D., Ben-Shaul, Y. & Spehr, M. Signal detection and coding in the accessory olfactory system. *Chem. Senses* 43, 667–695 (2018).

5. Wilson, K. C. P. & Raisman, G. Age-related changes in the neurosensory epithelium of the mouse vomeronasal organ: Extended period of post-natal growth in size and evidence for rapid cell turnover in the adult. *Brain Res.* 185, 103–113 (1980).

6. Luo, M., Fee, M. S. & Katz, L. C. Encoding Pheromonal Signals in the Accessory Olfactory Bulb of Behaving Mice. *Science* 299, 1196–1201 (2003).

7. Wyatt, T. D. Pheromones. *Curr. Biol.* 27, R739–R743 (2017).

8. Albone, E. S. *Mammalian semiochemistry : the investigation of chemical signals between mammals*. (John Wiley & Sons, Inc., 1984).

9. Nodari, F. *et al.* Sulfated steroids as natural ligands of mouse pheromone-sensing neurons. *J. Neurosci.* 28, 6407–18 (2008).

10. Hurst, J. L. *et al.* Individual recognition in mice mediated by major urinary proteins. *Nature* 414, 631–634 (2001).

11. Novotny, M. V. Pheromones, binding proteins and receptor responses in rodents. *Biochem. Soc. Trans.* 31, 117–122 (2003).

12. Röck, F., Mueller, S., Weimar, U., Rammensee, H.-G. & Overath, P. Comparative analysis of volatile constituents from mice and their urine. *J. Chem. Ecol.* 32, 1333–46 (2006).

13. Sturm, T. *et al.* Mouse urinary peptides provide a molecular basis for genotype discrimination by nasal sensory neurons. *Nat. Commun.* 4, 1616 (2013).

14. Overath, P., Sturm, T. & Rammensee, H.-G. Of volatiles and peptides: in search for MHC-dependent olfactory signals in social communication. *Cell. Mol. Life Sci.* (2014) doi:10.1007/s00018-014-1559-6.

15. Leinders-Zufall, T. *et al.* Ultrasensitive pheromone detection by mammalian vomeronasal neurons. *Nature* 405, 792–796 (2000).

16. Rivière, S., Challet, L., Fluegge, D., Spehr, M. & Rodriguez, I. Formyl peptide receptor-like proteins are a novel family of vomeronasal chemosensors. *Nature* 459, 574–577 (2009).

17. Kimoto, H., Haga, S., Sato, K. & Touhara, K. Sex-specific peptides from exocrine glands stimulate mouse vomeronasal sensory neurons. *Nature* 437, 898–901 (2005).

18. Leinders-Zufall, T. *et al.* MHC class I peptides as chemosensory signals in the vomeronasal organ. *Science* 306, 1033–1037 (2004).

19. Chamero, P. *et al.* Identification of protein pheromones that promote aggressive behaviour. *Nature* 450, 899–902 (2007).

20. Doyle, W. I. *et al.* Faecal bile acids are natural ligands of the mouse accessory olfactory system. *Nat. Commun.* 7, 11936 (2016).

21. Fu, X. *et al.* A Molecular Code for Identity in the Vomeronasal System. *Cell* 163, 313–323 (2015).

22. Turaga, D. & Holy, T. E. Organization of vomeronasal sensory coding revealed by fast volumetric calcium imaging. *J. Neurosci.* 32, 1612–21 (2012).

23. Haga-Yamanaka, S. *et al.* Integrated action of pheromone signals in promoting courtship behavior in male mice. *Elife* 3, e03025–e03025 (2014).

24. Haga-Yamanaka, S., Ma, L. & Yu, C. R. Tuning properties and dynamic range of type 1 vomeronasal receptors. *Front. Neurosci.* 9, 1–15 (2015).

25. Celsi, F., D'Errico, A. & Menini, A. Responses to Sulfated Steroids of Female Mouse Vomeronasal Sensory Neurons. *Chem. Senses* 1–10 (2012) doi:10.1093/chemse/bjs068.

26. Isogai, Y. *et al.* Molecular organization of vomeronasal chemoreception. *Nature* 478, 241–5 (2011).

27. Bansal, R. *et al.* Do all mice smell the same? Chemosensory cues from inbred and wild mouse strains elicit stereotypic sensory representations in the accessory olfactory bulb. *BMC Biol.* 19, 133 (2021).

28. He, J., Ma, L., Kim, S., Nakai, J. & Yu, C. R. Encoding Gender and Individual Information in the Mouse Vomeronasal Organ. *Science* 320, 535–538 (2008).

29. Phifer-Rixey, M. & Nachman, M. W. Insights into mammalian biology from the wild house mouse Mus musculus. *Elife* 4, 1–13 (2015).

30. Wade, C. M. & Daly, M. J. Genetic variation in laboratory mice. *Nat. Genet.* 37, 1175–1180 (2005).

31. Duvaux, L., Belkhir, K., Boulesteix, M. & Boursot, P. Isolation and gene flow: inferring the speciation history of European house mice. *Mol. Ecol.* 20, 5248–5264 (2011).

32. Cheetham, S. A., Smith, A. L., Armstrong, S. D., Beynon, R. J. & Hurst, J. L. Limited variation in the major urinary proteins of laboratory mice. *Physiol. Behav.* 96, 253–261 (2009).

33. Duyck, K., DuTell, V., Ma, L., Paulson, A. & Yu, C. R. Pronounced strain-specific chemosensory receptor gene expression in the mouse vomeronasal organ. *BMC Genomics* 18, 965 (2017).

34. Xu, P. S., Lee, D. & Holy, T. E. Experience-Dependent Plasticity Drives Individual Differences in Pheromone-Sensing Neurons. *Neuron* 91, 878–92 (2016).

35. Ibarra-Soria, X. *et al.* Variation in olfactory neuron repertoires is genetically controlled and environmentally modulated. *Elife* 6, 1–29 (2017).

36. Wynn, E. H., Sánchez-Andrade, G., Carss, K. J. & Logan, D. W. Genomic variation in the vomeronasal receptor gene repertoires of inbred mice. *BMC Genomics* 13, 415 (2012).

37. He, J. *et al.* Distinct signals conveyed by pheromone concentrations to the mouse vomeronasal organ. *J. Neurosci.* 30, 7473–7483 (2010).

38. Wong, W. M. *et al.* Sensory Adaptation to Chemical Cues by Vomeronasal Sensory Neurons. *Eneuro* 5, ENEURO.0223-18.2018 (2018).

39. Ishii, K. *et al.* A Labeled-Line Neural Circuit for Pheromone-Mediated Sexual Behaviors in Mice. *Neuron* 1, 1–15 (2017).

40. Kaur, A. W. *et al.* Murine Pheromone Proteins Constitute a Context-Dependent Combinatorial Code Governing Multiple Social Behaviors. *Cell* 157, 676–688 (2014).

41. Lê Cao, K. A., Boitard, S. & Besse, P. Sparse PLS discriminant analysis: Biologically relevant feature selection and graphical displays for multiclass problems. *BMC Bioinformatics* 12, 0–16 (2011).

42. Gromski, P. S. *et al.* A tutorial review: Metabolomics and partial least squares-discriminant analysis – a marriage of convenience or a shotgun wedding. *Anal. Chim. Acta* 879, 10–23 (2015).

43. Breiman, L. Random Forests. *Mach. Learn. 2001 451* 45, 5–32 (2001).

44. Roberts, S. A. *et al.* Darcin: a male pheromone that stimulates female memory and sexual attraction to an individual male's odour. *BMC Biol.* 8, 75 (2010).

45. Furuhashi, M. & Hotamisligil, G. S. Fatty acid-binding proteins: role in metabolic diseases and potential as drug targets. *Nat. Rev. Drug Discov.* 7, 489–503 (2008).

46. Hori, T. *et al.* Mucopolysaccharidosis type IVA: common double deletion in the N-acetylgalactosamine-6-sulfatase gene (GALNS). *Genomics* 26, 535–542 (1995).

47. Lev, S. *et al.* Protein tyrosine kinase PYK2 involved in Ca2+-induced regulation of ion channel and MAP kinase functions. *Nature* 376, 737–745 (1995).

48. Thorn, A. *et al.* Structure and activity of the only human RNase T2. *Nucleic Acids Res.* 40, 8733–8742 (2012).

49. O'Brien, J. S. & Kishimoto, Y. Saposin proteins: structure, function, and role in human lysosomal storage disorders. *FASEB J.* 5, 301–308 (1991).

50. van de Rijn, M., Heimfeld, S., Spangrude, G. J. & Weissman, I. L. Mouse hematopoietic stem-cell antigen Sca-1 is a member of the Ly-6 antigen family. *Proc. Natl. Acad. Sci.* 86, 4634–4638 (1989).

51. McCord, J. M. & Fridovich, I. Superoxide Dismutase. *J. Biol. Chem.* 244, 6049–6055 (1969).

52. Roberts, S. A., Davidson, A. J., McLean, L., Beynon, R. J. & Hurst, J. L. Pheromonal Induction of Spatial Learning in Mice. *Science (80-.).* 338, 1462–1465 (2012).

53. Hurst, J. L. Scent marking and social communication. in *Animal Communication Networks* 219–244 (Cambridge University Press, 2005). doi:10.1017/CBO9780511610363.014.

54. Mucignat, C., Redaelli, M. & Caretta, A. One nose one brain: contribution of the main and accessory olfactory system to chemosensation. *Front. Neuroanat.* 6, 1–9 (2012).

55. Demir, E. *et al.* The pheromone darcin drives a circuit for innate and reinforced behaviours. *Nature* 578, 137–141 (2020).

56. Kahan, A. & Ben-Shaul, Y. Extracting Behaviorally Relevant Traits from Natural Stimuli: Benefits of Combinatorial Representations at the Accessory Olfactory Bulb. *PLOS Comput. Biol.* 12, e1004798 (2016).

57. Ben-Shaul, Y. Extracting Social Information from Chemosensory Cues: Consideration of Several Scenarios and Their Functional Implications. *Front. Neurosci.* 9, 1–15 (2015).

58. Beynon, R. J. *et al.* Polymorphism in major urinary proteins: Molecular heterogeneity in a wild mouse population. *J. Chem. Ecol.* 28, 1429–1446 (2002).

59. Moudra, A. *et al.* Phenotypic and Clonal Stability of Antigen-Inexperienced Memory-like T Cells across the Genetic Background, Hygienic Status, and Aging. *J. Immunol.* ji2001028 (2021) doi:10.4049/jimmunol.2001028.

60. Uchida, N., Poo, C. & Haddad, R. Coding and Transformations in the Olfactory System. *Annu. Rev. Neurosci.* 363–385 (2013) doi:10.1146/annurev-neuro-071013-013941.

61. Cichy, A. *et al.* Extracellular pH Regulates Excitability of Vomeronasal Sensory Neurons. *J. Neurosci.* 35, 4025–4039 (2015).

62. Hurst, J. L., Robertson, D. H. L., Tolladay, U. & Beynon, R. J. Proteins in urine scent marks of male house mice extend the longevity of olfactory signals. *Anim. Behav.* 55, 1289–1297 (1998).

63. Miller, C. H., Campbell, P. & Sheehan, M. J. Distinct evolutionary trajectories of V1R clades across mouse species. *BMC Evol. Biol.* 20, 99 (2020).

64. Park, S. H., Podlaha, O., Grus, W. E. & Zhang, J. The microevolution of V1r vomeronasal receptor genes in mice. *Genome Biol. Evol.* 3, 401–12 (2011).

65. Lilue, J. *et al.* Sixteen diverse laboratory mouse reference genomes define strain-specific haplotypes and novel functional loci. *Nat. Genet.* 50, 1574–1583 (2018).

66. Dey, S. *et al.* Cyclic Regulation of Sensory Perception by a Female Hormone Alters Behavior. *Cell* 161, 1334–1344 (2015).

67. Veitinger, S. *et al.* Purinergic signalling mobilizes mitochondrial Ca^{2+} in mouse Sertoli cells. *J. Physiol.* 589, 5033–55 (2011).

68. Hagendorf, S., Fluegge, D., Engelhardt, C. H. & Spehr, M. Homeostatic control of sensory output in basal vomeronasal neurons: activity-dependent expression of ether-à-go-go-related gene potassium channels. *J. Neurosci.* 29, 206–21 (2009).

69. Černá, M., Kuntová, B., Talacko, P., Stopková, R. & Stopka, P. Differential regulation of vaginal lipocalins (OBP, MUP) during the estrous cycle of the house mouse. *Sci. Rep.* 7, 1–10 (2017).

70. Stopkova, R., Klempt, P., Kuntova, B. & Stopka, P. On the tear proteome of the house mouse (Mus musculus musculus) in relation to chemical signalling. *PeerJ* 2017, (2017).

71. Voukali, E., Veetil, N. K., Němec, P., Stopka, P. & Vinkler, M. Comparison of plasma and cerebrospinal fluid proteomes identifies gene products guiding adult neurogenesis and neural differentiation in birds. *Sci. Rep.* 11, 1–16 (2021).

72. Stopková, R., Otčenášková, T., Matějková, T., Kuntová, B. & Stopka, P. Biological Roles of Lipocalins in

Chemical Communication, Reproduction, and Regulation of Microbiota. *Front. Physiol.* 12, (2021).

73. Rodriguez, J., Gupta, N., Smith, R. D. & Pevzner, P. A. Does Trypsin Cut Before Proline? *J. Proteome Res.* 7, 300–305 (2007).

74. Cox, J. *et al.* Accurate Proteome-wide Label-free Quantification by Delayed Normalization and Maximal Peptide Ratio Extraction, Termed MaxLFQ. *Mol. Cell. Proteomics* 13, 2513–2526 (2014).

75. Hamacher, C. *et al.* A revised conceptual framework for mouse vomeronasal pumping and stimulus sampling. *Curr. Biol.* 1–16 (2024) doi:10.1016/j.cub.2024.01.036.

76. Thevenaz, P., Ruttimann, U. E. & Unser, M. A pyramid approach to subpixel registration based on intensity. *IEEE Trans. Image Process.* 7, 27–41 (1998).

77. Fluegge, D. *et al.* Mitochondrial Ca2+ mobilization is a key element in olfactory signaling. *Nat. Neurosci.* 15, 754–762 (2012).

78. Rohart, F., Gautier, B., Singh, A. & Lê Cao, K.-A. mixOmics: An R package for 'omics feature selection and multiple data integration. *PLOS Comput. Biol.* 13, 1–19 (2017).

79. Pavelka, N. *et al.* A power law global error model for the identification of differentially expressed genes in microarray data. *BMC Bioinformatics* 5, 203 (2004).

80. Perez-Riverol, Y. *et al.* The PRIDE database resources in 2022: a hub for mass spectrometry-based proteomics evidences. *Nucleic Acids Res.* 50, D543–D552 (2022).

Supplementary Figures

Figure 1 – figure supplement 1: (**a**) Area under curve response reliability index histograms of all generalist VSNs (gray; n = 10,258) and all specialist VSNs (red; n = 6,457) recorded from in this study. Note that for both response types, indices are normally distributed. While histograms peak around zero, distributions are much broader as compared to amplitude-based reliability index histograms (***Fig. 1d***). (**b** & **c**) Quantification of VSN responses recorded from male C57BL/6 mice challenged with urine stimuli from two different male C57BL/6 mouse cohorts (**b**) or two different female C57BL/6 mouse cohorts (**c**). Pie charts (top) illustrate the proportions of generalist (gray; 33 % (**b**) and 16 % (**c**), respectively) and specialist neurons (≤1%; dark *versus* light blue (**b**) and dark *versus* light red (**c**), respectively) among all K⁺-sensitive VSNs (n = 1867 (**b**) and n = 1946 (**c**), respectively). Box-and-whisker plots (bottom, left) break down the urine-sensitive neurons by categories and illustrate the generalists-to-specialists ratios over individual experiments (n = 21). Boxes

represent the first-to-third quartiles. Whiskers represent the 10th and 90th percentiles, respectively. Outliers (1.5 IQR; red x) are plotted individually. The central red band represents the population median (P$_{0.5}$). Response index histograms (bottom, left) depict signal amplitude strength distributions from generalist VSNs that responded to both paired stimuli (i.e., cohort / group 1 (g1) and cohort / group 2 (g2)). Multi-peak fitting with Gaussian curves (dashed lines) resulted either in a narrow single Gaussian that centers close to zero (**b**; peak = -0.06, σ = 0.16) or two additive Gaussian curves that best represented the calculated index distribution (**c**; peak 1 = -0.01, peak 2 = 0.25). Asterisk (*) indicates statistical significance, *p* < 0.05; Wilcoxon signed-rank test. (**d**) Concentration index histograms that outline individual compound concentration imbalances of VOCs (yellow; top) and proteins (red; bottom) that are found in both male and female urine samples of either BALB/c (left) or wild (right) mice. Histograms are heterogeneous and data are not normally distributed.

Figure 2 – figure supplement 1: (*a*&*b*) After training Random Forest classifiers [43], lists of "Gini importance" [43] scores rank urine VOCs (*a*; yellow) and proteins (*b*; red) that are likely most informative to decode sex and strain, respectively. (*c* - *f*) Box-and-whisker plots quantify concentrations of the respective five VOCs (*c*&*e*) and proteins (*d*&*f*) that are ranked most important / informative (*a*&*b*). Data are plotted as log2 signal intensity for each of the six sex/strain combinations (10 individual samples each). Red (female) and blue (male) boxes represent sex; strain as indicated. Boxes represent the first-to-third quartiles. Whiskers represent the 10th and 90th percentiles, respectively. Outliers (1.5 IQR; black dots) are plotted individually. Central bands represent the population median ($P_{0.5}$).

Figure 2 – figure supplement 2: (*a*) Box-and-whisker plots illustrating mean protein and VOC content (left) as well as individual compound concentration variance (right) for the six sex/strain combinations (10 individual samples each). Asterisks (∗) indicate statistical significance, *p* < 0.05, Kruskal-Wallis test. (*b*&*c*) Box-and-whisker plots quantify concentrations of previously reported VOC (*b*) and Mup (*c*) semiochemicals, respectively. Data are plotted as log2 signal intensity for each of the six sex/strain combinations (10 individual samples each). Red (female) and blue (male) boxes represent sex; strain as indicated. Boxes depict the first-to-third quartiles. Whiskers represent the 10^{th} and 90^{th} percentiles, respectively. Outliers (1.5 IQR; black dots) are plotted individually. Central bands represent the population median ($P_{0.5}$).

Figure 2 – figure supplement 3: Hierarchical clustering of mouse urine lipocalin content reveals an unexpectedly diverse repertoire of 27 lipocalins. The horizontal tree clusters lipocalins according to proteomic homology based on similarities in protein abundances. The vertical tree clusters the 60 individual urine samples (10 samples for each of the 6 sex/strain combinations) according to patterns of lipocalin abundance (pseudocolors).

Figure 5 – figure supplement 1: (*a*&*b*) Comparison of male BALB/c generalist VSN response preferences, upon exposure to paired male (*a*) and female (*b*) urine stimuli from different strains, with strain-dependent concentration imbalances among VOCs and proteins, respectively. Response index histograms (top rows) depict distributions of generalist data outlined in Figure 5 (gray bars). With one exception (*a* (middle), male C57BL/6 *versus* male wild), histograms are not well fitted by single Gaussian curves. Rather, histogram shape is best represented by additive multi-peak fitting (dashed lines) with two (*a* (right), *b* (left)), three (*a* (left), *b* (right)), or even four (*b* (middle)) Gaussians. Individual Gaussian curves center at -0.37, 0.07, and 0.3 (*a*; left), -0.06 (*a*; middle), -0.39 and -0.02 (*a*; right), -0.32 and 0.04 (*b*; left), -0.37, -0.18, 0.01, and 0.37 (*b*; middle), -0.38, -0.1, and 0.11 (*b*; right), respectively. For comparison, transparent and overlapping concentration index histograms (bottom rows) for VOCs (yellow) and proteins (red) are also shown (data correspond to results shown in Figure 4). (*c*&*d*) Comparative overlay of generalist VSN response index histograms (top rows) and corresponding fits (bottom rows) calculated for recordings from male BALB/c (white transparent bars) *versus* male C57BL/6 (black bars) animals that were stimulated with male (*c*) and female (*d*) urine, respectively. Vertical lines represent mean values. Asterisks (∗) indicate statistical significance between BALB/c and C57BL/6 response index distributions, *p* < 0.05, two-sample Kolmogorov-Smirnov test.

int ID	MS-DB-ID	bias / sex	comp	Chem-ID	alt. name	ChEBI / CID	ori
A140	Methane, nitro-	male	1	CH_3NO_2	Nitrocarbol	77701	?
A336	Thiazole, 2-ethyl-4,5-dihydro-	male	1	C_5H_9NS	2-Ethylthiazoline	86896	rodents
A220	Butanenitrile, 2-methyl-	male	1	C_5H_9N	2-Cyanobutane	51937	insects
A305	2-sec-Butylthiazole	male	1	$C_7H_{11}NS$	2-(1-Methylpropyl)thiazole	519539	mouse
A209	2-Butenal, 2-methyl-	male	1	C_5H_8O	Tiglic aldehyde	88419	mammals
A278	Furan, 2,3-dihydro-4-(1-methylethyl)-	male	1	$C_7H_{12}O$	Furan	35559	mammals
A102	2-Butanone	female	1	C_4H_8O	Butan-2-one	28398	mammals
A497	Benzyl alcohol	female	1	C_7H_8O	Phenylmethanol	17987	mammals
A28	Acetaldehyde	female	1	C_2H_4O	Ethanal	15343	mammals
A248	Butyric acid, 3-tridecyl ester	female	1	$C_{17}H_{34}O_2$	Tridecyl butyrate	169401	rodents
A307	1-Octen-3-ol	female	1	$C_8H_{16}O$	1-Vinylhexanol	34118	ubiquitous
A765	1,2-Benzenedicarboxylic acid, butyl 2-methylpropyl ester	female	1	$C_{16}H_{22}O_4$	Butyl isobutyl phthalate	519539	mammals
int ID	MS-DB-ID	bias / strain	comp	Chem-ID	alt. name	ChEBI / CID	ori
A51	Trichloromethane	C57BL/6	2	$CHCl_3$	Chloroform	6212	ubiquitous
A14	Cyclopropane, ethyl-	C57BL/6	2	C_5H_{10}	Ethylcyclopropane	70933	ubiquitous
A74	Ethyl Acetate	C57BL/6	2	$C_4H_8O_2$	Acetoxyethane	8857	yeast
A10	Ethylenimine	C57BL/6	2	C_2H_5N	Aziridine	9033	ubiquitous
A123	n-Propyl acetate	wild	2	$CH_3COOCH_2CH_2CH_3$	Propyl ethanoate	40116	insects
A355	Pyrazine, 2-ethenyl-6-methyl-	wild	2	$C_7H_8N_2$	Pyrazine, 2-methyl-6-vinyl-	518838	ubiquitous
A659	Dodecanoic acid, isooctyl ester	wild	2	$C_{20}H_{40}O_2$	Isooctyl laurate	2216918	plants
A165	2-Pentanone	wild	2	$C_5H_{10}O$	Ethyl acetone	7895	ubiquitous
A372	Pyridine, 2,3,4,5-tetrahydro-	wild	1	C_5H_9N	Piperideine	47858	ubiquitous
A486	Benzenamine, 3-methyl-	wild	1	$C_6H_4CH_3NH_2$	3-Toluidine	7934	?
A156	Pentanal	BALB/c	2	$C_5H_{10}O$	Valeraldehyde	8063	insects
A181	Isopropylsulfonyl chloride	BALB/c	2	$C_3H_7ClO_2S$	2-Propanesulfonyl chloride	82408	bacteria
A456	Cyclohexane, isothiocyanato-	BALB/c	2	$C_7H_{11}NS$	Isothiocyanocyclohexane	14289	?
A45	Heptane, 2,4-dimethyl-	BALB/c	2	C_9H_{20}	2,4-dimethylheptane	16656	insects
A265	Benzene, 1,2,3-trimethyl-	BALB/c	1	C_9H_{12}	Hemellitol	1797279	insects
A251	Benzene, 1,2,4-trimethyl-	BALB/c	1	C_9H_{12}	Pseudocumol	7247	insects
A977	Propene	BALB/c	1	CH_2CHCH_3	Methylethylene	6378	?

Table 1: List of VOCs that – according to sPLS-DA – display the most discriminative power to classify samples by sex (top; related to Figure 2c) or strain (bottom; related to Figure 2d). Note that for all VOCs that facilitate sex discrimination, x-variate 1 is most informative. VOCs that are potentially important for strain differentiation, however, are best separated by either x-variate 1 or 2. Entries list internal mass spectrometry identifiers (int ID), identifiers extracted from MS analysis database (MS-DB-ID), the sex or strain that drives separation (bias), which two-dimensional component / x-variate represents the most discriminative variable (comp), PubChem chemical formula (Chem-ID), PubChem common or alternative name (alt. name), Chemical Entities of Biological Interest (ChEBI) or PubChem Compound Identification (CID), and putative origin (ori).

int ID	MS-DB-ID	bias / sex	Chem-ID	alt. name	ChEBI / CID	ori
A140	Methane, nitro-	male	CH_3NO_2	Nitrocarbol	77701	?
A220	Butanenitrile, 2-methyl-	male	C_5H_9N	2-Cyanobutane	51937	insects
A235	3-Penten-2-one	male	C_5H_8O	(E)-pent-3-en-2-one	637920	insects
A336	Thiazole, 2-ethyl-4,5-dihydro-	male	C_5H_9NS	2-Ethylthiazoline	86896	rat
A13	Methanethiol	male	CH_3SH	MTMT	878	mammals
A20	Methylamine, N,N-dimethyl-	male	C_3H_9N	Trimethylamine	1146	mouse
A765	1,2-Benzenedicarboxylic acid, butyl	female	$C_{16}H_{22}O_4$	Butyl isobutyl phthalate	519539	mammals
A240	2-Penten-1-ol, (Z)-	male	$C_5H_{10}O$	trans-2-Pentenol	5364920	insects
A209	2-Butenal, 2-methyl-	male	C_5H_8O	Tiglic aldehyde	88419,	mammals
A305	2-sec-Butylthiazole	male	$C_7H_{11}NS$	2-(1-Methylpropyl)thiazole	519539	mouse
A337	2-sec-Butylthiazole (2nd variant)	male	$C_7H_{11}NS$	2-(1-Methylpropyl)thiazole	519539	mouse
A102	2-Butanone	female	C_4H_8O	Butan-2-one	28398	mammals
A152	2,4-Dimethyl-1-hexene	male	C_8H_{16}	2,4-Dimethylhex-1-ene	519301	?
A618	6-Methyl-2-pyridinecarbaldehyde	male	C_7H_7NO	6-Methylpicolinaldehyde	70737	?
A84	4-Hexen-3-one, 5-methyl-	male	$C_7H_{12}O$	5-Methylhex-4-en-3-one	256081	insects

int ID	MS-DB-ID	bias / strain	Chem-ID	alt. name	ChEBI / CID	ori
A51	Trichloromethane	C57BL/6	$CHCl_3$	Chloroform	6212	ubiquitous
A265	Benzene, 1,2,3-trimethyl-	BALB/c	C_9H_{12}	Hemellitol	1797279	insects
A63	Acetone	C57BL/6	C_3H_6O	Dimethyl ketone	180	ubiquitous
A14	Cyclopropane, ethyl-	C57BL/6	C_5H_{10}	Ethylcyclopropane	70933	ubiquitous
A172	(S)-(+)-2-Pentanol	wild	$C_5H_{12}O$	2-Pentanol	22386	mouse
A222	2-Hexanone	wild	$C_6H_{12}O$	Hexanone	11583	mouse
A165	2-Pentanone	wild	$C_5H_{10}O$	Ethyl acetone	7895	ubiquitous
A251	Benzene, 1,2,4-trimethyl-	BALB/c	C9H12	Pseudocumol	7247	insects
A123	n-Propyl acetate	wild	$C_5H_{10}O_2$	Propyl acetate	7997	insects
A355	Pyrazine, 2-ethenyl-6-methyl-	wild	$C_7H_8N_2$	Pyrazine, 2-methyl-6-vinyl-	518838	mammals
A74	Ethyl Acetate	C57BL/6	$C_4H_8O_2$	Acetoxyethane	8857	yeast
A156	Pentanal	BALB/c	$C_5H_{10}O$	Valeraldehyde	8063	insects
A146	Cyclohexene,3-(1-methylpropyl)-	wild	$C_{10}H_{18}$	3-Sec-butyl-1-cyclohexene	139917	insects
A194	o-Xylene	wild	$C_6H_4(CH_3)_2$	Ortho-Xylene	7237	?
A83	Benzene	wild	C_6H_6	Benzole	241	insects

Table 2: List of VOCs that – according to Random Forest classification and resulting Gini importance scores – display the most discriminative power to classify samples by sex (top; related to Figure 2 - figure supplement 1a) or strain (bottom; related to Figure 2 - figure supplement 1b). As for several compounds listed in Table 1, many VOCs that display high relative rankings of individual variable relevance are common metabolites. Generally, it is reassuring that several VOCs are listed in both Table 1 and Table 2, emphasizing that two different supervised machine learning algorithms (i.e., sPLS-DA (Table 1) and Random Forest (Table 2)) yield largely congruent results. Here, entries (in blue if identified by both sPLS-DA and Random Forest) list internal mass spectrometry identifiers (int ID), identifiers extracted from MS analysis database (MS-DB-ID), the sex or strain that drives separation (bias), PubChem chemical formula (Chem-ID), PubChem common or alternative name (alt. name), Chemical Entities of Biological Interest (ChEBI) or PubChem Compound Identification (CID), and putative origin (ori).