A computational model predicts sex-specific responses to calcium channel blockers in mammalian mesenteric vascular smooth muscle

Gonzalo Hernandez-Hernandez, Samantha C. O'Dwyer, Collin Matsumoto,
Mindy Tieu, Zhihui Fong, Pei-Chi Yang, Timothy J. Lewis[&],
L. Fernando Santana*, and Colleen E. Clancy*^

Department of Physiology & Membrane Biology, ^Center for Precision Medicine and
Data Science, University of California School of Medicine, Davis, California, 95616
[&] Department of Mathematics, University of California, Davis, California, 95616

*Correspondence to: ceclancy@ucdavis.edu and lfsantana@ucdavis.edu

Running title: sex-specific models of arterial myocytes

Abstract

The function of the smooth muscle cells lining the walls of mammalian systemic arteries and arterioles is to regulate the diameter of the vessels to control blood flow and blood pressure. Here, we describe an *in-silico* model, which we call the "Hernandez-Hernandez model", of electrical and Ca^{2+} signaling in arterial myocytes based on new experimental data indicating sex-specific differences in male and female arterial myocytes from murine resistance arteries. The model suggests the fundamental ionic mechanisms underlying membrane potential and intracellular Ca^{2+} signaling during the development of myogenic tone in arterial blood vessels. Although experimental data suggest that $K_V1.5$ channel currents have similar amplitudes, kinetics, and voltage dependencies in male and female myocytes, simulations suggest that the $K_V1.5$ current is the dominant current regulating membrane potential in male myocytes. In female cells, which have larger $K_V2.1$ channel expression and longer time constants for activation than male myocytes, predictions from simulated female myocytes suggest that $K_V2.1$ plays a primary role in the control of membrane potential. Over the physiological range of membrane potentials, the gating of a small number of voltage-gated K^+ channels and L-type Ca^{2+} channels are predicted to drive sex-specific differences in intracellular Ca^{2+} and excitability. We also show that in an idealized computational model of a vessel, female arterial smooth muscle exhibits heightened sensitivity to commonly used Ca^{2+} channel blockers compared to male. In summary, we present a new model framework to investigate the potential sex-specific impact of anti-hypertensive drugs.

70 **Introduction**
71
Our primary objective was to develop and implement a novel computational model that
comprehensively describes the essential mechanisms underlying electrical activity and
Ca^{2+} dynamics in arterial myocytes. We aimed to uncover the key components necessary
and sufficient to fully understand the behavior of arterial vascular smooth muscle
myocytes and the cellular response to variations in pressure. The model represents the
first-ever integration of sex-specific variations in voltage-gated $K_V2.1$ and $Ca_V1.2$
channels, enabling the prediction of sex-specific disparities in membrane potential and
the regulation of Ca^{2+} signaling in smooth muscle cells from systemic arteries. To further
investigate sex-specific responses to antihypertensive medications, we extended our
investigation to include a one-dimensional (1D) representation of tissue. This approach
enabled us to simulate and forecast the effects of Ca^{2+} channel blockers within the
controlled environment of an idealized mesenteric vessel. It is worth noting that this
computational framework can be expanded to predict the consequences of
antihypertensive drugs and other perturbations, transitioning seamlessly from single-cell
to tissue-level simulations.
87
Previous mathematical models[1–4] of vascular smooth muscle myocytes generated to
describe the membrane potential and Ca^{2+} signaling in vascular smooth muscle cells
have described the activation of G-protein-coupled receptors (GPCRs) by endogenous or
pharmacological vasoactive agents activating inositol 1,4,5-trisphosphate (IP_3) and
ryanodine (RyR) receptors resulting in the initiation of calcium waves. Earlier models
have also provided insights into the contraction activation by agonists and the behavior
of vasomotion. In a major step forward, the Karlin model[5] incorporated new cell structure
data and electrophysiology experimental data in a computational model that predicted the
essential behavior of membrane potential and Ca^{2+} signaling arising from intracellular
domains found in arterial myocytes. One notable limitation of earlier models is that they
are based entirely on data from male animals. Furthermore, many data used to
parameterize the Karlin model were obtained from smooth muscle from cerebral arteries.
While cerebral arteries are important for brain blood flow, they do not control systemic

blood pressure. Furthermore, they do not take into consideration the role of $K_V2.1$
channels in the regulation of smooth muscle cell membrane potential.

The function of the smooth muscle cells that wrap around small arteries is to regulate the
diameter of these vessels. Arterial myocytes contract in response to increases in
intravascular pressure[6]. Based on work largely done using cerebral arterial smooth
muscle, a model has been proposed in which this myogenic response is initiated when
membrane stretch activates Na^+-permeable canonical TRPC6[7,8] and melastatin-type
TRPM4[9,10]. A recent study in smooth muscle from mesenteric arteries identified two
additional TRP channels to the chain of events that link increases in intravascular
pressure to arterial myocyte depolarization: TRPP1 (PKD1) and TRPP2(PKD2)
channels[11,12]. Together, these studies point to an elaborate multiprotein complex that
plays a critical role in sensing pressure and initiating the myogenic response by inducing
membrane depolarization and activating voltage-gated, dihydropyridine-sensitive L-type
$Ca_V1.2$ Ca^{2+} channels[13,14]. Ca^{2+} entry via a single or small cluster of $Ca_V1.2$ channels
produces a local increase in intracellular free Ca^{2+} ($[Ca^{2+}]_i$) called a "$Ca_V1.2$ sparklet"[15–
18]. Activation of multiple $Ca_V1.2$ sparklets produces a global increase in $[Ca^{2+}]_i$ that
activates myosin light chain kinase, which initiates actin-myosin cross-bridge cycling and
thus contraction[19].

Negative feedback regulation of membrane depolarization and Ca^{2+} sparklet activation
occurs through the activation of large-conductance, Ca^{2+}-activated K^+ (BK_{Ca}) channels as
well as voltage-dependent $K_V2.1$ and $K_V1.5/1.2$ K^+ channels[20–23]. BK_{Ca} channels are
organized into clusters along the sarcolemma of arterial myocytes[24] and are activated by
Ca^{2+} sparks resulting from the simultaneous opening of ryanodine receptors type 2
(RyR2) located in a specialized junctional sarcoplasmic reticulum (SR) [22,25–28]. Because
the input resistance of arterial myocytes is high[29,30](about 2-10 GΩ), even relatively small
currents (10-30 pA) produced by the activation of a small cluster[22,31,32] of 6-12 BK_{Ca}
channels by a Ca^{2+} spark can transiently hyperpolarize the membrane potential of these
cells by 10-30 mV. Accordingly, decreases in BK_{Ca}, $K_V1.2$, $K_V1.5$, and/or $K_V2.1$ channels

depolarize arterial myocytes, increasing $Ca_V1.2$ channel activity, $[Ca^{2+}]_i$, and contraction

of arterial smooth muscle[21,33–36].

A recent study by O'Dwyer *et al.*[20] suggested that $K_V2.1$ channels have dual conducting

and structural roles in mesenteric artery smooth muscle with opposing functional

consequences. Conductive $K_V2.1$ channels oppose vasoconstriction by inducing

membrane hyperpolarization. Paradoxically, by promoting the structural clustering of the

$Ca_V1.2$ channel, $K_V2.1$ enhances Ca^{2+} influx and induces vasoconstriction. Interestingly,

$K_V2.1$ protein is expressed to a larger extent in female than in male arterial smooth

muscle. This induced larger $Ca_V1.2$ clusters and activity in female than in male arterial

myocytes.

Here, we describe a new model, which we call the "Hernandez-Hernandez model", of

mesenteric smooth muscle myocytes that incorporates new electrophysiological and Ca^{2+}

signaling data suggesting key sex-specific differences in male and female arterial

myocytes. The model simulates membrane currents and their impact on membrane

potential as well as local and global $[Ca^{2+}]_i$ signaling in male and female myocytes. The

Hernandez-Hernandez model predicts that $K_V2.1$ channels play a critical, unexpectedly

large role in the control of membrane potential in female myocytes compared to male

myocytes. Importantly, our model predicts that clinically used antihypertensive $Ca_V1.2$

channel blockers cause larger reductions in $Ca_V1.2$ currents in female than in male

arterial myocytes.

Finally, we present a one-dimensional (1D) vessel representation of electrotonically

coupled arterial myocytes connected in series. Predictions from the idealized vessel

suggest that Ca^{2+} channel blockers are more potent in females resulting in a more

substantial $[Ca^{2+}]_i$ reduction in female arterial smooth muscle compared to male. The

Hernandez-Hernandez model demonstrates the importance of sex-specific differences in

$Ca_V1.2$ and $K_V2.1$ channels and suggests the fundamental electrophysiological and Ca^{2+}

linked mechanisms of the myogenic tone. The model also points to testable hypotheses

underlying differential sex-based pathogenesis of hypertension and distinct responses to

antihypertensive agents.

RESULTS

In this study, we developed a computational model of the electrical activity of an isolated vascular smooth muscle cell (**Figure 1**). A key goal was to optimize and validate the model with experimental data and then use the model to predict the effects of measured sex-dependent differences in the electrophysiology of smooth muscle myocytes.

In constructing the model, we first set out to measure the kinetics of the voltage-gated L-type $Ca_V1.2$ currents (I_{Ca}) in male and female myocytes using Ca^{2+} as the charge carrier as shown in **Figure 2**. These data provided information on the kinetics of Ca^{2+}-dependent activation and inactivation of I_{Ca}. I_{Ca} is critical in determining cytosolic concentration $[Ca^{2+}]_i$ in vascular mesenteric smooth muscle cells and is the predominant pathway for Ca^{2+} entry[13,15,16,18,28,52]. Experiments using whole-cell patch-clamp were undertaken to measure the time constants of activation and deactivation (**panel 2A**) and inactivation (**panel 2B)** in male and female mesenteric artery smooth muscle cells shown as black and blue symbols, respectively. While the data from male (n = 10) and female (n = 12) myocytes showed comparable activation time constants, there was an observable trend of faster inactivation in the female cells in the lower voltage range, but the differences were not statistically significant. Steady-state activation and inactivation were also measured as shown in **panel 2C**, with male data in black symbols and female as blue symbols. No observable differences in the gating characteristics of the male and female I_{Ca} were measured. Finally, the current-voltage relationship is shown from measurements in female (blue) and male (black) in **panel 2D.** This analysis suggests that the amplitude of I_{Ca} was larger in female than in male myocytes over a wide range of membrane potentials.

We next used the experimental measurements to build and optimize a Hodgkin-Huxley model based on the data described above. The model includes voltage-dependent activation and inactivation gating variables, dL and dF, respectively. We modeled both gates following the approach by Kernik *et al.* [45]. It is important to note that smooth muscle cells operate within a voltage regime defined by the window current, which ranges between -45 mV and -20 mV. Under these conditions, $[Ca^{2+}]_i$ remains below 1 µM.

Therefore, we did not consider the Ca^{2+}-dependent inactivation gating mode of the
channel[2,53].

The model of I_{Ca} is described by:

(1) $$I_{Ca} = P_{Ca} * dL * dF * \frac{z_{Ca}^2 F^2 V}{RT} \left(\frac{[Ca]_i e^{\frac{z_{ca} F V}{RT}} - [Ca]_{out}}{e^{\frac{z_{ca} F V}{RT}} - 1} \right)$$

Where P_{Ca} is the ion permeability, R is the gas constant, F is the Faraday's constant, and
z_{Ca} is the valence of the Ca^{2+} ion. Parameters were optimized to male and female
experimental data as shown for activation time constants ($\tau_{activation}$) and inactivation time
constants ($\tau_{inactivation}$) as solid lines in **Figure 2A** and **Figure 2B**, respectively. Model
optimization to male and female activation and inactivation curves are shown in **Figure**
2C. The model was also optimized to the I_{Ca} current-voltage (I-V) relationships shown as
solid lines in **Figure 2D**.

We next set out to determine sex differences in voltage-gated K^+ currents (I_K) in male and
female mesenteric smooth muscle cells. I_K is produced by the combined activation of K_V
and BK_{Ca} channels. Following the approach previously published by O'Dwyer *et al.,*[20] we
quantified the contribution of K_V (I_{KV}) and BK_{Ca} (I_{BKCa}) current to I_K. K^+ currents were
recorded before and after the application of the channel blocker iberiotoxin (IBTX;
100nm). Once identified the contribution I_{BKCa} current, we isolated the voltage-gated
potassium currents (I_{KV}) whose contributors include the voltage-gated potassium
channels $K_V1.5$ and $K_V2.1$. The presumed function of $K_V1.5$ and $K_V2.1$ channels on
membrane potential is to produce delayed rectifier currents to counterbalance the effect
of the inward currents[19,20].

Having isolated I_{KV}, $K_V2.1$ currents were identified using the application of the $K_V2.1$
blocker ScTx1 (100 nM). As a result, the remaining ScTx1-insensitive component of the
I_{KV} current was attributed to $K_V1.5$ channels. The results are shown in **Figure 3**.
Experiments using whole-cell patch-clamp were undertaken to measure the steady-state

activation G/G_{max} of the $K_V2.1$ current ($I_{Kv2.1}$) as shown in **panel 3A** in female (blue) and

male (black) myocytes and no significant differences were observed. Measurements of

time constants of activation (**panel 3B**) of $I_{Kv2.1}$ in the voltage range of -30 to +40 mV in

female (blue, n=10) and male (black, n=7) myocytes exhibited significant differences.

Notably, activation time constants in male myocytes were smaller than those in female

myocytes, corresponding to a faster activation rate in males. The current-voltage

relationship of $I_{Kv2.1}$ is shown from measurements in female (blue, n=20) and male (black,

n=10) myocytes in **panel 3C**. Significant differences were observed in $I_{Kv2.1}$ at various

voltages. In **panel 3C**, data points without asterisks are not considered significant.

Similarly, we measured the steady-state activation of the $K_V1.5$ current ($I_{Kv1.5}$) as shown

in **panel 3D** where male and female experimental data in myocytes are shown with blue

and black symbols. Properties of $I_{Kv1.5}$ steady-state activation G/G_{max} show minimal sex-

specific differences. The current-voltage relationship of $I_{Kv1.5}$ is shown from

measurements in female (blue, n=10) and male (black, n=7) myocytes in **panel 3E.**

Finally, the current-voltage relationship of the contribution from $I_{Kv1.5}$ and $I_{Kv2.1}$ to the total

voltage-gated current (I_{KVTOT}) is shown in **panel 3F** with male and female data shown with

black and blue symbols, respectively. Data points in **panel D**, **panel E**, and **panel F**

without asterisks are not significant. The table in **panel 3H** summarizes the sex-

dependent maximal conductance and the current response at specific voltages of -50 mV,

-40 mV, -30 mV, and -20 mV for both $I_{Kv1.5}$ and $I_{Kv2.1}$.

To understand the contribution of each K^+ current to the total voltage-gated current

(I_{KVTOT}) in mesenteric vascular smooth muscle cells we built and optimized a Hodgkin-

Huxley model to the data described above. First, we developed a model to describe the

$K_V2.1$ current. The optimized model to $K_V2.1$ experimental data contains only a voltage-

dependent activation gating variable ($X_{2.1act}$). Since inactivation time is slow and is well

estimated by steady-state[3], we did not consider its effects in our model. The model of

$I_{Kv2.1}$ is described by:

(2) $I_{Kv2.1} = G_{Kv2.1} * X_{Kv2.1act} * (V - E_K)$

Where $G_{K2.1}$ is the maximal conductance of $K_V2.1$ channels and E_K is the Nernst potential

for potassium. Parameters were optimized to male and female experimental data as

shown for activation curves in **Figure 3A**. Model optimization to male and female time

constants of activation ($K_V2.1$ $\tau_{Activation}$) are shown as solid lines in **Figure 3B**. The model

was also optimized to the $I_{Kv2.1}$ current-voltage (I-V) relationships shown as solid lines in

Figure 3C.

Similarly, we developed a model for $K_V1.5$. The model was optimized to the $K_V1.5$

experimental data and contains only a voltage-dependent activation gating variable

($X_{Kv1.5act}$). The model of $I_{Kv1.5}$ is described by:

(3) $I_{KV1.5} = G_{Kv1.5} * X_{Kv1.5act} * (V - E_K)$

$G_{K1.5}$ is the maximal conductance of $K_V1.5$ channels and E_K is the Nernst potential for

potassium. Parameters were optimized to male and female experimental data as shown

for activation curves in **Figure 3D**. The model was also optimized to the $I_{Kv1.5}$ current-

voltage (I-V) relationships shown as solid lines in **Figure 3E**. From experiments, we

optimized the model to reproduce the overall time traces of K_V currents. The model

predicted that male and female myocytes have comparable time constants of activation

in $I_{Kv1.5}$ as shown in **Figure 3G.** Finally, the optimized model of the total voltage-gated

current (I_{KVTOT}) is shown in **Figure 3F**. The total voltage-gated K^+ current (I_{KvTOT}) is the

sum of $I_{KV1.5}$ and $I_{Kv2.1}$ mathematically described as:

(4) $I_{KvTOT} = I_{Kv2.1} + I_{Kv1.5}$

Notably, the main specific sex-specific differences observed in the total voltage-gated K^+

current (I_{KvTOT}) is attributable to the sex-specific differences in the current produced by

$K_V2.1$ channels.

We next analyzed the contribution of large-conductance calcium-activated potassium

(BK_{Ca}) channels to vascular smooth muscle cell electrophysiology. BK_{Ca} channels are

activated by membrane depolarization or increased $[Ca^{2+}]_i$ and are expressed in the

membrane of vascular smooth muscle cells with α and β1 subunits[22,54,55]. In smooth

muscle cells, Ca^{2+} sparks are the physiological activators of BK_{Ca} channels. We relied on

the assumption by Tong *et al.*[56] that BK_{Ca} currents (I_{BKCa}) are produced by two current

subtypes, one consisting of α subunits ($I_{BK\alpha}$) and the other consisting of α and β1 subunits

($I_{BK\alpha\beta1}$). Experimental evidence indicates that BK_{Ca} channels with αβ1 subunits form

clusters in the plasma membrane in specialized junctional domains formed by the

sarcoplasmic reticulum and the sarcolemma. BK_{Ca} channels with αβ1 subunits colocalize

with ryanodine receptors (RyRs) to in the junctional domains. During a Ca^{2+} spark, $[Ca^{2+}]_i$

elevations ranging from 10 to 100 µM activate BK_{Ca} channels[38,39,52,57,58]. In our model,

Ca^{2+} sparks are the physiological activators of BK_{Ca} channels.

The mathematical formulation of the BK_{Ca} with αβ1 current ($I_{BK\alpha\beta1}$) was optimized to fit the

experimental whole-cell electrophysiological data from Bao and Cox[54] obtained at room

temperature with a BK_{Ca} channel α subunit clone from mSlo-mbr5 and a β1 subunit clone

from bovine expressed in *Xenopus laevis* oocytes[54]. Experimental data for steady-state

activation and time constants of activation are shown in **Panel 4A** and **Panel 4B**

respectively. The activation gating variable (X_{ab}) depends on both voltage and junctional

calcium ($[Ca^{2+}]_{Jun}$). The activation gate was adapted from the Tong-Taggart model[56]. The

model of $I_{BK\alpha\beta1}$ is described by:

(5) $I_{BK\alpha\beta1} = P_{BKca} * X_{ab}(V, [Ca]_{Jun}) * \frac{z_K^2 F^2 V}{RT}\left(\frac{[K]_{in}e^{\frac{z_K FV}{RT}} - [K]_{out}}{e^{\frac{z_K FV}{RT}} - 1}\right)$

Where P_{BKCa} is the BK_{Ca} ion permeability, R is the gas constant, F is Faraday's constant,

and z_K is the valance of the potassium ions. Model optimization to activation curves are

shown with solid lines in **Figure 4A** at three different $[Ca^{2+}]_{Jun}$ concentrations 1 µM, 10

µM, and 100 µM. The results from the steady-state activation measurements at 10 µM

are also in agreement with the experimental data in vascular myocytes in *bufo marinus*[58]

(green symbols) which suggests that BK_{Ca} channels are exposed to a mean junctional

Ca^{2+} concentration ($[Ca^{2+}]_{Jun}$) of 10 µM. Time constants of activation were measured

experimentally at $[Ca^{2+}]_{Jun} = 0.003$ µM, our model was optimized and fit under the same

conditions shown in **Figure 4B** as solid lines. Notably when the model was run under

predicted $[Ca^{2+}]_{Jun} = 10$ µM conditions as shown in **Figure 4B** dashed lines, there was no

effect of the change in $[Ca^{2+}]_{Jun}$ on the time constant. The predicted current-voltage (I-V)

relationships of $I_{BK\alpha\beta1}$ are shown in **Figure 4C** using three different $[Ca^{2+}]_{Jun}$

concentrations; 1 µM, 10 µM, and 100 µM. We observed that the I-V curves are similar at

$[Ca^{2+}]_{Jun}$ concentrations of 10 µM (black trace) and 100 µM (orange trace) but markedly

reduced when $[Ca^{2+}]_{Jun} = 1$ µM (blue trace). As expected, the amplitude of the current

shown in the I-V curves in **Figure 4D** is sensitively dependent on the number of BK_{Ca}

channels as shown, we set $[Ca^{2+}]_{Jun} = 10$ µM and simulated the I-V curves using a BK_{Ca}

cluster size of 4, 6, 8 and 10 channels.

In vascular smooth muscle cells, the membrane potential over the physiological range of

intravascular pressures is less negative than the equilibrium potential of potassium ($E_K =$

-84 mV), suggesting active participation of inward currents regulated by sodium

conductance[19,59,60]. It has been postulated that basally activating TRP channels generate

nonselective cations currents (I_{NSC}) that depolarize the membrane potential. We built a

model for I_{NSC} as linear and time-independent cation current permeable to K^+ and Na^+

with permeability ratios $P_{Na}: P_K = 0.9:1.3$ adapted from Tong-Taggart model with a

reversal potential (E_{NSC}) described by:

(6) $E_{NSC} = \left(R * \frac{T}{F}\right) * log\left(\frac{P_K * K_{out} + P_{Na} * Na_{out}}{P_K * K_{in} + P_{Na} * Na_{in}}\right)$;

where R is the gas constant, F is the Faraday's constant, T is the temperature, Na_{in} and

K_{in} are the intracellular sodium and potassium intracellular concentrations. Similarly,

Na_{out} and Na_{out} are the extracellular sodium and potassium concentrations. The model

of I_{NSC} is described by

(7) $I_{NSC} = I_{NaNSC} + I_{KNSC}$

(8) $I_{NaNSC} = G_{NaNSC} * (V - E_{NSC})$

(9) $I_{KNSC} = G_{KNSC} * (V - E_{NSC})$

Where I_{NaNSC} represents sodium current contribution, I_{KNSC} represents potassium current

contribution, and G_{NaNSC} and G_{KNSC} are the maximal conductances of the contributing

sodium and potassium currents. In addition, we also included models for leak currents of

ion i calculated as

(10) $I_{i,b} = G_{i,b} (V - E_i)$

Where the Nernst potential of ion i with valance z_i is given by:

(11) $E_i = RT/z_i F \ln ([i]_{out}/[i]_{in})$, i = Na^+, K^+ and Ca^{2+}

Where R is the gas constant, F is the Faraday's constant, T is the temperature and $[i]_{out}$

denotes the extracellular concentration of ion i. FF

The remaining ionic currents, pumps, and transporters were optimized to data available

in the experimental literature and/or taken from computational models of vascular smooth

muscle and cardiac cells. The sodium-potassium pump (I_{NaK}) current was modeled using

data from smooth muscle cells from mesenteric resistance arteries of the guinea pig[56,61]

and the voltage dependency was adapted from the Luo-Rudy II model[41]. The sodium-

calcium exchanger current (I_{NCX}) was adapted from the formulation in the ten Tusscher

model[62] and the Luo-Rudy II model[41]. Finally, the sarcolemma calcium pump (I_{PMCA})

current was adapted from the Kargacin model[63].

(12) $N_{pow} = \left(Q10^{\frac{T-309.2}{10}} \right);$

(13) $N_1 = \frac{(K_{out}^{1.1})}{K_{out}^{1.1} + KmNaK_K^{1.1}};$

(14) $N_2 = \frac{(Na_{in}^{1.7})}{Na_{in}^{1.7} + KmNaK_{Na}^{1.7}};$

(15) $N_0 = \frac{1.0}{1 + \left(0.1245*exp\left(-0.1*V*\frac{F}{R*T}\right)\right) + \left(2.19e{-}3*\left(e^{\left(\frac{Na_{out}}{49.71}\right)}\right)*e^{\left(-1.9*V*\frac{F}{R*T}\right)}\right)};$

(16) $I_{NaK} = I_{NaK_{max}} * N_1 * N_2 * N_0 * N_{pow}$

(17) $phi_F = exp\left(gammax * V * \frac{F}{R*T}\right)$

(18) $phi_R = exp\left((gammax - 1) * V * \frac{F}{R*T}\right)$

(19) $X_{NCX} = \frac{(Na_{in}^3)*Ca_{out}*phi_F - (Na_{out}^3)*Ca_i*phi_R}{1+0.0003*\left((Na_{out}^3)Ca_{in}+(Na_{in}^3)*Ca_{out}\right)};$

(20) $I_{NCX} = P_{NCX} * X_{NCX}$

(21) $I_{PMCA} = I_{PMCAbar} * \frac{Ca_i^2}{Ca_i^2+K_{mPMCA}^2}$

We next set out to connect the ionic models and models of Ca^{2+} handling to make

predictions in the whole cell. In **Figure 5**, experimental data indicate that the electrical

activity of isolated mesenteric smooth muscle cells in male and female myocytes recorded

in current-clamp mode, is characterized by an oscillating membrane potential under

physiological conditions. The membrane potential is marked by repetitive spontaneous

transient hyperpolarization (TH), a ubiquitous feature of vascular smooth muscle

cells[57,64–66] as shown in **panel 5A.** Both male (black trace) and female (blue trace)

myocytes exhibited membrane hyperpolarizing transients in the potential range of -50 to

-20 mV. Notably, we observed that female myocytes always maintained a higher

depolarizing state between the hyperpolarization events compared to male myocytes.

We assessed the predictive capacity of our *in silico* model by comparing it to experimental

data. We first compared the morphology of the membrane potential in experiments **panel**

5A versus simulations **panel 5B** in male and female myocytes. Upon comparative

analysis between male and female experimental data and simulations, we noted that the

baseline membrane potential for male myocytes was around –40 mV, while female

myocytes exhibited a slightly more depolarized membrane potential at approximately –30

mV. Despite these variations in baseline membrane potential, both male and female

myocytes presented similar peak hyperpolarization values of approximately 10-15 mV,

ranging from –50 mV to –30 mV. Similarly, the frequency of THs from multiple myocytes

was calculated to be 1 to 2.8 Hz in the range of –50 mV to –30 mV which is identical to

the simulated frequency.

In the physiological range in which smooth muscle cells operate (-50 to -20 mV), ionic

currents are small and produced by the activation of a small number of ion channels.

Local fluctuations in the function of ion channels lead to noisy macroscopic signals that

are important to the variability of vascular smooth muscle cells[19]. In addition, smooth

muscle cells are subject to high input resistance where small perturbations can lead to

large changes in the membrane potential[19,29]. To approximate the physiological realities,

we applied two sources of noise to our deterministic *in silico* model to simulate the

stochastic fluctuations. The first source of the noise was introduced by adding a

fluctuating current term to the differential equations describing changes in membrane

potential (dV/dt) which represents the combined effect of the stochastic activity of ion

channels in the plasma membrane[46]. Second, we introduced noise into the $[Ca]_{SR}$ to

replicate the physiological responses consistent with those observed in experimental

studies[40]. Simulated whole-cell membrane potential with physiological noise is shown in

Figure 5B in male (black trace) and female (blue trace) myocytes.

We conducted a sensitivity analysis to determine which model parameters could underlie

the sex-specific differences observed in the experimental data. It is important to note that

we have experimental data indicating the amplitude and kinetics for a variety of currents

in male and female myocytes. For this reason, those model components were fit to the

data, fixed, and were not subject to sensitivity analysis. Our analysis, which focused solely

on variations in maximal conductance and maximal ion transport rates of the

transmembrane currents, indicated that the non-selective cation currents (I_{NSC}) and

delayed rectifier currents ($I_{KvTOT} = I_{Kv2.1} + I_{Kv1.5}$) interact to regulate the baseline membrane

potential in both male and female vascular smooth muscle myocytes (**Figure 5C**). Given

that I_{KvTOT} responds to depolarization, the primary stimulus that triggers depolarization

was determined to be attributable solely to the non-selective cation currents (I_{NSC}).

Indeed, when we adjusted the conductance of the non-selective cation currents and

implemented an increase in the conductance of I_{NSC} in the female model, we readily

reproduced the sex-specific baseline membrane potential observed experimentally

(**Figure 5A)**.

Next, using the whole-cell vascular smooth muscle myocyte computational model, we

investigated the sex-specific differences in the contribution to total voltage-gated current

(I_{KVTOT}) in mesenteric vascular smooth muscle cells. An interesting prediction from the *in*

silico simulations is that at different depolarizing states (-45, -40, and -35 mV) induced by

changing the conductance of nonselective cationic leak currents (I_{NSC}), the contribution

of $I_{Kv2.1}$ and $I_{Kv1.5}$ to I_{KVTOT} is different based on sex. In male vascular myocytes, the

contribution to total voltage-gated current (I_{KVTOT}) is largely attributable to the current

produced by $K_V1.5$ channels as shown in the lower panel in **Figure 6A**. Our results are

consistent with previous studies[35,67,68] in animal rodent male models showing the

characteristic behavior of $I_{Kv1.5}$ to control membrane potential. However, the model

predicts that in female myocytes, the contribution to total voltage-gated current (I_{KVTOT}) is

largely provided by the current produced by $K_V2.1$ channels as shown in the upper panel

in **Figure 6B**. To illustrate this point quantitatively, at a membrane potential of -40 mV,

the contribution of I_{KVTOT} from $I_{Kv1.5}$ and $I_{Kv2.1}$ is 86% and 14%, respectively, in male

myocytes compared to female myocytes in which the contribution from $I_{Kv1.5}$ and $I_{Kv2.1}$ is

23% and 77%, respectively. Regardless of the depolarization state at -45, -40, or -35 mV,

the profiles for male and female myocytes remain essentially the same as shown in

Figures 6C, 6D, and 6E. The *in silico* simulations suggest a distinctive sex-based

function of $K_V1.5$ and $K_V2.1$ channels that produce the delayed rectifier currents to

counterbalance the effect of inward currents causing graded membrane potential

depolarizations.

Having explored the regulation of graded membrane potential by the activation of I_{KVTOT}

to counterbalance the nonselective cations currents (I_{NSC}), we next explored the effects

of steady membrane depolarization in the *in silico* vascular smooth muscle cell myocyte

model on I_{Ca} in male and female myocytes. We predicted I_{Ca} in our male and female

simulations at steady-state membrane depolarization after simulation for 500 seconds.

We observed that as the membrane depolarizes from -55 to -35 mV, I_{Ca} in male myocytes

increased from 0 to 1.0 pA while in female myocytes I_{Ca} increased from 0 to 1.5 pA as

shown in **Figure 7A**, suggesting that I_{Ca} is larger in female compared to those of male

myocytes. We recorded the predicted $[Ca^{2+}]_i$ and observed that I_{Ca} led to a higher calcium

influx in female compared to male simulations as shown in **Figure 7B**. To illustrate in

detail, we show in **Figure 7C-D**, time traces of *in silico* predictions of membrane voltage

at -40 mV (*top panel*), I_{Ca} (*middle panel*), and $[Ca^{2+}]_i$ (*lower panel*) corresponding to the

male and female data points indicated by black and blue arrows respectively shown in

Figure 7A-B. In the male case (**Figure 7C**), at a steady membrane potential of -40 mV,

L-type calcium $Ca_V1.2$ channels produced a current of 0.5 pA. However, in female

simulations (**Figure 7D**), we observed that at a steady membrane potential of -40 mV, L-

type calcium $Ca_V1.2$ channels produced a current of 0.65 pA. We calculated that at -40

mV, two $Ca_V1.2$ channels are needed to sustain 0.5 pA of current in male myocytes while

three $Ca_V1.2$ channels are needed to sustain 0.65 pA of current in female myocytes.

Although the sex-specific differences in male and female simulations at -40 mV are small,

a 15 nM difference in the overall response of $[Ca^{2+}]_i$ can have a profound effect on the

constriction state of the myocytes. The predictions from the Hernandez-Hernandez model

provide a comprehensive picture of physiological conditions and support the idea that a

small number of $Ca_V1.2$ channels supply the steady Ca^{2+} influx needed to support a

maintained constricted state in small arteries and arterioles[53,69]. The differences between

males and females are notable in the context of observations indicating varied sex-based

responses to antihypertensive agents that target the Ca^{2+} handling system in vascular

smooth muscle cells.

Next, we simulated the effects of calcium channel blocker nifedipine on I_{Ca} at a steady

membrane potential of -40 mV in male and female simulations. Briefly, previous studies[70]

have shown that at the therapeutic dose of nifedipine (i.e., about 0.1 μM) L-type Cav1.2

channel currents are reduced by about 60-70%. Accordingly, we decreased I_{Ca} in our

mathematical simulations by the same extent. In **Figure 7C-D**, we show the predicted

male (gray) and female (pink) time course of membrane voltage at -40 mV (*top panel*),

I_{Ca} (*middle panel*), and $[Ca^{2+}]_i$ (*lower panel*). First, we observed that in both male and

females 0.1 μM nifedipine modifies the frequency of oscillation in the membrane potential,

by causing a reduction in oscillation frequency. Second, both male and female simulations

(middle panels) show that 0.1 μM nifedipine caused a reduction of I_{Ca} to levels that are

very similar in male and female myocytes following treatment. Consequently, the
reduction of I_{Ca} causes both male and female simulations to reach a very similar baseline
$[Ca^{2+}]_i$ of about 85 nM (lower panels). As a result, simulations provide evidence supporting
the idea that $Ca_V1.2$ channels are the predominant regulators of intracellular $[Ca^{2+}]$ entry
in the physiological range from -40 mV to -20 mV. Importantly, these predictions also
suggest that clinically relevant concentrations of nifedipine cause larger overall reductions
in Ca^{2+} influx in female than in male arterial myocytes.

Thus far, we have shown the development and application of models of vascular smooth
muscle myocytes incorporating measured sex-specific differences in currents from male
and female isolated cells. Given that hypertension is essentially a consequence of the
spatial organization and function of smooth muscle cells[71,72], we next expanded our study
to include a one-dimensional (1D) tissue representation of electrotonically coupled tissue
by connecting arterial myocytes in series.

A well-known phenomenon in excitable systems is that electrotonic coupling between
cells results in the minimization of individual cellular differences, thereby producing a
smoothing effect across the tissue[73–75]. We simulated 400 female or 400 male vascular
smooth muscle myocytes and set the gap junctional conductivity to zero to uncouple the
simulated cells. As expected, the uncoupled cells in both male and female cases
demonstrated the characteristic behavior of arterial myocytes, exhibiting spontaneous
hyperpolarization. Of the 400 cells, we show the simulated representative traces of Cell
1, Cell 50, and Cell 100 for female (**Figure 8A**) and male (**Figure 8B**).

Next, we modeled 400 cells but with electrotonic coupling by setting the simulated gap
junctional resistance to 71.4 Ωcm^2 [76]. In this case, we observed that the spontaneous
hyperpolarizations, previously observed in the uncoupled cells, diminished when cells
were coupled. The overall smoothing effect observed in **Figure 8C** is attributed to the
electrotonic coupling and consequential influence of neighboring cells. The electrical
response is consistent across the spatial domain for both male (**Figure 8C**; *black trace*)
and female (**Figure 8C**-*blue trace*) one-dimensional tissue representations. Notably, the

model predicts a more depolarized female membrane potential in the one-dimensional

tissue representations consistent with experimental measurements as shown in **Figure**

8D.

Having developed an idealized model of a vessel, we set out to validate the model

predictions of variable $[Ca^{2+}]_i$ between males and females by comparing the computed

calcium signaling in vascular smooth muscle with experimental recordings O'Dwyer *et*

al.[20]. Given that membrane potential predominantly governs calcium influx in vascular

smooth muscle[14], we varied the conductance of the nonselective cation currents (I_{NSCC})

in our simulations. Tuning of I_{NSCC} was performed to replicate the effects of pressure-

induced membrane depolarization, which results in activation of the voltage-gated L-type

Ca^{2+} channels and increases $[Ca^{2+}]_i$.

Our simulations (lines) are well validated by experimental recordings (symbols) in **Figure**

9A. A distinctive feature from the model prediction, which was validated by experimental

recordings is the observation that female (**Figure 9A**, blue trace and symbols) vessels

accommodate more $[Ca^{2+}]_i$ compared to male (**Figure 9A**, black trace and symbols)

vessels. Intriguingly, the mechanism of different $[Ca^{2+}]_i$ in male and female vessels was

revealed in single-cell simulations, which showed attributable sex-based differences in L-

type Ca^{2+} currents.

Finally, in our simulations, we computed the effects of $[Ca^{2+}]_i$ after the application of

clinically relevant calcium channel blocker nifedipine. We observed a substantial

reduction of $[Ca^{2+}]_i$ in both male (**Figure 9A**, *dashed black line*) and female (**Figure 9A**,

dashed blue line). Significant differences were found in the physiological range of

intravascular pressure from 40 to 120 mmHg. In the summary data (**Figure 9B**), we

quantified the relative change of $[Ca^{2+}]_i$ in male (*black*) and female (*blue*) after the

application of 0.1 mM L-type Ca^{2+} channel blocker nifedipine at 80 mmHg and 120 mmHg.

Our results show that nifedipine, when applied to male vessels, decreases $[Ca^{2+}]_i$ by 22%

and 25% at 80 mmHg and 120 mmHg, respectively. However, the same dose of nifedipine

when applied to female vessels decreases $[Ca^{2+}]_i$ by 38% and 45% at 80 mmHg and 120

mmHg. The results suggest that female arterial smooth muscle is more sensitive to

clinically used Ca^{2+} channel blockers than male smooth muscle.

Discussion

Here, we describe the development, validation, and application of an *in-silico* model to

simulate and understand the mechanisms of electrical activity and Ca^{2+} dynamics in a

single mesenteric vascular smooth muscle cell. The Hernandez-Hernandez model is the

first model to incorporate sex-specific differences in voltage-gated $K_V2.1$ and $Ca_V1.2$

channels and predicts sex-specific differences in membrane potential and Ca^{2+} signaling

regulation in the smooth muscle of both sexes from systemic arteries. In the pursuit of

stratifying sex-specific responses to antihypertensive drugs, we expanded our exploration

to encompass a one-dimensional (1D) tissue representation. Such an approach allowed

us to simulate and predict the impact of Ca^{2+} channel blockers within a mesenteric vessel.

Notably, the computational framework can be expanded to forecast the impact of

antihypertensives and other perturbations from single-cell to tissue-level simulations.

To specifically investigate the impact of sex-specific differences measured from ion

channel experiments and their impact on membrane potential and $[Ca^{2+}]_i$, we focused on

the isolated myocyte in the absence of complex signaling pathways. We first explored the

effects of $Ca_V1.2$ and $K_V2.1$ channels on membrane potential as experimental data

suggest key sex-specific differences in channel expression and kinetics. Notably, the

peak of the current-voltage (I-V) relationship of L-type $Ca_V1.2$ current is 40% smaller in

male compared to female myocytes (**Figure 2D**).

Similarly, the peak current-voltage (I-V) relationship of the voltage-gated $K_V2.1$ current

($I_{K_V2.1}$) is 70% smaller in male compared to female myocytes at +40 mV (**Figure 3C**).

O'Dwyer and coauthors[20] showed sex-dependent expression of $K_V2.1$ in the plasma

membrane, where male arterial myocytes have a total of about 75,000 channels

compared to 183,000 channels in female myocytes. Notably, less than 0.01% of channels

are conducting in male and female myocytes. In the computational model, we found that

to reproduce the experimentally measured amplitude of the $K_V2.1$ I-V curve (**Figure 3C**),

a maximum of ~44 male $K_V2.1$ channels was sufficient to reproduce the peak current

(68.8 pA at 40 mV). In contrast, ~143 channels were predicted to be needed in female

myocytes to reproduce the experimentally measured peak current (226.42 pA at +40 mV)

of the $K_V2.1$ I-V relationship. Modeling and simulation led to the prediction that in male

arterial myocytes, I_{KvTOT} is largely dictated by $K_V1.5$ channels. In contrast, in female

arterial myocytes, $K_V2.1$ channels dominate I_{KvTOT} **(Figure 3F)**.

An important aspect of the Hernandez-Hernandez model is that it includes Ca^{2+}-mediated

signaling between RyRs in the junctional SR and BK_{Ca} channel clusters in the nearby

sarcolemma membrane. This section of the model is similar to the one included in the

Karlin model[5] with some modifications. The Karlin model described how subcellular

junctional spaces influence membrane potential and $[Ca^{2+}]_i$ in response to intravascular

pressure, vasoconstrictors, and vasodilators. In this study, we reduced the complexity of

the model representation of subcellular Ca^{2+} signaling spaces to include just three

compartments: the cytosol, SR, and the SR-sarcolemma junction. Our model represents

on average, the behavior of a single junctional SR unit that is functional in a cell at a time.

The model uses a deterministic approach but mimics the process of production of Ca^{2+}

sparks that activate BK_{Ca} channel clusters[29]. We represented the activity of the RyRs in

the junctional domain deterministically in the model so that Ca^{2+} spark- BK_{Ca} currents

occur at a frequency of about 1 Hz at -40 mV in a space equivalent to 1% of the total cell

surface area of the plasma membrane[22].

Based on experimental observations, the Hernandez-Hernandez computational model

employs three key assumptions: First, Ca^{2+} sparks in the junctional domain are initiated

by activation of RyRs, where RyR gating opening probability is correlated with SR load.

Second, Ca^{2+}sparks lead to a $[Ca^{2+}]_{Jun}$ increase between 10-20 mM to match the

amplitude measured in experiments (**Figure 4A**)[54,58]. Third, activation of BK_{Ca} channels

and the resultant current amplitude derives from the experimentally observed

spontaneous outward currents (STOCs) in both amplitude and morphology.

Notably, model simulations revealed important mechanisms that may underlie
experimental observations in measurements of membrane potential (**Figure 5A**). The
model predicts that the mechanism of intrinsic oscillatory behavior in the vascular
myocytes results from a delicate balance of currents. Activation of non-selective cation
currents (I_{NSC}) likely causes membrane depolarization, but the delayed rectifier currents
(I_{KvTOT}) oppose them, resulting in membrane potential baseline in the physiological range
of -45 to -20 mV. Interestingly, the voltage-gated L-type $Ca_V1.2$ currents activation
threshold sits within this range at ~-45 mV. Therefore, small increases in I_{NSC} can
overwhelm I_{KvTOT} below -20 mV and result in sufficient depolarization to bring the
membrane potential to the threshold for activation of I_{Ca}. It is important to note that I_{KvTOT}
increases sharply upon depolarization from -45 to -20 mV, resulting in tight control of
membrane potential and prevention of large transient depolarization resulting from I_{NSC}.
Activation of L-type Ca^{2+} channels upon depolarization and subsequent Ca^{2+} release
within the small volume junction then activates the BK_{Ca} channels, which results in
hyperpolarization. Hyperpolarization reignites the oscillatory cascade as an intrinsic
resetting mechanism. Since vascular myocytes are subject to substantial noise from the
stochastic opening of ion channels in the plasma membrane, and fluctuations in the local
junctional domain components, such as the SR load, RyR opening, and BK_{Ca} channel
activity, we included noise in the simulation. To simulate the physiological noise in the
vascular smooth muscle cell (**Figure 5B**), we added Gaussian noise to the dV/dt
equations and $[Ca]_{SR}$.

Female mesenteric artery myocytes are more depolarized than male myocytes at
physiological intravascular pressures[20]. Our model suggests that female myocytes are
more depolarized than male myocytes due to larger non-selective cation currents in
female compared to male myocytes, most likely due to the activation of Na^+-permeable
TRP channels. To our knowledge, the only TRP channels found to regulate the
membrane potential of mesenteric artery smooth muscle are TRPP1 and TRPP2[11,12].
Future work will have to determine if TRPC6[8] and TRPM4[9,10], which have been shown to
mediate the myogenic response of cerebral artery smooth muscle, and/or other non-
selective cation channels also depolarize mesenteric artery smooth muscle[77].

The Hernandez-Hernandez model predicts that very few channels (based on total current

amplitude) are likely to control the baseline fluctuations in membrane potential in the

physiological range of -60 to -20 mV. The intrinsic oscillatory properties of the vascular

myocyte operating in the low voltage regime under conditions of high resistance

membrane are similar to other types of oscillatory electrical cells including cardiac

pacemaker cells.

As shown in (**Figure 6**), the model predicts that, at -40 mV, the amplitude of steady-state

$K_V2.1$ currents is about 0.8 pA in male and 3.3 pA in female arterial myocytes, indicating

that the contribution of $K_V2.1$ and $K_V1.5$ channels to membrane potential is different in

males and females. At -30 mV, it is 2.34 pA and 9.2 pA in male and female myocytes

respectively. Assuming a single channel current at -40 and -30 mV of 0.7 pA, we

calculated that, on average, in male myocytes a single channel is open at -40 mV and 3

channels are open at any particular time at -30 mV. In female myocytes, 6 channels are

predicted to be open at -40 mV, while 13 are predicted to be active at -30 mV.

The Hernandez-Hernandez model also allowed us to calculate the number of $Ca_V1.2$

channels needed to sustain the steady-state concentration of $[Ca^{2+}]_i$ in the physiological

range from -60 to -20 mV (**Figure 7**). The model predicts that at -40 mV in mouse male

myocytes, 2 channels were required to generate 0.5 pA of steady-state $Ca_V1.2$ current.

On the other hand, we found that in female myocytes, 3 channels were sufficient to

generate 0.65 pA of $Ca_V1.2$ current. These data are consistent with the work of Rubart et

al.[69], which suggested that steady-state Ca^{2+} currents at -40 mV were likely produced by

the opening of 2 $Ca_V1.2$ channels in rat cerebral artery smooth muscle cells.

The observation that a very small number of the conducting $K_V2.1$ and $Ca_V1.2$ channels

are involved in the regulation of membrane potential and Ca^{2+} influx in male and female

arterial myocytes at physiological membrane potentials is important for several reasons.

First, the analysis suggests that small differences in the number of $K_V2.1$ and $Ca_V1.2$

channels can translate into large, functionally important differences in membrane

potential and $[Ca^{2+}]_i$ and hence affect and control myogenic tone under physiological and
pathological conditions. *Second*, the small number of $K_V2.1$ and $Ca_V1.2$ channels gating
between -40 and -30 mV likely makes smooth muscle cells more susceptible to stochastic
fluctuations in the number and open probabilities of these channels than in cells where a
large number of channels regulate membrane excitability and Ca^{2+} influx (e.g., adult
ventricular myocyte[78]). This, at least in part, likely contributes to Ca^{2+} signaling
heterogeneity in vascular smooth muscle.

Hypertension fundamentally manifests through the spatial organization of cellular
components, particularly evident in the context of the tunica media, the middle layer of
vessels is predominantly constituted of smooth muscle cells which play a pivotal role in
vessel contraction and relaxation[71,72]. Such intricate biological machinery is imperative in
orchestrating the regulation of blood flow and blood pressure. Our approach began with
a process of distillation, aiming to shed light on cellular mechanisms within isolated
vascular myocytes from small systemic vessels and arterioles, which control blood
pressure, of both male and female mice.

Earlier research has confirmed that in mesenteric arteries, the pathogenesis leading to
hypertension is largely determined by the downregulation of $K_V2.1$[36] and/or $K_V1.5$[67,79] and
a concurrent increase in the activity of $Ca_V1.2$[80] channels. Building upon this knowledge,
we broadened our study to encompass a one-dimensional (1D) tissue model of
electrotonically linked tissue, achieved by connecting arterial myocytes in series. The 1D
cable model has anatomical relevance because the structure of third and fourth-order
mesenteric arteries have a singular layer of vascular myocytes encircling the lumen in a
cylindrical arrangement. The cable structure is analogous to an "unrolled" or lateral
arrangement of the vessel. Such an approach allowed a conceptual framework to bridge
the gap between understanding the combined effects of membrane potential and $[Ca^{2+}]_i$
in isolated cells and in the wider context of small vessels.

For instance, previous studies have proposed that gap junctions enable vessels to
function in a way that is analogous to a large capacitor[57,81]. The gap junctions actively

filter and transform single-cell electrical activity into sustained responses across the

tissue[81]. Recent studies add to this understanding by demonstrating that Connexin 37

(Cx37), a component of these gap junctions, seems to be expressed in the mesenteric

arteries[82]. In our simulations, we showed (**Figure 8A-B**) that indeed uncoupled cells

exhibit a spontaneous oscillatory behavior which studies have confirmed is not an artifact

due to isolation from the vessel but rather an intrinsic behavior required to sustain

electrical signals. When the cells are connected (**Figure 8C**) the spontaneous

hyperpolarization previously observed in the uncoupled cells diminished, the effect is

attributed to the electrotonic coupling and consequential influence of neighboring cells. In

addition, in our simulations, we found that it is required to have stochastic fluctuations to

allow the system to average the membrane potential behavior that dictates the amount of

$[Ca^{2+}]_i$ in the vessels.

Regarding, $Ca_V1.2$ channels, simulations forecast the clinically relevant concentrations

(0.1 µM) at which common Ca^{2+} channel blockers (e.g., nifedipine) effectively block

$Ca_V1.2$ channels in both male and female smooth muscle (**Figure 9**). Our simulations in

isolated arterial myocytes and in the one-dimensional (1D) tissue model suggest

heightened sensitivity to calcium channel blockers in the female compared to male.

The model predictions are aligned with documented sex-specific differences in

antihypertensive drug responses[83,84]. Previous studies, notably by Kloner et al., have

illustrated this point quantitatively, highlighting a more pronounced diastolic BP response

in women (91.4%) compared to men (83%) when treated with dihydropyridine-type

channel blockers, such as amlodipine. Importantly, this distinction persisted even after

adjusting for confounding factors such as baseline BP, age, weight, and dosage per

kilogram[84]. Another interesting observation from Kajiwara et al. emphasizes that

vasodilation-related adverse symptoms occur more frequently in younger women (<50

years) compared to their male counterparts, again suggesting a heightened sensitivity to

dihydropyridine-type calcium channel blockers[85].

Limitations

The model presented here describes the necessary and sufficient ion channels, pumps,

and transporters to describe the electrical activity and Ca^{2+} signaling of an isolated

mesenteric smooth muscle cell in the absence of complex signaling pathways. Such an

approach enabled us to perform a component dissection to analyze the sex-specific

differences observed in the fundamental electrophysiology of male and female myocytes.

However, it is well known that vascular smooth muscle cells are subject to a plethora of

stimuli from endothelial cells, neurotransmitters, endocrine, and paracrine signals[5]. The

next phase of the project includes an expansion of the model to incorporate receptor

mediated signaling pathways that are essential for blood pressure control.

Excitation-contraction coupling refers to an electrical stimulus that drives the release of

calcium from the sarcoplasmic reticulum and results in the physical translocation of fibers

that underlies muscle contraction. In the present model, we did not explicitly consider the

mechanical description of muscle contraction. Nevertheless, we can imply contractile

effects by tracking membrane potential and the elevation of $[Ca^{2+}]_i$ as a proxy.

To conclude, we developed and present the Hernandez-Hernandez model of male and

female isolated mesenteric vascular myocytes. An additional limitation of our study is the

reliance on predominantly murine data. Although mouse arteries do present numerous

parallels with human arteries—including analogous intravascular pressure-myogenic

tone relationships, resting membrane potentials, and the expression of typical ionic

channels like $Ca_V1.2$, BKCa channels, and RyRs[86–88]. Future research should assess the

direct applicability and implication of our findings in human subjects.

Conclusions

The Hernandez-Hernandez model of the isolated mesenteric vascular myocyte was

informed and validated with experimental data from male and female vascular myocytes.

We then used the model to reveal sex-specific mechanisms of $K_V2.1$ and $Ca_V1.2$ channels

in controlling membrane potential and Ca^{2+} dynamics. In doing so, we predicted that very

few channels are needed to contribute to and sustain the oscillatory behavior of the

membrane potential and calcium signaling. We expanded our computational framework

to include a one-dimensional (1D) tissue representation, providing a basis for simulating

the effect of drug effects within a vessel. The model predictions suggested differences in

the response of male and female myocytes to drugs and the underlying mechanisms for

those differences. These predictions may constitute the first step towards better

hypertensive therapy for males and females.

MATERIALS AND METHODS

Section 1. Experimental

1.1 Animals

This study was performed in strict accordance with the recommendations in the Guide for

the Care and Use of Laboratory Animals of the National Institutes of Health. All of the

animals were handled according to approved institutional animal care and use committee

(IACUC) protocols of the University of California Davis. IACUC protocol number is 22503.

8- to 12-week-old male and female mice C57BL/6J (The Jackson Laboratory,

Sacramento, CA) were used in this study. Animals were housed under standard light-dark

cycles and allowed to feed and drink ad libitum. Animals were euthanized with a single

lethal dose of sodium pentobarbital (250 mg/kg) intraperitoneally. All experiments were

conducted in accordance with the University of California Institutional Animal Care and

Use Committee guidelines.

8- to 12-week-old male and female mice C57BL/6J (The Jackson Laboratory,

Sacramento, CA) were used in this study. Animals were housed under standard light-dark

cycles and allowed to feed and drink ad libitum. Animals were euthanized with a single

lethal dose of sodium pentobarbital (250 mg/kg) intraperitoneally. All experiments were

conducted in accordance with the University of California Institutional Animal Care and

Use Committee guidelines.

1.2 Isolation of arterial myocytes from systemic resistance arterioles

Third and fourth-order mesenteric arteries were carefully cleaned of surrounding adipose

and connective tissues, dissected, and held in ice-cold dissecting solution (Mg^{2+}-PSS; 5

mM KCl, 140 mM NaCl, 2mM $MgCl_2$, 10 mM glucose, and 10 mM HEPES adjusted to pH

7.4 with NaOH). Arteries were first placed in dissecting solution supplemented with 1.23

mg/ml papain (Worthington Biochemical, Lakewood, NJ) and 1 mg/ml DTT for 14 minutes

at 37˚C. This was followed by a second 5-minute incubation in dissecting solution

supplemented with 1.6 mg/ml collagenase H (Sigma-Aldrich, St. Louis, MO), 0.5 mg/ml

elastase (Worthington Biochemical, Lakewood, NJ), and 1 mg/ml trypsin inhibitor from

Glycine max (Sigma-Aldrich, St. Louis, MO) at 37˚C. Arteries were rinsed three times with

dissection solution and single cells were obtained by gentle trituration with a wide-bore

glass pipette. Myocytes were maintained at 4˚C in dissecting solution until used.

1.3 Patch-clamp electrophysiology

All electrophysiological recordings were acquired at room temperature (22–25°C) with an

Axopatch 200B amplifier and Digidata 1440 digitizer (Molecular Devices, Sunnyvale, CA).

Borosilicate patch pipettes were pulled and polished to resistances of 3-6 MΩ for all

experiments using a micropipette puller (model P-97, Sutter Instruments, Novato, CA).

Voltage-gated Ca^{2+} currents (I_{Ca}) were measured using conventional whole-cell voltage-

clamp sampled at a frequency of 50 kHz and low-pass filtered at 2 kHz. Arterial myocytes

were continuously perfused with 115 mM NaCl,10 mM TEA-Cl, 0.5 mM $MgCl_2$, 5.5 mM

glucose, 5 mM CsCl, 20 mM $CaCl_2$, and 10 mM HEPES, adjusted to pH 7.4 with CsOH.

Micropipettes were filled with an internal solution containing 20 mM CsCl, 87 mM Aspartic

acid, 1 mM $MgCl_2$, 10 mM HEPES, 5 mM MgATP, and 10 mM EGTA adjusted to pH 7.2

using CsOH. Current-voltage relationships were obtained by exposing cells to a series of

300 ms depolarizing pulses from a holding potential of -70 mV to test potentials ranging

from -70 to +60 mV. A voltage error of 9.4 mV due to the liquid junction potential of the

recording solutions was corrected offline. Voltage dependence of Ca^{2+} channel activation

(G/G_{max}) was obtained from the resultant currents by converting them to conductance via

the equation G = I_{Ca}/(test potential – reversal potential of I_{Ca}); normalized G/G_{max} was

plotted as a function of test potential. Time constants of activation and inactivation of I_{Ca}

were fitted with a single exponential function.

I_{Kv} recordings were performed in the whole-cell configuration with myocytes exposed to

an external solution containing 130 mM NaCl, 5 mM KCl, 3 mM $MgCl_2$, 10 mM Glucose,

and 10 mM HEPES adjusted to 7.4 using NaOH. The internal pipette solution constituted

of 87 mM K-Aspartate, 20 mM KCl, 1 mM $CaCl_2$, 1 mM $MgCl_2$, 5 mM MgATP, 10 mM

EGTA, and 10 mM HEPES adjusted to 7.2 by KOH. A resultant liquid junction potential

of 12.7 mV from these solutions was corrected offline. To obtain current-voltage

relationships cells were subjected to a series of 500 ms test pulses increasing from -70

to +70 mV. To isolate the different K^+ channels attributed to composite I_K, cells were first

bathed in external I_K solution, subsequently exposed to 100 nM Iberiotoxin (Alomone,

Jerusalem, Israel) to eliminate any BK_{Ca} channel activity and finally immersed in an

external solution containing both 100 nM Iberiotoxin and 100 nM Stromatoxin (Alomone,

Jerusalem, Israel) to block both BK_{Ca} and $K_V2.1$ activity. Ionic current was converted to

conductance via the equation $G = I(V-E_K)$. E_K was calculated to be -78 mV. Activation

time constants for $K_V2.1$ currents were obtained by fitting the rising phase of these

currents with a single exponential function.

BK_{ca}-mediated spontaneous transient outward currents (STOCs) and membrane

potential were recorded using the perforated whole-cell configuration. To measure both,

myocytes were continuously exposed to a bath solution consisting of 130 mM NaCl, 5

mM KCl, 2 mM $CaCl_2$, 1 mM $MgCl_2$, 10 mM glucose, and 10 mM HEPES, pH adjusted to

7.4 with NaOH. Pipettes were filled with an internal solution containing 110 mM K-

aspartate, 30 mM KCl, 10 mM NaCl, 1 mM $MgCl_2$, 0.5 mM EGTA, and 10 mM HEPES

adjusted to a pH of 7.3 with KOH. The internal solution was supplemented with 250 µg/ml

amphotericin B (Sigma, St. Louis, MO). STOCs were measured in the voltage-clamp

mode and were analyzed with the threshold detection algorithm in Clampfit 10 (Axon

Instruments, Inc). Membrane potential was measured using the current clamp mode.

STOCs were recorded using the perforated whole-cell configuration. The composition of

the external bath solution consisted of 134 mM NaCl, 6 mM KCl, 1 mM $MgCl_2$, 2 mM

$CaCl_2$, 10 mM glucose, and 10 mM HEPES adjusted to a pH of 7.4 using NaOH. Pipettes

were filled with an internal solution of 110 mM K-aspartate, 10 mM NaCL, 30 mM KCl, 1

mM $MgCl_2$, 160 µg/ml amphotericin B, and 10 mM HEPES using NaOH to adjust to pH to

7.2. Myocytes were sustained at a holding potential of -70 mV before being exposed to a

400 ms ramp protocol from -140 to +60 mV. A voltage error of 12.8 mV resulting from the

liquid junction potential was corrected for offline. K_{ir} channels were blocked using 100 µM

Ba^{2+}.

1.4 Statistics

Data are expressed as mean ± SEM. All data sets were tested for normality. Normally distributed data were analyzed using T-tests or one-way analysis of variance (ANOVA). ANOVA analyses were followed by multiple comparison tests (i.e. Tuckey). $P<0.05$ was considered statistically significant.

Section 2. Computational modeling and simulation

2.1 Cell size and structure

The mean capacitance of the cells was experimentally calculated to be 16±3 pF based on all the male and female WT mesenteric C57BL/6J cells utilized in the experiments (N=45). Assuming the cells are roughly cylindrical in shape, the expected radius should be 2.485 μm and a length of 100 μm leading to a surface area of 1.6×10^{-5} cm^2 and a total volume of approximately 1.94×10^{-12} liters. The cell capacitance of excitable membranes is assumed to be 1.0×10^{-6} F/cm^2, with the calculated surface area the estimated total cell capacitance is $C_m = 16$ pF.

Because the total cell volume is roughly 2×10^{-12} liters, it is assumed that 50% of the total cell volume is occupied by organelles. There are three main compartments in vascular myocytes important to the regulation of membrane potential and calcium signaling: the cytosol, sarcoplasmic reticulum (SR), and specialized junctional domains formed by the SR and the plasma membrane. The cytosol occupies approximately 50% of total cell volume ($V_{cyt} = 1.0 \times 10^{-12}$ L). The sarcoplasmic reticulum occupies approximately 5% of cell cytosolic volume ($V_{SR} = 5.0 \times 10^{-14}$ L), and the junctional domain volume is approximately 1% of the cytosol volume ($V_{Jun} = 0.5 \times 10^{-14}$ L)[22,37–39].

2.2 Model development

The male and female *in silico* models are single whole-cell models based on the electrophysiology of isolated mesenteric vascular smooth muscle myocytes. A schematic of the proposed model is shown in **Figure 1**. The membrane electrophysiology can be described by the differential equation:

(22) $\dfrac{dV}{dt} = \dfrac{-I_{ion}}{C_m}$

Where V is voltage, t is time, C_m is membrane capacitance I_{ion} is the sum of

transmembrane currents. The contribution of each transmembrane current to the total

transmembrane ionic current can be described by the following equation:

(23) $I_{ion} = (I_{Kv1.5} + I_{Kv2.1} + I_{BKCa} + I_{K,b} + I_{Cav1.2} + I_{PMCA} + I_{Ca,b} + I_{NCX} + I_{NSC} + I_{NaK} + I_{Na,b})$

The eleven transmembrane currents are generated by ion channels, pumps, and

transporters. Currents from ion channels include the voltage-gated L-type calcium current

(I_{Ca}), the nonselective cation current (I_{NSC}), voltage-gated potassium currents ($I_{Kv1.5}$ and

$I_{Kv2.1}$), and the large-conductance Ca^{2+}-sensitive potassium current (I_{BKCa}). Additionally,

there are three background or leak currents ($I_{K,b}$, $I_{Ca,b}$, and $I_{Na,b}$). Currents from pumps

and transporters include the sodium-potassium pump current (I_{NaK}), plasma membrane

Ca-ATPase transport current (I_{PMCA}), and sodium-calcium exchanger current (I_{NCX}).

Cytosolic concentrations of sodium and potassium as a function of time are determined

by considering the sum of their respective fluxes into the cytosol.

(24) $\dfrac{d[K^+]_{cyt}}{dt} = -\dfrac{(I_{Kv2.1} + I_{Kv1.5} + I_{BKca} + I_{K,b} - 2I_{NaK} + I_{NSC-K})}{z_K Vol_{cyt} F}$

(25) $\dfrac{d[Na^+]_{cyt}}{dt} = -\dfrac{(3I_{NCX} + 3I_{NaK} + I_{Na,b} + I_{NSC-Na})}{z_{Na} Vol_{cyt} F}$

Where F is the Faraday's constant, Vol_{cyt} is the cytoplasmic volume and z_K and z_{Na} are

the valence of potassium and sodium ions, respectively.

The calcium dynamics is compartmentalized into three distinct regions: cytosol $[Ca^{2+}]_i$,

the sarcoplasmic reticulum $[Ca^{2+}]_{SR}$, and the junctional region $[Ca^{2+}]_{jun}$. The cytosol

includes a calcium buffer, which we assume can be described as a first-order dynamics

process.

Cytosolic calcium region ([Ca^{2+}]$_i$*):* calcium concentration in this region varies between

100-300 nM[13] and is mainly influenced by the following fluxes: transmembrane pumps

and transporters, the sarcoplasmic reticulum Ca-ATPase (J$_{SERCA}$), diffusion from the

junctional domain region (J$_{Jun-Cyt}$) and the calcium buffer calmodulin (BUF$_{CAM}$).

(26) $\frac{d[Ca^{2+}]_i}{dt} = -\frac{(I_{NCX}+I_{Cav1.2}+I_{Ca,b}-2I_{PMCA})}{z_{Ca}FV_{Cyt}} - J_{SERCA} + J_{jun-cyt} - \left(k_{BUF_{on}} * Ca_i * \right.$

$\left. (BUF_T - BUF_{CAM}) - k_{BUF_{off}} * BUF_{CAM} \right)$

Sarcoplasmic reticulum region ([Ca^{2+}]$_{SR}$): calcium concentration in this region varies

between 100-150 µM[40] and it is mainly influenced by the sarcoplasmic reticulum Ca-

ATPase (J$_{SERCA}$) and the flux from the ryanodine receptors (J$_{RyR}$).

(27) $\frac{d[Ca]_{SR}}{dt} = \left[\frac{Vol_{Cyt}}{Vol_{SR}}\right] J_{SERCA} - \left[\frac{Vol_{Cyt}}{Vol_{SR}}\right] [J_{Ryr}]$

Junctional region ([Ca^{2+}]$_{jun}$): calcium concentration in this region varies between 10-100

µM[37,38] and is mainly influenced by the flux from the ryanodine receptors (J$_{RyR}$), the

diffusion from the junctional region to the cytoplasm (J$_{Jun-Cyt}$)

(28) $\frac{d[Ca]_{jun}}{dt} = \left[\frac{Vol_{Cyt}}{Vol_{jun}}\right] J_{Ryr} - \left[\frac{Vol_{Cyt}}{Vol_{jun}}\right] J_{jun-cyt}$

In the model, the flux of J$_{SERCA}$ was adapted from the Luo-Rudy II model[41] and the flux

of J$_{RyR}$ was adapted from previous models of ryanodine receptors activation, originally

introduced in the field of cardiac electrophysiology[42–44].

2.3 Parameter optimization and reformulation of the gating ion channel models

The ionic current models of I$_{Ca}$, I$_{Kv2.1}$, and I$_{Kv1.5}$ were optimized using the approach

employed by Kernik *et al.*[45]. Here, the open probability P$_o$ of each voltage-dependent

gating variable "n" was defined by opening- and closing-rate voltage-dependent functions

α$_n$ and β$_n$ respectively, and were modeled by simple exponentials of the form:

(29) $\alpha_n(V) = x_1 e^{\left(\frac{V}{x_2}\right)}$

(30) $\beta_n(V) = x_3 e^{\left(\frac{V}{x_4}\right)}$

(31) $\tau_n(V) = \frac{1}{\alpha_n(V) + \beta_n(V)} + x_5$

The steady-state availability remains the same as the classical Hodgkin-Huxley

formulations and the time constant values follow a modified version formulation by

accommodating an extra parameter x_5 in equation 31. $(x_1, x_2, x_3, x_4, x_5)$ are parameters

to be optimized using experimental data. We used the parameter optimization employed

by Kernik et al.[45], which minimizes the error between model and experimental data using

the Nelder–Mead minimization of the error function. Random small perturbations (<10%)

were applied to find local minima, to improve data fit. The parameter fit with the minimal

error function value after 1000 to 10000 perturbations was used as the optimal model fit

to the data.

2.4 Cellular simulations with noise

The simulations encapsulate the cumulative effect of stochastic ion channel activity on

cell voltage dynamics through the fluctuating current term, ξ(t), into the membrane

potential (dV/dt) equation[46], as shown in equation 32. Here it is assumed ξ(t) is only a

function of time and it is implemented as Gaussian white noise[47].

(32) $\frac{dV}{dt} = -\frac{I_{total}(V)}{C_m} + \sigma\, \xi(t)$

We use the Euler-Maruyama numerical method for updating equation 33 and 34 as

follows.

(33) $V(t + \Delta t) = V(t) - \frac{I(V(t))}{C_m} \Delta t + \sigma * \text{randN} * \sqrt{\Delta t}$

(34) $\frac{d[Ca]_{SR}}{dt} = \left[\frac{Vol_{Cyt}}{Vol_{SR}}\right] J_{SERCA} - \left[\frac{Vol_{Cyt}}{Vol_{SR}}\right] [J_{Ryr}] + \sigma * \text{randN} * \sqrt{\Delta t}$

Where randN is a random number from a normal distribution (N(0,1)) with mean 0 and

variance 1. Δt is the time step and σ is the "diffusion coefficient", which represents the

amplitude of the noise. The numerical method for updating the voltage was forward Euler.

1000

2.5 One-dimensional simulations

The idealized one-dimensional representation of a vessel was developed by connecting

400 Hernandez-Hernandez model cells in series via simulated resistances to represent

gap junctions. For each cell in the cable, the Hernandez-Hernandez model was used to

compute ionic currents and concentration changes. The temporal transmembrane fluxes

of the Hernandez-Hernandez model are related to the spatial or current flow by a finite

difference approximation of the cable equation[48–50]

1008

$(35) \left[C_m \left(\frac{V_i^{(t+1)} - V_i^t}{\Delta t} \right) + I_{ion} + I_{stim} \right] = \frac{a}{4 \left(R_{myo} + \frac{R_g}{\Delta x} \right)} \frac{(V_{i-1}^t - 2V_{i-1}^t + V_{i-1}^t)}{\Delta x \Delta x}$

where I_{ion} represents the individual membrane ionic current densities (pA/pF) of the

Hernandez-Hernandez model, I_{stim} is the stimulus current density (pA/pF) set to zero in

our simulation, a is the radius of the fiber (5 µm), C_m is the membrane capacity (pA/pF),

V_{it} is membrane potential at segment i and time t, Δx is the discretization element (100

µm = 0.01 cm). Where R_{myo} is the myoplasmic resistance (R_{myo}=150 Ωcm) and R_g is the

gap junction resistance (R_g=71.4 Ωcm^2).

1016
2.6 Sensitivity Analysis

The baseline models in male and female vascular smooth muscle cells were analyzed

through a parameter sensitivity assessment using multivariable linear regression,

following the methodology introduced by Sobie[51]. The scope of the sensitivity analysis

encompassed variations in the maximal conductance and maximal ion transport rates of

the transmembrane currents, including $I_{Kv1.5}$, $I_{Kv2.1}$, I_{BKCa}, $I_{K,b}$, $I_{Cav1.2}$, I_{PMCA}, $I_{Ca,b}$, I_{NCX} ,

I_{NSC}, I_{NaK}, and $I_{Na,b}$. All other parameters, notably those defining model kinetics, remained

constant at the values established by the foundational model. Scaling factors were

randomly selected from a log-normal distribution characterized by a median value of 1

and a standard deviation of 0.1[45].

1027

2.7 Simulation protocols

Code for simulations and analysis was written in C++ and MATLAB 2018a. The single

vascular smooth muscle code was run on an Apple Mac Pro machine with two 2.7 GHz

12-Core Intel Xeon processors and an HP ProLiant DL585 G7 server with a 2.7 GHz 28-

core AMD Opteron processor. Vessel simulations were implemented in C++ and

parallelized using OpenMP. The C++ code was compiled with the Intel ICC compiler,

version 18.0.3. Numerical results were visualized using MATLAB R2018a by The

MathWorks, Inc. All codes and detailed model equations are available on GitHub

(https://github.com/ClancyLabUCD/sex-specific-responses-to-calcium-channel-blockers-

in-mesenteric-vascular-smooth-muscle)

1038

Acknowledgments

This work was supported by National Institutes of Health Common Fund Grant
OT2OD026580 (to C.E.C., L.F.S.), National Heart, Lung, and Blood Institute grants
R01HL128537 (C.E.C, L.F.S.,), R01HL152681 (to C.E.C., L.F.S.). UC Davis Department
of Physiology and Membrane Biology Research Partnership Fund (to C.E.C.) as well as
UC Davis T32 Predoctoral and Postdoctoral Training in Basic and Translational
Cardiovascular Medicine fellowship supported in part by NHLBI Institutional Training
Grant T32HL086350 (to C.M., G.H.H.)

1047

1048
1049
1050

References

1051   **References**
1052
1. Jacobsen, J. C. B. *et al.* Activation of a cGMP-sensitive calcium-dependent chloride
channel may cause transition from calcium waves to whole cell oscillations in smooth
muscle cells. *Am. J. Physiol. Heart Circ. Physiol.* **293**, H215-228 (2007).
2. Kapela, A., Bezerianos, A. & Tsoukias, N. M. A mathematical model of Ca2+
dynamics in rat mesenteric smooth muscle cell: agonist and NO stimulation. *J. Theor.*
Biol. **253**, 238–260 (2008).
3. Yang, J., Clark, J. W., Bryan, R. M. & Robertson, C. The myogenic response in
isolated rat cerebrovascular arteries: smooth muscle cell model. *Med. Eng. Phys.* **25**,
691–709 (2003).
4. Parthimos, D., Edwards, D. H. & Griffith, T. M. Minimal model of arterial chaos
generated by coupled intracellular and membrane Ca2+ oscillators. *Am. J. Physiol.*
277, H1119-1144 (1999).
5. Karlin, A. Membrane potential and Ca2+ concentration dependence on pressure and
vasoactive agents in arterial smooth muscle: A model. *J. Gen. Physiol.* **146**, 79–96
(2015).
6. Bayliss, W. M. On the local reactions of the arterial wall to changes of internal
pressure. *J. Physiol.* **28**, 220–231 (1902).
7. Welsh, D. G., Morielli, A. D., Nelson, M. T. & Brayden, J. E. Transient Receptor
Potential Channels Regulate Myogenic Tone of Resistance Arteries. *Circ. Res.* **90**,
248–250 (2002).
8. Spassova, M. A., Hewavitharana, T., Xu, W., Soboloff, J. & Gill, D. L. A common
mechanism underlies stretch activation and receptor activation of TRPC6 channels.
Proc. Natl. Acad. Sci. U. S. A. **103**, 16586–16591 (2006).
9. Earley, S., Waldron, B. J. & Brayden, J. E. Critical Role for Transient Receptor
Potential Channel TRPM4 in Myogenic Constriction of Cerebral Arteries. *Circ. Res.*
95, 922–929 (2004).
10. Earley, S., Straub, S. V. & Brayden, J. E. Protein kinase C regulates vascular
myogenic tone through activation of TRPM4. *Am. J. Physiol. Heart Circ. Physiol.* **292**,
H2613-2622 (2007).
11.   Sharif-Naeini, R. *et al.* Polycystin-1 and -2 Dosage Regulates Pressure Sensing.
Cell **139**, 587–596 (2009).
12. Bulley, S. *et al.* Arterial smooth muscle cell PKD2 (TRPP1) channels regulate
systemic blood pressure. *eLife* **7**, e42628 (2018).
13. Moosmang, S. *et al.* Dominant role of smooth muscle L-type calcium channel
Cav1.2 for blood pressure regulation. *EMBO J.* **22**, 6027–6034 (2003).
14. Knot, H. J. & Nelson, M. T. Regulation of arterial diameter and wall [Ca2+] in
cerebral arteries of rat by membrane potential and intravascular pressure. *J. Physiol.*
508, 199–209 (1998).
15. Navedo, M. F. & Santana, L. F. CaV1.2 sparklets in heart and vascular smooth
muscle. *J. Mol. Cell. Cardiol.* **58**, 67–76 (2013).
16. Navedo, M. F., Amberg, G. C., Votaw, V. S. & Santana, L. F. Constitutively active L-
type Ca2+ channels. *Proc. Natl. Acad. Sci.* **102**, 11112–11117 (2005).

17. Navedo, M. F., Amberg, G. C., Nieves, M., Molkentin, J. D. & Santana, L. F. Mechanisms Underlying Heterogeneous Ca2+ Sparklet Activity in Arterial Smooth Muscle. *J. Gen. Physiol.* **127**, 611–622 (2006).

18. Amberg, G. C., Navedo, M. F., Nieves-Cintrón, M., Molkentin, J. D. & Santana, L. F. Calcium sparklets regulate local and global calcium in murine arterial smooth muscle. *J. Physiol.* **579**, 187–201 (2007).

19. Nelson, M. T., Patlak, J. B., Worley, J. F. & Standen, N. B. Calcium channels, potassium channels, and voltage dependence of arterial smooth muscle tone. *Am. J. Physiol.* **259**, C3-18 (1990).

20. O'Dwyer, S. C. *et al.* Kv2.1 channels play opposing roles in regulating membrane potential, Ca2+ channel function, and myogenic tone in arterial smooth muscle. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 3858–3866 (2020).

21. Amberg, G. C. & Santana, L. F. Kv2 channels oppose myogenic constriction of rat cerebral arteries. *Am. J. Physiol. Cell Physiol.* **291**, C348-356 (2006).

22. Nelson, M. T. *et al.* Relaxation of arterial smooth muscle by calcium sparks. *Science* **270**, 633–637 (1995).

23. Plane, F. *et al.* Heteromultimeric Kv1 channels contribute to myogenic control of arterial diameter. *Circ. Res.* **96**, 216–224 (2005).

24. Sato, D. *et al.* A stochastic model of ion channel cluster formation in the plasma membrane. *J. Gen. Physiol.* **151**, 1116–1134 (2019).

25. Ledoux, J., Werner, M. E., Brayden, J. E. & Nelson, M. T. Calcium-Activated Potassium Channels and the Regulation of Vascular Tone. *Physiology* **21**, 69–78 (2006).

26. Brayden, J. E. & Nelson, M. T. Regulation of Arterial Tone by Activation of Calcium-Dependent Potassium Channels. *Science* **256**, 532–535 (1992).

27. Jaggar, J. H., Stevenson, A. S. & Nelson, M. T. Voltage dependence of Ca2+sparks in intact cerebral arteries. *Am. J. Physiol.-Cell Physiol.* **274**, C1755–C1761 (1998).

28. Knot, H. J., Standen, N. B. & Nelson, M. T. Ryanodine receptors regulate arterial diameter and wall [Ca2+] in cerebral arteries of rat via Ca2+-dependent K+ channels. *J. Physiol.* **508**, 211–221 (1998).

29. Pucovský, V. & Bolton, T. B. Localisation, function and composition of primary Ca(2+) spark discharge region in isolated smooth muscle cells from guinea-pig mesenteric arteries. *Cell Calcium* **39**, 113–129 (2006).

30. Yuan, X. J., Goldman, W. F., Tod, M. L., Rubin, L. J. & Blaustein, M. P. Ionic currents in rat pulmonary and mesenteric arterial myocytes in primary culture and subculture. *Am. J. Physiol.* **264**, L107-115 (1993).

31. Jaggar *et al.* Ca 2+ channels, ryanodine receptors and Ca 2+ -activated K + channels: a functional unit for regulating arterial tone: Ca2+ sparks in vascular smooth mucsle. *Acta Physiol. Scand.* **164**, 577–587 (1998).

32. Wang, S.-Q. *et al.* Imaging Microdomain Ca2+ in Muscle Cells. *Circ. Res.* **94**, 1011–1022 (2004).

33. Zhong, X. Z. *et al.* Stromatoxin-sensitive, heteromultimeric Kv2.1/Kv9.3 channels contribute to myogenic control of cerebral arterial diameter. *J. Physiol.* **588**, 4519–4537 (2010).

34. Archer, S. L. *et al.* Molecular identification of the role of voltage-gated K+ channels, Kv1.5 and Kv2.1, in hypoxic pulmonary vasoconstriction and control of resting

membrane potential in rat pulmonary artery myocytes. *J. Clin. Invest.* **101**, 2319–
2330 (1998).

35. Lu, Y., Hanna, S. T., Tang, G. & Wang, R. Contributions of Kv1.2, Kv1.5 and Kv2.1
subunits to the native delayed rectifier K(+) current in rat mesenteric artery smooth
muscle cells. *Life Sci.* **71**, 1465–1473 (2002).

36. Amberg, G. C., Rossow, C. F., Navedo, M. F. & Santana, L. F. NFATc3 regulates
Kv2.1 expression in arterial smooth muscle. *J. Biol. Chem.* **279**, 47326–47334
(2004).

37. Pérez, G. J., Bonev, A. D. & Nelson, M. T. Micromolar Ca2+ from sparks activates
Ca2+-sensitive K+ channels in rat cerebral artery smooth muscle. *Am. J. Physiol.-
Cell Physiol.* **281**, C1769–C1775 (2001).

38. Pérez, G. J., Bonev, A. D., Patlak, J. B. & Nelson, M. T. Functional coupling of
ryanodine receptors to KCa channels in smooth muscle cells from rat cerebral
arteries. *J. Gen. Physiol.* **113**, 229–238 (1999).

39. Kaßmann, M. *et al.* Role of Ryanodine Type 2 Receptors in Elementary Ca2+
Signaling in Arteries and Vascular Adaptive Responses. *J. Am. Heart Assoc.* **8**,
e010090 (2019).

40. ZhuGe, R. *et al.* The Influence of Sarcoplasmic Reticulum Ca2+ Concentration on
Ca2+ Sparks and Spontaneous Transient Outward Currents in Single Smooth Muscle
Cells. *J. Gen. Physiol.* **113**, 215–228 (1999).

41. Luo, C. H. & Rudy, Y. A dynamic model of the cardiac ventricular action potential. I.
Simulations of ionic currents and concentration changes. *Circ. Res.* **74**, 1071–1096
(1994).

42. Hernandez-Hernandez, G., Alvarez-Lacalle, E. & Shiferaw, Y. Role of connectivity
and fluctuations in the nucleation of calcium waves in cardiac cells. *Phys. Rev. E* **92**,
052715 (2015).

43. Greene, D. & Shiferaw, Y. Mechanistic link between CaM-RyR2 interactions and the
genesis of cardiac arrhythmia. *Biophys. J.* **120**, 1469–1482 (2021).

44. Shiferaw, Y., Watanabe, M. A., Garfinkel, A., Weiss, J. N. & Karma, A. Model of
Intracellular Calcium Cycling in Ventricular Myocytes. *Biophys. J.* **85**, 3666–3686
(2003).

45. Kernik, D. C. *et al.* A computational model of induced pluripotent stem-cell derived
cardiomyocytes incorporating experimental variability from multiple data sources. *J.*
Physiol. **597**, 4533–4564 (2019).

46. Goldwyn, J. H. & Shea-Brown, E. The What and Where of Adding Channel Noise to
the Hodgkin-Huxley Equations. *PLOS Comput. Biol.* **7**, e1002247 (2011).

47. Tanskanen, A. J. & Alvarez, L. H. R. Voltage noise influences action potential
duration in cardiac myocytes. *Math. Biosci.* **208**, 125–146 (2007).

48. Shaw, R. M. & Rudy, Y. Ionic Mechanisms of Propagation in Cardiac Tissue. *Circ.*
Res. **81**, 727–741 (1997).

49. Jack, J., Noble, D. & Tsien, R. Electric Current Flow In Excitable Cells. *Clarendon*
Press Oxf. (1975).

50. Hodgkin, A. L., Rushton, W. A. H. & Adrian, E. D. The electrical constants of a
crustacean nerve fibre. *Proc. R. Soc. Lond. Ser. B - Biol. Sci.* **133**, 444–479 (1997).

51. Sobie, E. A. Parameter Sensitivity Analysis in Electrophysiological Models Using
Multivariable Regression. *Biophys. J.* **96**, 1264–1274 (2009).

52. Hill-Eubanks, D. C., Werner, M. E., Heppner, T. J. & Nelson, M. T. Calcium
signaling in smooth muscle. *Cold Spring Harb. Perspect. Biol.* **3**, a004549 (2011).
53. Fleischmann, B. K., Murray, R. K. & Kotlikoff, M. I. Voltage window for sustained
elevation of cytosolic calcium in smooth muscle cells. *Proc. Natl. Acad. Sci. U. S. A.*
91, 11914–11918 (1994).
54. Bao, L. & Cox, D. H. Gating and ionic currents reveal how the BKCa channel's Ca2+
sensitivity is enhanced by its beta1 subunit. *J. Gen. Physiol.* **126**, 393–412 (2005).
55. Brenner, R. *et al.* Vasoregulation by the β1 subunit of the calcium-activated
potassium channel. *Nature* **407**, 870–876 (2000).
56. Tong, W.-C. *et al.* A computational model of the ionic currents, Ca2+ dynamics and
action potentials underlying contraction of isolated uterine smooth muscle. *PloS One*
6, e18685 (2011).
57. Jaggar, J. H., Porter, V. A., Lederer, W. J. & Nelson, M. T. Calcium sparks in
smooth muscle. *Am. J. Physiol. Cell Physiol.* **278**, C235-256 (2000).
58. Zhuge, R., Fogarty, K. E., Tuft, R. A. & Walsh, J. V. Spontaneous transient outward
currents arise from microdomains where BK channels are exposed to a mean Ca(2+)
concentration on the order of 10 microM during a Ca(2+) spark. *J. Gen. Physiol.* **120**,
15–27 (2002).
59. Dwyer, L. *et al.* Basally activated nonselective cation currents regulate the resting
membrane potential in human and monkey colonic smooth muscle. *Am. J. Physiol.*
Gastrointest. Liver Physiol. **301**, G287-296 (2011).
60. Setoguchi, M., Ohya, Y., Abe, I. & Fujishima, M. Stretch-activated whole-cell
currents in smooth muscle cells from mesenteric resistance artery of guinea-pig. *J.*
Physiol. **501 (Pt 2)**, 343–353 (1997).
61. Nakamura, Y., Ohya, Y., Abe, I. & Fujishima, M. Sodium-potassium pump current in
smooth muscle cells from mesenteric resistance arteries of the guinea-pig. *J. Physiol.*
519 Pt 1, 203–212 (1999).
62. ten Tusscher, K. H. W. J., Noble, D., Noble, P. J. & Panfilov, A. V. A model for
human ventricular tissue. *Am. J. Physiol. Heart Circ. Physiol.* **286**, H1573-1589
(2004).
63. Kargacin, G. J. Calcium signaling in restricted diffusion spaces. *Biophys. J.* **67**, 262–
272 (1994).
64. Désilets, M., Driska, S. P. & Baumgarten, C. M. Current fluctuations and oscillations
in smooth muscle cells from hog carotid artery. Role of the sarcoplasmic reticulum.
Circ. Res. **65**, 708–722 (1989).
65. Bychkov, R., Gollasch, M., Ried, C., Luft, F. C. & Haller, H. Regulation of
spontaneous transient outward potassium currents in human coronary arteries.
Circulation **95**, 503–510 (1997).
66. Bae, Y. M., Park, M. K., Lee, S. H., Ho, W. K. & Earm, Y. E. Contribution of Ca2+-
activated K+ channels and non-selective cation channels to membrane potential of
pulmonary arterial smooth muscle cells of the rabbit. *J. Physiol.* **514 (Pt 3)**, 747–758
(1999).
67. Sung, D. J. *et al.* Serotonin contracts the rat mesenteric artery by inhibiting 4-
aminopyridine-sensitive Kv channels via the 5-HT2A receptor and Src tyrosine
kinase. *Exp. Mol. Med.* **45**, e67 (2013).

68. Bratz, I. N., Swafford, A. N., Kanagy, N. L. & Dick, G. M. Reduced functional
expression of K(+) channels in vascular smooth muscle cells from rats made
hypertensive with N{omega}-nitro-L-arginine. *Am. J. Physiol. Heart Circ. Physiol.* **289**,
H1284-1290 (2005).
69. Rubart, M., Patlak, J. B. & Nelson, M. T. Ca2+ currents in cerebral artery smooth
muscle cells of rat at physiological Ca2+ concentrations. *J. Gen. Physiol.* **107**, 459–
472 (1996).
70. Davis, M. J., Meininger, G. A. & Zawieja, D. C. Stretch-induced increases in
intracellular calcium of isolated vascular smooth muscle cells. *Am. J. Physiol.-Heart*
Circ. Physiol. **263**, H1292–H1299 (1992).
71. Mulvany, M. J. & Aalkjaer, C. Structure and function of small arteries. *Physiol. Rev.*
70, 921–961 (1990).
72. Martinez-Lemus, L. A. The Dynamic Structure of Arterioles. *Basic Clin. Pharmacol.*
Toxicol. **110**, 5–11 (2012).
73. Heppner, D. B. & Plonsey, R. Simulation of electrical interaction of cardiac cells.
Biophys. J. **10**, 1057–1075 (1970).
74. Rudy, Y. & Quan, W. L. A model study of the effects of the discrete cellular structure
on electrical propagation in cardiac tissue. *Circ. Res.* **61**, 815–823 (1987).
75. Henriquez, C. S. & Plonsey, R. Effect of resistive discontinuities on waveshape and
velocity in a single cardiac fibre. *Med. Biol. Eng. Comput.* **25**, 428–438 (1987).
76. Yamamoto, Y., Klemm, M. F., Edwards, F. R. & Suzuki, H. Intercellular electrical
communication among smooth muscle and endothelial cells in guinea-pig mesenteric
arterioles. *J. Physiol.* **535**, 181–195 (2001).
77. Earley, S. & Brayden, J. E. Transient Receptor Potential Channels in the
Vasculature. *Physiol. Rev.* **95**, 645–690 (2015).
78. Guarina, L. *et al.* Biological noise is a key determinant of the reproducibility and
adaptability of cardiac pacemaking and EC coupling. *J. Gen. Physiol.* **154**,
e202012613 (2022).
79. Bae, Y. M. *et al.* Serotonin depolarizes the membrane potential in rat mesenteric
artery myocytes by decreasing voltage-gated K+ currents. *Biochem. Biophys. Res.*
Commun. **347**, 468–476 (2006).
80. Navedo, M. F. *et al.* Increased Coupled Gating of L-Type Ca2+ Channels During
Hypertension and Timothy Syndrome. *Circ. Res.* **106**, 748–756 (2010).
81. Diep, H. K., Vigmond, E. J., Segal, S. S. & Welsh, D. G. Defining electrical
communication in skeletal muscle resistance arteries: a computational approach. *J.*
Physiol. **568**, 267–281 (2005).
82. Earley, S., Resta, T. C. & Walker, B. R. Disruption of smooth muscle gap junctions
attenuates myogenic vasoconstriction of mesenteric resistance arteries. *Am. J.*
Physiol.-Heart Circ. Physiol. **287**, H2677–H2686 (2004).
83. Ueno, K. & Sato, H. Sex-related differences in pharmacokinetics and
pharmacodynamics of anti-hypertensive drugs. *Hypertens. Res. Off. J. Jpn. Soc.*
Hypertens. **35**, 245–250 (2012).
84. Kloner, R. A., Sowers, J. R., DiBona, G. F., Gaffney, M. & Wein, M. Sex- and age-
related antihypertensive effects of amlodipine. The Amlodipine Cardiovascular
Community Trial Study Group. *Am. J. Cardiol.* **77**, 713–722 (1996).

85. Kajiwara, A. *et al.* Younger females are at greater risk of vasodilation-related
adverse symptoms caused by dihydropyridine calcium channel blockers: results of a
study of 11,918 Japanese patients. *Clin. Drug Investig.* **34**, 431–435 (2014).
86. Yang, Y. *et al.* Function of BKCa channels is reduced in human vascular smooth
muscle cells from Han Chinese patients with hypertension. *Hypertens. Dallas Tex*
1979 **61**, 519–525 (2013).
87. Nystoriak, M. A. *et al.* Ser1928 phosphorylation by PKA stimulates the L-type Ca2+
channel CaV1.2 and vasoconstriction during acute hyperglycemia and diabetes. *Sci.*
Signal. **10**, eaaf9647 (2017).
88. Nieves-Cintrón, M. *et al.* Impaired BKCa channel function in native vascular smooth
muscle from humans with type 2 diabetes. *Sci. Rep.* **7**, 14058 (2017).

1288
1289
1290
1291 **FIGURE LEGENDS**
1292
1293
Figure 1. A schematic representation of the Hernandez-Hernandez model. The
components of the model include major ion channel currents shown in purple including
the voltage-gated L-type calcium current (I_{Ca}), nonselective cation current (I_{NSC}), voltage-
gated potassium currents ($I_{Kv1.5}$ and $I_{Kv2.1}$), and the large-conductance Ca^{2+}-sensitive
potassium current (I_{BKCa}). Currents from pumps and transporters are shown in red
including the sodium/potassium pump current (I_{NaK}), sodium/calcium exchanger current
(I_{NCX}), and plasma membrane ATPase current (I_{PMCA}). Leak currents are indicated in
green including the sodium leak current ($I_{Na,b}$), potassium leak current ($I_{K,b}$), and calcium
leak current ($I_{Ca,b}$). In addition, two currents in the sarcoplasmic reticulum are shown in
orange: the sarcoplasmic reticulum Ca-ATPase current (I_{SERCA}) and ryanodine receptor
current (I_{RyR}). Calcium compartments comprise three discrete regions including cytosol
($[Ca]_i$), sarcoplasmic reticulum ($[Ca]_{SR}$), and the junctional region ($[Ca]_{Jun}$). Red stars (*)
indicate measured sex-specific differences in ionic currents.
1307
Figure 2. Experimentally measured and modeled L-type calcium currents (I_{Ca}) from
male and female vascular smooth muscle cells. Properties of I_{Ca} are derived from
measurements in male and female vascular smooth muscle (VSM) cells isolated from the
mouse mesenteric arteries following voltage-clamp steps from -60 to 60 mV in 10 mV
steps from a -80 mV holding potential. Experimental data is shown in black circles for
male and blue squares for female. Model fits to experimental data are shown with black
solid lines for male and blue solid lines for female. (**A**) Male and female time constants of
I_{Ca} activation. (**B**) Male and female time constants of I_{Ca} inactivation. (**C**) Male and female
voltage-dependent steady-state activation and inactivation of I_{Ca}. (**D**) Current-voltage (I-
V) relationship of I_{Ca} from male and female vascular smooth muscle myocytes. *P < 0.05,
**P < 0.01, ***P<0.001. Error bars indicated mean ± SEM.
1319
Figure 3. Experimentally measured and modeled potassium currents (I_{KvTOT}) from
male and female vascular smooth muscle cells. The properties of $I_{Kv1.5}$ and $I_{Kv2.1}$ from
experimental measurements in male and female vascular smooth muscle cells isolated

from the mouse mesenteric arteries were recorded in response to voltage-clamp from -
60 to 40 mV in 10 mV steps (holding potential -80mV). Experimental data is shown as
black circles for male and blue squares for female. Model fits to experimental data are
shown with black solid lines for male and blue solid lines for female. (**A**) Male and female
voltage-dependent steady-state activation of $I_{Kv2.1}$. (**B**) Male and female time constants of
$I_{Kv2.1}$ activation. (**C**) Current-voltage (I-V) relationship of $I_{Kv2.1}$ from male and female
myocytes. (**D**) Male and female voltage-dependent steady-state activation of $I_{Kv1.5}$. (**E**)
Current-voltage (I-V) relationship of $I_{Kv1.5}$ from male and female myocytes. (**F**) Male and
female total voltage-gated potassium current $I_{KvTOT} = I_{Kv1.5} + I_{Kv2.1}$. (**G**) Predicted male
and female time constants of the $I_{Kv1.5}$ activation gate. (**H**) Table showing sex-specific
differences in conductance and steady-state total potassium current-voltage
dependence. *P < 0.05, **P < 0.01, ***P<0.001, ****P<0.0001. Data points without
asterisks are not significant. Error bars indicated mean ± SEM.
1336
Figure 4. Experimentally measured and modeled large-conductance Ca^{2+}-activated
K$^+$ currents ($I_{BK\alpha\beta1}$). The model was optimized to data from Bao and Cox (Bao & Cox,
2005). (**A**) Voltage-dependent activation of $I_{BK\alpha\beta1}$ from experiments performed with three
different [Ca]$_{Jun}$ concentrations (1 μM, 10 μM, 100 μM) shown in green circles is the data
from (Zhuge *et al.*, 2002) (**B**) Voltage-dependent activation time constants with
[Ca]$_{Jun}$=0.003 μM and simulations [Ca]$_{Jun}$=10 μM. (**C**) Simulated I-V curve at different
peak levels of [Ca]$_{Jun}$ levels. (**D**) Simulated I-V curve with different BK$_{ca}$ average cluster
sizes (N = 4,6, 8, and 10).
1345
Figure 5. Membrane potential from experiments and simulations in male and female
vascular smooth muscle myocytes. A) Whole-cell membrane potential recordings in
male and female myocytes showing spontaneous repeat transient hyperpolarization of
the membrane potential. **B**) Simulated whole-cell membrane potential with physiological
noise. **C**) Comparison of sensitivity analysis performed around the baseline membrane
potential in male and female models using multivariable regression.
1352
Figure 6. Differential effects of voltage-gated potassium current (I_{KVTOT}) block in
male and female myocytes. A) Simulated time course of male $I_{Kv2.1}$ (*top panel, solid*
traces) and $I_{Kv1.5}$ (*lower panel, dashed traces*) at three different baseline membrane
potentials (-45 mV *green*, -40 mV *blue*, and -35mV *black*). **B**) Simulated time course of
female $I_{Kv2.1}$ (*top panel solid traces*) and $I_{Kv1.5}$ (*lower panel, dashed traces*) at three
different baseline membrane potentials (-45 mV *light blue*, -40 mV *purple*, and -35mV
red). Current contribution to I_{KVTOT} from $K_V1.5$ (indicated by asterisks) and $K_V2.1$ in male
and female myocytes at a baseline membrane potential of -45 mV (**C**), -40 mV (**D**), and -
35 mV (**E**).
1362
Figure 7. Simulated L-type calcium currents (I_{Ca}) and calcium influx in male and
female vascular smooth muscle cells. A) Male and female whole-cell I_{Ca} membrane
potential relationship. (**B**) Male and female intracellular calcium concentration in the
cytosolic compartment at indicated membrane potential. **C**) Time course of membrane
potential in male vascular smooth muscle cells before (black) and after (gray) simulated
nifedipine application (*top panel*). Corresponding time course of L-type calcium current

I_{Ca} before (black) and after (gray) simulated nifedipine application (*middle panel*) and
intracellular calcium $[Ca^{2+}]_i$ concentration before (black) and after (gray) simulated
nifedipine application (*lower panel*). **D**) Time course of membrane potential in female
vascular smooth muscle cells before (blue) and after (pink) simulated nifedipine
application (*top panel*). Corresponding time course of L-type calcium current I_{Ca} before
(blue) and after (pink) simulated nifedipine application (*middle panel*) and intracellular
calcium $[Ca^{2+}]_i$ concentration before (blue) and after (pink) simulated nifedipine
application (*lower panel*).

1377
1378

Figure 8. **A one-dimensional tissue model representation of vascular smooth**
muscle cells connected in series. A) Uncoupled female vessel simulation showing cell
1, cell 50, and cell 100 at a baseline membrane potential of -35 mV. **B**) Uncoupled male
vessel simulations showing cell 1, cell 50, and cell 100 at a baseline membrane potential
of -45 mV. **C**) Composite female (blue trace) and male (black trace) membrane potential
of 400 coupled smooth muscle cells connected with gap junctional resistance of 71.4
Ωcm^2 in a one-dimensional tissue representation. (**D**) Sharp-electrode records of the
membrane potential of smooth muscle in pressurized (80-mmHg) female and male
arteries from O'Dwyer *et al.*, 2020.

1388
1389

Figure 9. **Experimentally measured and modeled intracellular calcium [Ca]$_i$ in male**
and female vessels and response to clinically used L-type Ca^{2+} channel blocker. A)
Intracellular calcium [Ca]$_i$ in female (blue symbols) and male (black symbols) arteries at
intravascular pressures ranging from 20 to 120 mmHg. Simulations showing [Ca]$_i$ in the
idealized female and male vessels are shown with blue and black solid lines, respectively.
Simulated [Ca]$_i$ after the application of clinically used L-type Ca^{2+} channel blocker
nifedipine is shown with dashed lines for male (black) and female (blue). **B**) Comparison
of the percentage change of [Ca]$_i$ in male (black) and female (blue) after the application
L-type Ca^{2+} channel blocker nifedipine at 80 mmHg and 120 mmHg. *P < 0.05, **P < 0.01.

1399
1400
1401

FIGURES AND LEGENDS

Figure 1

Figure 1. A schematic representation of the Hernandez-Hernandez model. The components of the model include major ion channel currents shown in purple including the voltage-gated L-type calcium current (I_{Ca}), nonselective cation current (I_{NSC}), voltage-gated potassium currents ($I_{Kv1.5}$ and $I_{Kv2.1}$), and the large-conductance Ca^{2+}-sensitive potassium current (I_{BKCa}). Currents from pumps and transporters are shown in red including the sodium/potassium pump current (I_{NaK}), sodium/calcium exchanger current (I_{NCX}), and plasma membrane ATPase current (I_{PMCA}). Leak currents are indicated in green including the sodium leak current ($I_{Na,b}$), potassium leak current ($I_{K,b}$), and calcium leak current ($I_{Ca,b}$). In addition, two currents in the sarcoplasmic reticulum are shown in orange: the sarcoplasmic reticulum Ca-ATPase current (I_{SERCA}) and ryanodine receptor current (I_{RyR}). Calcium compartments comprise three discrete regions including cytosol ($[Ca]_i$), sarcoplasmic reticulum ($[Ca]_{SR}$), and the junctional region ($[Ca]_{Jun}$). Red stars (*) indicate measured sex-specific differences in ionic currents.

Figure 2. Experimentally measured and modeled L-type calcium currents (I_{Ca}) from male and female vascular smooth muscle cells. Properties of I_{Ca} are derived from measurements in male and female vascular smooth muscle (VSM) cells isolated from the mouse mesenteric arteries following voltage-clamp steps from -60 to 60 mV in 10 mV steps from a -80 mV holding potential. Experimental data is shown in black circles for male and blue squares for female. Model fits to experimental data are shown with black solid lines for male and blue solid lines for female. (**A**) Male and female time constants of I_{Ca} activation. (**B**) Male and female time constants of I_{Ca} inactivation. (**C**) Male and female voltage-dependent steady-state activation and inactivation of I_{Ca}. (**D**) Current-voltage (I-V) relationship of I_{Ca} from male and female vascular smooth muscle myocytes. *P < 0.05, **P < 0.01, ***P<0.001. Error bars indicated mean ± SEM.

Figure 3

Figure 3. Experimentally measured and modeled potassium currents (I_{KvTOT}) from male and female vascular smooth muscle cells. The properties of $I_{Kv1.5}$ and $I_{Kv2.1}$ from experimental measurements in male and female vascular smooth muscle cells isolated from the mouse mesenteric arteries were recorded in response to voltage-clamp from -60 to 40 mV in 10 mV steps (holding potential -80mV). Experimental data is shown as black circles for male and blue squares for female. Model fits to experimental data are shown with black solid lines for male and blue solid lines for female. (**A**) Male and female voltage-dependent steady-state activation of $I_{Kv2.1}$. (**B**) Male and female time constants of $I_{Kv2.1}$ activation. (**C**) Current-voltage (I-V) relationship of $I_{Kv2.1}$ from male and female myocytes. (**D**) Male and female voltage-dependent steady-state activation of $I_{Kv1.5}$. (**E**) Current-voltage (I-V) relationship of $I_{Kv1.5}$ from male and female myocytes. (**F**) Male and female total voltage-gated potassium current $I_{KvTOT} = I_{Kv1.5} + I_{Kv2.1}$. (**G**) Predicted male and female time constants of the $I_{Kv1.5}$ activation gate. (**H**) Table showing sex-specific differences in conductance and steady-state total potassium current-voltage dependence. *P < 0.05, **P < 0.01, ***P<0.001, ****P<0.0001. Data points without asterisks are not significant. Error bars indicated mean ± SEM.

Figure 4. Experimentally measured and modeled large-conductance Ca²⁺-activated K⁺ currents (I_BKαβ1). The model was optimized to data from Bao and Cox (Bao & Cox, 2005). (**A**) Voltage-dependent activation of $I_{BK\alpha\beta1}$ from experiments performed with three different $[Ca]_{Jun}$ concentrations (1 µM, 10 µM, 100 µM) shown in green circles is the data from (Zhuge *et al.*, 2002) (**B**) Voltage-dependent activation time constants with $[Ca]_{Jun}$=0.003 µM and simulations $[Ca]_{Jun}$=10 µM. (**C**) Simulated I-V curve at different peak levels of $[Ca]_{Jun}$ levels. (**D**) Simulated I-V curve with different BK_ca average cluster sizes (N = 4,6, 8, and 10).

Figure 5. Membrane potential from experiments and simulations in male and female vascular smooth muscle myocytes. A) Whole-cell membrane potential recordings in male and female myocytes showing spontaneous repeat transient hyperpolarization of the membrane potential. **B)** Simulated whole-cell membrane potential with physiological noise. **C)** Comparison of sensitivity analysis performed around the baseline membrane potential in male and female models using multivariable regression.

Figure 6. Differential effects of voltage-gated potassium current (I_{KVTOT}) block in male and female myocytes. A) Simulated time course of male $I_{Kv2.1}$ (*top panel, solid traces*) and $I_{Kv1.5}$ (*lower panel, dashed traces*) at three different baseline membrane potentials (-45 mV *green*, -40 mV *blue*, and -35mV *black*). **B**) Simulated time course of female $I_{Kv2.1}$ (*top panel solid traces*) and $I_{Kv1.5}$ (*lower panel, dashed traces*) at three different baseline membrane potentials (-45 mV *light blue*, -40 mV *purple*, and -35mV red). Current contribution to I_{KVTOT} from $K_V1.5$ (indicated by asterisks) and $K_V2.1$ in male and female myocytes at a baseline membrane potential of -45 mV (**C**), -40 mV (**D**), and -35 mV (**E**).

Figure 7. Simulated L-type calcium currents (I$_{Ca}$) and calcium influx in male and female vascular smooth muscle cells. A) Male and female whole-cell I$_{Ca}$ membrane potential relationship. (**B**) Male and female intracellular calcium concentration in the cytosolic compartment at indicated membrane potential. **C**) Time course of membrane potential in male vascular smooth muscle cells before (black) and after (gray) simulated nifedipine application (*top panel*). Corresponding time course of L-type calcium current I$_{Ca}$ before (black) and after (gray) simulated nifedipine application (*middle panel*) and intracellular calcium [Ca^{2+}]$_i$ concentration before (black) and after (gray) simulated nifedipine application (*lower panel*). **D**) Time course of membrane potential in female vascular smooth muscle cells before (blue) and after (pink) simulated nifedipine application (*top panel*). Corresponding time course of L-type calcium current I$_{Ca}$ before (blue) and after (pink) simulated nifedipine application (*middle panel*) and intracellular calcium [Ca^{2+}]$_i$ concentration before (blue) and after (pink) simulated nifedipine application (*lower panel*).

Figure 8. **A one-dimensional tissue model representation of vascular smooth muscle cells connected in series.** **A**) Uncoupled female vessel simulation showing cell 1, cell 50, and cell 100 at a baseline membrane potential of -35 mV. **B**) Uncoupled male vessel simulations showing cell 1, cell 50, and cell 100 at a baseline membrane potential of -45 mV. **C**) Composite female (blue trace) and male (black trace) membrane potential of 400 coupled smooth muscle cells connected with gap junctional resistance of 71.4 Ωcm² in a one-dimensional tissue representation. (**D**) Sharp-electrode records of the membrane potential of smooth muscle in pressurized (80-mmHg) female and male arteries from O'Dwyer *et al.*, 2020.

Figure 9. **Experimentally measured and modeled intracellular calcium [Ca]$_i$ in male and female vessels and response to clinically used L-type Ca^{2+} channel blocker**. **A**) Intracellular calcium [Ca]$_i$ in female (blue symbols) and male (black symbols) arteries at intravascular pressures ranging from 20 to 120 mmHg. Simulations showing [Ca]$_i$ in the idealized female and male vessels are shown with blue and black solid lines, respectively. Simulated [Ca]$_i$ after the application of clinically used L-type Ca^{2+} channel blocker nifedipine is shown with dashed lines for male (black) and female (blue). **B**) Comparison of the percentage change of [Ca]$_i$ in male (black) and female (blue) after the application L-type Ca^{2+} channel blocker nifedipine at 80 mmHg and 120 mmHg. *P < 0.05, **P < 0.01.