https://www.overleaf.com/project/63dca913e106bf69d519d00d
https://www.overleaf.com/project/63dca913e106bf69d519d00d

Endocannabinoid system modulation alters fine motor behavior in mice: insights from 3D motion capture

Bogna M. Ignatowska-Jankowska[1*], Aysen Gurkan Ozer[1], Alexander Kuck[1], Micah J. Niphakis[2,1], Daisuke Ogasawara[3], Benjamin F. Cravatt[3] and Marylka Y. Uusisaari[1*]

[1*]Neuronal Rhythms in Movement Unit, Okinawa Institute of Science and Technology, 1919-1 Tancha, Onna-son, 904-0495, Okinawa, Japan.
[2] Lundbeck La Jolla Research Center, San Diego, 92121, CA, USA.
[3]Department of Chemical Physiology, Scripps Research Institute, La Jolla, 92037, CA, USA.

*Corresponding author(s). E-mail(s):
bogna.ignatowska@gmail.com; uusisaari@oist.jp;
Contributing authors: aysen.gurkan@oist.jp;
alexander.kuck@web.de; MIIP@lundbeck.com;
daisuke@scripps.edu; cravatt@scripps.edu;

Abstract

The neuromodulatory endocannabinoid system is a promising target for therapeutic interventions. One of well-known behavioral effects of cannabinoid CB_1 receptor activation with exogenous ligands such as THC is inhibition of locomotor activity. However, behavioral effects of the endogenous cannabinoids have not been extensively studied. Despite broad medicinal and recreational use of cannabinoids, little is known about their effects on fine movements, critical for therapeutic applications. Such fine motor function is only resolvable with the precision of marker-based motion capture, which so far has not been available in freely moving mice.

Therefore, we have adapted a high-speed, high-resolution marker-based 3D motion capture system to track movement (3D trajectories and speed of markers) during voluntary locomotor tasks in mice. Here we show that enhancing endocannabinoid signaling produces bidirectional effects on locomotor behavior that differ from those caused by exogenous cannabinoid receptor agonist, in unrestricted mice. Given our expectation of subtle rather than dramatic changes with the low and potentially therapeutic doses, we employed a sensitive and accurate marker-based 3D motion capture system to resolve the finest differences in full-body kinematic signatures of mice exploring horizontal and vertical environments. We found that selective upregulation of either of the two main endocannabinoids 2-arachidonoylglycerol (2-AG) or anandamide (AEA) with selective inhibitors of their degradation (MJN110 and PF3845, respectively), produced opposite effects: PF3845 suppressed locomotor activity whereas MJN110 enhanced it. Furthermore, while low doses of the synthetic cannabinoid agonist CP55,940 decreased locomotion in a an open field exploration task as expected, it surprisingly did not affect performance in a vertical climbing task. The results show that the effects of cannabinoid signaling on behavior are not predominantly inhibitory as commonly assumed. Furthermore, we found that the invariant microstructure of locomotory behavior remains unchanged under all treatments, pointing towards motivational rather than motor-related mechanisms of action.

Keywords: mouse, behavior, mouse models, motion capture, marker-based, movement trajectories, fine motor function, kinematics, freely moving, cannabinoid, endocannabinoid

1 Introduction

Despite its precision, robustness and common use in humans [1–4], use of marker-based 3D motion capture in small animals such as mice and rats has been limited by technical difficulties [3, 5–11] and markerless animal tracking has become the main tool for assessing behavior (e.g. Deep Lab Cut, DeepEthogram, EthoLoop, MoSeq [12–18]). These approaches have recently experienced significant advances in flexibility and ease-of-use, but achieving precise trajectory tracking remains challenging. Even in human studies, markerless techniques show only "reasonable agreement" with marker-based methods even under optimal conditions [3–6, 19, 20].

Presently, accurate kinematic tracking of rodents largely rely on restricting animals' movements [8] or on creating artificial environments (such as transparent floors or locomotion in narrow confines) that may distort behavior and limit translational potential of the results. As the current translational crisis has highlighted the weakness of animal experimental designs [10, 21–23],

their improvement is now urgently needed so that they would more accurately
reflect the complexity of behavior in natural environments.

In nature, it is rare for mice to explore flat, smooth surfaces such as those
commonly used in the context of laboratory experiments. Instead, their natural
behavior involves locomoting predominantly on uneven terrain with complex
demands for body kinematics: they balance, they climb, and they jump and
swim. Reaching towards precise and accurate kinematic recordings of unre-
stricted mice moving freely in three dimensions, we modified a marker-based
3D motion capture (MoCap) system commonly used for human and other large
animal tracking (Qualisys; [24–26]) to examine how mouse locomotor activity
and kinematics were affected by exogenous modulation of the endocannabinoid
system.

Broad therapeutic potential of approaches that target endocannabinoid
signaling as well as expanding medicinal and recreational use of exogenous
cannabinoids [27–29] call for thorough investigation of systemic effects of endo-
cannabinoid system modulation. The high expression of cannabinoid receptors
in cerebellum and basal ganglia [30–32] suggests that careful assessment of sub-
tle motor effects of cannabinoid receptor activation will be necessary. Presently
the research on effects of cannabinoids on precise movement in rodents is,
however, limited [33–37] but cannabinoids have been shown to have effects on
eyeblink [38] and kinematics of gait in humans[39].

Here we aimed to assess whether upregulation of local endocannabinoid
signaling by selective inhibition of their degradation [40–43] leads to behavioral
outcomes different from those seen after systemic administration of exogenous
ligands that bind to all available receptors in the body.

To tease apart effects of inhibiting degradation of two endocannabinoids,
2-arachidonoylglycerol (2-AG) and anandamide (AEA), we examine the effects
of selective inhibitors of their degradation in freely-behaving mice tracked with
the marker-based MoCap system.

The results reveal distinct bidirectional effects for the two
endocannabinoid-enhancing treatments, that are also different from those
seen after exogenous cannabinoid agonist administration. Our observations
suggest fundamentally different mechanisms of actions of distinct cannabi-
noids and call for revision of the current view of overwhelmingly inhibitory
effects of cannabinoids on behavior. The work demonstrates the strength of
marker-based 3D MoCap as a powerful and sensitive tool for examining subtle
differences in rodent behavior.

2 Results

2.1 Implementation and validation of the 3D Motion
Capture system for rodents

Realization of marker-based 3D motion capture on small rodents poses several
technical challenges that have been difficult to overcome [1, 2].

Fig. 1 Motion Capture system adapted to mice allows for tracking movement trajectories with unprecedented precision. a. Schematic of experimental setup: mouse with retroreflective markers screwed on stainless steel implants (barbel piercings) at strategic locations on the skin surface, moving freely within 30 x 30 x 30 cm arena surrounded by 7 cameras. b. 3D reconstruction of marker locations using Qualisys motion capture system software. c. Photograph of the recording arena. d. Marker visibility determines tracking reliability. Left:fraction of frames in which markers are missing with 7 or 4 cameras (open field and climbing, respectively). Right: missing trajectory length for markers. Less than 100 frames are lost even in least visible markers. e. Marker pairs connected under the skin surface allow measurement of tracking accuracy based on marker location variability (distance between markers). Tracking accuracy falls between 0.15 mm and 0.5 mm for all markers (mean±SEM). f. Hip trajectories with steps marked during horizontal exploration by a representative animal. g. Step detection method. i. Manual step verification shows over 60% of steps are detected automatically with neglible amount of false positives.

First, skin-glued markers are of limited use in unrestricted rodents, as they attract the animal's attention and distort behavior. Moreover, markers made of soft materials will not survive the attention and require frequent replacements. While in restricted and strongly task-motivated animals (e.g. under food deprivation) such markers may survive sufficiently long for reliable tracking [8], our preliminary trials revealed that examining whole-body kinematics in freely-moving animals was impossible with skin-glued, plastic markers. To address this problem we established a method for permanent under-skin implantation of stainless-steel piercings at landmark locatins, to which above-skin retroreflective markers are robustly attached (Fig. 1a, Suppl. Fig. 1 and 2, supplementary methods for details). After implantation of these markers at landmark locations, mice are habituated to them for two weeks prior to recordings. Implants do not cause noticeable discomfort to the animals, and they can wear them for over a year allowing multiple within-subject

tests. The under-skin implants are tightly attached into the connective tissue and thus their movement is less confounded by the loose rodent skin that obscures large ranges of joint movement.

Second, spurious reflections and distortions caused by any transparent surfaces between the camera and markers catastrophically degrade tracking performance that can only be compensated for by significant increase of tracking cameras. To allow reliable tracking with minimal number of cameras (6-7, placed at 60 degree angles around the recording arena in case of free horizontal exploration) we developed 2-week training protocol for the mice to learn not to leave the designated "arena" in the middle of the cameras despite absence of any physical barriers (Fig. 1a-c; Supp. Video 1). All animals used in the study learned this behavior without problems, allowing us to record their behavior not only on a horizontal, textured 30 x 30 cm surface (the "open field") but also several other settings. In this article we present the analysis of data of two tasks: a) exploration of the open field and b) vertical climbing behavior (Fig. 1c).

2.2 Performance of the marker-based mouse motion capture system

A key question in assessing the potential and limitations of a novel recording method relates to the reliability and accuracy of the measurements. Here, we quantify them in terms of marker visibility (fraction of frames in a recording where a marker is visible or occluded) and location accuracy (how far is the recorded location from real position of a marker). Similarly to the approach taken in previous work [7], we calculated how many markers are occluded in each frame. As shown in Fig. 1d, left, vast majority of the frames had no markers missing and around 90% of frames had not more than one marker missing. Notably, the tracking quality was high even in climbing trials that were recorded with only 4 cameras.

When assessing the quality of tracking, the durations of occasional occlusions are of great significance as short gaps are trivial to be interpolated over while longer gaps require sophisticated approaches such as employment of machine learning and skeleton model fitting processing. As shown in Fig. 1d, right, most of the gaps terminated within 100 frames (corresponding to 0.3 s with 300 fps recording). Even for the most challenging ankle markers that by necessity are fully occluded when the animal is in an idle, inactive position, the occlusions were on average resolved within 50 frames (0.15 s). We also could confirm that there were no differences in marker tracking performance for left and right sides of the animals, evidencing there were no biases in how the cameras were positioned with respect to the behavioral arena configuration (Student's paired t-test for left and right legs for 30 1-minute-long recordings, $p = 0.7$).

The accuracy of triangulation-based 3D tracking depends critically on the uniformity of marker shape and its projection into the camera image plane. To assess the performance of our system in this respect, we took advantage of the

fact that the visible marker pairs reside at the ends of metal rods of fixed and
known length, and the deviation of a marker's tracked position relative to it's
pair can be considered as its error. Fig. 1e shows that the distance between
paired markers varied on average no more than 0.5 mm even for the body
parts most occluded by hair (shoulder blades); notably, assuming independent
errors for markers, leg marker distance deviations of around 0.2 mm indicates
that a tracked position of a single marker could be expected to be within 100
micrometers of its real position.

For step detection (Fig. 1f,g), we considered that a swing consists of a
period of high-velocity ankle movement bordered with rapid acceleration and
deceleration events. Fig. 1g depicts schematically how speed (red trace) and
acceleration (green trace) relate to the vertical position of the ankle (blue trace)
and determine our "swing" segment. To validate this approach and define its
limitations, we compared the number of steps detected with those identified in
recordings by an experienced observer (BIJ). As shown in Fig. 1h (left), in 30
60-s trials mice took on average 92.9 ± 8.4 steps, out of which 62.1 ± 6.4 steps
were detected by the algorithm. Taking a step further in detailing the reliabil-
ity, we chose 3 animals representing the animals with average, highest (best)
and lowest (worst) percentage of steps automatically detected and determined
the number of missed steps and false positives in these samples (Fig. 1h, right,
Suppl. Fig. 3). In this representative set of tracked steps, on average 81 out of
108 steps were correctly detected; importantly, false positives were below 1%,
attesting to the validity of our step-detection principle.

Notably, post-processing of the trajectories with machine learning or model
fitting approaches could be used to further "deconvolve" the tracking data.
However, we chose to avoid unnecessary complexity in the analysis as the per-
formance was sufficient for simultaneous tracking of animal bodies in various
behavioral contexts and detecting a representative sample of step trajectories.

2.3 Validation of the 3D Motion Capture (MoCap) approach with well-known synthetic cannabinoid receptor agonist CP55,940

To validate the newly-developed 3D MoCap system, we first chose to replicate
previously-published results obtained with a well-known synthetic cannabi-
noid receptor agonist CP55,940 [30, 44], before using compounds of less-known
behavioral effects. We selected low doses (0.03, 0.1, 0.3 mg/kg), out of which
only the highest dose was expected to result in any noticeable somatic effects
[44], hypothesizing that the MoCap system might detect subtler behavioral
changes than standard methods. The doses were verified in our laboratory with
standard assay for triad of cannabimimetic effects (antinociception, hypother-
mia or catalepsy; Suppl. Fig. 4). The results were similar even though slightly
smaller than previously observed [44] indicating that the animals used in cur-
rent study might be less sensitive to the doses than we observed in our earlier
studies.

Fig. 2 Marker based 3D motion capture increases sensitivity of behavioral analysis. a. Example of ankle trajectories during horizontal open field exploration: swing starts and stops (upper panel) and average ankle trajectory during all swings of representative animal following vehicle or CP55,940 (0.3 mg/kg) treatment. b. Example traces of locomotor activity in the open field following vehicle or CP55,940 at 0.3 mg/kg in representative animal. c. Example of ankle trajectories during vertical climbing: swing starts and stops (upper panel) and average ankle trajectory during swing of all steps of a representative animal following vehicle or CP55,940 (0.3 mg/kg) treatment. d. Example traces of locomotor activity during climbing following vehicle or CP55,940 at 0.3 mg/kg in representative animal. e. Total distance traveled (top) and mean activity index (bottom) during open field (left) and climbing (right). CP55,940 significantly decreased mean activity index at 0.3 mg/kg compared to vehicle, *p<0.05 (1-way ANOVA F (2.19, 19.75) = 4.25, Dunnett post hoc test) in the open field but not during climbing. f. Distance traveled (top) and activity index (bottom) as function of time during open field exploration and climbing. CP55,940 significantly decreased activity index during open field exploration. Closed symbols denote significance in Tukey post hoc test vs. vehicle (2-way ANOVA). g. CP55,940 significantly reduced maximum and average speed of ankle during climbing (right) and decreased step height in both open field (left) and climbing: *p<0.05, **p<0.01, ***p<0.001 post hoc comparison vs. vehicle (2-way ANOVA). e-g. n = 10 mice per group (randomized, counterbalanced within-subject design) mean±SEM.

2.3.1 General activity and locomotion parameters.

We analyzed locomotor behavior in two behavioral tasks: horizontal exploration of the open field (Fig. 2a,b) and vertical climbing on a spoked wheel (Fig. 2c,d). We extracted measurements of changes in step kinematics as well as distance traveled, speed and general activity level, quantified as average of all markers' speeds ("activity index"; Fig. 2a-d show example recordings for a representative individual treated either with vehicle or CP55,940 at 0.3 mg/kg). The results confirmed previous observations of a slight reduction of locomotor activity in the open field (Fig. 2b,e,f, Suppl. Fig. 5a-e, Suppl. Video

2). However, even with the highest dose used, trends towards decreased distance travelled ($p = 0.36$, Fig. 2f) and time spent locomoting ($p = 0.18$; Supp. Fig. 5c) were not significant and speed of locomotion (above 40 mm/s; $p = 0.78$, Suppl. Fig. 5d) was not affected. Interestingly, CP55,940 induced significant dose-dependent decrease in the activity index (average speed of all markers) during 120 s recordings ($F (3, 27) = 4.245$, $p = 0.014$), which was most pronounced at 0.3 mg/kg (Fig. 2e,f), but also significant at 0.1 mg/kg (Suppl. Fig. 5a).

To our surprise, no decrease in locomotor activity was observed in the climbing task, even though the mice performed it immediately after completing an open field task (Fig. 2d,e,f, Suppl. Fig 5f-j, Suppl. Video 3). In fact, a trend toward increased activity index following CP55,940 at 0.1 mg/kg during climbing could be discerned (Suppl. Fig. 5f). The reduction of locomotor activity in the climbing task with time was significant in all groups but less pronounced compared to the open field (Fig. 2a-f, Suppl. Fig. 5a-l).

There was no significant effect on stance width in either open field or climbing assays (data not shown), in line with the previous studies [30] that reported widened stance only after a higher dose (1 mg/kg).

2.3.2 Step kinematics parameters.

The number of steps during trials was not significantly affected by CP55,940 in open field ($p=0.27$, Suppl. Fig. 5k) or climbing ($p=0.23$ Suppl. Fig. 5l). However, several step kinematic parameters were significantly altered (Fig. 2g, Suppl. Fig. 5m). Despite overall locomotion was not suppressed in the climbing task, significant slowing of ankle kinematics during swings was apparent with the highest dose (max ankle speed, $F(3, 27) = 8.77$, $p<0.001$; average ankle speed, $F (3, 27) = 3.295$, $p = 0.036$; Fig. 2g; Suppl. Fig. 5m shows similar measurements for knees). Moreover, CP55,940 significantly decreased height of ankle swing during climbing ($F (3, 27) = 16.03$, $p<0.001$), which was also observed in the open field ($F (3, 27) = 4.474$, $p = 0.011$); Fig. 2g) without other significant kinematic effects.

In summary, our results indicate that while CP55,940 inhibits exploratory locomotor behavior in the open field as previously reported [44] even at moderate doses, it does not show such inhibition in the climbing task. Moreover CP55,940 at moderate doses significantly altered step kinematics by affecting speed and height of the swing.

2.4 Effects of endocannabinoid signaling elevation on locomotor activity

Two best-known endogenous cannabinoids, 2-AG and AEA, are rapidly degraded by their respective catabolic enzymes, monoacylglycerol lipase (MAGL), and fatty acid amide hydrolase (FAAH)[27, 41, 42]. To elevate concentrations of locally-produced 2-AG and AEA we used selective inhibitors of MAGL and FAAH, MJN110 and PF3845, respectively. These inhibitors have

Fig. 3 Bidirectional effects of inhibition of endocannabinoid degradation on locomotor activity. Examples of locomotor activity during open field exploration (a, b) and climbing (c, d) following treatment with PF3845 (30 mg/kg) or MJN110 (2.5 mg/kg) and compared to vehicle. a and c: Examples of average ankle trajectories ±SEM of all ankle swings of a representative animal during horizontal exploration and climbing, respectively. b and d: Examples of locomotor activity of a representative animal during horizontal exploration and climbing, respectivtly. e. Total distance traveled (top) and mean activity index (bottom) were significantly affected by treatment in horizontal exploration (left) (F (1.67, 15.06) = 10.52, F (1.92, 17.28) = 9.95), but not in climbing (right), *p<0.05, vs. vehicle; ##p<0.01 vs. PF3845 in post hoc comparison. f, left: Distance traveled (top) and activity index (bottom) were reduced (PF3845) and increased (MJN110) during the 60-s trials compared to vehicle control during horizontal exploration. g. PF3845 and MJN110 affected swing speeds and direct distances in bidirectional manner, so that PF3845 decreased them and MJN110 increased during open field exploration (left). The effects observed during climbing (right) were smaller than in open field. n=10 mice per group (randomized, counterbalanced within-subject design), mean±SEM.

been shown to reliably elevate 2-AG and AEA levels in the mouse central nervous system and peripheral tissues [40–42]. To assess the behavioral effects of acute elevation of 2-AG and AEA in mice, we administred systemic MJN110 and PF3845 at relatively high doses that are effective in models of pain[40–42], and subjected mice to the same behavioral tasks as we used with CP55,940.

Inhibition of endocannabinoid degradation caused significant bidirectional changes in locomotor activity and step kinematics during horizontal exploration of the open field and smaller but significant changes during vertical climbing (Fig. 3, Suppl. Fig. 6, Suppl. Video 4,5). Fig. 3a-d shows representative examples demonstrating changes in average ankle swing trajectories (Fig. 3a,c) as well as distance traveled, speed and general activity (quantified as activity index (AI); Fig. 2b,d) in a mouse treated with vehicle, PF3845 (30 mg/kg) and MJN110 (2.5 mg/kg). As compared in 2-way ANOVA, effects of PF3845 and MJN110 were bidirectional, where PF3845 caused significant

inhibition of locomotor activity (Fig. 3a-f, Suppl. Fig. 6a-j) in all measures (distance traveled ($F_{(2, 18)} = 8.75$, p = 0.022), activity index ($F_{(2, 18)} = 9.96$, p = 0.012), time spent locomoting ($F_{(2, 18)} = 8.89$, p = 0.021) and locomotion speed ($F_{(2, 120)} = 13.56$, p<0.001)) compared to vehicle. In contrast, MJN110 induced significant increase in locomotor activity in the open field expressed as increase in distance traveled (Fig. 3a-f) and activity index (Fig. 3a-f) and increased time spent locomoting but had no significant effect on the speed as compared to vehicle. These bidirectional effects were also seen in climbing assay (Fig. 3d-f), as well as when lower doses were used (Suppl. Fig. 7). None of inhibitors affected stance width (data not shown).

2.5 Effects of endocannabinoid signaling elevation on step kinematics parameters

Similarly to effects on general locomotion, both inhibitors induced bidirectional effects on step kinematics. PF3845 (30 mg/kg) significantly decreased total number of ankle swings of both left and right leg during open field exploration ($F_{(1.75, 15.73)} = 10.45$, p=0.002) and climbing task ($F_{(1.79, 16.09)} = 4.59$, p=0.030) as compared to vehicle (Suppl. Fig. 6k,l). We have assessed differences in step kinematics parameters such as direct distance, total distance, height, duration, average speed, maximum speed in right and left leg under three treatment conditions: vehicle, PF3845, MJN110 (2-way ANOVA). PF3845 (30 mg/kg) significantly decreased direct ($F_{(2, 18)} = 3.75$, p=0.043) and total distance ($F_{(2, 18)} = 4.83$, p=0.021) as well as maximum ($F_{(2, 18)} = 24.09$, p<0.0001) and average speed ($F_{(2, 18)} = 8.69$, p=0.0023) of both right and left ankle (Fig. 3g). Similar effects of PF3845 were observed on swing of the knee (Suppl. Fig. 6m). MJN110 (2.5 mg/kg) significantly increased direct distance of the left ankle and knee swing compared to vehicle (Fig. 3g). Less pronounced but significant effects on swing total distance and speed were observed in the climbing task (Fig. 3g). There were no effects of treatment on duration or height of the swing of ankle or knee in open field or climbing (Fig. 3g). At lower doses (MJN110 1.25 and PF3845 10 mg/kg) we observed small but significant effects on step kinematics parameters of both compounds (Fig. 3g, Suppl. Fig 7m).

2.6 Endocannabinoid modulation of locomotory episode structure

Bidirectional effects of inhibition of endocannabinoid degradation on locomotor activity were also reflected in number of locomotory episodes defined by maintained displacement of the mouse at speeds over 40 mm/s. In the open field PF3845 induced significant decrease in total number of locomotory episodes as compared to vehicle (p = 0.013) and MJN110 (p=0.0026) (data not shown). In order to gain understanding of the mechanisms driving the bidirectional effects of MJN110 and PF3845 on locomotory activity, we analyzed the locomotory behavior of animals in more detail. Strikingly, we found that in all

mice under all pharmacological conditions, the variable-duration locomotory
episodes were composed of locomotory "bouts" during which the mice accel-
erate and decelerate in a stereotypical fashion. Comparing these bouts (shown
for a representative example mouse in Fig. 4a1) under different drug conditions
revealed that not only do the bouts maintain their speed profiles throughout
the trials (as shown in Fig. 4a2 by comparing the bouts speed averages from
first and last halves of the 60-s recordings), but also that the bouts are remark-
ably similar between different drug conditions (Fig. 4b1). Even in the PF3845
group, where the average locomotory speeds were significantly reduced in com-
parison to vehicle and MJN110 (Suppl. Fig. 6d), the duration of the bouts was
virtually indistinguishable between the groups (bout half widths 1.19 ± 0.003s,
1.36 ± 0.006s and 1.22 ± 0.003 s, ANOVA p = 0.06, Fig. 4b2). Indeed, the lower
average speeds in PF3845 might result from reduced acceleration during the
initial phase of locomotory bouts (quantified as the bouts slope; Fig. 4b3;
pooled averages over entire trials 62.6 ± 0.23 mm/s^2, 42.3 ± 0.22 mm/s^2 and
65.1 ± 0.21 mm/s^2 for vehicle, PF3845 and MJN110; vehicle vs. PF3845, p <
0.0001; vehicle vs. MJN110, p = 0.46).

Another property of the locomotory episode structure clearly evident in
the trial speed profiles (Fig. 4a1) was the invariance of bouts intervals; indeed,
throughout the entire trials and in all pharmacological conditions, sequential
bouts occurred at remarkably constant intervals (mean inter-bout intervals,
2.2 ± 0.004, 2.1 ± 0.007 and 2.1 ± 0.004 s for vehicle, PF3845 and MJN110;
ANOVA p > 0.99, Fig. 4b4) throughout the locomotory episodes. Thus,
the changes in time spent locomoting (Suppl. Fig. 6c) are reflected in the
larger number of bouts during the early parts of the trials under MJN110
(1.42 ± 0.008, 1.14 ± 0.006 1.7 ± 0.014 bouts per locomotory episode; 116, 92 and
105 episodes for vehicle, PF3845 nd MJN110 groups, respectively; Fig. 4b5).
In other words, the larger distance covered in MJN110 mice is due to them ini-
tiating more of these stereotypical bouts before pausing, while the mice under
PF3845 initiate slightly fewer of these 1-2 -second-long bouts.

The fact that MJN110 influences the stopping likelihood of a locomot-
ing mouse prompted us to ask whether it would also modulate the eagerness
of the mice to initiate a locomotory episode. If this were the case, it should
manifest as shortened duration of the pauses between locomotory episodes.
However, as shown in Fig. 4b6i, the distribution of pause durations was nearly
identical between vehicle and MJN110 (2.1 ± 0.03 and 1.9 ± 0.03 sek, K-S test
0.9), whereas the pauses were slightly longer in PF3845 (2.6 ± 0.03 s; K-S 0.047
for PF3845 vs. vehicle). Thus, the increased exploratory behavior in MJN110
involves changes in mechanisms that control stopping rather than starting loco-
motion. The slightly longer pause durations caused by PF3845 do not suffice to
explain the dramatic decrease in the distance the mice covered during PF3845
trials. Therefore we wondered if PF3845 would generally decrease muscle tone,
leading to weaker acceleration and thereby eventual failure of reaching suffi-
cient momentum for maintaining locomotion during the locomotory bout time
windows. To investigate this, we examined the posture and activity measures
552

Fig. 4 Invariant microstructure of locomotory episodes under modulation of endocannabinoid system. a, speed measurements during full 60-s trials in a single mouse in vehicle, PF3845 and MJN110 conditions. a1, speed during full trials; light and dark grey boxes denote locomotory episodes and pauses, respectively. Filled circles: speeds peaks (purple, first half of the trial; green, last half of the trial). Open circles: initiations of locomotory bouts. Dashed line indicates threshold for locomotion. a2, average ±SEM speeds in all locomotor bouts in all animals, divided into first and last-half events, aligned vertically to bout initiation (open circles). b, MJN110 does not cause changes in the shape of bouts (b1-3) or increase in likelihood of initiating locomotory episodes (b4) but increased number of bouts within the early episodes of locomotion (b5). PF3845 reduces the acceleration during a bout (b1, 3). Pause durations (b6i) and behavior during pauses (b6ii, iii) were not significantly affected. c, CP55,940 (0.3 mg/kg dose) did not cause changes in locomotory bout shape (c1-c3) nor their intervals or number within a locomotory episode (c4-c5). Instead, CP55,940 increased duration of locomotory pause durations (c6i), simultaneously lowering mouse posture and activity index (c6ii, iii). Closed symbols in panels b2-b5 denote significance in post hoc test vs.vehicle. Asterisk denote significant diffrence in distributions with K-S test vs. vehicle for the group indicated by color (b6) or ANOVA (c6) p* < 0.05, p***, < 0.001. n=10 mice per group, mean±SEM.

during the locomotory pauses (Fig. 4b6ii-iii). Somewhat surprisingly, no dramatic differences were found, despite slight lowering of the shoulder height in MJN110 could be seen (11.2±0.06, 12.7±0.07, 10.6±0.07 mm; ANOVA p = 0.062, K-S PF3845 vs. MJN110 p = 0.005). These results obtained from quantification of the invariant locomotory bouts suggest that the bidirectional effects of MJN110 and PF3845 are mainly realized by modulation of motivational parameters of behavior: PF3845 makes the mice slower in initiating a locomotory bout. In contrast, MJN110 upregulates the likelihood that a locomoting mouse will initiate another bout instead of stopping, leading to overall longer distance covered.

These results with the selective endocannabinoid degradation enzyme inhibitors are in stark contrast with those obtained with the exogenous 1 receptor agonist CP55,940 (Fig. 4c). While the bout shapes and intervals were virtually identical in mice under the influence of CP55,940 (panels c1 - c4; bout half-widths 1.25±0.006, and 1.39±0.013 sek; bout accelerations 54.9±2.6 and 55.3±3.7, p = 0.93; bout intervals 2.0±0.17 and 2.1±0.14 s for vehicle and 0.3 mg/kg CP55,940), the decrease of distances mice travelled was not caused

by shortening of locomotory episode duration or number of bouts per episode
(1.35±0.01 and 1.41±0.03 bouts in an episode in vehicle vs. CP55,940; n =
77 and 37 locomotory episodes; p > 0.99; Fig. 4c5). Instead, nonspecific acti-
vation of CB_1 receptors with CP55,940 increased duration of pauses between
locomotory episodes (Fig. 4c6i; 1.73±0.03 vs. 2.6±0.08 s; ANOVA p < 0.001;
n = 71 and 29 pauses in the first 60 s of the trials), concomitantly with lower-
ing of overall posture (shoulder height; Fig. 4c6ii, 10.8±0.07 vs. 8.0±0.1 mm;
ANOVA p < 0.001) and activity index (Fig. 4c6iii, 25.6±0.12 vs. 19.0±0.25,
ANOVA p < 0.001) during the pauses.

Taken together, even though selective FAAH inhibition with PF3845
affected behavior in a way that superficially resembles that of activation of
CB_1 receptor with exogenous agonist CP55,940, the underlying features are
different and likely involve different targets.

612
613 # 3   Discussion
614

Taking advantage of the high precision and accuracy of our markerless
3D MoCap system, we revealed that selective inhibition of degradation of
endocannabinoids 2-AG and AEA results in distinct, bidirectional kinematic
phenotypes that markedly differ from the effects of an exogenous synthetic lig-
and of cannabinoid receptor. To our surprise, we also found that the effects
were not universal to locomotory activity but differed between horizontal and
vertical locomotion.

The endocannabinoid system, consisting of endogenous cannabinoids, their
receptors and enzymes responsible for their biosynthesis and degradation [27],
is of high interest in development of treatments for various diseases including
neurodegenerative, autoimmune, and pain-related disorders [27–29, 45, 46].
The medicinal compounds targeting cannabinoid receptors have relatively mild
side effects (e.g. subjective intoxication, inhibition of locomotion and memory
impairment [47]) and future studies aiming at reduction of the cannabimimetic
side effects will further increase their attractiveness for widespread therapeu-
tic use [27, 40, 43, 44]. Thus in the present study we did not aim to evoke
dramatic behavioral distortions but focused on more subtle effects that might
be indicative distinct interoceptive states, difficult to observe with traditional
behavioral assessment methods[40, 43].

The profound locomotion-inhibiting effect of the CB_1 receptor activa-
tion by exogenous compounds is well known [30, 48]. However, even though
three decades have passed since the initial characterization of endogenous
cannabinoids [49, 50], their rapid degradation in living animals has limited
research into their roles in behavior. Moreover, effects of synthetic and phyto-
cannabinoid (e.g. delta-9-THC) on movement have mostly been examined
using coarse behavioral assays as well as high doses unsuitable for widespread
therapeutic use.

Recently-developed potent and selective inhibitors of the enzymes responsi-
ble for endocannabinoid degradation can be used to reveal effects of enhancing
644

endocannabnoid signaling produced naturally *in situ* [40–43, 51], unlike direct
exogenous compounds targeting the CB_1 receptor. We and others have shown
that administration of these highly selective FAAH and MAGL inhibitors have
effects similar to direct ligands, suggestive of therapeutic potential with less
cannabimimetic side effects [40–43, 51–54].

Endocannabinoids interact with multiple receptors besides CB_1 and CB_2
and manipulating their availability is expected to produce different effects than
compounds such as CP55,940, designed to target CB_1 and CB_2 receptors [27].
As the ligand-receptor interactions differ in a cell- and tissue-type-dependend
manner [55–57] and expression of receptors and enzymes is highly diverse
across animal bodies [27], it should not, in fact, be a surprise that each of
the treatments in this study led to a distinct behavioral phenotype. More
research is necessary to establish mechanisms underlying observed differences
and establish molecular targets involved, which was beyond the scope of this
article.

In the context of therapeutic use considerations, it is interesting to note
that the selective MAGL inhibitor MJN110, expected to enhance 2-AG sig-
naling, led to a clear stimulatory effects on locomotion, including increase
in hindlimb speed (Fig. 3). This is in line with our and others' observations
of mild stimulatory effects of MAGL inhibition [40, 58] as well as studies
suggesting that elevated 2-AG signaling might be associated with hyper-
dopaminergic and propsychotic outcomes [59, 60]. Considering the known
reduction of locomotion and emergence of suicidal ideation caused by CB_1
receptor antagonism, [61, 62], it seems possible that 2-AG, the most abun-
dant endogenous cannabinoid in the central nervous system, would play a
stimulatory or pro-dopaminergic role in animal behavior.

In contrast, enhancing AEA signaling with inhibition of FAAH resulted
in locomotory suppression, and the effects were consistent in the behavioral
tasks. However, while the cannabinoid receptor agonist CP55,940 with known
behavior-suppressing effects did dramatically reduce locomotion in horizontal
exploration, no such inhibition was seen in vertical climbing task (Fig. 2).
This observation clearly indicates that inhibition of exploration in the open
field is not a result of impaired locomotory coordination and suggests that
the suppression may consist of a significant motivational component, possibly
combined with effect on muscle tone. Furthermore, it highlights the importance
of using multiple behavioral tasks in order not to miss effects that might be
context-dependent.

In addition to observing the effects on general locomotory behavior that
could be in principle be assessed with simple whole-animal tracking, the
marker-based MoCap recordings revealed several more subtle signatures of
animal behavior. First, we note that the magnitude of animal movement
(quantified as "activity index"; Fig. 2-3) is more sensitive measurement of
behavioral suppression or enhancement than traditional measures such as dis-
tance traveled. Second, the bidirectional effects on general locomotor activity
were accompanied by significant changes in step kinematics, that were also

bidirectional (Fig. 3). While PF3845-treated mice generated less steps that
were shorter in distance and slower, MJN110-treated mice made steps that
were longer and faster. Interestingly, across all experiments the most pro-
nounced changes in step kinematics parameters were observed in swing speeds,
some changes were observed in swing distance or height, but step durations
were remarkably unchanged. Third, a comparison of behavioral microstructure
features (hindpaw swing duration, figures 2-3, and "bouts" of speed within
locomotory episodes, Fig. 4) revealed surprising invariance: across all animals,
all groups and throughout the trials, durations of swings and locomotory bouts
were virtually identical in both horizontal and vertical locomotion. Thus, the
CP55,940 and PF3845 did not make mice cover shorter distances by shortening
their steps or locomotory "bouts" durations, but by decreasing the acceler-
ation when initiating those events. Conversely, the striking enhancement of
locomotory activity induced by MJN110 was not accompanied by any changes
in locomotory "bout" in comparison to vehicle-treated condition, but by ele-
vated likelihood of initiating more bouts within a locomotory episode before
pausing the explorative activity. Finally, comparing animal posture and behav-
ior during the pauses between locomotory episodes suggests that while direct
activation of CB_1 receptors with a low dose of CP55,940 and enhancement
of AEA-mediated endocannabinoid signaling similarly suppressed locomotion,
the latter is associated with less cannabimimetic side effects even at high doses.

While further research into the systemic, cellular and molecular compo-
nents underlying these effects is needed, the work presented demonstrates
not only that marker-based 3D motion capture is an attainable method to
apply in mice and other laboratory rodents, but also that it allows investi-
gation of small movements (such as tremor; unpublished observations by the
authors) in entirely unrestricted animals. Such spatiotemporal precision may
reveal novel, subtle kinematic symptoms of degenerative diseases and support
targeted development and refinement of interventions. Finally, the simplicity
and precision of the method, combined with analysis that does not depend on
machine learning or model-fitting approaches opens doors for more creative
experimental design involving natural behaviors, such as hunting and social
interactions, and could identify kinematic signatures of interoceptive states
that currently are extremely challenging to define in rodent models ([40]).

4 Methods

4.1 Experimental subjects and drug administration

We used adult male C57BL/6A (CLEA, Japan) mice (10-12 weeks old at
the beginning of experiments) that were subjected to systemic intraperitoneal
(i.p.) administration of drugs followed by behavioral assessment. To modulate
endocannabinoid signaling we used the latest generation of pharmacological
tools including inhibitors of FAAH and MAGL enzymes (PF3845 and MJN110,
respectively, kindly provided by Dr. Benjamin Cravatt's laboratory, Scripps
Institute, USA). As a positive control we used potent agonist of CB_1 and CB_2

receptors CP55,940 (Tocris, UK). The drugs were administered in a vehicle solution consisting of ethanol, colliphor and saline (1:1:18 ratio). Drugs were given at a volume of 10 $\mu l/g$ of body mass. Enzyme inhibitors will be administered 120 min before testing to ensure endocannabinoid accumulation, direct agonist 15-30 min before testing and antagonists 10 min before inhibitors or agonists.

4.2 Permanent marker implantation on skin surface

A major obstacle in developing 3D motion capture for small animals has been lack of adequate markers that could be used in rodents. We have developed a new method of attachment of permanent stainless-steel markers on a surface of skin of mice. Our preliminary studies indicated that implantation of stainless steel markers is the safest and the least invasive method of marker placement, and it will ensure natural behavior of the animal, as commonly-used plastic markers attract attention of animals, are difficult to attach (require strong glue causing skin irritation) but easy for animals to remove, which leads to chewing and swallowing. In contrast, stainless steel piercings do not cause any pain or discomfort and do not cause any change in the behavior. Another advantage of permanent markers is that animals can be tracked over time with the same localization of markers which enables powerful within-subject studies and high level of measurement precision. 11-14 days before the start of behavioral experiments mice undergo a procedure of marker implantation that enables high-resolution behavioral monitoring with infrared (IR) cameras. Under full isofluorane anesthesia (2-3%), 5 pairs of stainless steel markers were implanted on the skin at landmark locations of the body (Fig. 1a, Suppl. Fig. 1 and 2.). Markers are <14 mm long, diameter of 0.9-1.27 mm (18-20 gauge) stainless steel rods ending with screw-on ball on both sides. The weight of all the markers combined does not exceed 10% of the animal body weight. The screw-on balls residing at the skin surface are covered with a strongly infrared (IR)-reflective coating enabling IR cameras to track the location of the markers and thereby position of the mouse body and limbs. To implant the markers, 2 small (<1.5 mm) holes are made in the skin with an 18 gauge (1.27 mm) sterile needle under full anesthesia complemented by local analgesic (3% lidocaine gel applied on the skin surface). The needle is then replaced with the piercing that has been disinfected by 70 % ethanol for at least 20 min. Markers (Fig.1, Suppl. Fig. 1, 2, 3) were placed in the landmark locations of animals' body: on the back, hips, knees, ankles and shoulder blades. These locations have been chosen to allow for precise calculation of body movements during locomotion, the posture of an animal as well as fine changes in body kinematics, such as tremor, miscoordination and subtle swaying.

4.3 Behavioral training

Another major obstacle in developing rodent 3D motion capture system is that during such 3D recordings there can be no obstruction between cameras

and an experimental subject, and it is not possible to have walls surrounding experimental arena. Therefore, animals require adequate training to a) habituate to naturally anxiogenic open field arena, b) train mice to remain in the arena during testing without any restriction of movement. We have developed a dedicated protocol of animal training allowing such behavioral assessment. Experiments started 11-14 days following marker implantation to ensure proper wound healing and stable marker position. During those days animals were subjected to handling, acclimating and training, in following order: 1) handling: a) taken out from the cage with the tube b) gently pulling by tail base on the outer part of hand or forearm c) scooping holding the tail gently d) scooping (2-3 days). 2) adaptation to arena: a) placement in circular 30 cm arena with 4 cm walls in brightly lit room (2-3 days) b) placement in circular area without walls and preventing mice from leaving the arena (taking back by the tail if 4 paws leave, tapping nose if 2 paws leave) (3-4 days). Mice are being kept in the arena for 2-5 min. 3) task training: a) vertical climbing on mesh b) horizontal climbing on string c) beam walking d) righting reflex (2-3 days, 1 min trial or less). Each animal is subjected to the same level of training. All animals are trained until they correctly perform task as expected. Markers that are worn by mice every day are being exchanged into retroreflective and larger markers 15 minutes before the start of recording. Mice are habituated to the process of manual exchange of markers that takes 3-5 min, so they do not need to be anesthetized, which could disturb the behavior. After finished recording markers are changed again and mouse is being returned to home cage.

4.4 Retroreflective marker preparation

Retroreflective markers are hand made by covering 4 mm diameter screw on stainless steel piercing ball (Fig. 1a, Suppl. Fig. 1a and 2) with highly retroreflective tape (3M, MN, USA). The tape is first cut into small pieces allowing covering ball with relatively smooth surface (Suppl. Fig. 3a). Then, tape is covered by layer of thin plastic (Linear Low Density Polyethylene (LLDPE) film, TRUSCO, Japan) followed by cover of liquid UV-cured plastic (Bondic, NY, USA) ensuring hard but transparent layer that will be sufficiently resistant to bites and other damage that may occur during testing. Layer of thin plastic protects tape from contact with liquid that would diminish its retroreflective properties. Other methods including various retroreflective paints and sprays did not provide strong enough retroreflective signal on such a small volume and/or were not sufficiently resistant to damage. Number of materials were tested before identifying most successful marker design providing strong signal with excellent detection by the cameras, lightweight, and hard surface resistant to damage. All retroreflective markers are quality tested before they are used on animals.

4.5 Assessing behavior using 3D motion capture

We adapted existing 3D motion capture system (Qualisys, Sweden) that is used for studying human (e.g. in sports medicine or diagnostics of movement disorders) and large animal biomechanics. Recordings were performed with 7 high-speed, high-resolution infrared-sensitive cameras (Qualisys Oqus 7, 12 megapixel sensor, 40 mm / f2.8 aperture) with integrated infrared-emitting LEDs, at 300 Hz allowing for highly precise recording and analysis of high-speed movement trajectories.

Behavioral assesment was taking place within 30 x 30 x 30 cm in the center of arena where cameras were focused. The open field arena is a 30 x 30 cm textured poliethylene floor (Suppl. Video 2,4). The climbing was performed on the outer surface of the metal mesh of the spoked running wheel for rats (25 cm in diameter). Wheel was moved manually according to voluntary mouse movement. Retroreflective markers were placed (spaced every 10 cm) on the outer side of the wheel to track it's movement (Suppl. Video 3,5). All elements of arena were cleaned in between trials.

4.6 Calibrating motion capture recording for optimal tracking

To ensure reliable tracking of the retroreflective markers in challenging behavioral conditions, we calibrated the camera system at the beginning of each experimental day and every 4 hours during the day to avoid erratic tracking due to minute shifts in experimental system arrangements. In brief, we used the procedure provided by the hardware maker (Qualisys; https://docs.qualisys. com/getting-started/content/getting_started/running_your_qualisys_system/ calibrating_your_system/calibrating_your_system.htm/) with slight modifications. First, to acquire uniform shapes of the markers while the animal moves in space, we focused the cameras at the outer perimeter of the behavioral arena instead of the center and used low aperture values (f = 2.8). This results in slight blurring of the small imperfections of the markers and more uniform image plane projection. Next, a L-shaped calibration bar and a T-shaped calibration wand are used according to the instructions provided to define the tracking volume so that X and Y directions are identical across all experiments. Great care was taken to ensure that the calibration volume was uniformly covered in the process. Calibration was considered acceptable when average triangulation residuals for all cameras were below 0.2 mm.

4.7 Source code and kinematic feature extraction

Kinematic parameters were extracted by custom made code in Matlab (2019a) run on MacOS Monterey 12.1. All code will be made available at github repository at the time of publication and link will be provided here. Key parameters computed from the 3D trajectories:

- mouse speed: displacement of mean hip markers in 1 second

• locomotory episodes: frames in which mouse speed remained over 40 mm/s
for longer than 100 frames. The threshold value was determined based on
observations of 10 randomly selected videos from the data pool.
• locomotory bouts: blocks with peaks in speed with minimal prominence of
15 mm/s, and the nearest matching local minima.
• Activity index: mean of all markers' speed (smoothed over 10 frames).
• Hip-shoulder height: distance of mean hip and shoulder markers in the
dimension orthogonal to locomotory plane (horizontal or vertical for open
field and climbing, respectively).
• Hip-shoulder distance: euclidean distance of the mean hip and shoulder
markers.
• Step detection: Swing segments of steps were delineated based on kinematic
rather than positional features of ankles, understanding that limb joints do
not simply move up and down during stepping. The underlying principle is
that during swings, ankles move (in any direction) with a high speed relative
to the ground (either floor or the wheel), and a fully-resolved swing must
be bordered with acceleration and deceleration events, reflecting take-off
and touch-down events. From detected high-speed ankle events, we excluded
those for which we could not determine precise acceleration and deceleration
peaks. While this approach necessarily omits steps with "creeping" take-off
or touch-down events (that could be more prominent in the moments lead-
ing to ceasing locomotion), we chose it over more lenient inclusion criteria
in order to reduce the probability of falsely-identified steps (such as those
related to scratching or postural shifts in place) from contaminating the
measurements.
• Step measurements (calculated for both left and right hind limb separately):

– swing height: maximal displacement of the ankle marker during a swing,
in dimension orthogonal to the locomotory plane (horizontal or vertical)
– swing direct distance: euclidean distance of the ankle or knee marker
position at frames defined as swing start and swing end;
– swing total distance: length of the 3D trajectory of the ankle or knee
marker between the frames defined as swing start and swing end;
– swing mean and max speed were calculated from the segments designated
as swings, without smoothing.
– stance width: sum of the mean distance of left and right ankles to the
centerpoint of hip and coordinate markers

4.8 Data analysis

914 Data are presented as mean ± standard error (SEM) and were analyzed using
915 one-way or two-way analysis of variance (ANOVA). In case when some val-
916 ues were missing, these data were analyzed by fitting a mixed model, rather
917 than by repeated measures ANOVA. Dunnett's was used for post-hoc analysis
918 in the dose–response experiments, and the Tukey or Sidak test was used for

post-hoc analyses comparing different treatment groups. Differences were considered significant at the level of $p<0.05$. Statistical analysis was performed with GraphPad Prism version 8.00 (San Diego, CA) or Matlab (ver. 2019b, Mathworks, Natick, MA).

4.9 Data availability

All motion capture tracking files including labeled data will be fully available for download at the time of publication.

Declarations

4.10 Authors' contributions:

BIJ: experimental design, conducting experiments, data analysis, manuscript writing, visualization

AK: conducting experiments, programming, manuscript revision

AGO: conducting experiments, manuscript revision

DO: drug design and synthesis, manuscript revision

MJN: drug design and synthesis, manuscript revision

BFC: drug design and synthesis, manuscript revision

MYU: Software design and programming, data analysis, manuscript writing, visualization

All authors have read and agreed to the published version of the manuscript.

4.11 Funding:

This research was supported by Japan Society for Promotion of Science (JSPS) Fellowship for Overseas Researchers (P17388), Kakenhi Grant-in-Aid for JSPS Fellows (17F17388), and Kakenhi Grant for Scientific Research (21K06399) awarded to Bogna M. Ignatowska–Jankowska.

4.12 Ethics approval:

The animal study protocol was conducted in accordance with procedures approved by the Okinawa institute of Science and Technology (OIST) Institutional Animal Care and Use Committee (IACUC) (Protocol IDs: 2017-188, 2020-305) in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals (National Research Council, 2011) Every effort was made to minimize suffering.

References

[1] Ceseracciu, E., Sawacha, Z., Cobelli, C.: Comparison of markerless and marker-based motion capture technologies through simultaneous data collection during gait: Proof of concept. PLoS One (2014). https://doi.org/

10.1371/journal.pone.0087640

[2] Liu, S.Q., Zhang, J.C., Zhu, R.: A wearable human motion tracking device using micro flow sensor incorporating a micro accelerometer. IEEE Trans Biomed Eng (2020). https://doi.org/10.1109/TBME.2019.2924689

[3] Nakano, N., Otsuka, Y., Yamamoto, S., Tanaka, S., Takenouchi, T.: Evaluation of 3d markerless motion capture accuracy using openpose with multiple video cameras. Frontiers in Sports and Active Living **2** (2020)

[4] Schmitz, A., Wulf, G., Hossner, E.-L., Weigelt, M., Schega, L.: The measurement of in vivo joint angles during a squat using a single camera markerless motion capture system as compared to a marker based system. Gait & Posture **42**, 196–202 (2015)

[5] Steinebach, T., Grosse, E.H., Glock, C.H., Wakula, J., Lunin, A.: Accuracy evaluation of two markerless motion capture systems for measurement of upper extremities: Kinect v2 and captiv. Human Factors and Ergonomics in Manufacturing & Service Industries **30**(4), 291–302 (2020)

[6] Topley, M., Richards, J.G.: A comparison of currently available opto-electronic motion capture systems. Journal of biomechanics **106**, 109820 (2020)

[7] Marshall, J.D., Aldarondo, D.E., Dunn, T.W., Wang, W.L., Berman, G.J., Ölveczky, B.P.: Continuous whole-body 3d kinematic recordings across the rodent behavioral repertoire. Neuron **109**(3), 420–437 (2021)

[8] Becker, M.I., Person, A.L.: Cerebellar control of reach kinematics for endpoint precision. Neuron **103**(5), 1048–10628 (2019). https://doi.org/10.1016/j.neuron.2019.05.007

[9] Becker, M.I., Calame, D.J., Wrobel, J., Person, A.L.: Online control of reach accuracy in mice. J Neurophysiol **124**, 1637–1655 (2020)

[10] Marshall, J.D., Li, T., Wu, J.H., Dunn, T.W.: Leaving flatland: advances in 3d behavioral measurement. Curr Opin Neurobiol **73**, 102522 (2022)

[11] von Ziegler, L., Sturman, O., Bohacek, J.: Big behavior: challenges and opportunities in a new era of deep behavior profiling. Neuropsychopharmacology **46**, 33–44 (2021). https://doi.org/10.1038/s41386-020-0751-7

[12] Mathis, A., Mathis, M., Billard, A., Maheswaranathan, N., Calinon, S., Dragan, A.D., Schaal, S.: Deeplabcut: markerless pose estimation of user-defined body parts with deep learning. Nature Neuroscience **21**(11), 1281–1289 (2018)

[13] Lauer, J., Mathis, A., Mathis, M., Billard, A., Maheswaranathan, N., Calinon, S., Dragan, A.D., Schaal, S.: Multi-animal pose estimation, identification and tracking with deeplabcut. Nature Methods **19**(3), 496–504 (2022)

[14] Nath, T., Mathis, A., Mathis, M., Billard, A., Maheswaranathan, N., Calinon, S., Dragan, A.D., Schaal, S.: Using deeplabcut for 3d markerless pose estimation across species and behaviors. Nature Protocols **14**(8), 2152–2176 (2019)

[15] Bohnslav, J.P., Babadi, B., Mathis, A., Mathis, M., Billard, A., Maheswaranathan, N., Calinon, S., Dragan, A.D., Schaal, S.: Deepethogram: a machine learning pipeline for supervised behavior classification from raw pixels. bioRxiv (2020)

[16] Nourizonoz, A., Mathis, A., Mathis, M., Billard, A., Maheswaranathan, N., Calinon, S., Dragan, A.D., Schaal, S.: Etholoop: automated closed-loop neuroethology in naturalistic environments. Nature Methods (2020)

[17] Pereira, T.D., Shaevitz, J.W., Murthy, M.: Quantifying behavior to understand the brain. Nature Neuroscience **23**, 1537–1549 (2020). https://doi.org/10.1038/s41593-020-00734-z

[18] Wiltschko, A.B., Tsukahara, T., Zeine, A., Anyoha, R., Gillis, W.F., Markowitz, J.E., Peterson, R.E., Katon, J., Johnson, M.J., Datta, S.R.: Revealing the structure of pharmacobehavioral space through motion sequencing. Nature neuroscience **23**(11), 1433–1443 (2020)

[19] Wade, L., Needham, L., McGuigan, P., Bilzon, J.: Applications and limitations of current markerless motion capture methods for clinical gait biomechanics. PeerJ **10**, 12995 (2022)

[20] Buckley, C., Alcock, L., McArdle, R., Rehman, R.Z.U., Del Din, S., Mazzà, C., Yarnall, A.J., Rochester, L.: The role of movement analysis in diagnosing and monitoring neurodegenerative conditions: Insights from gait and postural control. Brain sciences **9**(2), 34 (2019)

[21] Garner, J.P.: The significance of meaning: why do over 90% of behavioral neuroscience results fail to translate to humans, and what can we do to fix it? Ilar Journal **55**(3), 438–456 (2014)

[22] De Schutter, E.: Fallacies of mice experiments. Neuroinformatics **17**(2), 181–183 (2019)

[23] Kennedy, A.: The what, how, and why of naturalistic behavior. Current opinion in neurobiology **74**, 102549 (2022)

[24] Roepstorff, C., Dittmann, M.T., Arpagaus, S., Braganca, F.M.S., Hardeman, A., Persson-Sjödin, E., Roepstorff, L., Gmel, A.I., Weishaupt, M.A.: Reliable and clinically applicable gait event classification using upper body motion in walking and trotting horses. Journal of Biomechanics **114**, 110146 (2021)

[25] Dagenais, P., Hensman, S., Haechler, V., Milinkovitch, M.C.: Elephants evolved strategies reducing the biomechanical complexity of their trunk. Current Biology **31**(21), 4727–4737 (2021)

[26] O'Neill, M.C., Demes, B., Thompson, N.E., Larson, S.G., Stern Jr, J.T., Umberger, B.R.: Adaptations for bipedal walking: Musculoskeletal structure and three-dimensional joint mechanics of humans and bipedal chimpanzees (pan troglodytes). Journal of Human Evolution **168**, 103195 (2022)

[27] Di Marzo, V.: New approaches and challenges to targeting the endocannabinoid system. Nat Rev Drug Discov **17**, 623–639 (2018)

[28] Pacher, P., Kogan, N.M., Mechoulam, R.: Beyond thc and endocannabinoids. Annu Rev Pharmacol Toxicol **60**, 637–659 (2020)

[29] Mechoulam, R., Parker, L.A.: The endocannabinoid system and the brain. Annu Rev Psychol **64**, 21–47 (2013)

[30] Patel, S., Hillard, C.J.: Cannabinoid CB_1 receptor agonists produce cerebellar dysfunction in mice. Journal of Pharmacology and Experimental Therapeutics (2001)

[31] Rodriguez de Fonseca, F., del Arco, I., Martin-Calderon, J.L., Gorriti, M.A., Navarro, M.: Role of the endogenous cannabinoid system in the regulation of motor activity. Neurobiology of Disease **5**, 483–501 (1998)

[32] Tsou, K., Brown, S., Sañudo-Peña, M.C., Mackie, K., Walker, J.M.: Immunohistochemical distribution of cannabinoid CB_1 receptors in the rat central nervous system. Neuroscience **83**, 393–411 (1998)

[33] Albergaria, C., Silva, N.T., Darmohray, D.M., Carey, M.R.: Cannabinoids modulate associative cerebellar learning via alterations in behavioral state. Elife **9**, 61821 (2020)

[34] Steinmetz, A.B., Freeman, J.H.: Intracerebellar cannabinoid administration impairs delay but not trace eyeblink conditioning. Behavioural brain research **378**, 112258 (2020)

[35] Báez-Cordero, A.S., Pimentel-Farfan, A.K., Peña-Rangel, T., Rueda-Orozco, P.E.: Unbalanced inhibitory/excitatory responses in the substantia nigra pars reticulata underlie cannabinoid-related slowness of movements. Journal of Neuroscience **40**(30), 5769–5784 (2020)

[36] Steinmetz, A.B., Freeman, J.H.: Cannabinoid agonist administration within the cerebellar cortex impairs motor learning. Neurobiology of learning and memory **170**, 106896 (2020)

[37] Kishimoto, Y., Kano, M.: Endogenous cannabinoid signaling through the cb1 receptor is essential for cerebellum-dependent discrete motor learning. Journal of Neuroscience **26**(34), 8829–8837 (2006)

[38] Skosnik, P.D., Edwards, C.R., O'donnell, B.F., Steffen, A., Steinmetz, J.E., Hetrick, W.P.: Cannabis use disrupts eyeblink conditioning: evidence for cannabinoid modulation of cerebellar-dependent learning. Neuropsychopharmacology **33**(6), 1432–1440 (2008)

[39] Pearson-Dennett, V., Todd, G., Wilcox, R.A., Vogel, A.P., White, J.M., Thewlis, D.: History of cannabis use is associated with altered gait. Drug and alcohol dependence **178**, 215–222 (2017)

[40] Ignatowska-Jankowska, B., Wilkerson, J.L., Mustafa, M., Abdullah, R., Niphakis, M., Wiley, J.L., Cravatt, B.F., Lichtman, A.H.: Selective monoacylglycerol lipase inhibitors: antinociceptive versus cannabimimetic effects in mice. Journal of Pharmacology and Experimental Therapeutics **353**(2), 424–432 (2015)

[41] Ahn, K., Johnson, D.S., Mileni, M., Beidler, D., Long, J.Z., McKinney, M.K., Weerapana, E., Sadagopan, N., Liimatta, M., Smith, S.E., *et al.*: Discovery and characterization of a highly selective faah inhibitor that reduces inflammatory pain. Chemistry & biology **16**(4), 411–420 (2009)

[42] Niphakis, M.J., Cognetta III, A.B., Chang, J.W., Buczynski, M.W., Parsons, L.H., Byrne, F., Burston, J.J., Chapman, V., Cravatt, B.F.: Evaluation of nhs carbamates as a potent and selective class of endocannabinoid hydrolase inhibitors. ACS chemical neuroscience **4**(9), 1322–1332 (2013)

[43] Ignatowska-Jankowska, B., Ghosh, S., Crowe, M., Kinsey, S., Niphakis, M., Abdullah, R., Tao, Q., O'Neal, S., Walentiny, D., Wiley, J., *et al.*: In vivo characterization of the highly selective monoacylglycerol lipase inhibitor kml 29: antinociceptive activity without cannabimimetic side effects. British journal of pharmacology **171**(6), 1392–1407 (2014)

[44] Ignatowska-Jankowska, B.M., Baillie, G.L., Kinsey, S., Crowe, M., Ghosh, S., Owens, R.A., Damaj, I.M., Poklis, J., Wiley, J.L., Zanda, M., *et al.*: A cannabinoid cb1 receptor-positive allosteric modulator reduces

neuropathic pain in the mouse with no psychoactive effects. Neuropsychopharmacology **40**(13), 2948–2959 (2015)

[45] Marzo, V.D., Bifulco, M., Petrocellis, L.D.: The endocannabinoid system and its therapeutic exploitation. Nature reviews Drug discovery **3**(9), 771–784 (2004)

[46] Russo, E.B.: Cannabis therapeutics and the future of neurology. Frontiers in integrative neuroscience, 51 (2018)

[47] Russo, E.B.: Cannabinoids in the management of difficult to treat pain. Therapeutics and clinical risk management **4**(1), 245–259 (2008)

[48] Dewey, W.L.: Cannabinoid pharmacology. Pharmacological reviews **38**(2), 151–178 (1986)

[49] Devane, W.A., Hanuš, L., Breuer, A., Pertwee, R.G., Stevenson, L.A., Griffin, G., Gibson, D., Mandelbaum, A., Etinger, A., Mechoulam, R.: Isolation and structure of a brain constituent that binds to the cannabinoid receptor. Science **258**(5090), 1946–1949 (1992)

[50] Mechoulam, R., Ben-Shabat, S., Hanus, L., Ligumsky, M., Kaminski, N.E., Schatz, A.R., Gopher, A., Almog, S., Martin, B.R., Compton, D.R., *et al.*: Identification of an endogenous 2-monoglyceride, present in canine gut, that binds to cannabinoid receptors. Biochemical pharmacology **50**(1), 83–90 (1995)

[51] Long, J.Z., Nomura, D.K., Vann, R.E., Walentiny, D.M., Booker, L., Jin, X., Burston, J.J., Sim-Selley, L.J., Lichtman, A.H., Wiley, J.L., *et al.*: Dual blockade of faah and magl identifies behavioral processes regulated by endocannabinoid crosstalk in vivo. Proceedings of the National Academy of Sciences **106**(48), 20270–20275 (2009)

[52] Booker, L., Kinsey, S.G., Abdullah, R.A., Blankman, J.L., Long, J.Z., Ezzili, C., Boger, D.L., Cravatt, B.F., Lichtman, A.H.: The fatty acid amide hydrolase (faah) inhibitor pf-3845 acts in the nervous system to reverse lps-induced tactile allodynia in mice. British journal of pharmacology **165**(8), 2485–2496 (2012)

[53] Kinsey, S.G., Long, J.Z., O'Neal, S.T., Abdullah, R.A., Poklis, J.L., Boger, D.L., Cravatt, B.F., Lichtman, A.H.: Blockade of endocannabinoid-degrading enzymes attenuates neuropathic pain. Journal of Pharmacology and Experimental Therapeutics **330**(3), 902–910 (2009)

[54] Ghosh, S., Kinsey, S.G., Liu, Q.-s., Hruba, L., McMahon, L.R., Grim, T.W., Merritt, C.R., Wise, L.E., Abdullah, R.A., Selley, D.E., *et al.*:

Full fatty acid amide hydrolase inhibition combined with partial monoa-cylglycerol lipase inhibition: augmented and sustained antinociceptive effects with reduced cannabimimetic side effects in mice. Journal of Pharmacology and Experimental Therapeutics **354**(2), 111–120 (2015)

[55] Al-Zoubi, R., Morales, P., Reggio, P.H.: Structural insights into cb1 recep-tor biased signaling. International journal of molecular sciences **20**(8), 1837 (2019)

[56] Leo, L.M., Abood, M.E.: Cb1 cannabinoid receptor signaling and biased signaling. Molecules **26**(17), 5413 (2021)

[57] Mielnik, C.A., Lam, V.M., Ross, R.A.: Cb1 allosteric modulators and their therapeutic potential in cns disorders. Progress in Neuro-Psychopharmacology and Biological Psychiatry **106**, 110163 (2021)

[58] Wyatt, R.M., Fraser, I., Welty, N., Lord, B., Wennerholm, M., Sut-ton, S., Ameriks, M.K., Dugovic, C., Yun, S., White, A., *et al.*: Pharmacologic characterization of jnj-42226314,[1-(4-fluorophenyl) indol-5-yl]-[3-[4-(thiazole-2-carbonyl) piperazin-1-yl] azetidin-1-yl] methanone, a reversible, selective, and potent monoacylglycerol lipase inhibitor. Jour-nal of Pharmacology and Experimental Therapeutics **372**(3), 339–353 (2020)

[59] Mielnik, C., Lutelmowski, C., Johnson, C., Zebarth, J., Milenkovic, M., Horsfall, W., Swardfager, W., Bradshaw, H., Salahpour, A., Ross, R.: The unique effect of mjn110 (magl inhibitor) in hyperdopaminergic states: Implications for 2-ag modulation in psychosis. In: Abstract Booklet (2020)

[60] Catharine A. Mielnik, Lutelmowski, C., Salahpour, A., Bradshaw, H., Ross, R.: Monoacylglycerol Lipase (MAGL), But Not Fatty Amide Acid Hydrolase, Inhibition Exacerbates Hyperdopamin-ergic Phenotypes in Dopamine Transporter-Knockout (DATKO) Mice. In: The 31st Annual ICRS Symposium on the Cannabi-noids, International Convention Center Jerusalem, ISRAEL (2021). https://www.icrs.co/SYMPOSIUM.2021/ICRS2021.PROGRAMME.and.ABSTRACTS.pdf. Accessed 2021-05-12

[61] Christensen, R., Kristensen, P.K., Bartels, E.M., Bliddal, H., Astrup, A.: Efficacy and safety of the weight-loss drug rimonabant: a meta-analysis of randomised trials. The Lancet **370**(9600), 1706–1713 (2007)

[62] Topol, E.J., Bousser, M.-G., Fox, K.A., Creager, M.A., Despres, J.-P., Easton, J.D., Hamm, C.W., Montalescot, G., Steg, P.G., Pearson, T.A., *et al.*: Rimonabant for prevention of cardiovascular events (crescendo): a randomised, multicentre, placebo-controlled trial. The Lancet **376**(9740),

1243   517–523 (2010)
1288