SEVERE FIRE REGIMES DECREASE RESILIENCE OF ECTOTHERMIC POPULATIONS

A manuscript in preparation as a *Research Letter* for submission to the *Journal of Animal Ecology*

Heitor Campos de Sousa[1,*], Adriana Malvasio[1], Guarino Rinaldi Colli[4,**] & Roberto Salguero-Gómez[2,3,**]

[1] Universidade Federal do Tocantins – UFT, Quadra 109 Norte Av. NS-15, ALCNO-14, Plano Diretor Norte, CEP: 77001-090, Palmas, Tocantins, Brasil.

[2] Department of Zoology, University of Oxford, 11a Mansfield Rd, OX1 3SZ, UK.

[3] Evolutionary Demography Laboratory, Max Planck Institute for Demographic Research, Konrad Zuße straße 1, DE 18057, Germany.

[4] Universidade de Brasília – UnB, Departamento de Zoologia, Instituto de Ciências Biológicas, Avenida L4 Norte, Asa Norte, CEP: 70910-900, Brasília, Distrito Federal, Brasil.

* Contact author: heitor.sousa@uft.edu.br

** Shared senior authors

Type of article: Research article

Short running title: Demographic resilience of Cerrado lizards

Word count in abstract: 312; Word count in main text: 8,768; Number of figures: 5; Number of tables: 1; Number of references: 78.

Acknowledgments: This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001 and by a NERC Independent Research Fellowship to RSG (NE/M018458/1). HCS was supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) [grant numbers 88887.484511/2020-00 and 88881.623332/2021-01]. GRC thanks Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Fundação de Apoio à Pesquisa do Distrito Federal (FAPDF), and the USAID's PEER program under cooperative agreement AID-OAA-A-11-00012 for financial support.

Conflict of interest: Authors declare no conflict of interest.

Statement of authorship:

Conceptualization: HCS, AM, GRC, and RSG; Data curation: HCS and GRC; Formal analysis: HCS with assistance by RSG; Funding acquisition: GRC and RSG; Investigation: HCS and GRC; Methodology: HCS, GRC, and RSG; Project administration: GRC; Resources: GRC and RSG; Software: HCS and GRC; Supervision: AM, GRC, and RSG;

Validation: AM, GRC, and RSG; Visualization: HCS with assistance by RSG; Writing - original draft: HCS; Writing - reviewing and editing: AM, GRC, and RSG.

Data availability statement: We confirm that the data and code supporting the results will be archived in an appropriate public repository such as Dryad or Figshare, and the data DOI will be included at the end of the article.

Abstract (312 words)

1. Understanding populations' responses to environmental change is crucial for mitigating human-induced disturbances.

2. Here, we test hypotheses regarding how three essential components of demographic resilience (resistance, compensation, and recovery) co-vary along the distinct life histories of three lizard species exposed to variable, prescribed fire regimes.

3. Using a Bayesian hierarchical framework, we estimate vital rates (survival, growth, and reproduction) with 14 years of monthly individual-level data and mark-recapture models to parameterize stochastic Integral Projection Models from five sites in Brazilian savannas, each historically subjected to different fire regimes. With these models, we investigate how weather, microclimate, and ecophysiological traits of each species influence their vital rates, emergent life history traits, and demographic resilience components in varying fire regimes.

4. Overall, weather and microclimate are better predictors of the species' vital rates, rather than their ecophysiological traits. Our findings reveal that severe fire regimes increase populations' resistance, but decrease compensation or recovery abilities. Instead, populations have higher compensatory and recovery abilities at intermediate degrees of fire severity. Additionally, we identify generation time and reproductive output as predictors of resilience trends across fire regimes and climate. Our analyses demonstrate that the probability and quantity of monthly reproduction are the proximal drivers of demographic resilience across the three species.

5. Our findings suggest that populations surpass a tipping point in severe fire regimes and achieve an alternative stable state to persist. Thus, higher heterogeneity in fire regimes can increase the reproductive aspects and resilience of different populations and avoid high-severity regimes that homogenize the environment. Despite being

more resistant, species with long generation times and low reproductive output take

longer to recover and cannot compensate as much as species with faster paces of life.

We emphasize how reproductive constraints, such as viviparity and fixed clutch

sizes, impact the ability of ectothermic populations to benefit and recover from

disturbances, underscoring their relevance in conservation assessments.

Keywords: Demography, Disturbance, Ecophysiology, Environmental stochasticity, Environmental change,

Integral Projection Model, Life history, Reptile, Transient dynamics.

1 | INTRODUCTION

Environmental change is causing a fast decline in biodiversity worldwide (Newbold *et al.* 2020). The most impactful drivers of this loss include habitat degradation (Gonçalves-Souza, Verburg & Dobrovolski 2020) and the drastic modification of disturbance regimes (Hale *et al.* 2016), such as fires (Bowman *et al.* 2020). Understanding how species may respond to these impacts is at the core of ecology (Sutherland *et al.* 2013) and conservation biology (Sutherland *et al.* 2009). Ecological resilience research in the last decades has obtained variable success regarding how species traits can predict their responses to disturbances (Halpern 1989; Hu, Doherty & Jessop 2020). Indeed, some organismal traits confer species with higher tolerance or susceptibility to perturbations. Examples include mobility (Merrien *et al.* 2022), robustness (Dantas *et al.* 2013), rapid growth (Hoffmann *et al.* 2012), thermal performance (Hu, Doherty & Jessop 2020), and reproductive output (Capdevila *et al.* 2022). Therefore, deciphering which traits confer species the ability to resist and recover is critical to understanding and predicting how populations will respond to disturbances in the Anthropocene (Carmona *et al.* 2021).

Ecological resilience, the ability of an ecological system to resist and recover, or move to a new stable equilibrium from disturbances (Hodgson, McDonald & Hosken 2015), has been historically explored at the community and ecosystem levels (Kefi *et al.* 2019). Recently, though, key breakthroughs have been made in unraveling the mechanisms behind population responses to disturbances (Capdevila *et al.* 2020). Natural populations may respond to disturbances by avoiding decreases in size (resistance) or even by increasing in size (compensation). Following a transient period caused by the disturbance, populations may recover from said disturbance by transitioning back to the previous or a new stable stationary structure, at which point the composition of stages (*e.g.*, age, size classes) in the population remains relatively constant over time. Thus, we can quantify the full repertoire

of resilience responses to a disturbance by a natural population with three essential

components: (1) resistance, (2) compensation, and (3) recovery time. This novel

demographic resilience framework has the advantage of quantifying resilience based on the

species' vital rates (*i.e.*, survival, growth, and reproduction), allowing for inter- and intra-

specific comparisons, and providing a mechanistic understanding of the processes and trade-

offs involved in species' responses to disturbances (Capdevila *et al.* 2020).

Life history traits, key moments in an organism's life cycle (Stearns 1976), have

gained attention for their predictive potential in organismal responses (Allen, Street &

Capellini 2017). These traits, including generation time and net reproductive output, are

governed by survival, growth, and reproduction schedules and are subject to strong selection

for optimization (Stearns 2000). As these vital rates involve trade-offs, so do the emerging

life history traits (Stearns 1976). Understanding the combination of these traits, known as

life history strategies, is valuable in predicting population responses to environmental

changes and disturbances (Capdevila *et al.* 2021a). For instance, recent research has shown

that plant species with high reproductive output exhibit greater resistance and compensation

to disturbances (Iles *et al.* 2015), while longer-lived animals have higher resistance and

longer recovery times compared to fast-living species (Capdevila *et al.* 2022). Despite these

key findings, their generalization across the Tree of Life is challenging because organisms

present diverse life history traits (Salguero-Gómez *et al.* 2016a; Healy *et al.* 2019) and

because quantifying trade-offs in natural populations under control conditions is difficult

(Metcalf 2016), let alone in a context of disturbance regimes (Lytle 2001). Still, linking

mechanisms regulating vital rates to population-level responses to disturbances is crucial for

understanding how these trade-offs shape evolutionary processes.

Fire is a major disturbance that shapes ecosystems worldwide (Pausas & Parr 2018).

Fire consumes biomass, recycles nutrients, and triggers the renewal of vegetation, leading to

a burst of productivity and providing establishment opportunities for new individuals and

species (Pausas & Bond 2020). Despite regional heterogeneity in fire activity, the average

fire season had globally increased by 18.7% in the last three decades, and in many regions,

fire frequency, intensity, and spatial extent have also increased, presumably because of

climate change and human management (Bowman *et al.* 2020). In fire-prone ecosystems,

species typically develop adaptations to resist and recover from frequent fires (Keeley *et al.*

2011). Conversely, other species have evolved to take advantage of these disturbances.

Examples include fire-adapted plants that only resprout (Pausas, Keeley & Schwilk 2017)

or whose seed dormancy only breaks (Paniw *et al.* 2016) after a fire, thus benefiting from

the open space in temporarily less competitive environments. Historically, though, fire

ecology research has primarily focused on plant responses, leaving the mechanisms that

confer animals with resilience to fire poorly understood (but see Pausas, Keeley & Schwilk

2017).

To fill this knowledge gap, we investigate how the demographic resilience

components (resistance, compensation, and recovery time) co-vary with generation time,

reproductive output, and their underlying vital rates. To do so, we parameterize hierarchical

Bayesian Integral Projection Models (IPMs) with a 14-years of monthly mark-recapture data

from 3,538 individuals of three lizard species (*Copeoglossum nigropunctatum,*

Micrablepharus atticolus, and *Tropidurus itambere*) from the Brazilian Cerrado savannas

(Oliveira & Marquis 2002), with multiple populations exposed to prescribed fires with

varying frequencies and timings. Our three species have rather distinct life-history strategies

but co-exist in all of the study sites, thus creating a perfect *in situ* experimental laboratory to

explore how their life history traits and resilience components change in response to fire

regimes. *Copeoglossum nigropunctatum* (Scincidae) is a medium-sized (up to 150 mm

snout-vent length; SVL) viviparous skink found in Amazonia and Cerrado (Vitt & Blackburn

1991; Vitt, Zani & Lima 1997); its females can carry up to nine embryos for 10-12 months

(Vitt & Blackburn 1991). *Micrablepharus atticolus* (Gymnophthalmidae) is a small-sized

(up to 50 mm SVL) Cerrado endemic with a fixed clutch of 1-2 eggs (Vitt & Caldwell 1993;

Rodrigues 1996), but capable of producing multiple clutches (Vieira *et al.* 2000). *Tropidurus*

itambere (Tropiduridae) is a medium-sized (up to 100 mm SVL) species from central

Cerrado that lays multiple clutches of 5-8 eggs per season (Van Sluys & Henderson 1993).

All three species have generalist diets, primarily feeding on small invertebrates, with ants

being the main prey for *T. itambere* (Vitt & Blackburn 1991; Van Sluys 1993; Vieira *et al.*

2000). While most lizard species can withstand intense fires (Costa *et al.* 2013), our species

respond differently to fire-induced vegetation changes, influenced by microclimate and

microhabitat preferences (Costa *et al.* 2021).

Here, we test whether (H1) ecophysiological traits, such as thermal locomotor

performance and daily activity, and weather/microclimate shape how underlying vital rates

and emergent life history traits respond to fire regimes. We then quantify the extent to which

vital rates and life history traits drive demographic resilience. We hypothesize that: (H2)

more severe fire regimes decrease populations' resilience (*i.e.,* lower resistance, lower

compensation, and slower recovery); (H3) species' populations with greater generation time

will have higher resistance but slower recovery time; and (H4) populations with greater

reproductive output, higher body growth rate, and higher thermal performance will have

higher compensation and faster recovery time.

2 | MATERIALS AND METHODS

2.1 | Population monitoring

Our study site (*Reserva Ecológica do IBGE*, RECOR; see more details in the SI) is an ideal

natural experiment to test our hypotheses regarding the drivers of demographic resilience.

From 1972 to 1990, RECOR was protected from fires. In 1989, a long-term experiment was

initiated to evaluate the responses of animals and plants in 10 ha plots (200 x 500 m), each

submitted to unique prescribed fire regimes characterized by a combination of timing (early

dry season, in late June; mid dry season, in early August; and late dry season, in late

September) and frequency (biennial and quadrennial) (Miranda 2010). Plots were placed in

the cerrado *sensu stricto* physiognomy and shared a common recent fire history before the

experiment's onset. Thus, differences between plots over time are likely to have arisen due

to the prescribed fire regimes (Costa *et al.* 2021).

Here we focus on the three most abundant lizard species from the assemblage to

examine the traits that confer higher resilience to weather, microclimate, and fire regimes.

Of critical importance to test our hypotheses, each species presents different life history

strategies (driven by body sizes, reproductive modes, and clutch sizes). To acquire the

individual-level data used in our analyses, from November 2005 to December 2019, we

captured individual lizards in study plots using pitfall traps with drift fences in five plots

(Fig. S1): three under prescribed biennial fires (early biennial – EB, middle biennial – MB,

and late biennial – LB), one under prescribed quadrennial fires (middle quadrennial – Q),

and one control plot (C) (Fig. S1). Considering the number of fires each plot was submitted

and their canopy cover (Fig. S6), fire severity increased in the following order C < Q < EB

< MB < LB. We opened traps for six consecutive days every month and checked them daily

during this period. We recorded snout-vent length (SVL) with a ruler (1 mm precision) from

each captured individual, followed by marking by toe-clipping (after the appropriate

bioethics approval 33786/2016) and release at the capture site. Toe-clipping is a widely used

system to mark individuals in amphibians and reptiles, where each toe has a unique code to

set the identity of each individual (Hero 1989). We also assessed the reproductive state by

palpation in females. This assessment is possible because when female lizards bear eggs,

embryos, or enlarged follicles, they display a significant enlargement in their abdomens (Meiri, Brown & Sibly 2012).

Our approach considers that every plot comprises individual populations of each species, where individuals are subjected to different environments promoted by the long-term effects of fire regimes. Immigration and emigration, *i.e.*, movement into and movement out of a study area of arbitrary size and location, *sensu* Fryxell, Sinclair, and Caughley (2014), may bias vital rates estimates and the underlying effects of the experimental plots (Paquet *et al.* 2020), here fire regimes. However, considering the distance between transects in adjacent plots (*ca.* 200 m; Fig. S1A), the small size, and the territorial behavior of some species (Van Sluys 1997), we assumed no significant lizard movements among plots that could affect our results. Indeed, there were no inter-plot recaptures, supporting our assumption. In addition to the costs and difficulties of replication typical of large-scale ecological studies, our experiment could not be replicated because of legal issues associated with burning the vegetation inside protected areas. While this approach may reduce the statistical power of our study, doing so is the only way to ensure the availability of adequate treatment levels in such a large design (Driscoll *et al.* 2010).

2.3 | Ecophysiological traits, weather, and microclimate

To test (H1) whether ecophysiological traits and weather/microclimate shape how underlying vital rates and emergent life history traits respond to fire regimes, we estimated locomotor thermal performance curves (TPC) (Huey & Stevenson 1979) and hours of activity (*ha*) (Sinervo *et al.* 2010) based on laboratory experiments from at least 10 individuals by each species (see SI for more details). We initially had 14 microclimatic and weather variables. However, after excluding highly correlated variables (using the stepwise procedure *vifcor* from package USDM), we retained: mean air temperature at 200 cm height

(*tmed2m*), minimum air temperature at 0 cm height (*tmin0cm*), mean air temperature at 0 cm

height (*tmed0cm*), maximum relative air humidity (*RHmax*), solar radiation (*sol*), and

accumulated precipitation (*precip*). The *vifcor* algorithm finds a pair of variables highly

correlated (cor > 0.8) and excludes the one with a greater variance inflation factor (VIF)

repeatedly until no pair of variables highly correlated remains. We used this approach

because one of our primary goals is to understand the specific contributions of individual

environmental factors to the vital rates (H1) rather than composite axes of environmental

variation (such as from a principal component analysis). With the hourly microclimate

temperature estimates at 10 cm height, we predicted monthly mean locomotor performance

(*perf*) and the hours of activity (*ha*). SI presents the correlations and relationships among the

variables (Fig. S7) and the details of the microclimatic and thermal physiological estimates.

2.4 | Data analyses

We estimated each species' monthly vital rates (survival, growth, and reproduction) using

different datasets integrated into Bayesian hierarchical models and a combination of

Cormack-Jolly-Seber (CJS) and Pradel Jolly-Seber (PJS) mark-recapture models of open

populations (Cormack 1964; Pradel 1996; Tenan *et al.* 2014). Because the CJS model allows

for individuals' covariates, we related the survival (σ) to the individuals' body size (SVL),

while the PJS incorporated the monthly environmental variation (weather, ecophysiological,

microclimate, months with fire, and time since last fire – *TSLF*) into populations' survival

(σ) and recruitment (probability of recruitment, p_{rec}). The recruitment parameter (p_{rec}) is

based on the total population (males and females) and can be interpreted as mostly hatchling

recruitment since their peaks coincide with captures of small-sized hatchlings (Fig. S8).

Additionally, our mark-recapture models are sex-averaged since we have not found

differences in survival and growth between males and females (Fig. S9). For the changes in

the body size through time (growth, γ), we combined a Von Bertalanffy body growth

function with the individuals' records of SVL and capture history. For reproduction, we

considered three different processes: the probability of reproduction (p_{rep}), the production of

newborns (clutch size; n_b), and the probability of recruitment (p_{rec}). We used Bayesian

generalized linear models to relate p_{rep} and n_b to the individuals' SVL. We implemented the

models for each species with JAGS in R, using the packages JAGSUI, RUNJAGS, and RJAGS

(Kellner 2019; Plummer 2019). SI presents the details of the parameterizations estimated in

each model, including their formulas.

To test H2, H3, and H4 (see Introduction), we used the vital rates (σ, γ, p_{rep}, n_b, and

p_{rec}) parameter estimates to build stochastic Integral Projection Models (IPMs) using the

IPMR package (Levin *et al.* 2021). Although we have not parameterized the IPMs with

explicit density-dependent parameters constraining their vital rates, we assumed that the

environmental covariates and plots represented the primary resources the lizards use (habitat,

thermal niche, and food), thus indirectly modeling density dependence; this is a common

approach when analyzing long-term demographic data (Salguero-Gómez *et al.* 2016a), as it

is our case. IPMs are structured population models where the vital rates are influenced by a

continuously varying phenotypic trait (*e.g.*, body size) instead of discrete stages or ages

(Easterling, Ellner & Dixon 2000). In a population structured by a continuous variable body

size (SVL), as in our three study species, the IPM describing the number of individuals of

size y at time $t + 1$ is:

230
$$n(y, t+1) = \int_{L}^{U} [P(x,y) + F(x,y)]n(x,t)dx$$

231
$$= \int_{L}^{U} K(x,y)n(x,t)dx \qquad \text{(Eq. 1)}.$$

In equation 1, L and U specify the lower and upper body size limits in the model,

respectively; $P(x,y)$ describes the changes in size conditional on survival; $F(x,y)$ quantifies

fecundity; and $K(x,y) = P(x,y) + F(x,y)$ is the *kernel* surface, representing all possible

transitions and per-capita sexual contributions from size x to size y between t and $t+1$. The

P sub-kernel incorporates both survival and growth components, such that $P(x,y) = \sigma(x) \cdot$

$\gamma(x,y)$, where $\sigma(x)$ is the survival probability of a size-x individual and $\gamma(x, y)$ is the probability

of a size-x individual growing to size y (see SI for the IPMs' implementation). Considering

the five plots and 169 months, excluding the last month that mark-recapture models cannot

estimate (Cormack 1964; Pradel 1996), we generated 845 monthly IPM kernels for each

species.

To test whether populations' generation time (H3) and reproductive output (H4)

shape their demographic resilience, we estimated those two life history traits with the

monthly IPM kernels of each plot and species using the package RAGE (Jones *et al.* 2022).

We also estimated other life-history traits from these IPMs, but we decided to focus our

analyses on generation time and net reproductive output only, as they describe the principal

axes of the life history variation in our species (Figs. S13-S14), as well as in other animals

(Gaillard *et al.* 2005; Healy *et al.* 2019), including reptiles (Rodriguez-Caro *et al.* 2023).

To test our hypotheses regarding the demographic resilience components (H2-H4),

we estimated the resistance, compensation, and recovery time from our IPMs. To do so, we

used the package POPDEMO (Stott, Hodgson & Townley 2012) on our monthly, species- and

site-specific IPMs. We used the first time-step amplification (reactivity) and attenuation as

the measures for compensation and resistance, respectively, following Capdevila *et al.*

(2020). First time-step amplification and attenuation are the highest population increase and

the lowest population decrease possible in the next month, respectively. Attenuation values

vary from 0 to 1, where 0 means low resistance and 1 means high resistance. Thus, given the

species, sites, and monthly environmental covariates (weather, microclimate,

ecophysiological, fire-related, and randomness), resistance and compensation estimates

represent the monthly potential of each population to maintain and increase its size in the

next month, respectively. Recovery time was estimated as follows:

261
$$t_K = \frac{\log\left(\frac{\lambda_1}{\|\lambda_2\|}\right)}{\log(10)} \qquad \text{(Eq. 2)}.$$

In equation 2, λ_1 and λ_2 correspond to the dominant and subdominant eigenvalues of the K

kernel, respectively. This measure estimates the time (months) required to reach a stable

stationary structure every month.

Next, to test (H2) whether more severe fire regimes decrease populations' resilience

populations, and populations with (H3) greater generation time have higher resistance but

slower recovery, and (H4) populations with greater reproductive output have higher

compensation and faster recovery, we built Bayesian dynamic generalized additive models

using MVGAM (Clark & Wells 2023). We related the demographic resilience components

with the two life history traits and among plots. We used gamma error distributions for

compensation and recovery time, and for resistance, we used a beta distribution. For each

population time series (by plot and species), we included plot as a random variable to control

for pseudoreplication and non-independency among plots and repeated observations. We

also included a latent autoregressive and moving-average trend model ($ARIMA$) to capture

the first-lag autocorrelation. We $\log_{10}$-transformed the compensation values and z-scaled the

life history traits values prior to analyses.

To test specific aspects of H4, that populations with higher body growth rate and

higher thermal performance will have higher compensation and faster recovery time and, at

the same time, inspect which IPM vital rate parameters affected the different demographic

resilience components, we conducted a brute-force sensitivity analysis (Morris & Doak

2002). To implement the brute force sensitivity analysis, we added values from 0.01 to 0.1

to each of the parameters specifying our vital rates and recalculated the differences in the

demographic resilience components. Our parameter sensitivities (S_p) were estimated as in

equation 3, where the numerator describes the difference in the transient metric of choice (x;

e.g., reactivity for amplification) resulting from perturbing a vital rate parameter p at a time,

in the denominator, following Cant *et al.* (2023):

287
$$S_p = \frac{x_{original} - x_{perturbed}}{p_{original} - p_{perturbed}} \qquad \text{(Eq. 3)}.$$

3 | RESULTS

3.1 | Vital rates

Overall, we found that weather and microclimate shape the survival and recruitment of

Micrablepharus atticolus and *Tropidurus itambere*, but not of *Copeoglossum*

nigropunctatum (Fig. 1), partially supporting our hypothesis (H1). As opposed to H1,

ecophysiological traits were poor predictors of the vital rates, except for the recruitment of

M. atticolus. For instance, the vital rates of survival and recruitment of *C. nigropunctatum*

were better explained by random monthly variation than environmental covariates (Fig. 1;

Table S4). In contrast, survival of *M. atticolus* and *T. itambere* was only affected by weather

(*precip*, *tmed2m*, and *sol*; Fig. 1; Table S4). Partially supporting our predictions (H1),

weather (*tmed2m*), microclimatic (*tmin0cm*), and ecophysiological (*perf* and *ha*) conditions

affected the recruitment of *M. atticolus* (Fig. 1; Table S4); Recruitment of *T. itambere* only

varied with weather conditions (*tmed2m* and *sol*; Fig. 1; Table S4). These patterns resulted

in higher variation in the life-history space in these two species—*M. atticolus* and *T.*

itambere (Fig. S13). We found differences in these vital rates among plots, which had been

exposed to prescribed fire regimes for 18 years (Table S4). The differences in vital rates

were most apparent under both extremes of fire severity (absence of fire and frequent late-

dry season fires), suggestive of the long-term effects of the fire regimes (Table S4), in

addition to differences in microclimate and ecophysiological traits (H1). However, the other

fire-related variables (*fire* and *TSLF*) were poor predictors of the vital rates, except for the

recruitment of *M. atticolus* (Fig. 1; Table S4). See SI for mark-recapture summary statistics.

3.2 | Fire regimes and demographic resilience

Contrary to H2, populations of the three species presented high resistance ability in

severe fire regimes (Figs. 2-3). However, populations of *T. itambere* displayed low

compensation in the most severe fire regime (Figs. 2-3). On average, the three lizard species

recover rather fast to their stable stationary structure (Figs. 2-3; *C. nigropunctatum* - 1.844

$\pm$ 0.064 S.D. months; *M. atticolus* - 1.516 $\pm$ 0.291 S.D. months; *T. itambere* - 1.946 $\pm$ 0.200

S.D. months). However, they followed clearly differentiated strategies depending on the fire

severity, resulting in the lowest resistance, highest compensation, and fastest recovery

abilities at intermediate levels of severity (third and fourth levels–frequent early- and mid-

dry season fires; Fig. 3). The most striking differences, though, occurred at the extreme of

fire severity (frequent late-dry season fires): *M.atticolus* displayed higher resistance and

compensation, and *T. itambere* higher compensation and lower resistance (Fig. 3).

Interestingly, *M. atticolus* and *T. itambere* also lost the ability to recover fast (*i.e.*, < 1.5

months) from fire and climate disturbances in the most severe fire regime (Fig. 3).

The fire regime effects on the demographic resilience components highly agree with

the realized population growth rates and abundance trends in the three species. All

populations had mean positive population growth ($log(\lambda) > 0$) throughout the 14 years of

study (Fig. S11). However, *C. nigropunctatum* displayed lower stability (higher S.D. in

$log(\lambda)$) in the intermediate level of fire severity (third level–frequent early-dry season fires),

while *M. atticolus* and *T. itambere* populations varied more (higher S.D. in $log(\lambda)$) in the

most severe fire regimes (fourth and fifth levels, SI; Fig. S11). Overall, *C. nigropunctatum*

had fewer captures in intermediate fire severity levels (third and fourth levels–frequent early-

and mid-dry season fires), while *M. atticolus* and *T. itambere* displayed the opposite pattern

(Fig. S2). Both *M. atticolus* and *T. itambere* had low captures at the extremes of fire severity

(absence of fires and frequent late-dry season fires; Fig. S2).

3.3 | Life history traits and demographic resilience

Using Bayesian dynamic generalized additive models, we found that the differences

in the demographic resilience components between fire regimes were predicted by

generation time and reproductive output (Fig. S15), supporting H3 and H4. However, the

direction of some of our predictions differed for the species with fixed clutch size, *M.*

atticolus. We found that, in populations with higher generation times, resistance was higher,

compensation was lower, and recovery was longer (Fig. 4). This striking pattern was

consistent across the three examined species but with a weaker relationship in *C.*

nigropunctatum due to its constrained vital rate variation in response to the environment,

and thus lower variance in life history traits (see Figs. S13-S15 for different life history traits

among the species). An important exception to the overall pattern was again *M. atticolus*,

with its fixed clutch size, where populations with higher generation times displayed lower

resistance to fire and climate variation.

Reproductive output also shaped the demographic resilience components of the three

species, supporting H4. The opposite pattern regarding the relationship between resilience

components and life history traits was found with reproductive output (Fig. 5). Indeed, in

populations with higher reproductive output, resistance was lower, compensation was

higher, and recovery was faster (Fig. 5). This pattern was consistent across species, with the

exception of the fixed-clutch species *M. atticolus*, displaying higher resistance and lower

compensation with increasing reproductive output (Fig. 5).

3.4 | Vital rate parameters and demographic resilience

The brute-force sensitivity analysis revealed that the demographic resilience components are most affected by the reproductive parameters (Table 1) and, to a lesser extent, by body growth; but not by thermal physiology as we predicted (H4). The relationships between the probability of reproduction or the number of newborns and SVL regulated all the resilience components in the three species (Table 1). This result indicates that reproduction among the largest individuals (usually the oldest, (Halliday & Verrell 1988)) is critical for the demographic resilience of *C. nigropunctatum* and *T. itambere*. Higher vital rate parameter values describing allocation in reproduction led to lower resistance and higher compensation (Table 1). However, complex results emerged regarding the vital rate parameters shaping recovery time. In *T. itambere*, a higher probability of reproduction and a higher number of newborns resulted in a displacement of the population structure further away from the stable size distribution, leading to longer recovery times (Table 1). In *C. nigropunctatum*, we found the opposite pattern, except for the probability of reproduction (which presented complex non-linear relationships). In *M. atticolus,* a higher probability of reproduction of larger adult individuals shortened recovery times (Table 1). As we predicted, slower body growth rates led to slower recovery time, especially for the small-sized *M. atticolus* (Table 1). Other parameters had negligible effects on the demographic resilience components across species and fire regimes.

4 | DISCUSSION

Using 14 years of monthly mark-recapture data of three lizard species from the Brazilian Cerrado savannas, we show that severe disturbance regimes result in overall less resilient populations, supporting our primary hypothesis (H2). Indeed, our stochastic IPMs demonstrate that high-severity fire regimes increase resistance but decrease compensatory

or recovery abilities, leading to less stable populations. Accordingly, we also show that

generation time and reproductive output predict the differences in the demographic resilience

components (resistance, compensation, recovery time) among fire regimes and species.

Here, populations with higher generation times are characterized by a greater degree of

resistance and slower recovery time (H3), whereas those with higher reproductive output

tend to compensate more and recover faster (H4). The only species with different

relationships between these life history traits and demographic resilience components is the

small body-sized lizard *Micrablepharus atticolus,* which has a fixed clutch size of two eggs.

Despite these differences, all species present a strong negative correlation between resistance

and compensation to fires and climatic stochasticity. Partially supporting our hypothesis, the

species' probability and quantity of reproduction, but not its ecophysiology, are the main

drivers of the demographic resilience in our three ectothermic species (H1), with body

growth only being a strong predictor of recovery time in *M. atticolus*.

The most severe fire regime results in more resistant populations in our three species.

However, our lizards display longer recovery times (~ two months) or lower compensation

in response to this severe fire regime. This finding suggests that these populations surpass a

tipping point in severe fire regimes and achieve an alternative stable state to persist, as shown

in other systems, such as in shrubs and grasses from tallgrass prairies (Collins *et al.* 2021).

Here, the new state after intense fires is characterized by strategies that allocate resources to

either side along the fast-slow continuum (Stearns 1976) or the reproductive strategies

continuum (Salguero-Gómez *et al.* 2016b). For instance, populations of *T. itambere* display

longer generation times but lower reproductive output in the most severe fire regime.

Similarly, *M. atticolus* also presents lower reproductive output but a faster pace of life (*i.e.*,

lower generation time) in this regime. These findings further highlight the plasticity of

species resilience strategies to cope with different levels of disturbance so populations may

remain viable, an area that has received considerable attention in recent years (Vinton *et al.*

2022). Said plasticity seems particularly high in reptiles, which have not diminished their

ability to respond to disturbances in the last decades compared to other vertebrates, such as

mammals and birds (Capdevila *et al.* 2021a).

Our findings align with recent examinations of the relationship between the

demographic resilience of plants and animals with the fast-slow continuum (Stott *et al.* 2010;

Jiang *et al.* 2022). Indeed, recent studies have recently shown that populations with longer

generation times tend to display a lower ability to compensate and are slower to recover after

disturbances (Stott *et al.* 2010). In addition, here we show a clear relationship between

demographic resilience and reproductive strategy, an orthogonal axis to the fast-slow

continuum (Salguero-Gómez *et al.* 2016b; Healy *et al.* 2019). For instance, our populations

with higher reproductive output resist less, compensate more (except the species with fixed

clutch size, *M. atticolus*), and recover faster after disturbances. This reactive response to

disturbances agrees with work showing that the ability to compensate, linked to reproductive

output too, enhances colonizing and invasive potential among reptiles and amphibians

(Allen, Street & Capellini 2017), and even in plants (Iles *et al.* 2015).

Multiple calls have been made for a more mechanistic understanding of resilience in

ecological systems (Hodgson, McDonald & Hosken 2015; Kefi *et al.* 2019; Capdevila *et al.*

2020); these calls remain largely unanswered (Capdevila *et al.* 2021b). Here, we contribute

to this knowledge gap by mechanistically integrating into the local population dynamics of

each species their ecophysiology, size structure, environmental factors, and varying

disturbance regimes. Indeed, one of our key findings is that probability and quantity of

monthly reproduction are key drivers of demographic resilience. The lack of predictive

power of ecophysiological characteristics (*e.g.*, thermal performance, preferred temperature)

on resilience reinforces the idea that life history traits constitute an excellent approach to

understanding demographic resilience, as recently highlighted in a synthesis across 69

animal and 232 plant species (Capdevila *et al.* 2022). In addition, our results suggest that

resilience components exhibit strong intra- and inter-specific variation and that these

relationships are strongly modulated by species' vital rates and life history characteristics.

Thus, we argue that existing macroecological and comparative approaches (Capdevila *et al.*

2022) likely mask some of the intraspecific trade-offs we disentangle here using 14 years of

monthly mark-recapture data under five levels of fire disturbances. Future studies will likely

deliver important insights by testing the predictive power of various kinds of traits (*e.g.*,

morphological, physiological, life history) across a wider range of species and disturbance

levels.

Fast growth increases resilience after disturbances (Pausas & Bond 2020) and

enhances persistence in novel environments (Lytle 2001; Allen, Street & Capellini 2017). In

line with these findings, we show that strategies leading to earlier reproduction are effective

at conferring compensation and faster recovery against fires and climate variation in our

three lizard species. We suggest that this strategy may be particularly important for small

animals, as it may allow them to achieve sexual maturity earlier, like in the case of *M.*

atticolus. In contrast, we show that in months when individuals allocate more resources to

reproduction, populations lose the ability to resist disturbances. This pattern presumably

results from individuals' reproductive costs (Schwarzkopf & Shine 1992); thus, by delaying

reproduction, individuals of indeterminate growth—most reptiles (Halliday & Verrell

1988)—can continue to increase survival and reproductive rates at older ages (Halliday &

Verrell 1988; Healy *et al.* 2019), buffering the population against disturbances and

environmental stochasticity (Koons, Metcalf & Tuljapurkar 2008). The timing of

disturbances may affect population resilience and explain why the timing of reproduction

(*i.e.*, phenology) has a critical effect on the responses of lizards (Braithwaite 1987) and plants

(Zirondi, Ooi & Fidelis 2021) to fires prescribed in different seasons. This connection

between phenology and the timing of disturbances is increasingly relevant to other highly

seasonal environments, where species' vital rates are adapted to predictable environmental

changes (Prather *et al.* 2023).

Since reproduction is a critical vital rate conferring resilience, species with

morphological and physiological constraints on reproduction (*e.g.*, viviparity and fixed

clutch sizes) have limited ability to benefit from disturbances. This prediction is precisely

what we show in *C. nigropunctatum* and *M. atticolus*. However, both species relatively

sustained their resilience strategies in varying degrees of fire severity. Recently reported

links between demographic buffering and transient dynamics after disturbances

(demographic resilience) (Capdevila *et al.* 2020) may explain these patterns (Stott *et al.*

2010; McDonald *et al.* 2016). In reptiles, viviparity developed as an adaptation to increase

reproduction success in cold climates (Zimin *et al.* 2022), and might also confer species with

the ability to buffer populations against environmental stochasticity, including resilience to

disturbances. Therefore, demographically buffering species may outcompete others in the

most extreme fire regimes (total absence of fire *vs.* frequent and severe fires), as we show

here. However, because of the constraints on their reproductive schedules, individuals from

viviparous and fixed clutch-size species need to perform other strategies to increase their

reproductive output and to compensate after fires, such as reproducing more often (Cayuela

et al. 2022). Indeed, these results provide the possible mechanisms for why, among 75,000

species of vascular plants and tetrapods, long-lived and low-reproductive species present

higher extinction risks than fast-living and high-reproductive ones (Carmona *et al.* 2021).

Our findings have key management and conservation implications. First, easily

measurable traits such as probability of reproduction and clutch size predict species'

resilience to environmental stochasticity and fires. Despite being more resistant, species with

high generation time and low reproductive output take longer to recover and cannot

compensate as much as species with faster paces of life. In our study, populations have

higher resilience at low and intermediate degrees of fire severity. Therefore, these results

support the pyrodiversity hypothesis, whereby higher heterogeneity in fire regimes increases

the resilience of different populations and communities and avoids high-severity regimes

that homogenize the environment (Jones & Tingley 2021). Second, despite species persisting

in highly severe fire regimes, populations can attain alternative stable states (Collins *et al.*

2021), with limited compensatory and recovery abilities after disturbances. Thus, traditional

ecological community studies probably underestimate disturbance impacts since populations

may persist in severe regimes but not in optimal conditions. Third, reproduction is a key

mechanism conferring higher compensation and faster recovery time, both in our species

and others (Iles *et al.* 2015; Allen, Street & Capellini 2017). Accordingly, species with

constraints on reproduction (*e.g.*, viviparity, fixed clutch size) have higher resistance but

lower compensation and take longer to recover from disturbances. Taken together, these key

findings imply that, to enhance viable strategies after disturbances, managers should focus

on generating suitable and heterogeneous habitats to facilitate the reproductive aspects of

populations. The ability to quantify and predict the resilience components of natural

populations based on their vital rates and life history traits will be increasingly critical for

conservation biology in the coming decades when climate change and disturbances are

projected to increase in frequency and intensity (Bowman *et al.* 2020).

REFERENCES

Allen, W.L., Street, S.E. & Capellini, I. (2017) Fast life history traits promote invasion

success in amphibians and reptiles. *Ecology Letters,* **20,** 222-230.

Bowman, D.M.J.S., Kolden, C.A., Abatzoglou, J.T., Johnston, F.H., van der Werf, G.R. &

Flannigan, M. (2020) Vegetation fires in the Anthropocene. *Nature Reviews Earth*

& Environment, **1,** 500-515.

Braithwaite, R.W. (1987) Effects of fire regimes on lizards in the wet-dry tropics of

Australia. *Journal of Tropical Ecology,* **3,** 265-275.

Cant, J., Capdevila, P., Beger, M. & Salguero-Gomez, R. (2023) Recent exposure to

environmental stochasticity does not determine the demographic resilience of

natural populations. *Ecology Letters,* **26,** 1186-1199.

Capdevila, P., Noviello, N., McRae, L., Freeman, R. & Clements, C.F. (2021a) Global

patterns of resilience decline in vertebrate populations. *Ecology Letters***,** 1-12.

Capdevila, P., Stott, I., Beger, M. & Salguero-Gómez, R. (2020) Towards a comparative

framework of demographic resilience. *Trends in Ecology & Evolution,* **35,** 776-786.

Capdevila, P., Stott, I., Cant, J., Beger, M., Rowlands, G., Grace, M. & Salguero-Gómez,

R. (2022) Life history mediates the trade-offs among different components of

demographic resilience. *Ecology Letters,* **22,** 1566-1579.

Capdevila, P., Stott, I., Oliveras Menor, I., Stouffer, D.B., Raimundo, R.L.G., White, H.,

Barbour, M. & Salguero-Gómez, R. (2021b) Reconciling resilience across

ecological systems, species and subdisciplines. *Journal of Ecology,* **109,** 3102-

3113.

Carmona, C.P., Tamme, R., Pärtel, M., de Bello, F., Brosse, S., Capdevila, P., González-

M, R., González-Suárez, M., Salguero-Gómez, R., Vásquez-Valderrama, M. &

Toussaint, A. (2021) Erosion of global functional diversity across the tree of life.

Science Advances, **7,** eabf2675.

Cayuela, H., Monod-Broca, B., Lemaitre, J.F., Besnard, A., Gippet, J.M.W., Schmidt,

B.R., Romano, A., Hertach, T., Angelini, C., Canessa, S., Rosa, G., Vignoli, L.,

532   Venchi, A., Carafa, M., Giachi, F., Tiberi, A., Hantzschmann, A.M., Sinsch, U.,

533    Tournier, E., Bonnaire, E., . . . Lena, J.P. (2022) Compensatory recruitment allows

534    amphibian population persistence in anthropogenic habitats. *Proceedings of the*

535    *National Academy of Sciences, USA,* **119,** e2206805119.

536  Clark, N.J. & Wells, K. (2023) Dynamic generalised additive models (DGAMs) for

537    forecasting discrete ecological time series. *Methods in Ecology and Evolution,* **14,**

538    771-784.

539  Collins, S.L., Nippert, J.B., Blair, J.M., Briggs, J.M., Blackmore, P. & Ratajczak, Z. (2021)

540    Fire frequency, state change and hysteresis in tallgrass prairie. *Ecology Letters,* **24,**

541    636-647.

542  Cormack, R.M. (1964) Estimates of Survival from the Sighting of Marked Animals.

543    *Biometrika,* **51,** 429-438.

544  Costa, B.M., Pantoja, D.L., Sousa, H.C., de Queiroz, T.A. & Colli, G.R. (2021) Long-term,

545    fire-induced changes in habitat structure and microclimate affect Cerrado lizard

546    communities. *Biodiversity and Conservation,* **29,** 1659-1681.

547  Costa, B.M., Pantoja, D.L., Vianna, M.C.M. & Colli, G.R. (2013) Direct and short-term

548    effects of fire on lizard assemblages from a Neotropical savanna hotspot. *Journal of*

549    *Herpetology,* **47,** 502-510.

550  Dantas, V.d.L., Pausas, J.G., Batalha, M.A., de Paula Loiola, P. & Cianciaruso, M.V.

551    (2013) The role of fire in structuring trait variability in Neotropical savannas.

552    *Oecologia,* **171,** 487-494.

553  Driscoll, D.A., Lindenmayer, D.B., Bennett, A.F., Bode, M., Bradstock, R.A., Cary, G.J.,

554    Clarke, M.F., Dexter, N., Fensham, R., Friend, G., Gill, M., James, S., Kay, G.,

555    Keith, D.A., MacGregor, C., Russell-Smith, J., Salt, D., Watson, J.E.M., Williams,

556    R.J. & York, A. (2010) Fire management for biodiversity conservation: Key

research questions and our capacity to answer them. *Biological Conservation,* **143,**

1928-1939.

Easterling, M.R., Ellner, S.P. & Dixon, P.M. (2000) Size-specific sensitivity: Applying a

new structured population model. *Ecology,* **81,** 694-708.

Gaillard, J.M., Yoccoz, N.G., Lebreton, J.D., Bonenfant, C., Devillard, S., Loison, A.,

Pontier, D. & Allaine, D. (2005) Generation time: a reliable metric to measure life-

history variation among mammalian populations. *American Naturalist,* **166,** 119-

123.

Gonçalves-Souza, D., Verburg, P.H. & Dobrovolski, R. (2020) Habitat loss, extinction

predictability and conservation efforts in the terrestrial ecoregions. *Biological*

Conservation, **246,** 108579-108579.

Hale, S., Nimmo, D.G., Cooke, R., Holland, G., James, S., Stevens, M., De Bondi, N.,

Woods, R., Castle, M., Campbell, K., Senior, K., Cassidy, S., Duffy, R., Holmes,

B. & White, J.G. (2016) Fire and climatic extremes shape mammal distributions in

a fire-prone landscape. *Diversity and Distributions,* **22,** 1127-1138.

Halliday, T.R. & Verrell, P.A. (1988) Body size and age in amphibians and reptiles.

Journal of Herpetology, **22,** 253-265.

Halpern, C.B. (1989) Early successional patterns of forest species: interactions of life

history traits and disturbance. *Ecology,* **70,** 704-720.

Healy, K., Ezard, T.H.G., Jones, O.R., Salguero-Gómez, R. & Buckley, Y.M. (2019)

Animal life history is shaped by the pace of life and the distribution of age-specific

mortality and reproduction. *Nature Ecology and Evolution,* **3,** 1217-1224.

Hero, J.M. (1989) A simple code for toe clipping anurans. *Herpetological Review,* **20,** 66-

69.

Hodgson, D., McDonald, J.L. & Hosken, D.J. (2015) What do you mean, 'resilient'? *Trends*

in Ecology & Evolution, **30,** 503-506.

Hoffmann, W.A., Geiger, E.L., Gotsch, S.G., Rossatto, D.R., Silva, L.C.R., Lau, O.L.,

Haridasan, M. & Franco, A.C. (2012) Ecological thresholds at the savanna-forest

boundary: How plant traits, resources and fire govern the distribution of tropical

biomes. *Ecology Letters,* **15,** 759-768.

Hu, Y., Doherty, T.S. & Jessop, T.S. (2020) How influential are squamate reptile traits in

explaining population responses to environmental disturbances? *Wildlife Research,*

47, 249-259.

Huey, R.B. & Stevenson, R.D. (1979) Integrating thermal physiology and ecology of

ectotherms: a discussion of approaches. *American Zoologist,* **19,** 357-366.

Iles, D.T., Salguero-Gómez, R., Adler, P.B., Koons, D.N. & Zuidema, P. (2015) Linking

transient dynamics and life history to biological invasion success. *Journal of*

Ecology, **104,** 399-408.

Jiang, S., Jaggi, H., Zuo, W., Oli, M.K., Coulson, T., Gaillard, J.M. & Tuljapurkar, S.

(2022) Reproductive dispersion and damping time scale with life-history speed.

Ecology Letters, **25,** 1999-2008.

Jones, G.M. & Tingley, M.W. (2021) Pyrodiversity and biodiversity: A history, synthesis,

and outlook. *Diversity and Distributions*, 1-18.

Jones, O.R., Barks, P., Stott, I., James, T.D., Levin, S., Petry, W.K., Capdevila, P., Che-

Castaldo, J., Jackson, J., Römer, G., Schuette, C., Thomas, C.C. & Salguero-

Gómez, R. (2022) Rcompadre and Rage—Two R packages to facilitate the use of

the COMPADRE and COMADRE databases and calculation of life-history traits

from matrix population models. *Methods in Ecology and Evolution,* **13,** 770-781.

Keeley, J.E., Pausas, J.G., Rundel, P.W., Bond, W.J. & Bradstock, R.a. (2011) Fire as an

evolutionary pressure shaping plant traits. *Trends in Plant Science,* **16,** 406-411.

Kefi, S., Dominguez-Garcia, V., Donohue, I., Fontaine, C., Thebault, E. & Dakos, V.

(2019) Advancing our understanding of ecological stability. *Ecology Letters,* **22,**

1349-1356.

Kellner, K. (2019) jagsUI: a wrapper around 'rjags' to streamline 'JAGS' analyses.

Koons, D.N., Metcalf, C.J. & Tuljapurkar, S. (2008) Evolution of delayed reproduction in

uncertain environments: a life-history perspective. *American Naturalist,* **172,** 797-

805.

Levin, S.C., Childs, D.Z., Compagnoni, A., Evers, S., Knight, T.M. & Salguero-Gómez, R.

(2021) ipmr: Flexible implementation of Integral Projection Models in R. *Methods*

in Ecology and Evolution, **2021,** 1826-1834.

Lytle, D.A. (2001) Disturbance regimes and life-history evolution. *The American*

Naturalist, **157,** 525-525.

McDonald, J.L., Stott, I., Townley, S. & Hodgson, D.J. (2016) Transients drive the

demographic dynamics of plant populations in variable environments. *Journal of*

Ecology, **104,** 306-314.

Meiri, S., Brown, J.H. & Sibly, R.M. (2012) The ecology of lizard reproductive output.

Global Ecology and Biogeography, **21,** 592-602.

Merrien, T., Davis, K.J., Di Marco, M., Capdevila, P. & Salguero-Gómez, R. (2022)

Human disturbances erode the diversity of species resilience strategies. *BioRxiv*.

Metcalf, C.J. (2016) Invisible trade-offs: Van Noordwijk and de Jong and life-history

evolution. *American Naturalist,* **187,** iii-v.

Miranda, H.S. (2010) *Efeitos do regime do fogo sobre a estrutura de comunidades de*

cerrado: Resultados do Projeto Fogo. IBAMA, Brasília, Brasil.

Morris, W.F. & Doak, D.F. (2002) *Quantitative Conservation Biology: Theory and*

Practice of Population Viability Analysis. Sinauer Associates Inc., Sunderland,

USA.

Newbold, T., Bentley, L.F., Hill, S.L.L., Edgar, M.J., Horton, M., Su, G., Şekercioğlu,

Ç.H., Collen, B. & Purvis, A. (2020) Global effects of land use on biodiversity

differ among functional groups. *Functional Ecology,* **34,** 684-693.

Oliveira, P.S. & Marquis, R.J. (2002) *The cerrados of Brazil: Ecology and Natural History*

of a Neotropical Savanna. Columbia University Press, New York.

Paniw, M., Quintana-Ascencio, P.F., Ojeda, F. & Salguero-Gómez, R. (2016) Accounting

for uncertainty in dormant life stages in stochastic demographic models. *Oikos,*

126, 900-909.

Paquet, M., Arlt, D., Knape, J., Low, M., Forslund, P. & Pärt, T. (2020) Why we should

care about movements: Using spatially explicit integrated population models to

assess habitat source-sink dynamics. *Journal of Animal Ecology,* **89,** 2922-2933.

Pausas, J.G. & Bond, W.J. (2020) On the three major recycling pathways in terrestrial

ecosystems. *Trends in Ecology & Evolution,* **35,** 767-775.

Pausas, J.G., Keeley, J.E. & Schwilk, D.W. (2017) Flammability as an ecological and

evolutionary driver. *Journal of Ecology,* **105,** 289-297.

Pausas, J.G. & Parr, C.L. (2018) Towards an understanding of the evolutionary role of fire

in animals. *Evolutionary Ecology,* **32,** 113-125.

Plummer, M. (2019) rjags: bayesian graphical models using MCMC.

Pradel, R. (1996) Utilization of capture-mark-recapture for the study of recruitment and

population growth rate. *Biometrics,* **52,** 703-709.

Prather, R.M., Dalton, R.M., Barr, B., Blumstein, D.T., Boggs, C.L., Brody, A.K., Inouye,

D.W., Irwin, R.E., Martin, J.G.A., Smith, R.J., Van Vuren, D.H., Wells, C.P.,

Whiteman, H.H., Inouye, B.D. & Underwood, N. (2023) Current and lagged

climate affects phenology across diverse taxonomic groups. *Proceedings of the*

Royal Society of London, Series B: Biological Sciences, **290,** 20222181.

Rodrigues, M.T. (1996) A new species of lizard, genus *Microblepharus* (Squamata:

Gymnophtalmidae), from Brazil. *Herpetologica,* **52,** 535-541.

Rodriguez-Caro, R.C., Gracia, E., Blomberg, S.P., Cayuela, H., Grace, M., Carmona, C.P.,

Perez-Mendoza, H.A., Gimenez, A. & Salguero-Gomez, R. (2023) Anthropogenic

impacts on threatened species erode functional diversity in chelonians and

crocodilians. *Nature Communications,* **14,** 1542.

Salguero-Gómez, R., Jones, O.R., Archer, C.R., Bein, C., de Buhr, H., Farack, C.,

Gottschalk, F.n., Hartmann, A., Henning, A., Hoppe, G., R??mer, G., Ruoff, T.,

Sommer, V., Wille, J., Voigt, J., Zeh, S., Vieregg, D., Buckley, Y.M., Che-

Castaldo, J., Hodgson, D., . . . Vaupel, J.W. (2016a) COMADRE: A global data

base of animal demography. *Journal of Animal Ecology,* **85,** 371-384.

Salguero-Gómez, R., Jones, O.R., Jongejans, E., Blomberg, S.P., Hodgson, D.J., Mbeau-

Ache, C., Zuidema, P.A., de Kroon, H. & Buckley, Y.M. (2016b) Fast–slow

continuum and reproductive strategies structure plant life-history variation

worldwide. *Proceedings of the National Academy of Sciences, USA,* **113,** 230-235.

Schwarzkopf, L. & Shine, R. (1992) Costs of reproduction in lizards: escape tactics and

susceptibility to predation. *Behavioral Ecology and Sociobiology,* **31,** 17-25.

Sinervo, B., Méndez-de-la-Cruz, F., Miles, D.B., Heulin, B., Bastiaans, E., Villagrán-Santa

Cruz, M., Lara-Resendiz, R., Martínez-Méndez, N., Calderón-Espinosa, M.L.,

Meza-Lázaro, R.N., Gadsden, H., Avila, L.J., Morando, M., De la Riva, I.J.,

Victoriano Sepulveda, P., Rocha, C.F.D., Ibargüengoytía, N., Aguilar Puntriano,

C., Massot, M., Lepetz, V., . . . Sites, J.W. (2010) Erosion of lizard diversity by

climate change and altered thermal niches. *Science,* **328,** 894-899.

Stearns, S.C. (1976) Life-history tactics: a review of the ideas. *The Quarterly review of*

biology, **51,** 3-47.

Stearns, S.C. (2000) Life history evolution: successes, limitations, and prospects.

Naturwissenschaften, **87,** 476-486.

Stott, I., Franco, M., Carslake, D., Townley, S. & Hodgson, D. (2010) Boom or bust? A

comparative analysis of transient population dynamics in plants. *Journal of*

Ecology, **98,** 302-311.

Stott, I., Hodgson, D.J. & Townley, S. (2012) Popdemo: An R package for population

demography using projection matrix analysis. *Methods in Ecology and Evolution,*

3, 797-802.

Sutherland, W.J., Adams, W.M., Aronson, R.B., Aveling, R., Blackburn, T.M., Broad, S.,

Ceballos, G., CÔtÉ, I.M., Cowling, R.M., Da Fonseca, G.A.B., Dinerstein, E.,

Ferraro, P.J., Fleishman, E., Gascon, C., Hunter, M., Hutton, J., Kareiva, P., Kuria,

A., MacDonald, D.W., MacKinnon, K., . . . Watkinson, A.R. (2009) One hundred

questions of importance to the conservation of global biological diversity.

Conservation Biology, **23,** 557-567.

Sutherland, W.J., Freckleton, R.P., Godfray, H.C.J., Beissinger, S.R., Benton, T.,

Cameron, D.D., Carmel, Y., Coomes, D.A., Coulson, T., Emmerson, M.C., Hails,

R.S., Hays, G.C., Hodgson, D.J., Hutchings, M.J., Johnson, D., Jones, J.P.G.,

Keeling, M.J., Kokko, H., Kunin, W.E., Lambin, X., . . . Gibson, D. (2013)

Identification of 100 fundamental ecological questions. *Journal of Ecology,* **101,**

58-67.

Tenan, S., Pradel, R., Tavecchia, G., Igual, J.M., Sanz-Aguilar, A., Genovart, M. & Oro,

D. (2014) Hierarchical modelling of population growth rate from individual

capture-recapture data. *Methods in Ecology and Evolution,* **5,** 606-614.

Van Sluys, M. (1993) Food habits of the lizard *Tropidurus itambere* (Tropiduridae) in

Southeastern Brazil. *Journal of Herpetology,* **27,** 347-351.

Van Sluys, M. (1997) Home range of the saxicolous lizard *Tropidurus itambere*

(Tropiduridae) in Southeastern Brazil. *Copeia,* **1997,** 623-628.

Van Sluys, M. & Henderson, R.W. (1993) The reproductive cycle of *Tropidurus itambere*

(Sauria: Tropiduridae) in Southeastern Brazil. *Journal of Herpetology,* **27,** 28-32.

Vieira, G.H.C., Mesquita, D.O., Péres Jr, A.K., Kitayama, K. & Colli, G.R. (2000)

Micrablepharus atticolus (NCN). Natural History. *Herpetological Review,* **31,** 241-

242.

Vinton, A.C., Gascoigne, S.J.L., Sepil, I. & Salguero-Gomez, R. (2022) Plasticity's role in

adaptive evolution depends on environmental change components. *Trends in*

Ecology & Evolution, **37,** 1067-1078.

Vitt, L.J. & Blackburn, D.G. (1991) Ecology and life-history of the viviparous lizard

Mabuya bistriata (Scincidae) in the Brazilian Amazon. *Copeia*, 916-927.

Vitt, L.J. & Caldwell, J.P. (1993) Ecological observations on Cerrado lizards in Rondônia,

Brazil. *Journal of Herpetology,* **27,** 46-52.

Vitt, L.J., Zani, P.A. & Lima, A.C.M. (1997) Heliotherms in tropical rain forest: The

ecology of *Kentropyx calcarata* (Teiidae) and *Mabuya nigropunctata* (Scincidae)

in the Curua-Una of Brazil. *Journal of Tropical Ecology,* **13,** 199-220.

Zimin, A., Zimin, S.V., Shine, R., Avila, L., Bauer, A., Böhm, M., Brown, R., Barki, G.,

de Oliveira Caetano, G.H., Castro Herrera, F., Chapple, D.G., Chirio, L., Colli,

G.R., Doan, T.M., Glaw, F., Grismer, L.L., Itescu, Y., Kraus, F., LeBreton, M.,

Martins, M., . . . Meiri, S. (2022) A global analysis of viviparity in squamates

highlights its prevalence in cold climates. *Global Ecology and Biogeography,* **31,**

2437-2452.

Zirondi, H.L., Ooi, M.K.J. & Fidelis, A. (2021) Fire-triggered flowering is the dominant

post-fire strategy in a tropical savanna. *Journal of Vegetation Science,* **32,** 1-11.

Table 1. Sensitivity (mean ± S.D.) of the demographic resilience components to parameters' perturbation in three species of lizards from the Brazilian Cerrado savannas: *Copeoglossum nigropunctatum*, *Micrablepharus atticolus*, and *Tropidurus itambere*. Cell colors vary from white to dark orange, indicating absolute sensitivity values of demographic resilience component concerning different plausible drivers: $\alpha_{vitalrate}$ = intercept of the vital rate function; $\alpha_{vitalrateSVL}$: slope of the linear relationship with body size (snout-vent length—SVL); $\beta_{vitalrateSVL}^2$: slope of the quadratic relationship with body size (SVL); μ_K = mean body growth constant, *i.e.*, time required to reach the asymptotic size, thus higher μ_K values means slower body growth rates. Ony parameters related to probability of reproduction (*prep*), number of newborns (*nb*), and body growth were higher than zero. The sensitivity values of the other vital rates were null and not shown. Negative values, which mean negative effects on the demographic resilience components, are biologically meaningful and mathematically possible.

Parameter	Resistance	Compensation	Recovery time
Copeoglossum nigropunctatum			
α_{prep}	-0.026 ± 0.002	0.111 ± 0.004	0.052 ± 0.001
$\beta_{prepSVL}$	-2.909 ± 0.315	13.836 ± 0.784	8.023 ± 0.264
$\beta_{prepSVL}^2$	-71.366 ± 69.583	86.673 ± 102.262	-173.243 ± 108.166
α_{nb}	-0.067 ± 0.002	0.2 ± 0.001	-0.21 ± 0.012
β_{nbSVL}	-9.58 ± 0.219	27.562 ± 1.194	-6.326 ± 0.745
μ_K	0.007 ± 0.001	0.004 ± 0.001	0.03 ± 0.005
Micrablepharus atticolus			
α_{prep}	-0.037 ± 0.011	0.035 ± 0.005	-0.103 ± 0.182
$\beta_{prepSVL}$	-1.475 ± 0.462	1.603 ± 0.219	-4.227 ± 8.933

μ_K	0.012 ± 0.003	0.005 ± 0.001	0.654 ± 0.251

Tropidurus itambere

α_{prep}	-0.036 ± 0.008	0.127 ± 0.02	0.059 ± 0.005
$\beta_{prepSVL}$	-2.740 ± 0.608	10.516 ± 1.557	5.573 ± 0.342
α_{nb}	-0.128 ± 0.005	0.296 ± 0.003	-0.033 ± 0.004
β_{nbSVL}	-8.806 ± 0.201	24.955 ± 0.319	5.312 ± 0.914
β_{nbSVL}^2	-186.827 ± 141.564	1655.239 ± 163.511	$3.747 \times 10^{10} \pm 9.308 \times 10^{10}$
μ_K	0.008 ± 0.006	0.005 ± 0.004	0.005 ± 0.004

Figure legends

Fig. 1. Model coefficients (mean and 95% credible intervals) of the effects of environmental covariates on vital rates of three species of lizards corresponding to *tmed2m* = mean air temperature at 200 cm height; *tmin0cm* = minimum air temperature at 0 cm height; *tmed0cm* = mean air temperature at 0 cm height; *RHmax* = maximum relative air humidity; *sol* = solar radiation; *precip* = accumulated precipitation; *perf* = mean locomotor performance; *ha* = hours of activity; *fire* = month with fire; *TSLF* = time since last fire; *random* = monthly random variation. We group-colored the covariates related to weather, microclimate, ecophysiological traits, fire, and randomness. Species' silhouettes from photos taken by Nicolás Pelegrin (*C. nigropunctatum* and *M. atticolus*) and Carlos Morais (*T. itambere*).

Fig. 2. Three-dimensional space of the demographic resilience components (resistance, compensation, and recovery time) for three species of lizards from the Brazilian Cerrado savannas, showing distinct demographic resilience strategies. With a 14-year mark-recapture dataset, we derived demographic resilience components from monthly stochastic Integral Projection Models shaped by weather, microclimate, ecophysiological traits, and fire covariates.

Fig. 3. Raw data, density, mean estimates, and 95% credible intervals of the demographic resilience components: (A, D & G) resistance; (B, E & H) compensation, and (C, F & I) recovery time of three species of lizards: (A-C) *Copeoglossum nigropunctatum*, (D-F) *Micrablepharus atticolus*, and (G-I) *Tropidurus itambere* from the Brazilian Cerrado savannas under different fire regime severity. Within each panel, we show high posterior density (95% probability) contrast letters corresponding to groups that are not significantly

different (overlapping zero difference). Later letters in the alphabet mean higher values (e >

d > c > b > a).

Fig. 4. Relationship between generation time (z-scaled) and the demographic resilience

components: (A, D & G) resistance, (B, E & H) compensation, and (C, F & I) recovery time

of three lizard species: (A-C) *Copeoglossum nigropunctatum*, (D-F) *Micrablepharus*

atticolus, and (G-I) *Tropidurus itambere* under fire regimes of varying severity. Lines and

shades represent mean estimates and 95% credible intervals derived from Bayesian dynamic

generalized additive models, respectively. To offer a higher resolution for *M. atticolus'*

resistance and compensation, we also provide inserts in D and E with a narrower y-axis

range.

Fig. 5. Relationship between reproductive output (z-scaled) and the demographic resilience

components (A, D & G) resistance, (B, E & H) compensation, and (C, F & I) recovery time

of three lizard species (A-C) *Copeoglossum nigropunctatum*, (D-F) *Micrablepharus*

atticolus, and (G-I) *Tropidurus itambere* under fire regimes of varying severity. Lines and

shades represent mean estimates and 95% credible intervals derived from Bayesian dynamic

generalized additive models, respectively. To offer a higher resolution for *M. atticolus'*

compensation, we also provide an insert in E with a narrower y-axis range.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fire severity
1 2 3 4 5