KSHV vIL-6 Enhances Inflammatory Responses by Epigenetic Reprogramming

Tomoki Inagaki[1], Kang-Hsin Wang[1], Ashish Kumar[1], Chie Izumiya[1], Hiroki Miura[1],

Somayeh Komaki[1], Ryan R. Davis[2], Clifford G. Tepper[3], Harutaka Katano[4],

Michiko Shimoda[1], & Yoshihiro Izumiya[1, 3, *]

[1] Department of Dermatology, School of Medicine, the University of California Davis (UC Davis),
Sacramento, California USA
[2] Department of Pathology and Laboratory Medicine, School of Medicine, UC Davis, Sacramento,
California USA
[3] Department of Biochemistry and Molecular Medicine, School of Medicine, UC Davis, Sacramento,
California USA
[4] Department of Pathology, National Institute of Infectious Diseases, Shinjuku, Tokyo, Japan

*Correspondence: Yoshihiro Izumiya DVM, PhD

Address: UCDMC Research III, Room 2200B, 4645 2nd Avenue, Sacramento CA 95817
Email: yizumiya@ucdavis.edu
Phone: 916-734-7253

Abstract

Kaposi sarcoma-associated herpesvirus (KSHV) inflammatory cytokine syndrome (KICS) is

a newly described chronic inflammatory disease condition caused by KSHV infection and

is characterized by high KSHV viral load and sustained elevations of serum KSHV-

encoded IL-6 (vIL-6) and human IL-6 (hIL-6). KICS has significant immortality and

possesses greater risks of having other complications, which include malignancies.

Although prolonged inflammatory vIL-6 exposure by persistent KSHV infection is

expected to have key roles in subsequent disease development, the biological effects of

prolonged vIL-6 exposure remain elusive. Using thiol-Linked Alkylation for the Metabolic

Sequencing and Cleavage Under Target & Release Using Nuclease analysis, we studied

the effect of prolonged vIL-6 exposure in chromatin landscape and resulting cytokine

production. The studies showed that prolonged vIL-6 exposure increased Bromodomain

containing 4 (BRD4) and histone H3 lysine 27 acetylation co-occupancies on chromatin,

and the recruitment sites were frequently co-localized with poised RNAPII with associated

enzymes. Increased BRD4 recruitment on promoters was associated with increased and

prolonged NF-κB p65 binding after the lipopolysaccharide stimulation. The p65 binding

resulted in quicker and sustained transcription bursts from the promoters; this mechanism

increased total amounts of hIL-6 and IL-10 in tissue culture. Pretreatment with the BRD4

inhibitor, OTX015, eliminated the enhanced inflammatory cytokine production. These

findings suggest that persistent vIL-6 exposure may establish a chromatin landscape

favorable for the reactivation of inflammatory responses in monocytes. This epigenetic

memory may explain the greater risk of chronic inflammatory disease development in

KSHV-infected individuals.

Author summary

Combined and continuous cytokine stimulation triggers transcription reprogramming and

is often used for specific tissue development. Continuous vIL-6 exposure occurs in KSHV-

infected patients and leads to inflammatory cytokine storm with high mortality. However,

possible epigenetic reprogramming by the vIL-6 and its association with pathogenesis

remain unclear. Here we demonstrate the establishment of a new chromatin landscape

mediated by BRD4 through prolonged vIL-6 exposure which contributes to more robust

and rapid transcription and increased cytokines production. Inhibition of BRD4

suppressed this inflammatory response. Our results indicate that targeting the epigenetic

effect of viral cytokines may lead to novel therapies for KSHV-induced inflammatory

cytokine storms.

Introduction

Controlled inflammatory response enhances host immunity against external insults, whereas uncontrolled inflammatory responses resulting from excessive cytokine production leads to disruption of such balance. In particular, monocytes and macrophages play major roles in controlling pro-inflammatory cytokines expression and their tightly controlled activation is critical for regulating systemic inflammation in our body. Previous reports have shown that continuous inflammatory stimulation enhances [1, 2] or sometimes impairs [3, 4] responses to subsequent stimulus. In either case, continuous inflammatory stimulation appears to alter the cellular phenotype and affect inflammatory cytokine production.

Kaposi's sarcoma-associated herpesvirus (KSHV) was first identified in Kaposi's sarcoma (KS) lesions in 1994 [5]. KSHV is also associated with lymphoproliferative diseases, primary effusion lymphoma (PEL) and multicentric Castleman's disease (MCD) [6, 7]. Recent studies showed that KSHV infection also causes severe systemic inflammation, which is categorized as Kaposi's sarcoma-associated herpesvirus inflammatory cytokine syndrome (KICS) [8]. KICS is characterized by increased viral loads and inflammatory cytokines such as human IL-6 (hIL-6), IL-10, and KSHV-encoded

interleukin-6 homolog, viral IL-6 (vIL-6). Individuals with KICS are known to have a higher

risk of developing KSHV-associated cancers and other malignancies [8].

The vIL-6, encoded by KSHV ORF-K2, is expressed during the lytic replication

phase, and single cell transcriptomics studies also suggested that vIL-6 may be

expressed during latency in selected cell populations [9]. Similar to hIL-6, vIL-6 is known

to enhance cell proliferation, endothelial cell migration, and angiogenesis by upregulating

vascular endothelial growth factor (VEGF) [10] and downregulates the caveolin 1

expression [11-13]. Moreover, vIL-6 transduction also leads to cell transformation in the

3T3 cells model and induces tumors in xenograft mice [10]. Increased metastasis is also

seen in a murine xenograft model with transgenic vIL-6 mice [14]. The vIL-6 activates

downstream signaling pathways by binding to the cellular hIL-6 receptor, gp130.

Dimerization of gp130 induced by hIL-6 binding activates Janus tyrosine kinases and

phosphorylates the SH2-containing cytoplasmic protein STAT3 (signal transducer and

activator of transcription 3). The phosphorylated STAT3 forms a dimer and translocates

to the nucleus to activate the downstream genes that play essential roles in inducing the

inflammatory response, cell survival, and immune responses [15-17].

Synergistic interactions between another inflammatory-associated transcription

factor, nuclear factor-kappa B (NF-κB) and STAT3 are known to induce the

hyperactivation of NF-κB followed by the production of various inflammatory cytokines

including TNFα and hIL-6. Because hIL-6 expression is regulated by NF-κB pathway

activation, simultaneous activation of NF-κB and STAT3 triggers a positive feedback loop

of NF-κB activation in the hIL-6/STAT3 axis. This positive feedback loop is called the IL-

6 amplifier (IL-6 Amp) and is a key element in inflammatory disease development [18]. In

the IL-6 Amp disease model, obesity, injury, and infection trigger chronic inflammation

and/or a systemic cytokine storm, which is enhanced by interactions between local non-

immune cells and infiltrating immune cells [18]. A prominent example of IL-6 Amp is also

seen in Coronavirus disease 2019 (COVID-19), which is triggered by severe acute

respiratory syndrome coronavirus 2 (SARS-CoV2) infection [18]. Consistent with the

significance of hIL-6 in disease development, a humanized monoclonal antibody against

the hIL-6 receptor, tocilizumab, has proven successful for treating Castleman's disease

as well as COVID-19 [19, 20]. In the case of KSHV-associated diseases, the IL-6 Amp

may be triggered by KSHV infection and is likely to be associated with vIL-6 expression

from KSHV-infected cells. However, how persistent exposure of vIL-6 alters cellular

responses and enhances inflammation cytokine production remain elusive.

The bromodomain and extraterminal domain (BET) family proteins such as

bromodomain containing 4 (BRD4) are transcription regulators, which localize to cellular

enhancer and promoter loci and control the transcription of a wide range of

proinflammatory genes. BRD4 regulates transcription by recognizing acetylated histone

tail and interacts with transcription factors such as NF-κB p65 and transcription elongation

complex [21]. This mechanism facilitates the phosphorylation of RNAPII and therefore

promotes transcription initiation and elongation. In the context of proinflammatory gene

activation, interaction with NF-κB p65 is especially important, because p65 itself is

acetylated at K218, K221, and K310, and the acetylated K310 is recognized by the BRD4

bromodomain to form an active transcription complex [22]. In addition to the specific

protein-protein interactions, BRD4 also forms nuclear puncta on super-enhancer genomic

regions that exhibit properties of liquid-like condensates [23]. The intrinsically disordered

regions (IDRs) of BRD4 are responsible for phase-separated condensates formation with

other coactivators on the chromatin loci for transcription elongation [23], which suggests

that enrichment of BRD4 at specific enhancers or promoters are key for selection of genes

for activation.

Here we demonstrate that a biological effect of prolonged vIL-6 exposures alters

the chromatin landscape changes which provides a putative mechanism of

overexpression of inflammatory cytokines in vIL-6 pre-exposed cells. We show that an

increased amount of BRD4 on chromatin by the prolonged vIL-6 exposure is responsible

for transcription deregulation, and repetitive activation of STAT3 appeared to be

responsible for the change. BRD4 accumulation prolongs transcription activity at targeted

sites with a secondary stimulus by prolonging transcription factor binding and duration of

transcriptions. This epigenetic memory may explain the greater risk of chronic

inflammatory disease development in KSHV-infected individuals.

Results

Biological effects of prolonged vIL-6 exposure on gene transcription and

phenotypes

In contrast to temporally regulated hIL-6 expression during wound healing, persistent

KSHV infection leads to continuous vIL-6 production from infected cells. Especially, vIL-

6 is detected in the serum of KICS patients and the hematopoietic cells in Castleman's

disease patients with a high KSHV viral load [24]. To study the biological consequences

of vIL-6 exposure, we first purified recombinant vIL-6 protein from recombinant

baculovirus-infected cells and confirmed its biological activity by STAT1 and STAT3

activation [25]. In the study, we showed that longevity and strong phenotypic changes in

primary monocytes were induced by vIL-6 exposure for at least 9 days [25]. Here, we

prepared for a cell culture model with THP-1 cells which were exposed to vIL-6

continuously for two weeks (we refer to vIL-6/THP-1 hereafter) **(Fig. 1A)**. THP-1 cells, a

representative monocyte cell line, was employed because monocytes play a major role

in inflammatory responses [26]. Using this cell culture model, we compared the overall

transcriptional profiles between vIL-6/THP-1 and parental THP-1 cells followed by

secondary stimulation with selected cytokines. We used vIL-6 (100ng/ml) and hIL-6

(100ng/ml), which can activate common downstream signaling pathways, and TGF-

β (100ng/ml), which has a non-overlapping pathway, as a secondary stimulation.

To evaluate the newly synthesized RNAs by cytokines stimulation, thiol (SH)-

Linked Alkylation for the Metabolic sequencing of RNA (SLAM-seq) was employed [27].

vIL-6/THP-1 and parental THP-1 cells were unstimulated or stimulated by vIL-6, hIL-6

or TGF-β for 30 minutes, followed by incubation with 4-thiouracil (4sU) for 60 minutes to

label transcribing RNAs during the incubation periods. In this experimental setting,

sequence reads with mutations T>C identify the only nascent synthesized mRNA

because 4sU incorporation with chemical alkylation forces incorporation of G instead of

A at 4sU positions during the reverse transcription step [27]. Isolated sequence reads

with T>C mutation(s) were then compared among control (non-stimulation), vIL-6, hIL-6,

and TGF-β **(Fig.1A)**. As shown in **Fig. 1B**, principal component analysis (PCA)

demonstrated clearly distinct cellular responses in vIL-6/THP-1 and also among stimuli.

In response to vIL-6 stimulation, we observed that the number of upregulated genes (log2

fold change >1, adjusted *p*-value < 0.01) in vIL-6/THP-1 was about twice as many as

those with parental THP-1 cells (**Fig. 1C, left**). Similar responses were also seen with

hIL-6 as a secondary stimulation (**SFig. 1A**). The vIL-6 and hIL-6 secondary stimulation

showed very similar transcriptional profiles (**SFig. 1B**) and downstream pathways (**SFig.**

1C), suggesting that vIL-6 is clearly a functional homolog of hIL-6. On the other hand, the

number of upregulated genes with the TGF-β was slightly decreased in vIL-6/THP-1

compared to parental THP-1 cells (**Fig. 1C, right**). In addition, the degree of up-regulated

gene expression by vIL-6 or hIL-6 was also increased to a greater extent in vIL-6/THP1

cells compared to parental THP-1 cells (**Fig. 1D, SFig. 1D**), while there was no significant

differences with TGF-β (**Fig. 1D**). The results suggested that the enhanced transcription

response is a pathway dependent. Gene Ontology (GO) analysis indicated that genes

related to immune response, Th2 cell differentiation and inflammatory response were

commonly enriched by secondary vIL-6 stimulation in parental and vIL-6/THP-1 cells (**Fig.**

1E: light green), while cell cycle and cell division pathway were additionally activated in

vIL-6/THP-1 cells (**Fig. 1E: green**). Consistent with the GO analysis, cell proliferation was

increased in vIL-6/THP-1 cells measured by MTT assay (**SFig. 1E**). Despite an increased

number of responding genes with the prolonged vIL-6 exposure, the degree of STAT3

activation and gp130 expression (cell surface receptor for vIL-6) was not changed (**SFig.**

1F, G).

Since KICS patients are characterized not only by increased high vIL-6 or hIL-6

but also by increased viral loads, we also examined the effect in KSHV genes expression

during de novo infection in vIL-6/THP-1 cells [8]. For this, we infected THP-1 and vIL-

6/THP-1 cells with r219.KSHV and performed total RNA-sequencing (RNA-seq). As

shown in **Fig. 1F**, overall KSHV gene expression was higher in vIL-6/THP-1 cells

compared to parental THP-1 cells. Moreover, vIL-6 and PAN-RNA expression were

significantly higher in vIL-6/THP-1 compared to parental THP-1 cells (**Fig. 1G**).

Surprisingly, vIL-6 was transcribed as much as PAN RNA in THP-1 cells (**Fig. 1G**). These

results suggested that prolonged vIL-6 exposure enhances transcriptional response to

secondary vIL-6 and hIL-6 stimulation, and further assists KSHV genes transcription in

vIL-6 pre-treated THP-1 cells.

Prolonged vIL-6 stimulation increased BRD4 occupancies on chromatin

The studies above demonstrated that prolonged vIL-6 exposure increased the number of

responding genes followed by vIL-6 or similar (hIL-6) stimuli without altering the amounts

of signal-dependent transcription factor (STAT3). These results led us to examine

chromatin modification changes induced by prolonged vIL-6 exposure. For this, we first

used CHIP-Atlas, a data mining tool for predicting proteins bound to promoters of

upregulated genes based on a comprehensive epigenetic database to narrow down

possible epigenetic factors. As shown in **Fig. 2A**, STAT1 and STAT3 were commonly

extracted as putative responsible transcriptional factors with vIL-6 stimuli both in parental

THP-1 and vIL-6/THP-1 cells, demonstrating the validity of this enrichment analysis.

Interestingly, transcription coactivators and several transcription factors such as Cyclin

dependent kinase 8 (CDK8), Mediator complex subunit 12 (MED12) and Bromodomain-

containing protein 4 (BRD4), are likely to be enriched only in the genomic loci in vIL-

6/THP-1 cells (**Fig 2A**). BRD4 is a member of the BET (bromodomain and extra terminal

domain) family and an important coactivator that binds to H3K27Ac to mediate

transcription regulation together with CDK8, MED12 and RNA polymerase II (RNAPII)

[28]. Because BRD4 was identified as the most enriched transcription regulator with

prolonged vIL-6 exposure (**Fig. 2B**), we next performed Cleavage Under Targets &

Release Using Nuclease (CUT&RUN) for BRD4 and RNAPII. In addition, two active

histone modification marks (H3K27Ac, H3K4me3) and CTCF were examined to reveal

changes in chromatin landscape [29]. Parental THP-1 cells and vIL-6/THP-1 cells were

compared in duplicate samples for statistical analyses. Consistent with the CHIP-Atlas

results, BRD4 was more enriched in the transcription start sites of up-regulated genes in

vIL-6/THP-1 cells (**SFig. 2A**). We also found that BRD4 and H3K27Ac peaks were

increased across the genome in vIL-6/THP-1 cells while RNAPII and H3K4me3 were

unchanged (**Fig. 2C**). Immunoblotting showed that the total amount of BRD4 modification

was not affected with prolonged vIL-6 exposure, as well as H3K27Ac or H3K4me3,

demonstrating that increased BRD4 in vIL-6/THP-1 was from the non-chromatin bound

fraction **(SFig. 2B)**. HOMER motif analysis for newly accumulated BRD4 genomic regions

identified several key monocyte transcription factors such as E74, ETS Transcription

Factor 4, and PU.1（**SFig. 2C**). PU.1 was also identified as a transcriptional factor likely

to be involved in gene activation seen in prolonged vIL-6 exposure by SLAM-seq **(Fig.**

2A).

We next examined the position and association of the newly accumulated BRD4

with H3K27Ac to reveal the changes of active genomic hubs. The CUT&RUN studies

identified 31,929 new H3K27Ac regions, while 21,124 regions found in parental cells were

no longer modified by H3K27Ac in vIL-6/THP-1 cells **(Fig. 2D).** As shown in **Fig. 2E**,

BRD4 was enriched more significantly and sharply at the centers of H3K27Ac peaks in

vIL-6/THP-1 cells. The genomic regions, where BRD4 and H3K27Ac peaks overlap, were

also increased from 18,886 to 27,509 by prolonged vIL-6 exposure（**SFig. 2D**), suggesting

that newly accumulated BRD4 was recruited in newly established H3K27Ac loci in vIL-

6/THP-1 cells（**Fig. 2F**). RNAPII and H3K4me3 were also enriched in the genomic regions,

where BRD4 and H3K27Ac peaks overlapped（**SFig. 2E**). Closer examination of these

enriched sites demonstrated that while the overall number of BRD4/H3K27Ac overlapped

peaks around TSS was increased (**Fig. 2G**), the relative abundance of the peaks at the

promoter region decreased from 15.4% to 13.5 % in vIL-6/THP-1 cells (**Fig. 2H**). These

results may suggest that a larger fraction of newly generated active hubs is located at

intragenic regions, perhaps at newly activated enhancers. The CUT&RUN results with

the CTCF antibody also suggested that prolonged vIL-6 stimulation increased overall co-

occupancies between CTCF and BRD4 across the genome (**Fig. 2I**), suggesting the

establishment of activated chromatin hubs in vIL-6/THP-1 cells. The new BRD4 and

H3K27Ac peaks were co-localized to promoter regions of genes associated with cellular

response to DNA damage stimulus and cell division, such as cyclin E1, which is consistent

with the results from the SLAM-seq analysis **(SFig. 2F, G)**.

Enhanced response to LPS via prolonged vIL-6 stimulation

Synergistic interactions between NF-κB and STAT3 induce the hyperactivation of NF-κB

followed by the production of various inflammatory cytokines such as hIL-6 [18]. KSHV-

infected cells are known to have elevated NF-κB activation and STAT pathways are

constitutively activated in primary effusion lymphomas [30, 31]. Accordingly, we examined

if repetitive STAT activation by vIL-6 exposure alters NF-κB mediated cell responses in

cis. To study this, we stimulated vIL-6/THP-1 cells with LPS, a strong inducer of NF-κB

pathway activation, and examined the cellular response. We first confirmed that 100 ng/ml

of LPS stimuli for 6 hours is sufficient for detecting hIL-6 production by ELISA assay **(SFig.**

3A), and selected this as a time point to collect samples in order to minimize the

secondary effects from the newly produced cytokines with LPS stimuli. We measured a

series of downstream inflammatory cytokines produced with Olink, a multiplex proximity

extension assay (PEA) [32]. The results showed that LPS upregulated the expression of

21/41 inflammatory cytokines (log2 fold change >1) in parental THP-1 cells (**Fig. 3A**).

Among them, the expression of 15 cytokines, such as hIL-6, IL-10, IL-1β, and CCL8, was

enhanced in vIL-6/THP-1 cells compared to parental THP-1 cells (**Fig. 3A, B**). IFNα

stimulation also induced more cytokines in vIL-6/THP-1 cells similarly to that of LPS

stimulation (**SFig. 3B, C**), suggesting that prolonged vIL-6 exposure to monocytes

sensitized cells to secondary stimuli, reinforcing the notion that enhanced activation is

due to the formation of active chromatin complexes at local chromatin.

The transcription profile in response to LPS was also accompanied by SLAMseq

analysis. LPS (100ng/ml) and 4sU were added to parental THP-1 cells or vIL-6/THP-1

cells at the same time and incubated for 6 hours to label nascent transcribed RNAs. PCA

demonstrated that LPS strongly changed the gene transcription profile, and the variance

between THP-1 and vIL-6/THP-1 cells was further increased by LPS stimulation (**Fig. 3C**).

As expected, the inflammatory response was identified as the most enriched pathway in

commonly up-regulated genes (N = 1626) (**Fig. 3D**). Next, we investigated whether the

increased BRD4 and H3K27Ac occupancies in vIL-6/THP-1 cells contributed to the

increased inflammatory gene transcription by LPS stimulation. Commonly up-regulated

genes were extracted and the peak scores of the BRD4 and H3K27Ac overlapping

regions in THP-1 and vIL-6/THP-1 cells were compared (**Fig. 3E**). As shown in **Fig. 3F**,

co-occupancy of BRD4 and H3K27Ac was indeed associated with increased transcription

activation with LPS. The increased responses to LPS stimuli were also associated with

an increased BRD4 and H3K27Ac co-occupancy compared to genes whose expression

was not changed (**Fig. 3G**). These results suggested that prolonged vIL-6 exposure

enhances the inflammatory response via increased acetylation of H3K27 at BRD4

recruited sites.

BRD4 is responsible for prolonging the hIL-6 transcription burst with LPS

Because BRD4 was enriched in the promoter regions of hIL-6, hIL-10 and IL-1β with the

prolonged vIL-6 exposure **(SFig. 4A)** and are associated with increased production of

cytokines in culture media **(Fig. 3A)**, we next studied how BRD4 recruitment contributes

to increasing transcripts. We speculated that increased occupancies of BRD4, which

contains an intrinsically disordered domain to form liquid-liquid phase separation (LLPS)

[33], may increase the recruitment and duration of p65 binding at promoter regions. This

mechanism may extend the duration or robustness of transcription after stimulation.

Accordingly, we examined the recruitment of p65 after LPS stimuli in vIL6/THP-1 cells

and compare it with parental THP-1 cells. The proximity ligation assay showed that the

association between BRD4 and p65 was increased after LPS stimulation in vIL6/THP-1

cells (**Fig. 4A**), while the total p65 amount in the nucleus after LPS stimulation was not

significantly changed in vIL-6/THP-1 cells (**Fig. 4B**). Chip-qPCR also showed increased

p65 recruitment on the hIL-6, IL10, and IL-1β promoters by LPS stimuli in vIL6/THP-cells

(**Fig. 4C**). Transcription frequencies were also determined by qRT-PCR analysis of

nascent RNA **(SFig. 4B)**. As shown in **Fig. 4D**, hIL-6, IL-10 and IL-1β transcripts were

produced more rapidly (hIL-6 and IL-10) and/or continuously (IL-1β) in vIL-6/THP-1 cells

after stimulation. Importantly, BRD4 occupancies are responsible for the increased

transcription, because OTX-015, a BRD4 inhibitor, drastically suppressed the

transcription in both parental THP-1 and vIL-6/THP-1 cells, and the effect was more

pronounced in vIL-6/THP-1 cells (**Fig. 4E**). Taken together, prolonged vIL-6 exposure

changed the chromatin landscape to be favorable to the response of inflammatory

secondary stimuli, and establishment of active chromatin hubs with BRD4 recruitment is

critical for this response (**Fig. 5**).

Discussion

In this study, we applied SLAM-seq for most of the RNA-seq experiments to increase the

resolution to identify direct target genes (**Fig. 1, 3C**). A short window after the respective

stimuli was utilized to examine robustness and differences made in direct target gene

expression with prolonged vIL-6 stimuli. By measuring nascent RNA that was labelled

shortly after stimulation, we also avoid indirect, or secondary activation. The study clearly

demonstrated that direct target genes are significantly overlapped with that of hIL-6

(**Fig.1B**). This also suggests that these two signaling ligands are interchangeable, making

it easier to have a continuously stimulated IL-6 signaling pathway in KSHV infected

individuals.

Cytokines that activate STAT3 (e.g. hIL-6) and NF-κB (e.g. TNF-α)

synergistically increase the production of various inflammatory cytokines, called IL6-Amp.

Since vIL-6 has the same biological activity as hIL-6 **(Fig.1B, SFig. 1B, C),** it is reasonable

to speculate that vIL-6 bypasses hIL-6 expression to trigger the IL6-Amp in vivo. The vIL-

6 can bind to gp130 alone and does not require the high affinity IL-6 receptor [34, 35].

This is in clear contrast to hIL-6, which must bind to its high affinity receptor before a

signaling complex can be formed with its signal transducer, gp130 [36, 37]. Importantly,

the gp130 is widely expressed in human tissues, and the notion that vIL-6 activates

downstream JAK/STAT pathway without requiring binding to the classical gp80 receptor

[38-41] suggests that vIL-6 would initiate signaling in a wider variety of cell types [40, 42].

In addition, previous studies by Dr. Nicholas's group further demonstrated that vIL-6 can

function though intracellularly via activation of gp130 at the endoplasmic reticulum in

infected cells [43, 44]. The study suggested that vIL-6 may support the growth and

survival of KSHV-infected cells in an autocrine manner and such unique signaling

activities contribute to the maintenance of latently infected cells without alarming

bystander cells. Our recent study with primary monocyte infection as well as this study

agreed very well with the observation, in which prolonged vIL-6 stimuli shifted the cell

transcription program towards cell growth and anti-apoptotic signaling (**Fig. 1E, SFig. 1E**).

More importantly, similar phenotypes were also seen in wild-type KSHV primary infection

but not infection with vIL-6 stop virus [25].

Prolonged vIL-6 stimuli similarly increased the number of responding genes to

secondary vIL-6 and hIL-6 stimulation whereas TGF-β did not show enhanced responses

(**Fig. 1C, D**). Generally, chronic inflammation leads to a stronger inflammatory response,

which is described as immunological training [45]. In immunological training, the primary

stimulus will prime specific regulatory elements in enhancer regions. This established

chromatin landscape allows faster and increased transcriptional activation of the

secondary stimulus, leading to an enhanced inflammatory response. At the chromatin

level, immunological training has been shown to induce long-lasting epigenetic

remodeling of regulatory elements [46]. This remodeling includes changes in activating

histone marks such as H3K27Ac and H3K4me3 [46, 47]. In this study, signal-dependent

transcription factors such as STAT1 and STAT3 motifs were enriched at up-regulated

gene promoters, and BRD4 and H3K27Ac were increasingly co-occupied at those

promoters **(Fig. 2C)**. After removal of the stimulus, STAT3 is likely to be left from the

promoters; however, our study suggested that BRD4 and the activated H3K27Ac

modification remained and established a "primed" chromatin state for subsequent

activation **(Fig. 5)**. As a result, LPS activated selected genes that had been primed by

STAT3, more quickly and strongly **(Fig. 4E)**. Indeed, cellular enzymes that are known to

be enriched at enhancers such as PU.1, MED1, SWI/SNF complexes are predicted to be

co-occupied at newly responding gene promoters **(Fig. 2A, SFig. 2C)**. These

observations support that vIL-6 stimulation induced transcription reprogramming, which

favors a subsequent inflammation response.

What would be the evolutional advantage for maintaining vIL-6 in KSHV genome?

Interestingly, vIL-6 is reported to be constitutively expressed in some cell populations

even during latent infection [13, 48]. We think that immunological training effects are

utilized for the establishment of active latent chromatin for inducible lytic gene expression.

Consistent with this hypothesis, BRD4, H3K27Ac, and H3K4me3 are all highly

accumulated at the KSHV ori-Lyt region in latent chromatin [49]. We think that KSHV is

likely to utilize host defense mechanisms to recruit BRD4 on its genome to establish a

"primed" latent chromatin for further reactivation, which is partly seen in increased KSHV

transcription in vIL-6/THP-1 cells (**Fig. 1F, G**). Perhaps, related to this, we recently

showed that the vIL-6KO virus could not reprogram primary infected monocytes and could

not replicate continuously [25]. These cellular and virological vIL-6 functions are likely to

contribute to the highly inflammatory phenotypes seen in KSHV-infected patients.

In summary, we reported that long-term vIL-6 exposure increased BRD4 and

H3K27Ac co-occupancies at inflammatory-related genes, resulting in enhanced

inflammatory responses to LPS. BRD4 is clearly the center of the altered cell responses

caused by prolonged vIL-6 exposure, in which the pre-located BRD4 on the promoter

enabled transcription factors and cofactors that are brought in proximity by p65 to interact

more rapidly and steadily upon LPS stimulation. Changes in the enhancer-promoter

genomic interactions by long-term exposure to vIL-6 remain a promising avenue for

further research. Moreover, animal models with continuous vIL-6 exposure should reveal

how vIL-6 trained memory polarizes immune cell communications and react to secondary

stimuli, which will provide valuable information for a better understanding the

pathogenesis of KICS. Although continued treatment with BET inhibitors is known to

cause large genomic structural changes and rapidly induce drug-resistant cells [50, 51],

short-term treatment with well-scheduled drug holidays may be beneficial to reverse the

immunological training memory established by KSHV vIL-6.

MATERIALS AND METHODS

Chemicals, reagents and antibodies

Dulbecco's modified minimal essential medium (DMEM), fetal bovine serum (FBS),

phosphate-buffered saline (PBS), Trypsin-EDTA solution, 100 X penicillin–streptomycin–

L-glutamine solution were purchased from Thermo Fisher (Waltham, MA). Puromycin and

G418 solution were obtained from InvivoGen (San Diego, CA). Hygromycin B solution

was purchased from Enzo Life Science (Farmingdale, NY).

The following antibodies were used for CUT & RUN, immunoblotting and flow cytometry:

rabbit anti-BRD4 (Cell Signaling, E2A7X), rabbit anti-RNAPII (Millipore, clone CTD4H8),

rabbit anti-H3K27ac (CST, clone D5E4), rabbit anti-H3K4me3 (Cell Signaling, clone

C42D8), mouse anti-β-actin (Santa Cruz 47778), rabbit anti-gp130 (CST), rabbit IgG

(CST, clone DA1E). Rabbit anti-vIL-6 was a gift from Dr. Robert Yarchoan (NIH/NCI).

Cells

THP-1 cell lines was obtained from ATCC. Cells were maintained in an RPMI 1640

medium supplemented with 1% fetal bovine serum (FBS; Invitrogen), 2 mM L-glutamine

and 100 U/ml penicillin and 100 µg/ml streptomycin at 37°C in a humidified 5% CO_2. THP-

1 cells were passaged two to three times per week when reaching 1 x 10^6 cells/ml.

Purification of recombinant protein

Spodoptera frugiperda Sf9 cells (Millipore) were maintained in Ex-Cell 420 medium

(Sigma), and recombinant baculoviruses were generated with the BAC-to-Bac system as

previously described [52] [53].

SLAM-seq

SLAM-seq [27] was performed using the SLAMseq Kinetics Kit (Lexogen GmbH, Vienna,

Austria) according to the manufacturer's standard protocol.

Briefly, biological replicate cultures of THP-1 cells were incubated with vIL-6 (100μg/ml)

for 30 min or 2 weeks. Subsequently, 4-Thiouridine (s4U; 300 μM) was added to the

culture media and the cells were incubated for 1 h in order to label newly synthesized

RNA. As for the LPS (100ng/ml) stimulation, s4U was added to the culture media

simultaneously with LPS, and cells were incubated for 6 hours. Total RNA was isolated

and then the 4-thiol groups in the s4Uracil-labeled transcripts were alkylated with

iodoacetamide (IAA). QuantSeq 3′ mRNA-Seq (FWD) (Lexogen, Inc.) Illumina-compatible,

indexed sequencing libraries were prepared from alkylated RNA samples (100 ng)

according to the manufacturer's protocol for oligo(dT)-primed first-strand cDNA synthesis,

random-primed second-strand synthesis, and library amplification. Libraries were

multiplex sequenced (1 × 100 bp, single read) on an Illumina HiSeq 4000 sequencing

system. SLAM-Seq datasets were analyzed using the T > C conversion-aware SLAM-

DUNK (Digital Unmasking of Nucleotide conversion-containing k-mers) pipeline utilizing

the default parameters [27, 54]. Briefly, nucleotide conversion-aware read mapping of

adapter- and poly(A)-trimmed sequences to the human GRCh38/hg38 reference genome

assembly was performed with NextGenMap [55]. Alignments were filtered for those with

a minimum identity of 95% and minimum of 50% of the read bases mapped. For multi-

mappers, ambiguous reads and non-3′ UTR alignments were discarded, while one read

was randomly selected from multimappers aligned to the same 3′ UTR. SNP calling

(coverage cut-off of 10X and variant fraction cut-off of 0.8) with VarScan2 [56] was

performed in order to mask actual T > C SNPs. Non-SNP T > C conversion events were

then counted and the fraction of labeled transcripts was determined. All results were used

for downstream analyses, such as nascent transcript analysis and differential expression

analysis (DESeq2) [57].

Bioinformatics analysis of SLAM-seq data

The UCSC Genome Browser was used to convert RefSeq IDs to gene symbols (refGene).

The resulting data were first filtered by log2 fold change and sorted by adjusted p-value

from lowest to highest. The resulting up-regulated genes (log2 fold change >1 and

adjusted *p*-value <0.01) were extracted. For functional enrichment analysis (KEGG

pathway analysis), the official gene symbols were applied in DAVID web service tools

[58]. For transcription factor analysis, gene symbols were submitted to ChIP-Atlas [59] to

analyze common regulators and to predict transcription factor binding. The following

parameters were used; Organism (annotation): Homo sapiens (hg38), Experiment type;

TFs and others, Cell type Class; Blood, Threshold; 100. The output factors were then

listed and plotted in a Venn diagram.

RNA-sequencing

vIL-6/THP-1 and parental THP-1 cells were infected with or without r219.KSHV for 72h

and RNA was purified using the Quick-RNA miniprep kit (Zymo Research, Irvine, CA,

USA). Indexed, stranded mRNA-seq libraries were prepared from total RNA (100 ng)

using the KAPA Stranded mRNA-Seq kit (Roche) according to the manufacturer's

standard protocol. Libraries were pooled and multiplex sequenced on an Illumina

NovaSeq 6000 system (150-bp, paired-end, >30 × 106 reads per sample). RNA-Seq data

was analyzed using a Salmon-tximport-DESeq2 pipeline. Raw sequence reads (FASTQ

format) were mapped to the reference human genome assembly (GRCh38/hg38,

GENCODE release 36) and quantified with Salmon [60]. Gene-level counts were

imported with tximport [61] and differential expression analysis was performed by

DESeq2 [57], which are visualized by Venn diagram.

Cleavage under targets and release using nuclease (CUT&RUN)

CUT&RUN [29] was performed essentially by following the online protocol developed by

Dr. Henikoff's lab with a few modifications to fit our needs. Cells were washed with PBS

and wash buffer [20 mM HEPES-KOH pH 7.5, 150 mM NaCl, 0.5 mM Spermidine (Sigma,

S2626), and proteinase inhibitor (Roche)]. After removing the wash buffer, cells were

captured on magnetic concanavalin A (ConA) beads (Polysciences, PA, USA) in the

presence of CaCl2. Beads/cells complexes were washed three times with digitonin wash

buffer (0.02% digitonin, 20 mM HEPES-KOH pH 7.5, 150 mM NaCl, 0.5 mM Spermidine

and 1x proteinase inhibitor), aliquoted, and incubated with specific antibodies (1:50) in

250 μL volume at 4℃ overnight. After incubation, unbound antibody was removed by

washing with digitonin wash buffer three times. Beads were then incubated with

recombinant Protein A/G–Micrococcal Nuclease (pAG-MNase), which was purified from

E.coli in 250 μl digitonin wash buffer at 1.0 μg/mL final concentration for 1 h at 4 °C with

rotation. Unbound pAG-MNase was removed by washing with digitonin wash buffer three

times. Pre-chilled digitonin wash buffer containing 2 mM CaCl$_2$ (200 μL) was added to the

beads and incubated on ice for 30 min. The pAG-MNase digestion was halted by the

addition of 200 μl 2× STOP solution (340 mM NaCl, 20 mM EDTA, 4 mM EGTA, 50 μg/ml

RNase A, 50 μg/ml glycogen). The beads were incubated with shaking at 37 °C for 10 min

in a tube shaker at 300 rpm to release digested DNA fragments from the insoluble nuclear

chromatin. The supernatant was then collected by removing the magnetic beads. DNA in

the supernatant was purified using the NucleoSpin Gel & PCR kit (Takara Bio, Kusatsu,

Shiga, Japan). Sequencing libraries were then prepared from 3 ng DNA with the Kapa

HyperPrep Kit (Roche) according to the manufacturer's standard protocol. Libraries were

multiplex sequenced (2 × 150 bp, paired-end) on an Illumina NovaSeq 6000 system to

yield ~15 million mapped reads per sample. With separate replicated experiments, qPCR

was used to examine enrichment at selected genomic regions. Primer sequences are

provided in **Table.S1.**

CUT&RUN sequence reads were processed with fastp [62] and aligned to the human

GRCh38/hg38 reference genome assembly with Bowtie2 and yielding mapped reads in

BAM files [63]. Hypergeometric Optimization of Motif EnRichment (HOMER) v4.11 was

used for peak detection and their annotation utilizing the defalt parameters described in

the developer's manual [31]. Deeptools was used for making plotprofiles utilizing the

defalt parameters [64]. Peaks and read alignments were visualized using the Integrated

Genome Browser [65].

Olink analysis

1.5×10^6 THP-1 cells or vIL-6/THP-1 cells were prepared in triplicate and washed twice

by PBS10ml and then suspended with 1ml fresh RPMI medium in 12 well plates.

Inflammatory cytokines such as LPS (100ng/ml), IFNα (100ng/ml), vIL-6 (100ng/ml) and

hIL-6 (100ng/ml) were then added to each well and cultured at 37℃, 5% CO_2. 6 hours

after incubation, supernatants were harvested and centrifuged (3,000 rpm, 3min) to

remove the remaining cells. The resulting supernatant samples were stored at -20℃ for

approximately two weeks until they were sent for analysis. Samples wrapped in dry ice

were then sent to Olink Proteomics (Temple City, CA). Olink's Proximity Extension Assay

(PEA) technology uses antibody pairs conjugated to unique oligonucleotides and is

quantified via PCR. When both antibodies of a pair bind the target protein simultaneously,

their respective conjugated oligonucleotides are brought into proximity, facilitating

hybridization. The oligonucleotide sequence is then extended by DNA polymerase,

amplified, and measured by qPCR to determine the sample's initial protein abundance.

Raw analyte expression values after PCR underwent multiple rounds of transformation

by Olink, including a log2 transformation, and were returned as normalized protein

expression (NPX) values [66]. For evaluating inflammatory cytokine production in the

culture medium, the Olink® Target 48 Cytokine panel was selected by multiple

investigators. The data was processed with Olink Insight Stat Analysis software.

RT-qPCR

Total RNA was extracted using the Quick-RNA miniprep kit (Zymo Research, Irvine, CA,

USA). A total of 1 µg of RNA was incubated with DNase I for 15min and reverse

transcribed with the High Capacity cDNA Reverse Transcription Kit (Thermo Fisher,

Waltham, MA USA). The resulting cDNA was used for qPCR. 10 μl SYBR Green

Universal master mix (Bio-Rad) was used for qPCR according to the manufacturer's

instructions. Each sample was normalized to 18S ribosomal RNA, and the ddCt fold

change method was used to calculate relative quantification. All reactions were run in

triplicate. Primer sequences used for qRT-PCR are provided in the **Table. S1.**

Immunoblotting

Cells were washed with PBS, lysed in lysis buffer (50 mM Tris-HCl [pH 6.8], 2% SDS,

10% glycerol) and boiled for 3min. The protein concentrations of the lysates were

quantified with a BCA Protein Assay Kit (Thermo Fisher). Protein samples were

separated by SDS-PAGE using 10% agarose gel and transferred to transfer membranes

(Millipore-Sigma, St. Louis, MO, USA), which were incubated in 5% nonfat milk (A600669,

Sangon) at room temperature for 2 hours. The membrane was incubated with the primary

antibody at 4°C overnight or at room temperature for 2 hours. The membrane was then

incubated with horseradish-peroxidase-conjugated secondary antibody or Alexa-647-

conjugated secondary antibody at 25°C for 1 hour. For cell fractionation, cells were

suspended with hypotonic buffer (20mM Tris-HCl(pH 7.4), 10mM NaCl, 3mM MgCl2,

0.5mM DTT and proteinase inhibitor cocktail) for 15min on ice and 0.5% NP-40 was

added. Cells were then centrifuged for 10 minutes at 3000rpm to collect the supernatants

(cytoplasm fraction). Pellets were washed with PBS 2 times and suspended with protein

lysis buffer (nuclear fraction).

Flow cytometry

Cells in culture were washed twice with PBS and resuspended in FACS buffer (PBS

supplemented with 1% FBS) at 1×10^6 cells/ml. Cells were then stained for 2 hours at room

temperature with the gp130 antibodies (1:1000) followed by Alexa-647-conjugated

secondary antibody for 1 hour. For intracellular nuclear staining to measure cell cycle,

cells were washed twice with PBS and incubated with propidium iodide (PI) solution

(50ng/ml PI, 0.2% NP-40, 0,25mg/ml RNase) for 15 min at 4°C and then for 30 min at

37°C in the dark. Flow cytometry was carried out by using a BD Acuri instrument (BD

Biosciences) and data analysis was performed using FlowJo v10.8.1 (Tree Star) by gating

on live cells based on forward versus side scatter profiles. Cell cycle was calculated by

Watson model implemented in FlowJo according to the manufacture's protocol.

Capture of nascent RNAs

To capture nascent RNAs, 0.4 mM EdU was added into the culture medium and was

incorporated into the cells for 2 hours. Total RNA was prepared with (). The EdU-labeled

RNAs were then biotinylated and captured by using the Click-it Nascent RNA Capture Kit

(Life Technologies), in accordance with the manufacturer's instructions. Briefly, 1 µg of

EU-labeled RNA was biotinylated with 0.5 mM biotin azide in Click-iT reaction buffer. The

biotinylated RNAs were precipitated with ethanol at 4°C overnight and resuspended in 50

µl distilled water. The biotinylated RNAs were then mixed with 12 µl Dynabeads MyOne

Streptavidin T1 magnetic beads in Click-iT RNA binding buffer and heated at 68°C for 5

min, followed by incubation at room temperature for 30 min while gently vortexing. The

beads were immobilized using the DynaMag-2 magnet and were washed with Click-iT

wash buffers 1 and 2. The washed beads were resuspended in Click-iT wash buffer 2

and used for cDNA synthesis. cDNA synthesis was performed by the High Capacity cDNA

Reverse Transcription Kit as described above.

Statistical analysis

Experimental replicates of at least 3 for each sample, including negative controls, were

prepared whenever applicable. Results are shown as mean ± SD from at least three

independent experiments. Statistical analyses were performed using GraphPad Prism

9.4.1 software. Statistical significance was determined by appropriate testing with

Student's t-test and one-way ANOVA with Tukey's multiple comparison test. A value of p

< 0.05 was considered statistically significant.

Data availability

The RNA-seq, SLAMseq, and CUT and RUN data were deposited in NCBI Gene

Expression Omnibus (GEO) database under accession numbers; GSE232843,

GSE233119, and GSE233323.

Acknowledgments

We would like to thank Dr. Robert Yarchoan (NIH/NCI) for the generous gift of rabbit monoclonal antibody against vIL-6. We would like to thank the Flow Cytometry and Genomics Shared Resources at the UC Davis Comprehensive Cancer Center for expert support. This research was supported by public health grants from the National Cancer Institute (CA225266, CA232845) and the National Institute of Allergy and Infectious Disease (AI155515 and AI167663) to Y.I. The Genomics and Flow Cytometry Shared Resources are supported by the UC Davis Comprehensive Cancer Center Support Grant (CCSG) awarded by the National Cancer Institute (NCI P30CA093373).

Figure legends

Figure 1. Biological effects of prolonged vIL-6 exposure **(A)** A schematic diagram of the THP-1 cell culture model. THP-1 cells were treated with or without vIL-6 every other day for 2 weeks (THP-1, vIL-6/THP-1). After withdrawal of the cytokine for 24 hours. vIL-6, hIL-6 and TGF-β were added for 30 minutes for secondary stimulation. Subsequently, 4-Thiouridine (4sU, 300 µM) was added to the culture media and the cells were incubated for 1 hr to label newly synthesized RNA. Newly synthesized RNA was then alkylated by iodoacetamide (IAA, 100mM). Total RNA was isolated for the following analyses. **(B)** Principal component analysis (PCA). Nascent transcribed gene expression in THP-1 and vIL-6/THP-1 cells was shown. vIL-6, hIL-6 and TGF-β were used as secondary stimulation. Untreated THP-1 cells were used as a control. The samples were represented by three biological replicates. The x and y axes show the percentage of variance explained by PC1 (55% variance) and PC2 (22% variance). **(C)** The number of upregulated genes (log2 fold change >1, adj p-value < 0.01) after vIL-6 stimulation (left) and TGF-β stimulation (right). Red and green circles represent THP-1 and vIL-6/THP-1 cells, respectively. **(D)** Comparison of up-regulated gene expression between parent THP-1 and vIL-6/THP-1 cells after stimulation with vIL-6 (left, N = 344) or TGFβ (right, N = 484). Data were analyzed using Wilcoxon matched-pairs signed ranked test and shown as median. **(E)**

KEGG pathway analysis performed on up-regulated genes (log2 fold change >1, adj *p*-

value < 0.01) in either THP-1 or vIL-6/THP-1 cells with vIL-6 stimulation. Each bar

represents the pathways that were enriched only in parental THP-1 cells (red), only in

vIL-6/THP-1 cells (green), or commonly enriched (light green). Results are presented in

descending order of the analysis. **(F)** Comparison of all KSHV gene expression between

parent THP-1 and vIL-6/THP-1 cells after KSHV infection (N = 88). Data were analyzed

using Wilcoxon matched-pairs signed ranked test and shown as median. **(G)** RNA-

sequence reads aligned to the KSHV genome (NC 009333.1) in parent THP-1 (red) and

vIL-6/THP-1 cells (blue) after KSHV infection.

Figure 2. Newly BRD4 recruitment and H3K27Ac translocation in vIl-6/THP-1 cells

(A) Enrichment analysis by CHIP-Atlas.   Enrichment analysis was performed on the list

of genes whose expression were up-regulated by vIL-6 stimulation. The parameters used

were as follows; cell type class: blood, threshold for significance: 100, threshold for log10

adjusted *p* value < -10. The Venn diagram shows the relatedness and number of genes

up-regulated after vIL-6 stimulation in parental THP-1 (red circle) and vIL-6/THP-1 cells

(green circle). **(B)** The predicted transcription factors or mediators selectively enriched

only in vIL-6/THP-1 cells. The results were limited to THP-1 cells in Chip-Atlas database.

(C) CUT&RUN signals in ±5-kbp windows around the center of peaks in vIL-6/THP-1 cells.

The list of peaks (p-value < 10^{-4}) was extracted using findPeaks (HOMER with default

parameters). Enrichment is shown in vIL-6/THP-1 (blue line) and in THP-1 (red line) cells.

Images were drawn by plotProfile (HOMER). **(D)** H3K27Ac marks in parental THP-1 and

vIL-6/THP-1 cells. The number of H3K27Ac peaks in vIL-6/THP-1 and parental THP-1

cells are depicted by a Venn diagram. **(E)** BRD4 CUT&RUN signals in vIL-6/THP-1 cells.

The green line indicates BRD4 peaks±5-kbp around the center of H3K27Ac peaks in vIL-

6/THP1 cells while the pink line indicates those in parental THP-1 cells. **(F)** Schematic

model of H3K27Ac translocation and BRD4 recruitment at newly emerged H3K27Ac

regions in vIL-6/THP-1 cells. **(G)** Distances of the nearest TSS to overlapping peaks of

BRD4 binding and H3K27Ac enrichment sites. Total peaks counts are shown. *P* values

were calculated by the Kolmogorov–Smirnov test. **(H)** Annotation of overlapping peaks

between BRD4 and H3K27Ac enrichment sites. Each annotation and its proportion were

calculated by annotatePeaks.pl (Homer; parameters; default). Promoter regions are

defined as ± 1-kbp from TSS. **(I)** BRD4 CUT&RUN signals in ±5-kbp windows around the

center of CTCF peaks in vIL-6/THP-1 cells (blue) and parental THP-1 cells (pink).

Figure 3. Prolonged vIL-6 exposure enhances inflammatory response to LPS through the novel accumulation of BRD4 and H3K27Ac (A) Heatmap showing the results of Olink® Target 48 Cytokine panel analysis. LPS (100ng/ml) was added to parental THP-1 or vIL-6/THP-1 cells for 6 hours. Cytokine production in untreated THP-1 cells was set as 1 and $\log_2$ fold activation relative to untreated THP-1 cells is shown. Samples were prepared in triplicate and the mean value were shown. **(B)** Individual inflammatory cytokine production determined by Olink proximity extension assay. Data was analyzed using two-sided unpaired Student's *t* test and shown as mean ± SD. **(C)** PCA based on nascent transcribed gene expression in THP-1 cells and vIL-6/THP-1 cells after LPS stimulation. The samples were represented by 3 biological replicates. The x and y axes show the percentage of variance explained by PC1 (98% variance) and PC2 (1% variance). **(D)** The number of up-regulated genes (log2 fold change >1, adj *p*-value < 0.01) in THP-1 and vIL-6/THP-1 cells after LPS stimulation (left) and KEGG pathway analysis performed on commonly up-regulated genes   (right). **(E)** Schematic diagram for comparing peaks score of up-regulated genes between THP-1 and vIL-6/THP-1 cells. Commonly up-regulated genes by LPS stimulation were obtained from SLAMseq data and BRD4/H3K27Ac merge peaks score were obtained from CUT&RUN data. The peak scores were associated with each up-regulated gene. **(F)** BRD4/H3K27Ac merge peak

score for common upregulated genes in THP-1 and vIL-6/THP-1 cells (N = 1626). Data

were analyzed using Wilcoxon matched-pairs signed ranked test and shown as median.

(G) BRD4/H3K27Ac peak score between up-regulated genes (N = 1626) and unchanged

genes (log2 fold change >1 and < -1, adj *p*-value > 0.01) (N = 4582) in THP-1 (left) and

vIL-6/THP-1 cells (right). Data were analyzed using Mann-Whitney test and shown as

median.

Figure 4. Association of BRD4 and p65 is responsible for prolonging hIL-6

transcription burst by LPS **(A)** Proximity extension assay (PLA) to visualize BRD4 and

p65 interaction. THP-1 and vIL-6/THP-1 cells were incubated with LPS (100ng/ml) for 6

hours, fixed by 4% PFA/PBS for 15min, permeabilized by 0.15% TritonX/PBS for 10min

and PLA was performed. Representative images of PLA (red dots) and quantification of

PLA are shown (right). Data were analyzed using a one-way ANOVA test. **(B)**

Immunoblotting of p65, H3K27me3 and β-actin in the nuclear and cytoplasmic fractions.

THP-1 and vIL-6/THP-1 cells were incubated with LPS (100ng/ml) for 6 hours and cell

fractionation was performed. **(C)** The levels of p65 enrichment in the hIL-6, hIL-10 and IL-

1β promoter regions were detected using ChIP-qPCR. Normal Rabbit IgG antibody was

used as negative control. Data was analyzed using two-sided unpaired Student's *t* test

and shown as mean ± SD. **(D)** Nascent RNA expression in THP-1 and vIL-6/THP-1 cells

after LPS stimulation. Cells were incubated with LPS and ethynyl uridine (EU) was added

0h, 1h 2h, 4h, 8h, 12h after LPS stimulation for 1 hour. Mock samples were prepared

without EU for 1 hour incubation with LPS. **(E)** The level of hIL-6, IL-10 and IL-1β gene

expression in response to OTX-015 treatment. THP-1 and vIL-6/THP-1 cells were

incubated with OTX-015 (10nM) for 4 hours followed by LPS (100ng/ml) for 1 hour. RNA

was then collected and transcribed into cDNA for RT-qPCR. 18S rRNA expression was

used for internal control. Data were analyzed using a one-way ANOVA test.

Figure. 5 Schematic model of transcription reprogramming by prolonged vIL-6

exposure Signal-dependent transcription factors such as STAT3 are recruited at up-

regulated gene promoters by vIL-6, and BRD4 and H3K27Ac are increasingly co-

occupied at those promoters by prolonged vIL-6 exposure. After the removal of the

stimulus, STAT3 is removed from the promoters; however, the activated H3K27Ac

modification and BRD4 remained and established a "primed" chromatin state for

subsequent activation. Subsequent LPS stimulation activated hIL-6 or IL-10 genes that

had been primed by vIL-6, more quickly and robustly.

References

1.   Mizobuchi H, Yamamoto K, Tsutsui S, Yamashita M, Nakata Y, Inagawa H, et al. A unique hybrid characteristic having both pro- and anti-inflammatory phenotype transformed by repetitive low-dose lipopolysaccharide in C8-B4 microglia. Sci Rep. 2020;10(1):8945. Epub 20200602. doi: 10.1038/s41598-020-65998-8. PubMed PMID: 32488176; PubMed Central PMCID: PMCPMC7265460.

2.   Morein D, Rubinstein-Achiasaf L, Brayer H, Dorot O, Pichinuk E, Ben-Yaakov H, et al. Continuous Inflammatory Stimulation Leads via Metabolic Plasticity to a Prometastatic Phenotype in Triple-Negative Breast Cancer Cells. Cells. 2021;10(6). Epub 20210531. doi: 10.3390/cells10061356. PubMed PMID: 34072893; PubMed Central PMCID: PMCPMC8229065.

3.   Ajmone-Cat MA, Nicolini A, Minghetti L. Prolonged exposure of microglia to lipopolysaccharide modifies the intracellular signaling pathways and selectively promotes prostaglandin E2 synthesis. J Neurochem. 2003;87(5):1193-203. doi: 10.1046/j.1471-4159.2003.02087.x. PubMed PMID: 14622099.

4.   Twayana KS, Chaudhari N, Ravanan P. Prolonged lipopolysaccharide exposure induces transient immunosuppression in BV2 microglia. J Cell Physiol. 2019;234(2):1889-903. Epub 20180727. doi: 10.1002/jcp.27064. PubMed PMID: 30054903.

5.   Chang Y, Cesarman E, Pessin MS, Lee F, Culpepper J, Knowles DM, et al. Identification

of herpesvirus-like DNA sequences in AIDS-associated Kaposi's sarcoma. Science.

1994;266(5192):1865-9. doi: 10.1126/science.7997879. PubMed PMID: 7997879.

6.   Antman K, Chang Y. Kaposi's sarcoma. N Engl J Med. 2000;342(14):1027-38. doi:

10.1056/NEJM200004063421407. PubMed PMID: 10749966.

7.   Giffin L, Damania B. KSHV: pathways to tumorigenesis and persistent infection. Adv

Virus Res. 2014;88:111-59. doi: 10.1016/B978-0-12-800098-4.00002-7. PubMed PMID:

24373311; PubMed Central PMCID: PMCPMC4104069.

8.   Polizzotto MN, Uldrick TS, Wyvill KM, Aleman K, Marshall V, Wang V, et al. Clinical

Features and Outcomes of Patients With Symptomatic Kaposi Sarcoma Herpesvirus (KSHV)-

associated Inflammation: Prospective Characterization of KSHV Inflammatory Cytokine

Syndrome (KICS). Clin Infect Dis. 2016;62(6):730-8. Epub 2015/12/15. doi: 10.1093/cid/civ996.

PubMed PMID: 26658701; PubMed Central PMCID: PMCPMC4772848.

9.   Chandriani S, Ganem D. Array-based transcript profiling and limiting-dilution reverse

transcription-PCR analysis identify additional latent genes in Kaposi's sarcoma-associated

herpesvirus. J Virol. 2010;84(11):5565-73. Epub 20100310. doi: 10.1128/JVI.02723-09. PubMed

PMID: 20219929; PubMed Central PMCID: PMCPMC2876603.

10.   Aoki Y, Jaffe ES, Chang Y, Jones K, Teruya-Feldstein J, Moore PS, et al. Angiogenesis

and hematopoiesis induced by Kaposi's sarcoma-associated herpesvirus-encoded interleukin-6.

Blood. 1999;93(12):4034-43. Epub 1999/06/11. PubMed PMID: 10361100.

11.   Li W, Wang Q, Qi X, Guo Y, Lu H, Chen Y, et al. Viral interleukin-6 encoded by an

oncogenic virus promotes angiogenesis and cellular transformation by enhancing STAT3-

mediated epigenetic silencing of caveolin 1. Oncogene. 2020;39(23):4603-18. Epub 20200511.

doi: 10.1038/s41388-020-1317-1. PubMed PMID: 32393833; PubMed Central PMCID:

PMCPMC7970339.

12.   Zhu X, Guo Y, Yao S, Yan Q, Xue M, Hao T, et al. Synergy between Kaposi's sarcoma-

associated herpesvirus (KSHV) vIL-6 and HIV-1 Nef protein in promotion of angiogenesis and

oncogenesis: role of the AKT signaling pathway. Oncogene. 2014;33(15):1986-96. Epub

20130422. doi: 10.1038/onc.2013.136. PubMed PMID: 23604117.

13.   Giffin L, West JA, Damania B. Kaposi's Sarcoma-Associated Herpesvirus Interleukin-6

Modulates Endothelial Cell Movement by Upregulating Cellular Genes Involved in Migration. mBio.

2015;6(6):e01499-15. Epub 20151208. doi: 10.1128/mBio.01499-15. PubMed PMID: 26646010;

PubMed Central PMCID: PMCPMC4676281.

14.   Fullwood RA, Low GM, Chase EP, Grasley M, Beal SS, McCrary IM, et al. The Kaposi's

sarcoma-associated herpesvirus viral interleukin 6 gene affects metastasis and expression of B

cell markers in a murine xenograft model. PLoS One. 2018;13(9):e0204947. Epub 20180928. doi:

10.1371/journal.pone.0204947.   PubMed   PMID: 30265712;   PubMed   Central   PMCID:

PMCPMC6161906.

15.　　Yadav A, Kalita A, Dhillon S, Banerjee K. JAK/STAT3 pathway is involved in survival of

neurons in response to insulin-like growth factor and negatively regulated by suppressor of

cytokine signaling-3. J Biol Chem. 2005;280(36):31830-40. Epub 20050705. doi:

10.1074/jbc.M501316200. PubMed PMID: 15998644.

16.　　Haura EB, Turkson J, Jove R. Mechanisms of disease: Insights into the emerging role of

signal transducers and activators of transcription in cancer. Nat Clin Pract Oncol. 2005;2(6):315-

24. doi: 10.1038/ncponc0195. PubMed PMID: 16264989.

17.　　Yu H, Jove R. The STATs of cancer--new molecular targets come of age. Nat Rev Cancer.

2004;4(2):97-105. doi: 10.1038/nrc1275. PubMed PMID: 14964307.

18.　　Hirano T. IL-6 in inflammation, autoimmunity and cancer. Int Immunol. 2021;33(3):127-

48. doi: 10.1093/intimm/dxaa078. PubMed PMID: 33337480; PubMed Central PMCID:

PMCPMC7799025.

19.　　Fu B, Xu X, Wei H. Why tocilizumab could be an effective treatment for severe COVID-

19? J Transl Med. 2020;18(1):164. doi: 10.1186/s12967-020-02339-3. PubMed PMID: 32290839;

PubMed Central PMCID: PMCPMC7154566.

20.　　Ramaswami R, Lurain K, Peer CJ, Serquina A, Wang V, Widell A, et al. Tocilizumab in

patients with symptomatic Kaposi sarcoma herpesvirus-associated multicentric Castleman

disease. Blood. 2020;135(25):2316-9. doi: 10.1182/blood.2019004602. PubMed PMID:

32276276; PubMed Central PMCID: PMCPMC7316216.

21.   Nord JA, Wynia-Smith SL, Gehant AL, Jones Lipinski RA, Naatz A, Rioja I, et al. N-

terminal BET bromodomain inhibitors disrupt a BRD4-p65 interaction and reduce inducible nitric

oxide synthase transcription in pancreatic beta-cells. Front Endocrinol (Lausanne).

2022;13:923925. Epub 20220913. doi: 10.3389/fendo.2022.923925. PubMed PMID: 36176467;

PubMed Central PMCID: PMCPMC9513428.

22.   Huang B, Yang XD, Zhou MM, Ozato K, Chen LF. Brd4 coactivates transcriptional

activation of NF-kappaB via specific binding to acetylated RelA. Mol Cell Biol. 2009;29(5):1375-

87. Epub 2008/12/24. doi: 10.1128/MCB.01365-08. PubMed PMID: 19103749; PubMed Central

PMCID: PMCPMC2643823.

23.   Sabari BR, Dall'Agnese A, Boija A, Klein IA, Coffey EL, Shrinivas K, et al. Coactivator

condensation at super-enhancers links phase separation and gene control. Science.

2018;361(6400). Epub 2018/06/23. doi: 10.1126/science.aar3958. PubMed PMID: 29930091;

PubMed Central PMCID: PMCPMC6092193.

24.   Parravicini C, Corbellino M, Paulli M, Magrini U, Lazzarino M, Moore PS, et al.

Expression of a virus-derived cytokine, KSHV vIL-6, in HIV-seronegative Castleman's disease.

Am J Pathol. 1997;151(6):1517-22. PubMed PMID: 9403701; PubMed Central PMCID:

PMCPMC1858372.

25.   Shimoda M, Inagaki T, Davis R, Merleev A, Tepper CG, Maverakis E, et al. KSHV uses

viral IL6 to expand infected immunosuppressive macrophages. bioRxiv. 2023. Epub 2023/03/23.

doi: 10.1101/2023.03.05.531224. PubMed PMID: 36945595; PubMed Central PMCID:

PMCPMC10028810.

26.   Shi C, Pamer EG. Monocyte recruitment during infection and inflammation. Nat Rev

Immunol. 2011;11(11):762-74. Epub 2011/10/11. doi: 10.1038/nri3070. PubMed PMID: 21984070;

PubMed Central PMCID: PMCPMC3947780.

27.   Herzog VA, Reichholf B, Neumann T, Rescheneder P, Bhat P, Burkard TR, et al. Thiol-

linked alkylation of RNA to assess expression dynamics. Nat Methods. 2017;14(12):1198-204.

Epub 2017/09/26. doi: 10.1038/nmeth.4435. PubMed PMID: 28945705; PubMed Central PMCID:

PMCPMC5712218.

28.   Kanno T, Kanno Y, LeRoy G, Campos E, Sun HW, Brooks SR, et al. BRD4 assists

elongation of both coding and enhancer RNAs by interacting with acetylated histones. Nat Struct

Mol Biol. 2014;21(12):1047-57. Epub 20141110. doi: 10.1038/nsmb.2912. PubMed PMID:

25383670; PubMed Central PMCID: PMCPMC4720983.

29.   Skene PJ, Henikoff S. An efficient targeted nuclease strategy for high-resolution mapping

of DNA binding sites. Elife. 2017;6. Epub 2017/01/13. doi: 10.7554/eLife.21856. PubMed PMID:

28079019; PubMed Central PMCID: PMCPMC5310842.

30.   Aoki Y, Feldman GM, Tosato G. Inhibition of STAT3 signaling induces apoptosis and

decreases survivin expression in primary effusion lymphoma. Blood. 2003;101(4):1535-42. Epub

2002/10/24. doi: 10.1182/blood-2002-07-2130. PubMed PMID: 12393476.

31.   Grossmann C, Ganem D. Effects of NFkappaB activation on KSHV latency and lytic

reactivation are complex and context-dependent. Virology. 2008;375(1):94-102. Epub 20080305.

doi: 10.1016/j.virol.2007.12.044. PubMed PMID: 18321555; PubMed Central PMCID:

PMCPMC2822626.

32.   Lundberg M, Eriksson A, Tran B, Assarsson E, Fredriksson S. Homogeneous antibody-

based proximity extension assays provide sensitive and specific detection of low-abundant

proteins in human blood. Nucleic Acids Res. 2011;39(15):e102. Epub 20110606. doi:

10.1093/nar/gkr424. PubMed PMID: 21646338; PubMed Central PMCID: PMCPMC3159481.

33.   Han X, Yu D, Gu R, Jia Y, Wang Q, Jaganathan A, et al. Roles of the BRD4 short isoform

in phase separation and active gene transcription. Nat Struct Mol Biol. 2020;27(4):333-41. Epub

20200316. doi: 10.1038/s41594-020-0394-8. PubMed PMID: 32203489.

34.   Mullberg J, Geib T, Jostock T, Hoischen SH, Vollmer P, Voltz N, et al. IL-6 receptor

independent stimulation of human gp130 by viral IL-6. J Immunol. 2000;164(9):4672-7. doi:

10.4049/jimmunol.164.9.4672. PubMed PMID: 10779772.

35.   Chow D, He X, Snow AL, Rose-John S, Garcia KC. Structure of an extracellular gp130

cytokine   receptor   signaling   complex.   Science.   2001;291(5511):2150-5.   doi:

10.1126/science.1058308. PubMed PMID: 11251120.

36.   Mizobuchi H, Yamamoto K, Tsutsui S, Yamashita M, Nakata Y, Inagawa H, et al. A unique

hybrid characteristic having both pro- and anti-inflammatory phenotype transformed by repetitive

low-dose   lipopolysaccharide   in   C8-B4   microglia.   Scientific   Reports.   2020;10(1).   doi:

10.1038/s41598-020-65998-8.

37.   Boulanger MJ, Chow DC, Brevnova EE, Garcia KC. Hexameric structure and assembly

of the interleukin-6/IL-6 alpha-receptor/gp130 complex. Science. 2003;300(5628):2101-4. doi:

10.1126/science.1083901. PubMed PMID: 12829785.

38.   Heinrich PC, Behrmann I, Muller-Newen G, Schaper F, Graeve L. Interleukin-6-type

cytokine signalling through the gp130/Jak/STAT pathway. Biochem J. 1998;334 (Pt 2)(Pt 2):297-

314. Epub 1998/08/26. doi: 10.1042/bj3340297. PubMed PMID: 9716487; PubMed Central

PMCID: PMCPMC1219691.

39.   Boulanger MJ, Chow DC, Brevnova E, Martick M, Sandford G, Nicholas J, et al.

Molecular mechanisms for viral mimicry of a human cytokine: activation of gp130 by HHV-8

interleukin-6. J Mol Biol. 2004;335(2):641-54. Epub 2003/12/16. doi: 10.1016/j.jmb.2003.10.070.

PubMed PMID: 14672670.

40.   Meads MB, Medveczky PG. Kaposi's sarcoma-associated herpesvirus-encoded viral

interleukin-6 is secreted and modified differently than human interleukin-6: evidence for a unique

autocrine signaling mechanism. J Biol Chem. 2004;279(50):51793-803. Epub 2004/07/20. doi:

10.1074/jbc.M407382200. PubMed PMID: 15258150.

41.   Hu F, Nicholas J. Signal transduction by human herpesvirus 8 viral interleukin-6 (vIL-6)

is modulated by the nonsignaling gp80 subunit of the IL-6 receptor complex and is distinct from

signaling induced by human IL-6. J Virol. 2006;80(21):10874-8. Epub 2006/09/08. doi:

10.1128/JVI.00767-06. PubMed PMID: 16956953; PubMed Central PMCID: PMCPMC1641745.

42.   Osborne J, Moore PS, Chang Y. KSHV-encoded viral IL-6 activates multiple human IL-6

signaling pathways. Hum Immunol. 1999;60(10):921-7. Epub 1999/11/24. doi: 10.1016/s0198-

8859(99)00083-x. PubMed PMID: 10566591.

43.   Chen D, Choi YB, Sandford G, Nicholas J. Determinants of secretion and intracellular

localization of human herpesvirus 8 interleukin-6. J Virol. 2009;83(13):6874-82. Epub 2009/04/24.

doi: 10.1128/JVI.02625-08.   PubMed   PMID: 19386721;   PubMed   Central   PMCID:

PMCPMC2698573.

44.   Chen D, Sandford G, Nicholas J. Intracellular signaling mechanisms and activities of

human   herpesvirus   8   interleukin-6.   J   Virol.   2009;83(2):722-33.   Epub   2008/11/07.   doi:

10.1128/JVI.01517-08. PubMed PMID: 18987143; PubMed Central PMCID: PMCPMC2612405.

45.   Schultze JL, Schmieder A, Goerdt S. Macrophage activation in human diseases. Semin

Immunol. 2015;27(4):249-56. Epub 20150822. doi: 10.1016/j.smim.2015.07.003. PubMed PMID:

26303100.

46.   Novakovic B, Habibi E, Wang SY, Arts RJW, Davar R, Megchelenbrink W, et al. beta-

Glucan Reverses the Epigenetic State of LPS-Induced Immunological Tolerance. Cell.

2016;167(5):1354-68 e14. doi: 10.1016/j.cell.2016.09.034. PubMed PMID: 27863248; PubMed

Central PMCID: PMCPMC5927328.

47.   Saeed S, Quintin J, Kerstens HH, Rao NA, Aghajanirefah A, Matarese F, et al. Epigenetic

programming of monocyte-to-macrophage differentiation and trained innate immunity. Science.

2014;345(6204):1251086. doi: 10.1126/science.1251086. PubMed PMID: 25258085; PubMed

Central PMCID: PMCPMC4242194.

48.   Moore PS, Boshoff C, Weiss RA, Chang Y. Molecular mimicry of human cytokine and

cytokine response pathway genes by KSHV. Science. 1996;274(5293):1739-44. doi:

10.1126/science.274.5293.1739. PubMed PMID: 8939871.

49.   Kumar A, Lyu Y, Yanagihashi Y, Chantarasrivong C, Majerciak V, Salemi M, et al. KSHV

episome tethering sites on host chromosomes and regulation of latency-lytic switch by CHD4.

Cell Rep. 2022;39(6):110788. doi: 10.1016/j.celrep.2022.110788. PubMed PMID: 35545047;

PubMed Central PMCID: PMCPMC9153692.

50.   Fong CY, Gilan O, Lam EY, Rubin AF, Ftouni S, Tyler D, et al. BET inhibitor resistance

emerges from leukaemia stem cells. Nature. 2015;525(7570):538-42. Epub 20150914. doi:

10.1038/nature14888. PubMed PMID: 26367796; PubMed Central PMCID: PMCPMC6069604.

51.   Zhou S, Zhang S, Wang L, Huang S, Yuan Y, Yang J, et al. BET protein inhibitor JQ1

downregulates chromatin accessibility and suppresses metastasis of gastric cancer via

inactivating RUNX2/NID1 signaling. Oncogenesis. 2020;9(3):33. Epub 20200310. doi:

10.1038/s41389-020-0218-z. PubMed PMID: 32157097; PubMed Central PMCID:

PMCPMC7064486.

52.   Izumiya Y, Izumiya C, Hsia D, Ellison TJ, Luciw PA, Kung HJ. NF-kappaB serves as a

cellular sensor of Kaposi's sarcoma-associated herpesvirus latency and negatively regulates K-

Rta by antagonizing the RBP-Jkappa coactivator. J Virol. 2009;83(9):4435-46. Epub 2009/02/27.

doi: 10.1128/JVI.01999-08. PubMed PMID: 19244329; PubMed Central PMCID:

PMCPMC2668470.

53.   Kim KY, Huerta SB, Izumiya C, Wang DH, Martinez A, Shevchenko B, et al. Kaposi's

sarcoma-associated herpesvirus (KSHV) latency-associated nuclear antigen regulates the KSHV

epigenome by association with the histone demethylase KDM3A. J Virol. 2013;87(12):6782-93.

Epub 2013/04/12. doi: 10.1128/JVI.00011-13. PubMed PMID: 23576503; PubMed Central

PMCID: PMCPMC3676133.

54. Neumann T, Herzog VA, Muhar M, von Haeseler A, Zuber J, Ameres SL, et al.

Quantification of experimentally induced nucleotide conversions in high-throughput sequencing

datasets. BMC Bioinformatics. 2019;20(1):258. Epub 20190520. doi: 10.1186/s12859-019-2849-

7. PubMed PMID: 31109287; PubMed Central PMCID: PMCPMC6528199.

55. Sedlazeck FJ, Rescheneder P, von Haeseler A. NextGenMap: fast and accurate read

mapping in highly polymorphic genomes. Bioinformatics. 2013;29(21):2790-1. Epub 20130823.

doi: 10.1093/bioinformatics/btt468. PubMed PMID: 23975764.

56. Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, et al. VarScan 2: somatic

mutation and copy number alteration discovery in cancer by exome sequencing. Genome Res.

2012;22(3):568-76. Epub 20120202. doi: 10.1101/gr.129684.111. PubMed PMID: 22300766;

PubMed Central PMCID: PMCPMC3290792.

57. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550. doi: 10.1186/s13059-014-0550-8.

PubMed PMID: 25516281; PubMed Central PMCID: PMCPMC4302049.

58. Sherman BT, Hao M, Qiu J, Jiao X, Baseler MW, Lane HC, et al. DAVID: a web server

for functional enrichment analysis and functional annotation of gene lists (2021 update). Nucleic

Acids Res. 2022;50(W1):W216-21. Epub 20220323. doi: 10.1093/nar/gkac194. PubMed PMID:

35325185; PubMed Central PMCID: PMCPMC9252805.

59.   Zou Z, Ohta T, Miura F, Oki S. ChIP-Atlas 2021 update: a data-mining suite for exploring

epigenomic landscapes by fully integrating ChIP-seq, ATAC-seq and Bisulfite-seq data. Nucleic

Acids Res. 2022;50(W1):W175-82. Epub 20220324. doi: 10.1093/nar/gkac199. PubMed PMID:

35325188; PubMed Central PMCID: PMCPMC9252733.

60.   Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-

aware quantification of transcript expression. Nat Methods. 2017;14(4):417-9. Epub 20170306.

doi: 10.1038/nmeth.4197.   PubMed   PMID: 28263959;   PubMed   Central   PMCID:

PMCPMC5600148.

61.   Soneson C, Love MI, Robinson MD. Differential analyses for RNA-seq: transcript-level

estimates   improve   gene-level   inferences.   F1000Res.   2015;4:1521.   Epub   20151230.   doi:

10.12688/f1000research.7563.2.   PubMed   PMID: 26925227;   PubMed   Central   PMCID:

PMCPMC4712774.

62.   Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics.   2018;34(17):i884-i90.   doi: 10.1093/bioinformatics/bty560.   PubMed   PMID:

30423086; PubMed Central PMCID: PMCPMC6129281.

63.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.

2012;9(4):357-9. Epub 20120304. doi: 10.1038/nmeth.1923. PubMed PMID: 22388286; PubMed

Central PMCID: PMCPMC3322381.

64.   Ramirez F, Ryan DP, Gruning B, Bhardwaj V, Kilpert F, Richter AS, et al. deepTools2: a

next generation web server for deep-sequencing data analysis. Nucleic Acids Res.

2016;44(W1):W160-5. Epub 20160413. doi: 10.1093/nar/gkw257. PubMed PMID: 27079975;

PubMed Central PMCID: PMCPMC4987876.

65.   Freese NH, Norris DC, Loraine AE. Integrated genome browser: visual analytics platform

for genomics. Bioinformatics. 2016;32(14):2089-95. Epub 20160316. doi:

10.1093/bioinformatics/btw069. PubMed PMID: 27153568; PubMed Central PMCID:

PMCPMC4937187.

66.   Assarsson E, Lundberg M, Holmquist G, Bjorkesten J, Thorsen SB, Ekman D, et al.

Homogenous 96-plex PEA immunoassay exhibiting high sensitivity, specificity, and excellent

scalability. PLoS One. 2014;9(4):e95192. Epub 20140422. doi: 10.1371/journal.pone.0095192.

PubMed PMID: 24755770; PubMed Central PMCID: PMCPMC3995906.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

SFig.1 hIL-6 is a functional homolog of vIL-6 (A) The number of up-regulated genes (log2 fold change >1, adj *p*-value < 0.01) after hIL-6 stimulation. Red circle represents THP-1 cells and green circle represents vIL-6/THP-1 cells. **(B)** Individual gene expression in THP-1 cells with vIL-6 stimulation (left, N = 8206) and hIL-6 stimulation (right, N = 8210). Representative gene names were labelled adjacent to dots. The red dashed line indicated log2 fold change = ± 1 (vertical) and -log10 adj p-value = 2 (horizontal). **(C)** KEGG pathway analysis performed on up-regulated genes (log2 fold change >1, adj *p*-value < 0.01) in THP-1 cells with vIL-6 and hIL-6 stimulation. The result showed the top three pathways each. **(D)** Individual up-regulated gene expression (N = 348) in parent THP-1 and vIL-6/THP-1 cells after hIL-6 stimulation. Data were analyzed using Wilcoxon matched-pairs signed ranked test and shown as median. **(E)** Measurement of cell proliferation with MTT assays. 1 X 10^4 THP-1 or vIL-6/THP-1 cells were cultured in triplicate in a 96 well plate. vIL-6 was added to vIL-6/THP-1 cells every other day. OD (570-690nm) was measured on day 0,1,2,4 and 7. Data were analyzed using unpaired Student's *t* test and shown as mean ± SD. **(F)** Immunoblotting with antibodies directed against STAT3, pshopho-STAT3 (Y705) and β-Actin (loading control) protein in THP-1 and vIL6/THP-1 cells. (G) FACS analysis showing the gp130 expression on cell surface.

SFig.1

SFig.2 Prolonged vIL-6 exposure enhances the association of BRD4 and H3K27Ac
(A) BRD4 CUT &RUN signals in ±5kbp windows around the transcription start sites (TSS) of up-regulated genes in vIL-6/THP-1 cells (N = 303). **(B)** BRD4, H3K27Ac and H3K4me3 protein expression before and after vIL-6 stimulation in parental THP-1 and vIL-6/THP-1 cells. **(C)** DNA binding motif analysis of new BRD4 accumulation sites in vIL-6/THP-1 cells. Images were drawn by findMotif (HOMER). **(D)** The number of BRD4 and H3K27Ac peaks and their association in parental THP-1 and vIL-6/THP-1 cells. The overlapping peaks were extracted using mergepeaks (HOMER). **(E)** RNA pol II and H3K4me3 CUT &RUN signals in ±5kbp windows around the center of BRD4 and H3K27Ac peaks. **(F)** KEGG pathway analysis performed on genes at BRD4 and H3K27 overlapping peaks in promoter regions in vIL-6/THP-1 cells. Results are presented in descending order. **(G)** BRD4, RNA pol II, H3K27Ac and H3K4me3 enrichment in the *Cyclin E1* promoter region in parental THP-1 cells (pink) and vIL-6/THP-1 cells (blue). The peaks were visualized by importing the BAM files into the Integrative Genomics Viewer (IGV).

SFig.2

SFig.3 Inflammatory response to IFNα after vIL-6 prolonged exposure (A) hIL-6 production in THP-1 cells by LPS stimulation. THP-1 cells was incubated with LPS 100ng/ml or 1µg/ml for various time periods. Supernatants were harvested and incubated in triplicate in ELISA plate coated with hIL-6 antibody, Human IL-6 Uncoated ELISA kit (Invitrogen) was then used to evaluate the hIL-6 production by following the manufacturer's guideline. The protein binding measured as OD values at 450nm was shown. Results are presented as mean percentage viability ±SD ($n = 3$ samples/group). Data was analyzed by a one-way ANOVA test. **(B)** Heatmap showing the results of Olink® Target 48 Cytokine panel . IFNα (100ng/ml) was added to parent THP-1 cells or vil-6/THP-1 cells for 6 hours. Cytokine production in untreated THP-1 cells was set as 1 and $\log_2$ fold activation relative to untreated cells are shown. Samples were prepared in triplicate and the mean value were shown. **(C)** Inflammatory cytokine production determined by Olink proximity extension assay. Data was analyzed using two-sided unpaired Student's *t* test and shown as mean ± SD.

SFig.3

SFig.4 BRD4 enrichment in the promoter region of inflammatory genes (A) BRD4, RNA pol II, H3K27Ac and H3K4me3 enrichment in hIL-6, IL-10, IL-1β promoter region in parental THP-1 cells (pink) and vIL-6/THP-1 cells (blue). The promoter region is enclosed by a black line. Each CUT&RUN peak was visualized by importing the BAM files into Integrative Genomics Viewer (IGV). **(B)** Schematic diagram of nascent RNA labeling after LPS stimulation. LPS were added to culture medium in parental THP-1 and vIL-6/THP-1 cells and cells were incubated with EU for 1 hour in 0, 1, 2, 4 8, 12 h LPS stimulation.

SFig.4

Table. S1. Primer used for RT-qPCR and Chip-qPCR

RT-qPCR primer		
hIL-6	Forward	GGTACATCCTCGACGGCATCT
	Reverse	GTGCCTCTTTGCTGCTTT
IL-10	Forward	GGTTGCCAAGCCTTGTCTGA
	Reverse	AGGGAGTTCACATGCGCCT
IL-1β	Forward	AAACAGATGAAGTGCTCCTTCCAGG
	Reverse	TGGAGAACACCACTTGTTGCTCCA
18S rRNA	Forward	TTCGAACGTCTGCCCTATCAA
	Reverse	ATGGTAGGCACGGCGACTA
Chip-qPCR primer		
hIL-6 promoter	Forward	AACTTGGTTCAGGAGTGTCTAC
	Reverse	AGGATGGCTGGATGGTTTC
IL-10 promoter	Forward	GGAGATCTCGAAGCATGTTAGG
	Reverse	GGACAGAGAGGTGAAGGTCTA
IL-1β promoter	Forward	CAGCCAATCTTCATTGCTCAAG
	Reverse	CATACACACAAAGAGGCAGAGA