Glucose-stimulated KIF5B-driven microtubule sliding organizes microtubule networks in pancreatic beta cells

Kai M. Bracey[1], Pi'illani Noguchi[1], Courtney Edwards[2], Alisa Cario[1], Guoqiang Gu[1,3,4], Irina Kaverina[1,3]

Affiliations:

[1]Department of Cell and Developmental Biology, Vanderbilt University, Nashville, TN 37235, USA

[2]Department of Medicine, Vanderbilt University Medical Center, Nashville, TN 37232, USA.

[3]Program of Developmental Biology, Vanderbilt University, Nashville, TN 37232, USA.

[4]Center for Stem Cell Biology, Vanderbilt University, Nashville, TN 37232, USA.

Corresponding author: Irina Kaverina

Abstract

In pancreatic islet beta cells, molecular motors use cytoskeletal polymers microtubules as tracks for intracellular transport of insulin secretory granules. Beta-cell microtubule network has a complex architecture and is non-directional, which provide insulin granules at the cell periphery for rapid secretion response, yet to avoid over-secretion and subsequent hypoglycemia. We have previously characterized a peripheral sub-membrane microtubule array, which is critical for withdrawal of excessive insulin granules from the secretion sites. Microtubules in beta cells originate at the Golgi in the cell interior, and how the peripheral array is formed is unknown. Using real-time imaging and photo-kinetics approaches in clonal mouse pancreatic beta cells MIN6, we now demonstrate that kinesin KIF5B, a motor protein with a capacity to transport microtubules as cargos, slides existing microtubules to the cell periphery and aligns them to each other along the plasma membrane. Moreover, like many physiological beta-cell features, microtubule sliding is facilitated by a high glucose stimulus. These new data, together with our previous report that in high glucose sub-membrane MT array is destabilized to allow for robust secretion, indicate that MT sliding is another integral part of glucose-triggered microtubule remodeling, likely replacing destabilized peripheral microtubules to prevent their loss over time and beta-cell malfunction.

Introduction

The precise level of glucose-stimulated insulin secretion (GSIS) from pancreatic beta cells is crucial for glucose homeostasis. On one hand, insufficient insulin secretion decreases glucose uptake by peripheral tissues, leading to diabetes. On the other hand, excessive secretion causes glucose depletion from the bloodstream and hypoglycemia. Not surprisingly, multiple levels of cellular regulation control the amount of insulin secretory granules (IGs) released on every stimulus. One level of this control is facilitated by microtubules (MTs), intracellular polymers which serve as tracks for intracellular transport of IGs and define how many IGs are positioned at the secretion sites (Desai & Mitchison, 1997; Heaslip et al., 2014; Varadi, Ainscow, Allan, & Rutter, 2002).

Microtubules have a dual role in regulating the availability of IGs for secretion. Microtubules are necessary for efficient IG generation at the trans Golgi network (TGN). They are also required for IG transportation throughout the cell (Trogden et al., 2019; Zhu et al., 2015), which includes non-directional, diffusion-like redistribution in the cytoplasm (Tabei et al., 2013; Zhu et al., 2015) and directional runs of secretion-competent granules toward periphery (Hoboth et al., 2015; Muller et al., 2021). At the same time, peripheral IGs undergo MT-dependent withdrawal from the secretion sites, which prevents IG docking and acute over-secretion upon a given stimulus (Hu et al., 2021; Zhu et al., 2015). Such multi-faceted involvement of MT transport in secretion regulation is made possible by a complex architecture of MT networks in beta cells. To this end, interior beta-cell MTs are twisted and interlocked (Varadi, Tsuboi, Johnson-Cadwell, Allan, & Rutter, 2003; Zhu et al., 2015), which makes them dramatically distinct from radially organized MT arrays well-studied in generic cultured cell models and explains the predominantly non-

directional nature of IG transport (Bogan, 2021; Bracey, Gu, & Kaverina, 2022). Importantly,

withdrawal of IGs and secretion restriction is achieved by a prominent array of MTs underlying

cell membrane (Bracey et al., 2020). Under basal conditions, IGs are robustly withdrawn from

the secretion sites along submembrane MTs, which are stabilized by MT-associated proteins

(MAPs), including a well-known neuronal MAP tau (Ho et al., 2020). Upon a glucose stimulus,

tau is phosphorylated and submembrane MTs become more dynamic (Ho et al., 2020) and

fragmented, possibly via a MT-severing activity (Muller et al., 2021). Destabilization and partial

depolymerization of submembrane MTs leads to IG docking and allows for secretion (Ho et al.,

2020; Hu et al., 2021), probably as a consequence of decreased IG withdrawal by MT-

dependent transport. Thus, existing data provide at least initial understanding of the

mechanisms whereby beta-cell MT architecture allows for fine-tuning of secretion levels.

However, it is yet unclear how the complex beta-cell MT network forms. As in several other

eukaryotic cells, MTs in beta cells are nucleated at MT-organizing centers (MTOCs) in the cell

interior, partially at the centrosome and to a large extent at the Golgi membranes (Trogden et

al., 2019; Zhu et al., 2015). Conventionally, this should be followed by MT plus-end

polymerization toward the cell periphery and result in a radial MT array with high MT density in

the center rather than in the periphery. The beta cell lacks such well-characterized MT polarity

(Bracey et al., 2022). Thus, it is puzzling that the actual resulting MT system is non-radial and

consists of an interior mesh a peripheral array (Bracey et al., 2020; Heaslip et al., 2014). How

the beta cell organizes its cytoskeletal network for efficient trafficking of granules, and what

factors contribute to the maintenance of the sub-membrane array, are important questions.

One of the established ways to modify the MT network without changing the location of MTOCs

is to relocate already polymerized MTs by active motor-dependent transport. This phenomenon

is called "MT sliding" (Straube, Hause, Fink, & Steinberg, 2006). Several MT-dependent

molecular motors have been implicated in driving MT sliding (Lu & Gelfand, 2017). In some

cases, a motor facilitates MT sliding by walking along a MT while its cargo-binding domain is

stationary being attached to a relatively large structure, e.g. plasma membrane. This causes

sliding of a MT which served as a track for the stationary motor. This mechanism has been

described for dynein-dependent MT sliding (Grabham, Seale, Bennecib, Goldberg, & Vallee,

2007; He et al., 2005). MTs can also be efficiently slid by motors which have two functional

motor assemblies, such as a tetrameric kinesin-5/Eg5 (Acar et al., 2013; Vukusic, Ponjavic,

Buda, Risteski, & Tolic, 2021), or which carry a MT as a cargo while walking along another MT.

For the latter mechanism, a motor needs a non-motor domain with a capacity to bind either a

MT itself, or a MT-associated protein as an adapter (Cao et al., 2020; Kurasawa, Earnshaw,

Mochizuki, Dohmae, & Todokoro, 2004; Vukusic et al., 2021).

Out of these MT-sliding factors, kinesin-1 is known to be critical for organizing unusual MT

architecture in specialized cells. In oocytes, kinesin-1-dependent MT sliding empowers

cytoplasmic streaming (Barlan, Lu, & Gelfand, 2013). In differentiating neurons, kinesin-1 moves

organelles and MTs into emerging neurites, which is a defining step in developing branched MT

networks and long-distance neuronal transport (Jolly et al., 2010; Lu, Fox, Lakonishok,

Davidson, & Gelfand, 2013). With these data in mind, kinesin-1 presents itself as the most

attractive candidate for organizing MTs in beta cells. This motor highly expressed in beta cells

and is well known to act as a major driving force in IG transport (Varadi et al., 2003).

Here, we show that KIF5B, encoding one of the three subunits of kinesin 1, actively slides MTs

in beta cells and that this phenomenon defines MT network morphology and supplies MTs for

the submembrane array. Moreover, we find that MT sliding in beta cells is a glucose-dependent

process and thus likely participates in metabolically driven cell reorganization during each

secretion cycle.

RESULTS

Identification of KIF5B as MT-sliding motor in beta cells

To address the factors that shape the configuration of MT networks in beta cells, we tested for a

potential involvement of motors-dependent MT sliding. Not surprisingly, analysis of existing

RNA-sequencing data in functional mouse islet beta cells highlighted kinesin-1 KIF5B as the

highest expressing beta-cell motor protein (Fig. 1A) (Sanavia et al., 2021). Since this kinesin

has been reported to have MT sliding activity in many types of interphase cells, we tested its

potential ability to slide MTs in beta cells.

Efficient depletion of KIF5B was achieved by utilizing two independent lentiviral-based shRNA

against mouse KIF5B in mouse insulinoma cell line MIN6 (Fig. 1B, Fig. 1- Suppl. Fig. 1A). To

visualize MT sliding, shRNA-treated MIN6 cells expressing mEmerald-tubulin were imaged by

live-cell spinning disk confocal microscopy. We photobleached MTs in two large cell regions

leaving a thin unbleached band ("fluorescent belt") and analyzed relocation of MTs from the

"fluorescent belt" into the bleached areas over time. To minimize the effects of plausible MT

polymerization and to reduce photobleaching, MTs were imaged for short time periods (5

mins). Strikingly, in control cells (treated with scrambled control shRNA) MTs were efficiently

translocated from the "fluorescent belt" into the photobleached area, indicating that MT sliding

events are prominent in this cell type (Fig. 1C,D,). In contrast, MIN6 cells expressing either

KIF5B shRNA variants displayed a significant loss of MT sliding ability (Fig. 1C,E,F, Fig.1-Video

1), indicating that the loss of KIFB leads to the loss of MT sliding.

While the assay described above provides an easy visualization of MT sliding, it allows for

visualization of only a subset of the MT network. To further corroborate the above findings, we

used a less photodamaging system to visualize MT sliding that does not involve photobleaching

and allows for evaluation of displacements within the whole MT network. To this end we applied

a microtubule probe of fiducial marks, K560Rigor^{E236A}-SunTag (Tanenbaum, Gilbert, Qi,

Weissman, & Vale, 2014) in MIN6 cells (Fig 1G-H). This probe contains the human kinesin-1

motor domain (residues 1–560) with a rigor mutation in the motor domain (K560Rigor^{E236A}) and

fused to 24 copies of a GCN4 peptide. The rigor mutation in the motor domain causes it to bind

irreversibly to microtubules (Rice et al., 1999). When co-expressed with a pHalo-tagged anti-

GCN4 single-chain antibody (ScFv-GCN4-HaloTag-GB1-NLS), K560Rigor^{E236A} can recruit up to

24 of the Halo ligands to a single position on a microtubule. The pHalo- tagged anti GCN4

construct also contains a nuclear localization signal (NLS) which lends itself to reduce

background of the unbound dye. This enables visualization of MT sliding events via single

molecule tracking of the fiducial marks along the MT lattice, allowing us to analyze MT sliding

behavior within the whole network with high temporal and spatial resolution (Fig. 1G-K, Fig. 1-

Suppl. Fig. 1B, Fig.1-Video 2).

Our data indicate that in cells treated with scrambled control shRNA, a subset of

K560Rigor^{E236A}-SunTag fiducial marks underwent rapid directional movements, interpreted as

MT sliding events (Fig 1G, K). In contrast, majority of fiducial marks in cells expressing KIF5B-

sepcific shRNAs were stationary (Fig. 1 H, I, J), indicating the lack of MT sliding. Collectively,

these results indicate that KIF5B is necessary for MT sliding in MIN6 cells.

KIF5B is required for beta-cell MT organization

Because MT sliding mediated by KIF5B is a prominent phenomenon in beta cells, we sought to

test whether it has functional consequences for MT networks in these cells. Tubulin

immunostaining revealed striking differences in MT organization between MIN6 cells treated

with scrambled control shRNA versus KIF5B-specific shRNAs. While control cells had

convoluted non-radial MTs with a prominent sub-membrane array, typical for beta cells (Fig.

2A), KIF5B-depleted cells featured extra-dense MTs in the cell center and sparse reseeding

MTs at the periphery (Fig. 2B, C). Significant reduction of tubulin staining intensity at the cell

periphery (Fig. 2D) confirms the robustness of this phenotype. This indicated that loss of KIF5B

leads to a strong defect in MT location to the cell periphery.

KIF5B is required for beta-cell sub-membrane MT array alignment

Given the known significance of the peripheral MT array, which normally consists of well-

organized MTs parallel to the cell membrane (Bracey et al., 2020), we have further analyzed

directionally of MTs remaining at the cell periphery after KIF5B depletion. Previously we

published a custom image analysis algorithm (Bracey et al., 2020) allowing for detailed

quantitative characterization of MTs directionality in relation to the nearest cell border (Fig. 2

Supplemental Fig. 1). Here, we applied the same computational analysis to MT imaging data in

MIN6 cells with perturbed KIF5B level and/or function. After deconvolution for increased signal-

to-noise ratio, single 2D slices of MT images were subjected to thresholding optimized for the

peripheral MT array (Fig. 2 Supplemental Fig. 2) and the directionality of MTs was determined in

respect to the cell border. Every pixel of the image was analyzed with inconclusive pixels

disregarded. Subsequently, MT directionality was quantified as a function of the distance from

the cell border and directionality of peripheral MTs within 1µm of the cell border quantified. Our

results indicate that in cells treated with non-targeting control shRNA (Fig. 2 F), the distribution

of MT angles in the cell periphery are vastly parallel and co-aligned with the cell boundary, as

previously reported for islet beta cells (Bracey et al. 2019). In contrast, the loss of KIF5B via

shRNA depletion resulted in a significant loss of parallel MTs at the periphery (Fig. 2 G, H). This

indicated that MT sliding by KIF5B acts to align MT at the cell periphery in addition to delivering

MTs to this cell location.

Combined, our data demonstrate a dramatic effect of KIF5B perturbation on both the distribution

of MTs to the cell periphery and their orientation along the cell boundary. These data suggest

that KIF5B-driven MT sliding is a decisive mechanism of the sub-membrane MT array

generation, likely via redistribution of centrally nucleated MTs and subsequent aligning them at

the cell edge. Thus, MT sliding is likely a critical component in functional MT organization in

beta cells.

Beta-cell kinesin-1 drives MT sliding through the C-terminal MT-binding domain

While membrane cargo transport by KIF5s requires association of the heavy chain with the

kinesin light chains (KLCs) and/or other adaptors, transportation of MTs as cargos occurs due to

direct binding of KIF5 to MTs through the ATP-independent MT binding domain in heavy chain

tail (C-terminus) (Jolly et al., 2010; Seeger & Rice, 2010).

To specifically establish the role of MT sliding by KIF5B in beta cells, we sought to evaluate the

effects of suppressing the binding of KIF5B tail to MTs. To this end we used a previously

generated construct (Ravindran, Engelke, Verhey, & Tsai, 2017), which is a motor-less version

of wild-type (WT) kinesin-1 motor KIF5B containing the cargo-binding and ATP independent MT

binding domain and tagged with mCherry (mCh) at the amino terminus (Fig 3A). When

overexpressed, this construct acts as a dominant-negative (DN) tool preventing association of

the tail of endogenous KIF5B with MTs. This tool is referred to as KIFDN^{wt} (KIF5B dominant

negative wild-type) moving forward (Fig. 3A).

To confirm that KIF5B tail domain binds to MTs in MIN6 and acts a dominant negative we co-

expressed KIFDN^{wt} and mEmerald-tubulin. When subjected to the FRAP assay we detected a

complete loss of MT sliding events as compared to a control (Fig. 3B-D). To prevent tail

engagement of the MT lattice through the ATP-independent binding domain we opted to make

point mutations in the tail domain to change the residues 892-DRKRYQ to 892-DAAAYA, thus

generating KIFDN^{MUT} (Fig. 3A). Photobleaching assay in cells co-expressing of the KIFDN^{MUT}

with mEmerald-tubulin indicated that the MT sliding activity was not blocked in the presence of

the mutated construct (Fig. 3B, E, Fig.3-Video 1), confirming that KIF5B tail domain binding to

MTs is needed for MT sliding in beta cells.

The dominant negative constructs are also tagged with the FK506-rapamycin-binding protein

(FKBP), as indicated in Fig. 3A. This allows to heterodimerize them with a motor domain fused

with the FKBP-rapamycin binding (FRB) domain using A/C Heterodimerizer (rapalog) and

reconstitute a functional motor (Inobe & Nukina, 2016). We restored kinesin-1 activity by

connecting the motor-less KIF5B, KIFDN^{wt}, to kinesin-1 motor domain as a way to rescue the

effects of DN approach of KIF5B tail overexpression. To this end, we co-expressed MIN6 cells

with the tail domain, mEmerald-tubulin, and the KIF5C motor domain fused to FRB domain (Fig.

3A). Once the tail and motor domain were dimerized with rapalog, we saw that the once blocked

MT sliding events of the KIFDN^{wt} tail alone were now reversed (Fig. 3B, F, Fig.3-Video 2). In

contrast, under conditions of heterodimerization of KIFDN^{MUT} with the motor, MT sliding was

greatly impaired (Fig. 3B, G, Fig.3-Video 2), indicating that the motor with mutated ATP-

independent binding domain cannot use MTs as cargos. Interestingly, endogenous motor in this

case was unable to efficiently transport MTs, suggesting that the endogenous motor pool

engaged in MT sliding was significantly smaller than the overexpressed non-functional motor.

Overall, the results of the DN approach confirm that MT sliding in beta cells is driven by KIF5B

through direct kinesin-1 tail binding to cargo MTs.

195   **Effects of C-terminal MT-binding of kinesin-1 on beta-cell MT organization**

196   Keeping in mind that KIF5B has additional major functions in addition to MT sliding, we sought

197   to test the consequence of MT sliding more directly by turning to overexpression of the DN

198   constructs. Thus, we took advantage of our heterodimerization approach to analyze MT patterns

199   in cells with active kinesin-1 which is able or unable to slide MTs (see Fig.3F vs Fig.3G). We

200   analyzed MIN6 cells that express either the KIFDN^{wt} or KIFDN^{mut} tail domains alone (Fig. 4

201   Supplemental Fig. 1) or co-expressing and heterodimerized with the motor domain (Fig. 4B,C).

202   Cells were fixed and immunostained for tubulin to identify the MT network. As expected,

203   overexpression of the KIFDN^{WT} tail construct alone acted as dominant negative toward MT

204   distribution to the cell periphery, resulting in decreased peripheral tubulin intensity (Fig. 4

205   Supplemental Fig. 1A,C), while in cells expressing KIFDN^{MUT} MT patterns were comparable to

206   control (Fig. 4 Supplemental Fig. 1B,C).

207   Interestingly, expression of heterodimerized kinesin motors led to impaired MT network

208   configurations compared to NT control (Fig. 4). Specifically, blocking of MT sliding by

209   overexpression KIFDN^{mut} heterodimerized with the motor, led to the decrease in peripheral

210   tubulin intensity (Fig. 4C,D) and impaired MT aligning along the cell border (Fig. 4E,G). These

211   data indicate that KIF5B-driven relocation of centrally nucleated MTs to beta cell periphery

212   requires kinesin tail domain binding to "cargo" MTs. Strikingly, overexpression of functional

213   heterodimerized motor, which was capable of MT sliding and populating of the cell periphery

214   with MTs as detected by tubulin intensity readings (KIFDN^{wt} heterodimerized with the motor, Fig.

215   4B,D), also led to a deficient MT aligning at the periphery (Fig. 4E,H). This can be interpreted as

216   a result of unregulated sliding in these experimental conditions, since excessive kinesin-1-

217   dependent sliding can lead to MT bending (Straube et al., 2006). This suggests that proper

218   organization of MTs within the sub-membrane array requires fine tuning of MT sliding activity.

219

Collectively, this indicates that regulated KIF5B activity is essential for redistributing MTs to the

cell border and sustaining an aligned peripheral MT array.

Exaggerated MT sliding leads to defects in peripheral array alignment

Our data discussed above suggest that overexpression of functional kinesin-1 disrupts MT

alignment at the cell periphery, inducing their bending and buckling (Fig. 4E, G). To test if this

defect is a result of excessive MT sliding, we employed a small molecule, kinesore, which is

known to dramatically promote MT sliding by kinesisn-1 (Randall et al., 2017). Kinesore targets

kinesin cargo adaptor function, by impairing KLC from binding kinesin heavy chain. As a result,

kinesin heavy chain will excessively engage MTs through the C-terminal, ATP independent MT

binding domain, leading to exaggerated MT sliding and the loss of membrane cargo transport by

kinesin-1 (Randall et al., 2017).To this end we pretreated MIN6 cells with 50µm kinesore and

stained for MTs (Fig. 5). MT networks in MIN6 cells became over corrected and looped (Fig. 5B)

as previously seen in other cell types (Randall et al., 2017). This exaggerated MT looping

resulting in a slight decrease of peripheral MT intensity (Fig. 5C). Further analysis of the

peripheral bundle indicated that MT alignment was strongly impaired upon kinesore-driven MT

remodeling as compared with vehicle (DMSO) treatment (Fig. 5D-G). The loss of coaligned MTs

and loss of tubulin density at the periphery indicate that MT sliding must be gated to prevent

over corrected MT networks.

MT sliding in beta cells is activated by glucose stimulation

It has previously been reported that kinesin-1 switches activity level in the presence of glucose

stimuli (Donelan et al., 2002). We predicted that as KIF5B activity modulates the MT sliding

events would also change depending on the glucose concentration. To test this, we pre-

incubated MIN6 cells with media containing a low concentration of 2.8mM glucose (Fig. 6A). We

applied the photobleaching assay at these conditions and detected little to no MT sliding events.

When switching glucose to a high concentration of 20mM, MT sliding and remodeling events

were significantly increased (Fig. 6B, Fig.6-Video 1). Quantification of the sliding events

demonstrated that MIN6 displaced MTs via MT sliding significantly more efficient upon glucose

stimulation (Fig. 6C). We then turned to single molecule tracking of MT lattice fiducial marks

(K560Rigor^{E236A}-SunTag) to further investigate this observation. Consistent with the

photobleaching assay, the fiducial marks were predominantly stationary in cells pre-incubated in

2.8mM glucose (Fig. 6D) but frequently underwent directed relocation events indicative of MT

sliding in cells after stimulated with 20mM glucose (Fig. 6E, Fig.6-Video 2).

These data demonstrate that glucose-stimulated remodeling of the MT network involves

regulated MT sliding. Given the importance of MT sliding for peripheral MT organization (Figs. 2,

4), this effect may be essential to restore peripheral MT array after glucose-dependent

destabilization or regulate other aspects of MT-dependent tuning of GSIS.

Discussion

Since the first description of convoluted MT network in MIN6 cells by the Rutter group (Varadi et

al., 2003), our views on regulation, function, and dynamics of pancreatic beta cell MT network

have been gradually evolving (Bracey et al., 2022). However, the field is still far from the

understanding of the mechanisms underlying the network architecture. Here, we show that MT

sliding is a prominent phenomenon in beta cells, and that it is driven by kinesin KIF5B. This

kinesin-1-dependent MT sliding is a critical mechanism needed to formation and a long-term

maintenance of beta cell MT network, especially the peripheral MT arrays, and that glucose

stimulation facilitates MT sliding activity. Overall, our study establishes MT sliding as an

essential regulator of beta cell architecture and function.

Our data indicate that MT sliding is activated on a short-term basis after stimulation. It is

plausible to suggest that this is needed to replace MTs at the cell periphery that are destabilized

in high glucose after MT-stabilizing protein tau is phosphorylated and detached from the sub-

membrane MTs (Ho et al., 2020). However, the amount of MT polymer on every glucose

stimulation changes only slightly, often undetectable (Muller et al., 2021; Zhu et al., 2015). In

fact, we observe a prominent effect of peripheral MT loss only after a long-term kinesin

depletion (three-four days). This is consistent with our observation that only a minor subset of

MT is being moved at every stimulation. We assume that the loss of peripheral MT array in

KIF5B-depleted cells in a manifestation of accumulated lack of sliding over an extended period.

We also found that increasing MT sliding will also not yield a properly configured MT array:

kinesore-treated cells lack aligned peripheral MTs. This indicates that, similar to other part of

beta cell physiology, the dose of MT sliding has to be precisely tuned to achieve physiologically

relevant architecture. It was shown before that exaggerated kinesin-dependent MT sliding

causes MT bundling and buckling into aberrant configuration (Straube et al., 2006). We predict

that a fine-tuning regulatory pathway must exist to restrict the number of MT sliding events to

the cell needs.

Interestingly, blocking kinesin results in a striking accumulation of MT in the cell center where

they are normally nucleated at MTOCs, which include the centrosome and the Golgi, in

differentiated beta cells the latter being the main MTOC. Thus, sliding MTs originate from the

MTOC area. At the same time, FIB-SEM analysis did not detect many MTs associated with

MTOCs in physiologically normal beta cells (Muller et al., 2021). This implicates that MTs are

normally rapidly dissociated from MTOCs so that they become available for transport by sliding.

It is worth mentioning that for long-distance transport by sliding, cargo MTs must be short,

otherwise MT buckling and not long-distance transport will occur (Straube et al., 2006).

Interestingly, Mueller shorter MTs have been observed in high glucose conditions (Muller et al.,

2021), when MT are nucleated more actively (Trogden et al., 2019) and transported more

frequently (this paper). Possibly, nucleated MTs are detached from MTOCs before they achieve

a length that would prevent their transport. There is a possibility suggested that MTs are being

severed by katanin in high glucose (Muller et al., 2021), which would generate MT fragments

that can serve as cargos more easily. It is also possible that sliding MT subpopulation has

some additional specific features that make them preferred cargos, since it is becoming

increasingly clearer in the field that there is immense heterogeneity among MTs. Post-

translational modifications and MT associated proteins, which vastly alter stability and

coordination of motor proteins (Hammond, Cai, & Verhey, 2008; McKenney, Huynh, Vale, &

Sirajuddin, 2016; Monroy et al., 2018; Yu, Garnham, & Roll-Mecak, 2015), might also influence

which MTs serve as cargos versus transportation tracks in beta cells.

On a final note, it is important to evaluate the phenomenon reported here in light of the dual role

of KIF5B as IG transporter and MT transporter and the coordination of those two roles in IG

transport and availability for secretion. Our results indicate that KIF5B is needed for the

patterning of peripheral MTs which we have shown to restrict secretion (Bracey et al., 2020; Ho

et al., 2020). At the same time, it is well established that KIF5B transports IGs (Varadi et al.,

2002) and KIF5B loss of function impairs insulin secretion (Cui et al., 2011). After a prolonged

KIF5B inactivation, a loss of peripheral readily-releasable IG should be expected due to two

factors: because there is no MT bundle to prevent over-secretion and IG depletion, and because

there is no new IGs being transported from the Golgi area. In contrast, physiological activation

of kinesin by glucose (Donelan et al., 2002; Varadi et al., 2003) would both promote

replenishment of IG through non-directional transport through the cytoplasm and restoration of

peripheral MT array to prevent over-secretion on each stimulus.

In conclusion, here we add another very important cell type to the list of systems that employ

KIF5-dependent MT sliding to build functional MT networks. This system is unique because in

this case MT sliding is metabolically regulated and activated on a single-minute time scale by

nutrition triggers.

Materials and Methods

1. Key reagents

Reagent type or resource	Designation	Source or reference	Additional information
Cell line	MIN6		RRID:CVCL_0431
Chemical compound, drug	Kinesore	Tocris, Cat#: 6664	Final concentration (50µm)
Chemical compound, drug	A/C Heterodimerizing Drug	Takara, Cat#: 635056	Final concentration (25µm)
Antibody	Anti-KIF5B antibody	Abcam, Cat#: Ab167429	(1:500 dilution)
Antibody	Anti-alpha-Tubulin antibody	Sigma-Aldrich, Cat#: T9026	(1:500 dilution)
Halo Dye-585, 647	HaloTag® Ligands	CS315105, GA1120	2ul/mL

2. Cell Lines

MIN6 cells between passage 40-60 were utilized (Ishihara et al., 1993; Miyazaki et al., 1990).

Cells were maintained in 25 mM glucose Dulbecco's modified eagle medium (DMEM) (Life

Technologies, Frederick, MD) supplemented with 10% fetal bovine serum (FBS), 0.001% β-

mercaptoethanol, 0.1 mg/ml penicillin, and 0.1 mg/ml streptomycin in 5% CO_2 at 37 degrees C.

3. Reagents and antibodies

Primary antibodies for immunofluorescence were: mouse anti-β-tubulin (Sigma-Aldrich, 1:1000),

rabbit anti-β-tubulin (Sigma-Aldrich, 1:1000) and rabbit anti-KIF5B (Abcam), Alexa488-,

Alexa568-, and Alexa647-conjugated highly cross-absorbed secondary antibodies (Invitrogen).

Coverslips were mounted in Vectashield Mounting Medium (Vector Labaratories). Cells were

treated with indicated drugs for three hours unless otherwise indicated. Drugs used were:

Kinesore (Tocris Bioscience).

4. shRNA sequence

The KIF5B-targeting shRNA [shRNA KIF5B] #1, [TL510740B, 5'-

ACTCTACGGAACACTATTCAGTGGCTGGA] and [shRNA KIF5B] #2, [TL51074CB 5' –

AGACCGTAAACGCTATCAGCAAGAAGTAG] are in the plasmid backbone pGFP-C-shLenti

and were from Origene (Rockville, MD). The non-targeting shRNA control, was pGFP-C-shLenti

also from Origene.

5. DNA Constructs

Plasmid Construct	Source	Catalog #
SCR-TGFP	Origene	Custom
KIF5B shRNA#1-TGFP	Origene	Custom
KIF5B shRNA#2-TGFP	Origene	Custom
Scr shRNA mEmerald-Tubulin	This paper	Custom
KIF5B shRNA#1-mEmerald-tubulin	This paper	Custom
KIF5B shRNA#2-mEmerald-tubulin	This paper	Custom
mEmerald-Tubulin-C-18	Addgene	#54292
pcDNA4TO-K560-E236A-24xGCN4	Addgene	#60909
ScFv-GCN4-HaloTag-GB1-NLS	Addgene	#106303
FKBP-mCherry-KIF5B(568-964)	Kristen Verhey	((Ravindran et al., 2017))
p205ME_RnKIF5C(1-559)-TagBFP-FRB	Kristen Verhey	((Ravindran et al., 2017))
FKBP-mCherry-KIF5B(568-964)-AAAYA (MUT)	This paper	

6. Cloning

The Scr shRNA mEmerald-Tubulin, KIF5B shRNA#1-mEmerald-tubulin, KIF5B shRNA#2-

mEmerald-tubulin were all generated from their respective TGFP containing constructs. Using

the NotI and PmeI sites the TGFP was swapped for mEmerald-Tubulin.

The FKBP-mCherry-KIF5B(568-964) construct (gift from Kristen Verhey, University of

Michigan), has previously been previously described (Ravindran et al., 2017).

By using site directed mutagenesis, we made 8 point mutations in the tail domain to change

residues RKRYQ to AAAYA in the ATP independent MT binding domain. The point mutations

were sufficient to rescue MT sliding in the cell. As previously published, this disrupts the tail

domain to bind to the acidic e hook of the MT tail. Point mutations were introduced using a site

directed mutagenesis kit, In-Fusion® Snap Assembly (Takara).

7. Lentiviral Transduction and Transfection

Lentivirus production and infection followed standard methods (Huang et al., 2018). MIN6 were

treated with a given shRNA expressing a mEmerald-tubulin/cytosolic tgfp marker for 96hrs prior

to imaging to achieve KD efficiency. For non-viral vectors, MIN6 cells were transfected using

Amaxa Nucleofection (Lonza).

8. Western blotting

Cell lysates from MIN6 cells were lysed using 1% CHAPs buffer on ice for 5 minutes. For KIF5B

knockdown, cells were first sorted for GFP expression after 72 h. Protein lysate (20µg) was

loaded onto an 8% SDS-PAGE gel under reducing conditions and transferred to nitrocellulose

membranes. Membranes were probed with antibodies against KIF5B, and α-tubulin. The

membranes were then blocked with 5% nonfat dried milk (Sigma-Aldrich) for 1 h and incubated

overnight with primary antibodies: rabbit anti- KIF5B (Abcam), and mouse anti-GAPDH (Santa

Cruz). anti-rabbit HRP and anti-mouse HRP were used as secondary antibodies and imaged on

an ChemiDoc (Bio-Rad).

9. Image Acquisition

Immunofluorescence microscopy of fixed samples

Fixed samples were imaged using a laser scanning confocal microscope Nikon A1r based on a

TiE Motorized Inverted Microscope using a 100X lens, NA 1.49, run by NIS Elements C

software. Cells were imaged in 0.05µm slices through the whole cell.

Live cell imaging

Cells were cultured on 4-chamber MatTek dishes coated with 10 µg/µl fibronectin and

transduced 96hrs or transfected 48 h before experiment. For live-cell imaging of MT sliding,

cells were transfected with Emerald-Tubulin and imaged using a Nikon TiE inverted microscope

equipped with 488- and 568-nm lasers, a Yokogawa CSU-X1 spinning disk head, a PLAN APO

VC 100x NA1.4 oil lens, intermediate magnification 1.5X, and CMOS camera (Photometrics

Prime 95B), 405 Burker mini-scanner, all controlled by Nikon Elements software.

Photobleaching Assay

~1×10^{6} MIN6 cells were transfected with 1µg of mEmerald-tubulin or transduced with lentiviral

KIF5B shRNA with mEmerald-tubulin as a reporter and attached to glass dishes coated with

fibronectin for up to 96hrs. On the SDC microscope, the ROI tool in NIKON elements was used

to place two ROI's ~5μm apart at either end of the cell. These regions were assigned to be

photobleached with the equipped 405nm mini scanner laser leaving a fluorescent patch over the

middle which we termed the "fluorescent belt". After the regions were photobleached cells were

then acquired for 5mins, across 7 optical slices (0.4μm step size) in 10 second interval between

frames.

Sun Tag Rigor Kinesin and Tracking of MT sliding

SunTag system for MT lattice fiducial marks was adapted from (Lu, Winding, Lakonishok,

Wildonger, & Gelfand, 2016). ~$1 \times 10^{\wedge 6}$ MIN6 cells were co transfected with 1μg of the ScFv-

GCN4-HaloTag-GB1-NLS, and 0.5μg of the pcDNA4TO-K560-E236A-24xGCN4 plasmid

(K560Rigor^{E236A}-SunTag (Tanenbaum et al., 2014)). After 24hrs the cells were washed with 1x

PBS and the media replaced with KRB containing 2.8mM glucose for 1 hour, following a second

incubation with HALO dye of choice (Promega) for 30mins. Cells were imaged in 1 focal plane

for 2 mins with 100ms exposure time and no delay in acquisition. The acquired image was

processed through Imaris Microscopy Image Analysis Software (Oxford Instruments), where the

fiducial marks were tracked.

Sun Tag Rigor Kinesin Tracking Analysis

We sought to normalize the behaviors by comparing the MT movements over 5 second intervals

and calculated the displacement of a given fiducial mark. In total over 60,000 tracks were

detected, and the segmented displacement of ~25,000 of those tracks were calculated. The 5s

displacement was binned at 0.05um intervals and the % of distribution for each bin was

calculated for each cell and summarized in histograms (Fig 1, Fig 6).

10. Image Processing, and Analysis

Image processing

Figure 1: MaxIP (D-F), Single focal plane (G-I)

Figure 2: Cells shown are projections of the bottom 1µm of the cell. The tubulin channel was separated processed as gray value and inverted. The KIF5B channel was pseudo colored cyan and adjusted equally across all conditions (A-C) Single focal plane images under the nucleus of the cell were selected for analysis (F-H)

Figure 3: MaxIP of the tubulin channel for each time point (C-G)

Figure 4: Cells shown are projections of the bottom 1µm of the cell. The tubulin channel was separated processed as gray value and inverted (A-C).

Single focal plane images under the nucleus of the cell were selected for analysis (F-H) (see supplement Figure 2-1)

Figure 5: Cells shown are projections of the bottom 1µm of the cell. The tubulin channel was separated processed as gray value and inverted (A-B). Single focal plane images under the nucleus of the cell were selected for analysis (D-E)

Figure 6: MaxIP of the tubulin channel for each time point (A-B) Single plane (D-E)

MATLAB Script: MT Directionality

Oversampled images were deconvolved using the Richardson and Lucy Deconvolution

algorithm. Images were masked and threshold (IsoData) in ImageJ. The MT directionality script

was applied in MATLAB. Only the outer 1µm of MTs were taken for binning and quantification

purposes.

MATLAB Script: Msdanalyzer, Segmentation

The position of all tracked fiducial spots were exported from Imaris to excel. The MSDanalyzer

was developed by Nadine Tarantino et al, and adapted by Kai Bracey, Pi'Illani Noguchi, and

Alisa Cario (Vanderbilt University) to normalize the tracks in time. Tracks were segmented into

displacements over 5s and binned as shown in the results.

11. Statistics and reproducibility

For all experiments, *n* per group is as indicated by the figure legend and the scatter dot plots

indicate the mean of each group and error bars indicate the standard error of the mean. All

graphs and statistical analyses were generated using Excel (Microsoft) and Prism software

(Graphpad). Statistical significance for all *in vitro* and *in vivo* assays was analyzed using an

unpaired t-test, one-way ANOVA with Sidak's multiple comparisons test, Kolmogorov-Smirnov

test as indicated in the figure legends. For each analysis $p < 0.05$ was considered statistically

significant, and $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$.

Author contributions

KMB performed most of the experiments and a large part of data analysis and wrote the

manuscript. PN and AC provided scripts for data analysis and analyzed data. CE performed

Western Blotting experiments. GG provided conceptual insight and molecular cloning strategy.

I.K. supervised the study, provided conceptual insight and wrote the manuscript.

Acknowledgements

This work was supported by National Institutes of Health (NIH) grants F31DK122650 T32 (to

KMB), R35-GM127098 (to I.K.), R01-DK106228 (to I.K. and G.G.), R01-DK65949 (to G. G.),

R01-DK125696 (to G.G.). KMB was supported by an NIH training grant R25-GM062459

"Initiative for Maximize Student Diversity" (Sealy, PI), AC by an NIH training grant T32-

CA119925 "Integrated Biological Systems Training in Oncology" (Tansey, PI), CME was

supported by NIGMS of the NIH under award number T32GM007347, and PN by an NIH

training grant T32 DK101003 "Integrated Training in Engineering and Diabetes" (Young, PI). We

thank Hamida Ahmed for technical help. The authors also thank Margret Fye for critical reading

of the manuscript.

References

Acar, S., Carlson, D. B., Budamagunta, M. S., Yarov-Yarovoy, V., Correia, J. J., Ninonuevo, M. R., . . . Scholey, J. M. (2013). The bipolar assembly domain of the mitotic motor kinesin-5. *Nat Commun, 4*, 1343. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/23299893. doi:10.1038/ncomms2348

Barlan, K., Lu, W., & Gelfand, V. I. (2013). The microtubule-binding protein ensconsin is an essential cofactor of kinesin-1. *Curr Biol, 23*(4), 317-322. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/23394833. doi:10.1016/j.cub.2013.01.008

Bogan, J. S. (2021). Granular detail of beta cell structures for insulin secretion. *J Cell Biol, 220*(2). Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/33427875. doi:10.1083/jcb.202012082

Bracey, K. M., Gu, G., & Kaverina, I. (2022). Microtubules in Pancreatic beta Cells: Convoluted Roadways Toward Precision. *Front Cell Dev Biol, 10*, 915206. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/35874834. doi:10.3389/fcell.2022.915206

Bracey, K. M., Ho, K. H., Yampolsky, D., Gu, G., Kaverina, I., & Holmes, W. R. (2020). Microtubules Regulate Localization and Availability of Insulin Granules in Pancreatic Beta Cells. *Biophys J, 118*(1), 193-206. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/31839261. doi:10.1016/j.bpj.2019.10.031

Cao, Y., Lipka, J., Stucchi, R., Burute, M., Pan, X., Portegies, S., . . . Hoogenraad, C. C. (2020). Microtubule Minus-End Binding Protein CAMSAP2 and Kinesin-14 Motor KIFC3 Control Dendritic Microtubule Organization. *Curr Biol, 30*(5), 899-908 e896. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/32084403. doi:10.1016/j.cub.2019.12.056

Cui, J., Wang, Z., Cheng, Q., Lin, R., Zhang, X. M., Leung, P. S., . . . Huang, J. D. (2011). Targeted inactivation of kinesin-1 in pancreatic beta-cells in vivo leads to insulin secretory deficiency. *Diabetes, 60*(1), 320-330. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/20870970. doi:10.2337/db09-1078

Desai, A., & Mitchison, T. J. (1997). Microtubule polymerization dynamics. *Annu Rev Cell Dev Biol, 13*, 83-117. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/9442869. doi:10.1146/annurev.cellbio.13.1.83

Donelan, M. J., Morfini, G., Julyan, R., Sommers, S., Hays, L., Kajio, H., . . . Rhodes, C. J. (2002). Ca2+-dependent dephosphorylation of kinesin heavy chain on beta-granules in pancreatic beta-cells. Implications for regulated beta-granule transport and insulin exocytosis. *J Biol Chem, 277*(27), 24232-24242. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/11978799. doi:10.1074/jbc.M203345200

Grabham, P. W., Seale, G. E., Bennecib, M., Goldberg, D. J., & Vallee, R. B. (2007). Cytoplasmic dynein and LIS1 are required for microtubule advance during growth cone remodeling and fast axonal outgrowth. *J Neurosci, 27*(21), 5823-5834. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/17522326. doi:10.1523/JNEUROSCI.1135-07.2007

Hammond, J. W., Cai, D., & Verhey, K. J. (2008). Tubulin modifications and their cellular functions. *Curr Opin Cell Biol, 20*(1), 71-76. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/18226514. doi:10.1016/j.ceb.2007.11.010

He, Y., Francis, F., Myers, K. A., Yu, W., Black, M. M., & Baas, P. W. (2005). Role of cytoplasmic
dynein in the axonal transport of microtubules and neurofilaments. *J Cell Biol, 168*(5),
697-703. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/15728192.
doi:10.1083/jcb.200407191
Heaslip, A. T., Nelson, S. R., Lombardo, A. T., Beck Previs, S., Armstrong, J., & Warshaw, D. M.
(2014). Cytoskeletal dependence of insulin granule movement dynamics in INS-1 beta-
cells in response to glucose. *PLoS One, 9*(10), e109082. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/25310693. doi:10.1371/journal.pone.0109082
Ho, K. H., Yang, X., Osipovich, A. B., Cabrera, O., Hayashi, M. L., Magnuson, M. A., . . . Kaverina,
I. (2020). Glucose Regulates Microtubule Disassembly and the Dose of Insulin Secretion
via Tau Phosphorylation. *Diabetes, 69*(9), 1936-1947. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/32540877. doi:10.2337/db19-1186
Hoboth, P., Muller, A., Ivanova, A., Mziaut, H., Dehghany, J., Sonmez, A., . . . Solimena, M.
(2015). Aged insulin granules display reduced microtubule-dependent mobility and are
disposed within actin-positive multigranular bodies. *Proc Natl Acad Sci U S A, 112*(7),
E667-676. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/25646459.
doi:10.1073/pnas.1409542112
Hu, R., Zhu, X., Yuan, M., Ho, K. H., Kaverina, I., & Gu, G. (2021). Microtubules and Galphao-
signaling modulate the preferential secretion of young insulin secretory granules in islet
beta cells via independent pathways. *PLoS One, 16*(7), e0241939. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/34292976. doi:10.1371/journal.pone.0241939
Huang, C., Walker, E. M., Dadi, P. K., Hu, R., Xu, Y., Zhang, W., . . . Gu, G. (2018). Synaptotagmin
4 Regulates Pancreatic beta Cell Maturation by Modulating the Ca(2+) Sensitivity of
Insulin Secretion Vesicles. *Dev Cell, 45*(3), 347-361 e345. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/29656931. doi:10.1016/j.devcel.2018.03.013
Inobe, T., & Nukina, N. (2016). Rapamycin-induced oligomer formation system of FRB-FKBP
fusion proteins. *J Biosci Bioeng, 122*(1), 40-46. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/26777239. doi:10.1016/j.jbiosc.2015.12.004
Ishihara, H., Asano, T., Tsukuda, K., Katagiri, H., Inukai, K., Anai, M., . . . Oka, Y. (1993).
Pancreatic beta cell line MIN6 exhibits characteristics of glucose metabolism and
glucose-stimulated insulin secretion similar to those of normal islets. *Diabetologia,*
36(11), 1139-1145. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/8270128.
doi:10.1007/BF00401058
Jolly, A. L., Kim, H., Srinivasan, D., Lakonishok, M., Larson, A. G., & Gelfand, V. I. (2010). Kinesin-
1 heavy chain mediates microtubule sliding to drive changes in cell shape. *Proc Natl*
Acad Sci U S A, 107(27), 12151-12156. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/20566873. doi:10.1073/pnas.1004736107
Kurasawa, Y., Earnshaw, W. C., Mochizuki, Y., Dohmae, N., & Todokoro, K. (2004). Essential
roles of KIF4 and its binding partner PRC1 in organized central spindle midzone
formation. *EMBO J, 23*(16), 3237-3248. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/15297875. doi:10.1038/sj.emboj.7600347
Lu, W., Fox, P., Lakonishok, M., Davidson, M. W., & Gelfand, V. I. (2013). Initial neurite
outgrowth in Drosophila neurons is driven by kinesin-powered microtubule sliding. *Curr*

Biol, 23(11), 1018-1023. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/23707427. doi:10.1016/j.cub.2013.04.050

Lu, W., & Gelfand, V. I. (2017). Moonlighting Motors: Kinesin, Dynein, and Cell Polarity. *Trends Cell Biol, 27*(7), 505-514. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/28284467. doi:10.1016/j.tcb.2017.02.005

Lu, W., Winding, M., Lakonishok, M., Wildonger, J., & Gelfand, V. I. (2016). Microtubule-microtubule sliding by kinesin-1 is essential for normal cytoplasmic streaming in Drosophila oocytes. *Proc Natl Acad Sci U S A, 113*(34), E4995-5004. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/27512034. doi:10.1073/pnas.1522424113

McKenney, R. J., Huynh, W., Vale, R. D., & Sirajuddin, M. (2016). Tyrosination of alpha-tubulin controls the initiation of processive dynein-dynactin motility. *EMBO J, 35*(11), 1175-1185. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/26968983.
doi:10.15252/embj.201593071

Miyazaki, J., Araki, K., Yamato, E., Ikegami, H., Asano, T., Shibasaki, Y., . . . Yamamura, K. (1990). Establishment of a pancreatic beta cell line that retains glucose-inducible insulin secretion: special reference to expression of glucose transporter isoforms. *Endocrinology, 127*(1), 126-132. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/2163307. doi:10.1210/endo-127-1-126

Monroy, B. Y., Sawyer, D. L., Ackermann, B. E., Borden, M. M., Tan, T. C., & Ori-McKenney, K. M. (2018). Competition between microtubule-associated proteins directs motor transport. *Nat Commun, 9*(1), 1487. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/29662074. doi:10.1038/s41467-018-03909-2

Muller, A., Schmidt, D., Xu, C. S., Pang, S., D'Costa, J. V., Kretschmar, S., . . . Solimena, M. (2021). 3D FIB-SEM reconstruction of microtubule-organelle interaction in whole primary mouse beta cells. *J Cell Biol, 220*(2). Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/33326005. doi:10.1083/jcb.202010039

Randall, T. S., Yip, Y. Y., Wallock-Richards, D. J., Pfisterer, K., Sanger, A., Ficek, W., . . . Dodding, M. P. (2017). A small-molecule activator of kinesin-1 drives remodeling of the microtubule network. *Proc Natl Acad Sci U S A, 114*(52), 13738-13743. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/29229862. doi:10.1073/pnas.1715115115

Ravindran, M. S., Engelke, M. F., Verhey, K. J., & Tsai, B. (2017). Exploiting the kinesin-1 molecular motor to generate a virus membrane penetration site. *Nat Commun, 8*, 15496. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/28537258.
doi:10.1038/ncomms15496

Rice, S., Lin, A. W., Safer, D., Hart, C. L., Naber, N., Carragher, B. O., . . . Vale, R. D. (1999). A structural change in the kinesin motor protein that drives motility. *Nature, 402*(6763), 778-784. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/10617199.
doi:10.1038/45483

Sanavia, T., Huang, C., Manduchi, E., Xu, Y., Dadi, P. K., Potter, L. A., . . . Gu, G. (2021). Temporal Transcriptome Analysis Reveals Dynamic Gene Expression Patterns Driving beta-Cell Maturation. *Front Cell Dev Biol, 9*, 648791. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/34017831. doi:10.3389/fcell.2021.648791

Seeger, M. A., & Rice, S. E. (2010). Microtubule-associated protein-like binding of the kinesin-1
tail to microtubules. *J Biol Chem, 285*(11), 8155-8162. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/20071331. doi:10.1074/jbc.M109.068247
Straube, A., Hause, G., Fink, G., & Steinberg, G. (2006). Conventional kinesin mediates
microtubule-microtubule interactions in vivo. *Mol Biol Cell, 17*(2), 907-916. Retrieved
from https://www.ncbi.nlm.nih.gov/pubmed/16339079. doi:10.1091/mbc.e05-06-0542
Tabei, S. M. A., Burov, S., Kim, H. Y., Kuznetsov, A., Huynh, T., Jureller, J., . . . Scherer, N. F.
(2013). Intracellular transport of insulin granules is a subordinated random walk.
Proceedings of the National Academy of Sciences of the United States of America,
110(13), 4911-4916. Retrieved from <Go to ISI>://WOS:000318031900023.
doi:10.1073/pnas.1221962110
Tanenbaum, M. E., Gilbert, L. A., Qi, L. S., Weissman, J. S., & Vale, R. D. (2014). A protein-tagging
system for signal amplification in gene expression and fluorescence imaging. *Cell,*
159(3), 635-646. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/25307933.
doi:10.1016/j.cell.2014.09.039
Trogden, K. P., Zhu, X., Lee, J. S., Wright, C. V. E., Gu, G., & Kaverina, I. (2019). Regulation of
Glucose-Dependent Golgi-Derived Microtubules by cAMP/EPAC2 Promotes Secretory
Vesicle Biogenesis in Pancreatic beta Cells. *Curr Biol, 29*(14), 2339-2350 e2335.
Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/31303487.
doi:10.1016/j.cub.2019.06.032
Varadi, A., Ainscow, E. K., Allan, V. J., & Rutter, G. A. (2002). Involvement of conventional
kinesin in glucose-stimulated secretory granule movements and exocytosis in clonal
pancreatic beta-cells. *J Cell Sci, 115*(Pt 21), 4177-4189. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/12356920

https://jcs.biologists.org/content/joces/115/21/4177.full.pdf. doi:10.1242/jcs.00083
Varadi, A., Tsuboi, T., Johnson-Cadwell, L. I., Allan, V. J., & Rutter, G. A. (2003). Kinesin I and
cytoplasmic dynein orchestrate glucose-stimulated insulin-containing vesicle
movements in clonal MIN6 beta-cells. *Biochem Biophys Res Commun, 311*(2), 272-282.
Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/14592410.
doi:10.1016/j.bbrc.2003.09.208
Vukusic, K., Ponjavic, I., Buda, R., Risteski, P., & Tolic, I. M. (2021). Microtubule-sliding modules
based on kinesins EG5 and PRC1-dependent KIF4A drive human spindle elongation. *Dev*
Cell, 56(9), 1253-1267 e1210. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/33910056. doi:10.1016/j.devcel.2021.04.005
Yu, I., Garnham, C. P., & Roll-Mecak, A. (2015). Writing and Reading the Tubulin Code. *J Biol*
Chem, 290(28), 17163-17172. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/25957412. doi:10.1074/jbc.R115.637447
Zhu, X., Hu, R., Brissova, M., Stein, R. W., Powers, A. C., Gu, G., & Kaverina, I. (2015).
Microtubules Negatively Regulate Insulin Secretion in Pancreatic beta Cells. *Dev Cell,*
34(6), 656-668. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/26418295.
doi:10.1016/j.devcel.2015.08.020

Figures

Figure 1. MTs in pancreatic beta cells undergo extensive sliding driven by kinesin

KIF5B.

A subset of RNA-sequencing data from primary mouse beta cells showing highly

expressed kinesins as indicated by mRNA counts. KIF5B (most-right bar, red data

points) is the most abundant kinesin motor in this cell type. N=3. Note that this is a

subset of the RNA sequencing sets published in (Sanavia et al., 2021). (**B**) Efficient

depletion of KIF5B in MIN6 cells using two alternative shRNA sequences, as compared

to a scrambled shRNA control. GAPDH, loading control. Relevant ROIs from a Western

blotting is shown. A representative example out of 3 repeats. (**C**) Quantification of MT

sliding FRAP assay in cells treated with scrambled control or one of the two KIF5B-

specific shRNAs (see representative data in D-F). MT displacement is shown as area of

MTs displaced into the bleached area after 5 minutes of recovery. One-way ANOVA test

was performed for statistical significance (p-value <0.0001). N=9-20 cells per set. (**D-F**)

Frames from representative FRAP live-cell imaging sequences. mEmerald-tubulin-

expressing MIN6 cells. Inverted grayscale images of maximum intensity projections

over 1 µm-thick stacks by spinning disk confocal microscopy. (D1-F1) The last frame

prior to photobleaching. (D2-F2) The first frame after photobleaching. (D3-F3) A frame 5

minutes (300 seconds) after photobleaching. Light-blue dotted lines indicate the edges

of the photobleached areas. Red arrows indicate MTs displaced into the bleached area.

Scale bars, 5 µm. (**G-I**) MIN6 cells featuring fiducial marks at MTs due to co-expression

of SunTag-KIF5B-560Rigor construct and Halo-SunTag ligand. Representative

examples for scrambled control shRNA-treated cell (G), KIF5B shRNA #1-treated cell

(H) and KIF5B shRNA #2-treated cell (I) are shown. Single-slice spinning disk confocal

microscopy. Halo-tag signal is shown as inverted gray-scale image. Top panels show

cell overviews (scale bars 5µm). Below, boxed insets (scale bars 2 µm) are enlarged to

show dynamics of fiducial marks (color arrows) at 1 second intervals (1-5 seconds). 0-

to 5-second tracks of fiducial mark movement are shown in the bottom panel, each track

color-coded corresponding to the arrows in the image sequences. (**J**) Summarized

quantification of stationary fraction of fiducial marks along MT lattice. Scrambled shRNA

control N=1,421 tracks across 6 cells, shRNA#1 N=852 tracks across 5 cells, shRNA#2

N=679 tracks across 5 cells. P One-way ANOVA, p<0.001 (**K**) Summarized

quantification of motile fraction of fiducial marks along the MT lattice. Scrambled shRNA

control N=2,066 tracks across 6 cells, shRNA#1 N=390 tracks across 5 cells, shRNA#2

N=307 tracks across 5 cells. P One-way ANOVA, p<0.001

Figure 1 – Supplemental Figure 1.

(A) Full western blot of anti-kif5b from Fig. 1B. (**B**) Histogram of displacements over 5-second intervals for all fiducial marks in scrambled shRNA control vs KIF5B shRNA #1 and KIF5B shRNA #2.

Figure 2. Microtubule abundance and alignment at the cell periphery depend on KIF5B.

(**A-C**) MT organization in MIN6 cells expressing scrambled control shRNA (A), KIF5B-

targeting shRNA #1 (B), or KIF5B-targeting shRNA #2 (C). Top, immunofluorescence

staining for tubulin (grayscale, inverted). Bottom, immunofluorescence staining for

KIF5B (cyan). Laser scanning confocal microscopy maximum intensity projection of

1µm at the ventral side of the cell. N=12. Scale bars: 5um. (**D**) Quantification of mean

tubulin intensity within the outer 2µm peripheral area of a cell, in data represented in (A-

C). Mean values, black bars. One-way ANOVA, $p<0.0001$. N=5-9 cells. (**E**) Histograms

of MT directionality within 1um of cell boundary using perfected thresholds (see

Supplemental figure 2-1 for the analysis workflow) in cells treated with scrambled

control versus KIF5B-targeting shRNA. Data are shown for the summarized detectable

tubulin-positive pixels in the analyzed shRNA-treated cell population, as represented in

(F-H). Unpaired t-test were performed across each bin for all cells, and a K-S test was

performed on the overall distribution. The share of MTs parallel to the edge (bin 0-10) is

significantly higher in control as compared to KIF5B depletions. Pixel numbers in the

analysis: SCR N=106,741 pixels across 9 cells, shRNA #1 N=71,243 across 7 cells,

shRNA #2 N= 54,101 across 7 cells. (**F-H**) Representative examples of MT

directionality analysis quantified in (E). (**F**) Scrambled control shRNA-treated cell. (**G**)

KIF5B shRNA #1-treated cell. (**H**) KIF5B shRNA#1-treated cell. Overviews of cellular

MT networks are shown as threshold to detect individual peripheral MTs (see

Supplemental figure 2-1 panel A5). (**F1-H2**) Directionality analysis outputs of regions

from yellow boxes in (F-H) are shown color-coded for the angles between MTs and the

nearest cell border. (**I**) Color code for (F1-H2): MTs parallel to the cell edge, blue; MTs

perpendicular to the cell edge, red. Figure 2- Source Data 1

Figure 2 – Supplemental Figure 1. Workflow of MT directionality analysis.

(**A**) Representation of the analysis workflow using a control DMSO-treated cell
immunostained for tubulin as an example. (A1) An image of the original inverted
grayscale confocal slice. (A2) A deconvolved image. (A3) Mask of the cell boundary.
(A4) An image within the mask after application of standard % threshold. (A5) An image
within the mask after application of a threshold optimizing detection of peripheral MTs
for a particular cell. (A3-A5) are derivatives of (A2). (A6) A schematic illustrating map of
distances from the nearest cell (mask) border per pixel (not to scale). (A7) A schematic

733   illustrating map of angles per pixel (not to scale). A6 and A7 can be produced from A4

734   or A5. (A8) Color-coded output map of MT directionalities. A8 is a derivative of A7.

Figure 2-Supplemental Figure 2. Illustration of thresholding variations and their

influence on the output analysis.

(**A**) A KIF5B-depleted cell as an example of contrast/threshold optimization for different

MT densities and corresponding outputs. (A1) An deconvolved inverted grayscale

confocal slice image contrasted to highlight peripheral MTs. (A2) The same

deconvolved inverted grayscale confocal slice image contrasted to highlight central

MTs. (A3) The image from A1 after application of a threshold optimized to highlight

peripheral MTs. (A4) The image from A1 after application of a standard % threshold.

(A5) Analysis outcome from (A3). (A6) Analysis outcome from (A4). (**B**) Histograms of

MT directionality within 1um of cell boundary using standard thresholds (see

Supplemental figure 3/1 for the analysis workflow) in cells treated with scrambled

control versus KIF5B-targeting shRNA. Data are shown for the summarized detectable

tubulin-positive pixels in the analyzed shRNA-treated cell population, as represented in

(F-H). Unpaired t-test were performed across each bin for all cells, and a K-S test was

performed on the overall distribution. The share of MTs parallel to the edge (bin 0-10) is

significantly higher in control as compared to KIF5B depletions. Pixel numbers in the

analysis: SCR N=61,437 pixels across 9 cells, shRNA#1 N=15,215 pixels across 7

cells, shRNA#2 N= 21,125 pixels across 7 cells.

Figure 3. Microtubule sliding is facilitated through the ATP-independent MT-binding

domain of kinesin-1.

(**A**) Schematic of kinesin-1 (KIF5) and the Dominant Negative (KIFDN) and

heterodimerization strategy. Top schematic shows full length KIF5s, consisting of the

motor domain, stalk coil-coil domain and the tail. Three constructs utilized here include

(1) The KIF5C motor domain tagged with a blue fluorescent protein (BFP) and the FRB

for heterodimerization; (2) KIFDN^{wt} construct with KIF5B Tail domain tagged with the

mCherry fluorescent protein and the FKBP for heterodimerization. (3) KIFDN^{mut} construct

is the same as (2) but features a set of point mutations (magenta) making the ATP-

independent MT-binding domain unable to bind MT lattice. (**B**) Quantification of MT sliding

in FRAP assay in cells subjected to DN construct expression and heterodimerization

(shown as area of displaced MTs). MT displacement is shown as area of MTs displaced

into the bleached area after 5 minutes of recovery. See representative data (**C-G**). N= 5-

25 per condition. One-way ANOVA test was performed for statistical significance (p-value

<0.0001; ns, non-significant). **(C-G")** Frames from representative FRAP live-cell imaging

sequences. mEmerald-tubulin-expressing MIN6 cells. Inverted grayscale images of

maximum intensity projections over 1 µm-thick stacks by spinning disk confocal

microscopy. (C1-G1) The first frame after photobleaching. (C2-G2) A frame 5 minutes

(300 seconds) after photobleaching. Light-blue dotted lines indicate the edges of the

photobleached areas. Red arrows indicate MTs displaced into the bleached area. Scale

bars, 5 µm. (C3-G3) Schematics of experimental manipulation: green represents MTs,

blue represents endogenous KIF5B, magenta represents KIFDN^{wt}, purple represents

KIFDN^{mut}, gray represents KIF5C motor, orange bracket represents heterodimerizing

agent (rap, rapalog). Conditions: (**C1-C3**) Untreated control. Only endogenous KIF5B is

present. (**D1-D3**) KIFDN^{wt} overexpression. Endogenous KIF5B is unable to bind MTs.

(**E1-E3**) KIFDN^{mut} overexpression. It does not bind MTs and does not interfere with

endogenous KIF5B. (**F1-F3**) KIFDN^{wt} and KIF5C motor overexpression plus rapalog

treatment. Heterodimerization creates a large pool of motors capable of MT sliding. (**G-**

G") KIFDN^{mut} and KIF5C motor overexpression plus rapalog treatment.

Heterodimerization creates a large pool of the motor non-functional in MT sliding.

Figure 4. Effects of ATP-independent MT-binding domain of KIF5B on microtubule

abundance and alignment at the beta-cell cell periphery.

(**A-C**) MT organization in MIN6 cells expressing (**B**) KIFDN^{wt} and KIF5C motor

heterodimerized via rapalog treatment, (**C**) KIFDN^{mut} and KIF5C motor heterodimerized

via rapalog treatment and compared to a control cell with no ectopic expressions (**A**).

Top, immunofluorescence staining for tubulin (grayscale, inverted). Blue dotted line

indicates the borders of a cell expressing constructs of interest. Bottom in A, f-actin

(phalloidin, red) and DAPI (blue). Bottom in B and C, ectopically expressed mCherry-

labeled KIFDN constructs (magenta) and BFP-labeled KIF5C motor (green). Laser

scanning confocal microscopy maximum intensity projection of 1µm at the ventral side

of the cell. Scale bars: 5um. (**D**) Quantification of mean tubulin intensity within the outer

2µm peripheral area of a cell, in data represented in (A-C). Mean values, black bars.

One-way ANOVA, $p<0.0001$. N=5-15 cells. (**E**) Histograms of MT directionality within

1um of cell boundary (see Supplemental figure 2-1 for the analysis workflow) in control

cells compared to cells expressing heterodimerized KIFDN variants. Data are shown for

the summarized detectable tubulin-positive pixels in the analyzed cell population, as

represented in (F-H). Unpaired t-test were performed across each bin for all cells, and a

K-S test was performed on the overall distribution. The share of MTs parallel to the

edge (bin 0-10) is significantly higher in control as compared to the over-expressions.

NT control N=138,810 pixels across 9 cells, KIFDN^{wt} + motor N= 48,285 pixels across 9

cells, KIFDN^{mut} + motor N= 40,832 pixels across 10 cells. (**F-H**) Representative

examples of MT directionality analysis quantified in (E). (**F**) Control cell, no ectopic

expressions. (**G**) Cell expressing KIFDN^{wt}+ Motor. (**H**) Cell expressing KIFDN^{mut}+ Motor.

Overviews of cellular MT networks are shown as threshold to detect individual

peripheral MTs (see Supplemental figure 2-1 panel A5). (**F1-H2**) Directionality analysis

outputs of regions from yellow boxes in (F-H) are shown color-coded for the angles

between MTs and the nearest cell border (see Supplemental figure 2-1 panel A8). (**I**)

Color code for (F1-H2): MTs parallel to the cell edge, blue; MTs perpendicular to the cell

edge, red. Figure 4- Source Data 1

Figure 4 - Supplemental Figure 1. DN effect of KIF5B tail domain on MT abundance at

the beta-cell cell periphery requires ATP-independent MT-binding domain. (**A-B**) MT

organization in MIN6 cells expressing (**A**) KIFDN^{wt}, (**B**) KIFDN^{mut}, Top,

immunofluorescence staining for tubulin (grayscale, inverted). Blue dotted line indicates

the borders of a cell expressing constructs of interest. Bottom, ectopically expressed

mCherry-labeled KIFDN constructs (magenta). Laser scanning confocal microscopy

maximum intensity projection of 1μm at the ventral side of the cell. Scale bars: 5um. (**E**)

Quantification of mean tubulin intensity within the outer 2μm peripheral area of a cell, in

data represented in (A-B) as compared to untreated controls (see Figure 4A). Mean

values, black bars. One-way ANOVA, p<0.0001. N=7-11 cells.

Figure 4-Supplemental Figure 2. Influence of thresholding variations on the output

analysis of MT directionality in Figure 4 F-H. (**A**) Histograms of MT directionality within

1um of cell boundary using standard thresholds (see Supplemental figure 2-1 for the

analysis workflow) in cells expressing ectopic heterodimerized KIF5C motor. Data are

shown for the summarized detectable tubulin-positive pixels in the analyzed shRNA-

treated cell population, as represented in (Figure 4 F-H). Unpaired t-test were

performed across each bin for all cells, and a K-S test was performed on the overall

distribution. The share of MTs parallel to the edge (bin 0-10) is significantly higher in

control as compared to heterodimerized KIF5B over-expression. Pixel numbers in the

analysis: NT control N=120,959 pixels across 9 cells, KIFDN^{wt} + motor N= 26, 110

pixels across 7 cells, KIFDN^{mut} + motor N= 41,091 pixels across 10 cells.

Figure 5 Enhanced MT sliding results in loss of peripheral MT alignment at the border

(**A-B**) MT organization in MIN6 cells pretreated with DMSO and Kinesore respectively.

Immunofluorescence staining for tubulin (grayscale, inverted). Laser scanning confocal

microscopy maximum intensity projection of 1µm at the ventral side of the cell. Scale

bars: 5um. (**C**) Quantification of mean tubulin intensity within the outer 2µm peripheral

area of a cell, in data represented in (A-B). Mean values, black bars. One-way ANOVA,

p<0.0001. N=10 cells per condition. Histograms of MT directionality within 1um of cell

boundary (see Supplemental figure 2-1 for the analysis workflow) in DMSO treated

control cells compared to kinesore treated cells. Data are shown for the summarized

detectable tubulin-positive pixels in the analyzed cell population, as represented in (D-

E). Unpaired t-test were performed across each bin for all cells, and a K-S test was

performed on the overall distribution. The share of MTs parallel to the edge (bin 0-10) is

significantly higher in control as compared to the over-expressions. DMSO control

N=136,840 pixels across 10 cells, kinesore treated N= 87,361 pixels across 9 cells. (**D-**

E) Representative examples of MT directionality analysis quantified in (F). Directionality

analysis outputs of regions from yellow boxes in (D-E) are shown color-coded for the

angles between MTs and the nearest cell border (see Supplemental figure 2-1 panel

A8). (G) Color code for (D1-E2): MTs parallel to the cell edge, blue; MTs perpendicular

to the cell edge, red. Figure 5- Source Data 1

Figure 5-Supplemental Figure 1. Influence of thresholding variations on the output

analysis of MT directionality in Figure 5 D-E. (**A**) Histograms of MT directionality within

1um of cell boundary using standard thresholds (see Supplemental figure 2-1 for the

analysis workflow) in cells treated with DMSO or kinesore. Data are shown for the

summarized detectable tubulin-positive pixels in the analyzed respective cell population,

as represented in (Figure 5 D-E). Unpaired t-test were performed across each bin for all

cells, and a K-S test was performed on the overall distribution. The share of MTs

parallel to the edge (bin 0-10) is significantly higher in control as compared to kinesore

treated. Pixel numbers in the analysis: DMSO control N= 94,841pixels across 9 cells,

kinesore N= 29,796 pixels across 9 cells.

Figure 6. MT sliding in beta cells is stimulated by glucose.

(**A-B**) Frames from representative FRAP live-cell imaging sequences of MT sliding

response to glucose stimulation. mEmerald-tubulin-expressing MIN6 cells. Inverted

grayscale images of maximum intensity projections over 1 μm-thick stacks by spinning

disk confocal microscopy. (**A**) A cell pretreated with 2.8mM glucose before the assay.

(**B**) A cell pretreated with 2.8mM glucose and stimulated with 20 mM glucose before the

assay. (**A1-B1**) The first frame after photobleaching. (**A2-B2**) A frame 5 minutes (300

seconds) after photobleaching. Light-blue dotted lines indicate the edges of the

photobleached areas. Red arrows indicate MTs displaced into the bleached area. Scale

bars, 5 μm. (**C**) Quantification of MT sliding FRAP assay in cells in 2.8mM versus 20mM

glucose (see representative data in A-B). MT displacement is shown as area of MTs

displaced into the bleached area after 5 minutes of recovery. One-way ANOVA test was

performed for statistical significance (p-value <0.0001). N=16-24 cells per set. (**D-E**)

MIN6 cells featuring fiducial marks at MTs due to co-expression of SunTag-KIF5B-

560Rigor construct and Halo-SunTag ligand. Representative examples for cells in

2.8mM glucose (D) and a cell stimulated by 20mM glucose (E) are shown. Single-slice

spinning disk confocal microscopy. Halo-tag signal is shown as inverted gray-scale

image. Top panels show cell overviews (scale bars 5μm). Below, boxed insets are

enlarged to show dynamics of fiducial marks (color arrows) at 1 second intervals (1-5

seconds). 0- to 5-second tracks of fiducial mark movement are shown in the bottom

panel, each track color-coded corresponding to the arrows in the image sequences.

N=6-11 cells. (**F**) Histogram of all 5- second displacement of fiducial marks in Low vs

High glucose (**G**) Summarized quantification of stationary fiducial marks along MT

lattice. Low glucose N=5,615 tracks across 11 cells, High glucose 5min N=2,259 tracks

across 6 cells, High Glucose 20min N=3,059 tracks across 6 cells. P One-way ANOVA,

p<0.001 (**H**) Summarized quantification of moving fiducial marks along the MT lattice.

Low glucose N=2,595 tracks across 11 cells, High glucose 5min N=2,642 tracks across

6 cells, High Glucose 20min N=4,049 tracks across 6 cells. One-way ANOVA, p<0.001.

Video Titles

Figure 1-Video 1

KIF5B-mediated MT sliding visualization via FRAP. Cells treated with scrambled control

and two shRNAs against KIF5B are shown. Time, minutes:seconds.

Figure 1-Video 2

KIF5B-mediated MT sliding visualization via SunTag-KIF5B-560Rigor. Cells treated with

scrambled control and two shRNAs against KIF5B are shown. Sliding maximum

intensity projection (15 time frames projected in each video frame). Time, seconds.

Figure 3-Video 1

Dominant Negative MT sliding visualization via FRAP (Tails). Cells over-expressing

KIFDN^{wt} versus KIFDN^{mut} are shown. Time, minutes:seconds.

Figure 3-Video 2

Dominant Negative MT sliding via FRAP (Tails + Motors). Cells expressing

heterodimerized KIFDN^{wt} with motor versus heterodimerized KIFDN^{mut} with motor are

shown. Time, minutes:seconds.

Figure 6-Video 1

Glucose-Dependent MT sliding visualization via FRAP. Cells in low versus high glucose

are shown. Time, minutes:seconds

Figure 6-Video 2

Glucose-Dependent MT sliding visualization via SunTag-KIF5B-560Rigor. Cells in low

versus high glucose are shown. Sliding maximum intensity projection (15 time frames

projected in each video frame). Time, seconds.

Source Data

Figure 2- Source Data 1

MT Directionality Source Data: KIF5B Depletion

Figure 4- Source Data 1

MT Directionality Source Data: KIFDN Overexpression

Figure 5- Source Data 1

MT Directionality Source Data: Kinesore Treatment

Figure 6- Source Data 1

SunTag marks displacement 5s intervals across each cell. And population analysis referenced

in panels G and H.

943