1 **A Bioluminescent Activity Dependent Platform, BLADe, for Converting Intracellular**
2 **Activity to Photoreceptor Activation**
3

4 Emmanuel L. Crespo[1,2]†, Akash Pal[1,3]†, Mansi Prakash[1], Alexander D. Silvagnoli[1,3], Zohair
5 Zaidi[4], Manuel Gomez-Ramirez[5], Maya O. Tree[1], Nathan C. Shaner[6], Diane Lipscombe[5,7],
6 Christopher I. Moore[5,7]*, Ute Hochgeschwender[1-3]*
7
[1] College of Medicine, Central Michigan University, Mount Pleasant, MI 48859, USA
[2] Biochemistry, Cell and Molecular Biology Graduate Program, Central Michigan University,
Mount Pleasant, MI 48859, USA
[3] Program in Neuroscience, Central Michigan University, Mount Pleasant, MI 48859, USA
[4] Duke University, Undergraduate Neuroscience Program, Durham, NC 27710
[5] Department of Neuroscience, Brown University, Providence, RI 02912, USA
[6] University of California, San Diego, School of Medicine, Department of Neuroscience, 9500
Gilman Drive La Jolla, CA 92093-0662, USA
[7] Carney Institute for Brain Science, Brown University, Providence, RI 02906, USA
17
18 †These authors contributed equally to this work: Emmanuel L. Crespo [Present address:
19 University of Michigan, Ann Arbor, MI 48109, USA] and Akash Pal [Present address: Max
20 Planck Florida Institute for Neuroscience, Jupiter, FL 33458, USA]
21
22 *Corresponding authors: Ute Hochgeschwender (hochg1u@cmich.edu) and Christopher I.
23 Moore (christopher_moore@brown.edu)

24
25
26

Abstract

Genetically encoded sensors and actuators have advanced the ability to observe and manipulate cellular activity, yet few non-invasive strategies enable cells to directly couple their intracellular states to user-defined outputs. We promote a bioluminescent activity-dependent (BLADe) platform that facilitates programmable feedback through genetically encoded light generation. Using calcium (Ca^{2+}) flux as a model, we engineered a Ca^{2+}-dependent luciferase that functions as an activity-gated light source capable of photoactivating light-sensing actuators. As an initial demonstration of the versatility of this platform we present two separate use cases in neurons. In the first application, the presence of luciferin triggers Ca^{2+} dependent local illumination that provides activity dependent gene expression by activating a light-sensitive transcription factor. In the second application, neuronal activity-driven Ca^{2+} fluctuations via locally generated bioluminescence control neural dynamics through opsin activation in single cells, populations and intact tissue. BLADe can be expanded to couple any signal that bioluminescent enzymes can be engineered to detect with the wide variety of photosensing actuators. This modular strategy of coupling an activity dependent light emitter to a light sensing actuator offers, in principle, a generalizable framework for state dependent cell-autonomous control across biological systems.

Introduction

Bioluminescence, the emission of light generated when a luciferase enzyme oxidizes a small molecule luciferin, served as a non-invasive approach for *in vitro* and *in vivo* imaging for decades[1]. More recently, the development of brighter, blue light emitting luciferases in combination with light sensing elements, including opsins and non-opsin photoreceptors[2,3], transformed bioluminescence from a mere imaging reagent to a versatile tool for controlling molecular processes. We and others have combined luciferases with channelrhodopsins and pumps for bioluminescence-driven changes in membrane potential[4–6], with light-sensing transcription factors for gene expression[7–9], with light-sensing enzymes for cAMP production[10], as well as with genetically encoded photosensitizers for reactive oxygen species production[11,12]. The main advantages of using locally produced bioluminescence rather than external light sources for activating photoreceptors are the independence from optical fiber implants, thereby avoiding the physical damage to cells as well as from the limitation in the number of cells that can be simultaneously controlled.

A unique feature of genetically encoded light emitters is that they are amenable to molecular engineering. By inserting into the luciferase protein sensor moieties, light emission can be made dependent on the presence of various intracellular agents that report a cell's state. Several such bioluminescent sensors have been developed for reporting intracellular biochemical signals, such as intracellular Ca^{2+}, ATP, or cAMP, including for imaging studies *in vivo*[13]. Here, we are combining a bioluminescent sensor with light-dependent actuators to promote a general platform for converting biochemical signals within cells into photoreceptor activation and resulting downstream changes in cellular function. This Bioluminescence Activity Dependent (BLADe) paradigm serves to integrate cellular activity states with downstream consequences from transcription to membrane potential changes, depending on the optogenetic actuator coupled to the conditional luciferase. By engineering luciferases to respond to specific biochemical cues, this framework can, in principle, be extended to a broad range of intracellular states depending on the sensor domain used. Any bioluminescence indicator can, in principle, be transformed from a sensor into an activity integrator by using its light emission to activate photoreceptors.

78
As an initial demonstration of the versatility of this platform we employed a Ca²⁺ dependent
luciferase. By making light production dependent on Ca²⁺ binding in a split luciferase construct,
changes in Ca²⁺ levels are directly translated into bioluminescent light emission. Conversion of
intracellular calcium level fluctuations into photoreceptor activation was tested in two separate
applications in neurons. The Ca²⁺ dependent light emitter was paired with a light-dependent
transcription factor that detected this light and affected transcription *in vitro* and *in vivo*. In
another test case, when paired with channelrhodopsins Ca²⁺ dependent light emission enabled
neural activity-driven change in firing activity *in vitro* in single neurons and neuronal populations;
in mice, BLADe converted spontaneous activity and sensory evoked changes into activity
dependent control of neocortical network dynamics.
89
90
91 **Results**
92
93 **Engineering of a luciferase-based Ca²⁺ sensor**
94
The *Gaussia princeps* luciferase variant sbGLuc is an efficient transmitter of photons/energy to
opsin chromophores when tethered to opsins in luminopsins[5,14]. To make it Ca²⁺ dependent we
split sbGLuc as was done previously for wildtype GLuc between Q88 and G89[15] and introduced
the Ca²⁺ sensing moiety CaM-M13 employing the M13 sequence from the Ca²⁺ sensor YC3.6[16]
and the calmodulin sequence from GCaMP6f[17] (**Fig. 1a,b**). We generated a preliminary
structural model that is, however, limited due to the lack of evolutionary structure data similar to
the *Gaussia* luciferase protein structure (**Fig. 1c**). Nevertheless, we hypothesized that inserting
the CaM-M13 domain between Q88 and G89 would allosterically control enzymatic activity by
modulating the structural integrity of the native disulfide bond (C65–C77) which stabilizes the
rigid α4–α7 helical core of sbGLuc (**Supplementary Fig. S1 and S2**). Assuming this
mechanism, in the absence of Ca²⁺ the inserted calmodulin domain destabilizes this interaction,
whereas Ca²⁺ binding promotes reassembly of the split halves and restoration of luciferase
activity. We called this split sbGLuc based Ca²⁺ indicator Lumicampsin or LMC.
108
As the goal of LMC is to serve as a genetically encoded light source for activating blue light-
sensing photoreceptors, we first confirmed its light emission spectra and Ca²⁺ responsiveness.
HeLa cells were transfected with LMC, incubated with 5 µM luciferin (CTZ), and stimulated with
histamine to induce a physiologically relevant increase in intracellular Ca²⁺. LMC retained a
peak emission at roughly 490 nm (**Fig. 1d)** with a percent change in light emission before and
after histamine addition of 232% (**Fig 1e**, IQR: 215–262%; range: 172–318%; N = 12 wells).
Thus, LMC provides a blue-shifted optical report of intracellular calcium, enabling potential
coupling to blue light sensitive optogenetic proteins required for the BLADe platform.
117
Experiments in HeLa cells confirmed that LMC responds to histamine-induced Ca²⁺ release at
physiologic levels, but this system does not allow experimental control of intracellular Ca²⁺
levels in living cells. To address this, we transfected HEK293 cells with LMC and applied
ionomycin under defined extracellular Ca²⁺ concentrations ranging from 0.4875 µM to 39 µM
(**Fig. 1f**). LMC reported ionomycin-induced Ca²⁺ flux at all tested concentrations. The decrease
in bioluminescent signal at lower extracellular Ca²⁺ conditions (0.4785µM -1µM) reflects the loss
of intracellular Ca²⁺ as ionomycin moves Ca²⁺ from high (intracellular) to low (extracellular)
concentrations.
126

127 To test the ability of LMC to report Ca²⁺ dynamics in cultured cells, we transfected primary
128 cardiomyocytes with LMC and imaged them on DIV4 using an EMCCD camera. Cells were pre-
129 incubated with luciferin (100 µM CTZ) for ~10 min before imaging. Addition of 10 µM ATP
130 increased contractility and induced rapid Ca²⁺ transients with robust peaks (**Fig. 1g-j**).
131
Next, to test whether LMC could report neurotransmitter-evoked Ca²⁺ influx in neurons, we
transfected primary hippocampal cultures and perfused luciferin (CTZ) followed by luciferin with
glutamate. Bioluminescent signals were low with CTZ alone but increased after glutamate
application **(Fig. 1k)**. Time-locked traces showed a clear rise in bioluminescence following
glutamate onset (**Fig. 1l**). Across individual neurons, we observed an increase in normalized
peak responses with glutamate compared to luciferin alone (**Fig. 1m**). In addition, responses
also exhibited high contrast relative to baseline, with a median fold-change of ~2–4 across
neurons (**Fig. 1n**).
140
141 **Bioluminescence as an activity-gated light source**
142
We next compared LMC with other published bioluminescent Ca²⁺ indicators to evaluate its
baseline emission and suitability as an activity-gated light source. HeLa cells were transiently
transfected with the *Gaussia* luciferase based LMC, and the *Oplophorus* luciferase based
BlueCaMBI[18], GLICO[19], or GeNL[20] and light emission was assayed in a luminometer following
luciferin addition (CTZ for LMC, hCTZ for the other three sensors) and histamine stimulation.
Time-series traces show that BlueCaMBI, GLICO, and GeNL, produced higher overall photon
output than LMC (**Fig. 2a**). Although these sensors were brighter, they also exhibited substantial
baseline photon emission in the presence of luciferin under resting Ca²⁺ conditions, relative to
their response after histamine stimulation (**Fig. 2b**). For imaging Ca²⁺ fluctuations the baseline is
defined as the value immediately before histamine addition, and Ca²⁺ spikes are represented by
these before/after values. However, bioluminescent sensors will not be suitable as drivers of
photoreceptors unless their light emission in the initial presence of luciferin at baseline is below
the activation limit of photoreceptor activity. All three indicators—BlueCaMBI, GLICO, and
GeNL—have been reported to capture Ca²⁺ fluctuations, but in those studies, bioluminescence
signal traces were typically normalized following luciferin pre-incubation[18–20]. While this
normalization approach is appropriate for imaging relative Ca²⁺ dynamics, the high absolute
photon emission at resting levels of Ca²⁺ limits their use as activity-gated light sources required
for BLADe activation of photoreceptors.
161
High absolute brightness benefits imaging, but BLADe optogenetic applications require low
basal emission and a large absolute increase in light output during Ca²⁺ influx. When a sensor is
used for coupling light emission with photoreceptor activation, the critical baseline value is the
level of light emission once the luciferin enters the cell. If that initial level is too high, the
photoreceptor will be activated at baseline, rather than being activated as a consequence of
increased levels of Ca²⁺. Consistent with this, when normalized to the luciferin baseline, only
LMC exhibited a large relative change in photon emission following histamine-induced Ca²⁺
release (**Fig. 2c**). Accordingly, despite lower absolute brightness, the unique low background
and high dynamic range of LMC positions it as an activity-gated light source ideally suited for
converting intracellular signals into optical activation of downstream photoreceptors. At resting
calcium levels, we hypothesize LMC light emission remains dim and is below the threshold
needed to activate photoreceptor chromophores, minimizing spurious activation of light sensing
proteins. We next asked whether this activity-dependent light source could drive optogenetic
tools as the basis of the BLADe platform.
176

177
178
179 **Integrating Ca²⁺ flux with change in transcription**

180
LMC expressing cells showed low baseline bioluminescence in the presence of luciferin and a
strong increase in light output following Ca^{2+} influx. We reasoned that LMC's activity-gated light
emission could be used to photoactivate light-sensitive transcriptional effectors. To test this
prediction, we co-expressed LMC with the light sensing transcription factor EL222[21]. In the
presence of blue light, EL222 dimerizes and then binds to its promoter 5xC120, leading to
transcription of the reporter gene (**Fig. 3a**).

187
HeLa cells expressing EL222, the Firefly luciferase (FLuc) reporter, and each calcium sensor
were either left untreated, treated with luciferin for 15 minutes, or treated with luciferin and
histamine for 15 minutes. Transcription of the FLuc reporter was measured 18 hours later as
luminescence after addition of D-luciferin to the cells (**Fig. 3b**). Left untreated or in the absence
of a sensor, no EL222 activation was observed. With addition of luciferin (CTZ for LMC, hCTZ
for the other sensors) transcription at baseline Ca^{2+} levels is very low for unreconstituted LMC,
in contrast to the other three sensors. The addition of luciferin and histamine should reflect the
increased light emission of the reconstituted – due to the histamine-induced increase in
intracellular Ca^{2+} – luciferase resulting in higher EL222 activation and higher transcription of
FLuc. This prediction is seen with LMC, but not with the other three Ca^{2+} sensors, due to their
high baseline signal. Interestingly, the absolute FLuc reporter expression for BlueCaMBI,
GLICO, and GeNL is lower in cells exposed to luciferin and histamine than in cells exposed to
luciferin alone. This is likely a reflection of the increased consumption of the luciferin in the
presence of increased Ca^{2+}, resulting in decreasing the time of light emission for EL222
activation.

203
All calcium dependent indicators produced FLuc transcription under luciferin treatment alone,
indicating that transcription can be photoactivated even at very low levels of calcium. In the
presence of luciferin, baseline transcription was strongest for BlueCaMBI, GLICO, and GeNL,
which were markedly brighter than LMC — a trend consistent with their higher baseline
bioluminescence in luminometer traces (**Fig. 2b**). In contrast, only LMC showed a calcium-
dependent increase in transcription, with a significant increase in FLuc signal in the luciferin +
histamine condition relative to all other sensors (**Fig. 3c**). LMC maintains a low luciferin baseline
and undergoes a discrete calcium-dependent increase in light output, preserving a clear
calcium-state transition. Moreover, the overall transcriptional output from LMC remained modest
compared to the other sensors, which is consistent with its lower total light output in
luminometer assays (**Fig. 2b**). As a result, effective photoreceptor gating in the BLADe system
depends not on absolute light output but on preservation of a discrete calcium-state transition
above a low luciferin baseline. Having established that LMC produces calcium-dependent and -
state-specific light sufficient to activate EL222 *in vitro*, we next tested whether the BLADe
transcriptional system could function *in vivo*.

219
As a pilot study, we tested whether LMC could drive transcription in neurons *in vivo* when
network activity was chemically elevated with NMDA. We performed stereotactic injections of
AAV9-hSyn-LMC-P2A-dTomato, AAV9-CAG-EL222, and AAV9-5×C120-EYFP into the
prefrontal cortex of anesthetized mice. After virus delivery, the craniotomy was immediately
covered to further minimize unintended light exposure during recovery. After 2–3 weeks, to
allow for maximal expression, mice received intraventricular injections of either NMDA + vehicle,
luciferin (CTZ) alone, or luciferin + NMDA. These conditions allowed us to test: (1) non-specific
background transcriptional activity in the absence of light (NMDA + vehicle), (2) light-dependent

transcription in the context of spontaneous activity (luciferin alone), and (3) BLADe-mediated
transcription (luciferin + NMDA).
230
Confocal imaging of tissue collected ~20 hours after treatment showed increased EYFP reporter
labeling in individual LMC-expressing neurons from mice treated with luciferin + NMDA,
compared to the other treatment groups (**Fig. 3d,e**). Single-cell analysis confirmed that EYFP
reporter increase occurred across a range of dTomato expression levels (**Fig. 3f**), indicating that
the Ca²⁺-dependent enhancement of transcription was not due to differences in viral expression
of LMC. Quantification of EYFP reporter/dTomato tool expression ratios revealed higher
reporter expression in the luciferin + NMDA group compared to both the luciferin-only group and
the NMDA-only group (**Fig. 3g**, Wilcoxon rank-sum test, Bonferroni-corrected luciferin+NMDA
vs luciferin: P= 1.62×10^{-7}; luciferin+NMDA vs NMDA: P= 0.0057; luciferin vs NMDA: P =
0.196; pooled from FOVs of 197-350 cells/N=2 mice for each treatment group). The luciferin-
only and NMDA-only groups were not significantly different from each other suggesting that
reporter activation requires both intracellular calcium and bioluminescent light, as required with
LMC-BLADe gating. Together, these findings are consistent with BLADe functioning *in vivo* in
the mouse brain to drive calcium-dependent transcription, with the overall magnitude of reporter
labeling remaining modest. Increasing light output of the Ca²⁺ dependent luciferase, improving
the contrast between active and inactive cells, and simplifying viral expression strategies may
optimize this system for robust transcriptional labeling in future applications.
248
249 **Integrating Ca²⁺ flux with change in membrane potential**
250
To demonstrate the modularity of the BLADe platform, we tested whether LMC could drive light-
sensitive ion channels. Following an action potential, the influx of intracellular Ca²⁺ provides a
convenient and high-resolution proxy for BLADe neural activation. We reasoned that LMC could
detect these calcium transients and, through co-expression with an excitatory opsin, rapidly alter
spiking dynamics in an activity-dependent manner. To this end, we applied step current
injections to hippocampal neurons *in vitro* in the presence of synaptic blockers to isolate cell-
intrinsic responses. We then performed whole-cell patch-clamp recordings on neurons co-
transformed with LMC and an opsin (**Fig. 4a**). By visualizing co-localized expression of an
opsin-EYFP (green; here ChR2(C128S)) and the Ca²⁺-dependent luciferase LMC-P2A-dTomato
(red), we selected these double-positive neurons for testing the integration of cellular states via
BLADe opsin activation.
262
We predicted that LMC generates light directly in response to Ca²⁺ influx which occurs during
action potential generation induced by depolarizing current injections. We reasoned that
increasing levels of depolarizing current in the presence of the luciferin would produce enough
bioluminescence to further photoactivate ChR2(C128S)[22] via a cell-autonomous, positive
feedback optical loop. In line with our prediction, luciferin perfusion during the same depolarizing
current elicited significantly more detected action potentials than during vehicle perfusions (**Fig
4b, ii and iii,** N=9 neurons, p= 4.954×10^{-9}, paired sample Wilcoxon signed rank tests), indicating
a light-dependent, cell-autonomous control of membrane voltage potential.
271
These initial findings are consistent with LMC bioluminescence being sufficient to photoactivate
the opsin, but they do not directly prove BLADe control or reveal how the effect emerges over
the course of the 1-second depolarizing step. Given that LMC reports spontaneous intracellular
calcium dynamics with sub-second temporal resolution (**Fig. 1**), we hypothesized that any
optical feedback effect should manifest within the timescale of the depolarizing step. If BLADe
control depends on activity-coupled light production, then the timing of spiking should reflect this
coupling, i.e. the effect of LMC-driven opsin activation should evolve across the depolarizing

step. To this end, we first measured how spikes were distributed across the 1-second
depolarizing step using a normalized early–late comparison per trial to detect timing shifts rather
than changes in overall firing (see Methods). The spike-timing distribution shifted toward the end
of the 1-second depolarizing current during luciferin perfusion in neurons co-expressing LMC
and ChR2(C128S), in an opsin- and CTZ- dependent manner (**Supplementary Fig. S3**). Our
observations support a model in which LMC-driven bioluminescence engages the opsin through
a voltage-dependent feedback process and prompted us to characterize how LMC alters the
relationship between early and late spiking across the depolarizing stimulation.
287
We reasoned LMC-driven bioluminescence is not merely a tonic light source but may also act
as an activity-coupled light emitter, such that its optogenetic effect on neuronal firing scales with
the level of spiking activity within a trial. We therefore divided each trial into an early window (0–
200 ms following stimulation onset) and a late window (>200 ms), capturing initial versus
sustained spiking activity under continued current stimulation. To separate tonic light effects
from activity dependent modulation and quantify how LMC integrates ongoing spiking activity
into later responses, we applied a trial-level linear model relating late firing to early activity,
luciferin condition, and their interaction, pooled across neurons (see Methods). In this model, β_1
captures the relationship between early and late phase spiking in the vehicle condition,
reflecting the baseline relationship between early and late spiking within a trial. In contrast, the
main luciferin effect term (β_2) reflects an activity independent shift in late firing, consistent with
tonic optogenetic drive. The interaction term (β_3) captures activity dependent modulation (gain),
reflecting amplification (or suppression) of spiking that scales with ongoing activity.
301
Accordingly, a tonic optogenetic effect in the presence of luciferin would manifest as a uniform
shift in firing (β_2) with the direction of the shift (additive: $\beta_2 > 0$; subtractive: $\beta_2 < 0$) determined
by the biophysical properties of the co-expressed opsin. In contrast, activity-dependent
modulation would be expressed as a change in the slope linking early to late activity ($\beta_3 \neq 0$).
Thus, we predicted that LMC-driven bioluminescence during luciferin perfusion would rapidly
shape early–late epoch spike coupling, with positive β_3 values indicating multiplicative gain with
excitatory opsins and negative β_3 values indicating divisive suppression with inhibitory opsins.
309
For neurons expressing the excitatory opsin (**Fig. 4b iv**), luciferin increased the slope relating
early to late phase spiking, indicating that early epoch activity more strongly predicted late
responses and that this relationship was multiplicatively scaled across trials (β_1= 3.927, CI [1.49,
6.37], p= 0.002, β_2= 9.33, CI [3.63, 15.02] p= 0.002, β_3= 5.02, CI [2.30, 7.74], p = 0.001).
Across recordings, multiplicative gain was observed in most neurons (**Fig. 4b v**), with 6/9
recordings exhibiting increased slope relative to baseline and the remainder showing either no
change (1/9) or reduced slope (2/9). Together, in the presence of luciferin, LMC converts early
spiking activity into activity-dependent gain, amplifying late responses in proportion to the
magnitude of spiking activity established during the onset of stimulation.
319
In contrast, neurons co-expressing LMC and ChR2(C128S) that received two sequential vehicle
perfusions showed no changes in spiking dynamics. Overall firing rate and the temporal
distribution of spikes across the stimulus window was unaltered across trials **(Fig. 4c ii-iii**, N=14
neurons, *p*= 0.3117*,* paired sample Wilcoxon signed rank test). Consistent with this observation,
the linear model revealed no evidence of either tonic modulation or activity dependent changes
in gain (**Fig. 4c iv**, β_1= 4.54, CI [3.29, 5.78], p= 0.001, β_2= 0.55, 95% CI [-2.39, 3.49] p= 0.71,
β_3= 0.23, CI [-1.54, 2.00], p= 0.794). No consistent change in slope was observed, with most
recordings (10/14) showing no measurable change and a subset of neurons (4/14) exhibiting
reduced slopes (**Fig. 4c v**). Our result confirms that the observed bias in spike timing requires
luciferin, and by extension, LMC stimulus-dependent bioluminescence.

330
To confirm these changes depend on a functional opsin rather than bioluminescence alone, we
next tested a non-photoconductive ChR2 mutant, ChR2(C128S)-E97R-D253A[23,24] ("DUD"
opsin) alongside LMC (**Supplementary Fig. S4**). There were no observed luciferin-induced
shifts in spike timing or firing rate, indicating that BLADe control does not alter the intrinsic
excitability or temporal dynamics of spike emergence across trials in the absence of a functional
opsin.
337
Building on our findings with the excitatory opsin, we next tested whether LMC could suppress
firing in neurons expressing an inhibitory opsin. The same activity-coupled mechanism is
expected to produce divisive suppression, such that increases in early spiking reduce
subsequent firing later in the stimulus epoch. In line with our prediction, neurons co-expressing
hGtACR2[25] and LMC exhibited fewer total spikes (**Fig. 4d, ii and iii**, N=10 neurons, p=. 0.3117,
paired sample Wilcoxon signed rank test) and weakened the relationship between early and late
spiking after accounting for tonic optogenetic effects (**Fig. 4d, iv,** β_1= 7.22, CI [5.36, 9.07], p=
0.001, β_2= -11.44, CI [-20.89, -2.003], p=0.019, β_3= -6.489, CI [-20.89, -2.003], p= 0.019).
Across recordings, divisive suppression was observed in most neurons (**Fig. 4d v**), with 7/10
recordings exhibiting reduced slope relative to baseline and the remainder showed no
measurable slope change (3/10).
349
Moving beyond single-cell recordings, we next tested how LMC modulates firing rates at the
network level by performing extracellular recordings on cultured cortical neurons plated on multi-
electrode arrays (MEA)[26]. We virally co-transduced neurons with an excitatory opsin
ChR2(C128S)-EYFP and the Ca^{2+}-dependent luciferase LMC-P2A-dTomato (**Fig. 5a,b**). Time-
locked multi-unit activity (MUA) traces revealed rapid population-level activity following luciferin
or vehicle application (**Fig. 5c**). While vehicle addition alone can transiently increase network
activity in MEAs—likely due to mechanical or ionic effects of media perfusion—these responses
were brief and stereotyped across independent experiments. To enable comparison, we aligned
all trials to the application artifact, allowing time-locked analysis between vehicle and luciferin
treatments. In doing so, we could distinguish true activity-dependent changes from non-specific
network effects caused by vehicle addition. Vehicle and luciferin treatments were randomized
on the same MEA culture. All comparisons used the same electrode sites with recording ~4-5
hours apart.
363
In the presence of luciferin, MUA was enhanced relative to vehicle one second after application
(**Fig. 5d**). Burst firing rates were similar between conditions (**Fig. 5e**), yet burst duration was
significantly longer with luciferin (**Fig. 5f**). With burst firing rates unchanged but MUA elevated,
we next tested whether neural activity was prolonged over time due to co-expression of an
excitatory opsin with LMC. To this end, we analyzed the early phase as defined by the 0–50 ms
window beginning immediately after the application artifact, corresponding to the initial rise in
MUA relative to baseline (**Fig. 5g**). The time-locked window allowed for reliable comparisons of
immediate firing between treatment conditions. The late phase was defined as the remaining
portion of the 1-second window, from 100–1000 ms post-application, allowing us to test whether
LMC modulates both the onset and sustained components of network activity.
374
Early phase firing rates were similar between vehicle and luciferin conditions indicating that
initial responses to perfusion were comparable across treatments (**Fig. 5h**). Thus, the observed
effects are not due to media-driven excitability or non-specific stimulation. In contrast, the late-
phase response was significantly enhanced in the presence of luciferin (**Fig. 5i).** When
normalizing the late-phase MUA to the early-phase, we observed a clear enhancement in MUA
consistent with LMC activation with luciferin (**Fig. 5j**). The observed sustained effect on network

excitability in the presence of the luciferin suggests that LMC-driven light production prolongs
neuronal activity via local activation of the co-expressed excitatory opsin. Finally, to test whether
LMC can bidirectionally modulate population activity, we co-expressed LMC with the inhibitory
opsin hGtACR2 and observed luciferin-dependent suppression of MUA, confirming that
inhibition requires LMC expression (**Supplementary Fig. S5,** median ΔFR = +7 Hz, *n* = 19 vs
median ΔFR = –14 Hz, *n* = 51 with LMC, Wilcoxon rank-sum test, p= 1.576 × 10^{-10}). Our MEA
experiments extend our single-cell findings by testing whether LMC optogenetically modulates
neural activity at the population level in cultured neurons. However, because spiking dynamics
are not resolved on a trial-by-trial basis in this preparation, we next tested whether LMC
mediates activity-dependent gain *in vivo* using extracellular recordings.

Building on our *in vitro* electrophysiology observations, we next wanted to determine if LMC can
integrate physiological sensory inputs to alter neocortical networks. We leveraged the well-
characterized vibrissae circuit to ask whether LMC can detect sensory cues and reshape how
they are encoded within the neocortex. To this end, we injected Cre-dependent versions of LMC
and the excitatory opsin CheRiff[27] into the left cortex of Emx1-Cre mice to target both constructs
to excitatory pyramidal neurons within the vibrissa region of primary somatosensory cortex
(vSI). Three to four weeks later, we then performed extracellular recordings with laminar silicon
probes in vSI. As a control, a separate cohort of Emx1-Cre mice received identical surgical
procedures and recordings but did not express the opsin (N = 2 mice per group). During
recordings in head fixed anesthetized mice, we perfused ACSF or ACSF containing luciferin
(CTZ) over the cortical surface while delivering vibrissal deflections (**Fig. 6a, b**).

We first tested whether luciferin-driven bioluminescence alone was sufficient to enhance
spontaneous cortical activity in the absence of external light stimulation. By expressing CheRiff
in vSI pyramidal neurons, we predicted increased MUA either through external LED stimulation
or by luciferin-driven bioluminescence. CTZ application enhanced the MUA firing rates in opsin-
expressing mice relative to the null distribution derived from mice lacking opsin expression (**Fig.
6c**, No opsin: 21.38 Hz vs opsin: 50.20 Hz two-sided, unpaired Wilcoxon rank-sum test, p=
3.159 ×10^{-11}). At the end of each recording session during CTZ perfusion, a brief sinusoidal
LED stimulus (8 Hz, 500 ms) was delivered to confirm optogenetic responsiveness (**Fig. 6d**).
LED-evoked MUA increased in opsin-expressing mice relative to the null distribution derived
from the control cohort (No opsin: 2.456 Hz vs opsin: 86.420 Hz; two-sided, unpaired Wilcoxon
rank-sum test, p= 7.6 ×10^{-17}). Together, these results are consistent with bioluminescent
optogenetic activation of CheRiff selectively enhancing spontaneous cortical activity *in vivo*.

Following our observation that LMC enhances cortical activity during luciferin perfusion, we next
tested whether LMC can rapidly amplify sub-second neural responses to sensory inputs.
Consistent with BLADe activation of CheRiff, we tested for relationships between enhanced
neural activity and vibrissae stimulation by comparing MUA between vehicle and CTZ epochs.
We delivered 10 vibrissae deflections at 20 Hz over a 500 ms stimulus window, with deflections
randomized in amplitude. We observed evoked spikes within ~10–20 ms in vSI following low-
amplitude vibrissae deflections, consistent with known sensory latencies in vSI and validating
our electrode placement (**Fig. 6e**). To isolate BLADe-dependent modulation of sensory-evoked
cortical activity, we analyzed whisker-evoked multiunit activity (MUA) in a 500 ms post-stimulus
window. We then computed the CTZ-induced change in stimulus-locked MUA (POST–PRE) for
each channel and compared these responses between CheRiff-expressing and control mice.
Opsin-expressing mice displayed enhanced evoked MUA activity in the presence of CTZ
following continuous vibrissae deflections, consistent with BLADe dependent recruitment of
sensory-evoked cortical responses (**Fig. 6f**, p= 2.65 ×10^{-17}, two-sided, unpaired Wilcoxon rank-
sum test). In control mice, baseline-corrected evoked responses were centered near zero and

showed a slight bias toward decreased activity, indicating nonspecific CTZ-related modulation
that was not contingent on opsin expression. Together, these findings indicate that LMC rapidly
enhances stimulus-locked cortical activity by linking ongoing neural activity to additional
optogenetic drive, leading to increased sensory evoked responses.
436
We next asked whether the activity-dependent gain observed in the patch clamp recordings with
excitatory opsins extends to the circuit level *in vivo*. In CheRiff-expressing mice (**Fig. 6g**),
luciferin perfusion produced a tonic increase in late activity (β_1= 0.30, CI [0.28, 0.32], p= 1.29
$\times10^{-19}$, β_2= 7.01, CI [5.16, 8.85], p= 9.998 $\times10^{-14}$). After controlling for this optogenetic effect,
luciferin perfusion allowed sensory-evoked responses to be amplified as a function of early
spiking activity, such that trials with greater early activity exhibited larger responses during the
late epoch (β_3= 0.0647, CI [0.044, 0.086], p= 1.721 $\times10^{-9}$). This corresponded to a 22%
enhancement of late-phase firing (β_3/β_1 = 0.0647/0.301)**.** In contrast, mice not expressing
CheRiff (**Fig. 6h**) showed no evidence of gain modulation (β_3 = −0.024, 95% CI [−0.061, 0.012],
p = 0.189), corresponding to a negligible 4% decrease in slope. Luciferin perfusion produced a
non–opsin-specific tonic suppression of late activity (β_1= 0.6031, CI [0.568, 0.638], p= 8.38 $\times10^{-25}$,
β_2= −5.88, CI [−8.12, −3.64], p= 2.8 $\times10^{-7}$). Application of CTZ revealed a multiplicative gain
effect in late spiking activity as a function of early firing selectively in LMC and CheRiff co-
expressing mice (**Supplementary Fig. S6**).
451
452 Collectively, these findings are consistent with LMC enabling cell-autonomous, activity-gated
453 control of sensory-evoked cortical activity *in vivo*. Through co-expression of LMC with light
454 sensitive ion channels, BLADe photoactivation offers a genetically encoded strategy for sensing
455 and rapidly (sub-second) manipulating stimulus locked neural dynamics.
456
457 **Discussion**
458
In this study we showed evidence in support of a bioluminescent activity-dependent (BLADe)
platform that utilizes a luciferase split by a sensing moiety to drive a photoreceptor, allowing
integration of intracellular biochemical fluctuations with cellular outcomes. For an example
application, we engineered a Ca^{2+} dependent luciferase and applied it in two scenarios, the
neuronal activity driven activation of a) a light sensing transcription factor and of b) a light
sensing ion channel.
465
We split the *Gaussia* luciferase variant sbGLuc and inserted a Ca^{2+} sensing domain composed
of the M13 peptide from YC3.6 and calmodulin from GCaMP6f to generate lumicampsin (LMC).
When comparing LMC to other bioluminescent Ca^{2+} sensors we identified critical design
constraints for enabling activity dependent control of photosensing molecules: at resting
intracellular Ca^{2+} levels, the presence of luciferin should generate minimal light output – ideally
below the activation threshold of photoreceptor chromophores. Furthermore, elevation of
intracellular Ca^{2+} must produce a sufficient increase in bioluminescence to cross this threshold
and initiate photoreceptor activation.
474
Several excellent bioluminescent Ca^{2+} sensors have been developed by splitting luciferase
enzymes and inserting Ca^{2+} sensing moieties[16,18–20,28–30], and new sensors for reporting various
intracellular signals are being developed. Because bioluminescent probes have historically been
far dimmer than fluorescent probes, most efforts have focused on maximizing light output. As a
result, existing bioluminescent Ca^{2+} sensors successfully report Ca^{2+} fluctuations, but with Ca^{2+}
dependent increases in signal riding on top of high baseline light emission. For many imaging
applications, baseline bioluminescent signal in the absence of high Ca^{2+} is acceptable.

However, this is incompatible with the BLADe approach, as the background light emission by
itself is already high enough to activate a photoreceptor. Thus, an excellent bioluminescent
sensor cannot be automatically assumed to be a suitable driver for BLADe applications. To
avoid unintended photoactivation, the split luciferase must produce very low levels of light
emission at rest –when luciferin is present but intracellular calcium is low. Under these
conditions, the luciferase enzyme should remain incompletely reconstituted and emit minimal
light. Only upon calcium binding—or the presence of a specific co-factor needed for enzyme
reconstitution—should bioluminescence steeply increase to a level sufficient for activating
nearby photoreceptors. Comparison to other bioluminescent Ca^{2+} sensors revealed this critical
difference in SNR for the LMC construct used here to promote the BLADe paradigm.
Nonetheless, here we tested only a limited number of available sensors and there might be
suitable sensors among the already available ones. Furthermore, available sensors with
excellent performance in imaging applications should continue to be used for such but would
require further analysis if they were to be considered as drivers of optogenetic actuators.
496
BLADe depends on conditional light emission that stays *low* at baseline and *rises steeply* when
the target co-factor is present. LMC satisfied this criterion but is very dim and thus of limited
broad use, which prompted the systematic development of a novel Ca^{2+} dependent luciferase
with low baseline signal and high peak brightness, CaBLAM (Ca^{2+} BL Activity Monitor)[31].
Different from the *Gaussia* luciferase based LMC, CaBLAM was evolved from *Oplophorus*
gracilirostris luciferase [32–35]. Key differences to LMC in the CaBLAM sensor design are the
choice of a split in the loop that joins the final two β-strands for insertion of the Ca^{2+}-sensing
domains, a different M13-CaM pair, an N-terminal fluorescent partner for FRET enhancement
and restricting Ca^{2+} readout to the BL channel, and extensive affinity tuning that generated
variants covering much of the physiological range of Ca^{2+} concentrations. While LMC served to
demonstrate the BLADe concept and to reveal key aspects of BLADe, CaBLAM's superb
features position it as the Ca^{2+} indicator of choice for future iterations of BLADe, superseding
LMC. Moreover, the molecular evolution of CaBLAM incorporates a new luciferase scaffold,
SSLuc (Sensor Scaffold Luciferase)[31] that is highly amenable to insertion of sensor domains,
making it a flexible starting point for engineering an array of new BLADe indicators that couple
other biochemical events to light output for tailored optogenetic output.
513
In previous studies, two approaches have been used for integrating intracellular event sensing
with molecular outcomes by coupling bioluminescent light emission to downstream processes.
In the first, interaction between an intact luciferase and a light sensing protein is designed to be
contingent on an intracellular event, such as protein-protein interaction (SPARK2)[36], Ca^{2+}
increase (FLiCRE)[37], or GPCR activation (LOVTurbo)[38]. The second approach uses a two-
component split luciferase construct (LuCID)[39], where two halves of a split luciferase were fused
to two independent parts sensing Ca^{2+} (CaM and CaM binding protein). LuCID was also used to
couple protein-protein interaction to transcription, requiring new constructs that fuse the two
separate parts of the luciferase to two separate sequences detecting protein interaction[39]. The
above systems require multiple components, which may be a feature of a tool in some
applications but requires a lot of engineering for expanding its use to different contexts. In
contrast, BLADe is a highly modular tool that uses a single molecule containing both halves of
the luciferase split by a sensing moiety. Intracellular events that can be integrated with
optogenetic target effects are only limited by which sensors can be designed. And the same
bioluminescent sensor can be coupled with, in principle, any optogenetic actuator with matching
excitation wavelength.
530
These previous approaches of bioluminescent sensing and modulating, i.e. integrating, have
been applied to drive diverse cellular outcomes, including transcription of fluorescent reporters

to label cells that experienced particular intracellular events, the transcription of actuators to
allow manipulation of cells, and proximity labeling of nearby proteins. Bioluminescence has not
been used previously for coupling intracellular Ca^{2+} to opsin activation for changing the same
cell's membrane potential. However, a Ca^{2+} dependent luciferase has been used for synaptic
transmission of photons from an activated presynaptic neuron to a channelrhodopsin-
expressing postsynaptic neuron (PhAST)[40]. This application of bioluminescence-mediated cell-
to-cell communication supports the potential for using Ca^{2+} dependent bioluminescence
emission for activating photoreceptors, even over some distance without tethering.
541
Bioluminescence has been used for coupling intracellular acidification to opsin activation for
hyperpolarizing the cell[41]. Here, bioluminescence emitted from an intact luciferase was applied
to couple a sensor for neuronal hyperactivity, i.e. acidification, with a molecular actuator capable
of switching off neuronal activity in an all-molecular negative feedback loop at the single neuron
level[41]. To generate a pH sensitive inhibitory luminopsin (pHIL), three proteins were fused: a
hyperpolarizing opsin (eNpHR3,0), a pH sensing fluorescent protein (E2GFP), and a *Renilla*
luciferase. Through a two-step resonant energy transfer cascade triggered by acidification in the
presence of luciferin, pHIL translates the hyperactivity-induced intracellular acidosis into
silencing of neuronal activity. Cell-autonomous hyperpolarization was demonstrated *in vivo* by
mitigating acute tonic-clonic seizures induced by pilocarpine and by counteracting audiogenic
seizures in the PRoline-Rich Transmembrane protein 2 knockout (PRRT2 KO) mouse, a model
of genetic epilepsy. While pHIL is designed to specifically convert neuronal hyperactivity into
silencing, the BLADe approach includes and transcends this application of real-time feedback.
Co-expression of a Ca^{2+} BLADe integrator with an inhibitory or excitatory opsin results in neural
activity-driven hyperpolarization or additional depolarization, respectively.
557
The general motivation for developing closed-loop systems for modifying cells based on their
intracellular activity has led to a number of activity dependent regulatory approaches. The drive
for developing such approaches has been most prolific in neuroscience, challenged by the
question of how to identify and study the ensemble of active neurons underlying a specific
behavior and to therapeutically control selected subsets of overactive neurons. Some of the
current solutions are based on implanted hardware and computer-based detection algorithms to
conduct detection and deliver control. For example, signals are being tracked using electrodes,
and patterns detected by computer algorithms trigger countermanding stimulation through
implanted electrodes or fiber optics for electrical or optogenetic regulation[42–44]. However, these
methods require chronic implants to detect patterns and deliver stimuli, and electrical stimuli
impact all adjacent cells. The desire to replace problematic "externally closed-loop" systems by
"internal closed-loop sensor-actuators" working at the level of individual neurons motivated the
development of tools for all-molecular feedback regulation. Several approaches were developed
for coupling intracellular events to transcription of actuators using calcium sensitivity and
engineered TEV protease[45–47]. Cal-Light[45] and FLARE[46] sense Ca^{2+} and are temporally gated by
light exposure to drive transcription of actuators. Other systems, such as TRAP[48] and CANE[49],
use immediate early gene expression as a proxy for neuronal activity to drive transcription.
Refinements of these concepts have led to powerful preclinical tools, for example by using an
immediate early gene promoter to drive the expression of an engineered potassium channel in
hyperactive neurons for as long as they exhibit abnormal activity[50]. This activity-dependent gene
therapy was effective for on-demand cell-autonomous treatment for epilepsy in mice. In contrast
to sensor-actuator coupling via transcription, other approaches allow integration of biochemical
events with cellular outcomes in real time. Again, in the context of epilepsy therapy, biochemical
closed-loop treatments were achieved by employing pathologically elevated glutamate or
acetylcholine to activate engineered neurotransmitter-gated chloride and glycine channels,
respectively, to effectively control seizures in rodents[51,52].

584
Conceptually, the BLADe platform encompasses these above applications by allowing sensing
of fluctuations of any component for which a bioluminescent sensor can be designed and
integrating these with a variety of optogenetic actuators for a wide spectrum of cellular
outcomes. The BLADe design with a single-component sensor is very straightforward, different
from several of the above systems that require more complicated molecular designs, increasing
the potential for failure points in distinct cell types. Moreover, as the sensor can also be imaged,
BLADe enables simultaneous monitoring of sensor-actuator interactions. The need for luciferin
application places BLADe always under experimenter control. At the same time, it allows for
tuned effects through variation in luciferin concentrations, its route of administration with
different kinetics for intravenous versus intraperitoneal versus intraventricular application, use of
specific luciferins with short-term flash versus long-term glow kinetics, and timing of luciferin
administration throughout a behavioral paradigm.
597
Our study has important limitations. While our data are consistent with LMC integrating Ca^{2+}
fluctuations to drive transcription and change membrane potential at the functional level, i.e.
activity- and history-dependent coupling of cellular state change to output, we do not show a
causal biochemical mapping between intracellular Ca^{2+} and cellular output. Such formal proof
would require experiments employing Ca^{2+} chelators and Ca^{2+} titrations to establish a
quantifiable causal relationship between Ca^{2+} levels and optogenetic output. However, this
limitation does not negate the functional evidence from our *in vitro* and *in vivo* data that
demonstrate BLADe coupling past activity to future output in real time. In fact, we show
integration across two temporal regimes. First, in the opsin patch and *in vivo* sensory evoked
data, the β_3 interaction demonstrates that prior activity changes how subsequent activity is
generated within a single stimulus epoch—i.e., real-time, sub-second integration. Second, in the
transcriptional experiments the readout reflects prior cellular activity, as shown in both our HeLa
and mouse vSI experiments, where chemically enhancing intracellular activity (e.g., histamine or
NMDA) leads to produce corresponding changes in gene expression. Nevertheless, the above
biochemical studies should be done with the next generation of suitable Ca^{2+} sensors. Another
set of experiments should be a titration of Ca^{2+} sensor expression as overexpression of
luciferase-based sensors may lead to nonlinear signal behavior. While LMC served to establish
the design requirements for Ca^{2+} sensors to serve as integrators and allowed to demonstrate
applications of the BLADe approach, its light emission is very dim. For going forward, a new
Ca^{2+} sensor suitable for BLADe applications superseding LMC has recently been developed[31].
In future work, the BLADe platform could be improved and extended in a number of ways. The
luciferase can be split by different sensing domains whose light activation can be coupled to a
wide range of light-sensing actuators, providing experimental control over diverse molecular
outcomes depending on the user's goal. Further developed and improved versions of these
tools are expected to have broad applications. In basic research, this paradigm allows
investigators to selectively detect, and then suppress or amplify, specific biological events, a
unique approach to understanding the role of those processes. In potential therapeutic
applications it allows sensing and discontinuing aberrant activity before it can cause harm, or
stopping pathological processes from continuing, or amplifying and rescuing failing processes,
without the use of implanted devices and regulated by simple peripheral injection of a luciferin.
628
629 **METHODS**
630
631 **Chemicals**
632 All chemicals, including Ionomycin and Histamine, were purchased from Sigma. The Calcium
633 Calibration Buffer Kit #1 from Life Technologies was used to prepare Zero Ca^{2+} buffer and 39
634 µM Ca^{2+} buffer.

635
Luciferins
For *in vitro* experiments, Coelenterazine (CTZ) (NanoLight Technology; Pinetop, AZ; NanoLight
303) and Coelenterazine h (hCTZ) (NanoLight # 301) were dissolved in NanoFuel solvent
(NanoLight # 399) and stored in 50 mM stock solutions at -80 ºC. CTZ, hCTZ, or NanoFuel
Solvent (for vehicle controls) were freshly diluted in culture medium for final concentrations. D-
luciferin potassium salt was purchased from GoldBio (USA), dissolved in water and stored as 60
mg/ml stock aliquots at -80°C. For use in plate readings, stock solutions were diluted in
OptiMEM, then further diluted into wells for 150 µg/ml final concentration. For *in vivo*
experiments, water soluble Coelenterazine (Nanolight #3031) was freshly prepared by
dissolving in sterile water to a stock concentration of 1.6 mM for lateral ventricle injection to yield
a final concentration of 200 µM and to a stock concentration of 2.36 mM for dilution in ACSF to
yield a final concentration of 20 µM.
648
Plasmids
A pcDNA3.1 backbone with the CMV promoter was used to insert gBlocks (IDT) that encode for
N-sbGluc, CaM-M13 variants, C-sbGluc and p2a dTomato by Gibson cloning (New England
Biolabs HiFi DNA Master Mix). Coding sequences from plasmids for GCaMP6f, hGtACR2,
ChR2(C128S), and CheRiff (Addgene) and for BlueCaMBI and GLICO (synthesized by
Genscript) were cloned into the pcDNA3.1-CMV backbone. Expression plasmids for VP-EL222
and 5xC120-FireflyLuc were kindly provided by Dr. Kevin Gardner, CUNY, New York, NY, and
for GeNL(Ca2+) by Dr. Takeharu Nagai (Addgene). For viral vectors, coding sequences were
cloned into a pAAV vector downstream of a CAG or hSyn promoter, or into a pAAV-EfIa-DIO
construct. For list of plasmids used see Supplementary Table 1.
659
pcDNA3.1-CAG-VP-EL222 and pcDNA3.1-5xC120-FireflyLuc were kindly provided by Dr. Kevin
Gardner, CUNY, New York, NY
pAAV.Syn.GCaMP6f.WPRE.SV40 was a gift from Douglas Kim & GENIE Project (Addgene
plasmid # 100837 ; http://n2t.net/addgene:100837 ; RRID:Addgene_100837)
GeNL(Ca²⁺)_520/pcDNA3 was a gift from Takeharu Nagai (Addgene plasmid # 85204 ;
http://n2t.net/addgene:85204 ; RRID:Addgene_85204)
pLenti-CaMKIIa-hChR2(C128S)-EYFP-WPRE was a gift from Karl Deisseroth (Addgene
plasmid # 20294 ; http://n2t.net/addgene:20294 ; RRID:Addgene_20294)
pFUGW-hGtACR2-EYFP was a gift from John Spudich (Addgene plasmid # 67877 ;
http://n2t.net/addgene:67877 ; RRID:Addgene_67877)
pAAV-hSyn-CheRiff-eGFP was a gift from Adam Cohen (Addgene plasmid # 51697 ;
http://n2t.net/addgene:51697 ; RRID:Addgene_51697)
672
AAV
AAV2/9 preparations were generated by triple lipofection of HEK293-FT cells and harvesting
viral particles as previously described[5]. Briefly, triple plasmid transfection in HEK293 cells
(Agilent, Cat # 240073-41) was performed. The helper plasmid pAd delta F6 (24 µg), respective
pAAV-hSyn plasmids (12 µg), and the serotype plasmid AAV2/9 (20 µg) were transfected, using
Lipofectamine 2000, into subconfluent HEK293 cells grown in 10 cm culture dishes. Virus was
purified from supernatant and cells with an aqueous two-phase system after 72 hrs, dialyzed at
4°C overnight against PBS (Ca2+ and Mg2+ free). The next morning, virus was concentrated in
Amicon Ultra-0.5 mL Centrifugal Filters. Q-PCR targeting the WPRE element was used to
determine viral titers.
683
Animals

All mouse experiments described in this study were approved by the Central Michigan
University Institutional Animal Care and Use Committee (IACUC) and conducted in accordance
with all relevant guidelines and regulations, including the US National Research Council's Guide
for the Care and Use of Laboratory Animals, the US Public Health Service's Policy on Humane
Care and Use of Laboratory Animals, the Guide for the Care and Use of Laboratory Animals,
and the ARRIVE guidelines (Animal Research: Reporting of In Vivo Experiments, PLoS Bio
8(6), e1000412,2010) for reporting animal research. Mice were group-housed in ventilated
cages under 12-hour reverse light cycle, provided with tap water and standard chow and
allowed to feed ad libitum. C57BL/6 (JAX #000664), Emx1-Cre (JAX# 005628), and Swiss
Webster (Charles River) mice of both sexes were used. Mice were euthanized by CO_2
asphyxiation followed by cervical dislocation in accordance with the above guidelines for animal
euthanasia.
697
698
Cell Culture and Cell Transformation
To avoid any ambient light contamination, all tissue culture work was carried out in a dark room
with a red light source; this room also housed the cell culture incubator. Human embryonic
kidney fibroblasts (HEK293, RRID:CVCL_0045) and human cervical adenocarcinoma cells
(HeLa, RRID:CVCL_0030) were purchased from the American Type Culture Collection
(HEK293: ATCC, CRL-1573, Lot #70039815; HeLa: ATCC, CCL-2, Lot #70037076), expanded,
and aliquots at low passage numbers were frozen. Cells were grown in Gibco DMEM media
supplemented with 10% Fetal Bovine Serum, 1% Glutamax, 0.5 % Pen-Strep, 1% Non-
Essential Amino Acids and 1% sodium pyruvate. Cells were cultured at 37°C and 5%
atmospheric carbon dioxide. Cells were transfected with Lipofectamine 2000 or 3000 in
accordance with the manufacturer's instructions.
710
Primary E18 rat cortical and hippocampal neurons were prepared from tissue shipped from
BrainBits (Transnetyx) following the vendor's protocol. Neurons were grown in Gibco
Neurobasal Media supplemented with 2% B27 supplement and 0.1% gentamycin and 1%
Glutamax. Neurons were nucleofected with plasmids of interest using the Lonza 2b
Nucleofector and Rat Neuron Nucleofector Kit (Lonza, VPG-1003) according to the
manufacturer's instruction, then seeded. Or neurons were nucleofected, seeded, then
transduced with AAV the next day. Neurons were seeded on PDL-coated glass coverslip
(Neuvitro) in 12-well tissue culture plates (1 x 10^5 neurons per well) or were plated on the
electrode area of 1-well MEA dishes (60MEA200/30iR-Ti; Multi Channel Systems, Germany)
coated with PEI (0.1%) and laminin (50 μg/ml) ($1x10^5$ neurons/10 μL/well) as described in detail
in Prakash 2020[26].
722
Primary cardiomyocytes were harvested from E18 Swiss Webster mice, papain digested and
plated on 35mm glass bottom dishes (MatTek) for real time bioluminescence imaging. Primary
cardiomyocytes were grown in Claycomb Media supplemented with 10% HL-1 screened FBS,
100 μg/mL penicillin/streptomycin, 0.1mM norepinephrine and 2% Glutamax. Cells were
nucleofected with plasmids of interest using the Lonza 2b Nucleofector and Rat cardiomyocyte
Nucleofector Kit (Lonza, VAPE-1002) according to the manufacturer's instruction.
729
730　**Microscopy**
731　Initially, all transfected cells were imaged using a Zeiss Axio Observer A1 microscope with a LD
732　A-Plan 20x/0 air objective and a Hamamatsu Orca Flash 4.0 CMOS camera to confirm
733　expression by fluorescence microscopy at 50 ms exposure. Imaging of brain sections were
734　done on an Olympus Fluoview 300 CLSM confocal microscope with a 60x/1.3 Oil objective.
735

Bioluminescence Imaging
For photon counting HEK293 and HeLa cells were seeded on Costar 96 well clear bottom white plates and luminescence was measured in a SpectraMax® L luminometer (Molecular DevicesTM). For real-time bioluminescence imaging we used a Zeiss Axio Observer A1 microscope with a Fluar 40x/1.3 Oil objective with either a Hamamatsu Orca Flash 4.0 CMOS camera or an Andor iXon Ultra 888 EM-CCD camera with their respective softwares. HEK293 and HeLa cells were imaged in imaging solution ($CaCl_2$ 1.25 mM, HEPES 19.7 mM, KCl 4.7 mM, KH_2PO_4 1.2 mM, $MgSO_4$ 1 mM, NaCl 130 mM, dextrose 0.5, pH 7.2-7.4) with either 100 µM or 12.5 µM CTZ. Primary cardiomyocytes were imaged in Tyrode's solution (NaCl 137 mM, KCl 2.7 mM, $MgCl_2$ 1 mM, $CaCl_2$ 1.8 mM, Na_2HPO_4 0.2 mM, $NaHCO_3$ 12 mM, D-glucose 5.5 mM, pH 7.2-7.4) supplemented with 10 µM norepinephrine and 100 µM CTZ. Primary E18 rat hippocampal neurons were imaged in artificial cerebrospinal fluid, containing: 121 mM NaCl, 1.25 mM NaH_2PO_4, 26 mM $NaHCO_3$, 2.8 mM KCl, 15 mM d(+)-glucose, 2 mM $CaCl_2$, 2 mM $MgCl_2$ (maintained at ~37 ºC, pH 7.3-7.4, continuously bubbled with 95% O_2/5% CO_2 vol/vol). All imaging was conducted in the RC-49MFSH heated perfusion chamber where the temperature was continuously monitored (Warner Instruments).

Electrophysiology recordings from primary neurons
Whole Cell Recordings
Neurons (rat E18, cortical) were co-nucleofected with LMC4f and either an inhibitory opsin, hGtACR2, an excitatory opsin, ChR2(C128S), or a "Dud" opsin, ChR2(C128S)-E97R-D253A. Post nucleofection, neurons were plated on laminin-coated glass coverslips (15mm, Neuvitro) in culture medium consisting of Neurobasal Medium (Gibco # 21103-049), B-27 supplement (Gibco # 17504-044), 2 mM Glutamax (Gibco # 35050-061), and 5% Fetal Calf Serum (FCS). The following day, the medium was replaced with serum-free medium (NB-Plain medium). Half of the medium was replaced with fresh NB-Plain medium every 3–4 days thereafter. Neurons were used for whole cell patch clamp recordings between DIVs 21–25. For patch clamp recording, a coverslip was transferred to a recording chamber mounted on an upright microscope (BX51WI, Olympus) and perfused with aCSF containing (in mM): 121 NaCl, 2.8 KCl, 1 NaH_2PO_4, 26 $NaHCO_3$, 2 $CaCl_2$, 2 $MgCl_2$ and 15 D-glucose (310 mOsm/kg, pH 7.3-7.4) at a rate of 1.5 ml/min. All solutions were bubbled with a gas mixture of 95% O_2 and 5% CO_2. Whole-cell patch clamp recordings were performed using a Multiclamp 700b amplifier and Digidata 1440 digitizer together with the pClamp recording software (Molecular Devices). Borosilicate glass micropipettes were manufactured using a PC-100 puller (Narishige) and had resistances of 3–5 MΩ. In current clamp recordings, pipettes were filled with intracellular solution containing (in mM): 130 K-gluconate, 10 KCl, 15 HEPES, 5 Na_2-phosphocreatine, 4 Mg-ATP and 0.3 Na-GTP (310 mOsm/kg, pH 7.3). The aCSF was supplemented with D-AP5 (50 µM), CNQX (15 µM) and picrotoxin (100 µM) to block fast glutamatergic and GABAergic synaptic transmission. Rs was compensated by using bridge balance. Firing frequencies were calculated from the total number of action potentials produced per unit time during depolarizing current injections (10 sweeps, 1.5 sec duration) in episodic stimulation acquisition mode, at a membrane potential of -70mV, before and after CTZ (100 µM) or vehicle treatments.

To quantify activity-dependent changes in spike generation during sustained depolarization, we computed a spike count fractional difference for each trial. For each 1-s depolarizing current injection, spikes were binned into the first and second halves of the stimulus (0–0.5 s and 0.5–1.0 s). The number of spikes in the second half was subtracted from the number of spikes in the first half and normalized by the total spike count for that trial. This metric ranges from +1, indicating all spikes occurred in the first half of the stimulus, to −1, indicating all spikes occurred in the second half, with 0 reflecting an even distribution across the stimulus. This analysis was applied on a per-trial basis before and after CTZ or vehicle treatment to assess how LMC-

787 dependent opsin activation altered the temporal distribution of action potentials during
788 depolarization.
789

MEA Recordings
Once the neurons (rat E18, cortical) were matured (DIV14-DIV19), recordings were conducted
using a MEA2100-Lite-System (Multichannel Systems, Germany). As described in detail in
Prakash et al.[26], raw extracellular signals were acquired via the MCS system and stored in
HDF5 format. All-trans retinal (R2500; Sigma- Aldrich, St. Louis, MO) was added to the culture
medium to 1 µM final concentration before electrophysiological recordings. Prior to recording, all
reagents were pre-warmed to 37 °C. A micropipette was used to add reagents with the reagent
drop gently touching the liquid surface, creating a time-locked artifact in the recordings.
Recordings were carried out with a sample rate of 10 kHz. Raw samples were read directly from
the HDF5 files and analyzed in native ADC units. Conversion to physical units was defined by
MCS channel metadata (ADZero, ConversionFactor, Exponent) according to the manufacturer's
specification; for the recordings analyzed here, this corresponded to a scaling of 0.0596 µV per
ADC count.
803

For visualization and spike detection, voltage traces were temporally aligned to the timestamp of
chemical application (CTZ or vehicle). This timestamp was identified by a large, stereotyped
amplitude artifact produced by direct addition of solution to the MEA which provided a within
well experiment-wide temporal reference. A symmetric time window (±1000 ms) centered on
this chemical event was extracted, defining pre- and post-application analysis periods.
Chemically aligned voltage traces were high-pass filtered to isolate spiking activity from slower
components of the extracellular signal (e.g., baseline drift and low-frequency fluctuations).
Filtering was performed using a 4th-order Butterworth high-pass filter with a 300 Hz cutoff,
applied with zero-phase forward–reverse filtering (filtfilt) to preserve spike timing.
813

Detection thresholds were defined as five times the baseline noise level, where baseline noise
was estimated from the pre-chemical window using the median absolute deviation (MAD).In
brief, noise was computed as $1.4826 \times \text{median}(|yb - \text{median}(yb)|)$, where yb denotes the filtered
baseline signal for each channel, and the detection threshold was set to 5.0 × noise. Negative
spikes were identified as threshold-crossing deflections in the filtered extracellular signal using
peak detection. Detected events were constrained by a refractory period of 1.0 ms and a
minimum spike width of 0.3 ms; events occurring within 0.3 ms of one another were merged to
prevent double counting. Only spikes occurring within the post-chemical window were retained
for downstream analyses and were treated as multi-unit activity (MUA). Similar to *in vivo*
recordings, MUA was estimated by converting raw spike times into instantaneous firing rates
using 1 ms bins.
825

Transcription *in vitro*
HeLa cells were used to test whether LMC-driven light emission could activate transcription
through the light-responsive transcription factor EL222. Cells were plated in 6-well plates and
transfected at ~80–90% confluency using Lipofectamine 2000 (Thermo Fisher). Each well
received a total of 2 µg DNA, composed of the following: 666 ng EL222 (light-sensitive
transcription factor), 333 ng 5×C120-FLuc (EL222-responsive firefly luciferase reporter), and
1000 ng LMC4f-dTomato (calcium-dependent luciferase). Parallel wells were transfected with
other calcium-dependent bioluminescent sensors (GeNL, GLICO, BlueCaMBI) and matched
FLuc reporters.
835

Three hours after transfection, cells were trypsinized (300 µL trypsin per well) and replated into
PDL-coated, white-walled, clear-bottom 96-well plates (~100,000 cells per well). Cells were

allowed to adhere for 9–12 hours before stimulation. For transcriptional activation, 100 µL of
stimulation buffer was added per well (1:1 with cell media) for a final well volume of 200 µL. The
stimulation solution contained 100 µM CTZ or hCTZ (depending on the sensor), with or without
10 µM histamine. Vehicle controls received no treatment. After stimulation, cells were incubated
for 7–9 hours before transcriptional reporter readout.
843
For FLuc transcriptional measurement, activity was recorded 18 hours after stimulation using a
luminometer. Prior to measurement, cells were washed and incubated in phenol red–free media
(100 µL per well). D-luciferin (final concentration 150 µg/mL) was prepared from a frozen
60 mg/mL stock by serial dilution into Opti-MEM. For each well, 25 µL of 750 µg/mL D-luciferin
working stock was injected into 100 µL of media. FLuc bioluminescence was recorded
immediately after injection.
850
Transcription *in vivo*
To test bioluminescent dependent transcription *in vivo*, we co-injected three AAV constructs into
the prefrontal cortex of anesthetized mice (isoflurane at a rate maintained between 2 – 3% and
oxygen at 55 mL/min; SomnoSuite Low-Flow Anesthesia System, Kent Scientific Corporation):
AAV9-hSyn-LMC4f-P2A-dTomato (Ca^{2+}-sensitive luciferase), AAV9-CAG-EL222 (light-activated
transcription factor), and AAV9-5×C120-EYFP (EL222-responsive reporter). Viral solutions were
mixed prior to injection to a total volume of 1 µL per site, yielding final per-virus titers of: LMC4f
(1×10^{10} GC), EL222 (0.7×10^{10} GC), and C120-EYFP (0.3×10^{10} GC). The injection was
targeted to the left medial prefrontal cortex at +0.37 mm anterior to bregma, 3.0 mm lateral.
Following injection, the craniotomy was sealed to minimize ambient light exposure during
recovery.
862
At the time mice were expressing light-sensing constructs, experiments were conducted in a
dark room illuminated with red light. After 7 days, mice received intraventricular (ICV) injections
of one of three treatments: (1) NMDA + vehicle, (2) luciferin (CTZ) alone, or (3) luciferin +
NMDA. These conditions were used to assess background transcription, spontaneous activity-
driven transcription, and full BLADe-dependent transcriptional gating, respectively. Injections
were performed using a 26-gauge needle and 10 µL Hamilton syringe, targeting the right lateral
ventricle (AP: –0.5 mm, ML: –1.1 mm, DV: –2.0 mm from bregma). The luciferin was prepared
fresh by dissolving coelenterazine (NanoLight #3031) in sterile water to a stock concentration of
1.6 mM, then when injected was assumed to be diluted 1:8 in CSF within the ventricle to yield a
final injection concentration of 200 µM. NMDA was prepared at a working concentration of
75 ng/µL in sterile water. For combined delivery, 1 µL of NMDA solution (75 ng) was mixed with
4 µL of luciferin solution per injection (final 5 µL per mouse, injected over 5 mins). Solutions
were prepared under red light and protected from exposure throughout. After injection, mice
were removed from the stereotaxic apparatus and allowed to recover. Brains were collected for
histological analysis ~20 hours after ICV injection. Mice were anesthetized with isoflurane and
transcardially perfused with PBS followed by 4% paraformaldehyde. Coronal brain sections
were collected and imaged by confocal microscopy to quantify EYFP reporter expression.
Confocal imaging of EL222 was performed using a rabbit anti-VP16 tag primary antibody
(Abcam, ab4808), followed by a far-red Alexa Fluor 647-conjugated goat anti-rabbit secondary
antibody (Invitrogen, A-21245) to minimize spectral overlap with dTomato fluorescence from
LMC.
884
To analyze calcium- and light-dependent transcriptional activation *in vivo*, we quantified EYFP
reporter expression in dTomato-labeled neurons across experimental groups, with the individual
scoring sections blinded to experimental groups. Confocal images from all treatment conditions
(NMDA-only, luciferin-only, luciferin + NMDA) were analyzed. First, dTomato+ images were

pooled and segmented from maximum intensity projections using the MATLAB 'cell-segm'
thresholding and size-filtering algorithm[53], ensuring unbiased detection across all fields of view.
The resulting binary masks were then applied to the corresponding EYFP images to extract raw
per-cell reporter fluorescence from each dTomato+ neuron. Finally, cells were grouped by
treatment, and the ratio of EYFP to dTomato fluorescence was computed per cell to normalize
for expression variability.
895
Electrophysiology *in vivo*
To express LMC and CheRiff in excitatory cortical neurons, we injected AAVs into the left
vibrissa primary somatosensory cortex (vSI) of Emx1-Cre mice (3–4 month old). AAV1-EF1α-
DIO-LMC4f and AAV1-EF1α-DIO-CheRiff-EYFP were mixed 1:1 and delivered in a single
injection (1 µL total) using a 33-gauge Nanofil syringe. Using a dental drill, a burr hole was made
for AAV injection. The injection site was located 1.5 mm posterior and 3.0 mm lateral to bregma,
and the needle was lowered to a depth of 500–1000 µm below the pial surface. Virus was
infused at a rate of 50 nL/min over ~20 minutes. Following injection, the needle was held in
place for 10 min, then raised to 200 µm and held for an additional 10 min before full withdrawal
to allow the virus to spread across the layers of the cortex. During the same surgery, a custom
metal headpost was implanted for head fixation. After exposing the skull, a thin layer of clear
Metabond was applied to stabilize the surface while keeping bregma and surface vasculature
visible. The headpost was then affixed using additional Metabond, positioned off-center to leave
vSI accessible. The short arm was angled over the left ear to allow right-side whisker
stimulation. After injection and headpost implantation, the craniotomy was covered with mineral
oil and sealed with Metabond. Animals recovered for at least 3 weeks before recordings.
912
After 2–3 weeks of recovery, mice were anesthetized with isoflurane (1.2–1.4%) and head-fixed.
Again, these experiments were conducted in a dark room illuminated with red light. The
craniotomy used for AAV injection was reopened and expanded to 1 mm to expose the
underlying cortex and target the same vSI region as for the viral delivery. Throughout the
procedure, the brain remained covered in carbogenated ACSF (95% O_2 / 5% CO_2) under
constant perfusion. Animals were placed in a light-tight, electrically shielded chamber and
maintained under anesthesia. Mice were secured in place to allow perfusion tubing to be
positioned next to the craniotomy, allowing for constant ACSF inflow (for bioluminescence CTZ
was diluted to 20 µM in ACSF). All macrovibrissae on the right side of the mouse were secured
~3mm from the mystacial pad. A custom-made clamp was glued to a piezoelectric bender
(Noliac CMBP09), positioned to move all macrovibrissae in the caudorostral direction, with a
half-sine wave velocity profile with a rising phase (6ms) and a slower relaxation phase (20ms).
On each trial, 20 Hz vibratory vibrissal stimulus trains (10 deflections, 500 ms) were delivered
for 1000 trials total. On a trial by trial and randomized basis, the stimulus amplitude varied
between 0 to maximal amplitude (~1mm deflection). For each recording session, ACSF was
constantly perfused for the first 500 trials and luciferin (CTZ) was then perfused for the
remaining 500 trials.
930
All recordings in barrel cortex were conducted under light isoflurane anesthesia. Laminar probes
consisted of single shank,32-channel silicon probes with a fiber optic 50 µm above the highest
recording site (A1x32 Poly2-5mm-50s-177-OA32LP, Neuronexus Technologies; 0.15mm silver
wire reference). Data was sampled at 30 kHz and passed through a digital amplifier (Cereplex-
µ, Blackrock Microsystems), and directed through HDMI to the Cereplex Direct data acquisition
box (Blackrock Microsystems). The 32-channel linear silicon probe was inserted perpendicularly
into cortex using a micromanipulator at ~10 µm/s until the top contact of the electrode
disappeared beneath the pial surface. The probe was allowed to settle for ~30–60 minutes
before recording. Luciferin (coelenterazine; NanoLight #3031) was freshly prepared in sterile

water (1 µg/mL) and diluted into ACSF to a final concentration of 20 µM. Following an initial
recording period under vehicle ACSF, the perfusion line was switched to CTZ-containing ACSF
to enable continuous bath application during the second half of the session. Flow rate and
volume were monitored to maintain complete coverage of the craniotomy and probe without
overflow.
945
Raw electrophysiological data were downsampled to 10 kHz. For each recording, electrode
contacts with RMS values exceeding three times the interquartile range above the 75th
percentile or below the 25th percentile across the 32-channel array were flagged as noise and
removed for further analysis. All remaining electrodes were re-referenced to the common
average[54]. MUA was estimated by converting raw spike times into instantaneous firing rates
using 1 ms bins. For luciferin experiments, stimulus-aligned firing rates were extracted from
trials before and after the defined luciferin application time.

For sensory stimulation experiments, MUA peristimulus time histograms (PSTHs) were
constructed by aligning spike data to stimulus onset and binning at 1 ms. PSTHs were
smoothed using a Gaussian kernel (window = 20 ms, SD = 3 ms) and averaged across trials.
PSTHs were computed separately for pre- and post-luciferin trial blocks. Optogenetic responses
to LED stimulation were quantified using a pre-stimulus baseline window (−500 to 0 ms) and a
post-stimulus response window (0 to 500 ms). For each channel, the change in firing rate was
defined as the difference between mean firing during the response window and baseline. A null
distribution was constructed from no-opsin control recordings and used to define response
thresholds. CTZ-evoked responses were quantified using the same baseline window by
computing the change in firing rate between post-CTZ and pre-CTZ periods for each channel. A
corresponding null distribution was generated from no-opsin control recordings using the same
procedure and used for channel classification.

Quantification of BLADe-dependent modulation of sensory-evoked activity
To isolate BLADe-dependent modulation of sensory-evoked cortical activity, whisker-evoked
multiunit activity (MUA) was quantified on a trial-by-trial basis using a baseline-corrected metric.
For each recording channel, peri-stimulus time histograms (PSTHs) were generated by time-
locking spike times to stimulus onset within a 1500 ms window, pooling across stimulus
amplitudes, and converting spike counts to firing rate (Hz) by scaling by the bin width. The
alignment set stimulus onset to 0 ms across PSTHs and was used for visualization.

For each recording channel, spike trains were first aggregated across trials to compute firing
rates aligned to stimulus onset. The mean firing rate during a 500 ms post-stimulus analysis
window (500–1000 ms after stimulus onset) was then computed and baseline-corrected by
subtracting the mean firing rate during the 500 ms immediately preceding stimulus onset (−500
to 0 ms) from the same channel. This channel-wise baseline subtraction normalizes evoked
responses relative to ongoing activity within each recording channel, removing contributions
from spontaneous firing, slow temporal drift, and global CTZ-induced changes in baseline
excitability that are not time-locked to the sensory stimulus.

For each recording channel, CTZ-dependent modulation of sensory-evoked responses was
quantified by computing the difference between baseline-corrected evoked firing rates
measured during CTZ trials and those measured during vehicle trials. This within-recording
subtraction isolates changes in stimulus-evoked activity attributable to CTZ within each channel
after normalization to baseline firing. Therefore, the resulting channel-wise CTZ–vehicle

difference served as the primary quantitative measure used to compare CTZ-dependent
modulation of sensory-evoked activity between opsin-expressing and non-opsin control mice.
987

988   **Analysis of feed-forward effects in BLADe**
989   For the linear regression, late evoked activity was modeled as a linear function of early evoked
990   spiking activity and luciferin condition according to:
991

992
$$r_{\text{late}} = \beta_0 + \beta_1 \cdot r_{\text{early}} + \beta_2 \cdot \text{Luc} + \beta_3 \cdot \left(\text{Luc} \times r_{\text{early}} \right)$$

Here, (r_{late}) is the late evoked firing rate measured in the late window (>200 ms after stimulation
onset) while (r_{early}) is the early evoked firing rate measured in the 0–200 ms window following
stimulation onset. Luc indicates the absence (Luc = 0; vehicle) or presence (Luc = 1; luciferin,
CTZ). β_0 and β_1 define the intercept and slope relating early to late activity in the absence of
luciferin, whereas β_2 captures a luciferin-dependent shift in the intercept (offset) and β3 captures
a luciferin-dependent change in the slope relating early and late activity (gain). Early and late
responses were pooled across trials, with regression performed at the level of individual
neurons for patch-clamp recordings and at the level of individual recording channels for *in vivo*
recordings. In this model, β_2 reflects a tonic, activity-independent shift in firing during the late
phase window. As the model includes a tonic term (β_2), the interaction term (β_3) captures
changes in the slope of the early–late relationship after accounting for activity-independent
shifts. Thus, β_3 represents a luciferin-dependent change in the relationship between early and
late spiking (gain). A positive β_2 indicates a tonic increase in late firing independent of early
activity, whereas a negative value indicates tonic suppression. For β_3, positive values reflect
multiplicative gain by strengthening the early–late relationship, whereas negative values reflect
divisive neuromodulation.
1010

1011   **Statistics**
1012   Preprocessing and analysis of the data was performed with custom-written Matlab (R2020a,
1013   Mathworks) and Python scripts.
1014

In **Fig. 1m**, maximal LMC luminescence responses (Max $\Delta RLU/RLU_0$) with CTZ alone and
glutamate+CTZ were compared using paired Wilcoxon signed-rank tests within the same
neurons (same neuron, CTZ vs CTZ+glutamate). In **Fig. 2c**, histamine-specific changes in
bioluminescence normalized to the luciferin baseline were compared across calcium sensors
using a one-way ANOVA followed by Tukey's post hoc multiple-comparisons test as an initial
exploratory characterization of calcium-dependent luminescence responses. In **Fig. 3c**, overall
differences across sensors were assessed using a Kruskal–Wallis test (non-parametric
ANOVA), followed by targeted Wilcoxon rank-sum tests for hypothesis-driven comparisons
involving LMC. In **Fig. 3g**, EYFP reporter/dTomato expression ratios from three independent
conditions were compared using Wilcoxon rank-sum tests applied to all pairwise group
comparisons, with Bonferroni correction for multiple testing (CTZ+NMDA vs CTZ vs NMDA). In
Fig. S3 and S4, cumulative distributions of spike count fractional differences between the first
and second halves of depolarizing current injections were compared using two-sample
Kolmogorov–Smirnov tests across vehicle and CTZ perfusion conditions. In **Fig. S5,** luciferin-
dependent changes in multiunit activity (MUA) were compared between inhibitory-opsin
recordings with and without LMC co-expression using an unpaired Wilcoxon rank-sum test. In
Fig. 4b–d(iii), luciferin and vehicle action potential counts were compared using paired
Wilcoxon signed-rank tests, with measurements paired within the same neurons across
depolarizing steps under vehicle and CTZ conditions (same neuron, vehicle vs CTZ OR vehicle
vs vehicle). In **Fig. 4b–d(iv)**, early and late spike counts were related using linear regression

performed within individual neurons under vehicle and CTZ conditions. In **Fig. 5**, CTZ and vehicle responses were compared using paired Wilcoxon signed-rank tests, with each data pair corresponding to measurements from the same electrode recorded under vehicle and CTZ conditions at different time points on the same day (same electrodes, vehicle vs CTZ). In **Fig. 6 c,d, and f**, evoked firing changes in opsin-expressing and no-opsin channels were compared using an unpaired Wilcoxon rank-sum test. In **Fig. 6 g and h,** early and late evoked activity were related using linear regression performed at the level of individual recording channels under vehicle and luciferin conditions.

REFERENCES

1. Love, A. C. & Prescher, J. A. Seeing (and Using) the Light: Recent Developments in Bioluminescence Technology. *Cell Chemical Biology* **27**, 904–920 (2020).
2. Rost, B. R., Schneider-Warme, F., Schmitz, D. & Hegemann, P. Optogenetic Tools for Subcellular Applications in Neuroscience. *Neuron* **96**, 572–603 (2017).
3. Losi, A., Gardner, K. H. & Moglich, A. Blue-Light Receptors for Optogenetics. *Chem. Rev.* **118**, 10659–10709 (2018).
4. Berglund, K., Birkner, E., Augustine, G. J. & Hochgeschwender, U. Light-Emitting Channelrhodopsins for Combined Optogenetic and Chemical-Genetic Control of Neurons. *PLoS One* **8**, e59759 (2013).
5. Berglund, K. *et al.* Luminopsins integrate opto- and chemogenetics by using physical and biological light sources for opsin activation. *Proc. Natl. Acad. Sci. U. S. A.* **113**, E358–E367 (2016).
6. Tung, J. K., Gutekunst, C.-A. & Gross, R. E. Inhibitory luminopsins: genetically-encoded bioluminescent opsins for versatile, scalable, and hardware-independent optogenetic inhibition. *Sci. Rep.* **5**, 14366 (2015).
7. Crespo, E. L., Bjorefeldt, A., Prakash, M. & Hochgeschwender, U. Bioluminescent Optogenetics 2.0: Harnessing Bioluminescence to Activate Photosensory Proteins In Vitro and In Vivo. *J. Vis. Exp.* **174**; doi:10.3791/62850 (2021).
8. Li, T. *et al.* A synthetic BRET-based optogenetic device for pulsatile transgene expression enabling glucose homeostasis in mice. *Nat. Commun.* **12**, 615; 10.1038/s41467-021-20913-1 (2021).
9. Parag-Sharma, K. *et al.* Engineered BRET-Based Biologic Light Sources Enable Spatiotemporal Control over Diverse Optogenetic Systems. *ACS Synth. Biol.* **9**, 1-9 (2020).
10. Naim, N. *et al.* Luminescence-activated nucleotide cyclase regulates spatial and temporal cAMP synthesis. *J. Biol. Chem.* **294**, 1095-1103 (2019).
11. Kim, E. H. *et al.* Self-luminescent photodynamic therapy using breast cancer targeted proteins. *Sci. Adv.* **6**, eaba3009; 10.1126/sciadv.aba3009 (2020).
12. Shramova, E. I., Frolova, A. Y., Filimonova, V. P., Deyev, S. M. & Proshkina, G. M. System for Self-excited Targeted Photodynamic Therapy Based on the Multimodal Protein DARP-NanoLuc-SOPP3. *Acta Naturae* **15**, 100–110 (2023).
13. Yeh, H.-W. & Ai, H.-W. Development and Applications of Bioluminescent and Chemiluminescent Reporters and Biosensors. *Annu. Rev. Anal. Chem.* **12**, 129–150 (2019).
14. Welsh, J. P., Patel, K. G., Manthiram, K. & Swartz, J. R. Multiply mutated Gaussia luciferases provide prolonged and intense bioluminescence. *Biochem. Biophys. Res. Commun.* **389**, 563–8 (2009).
15. Kim, S. B., Sato, M. & Tao, H. Split Gaussia luciferase-based bioluminescence template for tracing protein dynamics in living cells. *Anal. Chem.* **81**, 67–74 (2009).
16. Saito, K. *et al.* Luminescent proteins for high-speed single-cell and whole-body imaging. *Nat. Commun.* **3**, 1262; 1038/ncomms2248 (2012).

17.   Chen, T.-W. *et al.* Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature*
499, 295–300 (2013).
18.   Oh, Y. *et al.* An orange calcium-modulated bioluminescent indicator for non-invasive
activity imaging. *Nat. Chem. Biol.* **15**, 433–436 (2019).
19.   Farhana, I., Hossain, M. N., Suzuki, K., Matsuda, T. & Nagai, T. Genetically Encoded
Fluorescence/Bioluminescence Bimodal Indicators for Ca(2+) Imaging. *ACS sensors* **4**,
1825–1834 (2019).
20.   Suzuki, K. *et al.* Five colour variants of bright luminescent protein for real-time multicolour
bioimaging. *Nat. Commun.* **7**, 13718; 10.1038/ncomms13718 (2016).
21.   Motta-Mena, L. B. *et al.* An optogenetic gene expression system with rapid activation and
deactivation kinetics. *Nat. Chem. Biol.* **10**, 196-202 (2014).
22.   Berndt, A., Yizhar, O., Gunaydin, L. A., Hegemann, P. & Deisseroth, K. Bi-stable neural
state switches. *Nat. Neurosci.* **12**, 229–34 (2009).
23.   Berglund, K., Fernandez, A. M., Gutekunst, C. A. N., Hochgeschwender, U. & Gross, R.
E. Step-function luminopsins for bimodal prolonged neuromodulation. *J. Neurosci. Res.*
98, 422–436 (2020).
24.   Prakash, M. *et al.* Selective control of synaptically-connected circuit elements by all-
optical synapses. *Commun. Biol.* **5**, 33; 10.1038/s42003-021-02981-7 (2022).
25.   Govorunova, E. G., Sineshchekov, O. A., Janz, R., Liu, X. & Spudich, J. L. Natural light-
gated anion channels: A family of microbial rhodopsins for advanced optogenetics.
Science **349**, 647–50 (2015).
26.   Prakash, M., Medendorp, W. E. & Hochgeschwender, U. Defining parameters of
specificity for bioluminescent optogenetic activation of neurons using in vitro multi
electrode arrays (MEA). *J. Neurosci. Res.* **98**, 437–447 (2020).
27.   Hochbaum, D. R. *et al.* All-optical electrophysiology in mammalian neurons using
engineered microbial rhodopsins. *Nat. Methods* **11**, 825–833 (2014).
28.   Yang, J. *et al.* Coupling optogenetic stimulation with NanoLuc-based luminescence
(BRET) Ca(++) sensing. *Nat. Commun.* **7**, 13268; 10.1038/ncomms13268 (2016).
29.   Qian, Y., Rancic, V., Wu, J., Ballanyi, K. & Campbell, R. E. A Bioluminescent Ca(2+)
Indicator Based on a Topological Variant of GCaMP6s. *Chembiochem* **20**, 516–520
(2019).
30.   Tian, X. *et al.* An Enhanced Red Bioluminescent Indicator for Responsive Detection of
Physiological Calcium Dynamics in Cells and Mice. *ACS sensors* **10**, 5826–5833 (2025).
31.   Lambert, G. G. *et al.* CaBLAM: a high-contrast bioluminescent Ca2+ indicator derived
from an engineered Oplophorus gracilirostris luciferase. *Nat. Methods* **23**, 205-215
(2026).
32.   Shimomura, O., Masugi, T., Johnson, F. H. & Haneda, Y. Properties and reaction
mechanism of the bioluminescence system of the deep-sea shrimp Oplophorus
gracilorostris. *Biochemistry* **17**, 994–8 (1978).
33.   Inouye, S., Watanabe, K., Nakamura, H. & Shimomura, O. Secretional luciferase of the
luminous shrimp Oplophorus gracilirostris: cDNA cloning of a novel imidazopyrazinone
luciferase(1). *FEBS Lett.* **481**, 19–25 (2000).
34.   Inouye, S. & Sasaki, S. Overexpression, purification and characterization of the catalytic
component of Oplophorus luciferase in the deep-sea shrimp, Oplophorus gracilirostris.
Protein Expr. Purif. **56**, 261–268 (2007).
35.   Inouye, S., Sato, J.-I., Sahara-Miura, Y., Yoshida, S. & Hosoya, T. Luminescence
enhancement of the catalytic 19 kDa protein (KAZ) of Oplophorus luciferase by three
amino acid substitutions. *Biochem. Biophys. Res. Commun.* **445**, 157–162 (2014).
36.   Kim, C. K., Cho, K. F., Kim, M. W. & Ting, A. Y. Luciferase-LOV BRET enables versatile
and specific transcriptional readout of cellular protein-protein interactions. *Elife* **8**, e43826;
10.7554/eLife.43826 (2019).

37. Kim, C. K. *et al.* A Molecular Calcium Integrator Reveals a Striatal Cell Type Driving
Aversion. *Cell* **183**, 2003-2019 (2020).
38. Lee, S.-Y. *et al.* Engineered allostery in light-regulated LOV-Turbo enables precise
spatiotemporal control of proximity labeling in living cells. *Nat. Methods* **20**, 908–917
(2023).
39. Erdenee, E. & Ting, A. Y. A Dual-Purpose Real-Time Indicator and Transcriptional
Integrator for Calcium Detection in Living Cells. *ACS Synth. Biol.* **11**, 1086–1095 (2022).
40. Porta-de-la-Riva, M. *et al.* Neural engineering with photons as synaptic transmitters. *Nat.*
Methods **20**, 761–769 (2023).
41. Merolla, A. *et al.* A pH-sensitive closed-loop nanomachine to control hyperexcitability at
the single neuron level. *Nat. Commun.* **15**, 5609 (2024).
42. Novitskaya, Y., Sara, S. J., Logothetis, N. K. & Eschenko, O. Ripple-triggered stimulation
of the locus coeruleus during post-learning sleep disrupts ripple/spindle coupling and
impairs memory consolidation. *Learn. Mem.* **23**, 238–248 (2016).
43. Lustenberger, C. *et al.* Feedback-Controlled Transcranial Alternating Current Stimulation
Reveals a Functional Role of Sleep Spindles in Motor Memory Consolidation. *Curr. Biol.*
26, 2127–2136 (2016).
44. Paz, J. T. *et al.* Closed-loop optogenetic control of thalamus as a tool for interrupting
seizures after cortical injury. *Nat. Neurosci.* **16**, 64–70 (2013).
45. Lee, D., Hyun, J. H., Jung, K., Hannan, P. & Kwon, H.-B. A calcium- and light-gated
switch to induce gene expression in activated neurons. *Nat. Biotechnol.* **35**, 858–863
(2017).
46. Wang, W. *et al.* A light- A nd calcium-gated transcription factor for imaging and
manipulating activated neurons. *Nat. Biotechnol.* **35**, 864–871 (2017).
47. O'Neill, B. K. & Laughlin, S. T. Neuronal Calcium Recording with an Engineered TEV
Protease. *ACS chemical biology* **13**, 1159–1164 (2018).
48. DeNardo, L. A. *et al.* Temporal evolution of cortical ensembles promoting remote memory
retrieval. *Nat. Neurosci.* **22**, 460–469 (2019).
49. Sakurai, K. *et al.* Capturing and Manipulating Activated Neuronal Ensembles with CANE
Delineates a Hypothalamic Social-Fear Circuit. *Neuron* **92**, 739–753 (2016).
50. Qiu, Y. *et al.* On-demand cell-autonomous gene therapy for brain circuit disorders.
Science **378**, 523–532 (2022).
51. Lieb, A. *et al.* Biochemical autoregulatory gene therapy for focal epilepsy. *Nat. Med.* **24**,
1324–1329 (2018).
52. Nguyen, Q.-A. *et al.* Acetylcholine receptor based chemogenetics engineered for
neuronal inhibition and seizure control assessed in mice. *Nat. Commun.* **15**, 601;
10.1038/s41467-024-44853-8 (2024).
53. Hodneland, E., Kögel, T., Frei, D. M., Gerdes, H.-H. & Lundervold, A. CellSegm - a
MATLAB toolbox for high-throughput 3D cell segmentation. *Source Code Biol. Med.* **8**,
16; 10.1186/1751-0473-8-16 (2013).
54. Ludwig, K. A. *et al.* Using a common average reference to improve cortical neuron
recordings from microelectrode arrays. *J. Neurophysiol.* **101**, 1679–1689 (2009).

1179

1180 **Acknowledgments**
1181 We would like to thank all members of Bioluminescence Hub laboratories for their feedback,
1182 discussions, and thoughtful comments throughout the progression of this work.
1183 (http://www.bioluminescencehub.org/)
1184
1185 **Funding**

1186   This work was supported by National Institutes of Health grants R21MH101525, R21EY026427,
1187   U01NS099709, F99NS129170, National Science Foundation grants CBET-1464686, DBI-
1188   1707352, and the W.M. Keck Foundation.
1189
Author contributions
Conceptualization, AP, ELC, NCS, CIM, UH; Validation, AP, ELC, CIM, DL, NCS, UH; Formal
Analysis, AP, ELC; Investigation, AP, ELC, MP, ADS, ZZ, MG-R, MOT; Data Curation, AP, ELC;
Writing—Original Draft, AP, ELC; Writing—Review & Editing, ELC, NCS, CIM, UH;
Visualization, AP, ELC; Supervision, CIM, NCS, UH; Project Administration, UH; Funding
Acquisition, CIM, DL, NCS, UH.
1196
1197   **Data availability**
1198   The source data supporting the conclusions of this research is provided in this article and in the
1199   supplement. All datasets and custom scripts generated during these studies are freely available
1200   via the Brown Digital Repository (DOI pending).
1201
1202   **Competing interests**
1203   The authors have no conflicts of interest to declare.
1204
FIGURE LEGENDS
Figure 1. **Design of a *Gaussia* luciferase based Ca^{2+} sensor. a**. sbGLuc (light blue) is split by
a CaM–M13 module (purple), producing a dim basal signal in the presence of its luciferin (CTZ).
Binding of Ca^{2+} (yellow) reconstitutes an active luciferase and increases light emission in the
presence of luciferin. **b**. Elements of the coding sequence for LMC: *Gaussia* luciferase signal
peptide (red), N- and C-terminal fragments of sbGLuc (light blue), calmodulin sequence from
GCaMP6f (purple), M13 sequence (purple), and various linker regions (gray). **c.** Structure model
of LMC based on Alphafold2. **d-n.** LMC Ca^{2+} sensing in various contexts. **d, e.** HeLa cells. **d.**
Emission wavelength for LMC. LMC construct was transiently expressed in HeLa cells and light
emission after luciferin and histamine addition was measured across the wavelength spectra
(data presented as mean ± SEM). **e.** Change in LMC luminescence in the presence of luciferin
in response to 10 µM histamine in HeLa cells (N=12 wells). Boxplots show median, 25th and
75th percentiles (box edges), whiskers to most extreme data points, and individual outliers. **f.**
HEK293 cells. Percent change in bioluminescence in HEK293 cells expressing LMC in
response to 2 µM ionomycin under increasing extracellular Ca^{2+} concentrations. Cells were pre-
incubated with 5 µM luciferin and varying concentrations of Ca^{2+}; acute Ca^{2+} flux was initiated by
injection of ionomycin before photon counting in a luminometer (mean ± s.e.m). **g - j.**
Cardiomyocytes. **g.** Representative bioluminescence trace from an individual cardiomyocyte
over time. Labels t1-t3 indicate troughs and peaks at two different time points (orange font, gray
font). Pseudocolor images below show the same cell at the selected time points with ROI
indicated by red box. Scale bar = 20 µm. **h.** Bioluminescence traces from regions of interest
(ROIs) from three independent cardiomyocytes in culture (ROI3 represented in images in (g)),
highlighting spontaneous and ATP-evoked activity. **i.** Individual bioluminescence transients from
each ROI aligned to their peak and z-scored, with mean responses shown in bold. **j.**
Quantification of peak z-scored bioluminescence responses across ROIs. **k - n.** Neurons. **k.**
Representative hippocampal neurons expressing LMC imaged with luciferin alone (left) or with
luciferin and glutamate (right). Scale bar = 100 µm. **l.** Mean (blue) overlaid with individual
bioluminescence traces (grey) of LMC responses to bath-applied luciferin and glutamate. **m.**
Box plot showing maximal LMC luminescence responses (Max ΔRLU/RLU$_0$) with luciferin alone
vs glutamate plus luciferin (N= 29 neurons from two independent cultures, Wilcoxon signed rank
test: p= 2.7023 x10^{-6}). **n.** Summarizes the maximal responses shown in (m), presented as the

max fold change which was calculated per neuron as the ratio of peak responses in the
CTZ+glutamate condition relative to the CTZ alone.
1238
Figure 2. Integrating Ca^{2+} flux with activation of photoreceptors. a. Luminometer time
series from HeLa cells transiently expressing BlueCaMBI, GLICO, GeNL, or LMC following
addition of luciferin and histamine. Inset shows LMC traces on an extended scale. **b.** Peak
bioluminescence measured after luciferin and histamine stimulation for each sensor from (a). **c.**
Histamine-specific percent change in bioluminescence normalized to luciferin baseline,
expressed as percent change for each calcium sensor. (N=4 wells/sensor, LMC vs GLICO: *P*=
4.65x10^{-6}, GeNL: *P*= 5.51x10^{-7}, BlueCAMBI: *P*= 1.46x10^{-6}; GeNL vs GLICO *P*= 0.38,
BlueCAMBI P= 0.996; BlueCAMBI vs GLICO: *P*=0.723, Tukey's post hoc multiple-comparison's
test following a one-way analysis of variance (ANOVA), F($_{3, 12}$) **=** 48.56, *P=* 5.51x10^{-7}). Boxplots
show median, 25th and 75th percentiles (box edges), whiskers to most extreme data points, and
individual outliers.
1250
Figure 3. Converting intracellular activity to transcription. a. Schematic of LMC leading to
bioluminescence mediated dimerization of the light sensing transcription factor EL222 in the
presence of Ca^{2+} and luciferin (CTZ). The photoactivated EL222 homodimer then binds to the
promoter 5xC120 and leads to transcription of the reporter gene. **b.** Firefly luciferase (FLuc)
transcriptional reporter measured following no treatment (no luciferin, no histamine), luciferin-
only, or luciferin and histamine treatments in HeLa cells transiently expressing BlueCaMBI,
GLICO, GeNL, or LMC. **c.** FLuc transcriptional reporter normalized to luciferin baseline
expressed as percent change for each sensor/condition. N = 4-5 wells per sensor; LMC vs
BlueCaMBI: P = 0.00998; vs GLICO: P = 0.00998; vs GeNL 520: P = 0.00609; vs no calcium
sensor: P = 0.00609; one-sided Wilcoxon rank-sum tests following a Kruskal–Wallis (non-
parametric ANOVA) test across sensors: H = 16.1, df = 4, P = 0.00287. **d.** Representative
confocal images showing EYFP transcriptional reporter expression (green) in mouse prefrontal
cortex following NMDA alone, luciferin (CTZ) alone, or combined CTZ and NMDA
administration. Red fluorescence marks neurons expressing the LMC-P2A-dTomato construct.
Scale bar = 100 μm. **e.** Individual channels are shown separately with a merged image below.
Insets highlight representative neuron demonstrating co-localization of EYFP (green), LMC
(red), and EL222 (magenta) expression. Scale bar = 50 μm. Merged image includes DAPI and
scale bar = 20 μm. **f.** EYFP reporter expression quantified as per-cell EYFP/dTomato ratio
across treatment groups. Each point represents a single cell derived from three treatment
groups: Luciferin-NMDA (N= 350 cells/2 mice), Luciferin-ONLY N= 197 cells/2 mice), NMDA-
ONLY (N= 147 cells/2 mice). Dotted line indicates the 90th percentile EYFP/dTomato ratio
observed in the NMDA-ONLY group, used as a threshold to define transcriptional activation
above baseline. **g.** EYFP reporter intensity plotted against dTomato fluorescence for each cell
and condition. Boxplots show median, 25th and 75th percentiles (box edges), whiskers to most
extreme data points, and individual outliers.
1276
Figure 4. Integrating Ca^{2+} flux with change in membrane potential in single neurons *in*
***vitro*. a.** Brightfield and fluorescence images of cultured cortical neurons nucleofected with
opsin (ChR2(C128S))-EYFP and LMC-P2A-dTomato and used for whole-cell recordings. The
patch electrode is visible in the brightfield image. Scale bar = 50 μm. **b.** (i) Schematic of
farnesylated LMC anchored to the inner cell membrane near the excitatory opsin ChR2(C128S)
enabling light-mediated activation in the presence of depolarization-induced Ca^{2+} flux and the
luciferin (CTZ). (ii) Representative membrane voltage responses to step current injections
before (gray) and after (blue) bath perfusion of luciferin in a neuron co-expressing LMC and
ChR2(C128S). (iii) Action potential counts under vehicle (gray) and luciferin (blue) conditions.
Data expressed as mean ± SEM. (iv) Trial-level regression relating late firing to early firing

across sweeps. (v) Normalized across-neuron early–late lines showing per-cell transformations before and during luciferin. **c.** (i) Schematic of a control condition, where LMC is co-expressed with ChR2(C128S) in the presence of Ca^{2+} but without luciferin, preventing light-mediated activation. (ii–v) Same analyses as in b. **d.** (i) Schematic of LMC co-expressed with the inhibitory opsin hGtACR2, enabling light-mediated suppression of activity in the presence of Ca^{2+} and luciferin. (ii–v) Same analyses as in b, showing inhibition rather than activation. Data are presented as mean ± SEM in panels.

Figure 5. Integrating Ca^{2+}+ flux with change in membrane potential in neural populations *in vitro*. a. Schematic of farnesylated LMC anchored to the inner cell membrane near the excitatory opsin ChR2(C128S) enabling light-mediated activation in the presence of depolarization-induced Ca^{2+} flux and the luciferin (CTZ). **b.** Fluorescence images of cultured cortical neurons co-transduced with LMC-P2A-dTomato (red) and ChR2(C128S)-EYFP (green). **c.** Representative filtered voltage traces from one channel before and after vehicle (top) or luciferin (bottom) application. Red dashed line indicates time of application. **d.** MUA firing rate was increased in the luciferin condition (Wilcoxon signed-rank test, $P= 3.93 \times 10^{-28}$, Holm-corrected). **e.** Burst rate was not significantly different between luciferin and vehicle conditions (Wilcoxon signed-rank test, $P= 0.63$, Holm-corrected). **f.** Burst duration was significantly prolonged in the luciferin condition (Wilcoxon signed-rank test, $P= 3.18 \times 10^{-49}$, Holm-corrected). **g.** Time-aligned MUA traces across all electrodes showing normalized population responses following vehicle (top) or luciferin (bottom) application. Individual electrode traces are shown in grey (vehicle) and blue (luciferin); thick lines represent the mean response per condition. MUA was smoothed using a sliding 5-point boxcar window. **h.** Early-phase (0–50 ms) MUA firing rates following application was not different between luciferin and vehicle (Wilcoxon signed-rank test, $P= 0.13$, Holm-corrected). **i.** Late-phase (150–1000 ms) MUA firing rate was significantly higher in the luciferin condition (Wilcoxon signed-rank test, $P= 2.61 \times 10^{-8}$, Holm-corrected). **j.** Late phase normalized MUA was significantly increased under luciferin (Wilcoxon signed-rank test, $P= 1.02 \times 10^{-11}$, Holm-corrected). For **d-f** and **h-j**, N=57-59 pairs of electrodes per recording pooled across three independent MEA cultures. Boxplots show median, 25th and 75th percentiles (box edges), whiskers to most extreme data points, and individual outliers.

Figure 6. Integrating spontaneous and sensory inputs into changes in firing rates *in vivo*. a. Schematic of the experimental design. Extracellular electrophysiology recordings from SI barrel cortex during varied vibrissal deflections (zero, low, mid, max intensities) for 500ms at 20Hz. **b**. Schematic of the experimental timeline. Each recording consisted of 1000 trials (~2.5 hour sessions) where ACSF containing vehicle or CTZ was constantly perfused over the craniotomy. During each treatment epoch, vibrissal deflections were randomly presented. In the presence of the sensory stimulus and vehicle (grey box), there is no activity dependent light emission of LMC. In contrast, in the presence of CTZ and the sensory stimulus (blue box), sensory recruitment leads to activity dependent bioluminescence and subsequently photoactivation of opsins in neurons. **c**. Luciferin (CTZ) application increases spontaneous multiunit activity (MUA) in opsin-expressing mice relative to no-opsin controls. **d.** LED-evoked MUA responses in opsin-expressing mice exceed the 95th percentile of the no-opsin null distribution. *Inset:* MUA traces from a non-responsive channel (NR; orange) in a −opsin cohort and from responsive (R; teal) and non-responsive (NR; green) channels in a +opsin cohort. The red dashed line indicates LED onset. **e.** Sensory-evoked responses in vibrissae S1 from LMC-expressing mice (N = 2 mice per group). *Left*, mean ± SEM baseline-corrected multiunit activity (MUA) responses to a single whisker deflection during vehicle (grey) and after CTZ perfusion (blue) in opsin-expressing mice; corresponding responses from no-opsin control mice are shown in grey (vehicle) and black (CTZ). *Middle*, peristimulus time histogram (PSTH) of MUA during the full whisker stimulation epoch (1500 ms window); the red dotted line indicates

stimulus onset, and the whisker stimulus consisted of a 500 ms deflection train; *inset (green*
box) indicates the analysis window used to quantify evoked responses. **f.** Baseline-corrected
whisker-evoked multiunit activity (MUA) quantified from the analysis window indicated in **e** and
compared between vehicle and CTZ epochs across stimulus amplitudes (pooled across low,
medium, high trials). **g.** Trial-level regression relating late firing to early firing across recording
channels in LMC and opsin–expressing mice. **h.** Same analysis as in g, performed in LMC-
expressing mice lacking opsins.

a Lumicampsin LMC

+Luciferin (CTZ)

+Ca²⁺

genetically encoded, calcium dependent blue-light emitter

b
N-sbGLuc | Calmodulin from GCaMP6f | M13 | C-sbGLuc

GLuc Signal Sequence

Linker #1   Linker #2   Linker #3

c
C65 C77 disulfide bond

C-sbGLuc

M13

N-sbGLuc

CaM

Ca²⁺ ions

d
LMC

e
LMC

f
$R^2 = 0.8146$

g *in vitro* - cardiomyocytes single-cell bioluminescent dynamics

t1 t2 t3   t1 t2 t3

10 RLU (a.u.)   3 s

t1   t2   t3

t1   t2   t3

RLU (a.u.)   1.2   4.7

h *in vitro* - cardiomyocytes bioluminescent population dynamics

ROI1

ROI2

ROI3

ATP Onset

2 ΔRLU/RLU₀ (z-score)   10 s

i
ROI1
ROI2
ROI3

j
ROI1
ROI2
ROI3

k *in vitro* - hippocampal neurons bioluminescent population dynamics

+CTZ   +CTZ +Glutumate

RLU (a.u.)   0   1

l
CTZ Onset   Glutamate Onset

0.25 RLU (a.u)

250 s

m
***

+CTZ   +Glutamate +CTZ

n

a

b

c

a

| Brightfield | LMC P2A dTom | Opsin-EYFP | Merged |

b

(i) ChR2(C128S)

(ii) +VEH   +CTZ

(iii)

(iv)

(v)

400 ms

c

(i) ChR2(C128S)

(ii) +VEH   +VEH

(iii)

(iv)

(v)

400 ms

d

(i) hGtACR2

(ii) +VEH   +CTZ

(iii)

(iv)

(v)

400 ms