A Bioluminescent Activity Dependent (BLADe) Platform for Converting Intracellular Activity to Photoreceptor Activation

Emmanuel L. Crespo[1,2]†, Akash Pal[1,3]†, Mansi Prakash[1], Alexander D. Silvagnoli[1,3], Zohair Zaidi[4], Manuel Gomez-Ramirez[5], Maya O. Tree[1], Nathan C. Shaner[6], Diane Lipscombe[5,7], Christopher I. Moore[5,7]*, Ute Hochgeschwender[1-3]*

[1] College of Medicine, Central Michigan University, Mount Pleasant, MI 48859, USA
[2] Biochemistry, Cell and Molecular Biology Graduate Program, Central Michigan University, Mount Pleasant, MI 48859, USA
[3] Program in Neuroscience, Central Michigan University, Mount Pleasant, MI 48859, USA
[4] Duke University, Undergraduate Neuroscience Program, Durham, NC 27710
[5] Department of Neuroscience, Brown University, Providence, RI 02912, USA
[6] University of California, San Diego, School of Medicine, Department of Neuroscience, 9500 Gilman Drive La Jolla, CA 92093-0662, USA
[7] Carney Institute for Brain Science, Brown University, Providence, RI 02906, USA

†These authors contributed equally to this work: Emmanuel L. Crespo and Akash Pal
*Corresponding authors: Ute Hochgeschwender (hochg1u@cmich.edu) and Christopher I. Moore (christopher_moore@brown.edu)

Abstract

Genetically encoded sensors and actuators have advanced the ability to observe and manipulate cellular activity, yet few non-invasive strategies enable cells to directly couple their intracellular states to user-defined outputs. We developed a bioluminescent activity-dependent (BLADe) platform that facilitates programmable feedback through genetically encoded light generation. Using calcium (Ca^{2+}) flux as a model, we engineered a Ca^{2+}-dependent luciferase that functions as both a reporter and an activity-gated light source capable of photoactivating light-sensing actuators. In neurons, the presence of luciferin triggers Ca^{2+} dependent local illumination that provides activity dependent gene expression by activating a light-sensitive transcription factor and control of neural dynamics through opsin activation in single cells, populations and intact tissue. BLADe can be expanded to couple any signal that bioluminescent enzymes can be engineered to detect with the wide variety of photosensing actuators. This modular strategy of coupling an activity dependent light emitter to a light sensing actuator offers a generalizable framework for state dependent cell-autonomous control across biological systems.

Introduction

Bioluminescence, the emission of light generated when a luciferase enzyme oxidizes a small molecule luciferin, served as a non-invasive approach for *in vitro* and *in vivo* imaging for decades[1]. More recently, with the development of brighter, blue light emitting luciferases in combination with light sensing elements, including opsins and non-opsin photoreceptors[2,3], transformed bioluminescence from a mere imaging reagent to a versatile tool for controlling molecular processes. We and others have combined luciferases with channelrhodopsins and pumps for bioluminescence-driven changes in membrane potential[4–6], light-sensing transcription factors for gene expression[7–9], light-sensing enzymes for cAMP production[10], as well as with genetically encoded photosensitizers for reactive oxygen species production[11,12]. The main advantages of using locally produced bioluminescence rather than external light sources for activating photoreceptors are the independence from optical fiber implants, thereby avoiding the physical

52 damage to cells as well as the limitation in the number of cells that can be simultaneously
53 controlled.
54
We now are exploring a unique feature of employing genetically encoded light emitters. By
engineering the luciferase protein to accommodate sensor moieties, light emission can be made
dependent on the presence of various intracellular agents that report a cell's state. Several such
bioluminescent sensors have been developed for reporting intracellular biochemical signals, such
as intracellular Ca^{2+}, ATP, or cAMP, including for imaging studies *in vivo*[13]. Here, we are
combining a bioluminescent sensor with light-dependent actuators to create an integrator,
opening a general platform for converting biochemical signals within cells into photoreceptor
activation and resulting downstream changes in cellular function. This Bioluminescence Activity
Dependent (BLADe) paradigm serves to integrate cellular activity states with downstream
consequences from transcription to membrane potential changes, depending on the optogenetic
actuator coupled to the conditional luciferase. By engineering luciferases to respond to specific
biochemical cues, this framework can, in principle, be extended to a broad range of intracellular
states depending on the sensor domain used. Any bioluminescence indicator can be transformed
from a sensor into an activity integrator by using its light emission to activate photoreceptors.
69
To validate this experimental framework, we developed a BLADe platform based on intracellular
calcium level fluctuations. Changes in Ca^{2+} levels are directly translated into bioluminescence
light emission by making light production dependent on Ca^{2+} binding in a split luciferase construct.
The Ca^{2+} dependent light emitter was paired with a light-dependent transcription factor that
detected this light and affected transcription *in vitro* and *in vivo*. Moreover, when paired with
channelrhodopsins Ca^{2+} dependent light emission enabled neural activity-driven change in firing
activity *in vitro* in single neurons and neuronal populations; in mice, BLADe converted
spontaneous activity and sensory evoked changes into activity dependent control of neocortical
network dynamics.
79
80
81 **Results**
82
Engineering of a luciferase-based Ca^{2+} sensor
The *Gaussia princeps* luciferase variant sbGLuc is an efficient transmitter of photons/energy to
opsin chromophores when tethered to opsins in luminopsins[5,14]. To make it Ca^{2+} dependent we
split sbGLuc as was done previously for wildtype GLuc between Q88 and G89[15] and introduced
the Ca^{2+} sensing moiety CaM-M13 employing the M13 sequence from the Ca^{2+} sensor YC3.6[16]
and the calmodulin sequence from GCaMP6f[17] (**Fig. 1a-c**). We reasoned that inserting the CaM-
M13 domain between Q88 and G89 would allosterically control enzymatic activity by modulating
the structural integrity of the native disulfide bond (C65–C77) which stabilizes the rigid α4–α7
helical core of sbGLuc (**Fig. S1 and S2**). In the absence of Ca^{2+}, the inserted calmodulin domain
destabilizes this interaction, whereas Ca^{2+} binding promotes reassembly of the split halves and
restoration of luciferase activity. We called this split sbGLuc based Ca^{2+} indicator Lumicampsin
or LMC, and LMC4 as this was the fourth split sbGLuc construct we generated.
95
To expand on LMC4, we engineered four additional variants: LMC5, LMC6, LMC7, and LMC8.
LMC5 and LMC6 incorporate calmodulin sequences from the fluorescent calcium indicators
GCaMP6s and GCaMP6m, respectively[17]. LMC7 and LMC8 incorporate calmodulin domains
derived from the Ca^{2+}-dependent luciferases Ca^{2+}-NL and Ca^{2+}-eNL[16,18]. We selected the
calmodulin domains based on their performance in either fluorescent Ca^{2+} indicators or prior

luciferase-based calcium sensors, with the goal of improving the dynamic range of our LMC
variants (refer to **Fig. S3** for the amino acid differences across variants).

As the goal of LMC is to serve as a genetically encoded light source for activating blue light-
sensing photoreceptors, we first confirmed its light emission spectra and Ca^{2+} responsiveness in
cells across variants. HeLa cells were transfected with each LMC variant, incubated with 5 µM
luciferin (CTZ), and stimulated with histamine to induce a physiologically relevant increase in
intracellular Ca^{2+}. All variants retained a peak emission at roughly 490 nm (**Fig. 1d).** The percent
change in light emission before and after histamine addition ranged from 232% to 288% across
LMC variants. All LMC variants provide a blue-shifted optical report of intracellular calcium,
enabling potential coupling to blue light sensitive optogenetic proteins required for the BLADe
platform. Overall, there was no significant difference between the variants when compared to
LMC4 (**Fig. 1e**).

Experiments in HeLa cells confirmed that LMC4 responds to histamine-induced Ca^{2+} release at
physiologic levels, but this system does not allow experimental control of intracellular Ca^{2+} levels
in living cells. To address this, we transfected HEK293 cells with LMC4 and applied ionomycin
under defined extracellular Ca^{2+} concentrations ranging from 0.4875 µM to 39 µM (**Fig. 1f**). LMC4
reported ionomycin-induced Ca^{2+} flux at all tested concentrations. The decrease in
bioluminescent signal at lower extracellular Ca^{2+} conditions (0.4785µM -1µM) reflects the loss of
intracellular Ca^{2+} as ionomycin moves Ca^{2+} from high (intracellular) to low (extracellular)
concentrations. We compared the response to 39 µM extracellular Ca^{2+} between HEK293 cells
expressing LMC4 or GCaMP6f. Under the same conditions, LMC4 reported a 43% change in
signal from baseline whereas GCaMP6f reported an 8.2% change, indicating a high dynamic
range of calcium dependent light emission by LMC4 (**Fig. 1g**).

Having established that LMC4 responds to flux in intracellular Ca^{2+} levels in populations of cells,
we next tested how subcellular localization affects its performance. We generated four LMC4
variants targeted to distinct subcellular compartments by adding varying localization sequences
(**Fig. S4a**). As one of the intended uses of LMC4 is to activate light sensing proteins, we compared
two ways of localizing the Ca^{2+}dependent light emitter to the inner cell membrane. We did this by
adding either a C-terminal farnesylation (LMC4f) or an N-terminal myristoylation (myLMC4) signal.
We compared Ca^{2+} induced light emission of the two constructs in HeLa cells and found the
farnesylated version to perform significantly better (**Fig. S4b**). We compared LMC4f to LMC4er
that localizes to the ER via a C-terminal KDEL sequence. HEK293 cells expressing either
construct were tested in the absence (0 µM) or presence (39 µM) of extracellular Ca^{2+} and 10 µM
CTZ. Acute Ca^{2+} flux was elicited by addition of 2 µM ionomycin. In the absence of extracellular
Ca^{2+} the signal from LMC4er expressing cells was higher than that from LMC4f expressing cells
due to ER mobilization of internal Ca^{2+} stores. In contrast, in 39 µM extracellular Ca^{2+} a high
bioluminescence signal was observed in LMC4f expressing HEK cells with very low signal from
LMC4er expressing cells (**Fig. S4c**). Lastly, we generated miLMC4 that targets the Ca^{2+} sensor
to mitochondrial membranes. Mitochondria localization was confirmed by co-expressing miLMC4
with a mitochondrial tag BacMam2.0-Mito-GFP in HeLa cells. Confocal images show that the
Mito-GFP tag and miLMC4 co-localize (**Fig. S4d**). Thus, we proceeded with LMC4f for all
subsequent experiments, designated henceforth as LMC.

To test the ability of LMC to report Ca^{2+} dynamics in culture, we transfected primary
cardiomyocytes with LMC and imaged them on DIV4 using an EMCCD camera. Cells were pre-
incubated with luciferin (100 µM CTZ) for ~10 min before imaging. Addition of 10 µM ATP
increased contractility and induced rapid Ca^{2+} transients with robust peaks (**Fig. 2a and b**). To

assess Ca²⁺ reporting in deep tissue, we transduced the hearts of Swiss Webster mice with
AAV2/9-CAG-LMC4 and performed *in vivo* imaging three weeks later. Following intraperitoneal
injection of 100 µg water-soluble luciferin, bioluminescent signals from the heart became visible
through the chest within 10 minutes, peaked around 15 minutes post-injection, and gradually
declined to baseline over the next hour (**Fig S5**).

156
Next, to test whether LMC could report neurotransmitter-evoked Ca²⁺ influx in neurons, we
transfected primary hippocampal cultures and perfused luciferin (CTZ) followed by luciferin with
glutamate. Bioluminescent signals were low with luciferin alone but increased after glutamate
application **(Fig. 2c)**. Time-locked traces showed a clear rise in bioluminescence following
glutamate onset (**Fig. 2d**). Across individual neurons, we observed an increase in normalized
peak responses with glutamate compared to luciferin alone (**Fig. 2e**). In addition, responses also
exhibited high contrast relative to baseline, with a median fold-change of ~2–4 across neurons
(**Fig. 2f**).

165
Having confirmed that LMC reports Ca²⁺ fluctuations in cardiomyocytes (*in vitro* and *in vivo*) and
glutamate-evoked responses in hippocampal neurons, we next tested its performance in the brain
in vivo. Mouse barrel cortex was injected in two distinct regions with AAV2/9-hSyn-LMC or
calcium-insensitive GLuc and imaged through a cranial window in head-fixed mice. Luciferin
(CTZ, 50 µL) was delivered into the lateral ventricle during continuous imaging, followed 5 min
later by direct cortical NMDA injection (1 µL). Bioluminescence increased selectively in the LMC-
expressing region after NMDA, whereas the calcium insensitive variant GLuc control region
remained at baseline (**Fig. 2 g and h**). Together, these results validate LMC as a bioluminescent
reporter of Ca²⁺ dynamics *in vitro* and *in vivo*. Further engineering of LMC is necessary for
acquiring higher spatial and temporal resolution, as the image acquisition rate was restricted to
~1-4 Hz in our imaging conditions.

177
Bioluminescence as an activity-gated light source
We next compared LMC with other published bioluminescent Ca²⁺ indicators to evaluate its
baseline emission and suitability as an activity-gated light source. HeLa cells were transiently
transfected with LMC, BlueCaMBI[19], GLICO[20], or GeNL[18] and light emission was assayed in a
luminometer following luciferin addition and histamine stimulation. Time-series traces show that
BlueCaMBI, GLICO, and GeNL, produced higher overall photon output than LMC (**Fig. 3a**).
Although these sensors were brighter, they also exhibited substantial baseline photon emission
in the presence of luciferin under resting Ca²⁺ conditions, relative to their response after histamine
stimulation. (**Fig. 3b**). All three indicators—BlueCaMBI, GLICO, and GeNL—have been reported
to capture Ca²⁺ fluctuations, but in those studies, bioluminescence signal traces were typically
normalized following luciferin pre-incubation[18–20]. While this normalization approach is appropriate
for imaging relative Ca²⁺ dynamics, the high absolute photon emission at resting levels of Ca²⁺
limits their use as activity-gated light sources required for BLADe activation of photoreceptors.

191
High absolute brightness benefits imaging, but BLADe optogenetic applications require low basal
emission and a large absolute increase in light output during Ca²⁺ influx. Consistent with this,
when normalized to the luciferin baseline, only LMC exhibited a large relative change in photon
emission following histamine-induced Ca²⁺ release (**Fig. 3c**). Accordingly, despite lower absolute
brightness, the unique low background and high dynamic range of LMC positions it as an activity-
gated light source ideally suited for converting intracellular signals into optical activation of
downstream photoreceptors. At resting calcium levels, we hypothesize LMC light emission
remains dim and is below the threshold needed to activate photoreceptor chromophores,
minimizing spurious activation of light sensing proteins. We next asked whether this activity-
dependent light source could drive optogenetic tools as the basis of the BLADe platform.

Integrating Ca²⁺ flux with change in transcription

203
LMC expressing cells showed low baseline bioluminescence in the presence of luciferin and a
204
strong increase in light output following Ca²⁺ influx. We reasoned that LMC's activity-gated light
205
emission could be used to photoactivate light-sensitive transcriptional effectors. To test this
206
prediction, we co-expressed LMC with a light sensing transcription factor EL222[21]. In the presence
207
of blue light, EL222 dimerizes and then binds to its promoter 5xC120, leading to transcription of
208
the reporter gene (**Fig. 4a**). Next, we treated HeLa cells expressing each calcium sensor with
209
luciferin for 15 minutes and measured the FLuc reporter 18 hours later in the following conditions:
210
dark, luciferin-only, and luciferin + histamine (**Fig. 4b**).
211

212
All calcium dependent indicators produced FLuc transcription under luciferin treatment alone,
213
indicating that transcription can be photoactivated even in the absence of calcium. In the presence
214
of luciferin, baseline transcription was strongest for BlueCaMBI, GLICO, and GeNL, which were
215
markedly brighter than LMC — a trend consistent with their higher baseline bioluminescence in
216
luminometer traces (Fig. 3b). In contrast, only LMC showed a calcium-dependent increase in
217
transcription, with a significant increase in FLuc signal in the luciferin + histamine condition
218
compared to luciferin alone (**Fig. 4c**, LMC vs BlueCaMBI: P= 0.00998, GeNL: P=0.00609, GLICO:
219
P=0. 00998, one sided Mann-Whitney U test). The overall transcriptional output from LMC
220
remained modest compared to the other sensors, which is consistent with its lower total light
221
output in luminometer assays (Fig. 3b). Having established that LMC produces calcium-
222
dependent light sufficient to activate EL222 *in vitro*, we next tested whether the BLADe
223
transcriptional system could function *in vivo*.
224

225
As a pilot study, we tested whether LMC could drive transcription in neurons *in vivo* when network
226
activity was chemically elevated with NMDA. We performed stereotactic injections of AAV9-hSyn-
227
LMC-P2A-dTomato, AAV9-CAG-EL222, and AAV9-5×C120-EYFP into the prefrontal cortex of
228
anesthetized mice. After virus delivery, the craniotomy was immediately covered to further
229
minimize unintended light exposure during recovery. After 2–3 weeks, to allow for maximal
230
expression, mice received intraventricular injections of either NMDA + vehicle, luciferin (CTZ)
231
alone, or luciferin + NMDA. These conditions allowed us to test: (1) non-specific background
232
transcriptional activity in the absence of light (NMDA + vehicle), (2) light-dependent transcription
233
in the context of spontaneous activity (luciferin alone), and (3) BLADe-mediated transcription
234
(luciferin + NMDA).
235

236
Confocal imaging of tissue collected ~20 hours after treatment showed increased EYFP reporter
237
labeling in individual LMC-expressing neurons from mice treated with luciferin + NMDA, compared
238
to the other treatment groups (**Fig. 4d**). Single-cell analysis confirmed that EYFP reporter
239
increase occurred across a range of dTomato expression levels (**Fig. 4e**), indicating that the Ca²⁺-
240
dependent enhancement of transcription was not due to differences in viral expression of LMC.
241
Quantification of EYFP reporter/dTomato tool expression ratios revealed higher reporter
242
expression in the luciferin + NMDA group compared to both the luciferin-only group and the
243
NMDA-only group (**Fig. 4f**, Wilcoxon rank-sum test, Bonferroni-corrected luciferin+NMDA vs
244
luciferin: P= 1.62×10^{-7}; luciferin+NMDA vs NMDA: P= 0.0057; luciferin vs NMDA: P = 0.196;
245
pooled from FOVs of 197-350 cells/N=2 mice for each treatment group). The luciferin-only and
246
NMDA-only groups were not significantly different from each other suggesting that reporter
247
activation requires both intracellular calcium and bioluminescent light, consistent with LMC-
248
BLADe gating. Together, these findings show that BLADe can function *in vivo* in the mouse brain
249
to drive calcium-dependent transcription, but the overall magnitude of reporter labeling remains
250
modest. Increasing light output of the Ca²⁺ dependent luciferase or improving the contrast
251

252 between active and inactive cells may help optimize this system for robust transcriptional labeling
253 in future applications.
254
Integrating Ca²⁺ flux with change in membrane potential
To demonstrate the modularity of the BLADe platform, we tested whether LMC could drive light-
sensitive ion channels. Following an action potential, the rapid influx of intracellular Ca²⁺ provides
a convenient and high-resolution proxy for BLADe neural activation. We reasoned that LMC could
detect these calcium transients and, through co-expression with an excitatory opsin, rapidly
modulate membrane potential in an activity-dependent manner. To this end, we applied step
current injections to hippocampal neurons *in vitro* in the presence of synaptic blockers to isolate
cell-intrinsic responses. We then performed whole-cell patch-clamp recordings on neurons co-
transformed with LMC and an opsin **(Fig. 5a).** By visualizing co-localized expression of an opsin-
EYFP (green; here ChR2(C128S)) and the Ca²⁺-dependent luciferase LMC-P2A-dTomato (red),
we selected these double-positive neurons for testing the integration of cellular states via BLADe
opsin activation.
267
We predicted that LMC generates light directly in response to Ca²⁺ influx which occurs during
spiking induced by depolarizing current injections. We reasoned that increasing levels of
depolarizing current in the presence of the luciferin would produce enough bioluminescence to
further photoactivate ChR2(C128S)[22] via a cell-autonomous, positive feedback optical loop. In
line with our prediction, luciferin perfusion during the same depolarizing current elicited
significantly more detected action potentials than during vehicle perfusions **(Fig 5b, ii and iii,**
N=16 neurons, *P= 4.954x10⁻⁹,* paired sample Wilcoxon signed rank tests), indicating a light-
dependent, cell-autonomous control of membrane voltage potential. Our findings demonstrate
LMC bioluminescence is sufficient to photoactivate the opsin, they do not directly confirm BLADe
control of opsins.
278
To isolate the activity-dependent component—which is expected to change how spikes are
generated over the course of the current stimulation—we calculated a spike count fractional
difference by evaluating the temporal distribution of spikes within each 1-second depolarizing
current across trials. We subtracted the number of spikes in the second half of the current from
the number of spikes in the first half, then normalized by the total spike count of a given trial. A
value of +1 indicates that all spikes occurred in the first half, a value of 0 reflects an even
distribution across both halves, and a value of -1 indicates that all spikes occurred in the second
half. Therefore, we can determine how LMC-driven opsin activation changes action potential
generation over the course of depolarization.
288
For neurons co-expressing the excitatory opsin under luciferin perfusion, we observed a
significant shift toward fraction-difference values in the presence of the luciferin **(Fig 5b iv**, N=16
neurons, Kolmogorov-Smirnov test, KS= 0.25, *P*= 0.01319). Neurons co-expressing LMC and
ChR2(C128S) generated action potentials in the initial 500 ms but produced additional spikes in
the latter half of the 1s depolarizing step in the presence of the luciferin. The observed shift in
firing patterns supports the activity dependent nature of LMC light emission and BLADe control of
opsins. We hypothesize that the observed effect arises from an intensifying positive feedback
loop mechanism, whereby sustained calcium influx over the course of the depolarizing current
promotes ongoing opsin activation and alters spike timing.
298
In contrast, neurons co-expressing LMC and ChR2(C128S) that received two sequential vehicle
perfusions—rather than luciferin in the second perfusion—showed no changes in both the
frequency and temporal shift of the firing rate distributions **(Fig 5c ii-iv,** N=14 neurons, *P*=0.3117*,*
paired sample Wilcoxon signed rank test, Kolmogorov-Smirnov test, KS=0.0597, *P*= 0.999). This

result confirms that the observed bias in spike timing requires luciferin, and by extension, BLADe's
stimulus-dependent bioluminescence. To further ensure these changes depend on a functional
opsin rather than bioluminescence alone, we next tested a non-photoconductive ChR2 mutant,
ChR2(C128S)-E97R-D253A[23,24] ("DUD" opsin) alongside LMC (**Fig. S6**). There were no observed
luciferin-induced shifts in spike timing or firing rate, confirming that BLADe control does not alter
the excitability in the absence of an active light-sensitive channel.

Building on our findings with the excitatory opsin, we next tested whether BLADe could suppress
firing in neurons expressing an inhibitory opsin. The same cell-autonomous bioluminescent
positive feedback loop is expected to drive hyperpolarization rather than depolarization, leading
us to predict a reduction in firing rate and spike timing. In line with our prediction, neurons co-
expressing hGtACR2[25] and LMC exhibited fewer total spikes (**Fig. 5d, ii and iii,** N=16 neurons**,**
P=0.3117*,* paired sample Wilcoxon signed rank test) and had decreased spikes in the second
half of the depolarizing current (**Fig. 5d, iv,** N=16 neurons, Kolmogorov-Smirnov test, KS=
0.3888, *P*= 0.00809).

By coupling a depolarizing current and the resulting Ca^{2+} influx to opsin activation, BLADe
provides an all-molecular strategy to sense and modulate neuronal excitability. This can be further
customized based on the biophysical characteristics of the co-expressed opsin. Therefore, LMC
integrates intracellular states to bidirectionally alter the magnitude and spiking dynamics in a cell-
autonomous manner.

Moving beyond single-cell recordings, we next tested how LMC modulates firing rates at the
network level by performing extracellular recordings on cultured cortical neurons plated on multi-
electrode arrays (MEA)[26]. As in our patch experiment, we virally co-transduced an excitatory opsin
ChR2(C128S)-EYFP and the Ca^{2+}-dependent luciferase LMC-P2A-dTomato (**Fig. 6a,b**). Time-
locked multi-unit activity (MUA) traces revealed rapid population-level activity following luciferin
or vehicle application (**Fig. 6c**). While vehicle addition alone can transiently increase network
activity in MEA—likely due to mechanical or ionic effects of media perfusion—these responses
were brief and stereotyped across independent experiments. To enable comparison, we aligned
all trials to the application artifact, allowing time-locked analysis between vehicle and luciferin
treatments. In doing so, we could distinguish true activity-dependent changes from non-specific
network effects caused by vehicle addition. Vehicle and luciferin treatments were randomized on
the same MEA culture. All comparisons used the same electrode sites with recording ~4-5 hours
apart.

In the presence of luciferin, MUA was enhanced relative to vehicle one second after application
(**Fig. 6d**)**.** Burst firing rates were similar between conditions (**Fig. 6e**), yet burst duration was
significantly longer with luciferin (**Fig. 6f**). With burst firing rates unchanged but MUA elevated,
we next tested whether neural activity was prolonged over time due to co-expression of an
excitatory opsin with LMC. To this end, we analyzed the early phase as defined by the 0–50 ms
window beginning immediately after the application artifact, corresponding to the initial rise in
MUA relative to baseline (**Fig. 6g**). The time-locked window allowed for reliable comparisons of
immediate firing between treatment conditions. The late phase was defined as the remaining
portion of the 1-second window, from 100–1000 ms post-application, allowing us to test whether
LMC modulates both the onset and sustained components of network activity.

Early phase firing rates were similar between vehicle and luciferin conditions indicating that initial
responses to perfusion were comparable across treatments (**Fig. 6h**). Thus, the observed effects
are not due to media-driven excitability or non-specific stimulation. In contrast, the late-phase
response was significantly enhanced in the presence of luciferin (**Fig. 6i).** When normalizing the

late-phase MUA to the early-phase, we observed a clear enhancement in MUA consistent with LMC activation with luciferin (**Fig. 6j**). The observed sustained effect on network excitability in the presence of the luciferin suggests that LMC-driven light production prolongs neuronal activity via local activation of the co-expressed excitatory opsin.

Lastly, we tested whether LMC could bidirectionally modulate firing rates in populations of cultured neurons. In agreement with our whole-cell patch-clamp recordings, we observed a reduction in MUA activity when LMC was co-transduced with the inhibitory opsin hGtACR2 (**Fig. S7**). To test whether the observed inhibition requires LMC bioluminescence, we compared firing rate changes before and after luciferin application in neurons expressing hGtACR2 alone (**Fig. S7, S8**) versus those co-expressing LMC. Neurons lacking LMC showed a modest increase in activity (median ΔFR = +7 Hz, n = 19), while LMC4-expressing neurons exhibited a strong suppression (median ΔFR = −14 Hz, n = 51). The difference in MUA was significant (Wilcoxon rank-sum test, P = 1.576 × 10^{-10}), confirming that luciferin-induced inhibition is specific to neurons expressing LMC.

Building on our *in vitro* electrophysiology observations, we next wanted to determine if LMC could integrate physiological sensory inputs to alter neocortical networks. To move from cellular responses and into tissue-level dynamics, we leveraged the well-characterized vibrissae circuit to ask whether LMC can detect organism-scale sensory cues and reshape how they are encoded within the neocortex. To this end, we injected Cre-dependent versions of LMC and the excitatory opsin CheRiff[27] into the left cortex of Emx1-Cre mice to target both constructs to excitatory pyramidal neurons within the vibrissa region of primary somatosensory cortex (vSI). Three to four weeks later, we then performed extracellular recordings with laminar silicon probes in vSI. During recordings in head fixed anesthetized mice, we constantly perfused ACSF or ACSF containing luciferin (CTZ) over the cortical surface while delivering vibrissal deflections (**Fig. 7a, b**).

We first tested whether spontaneous neural activity driven bioluminescence could alone modulate cortical activity via BLADe opsin activation. By expressing CheRiff in vSI pyramidal neurons, we predicted increased MUA either through external LED stimulation or via activity-driven bioluminescence. Consistent with our prediction, we observed increased spontaneous MUA in the presence of luciferin relative to vehicle epochs (**Fig. 7c**, Wilcoxon signed rank test p=2.32 x 10^{-10}, p < 0.05 with post hoc Bonferroni correction for multiple comparison, vehicle epoch: M= 26.4, SD= 16.61, median= 22.1, range= 5.4-80.4, CTZ epoch: M= 52.7, SD= 26.0, median= 43.6, range= 17.5-126.6). These results suggest that LMC detects spontaneous intracellular calcium transients and converts them into bioluminescent output sufficient to activate opsins and enhance cortical spiking.

To confirm that the luciferin mediated enhancement was through opsin activation, we delivered a brief sinusoidal LED stimulation (8 Hz sinusoidal pattern, 500 ms) at the end of each session. Light responsive channels were observed relative to baseline spontaneous activity (**Fig. 7d**, Kruskal-Wallis test p= 3.23 x 10^{-08}; p < 0.05 with post hoc Bonferroni correction for multiple comparison, during baseline: M= 24.13 Hz, SD= 8.51, median= 22.7, range: 14.4-54.8, during stimulation: M= 83.1 Hz, SD= 96.63, median= 45.2, range=16.9-406.83). In contrast, non-responsive channels had no change during LED stimulation (**Fig. 7d,** Kruskal-Wallis test p > 0.05, during baseline: M=40.5, SD= 32.4, median= 22.7, range=0.1-173.26, during stimulation: 0.1-173.07). Together with our luciferin epoch data, these results support that LMC increases spontaneous cortical activity through bioluminescent optogenetic activation of CheRiff.

Following our observation that LMC enhances spontaneous cortical activity *in vivo*, we next tested whether LMC can also rapidly amplify neural responses to external sensory inputs. To this end, we tested for relationships between enhanced neural activity and vibrissae stimulus amplitude by

comparing MUA between vehicle and CTZ epochs. We delivered 10 vibrissae deflections at 20 Hz
over a 500 ms stimulus window, with deflections randomized in amplitude. We observed evoked
spikes within ~10–20 ms in vSI following low-amplitude vibrissae deflections, consistent with
known sensory latencies in vSI and validating our electrode placement (**Fig. 7e**). There is an
increase in MUA in sensory responsive channels in low stimulus trials in the presence of luciferin
compared to vehicle (blue vs grey trace, **Fig. 7e**).

To isolate BLADe recruitment of cortical activity, we quantified evoked MUA in two non-
overlapping post-stimulus windows: an early phase (0–25 ms), corresponding to the immediate
sensory response, and a late phase (26–50 ms), where delayed or prolonged firing may reflect
enhanced excitability. During the early response window (0–25 ms), which corresponds to direct
thalamocortical input, firing rates were increased during the luciferin trials compared to vehicle
(**Fig. 7f**, Kruskal-Wallis test, low: p= 7.62 x 10^{-06}, mid: p= 1.9 x 10^{-06}, max: p= 3.72 x 10^{-09}, p <
0.05 with post hoc Bonferroni correction for multiple comparison). In the late window (26–50 ms),
likely reflecting recurrent or sustained neocortical activity, we also observed enhancement in the
presence of luciferin (**Fig. 7g**, Kruskal-Wallis test, low: p= 7.63 x 10^{-06}, mid: p= 3.81 x 10^{-06}, max:
p= 2.61 x 10^{-08}, p < 0.05 with post hoc Bonferroni correction for multiple comparison). We interpret
this as evidence that LMC contributes to feedforward amplification of thalamocortical sensory
inputs and delayed phases of cortical activity, likely driven by recurrent excitatory input through
BLADe opsin activation. Collectively, these findings demonstrate that LMC enables cell-
autonomous, activity-gated control of spontaneous and sensory evoked neocortical network
dynamics *in vivo*. Through co-expression of LMC with light sensitive ion channels, BLADe
photoactivation offers a genetically encoded strategy for sensing and manipulating neural
dynamics.

Discussion

By using the concept of a luciferase split by a sensing moiety, any bioluminescent sensor can be
made into an integrator by combining it with a photoreceptor. As a proof-of-concept and to
demonstrate the versatility of the approach, we engineered a Ca^{2+} dependent luciferase and
tested its ability to activate either a transcription factor or channelrhodopsins. We split the *Gaussia*
luciferase variant sbGLuc and inserted a Ca^{2+} sensing moiety between the luciferase fragments.
The engineered luciferase, called lumicampsin (LMC), contains a calcium sensing domain
composed of the M13 peptide from YC3.6 and calmodulin from GCaMP6f. We identified critical
design constraints for enabling activity dependent control of photosensing molecules: at resting
intracellular Ca^{2+} levels, the presence of luciferin should generate minimal light output – ideally
below the activation threshold of photoreceptor chromophores. Furthermore, elevation of
intracellular Ca^{2+} must produce a sufficient increase in bioluminescence to cross this threshold
and initiate photoreceptor activation. We successfully applied LMC to convert neural activity into
transcription of a reporter gene and into changes in membrane potential *in vitro* and *in vivo*.

Several bioluminescent Ca^{2+} sensors have been developed by splitting luciferase enzymes and
inserting Ca^{2+} sensing moieties[16,18–20,28,29]. Because bioluminescent probes have historically been
far dimmer than fluorescent probes, most efforts have focused on maximizing light output. As a
result, existing bioluminescent Ca^{2+} sensors successfully report Ca^{2+} fluctuations, but with Ca^{2+}
dependent increases in signal riding on top of high baseline light emission. For many imaging
applications, baseline bioluminescent signal in the absence of high Ca^{2+} is acceptable. However,
this is incompatible with the BLADe approach, as the background light emission by itself is already
high enough to activate a photoreceptor. To avoid unintended photoactivation, the split luciferase
must produce very low levels of light emission at rest –when luciferin is present but intracellular
calcium is low. Under these conditions, the luciferase enzyme should remain incompletely

reconstituted and emit minimal light. Only upon calcium binding—or the presence of a specific
co-factor needed for enzyme reconstitution—should bioluminescence steeply increase to a level
sufficient for activating nearby photoreceptors. Comparison to other bioluminescent Ca^{2+} sensors
revealed this critical difference in SNR for the LMC construct used here to establish the BLADe
paradigm. This also motivated the development of a much brighter Ca^{2+} dependent luciferase,
CaBLAM, that fulfills these criteria and will enable improved performance of BLADe-based
systems[30].

There is an increasing need to not only monitor biochemical fluctuations in real time, but to
permanently mark and modify cells based on intracellular activity, enabling their subsequent
identification and targeted manipulation. The drive for developing such integrators has been most
prolific in neuroscience, challenged by the question of how to identify and study the ensemble of
active neurons underlying a specific behavior: Which cells in which contexts form the necessary
chain of actors that selectively lead to perceptions, actions and cognition. A recent generation of
approaches was created that attempts to advance this area of study by using calcium sensitivity
and engineered TEV protease as the mechanism that creates activation-dependence[31–33]. The
leading examples of this approach, Cal-Light[31] and FLARE[32], employ timed light to target specific
behavioral epochs. In essence, the coincidence of light and neural activity results in release of a
transcription factor: light application leads a LOV domain to expose a protease cleavage site; the
protease will only be in proximity to the cleavage site in the presence of high Ca^{2+}, with
photostimulation providing a high degree of temporal and spatial control. This Ca^{2+} dependent
protease strategy has significant limitations in that it requires a fairly complicated molecular
design, increasing the potential for failure points in distinct cell types. More importantly, light
pulses from a fiber optic are required to drive expression, which does not allow whole ensemble
tagging across the brain. In contrast, BLADe relies on the straightforward design of a split
luciferase that reconstitutes with increasing levels of Ca^{2+} and emits light sufficient for activating
light-sensing transcription factors in the presence of luciferin, as demonstrated *in vivo* by
expression of fluorescent reporters in mouse neocortex. As the system is chemically gated,
systemic application of the luciferin will reach each cell expressing BLADe integrators based on
increasingly improved bioavailability of luciferins.

One of the most powerful applications of BLADe for molecular integration is as a tool for all-
molecular, real-time feedback regulation. The lives of cells, organs and organisms are dependent
on chemical fluctuations on the time scale of milliseconds to seconds. Failure to regulate these
ongoing dynamics is a common motif in disease. When the proper feedback control of a local
augmenting process fails, runaway activity can result—as observed in activity-dependent cell
death and spreading neural activity during a seizure. Conversely, failure to amplify local
fluctuations can devastate biological processes such as glucose-driven insulin production or
information processing in the brain. When paired with ion-moving optogenetic actuators, BLADe
enables the conversion of neural activity into real-time changes in membrane potential, supporting
closed-loop feedback control of neuronal activity. The efficacy of this self-regulation was
demonstrated *in vivo* where LMC converted spontaneous activity and sensory evoked responses
into activity dependent control of neocortical network dynamics.

Recently, bioluminescence emitted from an intact luciferase has been applied to couple a sensor
for neuronal hyperactivity with a molecular actuator capable of switching off neuronal activity in
an all-molecular negative feedback loop at the single neuron level[34]. When excited at ~400–410
nm, the ratiometric pH sensor E2GFP increases light emission at 510 nm in proportion to
acidification—a hallmark of neuronal hyperactivity in seizures. To generate a pH sensitive
inhibitory luminopsin (pHIL), three proteins were fused together: a hyperpolarizing opsin
(eNpHR3,0), E2GFP, and a *Renilla* luciferase. The N-terminus of the construct contained the

eNpH3.0, a chloride pump activated by 510 nm light. E2GFP was placed in the center, serving
as the pH-responsive optical relay for amplifying bioluminescence. The C-terminus of the E2GFP
was fused to the *Renilla* luciferase variant RLuc8, which emits at 405 nm when supplied with
bisdeoxycoelenterazine (CTZ 400a). Through a two-step resonant energy transfer cascade
triggered by acidification in the presence of luciferin, pHIL translates the hyperactivity-induced
intracellular acidosis into silencing of neuronal activity. This was demonstrated *in vivo* by
mitigating acute tonic-clonic seizures induced by pilocarpine and by counteracting audiogenic
seizures in the PRoline-Rich Transmembrane protein 2 knockout (PRRT2 KO) mouse, a model
of genetic epilepsy. While pHIL is designed to specifically convert neuronal hyperactivity into
silencing, BLADe includes and transcends this application of real-time feedback. Co-expression
of the BLADe LMC integrator with an inhibitory or excitatory opsin resulted in neural activity-driven
hyperpolarization or additional depolarization, respectively, of the individual cell and neural
networks.
520
The development of tools for all-molecular real-time feedback regulation in neuroscience is
motivated by the desire to replace problematic "externally closed-loop" systems by "internal
closed-loop sensor-actuators" working at the level of individual neurons. Current solutions are
based on implanted hardware and computer-based detection algorithms to conduct detection and
deliver control. For example, in neuroscience approaches to real-time detection and feedback
control for altering maladaptive activity patterns such as overly exuberant neural bursting rely on
signals being tracked using electrodes, and patterns detected by computer algorithms that trigger
countermanding stimulation through implanted electrodes or fiber optics for electrical or
optogenetic regulation[35,36]. This strategy has shown success, for example in stopping epileptic
seizures dependent on thalamic bursting with optogenetic drive[37]. These closed-loop approaches
for dynamic brain regulation are promising but have key limitations. These methods require
chronic implants to detect patterns and deliver stimuli, and electrical stimuli or injected chemicals
impact all adjacent cells. An ideal strategy would not require implants, would only regulate cells
when they express maladaptive patterns, and would specifically regulate only those cells
exhibiting aberrant patterns, not the entire field of cells near an electrode.
536
The BL-ADe platform provides a solution to dynamic control, detecting calcium increases to drive
feedback through local opsins with cellular precision. By providing real-time feedback control,
driving opsins only when a cell is active, it is a unique way to sense such dynamics and locally
and immediately alter their trajectory within a single neuron. This control mechanism can enhance
behaviorally relevant neural activity patterns (e.g., sensory responses), but only when the cells
are activated above a threshold. This method can allow triggering based on underlying dynamics
in specific cell types in specific foci, for example, briefly increasing the firing of only those
pyramidal neurons that burst, to test if their increased firing duration enhanced sensory detection.
This approach is also ideal for directly targeting dynamical processes. As an example, using
BLADe to trigger enhanced hyperpolarization (or, depolarization) only after individual neurons
show a large calcium event is ideal for altering the operative mechanism to extend or suppress
macro-scale phenomena within tissue.
549
Here we developed a highly versatile Bioluminescent Activity Dependent (BLADe) platform for
converting intracellular activity to photoreceptor activation. As proof-of-principle, our data show
that a Ca^{2+} dependent luciferase can integrate Ca^{2+} fluctuations to drive transcription and change
membrane potential. The BLADe platform can be furthered by engineering Ca^{2+} dependent
luciferases with higher light emission, faster kinetics, and expanded dynamic ranges for Ca^{2+}
sensing. Beyond these proof-of-principle applications, Ca^{2+} signals are ubiquitous in biology and
essential for life, driving, for example, contraction in muscles and insulin release in pancreatic
cells, in addition to information processing in neurons. Importantly, the modularity of the BLADe

framework allows Ca²⁺-dependent light activation to be coupled to a wide range of light-sensing
actuators, providing experimental control over diverse molecular outcomes depending on the
user's goal. Moreover, the luciferase can be split by different sensing domains, providing a
general mechanism for cells to detect their own biochemical state and then drive discrete user-
defined outcomes. Further developed and improved versions of these tools are expected to have
broad applications. In basic research, this paradigm allows investigators to selectively detect, and
then suppress or amplify, specific biological events, a unique approach to understanding the role
of those processes. In potential therapeutic applications it allows sensing and discontinuing
aberrant activity before it can cause harm, or stopping pathological processes from continuing, or
amplifying and rescuing failing processes, without the use of implanted devices and regulated by
simple peripheral injection of a luciferin.
569
570
571 **MATERIALS AND METHODS**
572
573 **Chemicals**
574 Coelenterazine (CTZ) for in vitro (#303) and in vivo (#3031) experiments was purchased from
575 Nanolight Technologies. The Calcium Calibration Buffer Kit #1 from Life Technologies was used
576 to prepare Zero Ca²⁺ buffer and 39 µM Ca²⁺ buffer. All other chemicals, including Ionomycin and
577 Histamine, were purchased from Sigma.
578
Plasmids
A pcDNA3.1 backbone with the CMV promoter was used to insert gBlocks (IDT) that encode for
N-sbGluc, CaM-M13 variants, C-sbGluc and p2a dTomato by Gibson cloning (New England
Biolabs HiFi DNA Master Mix). Coding sequences from plasmids for GCaMP6f, hGtACR2,
ChR2(C128S), and CheRiff (Addgene) and for BlueCaMBI and GLICO (synthesized by Genscript)
were cloned into the pcDNA3.1-CMV backbone. Expression plasmids for VP-EL222 and 5xC120-
FireflyLuc were kindly provided by Dr. Kevin Gardner, CUNY, New York, NY, and for GeNL(Ca2+)
by Dr. Takeharu Nagai (Addgene). For viral vectors, coding sequences were cloned into a pAAV
vector downstream of a CAG or hSyn promoter, or into a pAAV-EfIa-DIO construct. For list of
plasmids used see Supplementary Table 1.
589
pcDNA3.1-CAG-VP-EL222 and pcDNA3.1-5xC120-FireflyLuc were kindly provided by Dr. Kevin
Gardner, CUNY, New York, NY
pAAV.Syn.GCaMP6f.WPRE.SV40 was a gift from Douglas Kim & GENIE Project (Addgene
plasmid # 100837 ; http://n2t.net/addgene:100837 ; RRID:Addgene_100837)
GeNL(Ca²⁺)_520/pcDNA3 was a gift from Takeharu Nagai (Addgene plasmid # 85204 ;
http://n2t.net/addgene:85204 ; RRID:Addgene_85204)
pLenti-CaMKIIa-hChR2(C128S)-EYFP-WPRE was a gift from Karl Deisseroth (Addgene plasmid
20294 ; http://n2t.net/addgene:20294 ; RRID:Addgene_20294)
pFUGW-hGtACR2-EYFP was a gift from John Spudich (Addgene plasmid # 67877 ;
http://n2t.net/addgene:67877 ; RRID:Addgene_67877)
pAAV-hSyn-CheRiff-eGFP was a gift from Adam Cohen (Addgene plasmid # 51697 ;
http://n2t.net/addgene:51697 ; RRID:Addgene_51697)
602
603 **AAV**
604 AAV2/9 preparations were generated by triple lipofection of HEK293-FT cells and harvesting viral
605 particles as previously described[5].
606
607
608

Animals
All experiments involving animals were carried out following the guidelines and protocols
approved by the Institutional Animal Care and Use Committee at the participating universities and
were in compliance with the US National Research Council's Guide for the Care and Use of
Laboratory Animals, the US Public Health Service's Policy on Humane Care and Use of
Laboratory Animals, and Guide for the Care and Use of Laboratory Animals. Mice were group-
housed in ventilated cages under 12-hour reverse light cycle, provided with tap water and
standard chow and allowed to feed ad libitum. C57/BL6 (JAX #000664), Emx1-Cre (JAX#
005628), and Swiss Webster (Charles River) mice of both sexes were used.
618
Cell Culture and Cell Transformation
HEK293 and HeLa cells were obtained from ATCC (American Type Culture Collection),
expanded, and aliquots at low passage numbers were frozen. Cells were grown in Gibco DMEM
media supplemented with 10% Fetal Bovine Serum, 1% Glutamax, 0.5 % Pen-Sterp, 1% Non-
Essential Amino Acids and 1% sodium pyruvate. Cells were cultured at 37°C and 5% atmospheric
carbon dioxide. Cells were transfected with Lipofectamine 2000 or 3000 in accordance with the
manufacturer's instructions.
626
Primary E18 rat hippocampal neurons were prepared from tissue shipped from BrainBits
(Transnetyx) following the vendor's protocol. Neurons were grown in Gibco Neurobasal Media
supplemented with 2% B27 supplement and 0.1% gentamycin and 1% Glutamax. Neurons were
nucleofected with plasmids of interest using the Lonza 2b Nucleofector and Rat Neuron
Nucleofector Kit (Lonza, VPG-1003) according to the manufacturer's instruction, then seeded. Or
neurons were nucleofected, seeded, then transduced with AAV the next day. Neurons were
seeded on PDL-coated glass coverslip (Neuvitro) in 12-well tissue culture plates (1 x 10^5 neurons
per well) or were plated on the electrode area of 1-well MEA dishes (60MEA200/30iR-Ti; Multi
Channel Systems, Germany) coated with PEI (0.1%) and laminin (50µg/ml) (1x10^5 neurons/10
µL/well) as described in detail in Prakash 2020[26].
637
Primary cardiomyocytes were harvested from E18 Swiss Webster mice, papain digested and
plated on 35mm glass bottom dishes (MatTek) for real time bioluminescence imaging. Primary
cardiomyocytes were grown in Claycomb Media supplemented with 10% HL-1 screened FBS,
100µg/mL penicillin/streptomycin, 0.1mM norepinephrine and 2% Glutamax. Cells were
nucleofected with plasmids of interest using the Lonza 2b Nucleofector and Rat cardiomyocyte
Nucleofector Kit (Lonza, VAPE-1002) according to the manufacturer's instruction.
644
Microscopy
Initially, all transfected cells were imaged using a Zeiss Axio Observer A1 microscope with a LD
A-Plan 20x/0 air objective and a Hamamatsu Orca Flash 4.0 CMOS camera to confirm expression
by fluorescence microscopy at 50ms exposure except for LMCer, which was imaged at a 10
seconds exposure. Confirmation of subcellular localization and imaging of brain sections were
done on an Olympus Fluoview 300 CLSM confocal microscope with a 60x/1.3 Oil objective.
651
Bioluminescence Imaging
In vitro
For photon counting HEK293 and HeLa cells were seeded on Costar 96 well clear bottom white
plates and luminescence was measured in a SpectraMax® L luminometer (Molecular
DevicesTM). For real-time bioluminescence imaging we used a Zeiss Axio Observer A1
microscope with a Fluar 40x/1.3 Oil objective with either a Hamamatsu Orca Flash 4.0 CMOS
camera or an Andor iXon Ultra 888 EM-CCD camera with their respective softwares. HEK293
and HeLa cells were imaged in imaging solution (CaCl2 1.25mM, HEPES 19.7mM, KCl 4.7mM,

KH2PO4 1.2mM, MgSO4 1mM, NaCl 130mM, dextrose 0.5, pH 7.2-7.4) with either 100µM or 12.5
µM CTZ. Primary cardiomyocytes were imaged in Tyrode's solution (NaCl 137mM, KCl 2.7mM,
MgCl2 1mM, CaCl2 1.8mM, Na2HPO4 0.2mM, NaHCO3 12mM, D-glucose 5.5mM, pH 7.2-7.4)
supplemented with 10µM norepinephrine and 100 µM CTZ.
664
In vivo
Brain: AAV2/9-hSyn-LMC4f-P2A-dTomato (450 nl) was injected across three locations in left
primary somatosensory (SI) cortex (150 nl per site) as described in detail in Gomez-Ramirez
2020[38]. Water-soluble CTZ (Nanolight Technology #3031) was diluted to yield a concentration of
2.36 mM. CTZ and NMDA injections were done directly in cortex. Bioluminescence was measured
using an electron multiplier charge coupled device (EMCCD) camera (Ixon 888, Andor) attached
to a Navitar Zoom 6000 lens system (Navitar, 0.5× lens). Images were collected in a custom-
made light-tight chamber with an exposure time of 10 s, and the EM gain set to 30. Imaging data
were recorded using the Solis image acquisition and analysis software (Solis 4.29, Andor).
674
Electrophysiology recordings from primary neurons
Neurons (rat E18, cortical) were co-nucleofected with LMC4f and either an inhibitory opsin,
hGtACR2, an excitatory opsin, ChR2(C128S), or a "Dud" opsin, ChR2(C128S)-E97R-D253A.
Post nucleofection, neurons were plated on laminin-coated glass coverslips (15mm, Neuvitro) in
culture medium consisting of Neurobasal Medium (Gibco # 21103-049), B-27 supplement (Gibco
17504-044), 2 mM Glutamax (Gibco # 35050-061), and 5% Fetal Calf Serum (FCS). The
following day, the medium was replaced with serum-free medium (NB-Plain medium). Half of the
medium was replaced with fresh NB-Plain medium every 3–4 days thereafter. Neurons were used
for whole cell patch clamp recordings between DIVs 21–25. For patch clamp recording, a
coverslip was transferred to a recording chamber mounted on an upright microscope (BX51WI,
Olympus) and perfused with aCSF containing (in mM): 121 NaCl, 2.8 KCl, 1 NaH_2PO_4, 26
$NaHCO_3$, 2 $CaCl_2$, 2 $MgCl_2$ and 15 D-glucose (310 mOsm/kg, pH 7.3-7.4) at a rate of 1.5 ml/min.
All solutions were bubbled with a gas mixture of 95% O_2 and 5% CO_2. Whole-cell patch clamp
recordings were performed using a Multiclamp 700b amplifier and Digidata 1440 digitizer together
with the pClamp recording software (Molecular Devices). Borosilicate glass micropipettes were
manufactured using a PC-100 puller (Narishige) and had resistances of 3–5 MΩ. In current clamp
recordings, pipettes were filled with intracellular solution containing (in mM): 130 K-gluconate, 10
KCl, 15 HEPES, 5 Na_2-phosphocreatine, 4 Mg-ATP and 0.3 Na-GTP (310 mOsm/kg, pH 7.3).
The aCSF was supplemented with D-AP5 (50 µM), CNQX (15 µM) and picrotoxin (100 µM) to
block fast glutamatergic and GABAergic synaptic transmission. Rs was compensated by using
bridge balance. Firing frequencies were calculated from the total number of action potentials
produced per unit time during depolarizing current injections (10 sweeps, 1.5 sec duration) in
episodic stimulation acquisition mode, at a membrane potential of -70mV, before and after CTZ
(100 µM) or vehicle treatments.
699
MEA Recordings
Once the neurons were matured (DIV14-DIV19), only those were included in the analysis that
were consistently spiking and were positive for expression of their respective opsins as tested by
shining blue LED light (470nm) on the culture and recording increase or decrease of spiking. Total
recording time was 900 seconds. MEAs were subjected to 10µM CTZ at around 600 seconds.
Experimental cultures expressed both LMC4f and an opsin (ChR2(C128S) or hGtARC2), while
control MEAs only expressed the opsin. MC Rack software was used for data acquisition. All MEA
analysis was done offline with MC Rack software (MultiChannel Systems; RRID:SCR_014955)
and NeuroExplorer (RRID:SCR_001818) as described in detail in Prakash et al., 2020[26].
709
710

Transcription *in vitro*

HeLa cells were used to test whether LMC-driven light emission could activate transcription through the light-responsive transcription factor EL222. Cells were plated in 6-well plates and transfected at ~80–90% confluency using Lipofectamine 2000 (Thermo Fisher). Each well received a total of 2 µg DNA, composed of the following: 666 ng EL222 (light-sensitive transcription factor), 333 ng 5×C120-FLuc (EL222-responsive firefly luciferase reporter), and 1000 ng LMC4f-dTomato (calcium-dependent luciferase). Parallel wells were transfected with other calcium-dependent bioluminescent sensors (GeNL, GLICO, BlueCaMBI) and matched FLuc reporters.

Three hours after transfection, cells were trypsinized (300 µL trypsin per well) and replated into PDL-coated, white-walled, clear-bottom 96-well plates (~100,000 cells per well). Cells were allowed to adhere for 9–12 hours before stimulation. For transcriptional activation, 100 µL of stimulation buffer was added per well (1:1 with cell media) for a final well volume of 200 µL. The stimulation solution contained 100 µM CTZ or hCTZ (depending on the sensor), with or without 10 µM histamine. Vehicle controls received no treatment. After stimulation, cells were incubated for 7–9 hours before transcriptional reporter readout.

For FLuc transcriptional measurement, activity was recorded 18 hours after stimulation using a luminometer. Prior to measurement, cells were washed and incubated in phenol red–free media (100 µL per well). D-luciferin (final 150 µg/mL) was prepared from a frozen 60 mg/mL stock by serial dilution into Opti-MEM. For each well, 25 µL of 750 µg/mL D-luciferin working stock was injected into 100 µL of media. FLuc bioluminescence was recorded immediately after injection.

Transcription *in vivo*

To test bioluminescent dependent transcription *in vivo*, we co-injected three AAV constructs into the prefrontal cortex of anesthetized mice: AAV9-hSyn-LMC4f-P2A-dTomato (Ca²⁺-sensitive luciferase), AAV9-CAG-EL222 (light-activated transcription factor), and AAV9-5×C120-EYFP (EL222-responsive reporter). Viral solutions were mixed prior to injection to a total volume of 1 µL per site, yielding final per-virus titers of: LMC4f (1×10^{10} GC), EL222 (0.7×10^{10} GC), and C120-EYFP (0.3×10^{10} GC). The injection was targeted to the left medial prefrontal cortex at +0.37 mm anterior to bregma, 3.0 mm lateral. Following injection, the craniotomy was sealed to minimize ambient light exposure during recovery.

After 7 days, mice received intraventricular (ICV) injections of one of three treatments: (1) NMDA + vehicle, (2) luciferin (CTZ) alone, or (3) luciferin + NMDA. These conditions were used to assess background transcription, spontaneous activity-driven transcription, and full BLADe-dependent transcriptional gating, respectively. Injections were performed using a 26-gauge needle and 10 µL Hamilton syringe, targeting the right lateral ventricle (AP: –0.5 mm, ML: –1.1 mm, DV: –2.0 mm from bregma). The luciferin was prepared fresh by dissolving coelenterazine (NanoLight #3031) in sterile water to a stock concentration of 1.6 mM, then when injected was assumed to be diluted 1:8 in CSF within the ventricle to yield a final injection concentration of 200 µM. NMDA was prepared at a working concentration of 75 ng/µL in sterile water. For combined delivery, 1 µL of NMDA solution (75 ng) was mixed with 4 µL of luciferin solution per injection (final 5 µL per mouse, injected over 5mins). Solutions were prepared under minimal light and protected from exposure throughout. After injection, mice were removed from the stereotaxic apparatus and allowed to recover. Brains were collected for histological analysis ~20 hours after ICV injection. Mice were anesthetized and transcardially perfused with PBS followed by 4% paraformaldehyde. Coronal brain sections were collected and imaged by confocal microscopy to quantify EYFP reporter expression.

To analyze calcium- and light-dependent transcriptional activation *in vivo*, we quantified EYFP reporter expression in dTomato-labeled neurons across experimental groups. Confocal images from all treatment conditions (NMDA-only, luciferin-only, luciferin + NMDA) were analyzed. First, dTomato+ images were pooled and segmented from maximum intensity projections using the MATLAB 'cell-segm' thresholding and size-filtering algorithm[39], ensuring unbiased detection across all fields of view. The resulting binary masks were then applied to the corresponding EYFP images to extract raw per-cell reporter fluorescence from each dTomato+ neuron. Finally, cells were grouped by treatment, and the ratio of EYFP to dTomato fluorescence was computed per cell to normalize for expression variability.

Electrophysiology *in vivo*
To express LMC and CheRiff in excitatory cortical neurons, we injected AAVs into the left virbissa primary somatosensory cortex (vSI) of Emx1-Cre mice (3–4 month old). AAV1-EF1a-DIO-LMC4f and AAV1-EF1a-DIO-CheRiff-EYFP were mixed 1:1 and delivered in a single injection (1 µL total) using a 33-gauge Nanofil syringe. Using a dental drill, a burr holes was made for AAV injection. The injection site was located 1.5 mm posterior and 3.0 mm lateral to bregma, and the needle was lowered to a depth of 500–1000 µm below the pial surface. Virus was infused at a rate of 50 nL/min over ~20 minutes. Following injection, the needle was held in place for 10 min, then raised to 200 µm and held for an additional 10 min before full withdrawal to allow the virus to spread across the layers of the cortex. During the same surgery, a custom metal headpost was implanted for head fixation. After exposing the skull, a thin layer of clear Metabond was applied to stabilize the surface while keeping bregma and surface vasculature visible. The headpost was then affixed using additional Metabond, positioned off-center to leave vSI accessible. The short arm was angled over the left ear to allow right-side whisker stimulation. After injection and headpost implantation, the craniotomy was covered with mineral oil and sealed with Metabond. Animals recovered for at least 3 weeks before recordings.

After 2–3 weeks of recovery, mice were anesthetized with isoflurane (1.2–1.4%) and head-fixed. The craniotomy used for AAV injection was reopened and expanded to 1 mm to expose the underlying cortex and target the same vSI region as the viral delivery. Throughout the procedure, the brain remained covered in carbogenated ACSF (95% O_2 / 5% CO_2) under constant perfusion. Animals were placed in a light-tight, electrically shielded chamber and maintained under anesthesia. Mice were secured in place to allow perfusion tubing to be positioned next to the craniotomy, allowing for constant aCSF inflow (for bioluminescence CTZ was diluted to 50 µM in aCSF). All macrovibrissae on the right side of the mouse were secured ~3mm from the mystacial pad. A custom-made clamp was glued to a piezoelectric bender (Noliac CMBP09), positioned to move all macrovibrissae in the caudorostral direction, with a half-sine wave velocity profile with a rising phase (6ms) and a slower relaxation phase (20ms). On each trial, 20Hz vibratory vibrissal stimulus trains (10 deflections, 500ms) were delivered for 1000 trials total. On a trial by trial and randomized basis, the stimulus amplitude varied between 0 to maximal amplitude (~1mm deflection). For each recording session, aCSF was constantly perfused for the first 500 trials and luciferin (CTZ) was then perfused for the remaining 500 trials.

All recordings in barrel cortex were conducted under light isoflurane anesthesia. Laminar probes consisted of single shank,32-channel silicon probes with a fiber optic 50µm above the highest recording site (A1x32 Poly2-5mm-50s-177-OA32LP, Neuronexus Technologies; 0.15mm silver wire reference). Data was sampled at 30kHz and passed through a digital amplifier (Cereplex-µ, Blackrock Microsystems), and directed through HDMI to the Cereplex Direct data acquisition box (Blackrock Microsystems). The 32-channel linear silicon probe was inserted perpendicularly into cortex using a micromanipulator at ~10 µm/s until the top contact of the electrode disappeared beneath the pial surface. The probe was allowed to settle for ~30–60 minutes before recording.

Luciferin (coelenterazine; NanoLight #3031) was freshly prepared in sterile water (1 µg/mL) and
diluted into ACSF to a final concentration of 20 µM. Following an initial recording period under
vehicle ACSF, the perfusion line was switched to CTZ-containing ACSF to enable continuous
bath application during the second half of the session. Flow rate and volume were monitored to
maintain complete coverage of the craniotomy and probe without overflow.

Raw electrophysiological data were downsampled to 10 kHz. For each recording, electrode
contacts with RMS values exceeding three times the interquartile range above the 75th percentile
or below the 25th percentile across the 32-channel array were flagged as noise and removed for
further analysis. All remaining electrodes were re-referenced to the common average[40]. MUA was
estimated by converting raw spike times into instantaneous firing rates using 1 ms bins. For
luciferin experiments, stimulus-aligned firing rates were extracted from trials before and after the
defined luciferin application time. Channels were classified as luciferin-responsive if the 99%
confidence interval of the post-luciferin firing rate (bootstrapped, n = 1000) did not overlap with
the baseline distribution. For sensory stimulation experiments, MUA peristimulus time histograms
(PSTHs) were constructed by aligning spike data to stimulus onset and binning at 1 ms. PSTHs
were smoothed using a Gaussian kernel (window = 20 ms, SD = 3 ms) and averaged across trials.
PSTHs were computed separately for pre- and post-luciferin trial blocks.

Statistics
All analysis was carried out using MATLAB and SPSS software. Data are displayed as
mean±standard error of the mean (SEM). We ran two-way repeated measures and One-way
ANOVA. Tukey's post hoc was run where significant main effect was found.

Electrophysiological in vivo data was high pass filtered at 250Hz to extract spikes. Spike data was
thresholded at -63µV and sorted for each channel based on waveform characteristics using
Principal Components Analysis (PCA). Spikes were binned to calculate frequency of firing over
time. Differences between groups were assessed using two-way repeated measures ANOVAs
(repeat trials per mouse). For ex vivo electrophysiological data, statistical significance between
groups was determined using Student's T-test.

REFERENCES
1.   Love, A. C. & Prescher, J. A. Seeing (and Using) the Light: Recent Developments in
Bioluminescence Technology. *Cell Chemical Biology* vol. 27 904–920 at
https://doi.org/10.1016/j.chembiol.2020.07.022 (2020).
2.   Rost, B. R., Schneider-Warme, F., Schmitz, D. & Hegemann, P. Optogenetic Tools for
Subcellular Applications in Neuroscience. *Neuron* **96**, 572–603 (2017).
3.   Losi, A., Gardner, K. H. & Moglich, A. Blue-Light Receptors for Optogenetics. *Chem. Rev.*
118, 10659–10709 (2018).
4.   Berglund, K., Birkner, E., Augustine, G. J. & Hochgeschwender, U. Light-Emitting
Channelrhodopsins for Combined Optogenetic and Chemical-Genetic Control of
Neurons. *PLoS One* **8**, e59759 (2013).
5.   Berglund, K. *et al.* Luminopsins integrate opto- and chemogenetics by using physical and
biological light sources for opsin activation. *Proc. Natl. Acad. Sci. U. S. A.* **113**, E358–
E367 (2016).
6.   Tung, J. K., Gutekunst, C.-A. & Gross, R. E. Inhibitory luminopsins: genetically-encoded
bioluminescent opsins for versatile, scalable, and hardware-independent optogenetic
inhibition. *Sci. Rep.* **5**, 14366 (2015).
7.   Crespo, E. L., Bjorefeldt, A., Prakash, M. & Hochgeschwender, U. Bioluminescent
Optogenetics 2.0: Harnessing Bioluminescence to Activate Photosensory Proteins In
Vitro and In Vivo. *J. Vis. Exp.* (2021) doi:10.3791/62850.

8.   Li, T. *et al.* A synthetic BRET-based optogenetic device for pulsatile transgene
expression enabling glucose homeostasis in mice. *Nat. Commun.* **12**, (2021).

9.   Parag-Sharma, K. *et al.* Engineered BRET-Based Biologic Light Sources Enable
Spatiotemporal Control over Diverse Optogenetic Systems. *ACS Synth. Biol.* (2020)
doi:10.1021/acssynbio.9b00277.

10.   Naim, N. *et al.* Luminescence-activated nucleotide cyclase regulates spatial and temporal
cAMP synthesis. *J. Biol. Chem.* (2019) doi:10.1074/jbc.AC118.004905.

11.   Kim, E. H. *et al.* Self-luminescent photodynamic therapy using breast cancer targeted
proteins. *Sci. Adv.* **6**, (2020).

12.   Shramova, E. I., Frolova, A. Y., Filimonova, V. P., Deyev, S. M. & Proshkina, G. M.
System for Self-excited Targeted Photodynamic Therapy Based on the Multimodal
Protein DARP-NanoLuc-SOPP3. *Acta Naturae* **15**, 100–110 (2023).

13.   Yeh, H.-W. & Ai, H.-W. Development and Applications of Bioluminescent and
Chemiluminescent Reporters and Biosensors. *Annu. Rev. Anal. Chem. (Palo Alto. Calif).*
12, 129–150 (2019).

14.   Welsh, J. P., Patel, K. G., Manthiram, K. & Swartz, J. R. Multiply mutated Gaussia
luciferases provide prolonged and intense bioluminescence. *Biochem. Biophys. Res.*
Commun. **389**, 563–8 (2009).

15.   Kim, S. B., Sato, M. & Tao, H. Split Gaussia luciferase-based bioluminescence template
for tracing protein dynamics in living cells. *Anal. Chem.* **81**, 67–74 (2009).

16.   Saito, K. *et al.* Luminescent proteins for high-speed single-cell and whole-body imaging.
Nat. Commun. **3**, 1262 (2012).

17.   Chen, T.-W. *et al.* Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature*
499, 295–300 (2013).

18.   Suzuki, K. *et al.* Five colour variants of bright luminescent protein for real-time multicolour
bioimaging. *Nat. Commun.* **7**, 13718 (2016).

19.   Oh, Y. *et al.* An orange calcium-modulated bioluminescent indicator for non-invasive
activity imaging. *Nat. Chem. Biol.* **15**, 433–436 (2019).

20.   Farhana, I., Hossain, M. N., Suzuki, K., Matsuda, T. & Nagai, T. Genetically Encoded
Fluorescence/Bioluminescence Bimodal Indicators for Ca(2+) Imaging. *ACS sensors* **4**,
1825–1834 (2019).

21.   Motta-Mena, L. B. *et al.* An optogenetic gene expression system with rapid activation and
deactivation kinetics. *Nat. Chem. Biol.* (2014) doi:10.1038/nchembio.1430.

22.   Berndt, A., Yizhar, O., Gunaydin, L. A., Hegemann, P. & Deisseroth, K. Bi-stable neural
state switches. *Nat. Neurosci.* **12**, 229–34 (2009).

23.   Berglund, K., Fernandez, A. M., Gutekunst, C. A. N., Hochgeschwender, U. & Gross, R.
E. Step-function luminopsins for bimodal prolonged neuromodulation. *J. Neurosci. Res.*
98, 422–436 (2020).

24.   Prakash, M. *et al.* Selective control of synaptically-connected circuit elements by all-
optical synapses. *Commun. Biol.* **5**, 33 (2022).

25.   Govorunova, E. G., Sineshchekov, O. A., Janz, R., Liu, X. & Spudich, J. L. Natural light-
gated anion channels: A family of microbial rhodopsins for advanced optogenetics.
Science (80-.). **349**, 647–50 (2015).

26.   Prakash, M., Medendorp, W. E. & Hochgeschwender, U. Defining parameters of
specificity for bioluminescent optogenetic activation of neurons using in vitro multi
electrode arrays (MEA). *J. Neurosci. Res.* **98**, 437–447 (2020).

27.   Hochbaum, D. R. *et al.* All-optical electrophysiology in mammalian neurons using
engineered microbial rhodopsins. *Nat. Methods* **11**, 825–833 (2014).

28.   Yang, J. *et al.* Coupling optogenetic stimulation with NanoLuc-based luminescence
(BRET) Ca(++) sensing. *Nat. Commun.* **7**, 13268 (2016).

29.   Qian, Y., Rancic, V., Wu, J., Ballanyi, K. & Campbell, R. E. A Bioluminescent Ca(2+)

Indicator Based on a Topological Variant of GCaMP6s. *Chembiochem* **20**, 516–520
(2019).
30.   Lambert, G. G. *et al.* CaBLAM! A high-contrast bioluminescent Ca(2+) indicator derived
from an engineered Oplophorus gracilirostris luciferase. *bioRxiv* (2025)
doi:10.1101/2023.06.25.546478.
31.   Lee, D., Hyun, J. H., Jung, K., Hannan, P. & Kwon, H.-B. A calcium- and light-gated
switch to induce gene expression in activated neurons. *Nat. Biotechnol.* **35**, 858–863
(2017).
32.   Wang, W. *et al.* A light- A nd calcium-gated transcription factor for imaging and
manipulating activated neurons. *Nat. Biotechnol.* **35**, 864–871 (2017).
33.   O'Neill, B. K. & Laughlin, S. T. Neuronal Calcium Recording with an Engineered TEV
Protease. *ACS chemical biology* vol. 13 1159–1164 at
https://doi.org/10.1021/acschembio.8b00130 (2018).
34.   Merolla, A. *et al.* A pH-sensitive closed-loop nanomachine to control hyperexcitability at
the single neuron level. *Nat. Commun.* **15**, 5609 (2024).
35.   Novitskaya, Y., Sara, S. J., Logothetis, N. K. & Eschenko, O. Ripple-triggered stimulation
of the locus coeruleus during post-learning sleep disrupts ripple/spindle coupling and
impairs memory consolidation. *Learn. Mem.* **23**, 238–248 (2016).
36.   Lustenberger, C. *et al.* Feedback-Controlled Transcranial Alternating Current Stimulation
Reveals a Functional Role of Sleep Spindles in Motor Memory Consolidation. *Curr. Biol.*
26, 2127–2136 (2016).
37.   Paz, J. T. *et al.* Closed-loop optogenetic control of thalamus as a tool for interrupting
seizures after cortical injury. *Nat. Neurosci.* **16**, 64–70 (2013).
38.   Gomez-Ramirez, M., More, A. I., Friedman, N. G., Hochgeschwender, U. & Moore, C. I.
The BioLuminescent-OptoGenetic in vivo response to coelenterazine is proportional,
sensitive, and specific in neocortex. *J. Neurosci. Res.* **98**, 471–480 (2020).
39.   Hodneland, E., Kögel, T., Frei, D. M., Gerdes, H.-H. & Lundervold, A. CellSegm - a
MATLAB toolbox for high-throughput 3D cell segmentation. *Source Code Biol. Med.* **8**, 16
(2013).
40.   Ludwig, K. A. *et al.* Using a common average reference to improve cortical neuron
recordings from microelectrode arrays. *J. Neurophysiol.* **101**, 1679–1689 (2009).

Acknowledgments

We would like to thank all members of Bioluminescence Hub laboratories for their feedback, discussions, and thoughtful comments throughout the progression of this work. (http://www.bioluminescencehub.org/)

Competing interests

The authors have no conflicts of interest to declare.

Author contributions

Conceptualization, AP, ELC, NCS, CIM, UH; Validation, AP, ELC, CIM, DL, NCS, UH; Formal Analysis, AP, ELC; Investigation, AP, ELC, MP, ADS, ZZ, MG-R, MOT; Data Curation, AP, ELC; Writing—Original Draft, AP, ELC; Writing—Review & Editing, ELC, NCS, CIM, UH; Visualization, AP, ELC; Supervision, CIM, NCS, UH; Project Administration, UH; Funding Acquisition, CIM, DL, NCS, UH.

Funding

This work was supported by National Institutes of Health grants R21MH101525, R21EY026427, U01NS099709, National Science Foundation grants CBET-1464686, DBI-1707352, and the W.M. Keck Foundation.

Figure 1. **Design of a *Gaussia* luciferase based Ca²⁺ sensor. a**. sbGLuc (light blue) is split by a CaM–M13 module (purple), producing a dim basal signal in the presence of its luciferin (CTZ). Binding of Ca²⁺ (yellow) reconstitutes an active luciferase and increases light emission in the presence of luciferin. **b**. Elements of the coding sequence for LMC: *Gaussia* luciferase signal peptide (red), N- and C-terminal fragments of sbGLuc (light blue), calmodulin sequence from GCaMP6f (purple), M13 sequence (purple), and various linker regions (gray). **c**. Structure model of LMC. **d.** Emission wavelength across LMC variants (4-8; for color code see e.). The LMC constructs were transiently expressed in HeLa cells and light emission after luciferin and histamine addition was measured across the wavelength spectra (data presented as mean ± SEM). **e.** Change in luminescence in the presence of luciferin in response to 10 μM histamine in HeLa cells across LMC variants (N=12 wells per sensor, Unpaired *t*-test was used to determine *p*-values, corrected for multiple comparisons with Holm-Šídák method: LMC4 vs LMC5: *P*=0.3712; LMC4 vs LMC6: *P*=0.6193; LMC4 vs LMC7: *P*= 0.3977; LMC4 vs LMC8: *P*= 0.3712). Boxplots show median, 25th and 75th percentiles (box edges), whiskers to most extreme data points, and individual outliers. **f.** Percent change in bioluminescence in HEK293 cells expressing LMC4 in response to 2 μM ionomycin under increasing extracellular Ca²⁺ concentrations. Cells were pre incubated with 5 μM luciferin and varying concentrations of Ca²⁺; acute Ca²⁺ flux was initiated by injection of ionomycin before photon counting in a luminometer (mean ± s.e.m). **g.** Response of LMC4 and GCaMP6f to acute Ca²⁺ flux. HEK293 cells expressing LMC4 or GCaMP6f were held in 39 μM extracellular Ca²⁺ concentration and ionomycin was added to 2μM final concentration. LMC4 expressing HEK cells were pre-incubated with CTZ at 5 μM final concentration. Bioluminescence readings for LMC4 were taken on a SpectraMax-L luminometer and fluorescent images for GCaMP6f were taken on a Zeiss AxioObserver A1 microscope with a Hamamatsu Ocra Flash4.0 CMOS camera before and after the addition of ionomycin. Graph displays percent change in signal before and after ionomycin induced Ca²⁺ flux (mean ± SEM).

Figure 2. Reporting Ca²⁺ flux *in vitro* and *in vivo*. a. Primary cardiomyocytes nucleofected with LMC and incubated with luciferin (100 µM CTZ). Pseudocolored images show regions of interest (ROIs) at three time points (t1–t3) before and after ATP (10 µM) application. **b.** Representative bioluminescence traces from the ROIs in (a). Orange and grey arrows mark the time points corresponding to t1–t3 in (a). **c.** Representative hippocampal neurons expressing LMC imaged with luciferin alone (left) or with luciferin and glutamate (right). **d.** Mean (blue) overlaid with individual bioluminescence traces (grey) of LMC responses to bath-applied luciferin and glutamate. **e.** Box plot showing maximal luminescence responses (Max ΔRLU/RLU₀) of LMC-expressing cells exposed to luciferin alone or to glutamate in the presence of luciferin (N= 29 neurons from two independent cultures, Wilcoxon signed rank test: *P*= 2.7023 x10⁻⁶). **f.** Maximum fold change (contrast) in response across neurons. **g.** *in vivo* imaging of mouse barrel cortex (vSI) expressing LMC. *left*: brightfield image showing injection site. *right:* Relative luminescence heatmap with ROIs for LMC (red) and control GLuc (black). Luciferin (CTZ, 50 µL) was delivered via the lateral ventricle, and NMDA (1 µL) was applied directly to cortex. **h**. Comparison of bioluminescence emission over area transduced with Ca2+ dependent luciferase, AAV-LMC4f (red box in image, red line in graph) versus area transduced with non-Ca2+ responsive luciferase, AAV-GLuc (black box in image, black line in graph). Representative traces from ROIs in (g) showing luminescence after luciferin (CTZ) and NMDA delivery. Onset of CTZ and NMDA injections are indicated. Boxplots show median, 25th and 75th percentiles (box edges), whiskers to most extreme data points, and individual outliers.

Figure 3. Integrating Ca^{2+} flux with activation of photoreceptors. a. Luminometer time series from HeLa cells transiently expressing BlueCaMBI, GLICO, GeNL, or LMC following addition of luciferin and histamine. Inset shows LMC traces on an extended scale. **b.** Peak bioluminescence measured after luciferin and histamine stimulation for each sensor from (a). **c.** Histamine-specific percent change in bioluminescence normalized to luciferin baseline, expressed as percent change for each calcium sensor. (N=4 wells/sensor, LMC vs GLICO: P= 4.65x10^{-6}, GeNL: P= 5.51x10^{-7}, BlueCAMBI: P= 1.46x10^{-6}; GeNL vs GLICO P= 0.38, BlueCAMBI P= 0.996; BlueCAMBI vs GLICO: P=0.723, Tukey's post hoc multiple-comparison's test following a one-way analysis of variance (ANOVA), F$_{(3, 12)}$ **=** 48.56, P**=** 5.51x10^{-7}). Boxplots show median, 25th and 75th percentiles (box edges), whiskers to most extreme data points, and individual outliers.

Figure 4. Converting intracellular activity to transcription. a. Schematic of LMC leading to bioluminescence mediated dimerization of the light sensing transcription factor EL222 in the presence of Ca^{2+} and luciferin (CTZ). The photoactivated EL222 homodimer then binds to the promoter 5xC120 and leads to transcription of the reporter gene. **b.** Firefly luciferase (FLuc) transcriptional reporter measured under dark, luciferin-only, or luciferin + histamine treatments in HeLa cells transiently expressing BlueCaMBI, GLICO, GeNL, or LMC. **c.** FLuc transcriptional reporter normalized to luciferin baseline expressed as percent change for each sensor/condition. **d.** Representative confocal images showing EYFP transcriptional reporter (green), LMC P2A-tdTomato viral expression (red), and nuclear DAPI stain (blue) from a field of view in mouse prefrontal cortex following intraventricular injection of luciferin and/or NMDA. Mixtures of AAV9 preparations of the 3 constructs were stereotaxically injected into the medial prefrontal cortex of mice. Three or more weeks later mice received injections of NMDA alone, CTZ alone, or CTZ and NMDA. Brains were collected 18 hours later and sections imaged under a confocal microscope. **e.** EYFP reporter expression quantified as per-cell EYFP/tdTomato ratio across treatment groups. Each point represents a single cell derived from three treatment groups: Luciferin-NMDA (N= 350 cells/2 mice), Luciferin-ONLY N= 197 cells/2 mice), NMDA-ONLY (N= 147 cells/2 mice). Dotted line indicates the 90th percentile EYFP/dTomato ratio observed in the NMDA-ONLY group, used as a threshold to define transcriptional activation above baseline. **f.** EYFP reporter intensity plotted against tdTomato fluorescence for each cell and condition. Boxplots show median, 25th and 75th percentiles (box edges), whiskers to most extreme data points, and individual outliers.

Figure 5. Integrating Ca²⁺ flux with change in membrane potential in single neurons *in vitro*. **a.** Brightfield and fluorescence images of cultured cortical neurons nucleofected with opsin (ChR2(C128S))-EYFP and LMC-P2A-dTomato and used for whole-cell patch-clamp recordings. The patch electrode is visible in the brightfield image. **b.** (i) Schematic of farnesylated LMC anchored to the inner cell membrane near the excitatory opsin ChR2(C128S) enabling light-mediated activation in the presence of depolarization-induced Ca²⁺ flux and the luciferin (CTZ). (ii) Representative membrane voltage responses to step current injections before (gray) and after (blue) bath perfusion of luciferin in a neuron co-expressing LMC and ChR2(C128S). (iii) Action potential counts under vehicle (gray) and luciferin (blue) conditions. Data expressed as mean ± SEM. (iv) Empirical cumulative distribution of spike fraction changes across voltage traces for vehicle and luciferin conditions, represented as the difference in spike counts between the first and second halves of the recording. **c.** (i) Schematic of a control condition, where LMC is co-expressed with ChR2(C128S) in the presence of Ca²⁺ but without luciferin, preventing light-mediated activation. (ii–iv) Same analyses as in b. **d.** (i) Schematic of LMC co-expressed with the inhibitory opsin hGtACR2, enabling light-mediated suppression of activity in the presence of Ca²⁺ and luciferin. (ii–iv) Same analyses as in b, showing inhibition rather than activation. Data are presented as mean ± SEM in panels.

1092

Figure 6. Integrating Ca²⁺+ flux with change in membrane potential in neural populations *in vitro*. a. Schematic of farnesylated LMC anchored to the inner cell membrane near the excitatory opsin ChR2(C128S) enabling light-mediated activation in the presence of depolarization-induced Ca²⁺ flux and the luciferin (CTZ). **b.** Fluorescence images of cultured cortical neurons co-transduced with LMC-P2A-dTomato (red) and ChR2(C128S)-EYFP (green). **c.** Representative filtered voltage traces from one channel before and after vehicle (top) or luciferin (bottom) application. Red dashed line indicates time of application. **d.** MUA firing rate was increased in the luciferin condition (Wilcoxon signed-rank test, $P= 3.93 \times 10^{-28}$, Holm-corrected). **e.** Burst rate was not significantly different between luciferin and vehicle conditions (Wilcoxon signed-rank test, $P= 0.63$, Holm-corrected). **f.** Burst duration was significantly prolonged in the luciferin condition (Wilcoxon signed-rank test, $P= 3.18 \times 10^{-49}$, Holm-corrected). **g.** Time-aligned MUA traces

across all electrodes showing normalized population responses following vehicle (top) or luciferin (bottom) application.
Individual electrode traces are shown in grey (vehicle) and blue (luciferin); thick lines represent the mean response per
condition. MUA was smoothed using a sliding 5-point boxcar window. **h.** Early-phase (0–50 ms) MUA firing rates
following application was not different between luciferin and vehicle (Wilcoxon signed-rank test, P= 0.13, Holm-
corrected). **i.** Late-phase (150–1000 ms) MUA firing rate was significantly higher in the luciferin condition (Wilcoxon
signed-rank test, P= 2.61 × 10^{-8}, Holm-corrected). **j.** Late phase normalized MUA was significantly increased under
luciferin (Wilcoxon signed-rank test, P= 1.02 × 10^{-11}, Holm-corrected). For **d-f** and **h-j**, N=57-59 pairs of electrodes per
recording pooled across three independent MEA cultures. Boxplots show median, 25th and 75th percentiles (box
edges), whiskers to most extreme data points, and individual outliers.

1111

1112
1113
Figure 7. Integrating spontaneous and sensory inputs into changes in firing rates *in vivo*. a. Schematic of the
experimental design. Extracellular electrophysiology recordings from SI barrel cortex during varied vibrissal deflections
(zero, low, mid, max intensities) for 500ms at 20Hz. **b**. Schematic of the experimental timeline. Each recording consisted
of 1000 trials (~2.5 hour sessions) where ACSF containing vehicle or CTZ was constantly perfused over the craniotomy.
During each treatment epoch, vibrissal deflections were randomly presented. In the presence of the sensory stimulus
and vehicle (grey box), there is no activity dependent light emission of LMC. In contrast, in the presence of CTZ and
the sensory stimulus (blue box), sensory recruitment leads to activity dependent bioluminescence and subsequently
photoactivation of opsins in neurons. **c**. MUA firing rate before and after the bath application of CTZ over the craniotomy.
d. MUA firing rates for responsive and non-responsive channels to optogenetic stimulation. *Inset:* the pattern LED
signal (teal), percent change of firing rate for responsive (blue) and non-responsive channels (grey). **e.** Peristimulus
time histograms from two distinct recording sites reflect changes in spiking activity evoked by contralateral sensory
driven recruitment during the vehicle (grey) and CTZ (blue) treatment periods during low amplitude vibrissae
stimulation. **f.** Mean multiunit activity responses to vibrissae stimulation during the early phase window, vehicle (grey)
vs CTZ (blue) epoch as a function of low, medium, and max vibrissae stimulus amplitude. *inset:* schematic of the analog
signal controlling the piezo stimulator, red box is the window of time where the mean was quantified. **g.** Same as (f),
but during the late phase window. All data are plotted as mean +/- SEM.
1130
1131

1132
1133
1134 **Supplemental Figure S1. Ribbon diagrams of GLuc indicating 9 alpha helices**. Left: Of the 9 alpha helices α4 and
1135 α7 form the most rigid block of the luciferase stabilized by a disulphide bond between Cysteine 65 and 77. Right: The
1136 Calcium sensing moiety, CaM-M13 was inserted between Q88 and G89. In its unbound state, the CaM-M13
1137 destabilizes the C65-C77 disuphide bond just by pushing the two halves away from each other. When bound to Ca^{2+},
1138 the CaM-M13 pulls on both halves and stabilizes the C65-C77 disulphide bond to render the luciferase functional.
1139
1140

a

sbGluc(M43L_M110L):

K[1]PTENNEDFNIVAVASNFATTDLDADRGKLPGKKLPLEVLKELEANARKAGCTRGCLIC
LSHIKC[65]TPKMKKFIPGRC[77]HTYEGDKESAQ[88]G[89]GIGEAIVDIPEIPGFKDLEPLEQF
IAQVDLCVDCTTGCLKGLANVQCSDLLKKWLPQRCATFASKIQGQVDKIKGAGGD[168]

b

LMC4: Signal Sequence; Gluc N half; CaM-M13; Gluc C half;
Alpha Helix; Linker

MGVKVLFALICIAVAEAK[1]PTENNEDFNIVAVASNFATTDLDADRGKLPGKKLPLEVLKELE
ANARKAGCTRGCLICLSHIKC[65]TPKMKKFIPGRC[77]HTYEGDKESAQ[88]GGGGTMADQLTEE
QIAEFKEEFSLFDKDGDGTITTKELGTVMRSLGQNPTEAELQDMINEVDADGDGTIDFPEF
LTMMARKMKYRDTEEEIREAFGVFDKDGNGYISAAELRHVMTNLGEKLTDEEVDEMIREAD
IDGDGQVNYEEFVQMMTAKGGKRRWKKNFIAVSAANRFKKISSSGALGSGGGG[89]GIGEAIV
DIPEIPGFKDLEPLEQFIAQVDLCVDCTTGCLKGLANVQCSDLLKKWLPQRCATFASKIQG
QVDKIKGAGGD[168]

Supplemental Figure S2. Amino acid sequences of sbGLuc and LMC4. **a**. Amino acid sequence of sbGLuc with landmark mutations of sbGLuc versus wildtype GLuc color coded. **b**. Amino acid sequence of LMC4 with landmarks indicated by colors and underlines.

a

LMC Linker-CaM-M13 Comparisons

```
LMC4  1  GGGGTMADQLTEEQIAEFKEEFSLFDKDGDGTITTKELGTVMRSLGQNPTEAELQDMINEVDADGDGTIDFPEFLTMMAR  80
LMC5  1  GGGGTLPDQLTEEQIAEFKEAFSLFDKDGDGTITTKELGTVMRSLGQNPTEAELQDMINEVDADGDGTIDFPEFLTMMAR  80
LMC6  1  GGGGTLPDQLTEEQIAEFKEAFSLFDKDGDGTITTKELGTVMRSLGQNPTEAELQDMINEVDADGDGTIDFPEFLTMMAR  80
LMC7  1  GGGGTMHDQLTEEQIAEFKEAFSLFDKDGDGTITTKELGTVMRSLGQNPTEAELQDMINEVDADGNGTIYFPEFLTMMAR  80
LMC8  1  GGGGTMHDQLTEEQIAEFKEAFSLFDKDGDGTITTRELGTVMRSLGQNPTEAELQDMINEVDADGNGTIYFPEFLTMMAR  80

LMC4 81  KMKYRDTEEEIREAFGVFDKDGNGYISAAELRHVMTNLGEKLTDEEVDEMIREADIDGDGQVNYEEFVQMMTAKGGKRRW  160
LMC5 81  KMKYRDTEEEIREAFGVFDKDGNGYISAAELRHVMTNLGEKLTDEEVDEMIREADIDGDGQVNYEEFVQMMTAKGGKRRW  160
LMC6 81  KGSYRDTEEEIREAFGVFDKDGNGYISAAELRHVMTNLGEKLTDEEVDEMIREADIDGDGQVNYEEFVQMMTAKGGKRRW  160
LMC7 81  KMKDTDSEEEIREAFRVFDKDGNGYISAAQLRHVMTNLGEKLTDEEVDEMIREADIDGDGQVNYEEFVQMMTAKGGKRRW  160
LMC8 81  KMKDTDSEEEIREAFRVFDKDGNGYISAAQLRHVMTNLGVKLTDEEVDEMIREADIDGDGQVNYEEFEQMMTAKGGKRRW  160

LMC4 161 KKNFIAVSAANRFKKISSSGALGSGGG  187
LMC5 161 KKNFIAVSAANRFKKISSSGALGSGGG  187
LMC6 161 KKNFIAVSAANRFKKISSSGALGSGGG  187
LMC7 161 KKNFIAVSAANRFKKISSSGALGSGGG  187
LMC8 161 KKNFIAVSAANRFKKISSSGALGSGGG  187
```

Supplemental Figure S3. Different variants of Lumicampsin (LMC) based on different calmodulin moieties. **a**. All LMC derivatives have the same sbGluc split between residues Q88 and G89 and the same M13 moiety with different calmodulin moieties inserted: GCaMP6f (Chen et al., 2013; LMC4), GCaMP6s (Chen et al., 2013; LMC5), GCaMP6m (Chen et al., 2013; LMC6), Ca^{2+}-NL (Saito et al., 2012; LMC7), Ca^{2+}-eNL (Suzuki et al., 2016; LMC8). Dark blue font indicates amino acids varying between the five versions.

1157
1158
Supplemental Figure S4. Different versions of LMC4 based on different subcellular targeting sequences. **a**.
Schematics of four versions of LMC4 targeting the Ca2+ sensor to the inner cell membrane via a farnesylation or
myristoylation signal, to the ER via a KDEL signal, or to the mitochondrial membrane via a mitochondrial targeting
signal of cytochrome c oxidase. **b**. Mean fold change of light emission from HeLa cells expressing LMC4f or myLMC4
after histamine-induced Ca2+ flux. **c**. Bioluminescence response to ionomycin induced Ca^{2+} flux in HEK293 cells
expressing LMC4f or LMC4er in the presence of CTZ and in the absence or presence of extracellular Ca^{2+}. **d**. Confocal
images of HeLa cells expressing miLMC4-dTomato and BacMam2.0-Mito-GFP.
1166

1167

1168 **Supplementary Figure S5. LMC reports Ca²⁺ flux in cardiomyocytes *in vivo*. a.**Pseudocolored IVIS images
1169 showing bioluminescent signal from a mouse heart transduced with AAV2/9-CAG-LMC4f. Images correspond to
1170 baseline (left) and peak signal (right) following intraperitoneal injection of 100 µg water-soluble luciferin (CTZ). **b.** Peak
1171 signal was reached approximately 15 minutes post-injection and is plotted in the accompanying time course graph
1172 (right).

1173

1174
1175
Supplementary Figure S6. Lack of BLADe-mediated changes in neurons expressing a non-functional "DUD"
opsin. (i) Schematic of farnesylated LMC anchored to the inner cell membrane near the non-functional DUD opsin,
showing that in the presence of depolarization-induced Ca^{2+} flux and luciferin (CTZ), no light-mediated modulation is
expected. (ii) Representative membrane voltage responses to step current injections before (gray) and after (blue) bath
perfusion of luciferin in a neuron co-expressing the DUD opsin and LMC, showing no change in firing. (iii) Action
potential counts under vehicle (gray) and luciferin (blue) conditions remain similar. (iv) Empirical cumulative distribution
of spike fraction changes across voltage traces for vehicle and luciferin conditions, demonstrating no shift in spike
timing (Kolmogorov-Smirnov test, KS= 0.0545, *P*= 0.999) or firing rate in the absence of a functional opsin. Data
expressed as mean ± SEM from N=6 neurons.
1185

1186

1187

Supplementary Figure S7. LMC enables light-dependent suppression of cortical network activity via an
inhibitory opsin. a. Representative multi-unit activity (MUA) traces from cortical neurons recorded on multielectrode
arrays. Neurons co-expressing LMC and the inhibitory opsin hGtACR2 (top) show reduced spiking following luciferin
(CTZ) addition (red arrow). In contrast, neurons expressing the opsin alone (bottom) show no consistent change in
firing after CTZ. **b.** Quantification of firing rate changes before and after luciferin addition across individual electrodes.
n.c., no change in firing; inhibited, decrease in firing; enhanced, increase in firing

1194

1195

Supplementary Figure S8. MEA co-expression control: opsin-only cultures lack LMC. a. Schematic of multi-
electrode array (MEA) setup with an opsin-expressing neuron positioned above the recording site. **b.** Fluorescence
images of a representative opsin-only culture showing absence of dTomato signal (left, red channel) and strong opsin-
EYFP expression (middle, green channel). Merged image (right) confirms that these neurons express the opsin but not
LMC. Insets show magnified views of cell bodies overlaying electrode contacts. These cultures serve as a negative
control for luciferin-dependent changes in network activity.

1202

1203 **Supplementary Table 1. List of plasmids**

1204

Plasmid	Source
pcDNA3.1-CMV-LMC4-P2A-dTomato	This paper
pcDNA3.1-CMV-LMC5-P2A-dTomato	
pcDNA3.1-CMV-LMC6-P2A-dTomato	
pcDNA3.1-CMV-LMC7-P2A-dTomato	
pcDNA3.1-CMV-LMC8-P2A-dTomato	
pcDNA3.1-CMV-LMC4f-P2A-dTomato	
pcDNA3.1-CMV-myLMC4-P2A-dTomato	
pcDNA3.1-CMV-LMC4er-P2A-dTomato	
pcDNA3.1-CMV-miLMC4-P2A-dTomato	
pcDNA3.1-CMV-GCaMP6f	
pcDNA3.1-CMV-BlueCaMBI	
pcDNA3.1-CMV-GLICO	
pcDNA3.1-CMV-ChR2(C128S)-EYFP	
pcDNA3.1-CMV-hGtACR2-EYFP	
pcDNA3.1-CMV- ChR2(C128S)-E97R-D253A	
pcDNA3.1-5xC120-EYFP	
pAAV-hSyn-LMC4f-P2A-dTomato	
pAAV-CAG-VP-EL222	
pAAV-5xC120-EYFP	
pAAV-Efla-DIO-CheRiff-EYFP	
pAAV-Efla-DIO-LMC4f-P2A -dTomato	
pcDNA3.1-CAG-VP-EL222	Kevin Gardner, CUNY
pcDNA3.1-5xC120-FireflyLuc	Kevin Gardner, CUNY
pcDNA3-CMV-GeNL(Ca2+)_520	Addgene plasmid # 85204
pAAV.Syn.GCaMP6f.WPRE.SV40	Addgene plasmid # 10083
pFUGW-hGtACR2-EYFP	Addgene plasmid # 67877
pLenti-CaMKIIa-hChR2(C128S)-EYFP-WPRE	Addgene plasmid # 20294
pAAV-hsyn-CheRiff-eGFP	Addgene plasmid # 51697

1205

1206