A FRET-based excitatory luminopsin with improved efficacy of bioluminescence-driven channelrhodopsin activation

Andreas Björefeldt[1-3], Jeremy Murphy[2], Emmanuel L Crespo[1,4], Mansi Prakash[1],
Nina Friedman[2], Tariq M Brown[1], Diane Lipscombe[5], Christopher I Moore[5], Ute
Hochgeschwender[1,4,6], Nathan C Shaner[7]

1 College of Medicine, Central Michigan University, Mount Pleasant, MI 48859, USA

2 Department of Neuroscience, Brown University, Providence, RI 02912, USA

3 Department of Physiology, Institute of Neuroscience and Physiology, University of
Gothenburg, Box 432, 405 30, Gothenburg, Sweden

4 Biochemistry, Cell and Molecular Biology Graduate Program, Central Michigan
University, Mount Pleasant, MI 48859, USA

5 Carney Institute for Brain Science, Brown University, Providence, RI 02906, USA

6 Program in Neuroscience, Central Michigan University, Mount Pleasant, MI 48859,
USA

7 University of California, San Diego, School of Medicine, Department of
Neuroscience, 9500 Gilman Drive La Jolla, CA 92093-0662, USA

Corresponding authors:

andreas.bjorefeldt@gu.se
hochg1u@cmich.edu

Abstract

The ability to manipulate neuronal activity both opto- and chemogenetically with a single actuator molecule presents unique and flexible means to study neural circuit function. We previously developed methodology to enable such bimodal control using fusion molecules called luminopsins (LMOs), where a channelrhodopsin actuator can be activated using either physical (LED driven) or biological (bioluminescent) light. While activation of LMOs using bioluminescence has previously allowed manipulation of circuits and behavior in mice, further improvement would advance the utility of this technique. Thus, we here aimed to increase the efficiency of bioluminescent activation of channelrhodopsins by development of novel FRET-probes with bright and spectrally matched emission tailored to *Volvox* channelrhodopsin 1 (VChR1). We find that pairing of a molecularly evolved *Oplophorus* luciferase variant with mNeonGreen significantly improves the efficacy of bioluminescent activation when tethered to VChR1 (construct named LMO7) as compared to previous and other newly generated LMO variants. We proceed to extensively benchmark LMO7 against previous LMO standard (LMO3) and find that LMO7 outperforms LMO3 in the ability to drive bioluminescent activation of VChR1 both in vitro and in vivo, and efficiently modulates animal behavior following intraperitonial injection of fluorofurimazine. In conclusion, we demonstrate a rationale for improving bioluminescent activation of optogenetic actuators using a tailored molecular engineering approach and provide a new tool to bimodally manipulate neuronal activity with increased bioluminescence-driven efficacy.

Introduction

Understanding how specific cell types, circuits and brain areas are involved in generating percepts and behaviour requires approaches to experimentally manipulate spatiotemporal patterns of neural activity in genetically-defined populations. At present, optogenetic and chemogenetic techniques constitute the major workhorses used for this purpose, each featuring unique strengths and limitations.

Optogenetics notably offers the advantage of millisecond control of neuronal output (Boyden et al 2005, Fenno et al 2011) but requires invasive delivery of a light source. In comparison, chemogenetic approaches (among which designer receptors exclusively activated by designer drugs, DREADDs, are most widely used) feature low temporal resolution but provide non-invasive modulation of neuronal populations distributed across much larger areas of the brain (Sternson & Roth 2014). To combine the strengths of above techniques and increase user flexibility we previously developed a bimodal strategy (bioluminescent optogenetics, BL-OG) where a luciferase-opsin fusion molecule (a luminopsin, LMO) integrates opto- and chemogenetic features in a single actuator (Moore & Berglund 2020).

With BL-OG, chemogenetic modulation of neuronal activity is achieved via delivery of a small luciferase substrate, coelenterazine (CTZ), that when oxidized produces biological light (bioluminescence, BL) to activate an optogenetic actuator. So far, the efficiency of CTZ-driven (chemogenetic) activation has been adequate but not yet fully optimized. Thus, further improvements in this area would increase the utility of BL-OG and promote wider adoption of the technology in the neuroscience field.

We have previously shown that the coupling efficiency between bioluminescent emitter and optogenetic actuator is readily modifiable, e.g. by pairing different

luciferase and opsin variants (Berglund et al 2013, Berglund et al 2016, Park et al

2020). With bioluminescent activation of an opsin, the magnitude of the total

transmembrane current generated (i.e. the efficiency of activity modulation) depends

on parameters including the energy output and brightness of the emitter, spectral

compatibility (photon emission wavelength with respect to maximum opsin

absorption wavelength), photosensitivity of the recipient opsin chromophore, as well

as LMO expression level and bioavailability of CTZ.

We here explore a strategy for improving bioluminescence-driven opsin activation

based on development of bright and spectrally matched emitters featuring two novel

engineered luciferases optimized for förster resonance energy transfer (FRET) to the

fluorescent protein mNeonGreen. By fusing one such emitter to *Volvox*

Channelrhodopsin 1 (VChR1) we identify a new excitatory LMO (LMO7) with a

significantly improved efficiency of bioluminescence-driven opsin activation.

Results

Generation of FRET-based LMOs

As strategy to generate more efficient LMOs, we focused on new light-emitters

displaying brighter BL emission and/or improved spectral compatibility with the

Volvox channelrhodopsin 1 (VChR1) chromophore. To this end, we engineered

highly efficient FRET pairs, using mNeonGreen as the acceptor, by (i) designing a

novel bright copepod-derived luciferase and (ii) evolving an *Oplophorus gracilirostris*

luciferase (OLuc) (Inouye et al 2000) variant with improved performance compared

to NanoLuc (Hall et al 2012).

To generate an improved copepod-type luciferase, we designed several variants by

introducing subsets of mutations identified by other groups for *Gaussia princeps*

luciferase (Gluc) (Kim et al 2011, Welsh et al 2009) into consensus-derived "artificial"

luciferase (Aluc) sequences (Kim et al 2013). The clone yielding the highest

luminescent signal with CTZ when produced in *E. coli* was chosen for full

characterization and named "sbALuc1" to reflect its major derivation from "sbGluc"

mutations paired with an Aluc. We next optimized a fusion construct of sbALuc1 with

mNeonGreen following a similar approach to that taken to optimize the mVenus-

RLuc8 fusion Yellow NanoLantern (YNL) (Saito et al 2012). Ultimately, this

generated an optimized FRET construct code-named "NCS1" which was used as

light emitter to generate LMO6 (Fig. 1A-C, and Table 1).

We also generated an OLuc variant with favourable properties for fusion constructs

as part of a larger effort to generate bright luminescent probes, described separately

in Lambert et al 2023. Briefly, we performed multiple rounds of directed evolution

consisting of alternating error-prone and site-directed mutagenesis libraries in *E. coli*.

For each round, variants were selected based on higher bioluminescent emission

and solubility. We subsequently introduced a subset of mutations found in NanoLuc

(Hall et al 2012) and NanoBit (Dixon et al 2016) to generate the clone "eKL9h,"

which was fused to mNeonGreen using the same end truncations and linker

sequence as used for the mNeonGreen-NanoLuc pair GeNL (Suzuki et al 2016). A

final module code-named "NCS2" was obtained and used as light emitter to generate

LMO7 (Fig. 1A-C, Table 1). Notably, NCS2 is fully soluble when expressed in *E. coli*

while GeNL, which differs only in the luciferase domain (NanoLuc), is largely

insoluble (Lambert et al 2023). We therefore chose NCS2 over GeNL as the

bioluminescent component of our new LMO constructs.

Since the mNeonGreen FRET acceptor produces bioluminescence emission peaking

around 518 nm for both NCS1 and NCS2, we expected both to activate VChR1 more

efficiently than blue-emitting luciferases (Fig. 1B). To test the ability of NCS1 and NCS2 to drive opsin activation, we generated new LMOs by fusing them to the N-terminus of VChR1 alongside NanoLuc (Nluc) for comparison (Fig. 1C and see Table 1 for information on tested LMOs). We first examined the light output from LMOs in transiently transfected HEK293 cells by IVIS imaging to obtain values for the Average Radiance [p/s/cm²/sr] for each individual well (see Fig. 1D for an example). To baseline light emission to single molecules, RNA was prepared from duplicate plates, reverse transcribed and used for qPCRs to determine the copy number of VChR1 molecules per well. As the light emitters are tethered to the opsin, this allows to determine radiance per copy of LMO. Figure 1E shows the average radiance [p/s/cm²/sr] recorded for each LMO. Interestingly, radiance was lowest from LMO7, leading us to speculate that FRET from the light emitter (here the mNeonGreen-luciferase fusion protein) to the nearby VChR1 chromophore is more efficient for NCS2 than for the other light emitters. To explore this further, we tethered sbGLuc and NCS2 to the extracellular side of the B7 transmembrane sequence from the mouse CD80 antigen and compared their light emission when tethered to an opsin versus a transmembrane sequence. With light emission per copy number set at 100% for the B7 tethered constructs, radiance per copy decreased for both opsin tethered light emitters, with a higher decrease for LMO7 versus LMO3 (Fig. 1F). This is consistent with a loss of photon emission to direct energy transfer being more efficient for LMO7 than for LMO3.

FRET-based LMOs generate superior bioluminescence-driven inward currents in HEK293 cells

To compare the ability of new LMOs to drive VChR1-mediated inward currents we

generated stable expression HEK293 lines for LMO3, 5, 6 and 7, and used whole-cell patch clamp recordings to determine the coupling efficiency of each construct (i.e. the ratio of luciferin-induced versus maximum, LED-induced photocurrent amplitude response). Expressing cells were identified with epifluorescence and voltage-clamped at -60 mV. A brief light stimulation was first applied through the 40x water immersion objective to record the photocurrent response. Depending on the construct, either native or H-CTZ (100 μM) was then introduced via the ACSF perfusion to cells during continuous recording (Fig. 2). In non-expressing (control) HEK 293 cells, none of the two substrates had significant effect on membrane current (Fig. 2B, E and I). In LMO-expressing cells, however, CTZ-driven inward currents of variable magnitude were observed with all four constructs (Fig. 2C, D, F, G). Inward currents developed gradually over 3-4 minutes to reach plateau levels within 5 minutes of substrate entry into the bath (Fig. 2J). We found that LMO6 and LMO7 generated significantly higher CTZ-driven inward current amplitudes on average relative to LMO3 and LMO5 (LMO3: -131.1 ± 32.7 pA, n = 9; LMO5: -87.1 ± 33.3 pA, LMO6: -251.4 ± 44.4 pA, n = 10; LMO7: -659.1 ± 103.9 pA, n = 9; Fig. 2I, J), with average photocurrent amplitudes and densities being similar among groups (Fig. S1). As a result, the coupling efficiency of LMO6 and LMO7 was significantly higher (LMO3: 13.4 ± 3.4%, n = 9; LMO5: 7.2 ± 1.7%, LMO6: 24.8 ± 3.7%, n = 10; LMO7: 63.7 ± 7.2%, n=9; Fig. 2K). LMO7 showed the largest average CTZ-driven current amplitude and coupling efficiency that were highly statistically significant versus LMO3 (CTZ-induced current: p = <0.001, Mann–Whitney U test; coupling efficiency: p = <0.001, Mann–Whitney U test).

The excitation wavelength used to photoactivate VChR1 was matched to the peak emission wavelength of the BL emitter (see Table 1), and thus involved blue light

(480 nm) excitation of LMO3 and 5, and green light (540 nm) excitation of LMO6 and

7. To confirm that this protocol allowed for side-by-side comparison of LMOs we

recorded photocurrent responses to blue and green light from the same cell in a few

LMO3 and LMO7 expressing cells (Fig. S2). At the irradiance level used in the above

set of patch clamp experiments (480 nm: 31 mW/cm^2; 540 nm; 50 mW/cm^2), evoked

photocurrents showed no significant difference in voltage dependence or reversal

potential (Fig. S2A, B) and were of similar amplitude (Fig. S2C, D) in response to

blue and green light stimulation.

LMO7 generates stronger depolarization in neocortical brain slice neurons

We next tested the ability of LMO7 to drive membrane depolarization in neurons,

using LMO3 as reference. Postnatal day 2 (P2) mouse pups were bilaterally injected

with AAV encoding LMO3 or LMO7 under the human synapsin promoter (hSyn) in

fronto-parietal cortex, and coronal slices were prepared at 3-6 weeks. For both

LMOs, expression was observed primarily in layer 5, as verified with 480 nm

epifluorescence prior to experiments and confirmed with post hoc imaging (Fig. S3).

LMO-expressing cells were randomly targeted for whole cell recordings and showed

predominantly pyramidal morphological and electrophysiological properties (Fig. 3A,

D). To verify opsin expression, cells were exposed to a 1 s pulse of 480 nm (31

mW/cm2) and 540 nm (50 mW/cm2) light after break-in to record the maximum

photocurrent amplitude and confirm light-evoked firing responses (Fig. 3A, D, right).

Following a stable baseline recording period at resting membrane potential,

respective CTZ substrate was applied at equal concentration (~300 µM final

concentration in bath) through the ACSF perfusion. Upon CTZ delivery to the slice,

bioluminescent emission typically peaked during the first 60 s after onset, followed

by a reversible depolarizing shift in the membrane potential (Fig. 3B, E) of variable

magnitude (Fig. 3H). We found that LMO7-expressing neurons on average produced

a ~2-fold greater depolarization amplitude relative to LMO3 (10.5 ± 1.8 mV vs 5 ± 1

mV, Fig. 3G, H, p = 0.019, unpaired t-test). In control experiments, application of

vehicle had no effect on the membrane potential over the corresponding time period

(Fig. 3B,E,H), and average photocurrent amplitudes were similar between cell

populations (Fig. S4, LMO3: -1191.5 ± 186 pA, LMO7: -1112,9 ± 166 pA, p = 0.76,

unpaired t-test). In a small number of cells (LMO3: 1/10, LMO7: 3/11) CTZ

application produced suprathreshold depolarization (Fig. 3C, F). To verify that LMO7

increases the firing output of neurons, we also recorded the frequency-current (f-I)

response in a subset of neurons (n = 7) pre, during and post CTZ application (Fig.

3I-L). We found that CTZ application significantly lowered the rheobase current (100

± 7.7 pA vs 64.3 ± 5.1 pA, Fig. 3K, n = 7, p = 0.0004, paired t-test) and increased the

maximum number of action potentials produced (18 ± 1.8 vs 21 ± 2.2, Fig. 3L, n = 7,

p = 0.01, paired t-test), causing a left-shift of the f-I curve (Fig. 3J). Statistically

significant firing responses were observed in the 50–175 pA range (p < 0.05, paired

t-test).

LMO7 elicits larger bioluminescence-driven MUA responses in mouse

somatosensory cortex

The ability of LMO7 to drive neuronal spiking in vivo was next examined in the

primary somatosensory (S1) cortex of adult wild type mice in comparison to LMO3.

Seven weeks post viral injection (AAV9-hSyn-LMO3 or LMO7) all animals showed

clear MUA responses to the LED stimulus (Fig. 4, panel C, percent change from

baseline, LMO7: M = 729.21, SD = 353.48, median = 844.07, range = 322.9–1197.9;

LMO3: M = 1233.2, SD = 1130.9, median = 1047.7, range = 164–3285). Although

the LMO3 group demonstrated greater responses to the LED on average than the

LMO7 group, this difference did not reach statistical significance (t (11) = -1.12, p =

0.28). All animals additionally showed clear bioluminescence after CTZ delivery (Fig.

4, panel B; percent change from baseline, LMO7: M = 273.3, SD = 216.2, median =

237, range = 25.93–572.2; LMO3: M = 739.7, SD = 590.1, median = 741.3, range =

58.74–1517.8). Bioluminescent output was greater on average in the LMO3 group

than the LMO7 group, and this difference trended towards statistical significance

(paired t-test: t (11) = -1.96, p = 0.076). While both the LED MUA response as well

as bioluminescent output were greater in the LMO3 group, the LMO7 group

nevertheless demonstrated a 48.62% larger increase in MUA with CTZ delivery on

average (Fig. 4, panel D; LMO7: M = 51.5, SD LMO7 = 23, median = 54.3, range =

25.7–82.8; LMO3: M = 34.6, SD = 9.7, median = 33.6, range = 22.6–50.2). This

difference did not reach statistical significance (t (11) = 1.66, p = 0.12).

The differences in the LED MUA responses between the two groups concerned us in

that the MUA response to external LED drive should be equal between the groups

given that the optogenetic element in both constructs is VChR1. This difference is

most likely due to subtle differences in construct expression. Indeed, in support of

this across the two groups and within each group, there was a strong linear

relationship between CTZ driven MUA and LED driven MUA (Fig. 4, panel E). A

robust linear fit of the CTZ MUA to the LED driven MUA, collapsing across the two

groups showed a strong, statistically significant relationship (R^2 = 0.596, F (1,11) =

15.3, p = 0.002), which remained consistent when tested within each group (LMO7:

F (1,5) = 16.8, p = 0.009, R^2 = 0.775; LMO3: F(1,3) = 11.4, p = 0.04, R^2 = 0.79).

This suggests that any systematic imbalance, however slight, in the expression of

the two constructs will distort the true differences in the constructs' ability to drive MUA changes. With this in mind we chose a matched subset of animals (LMO3 N = 3, LMO7 N = 4) in which the LED driven responses clustered closely about the median LED responses of the two groups (LMO7 LED subset response: M = 999.5, SD = 146.8, median = 978, range = 844.1–1197.9; LMO3 LED subset response: M = 1215.1, SD = 298.7, median = 1119.1, range = 976.2–1549.9, Fig. 4; orange outlined dots throughout). When these subgroups were compared, the LMO7 animals showed a 64.2% greater CTZ driven MUA response on average (M = 68.6, SD = 11.7, median = 68.6, range = 54.3–82.8) than the LMO3 subgroup (M = 41.8, SD = 7.8, median = 40.2, range = 34.9–50.2), and this difference was statistically significant (t (5) = 3.41, p = 0.019).

LMO7 expression in the basal ganglia drives motor behavior in mice after intraperitoneal injection of fluorofurimazine

To demonstrate the capability of LMO7 for chemogenetic manipulation of behavior, we examined motor behavior in freely moving mice that unilaterally express LMO7 within the substantia nigra pars reticulata (SNr; Fig. 5A). While the native *Oplophorus* luciferase (OLuc) uses coelenterazine as its luciferin, engineered luciferases derived from OLuc emit more light with the coelenterazine analogue furimazine (Hall et al 2012). In addition, furimazine has been chemically modified to provide fluorinated analogs, such as fluorofurimazine (FFz), with improved pharmacokinetics and rapid onset of peak light emission *in vivo* (Gaspar et al 2021, Su et al 2020). FFz has been reported to reach maximal photon emission in mouse brain 5 minutes after i.p. injection (Su et al 2023). FFz administered by i.p. injections induced substantial rotational behavior compared to vehicle injections in LMO7

expressing mice (Fig. 5B-E). This is evident from the trajectories of the snout of the

same mouse after vehicle versus FFz injection (Fig. 5B) and from plotting cumulative

rotational angles of all three mice (Fig. 5C, D) within 5–10 minutes after injection

(160 seconds following the start of video recordings). Cumulative rotational angles

are significantly different between vehicle and FFz treated mice (Fig. 5E, paired t-

test, p = 0.04). As expected, a preference for ipsilateral rotation was detected

compared to contralateral in FFz treated mice, as can be seen in the significantly

different rotational ratios (Fig. 5F, paired t-test, p = 0.01). In addition to preference in

body orientation, increased grooming was observed after unilateral chemo-

optogenetic activation of SNr, presumably through subthalamic nuclei projections

(Guillaumin et al 2021). FFz i.p. injections increased the duration of grooming

significantly (Fig. 5G right panel, paired t-test, p = 0.006). There was a trend towards

an increased number of grooming bouts in FFz compared to vehicle treated mice,

but it did not reach statistical significance (Fig. 5G left panel, paired t-test, p = 0.06).

Discussion

Here we developed and thoroughly benchmarked a new excitatory LMO with

improved efficiency of bioluminescent (chemogenetic) activation in response to small

molecule luciferins (coelenterazines). We identified a novel luciferase variant

obtained through molecular evolution of OLuc whose pairing with mNeonGreen

resulted in bright FRET-based emission peaking at ~518 nm. By tethering this

emitter to VChR1 we generated a new LMO (LMO7) molecule that outperforms our

current standard (LMO3) in its ability to activate this opsin using bioluminescence.

We show that in presence of H-CTZ or FFz, LMO7 generates superior inward

currents in HEK cells, promotes increased neuronal depolarization and spiking in

brain slice and in vivo recordings, and controls motor and grooming behavior in mice

following viral injection.

We tested the ability of four different bioluminescent emitters (sbGluc, NanoLuc,

NCS1 and NCS2) to drive bioluminescence-dependent activation of VChR1 in HEK

cells. As compared to sbGluc and NanoLuc (both single luciferase emitters) we

found that the two FRET probes (NCS1 and NCS2) had higher coupling efficiency

and were able to generate larger inward currents, suggesting that, in addition to

maximum brightness, close spectral pairing of emitter and opsin may be important at

the comparably lower light intensity levels obtained with bioluminescence versus

LED light sources. Whereas both FRET-probes featured emission peaks at ~518

nm, the coupling efficiency of NCS2 to VChR1 proved to be significantly higher than

for NCS1, suggesting that the orientation of NCS2 relative to the channel may also

be favorable. As we observed substantially lower bioluminescent emission with

LMO7 versus other LMOs, it is possible that NCS2 may stimulate opsin activation

more efficiently through higher percentage of (non-radiant) FRET to the VChR1

chromophore.

Compared to previous versions of LMOs, LMO7 will allow more flexibility in

experimental applications. Due to its higher coupling efficiency, it can be used with

lower concentrations of luciferin, allowing intraperitoneal application rather than

intraventricular of intravenous. Moreover, as the luciferase in LMO7 utilizes

furimazine analogues, investigators can take advantage of the ongoing development

of variants with improved brain access (Gaspar et al 2021, Su et al 2023, Su et al

2020).

It should be readily possible to further enhance the efficacy of bioluminescent

activation of opsins in future studies. As bioluminescent emitters and highly light

sensitive opsins continue to evolve (Marshel et al 2019, Vierock et al 2022), and as

new CTZ analogues with increased brain permeability, half-life and brightness are

developed (Su et al 2023, Su et al 2020), the ability to manipulate neuronal activity

with biological light will continue to improve.

In summary, we present a new tool allowing combined opto-chemical manipulation of

neuronal excitation with improved bioluminescence-driven efficacy, which should

further increase the utility of the BL-OG technique.

Materials and methods

Animals

All experiments were conducted in accordance with protocols approved by the

Institutional Animal Care and Use Committee (IACUC) at respective universities

following National Institutes of Health (NIH) guidelines. C57/BL6 (JAX stock

#000664) and Swiss Webster (Charles River) mice of both sexes were used and

housed 4–5 animals per cage at a temperature and humidity controlled vivarium on a

12 hour light/dark cycle with ad libitum access to water and food.

Luciferase constructs and characterization of recombinant proteins

Bacterial expression plasmids were constructed using the pNCS backbone and

recombinant protein expression was performed in *E. coli* strain NEB 10-beta for all

experiments as previously described (Lambert et al 2020, Shaner et al 2013).

Directed evolution including error-prone and site-directed mutagenesis and

screening for luminescence intensity and wavelength was performed according to methodology described in Lambert et al 2023. Bioluminescence emission spectra were collected using a ClarioStar multimode plate reader (BMG).

Plasmids

The generation of LMO plasmids has been described in detail previously (Berglund et al 2013, Berglund et al 2016). For the new versions of LMOs, the sbGLuc sequence of LMO3 (Berglund et al 2016) was switched out with synthetized mammalian codon-optimized sequences (IDT) of NanoLuc (Hall et al 2012), NCS1 and NCS2 (Lambert et al 2023). For the B7 constructs pcDNA3.1-CMV plasmids were generated with sbGLuc and NCS2, respectively, tethered to the N-terminus of B7 (Chou et al 1999) via the same 15 amino acid linker used for tethering light emitters to opsins in LMOs, and EYFP fused to the C-terminus of B7. For viral vectors, the coding sequence of LMO7 was cloned into an AAV vector downstream of an hSyn promoter. For AAV LMO3 the published plasmid was used (Berglund et al 2016; Addgene #114099).

Virus

AAVs carrying LMOs were produced by transfecting subconfluent HEK293FT cells per 10-cm culture dish with 24 µg of the helper plasmid pAd delta F6, 20 µg of the serotype plasmid AAV2/9, and 12 µg of the LMO plasmid using Lipofectamine 2000. After 72 h, the supernatant was harvested from culture plates and filtered at 0.45 µm. Virus was purified from cells and supernatant according to previously described methodology (Guo et al 2012), but without the partitioning step in the aqueous two-phase system. Virus was dialyzed against PBS (w/o Ca, Mg) overnight at 4 °C, using FLOAT-A-LYZER G2, MWCO 50 kDa, followed by concentration in Amicon Ultra-0.5 mL Centrifugal Filters. Viral titers were determined by quantitative PCR for the

381   woodchuck hepatitis post-transcriptional regulatory element. Preparations with titers

382   around 1×10^{13} vg/mL were used in the study.

Cell culture

HEK293 cells were grown in Dulbecco's Modified Eagle Medium (DMEM,

ThermoFisher) containing 10% fetal calf serum and 0.5% penicillin/streptomycin.

Cells were cultured in 6-well plates at 37 °C in a 5% CO_2 atmosphere and split every

72 hours by trypsinization (0.25%, Gibco, ThermoFisher). Stable HEK293 lines were

generated by lipofection (Lipofectamine 2000, ThermoFisher) of ~80% confluent

cells with 2 µg of linearized expression plasmid overnight according to the

manufacturer's instructions. Cells expressing EYFP and mNeonGreen (LMO3 and

LMO5 versus LMO6 and LMO7, respectively) were purified with puromycin titration

(1-10 µg/mL, SigmaAldrich) and frozen at -80°C. The percentage of cells showing

stable expression ranged from 30–60% cells in mixed populations.

RNA isolation, cDNA synthesis and RT-qPCR

RNA was extracted from individual wells of 24-well culture plates using a RNeasy

Plus Mini Kit (Qiagen) per the manufacturer's instructions. Culture medium was

removed from wells and cells were lysed on the plate in situ. The lysate was

homogenized using a QIAshredder spin column, and the homogenized lysate was

further purified with a gDNA Eliminator spin column and an on-column DNase

digestion. Concentration was determined with a Nanodrop (A260/A280=1.8-2.2;

ThermoFisher) and integrity of RNA was assessed in a Bioanalyzer (Agilent). RNA

(100ng/sample) was reverse transcribed to cDNA using the High-Capacity RNA to

cDNA kit (Thermo Fisher Scientific) per the manufacturer's instructions. The

generated cDNA (5 ng) was used as a template for RT-qPCR, which was performed

with the Applied Biosystems Step One Plus using Power SYBR Green PCR Master

Mix (Applied Biosystems, Thermo Fisher 4367659). The forward and reverse primers

for VChR1 in LMO constructs were 5'- CGGATGGGAGGAGGTTTACG-3' and 5'-

AACTACGCCGTTCCCTGAAG-3' and for EYFP in B7 constructs were 5'-

TAAACGGCCACAAGTTCAGCGTGT-3' and 5'-

ATGTTGCCGTCCTCCTTGAAGTCGA-3'.

IVIS imaging

Luminescence was measured using an IVIS Lumina LT in vivo bioluminescence and

fluorescence imaging system (Perkin Elmer). The 4-well plate was moved from the

incubator to the imaging room. Medium was removed from each of the 4 wells and

replaced with the CTZ-containing medium. Plates were immediately imaged (1

second, no filters, medium binning). Images were displayed as a pseudo-color

photon count image. Regions of interest (ROIs) were defined using an automatic

intensity contour procedure to identify bioluminescent signals with intensities

significantly greater than background. ROIs were measured for each image to obtain

values for the Average Radiance [p/s/cm²/sr] for each individual well.

HEK cell electrophysiology

Stable HEK293 cell lines were seeded at low density on 15 mm poly-D-lysine coated

coverslips (Neuvitro) in 12-well plates and incubated for 48-72 hours pending

electrophysiological analysis. A coverslip was transferred to a recording chamber

(RC26-GLP, Warner Instruments) on the stage of an upright microscope (BX51WI,

Olympus) and perfused with ACSF (1–1.5 mL/min) containing (in mM): 150 NaCl, 3

KCl, 10 HEPES, 2 $CaCl_2$, 2 $MgCl_2$ and 20 D-glucose (pH 7.4, ~300 mOsm/kg) at a

temperature of 35 ± 1°C. The intracellular solution contained (in mM): 130 K-

gluconate, 8 KCl, 15 HEPES, 5 Na_2-phosphocreatine, 4 Mg-ATP, 0.3 Na-GTP (pH

7.25, ~300 mOsm/Kg). Borosilicate glass micropipettes were manufactured using a

vertical puller (PC-100, Narishige) and had resistances from 4–6 MΩ. Cells expressing EYFP or mNeonGreen were visually targeted in clusters of 5–15 cells using epifluorescence. VChR1 was excited through the objective (LUMPLFLN40XW, NA 0.8, Olympus) using a metal halide light source (130 W, U-HGLGPS, Olympus) together with filter cubes for blue (Ex/Em: 480/530 nm, U-MNIBA3, Olympus) and green (Ex/Em: 540/600, U-MWIGA3, Olympus) excitation. An electronic shutter (Lambda SC, Sutter Instruments) was used to control light delivery to cells. Photocurrents were evoked by 1 s exposure to blue or green light, respectively, according to peak bioluminescent emission of the evaluated construct (see Table 1). At maximum light intensity the irradiance was 30.8 and 50 mW/cm^2, respectively (photocurrent amplitudes saturated at lower irradiance for both wavelengths, Fig. S2).

Whole-cell voltage clamp recordings were performed at -60 mV using a Multiclamp 700b amplifier and Digidata 1440 digitizer together with the pCLAMP 10 recording software (Molecular Devices). Data was sampled at 10 kHz, filtered at 3 kHz and analyzed in Igor Pro (WaveMetrics). After break-in, the photocurrent response of the cell was determined using a gap free protocol followed by application of CTZ (Nanolight Technology, #301 and 303) through the perfusion. A fresh stock of CTZ (50 mM in DMSO) was aliquoted at 2 mM and diluted to 250 µM in 1 mL ACSF. Fresh CTZ aliquots were quickly thawed and immediately introduced into the perfusion (final concentration in bath ~100 µM) for each recording. VChR1 photocurrent-voltage plots were acquired in voltage steps of 10 mV (-80 to +80 mV) during 1 s windows of blue and green light illumination separated by 10 s windows of no light stimulation. The series resistance (Rs) was monitored throughout and recordings discarded in cases were fluctuations exceeded 25% from start to end.

456 The liquid junction potential was not corrected for.

Brain slice electrophysiology

Acute brain slices were prepared from 3–6 week-old Swiss Webster mice that
received intracortical injection of AAV9-hSyn-LMO virus at postnatal day 2. Mice
were deeply anaesthetized via inhalation of isoflurane (Abbott), decapitated, and
dissected. The cerebrum was quickly isolated and placed in an ice-cold slicing ACSF
containing (in mM): 92 NMDG, 2.5 KCl, 0.5 $CaCl_2$, 10 $MgSO_4$, 30 $NaHCO_3$, 1.25
NaH_2PO_4, 20 HEPES, 2 Thiourea, 5 Na-ascorbate, 3 Na-pyruvate and 25 D-glucose
(310 mOsm/kg, pH 7.3–7.4) (Ting et al 2018). Coronal sections (300 μm) were
prepared from the fronto-parietal cortex of both hemispheres using a vibratome
(VT1000s, Leica) and transferred to a chamber containing slicing ACSF at 34 ± 1 °C.
Following a NaCl spike-in protocol (Ting et al 2018), slices were transferred to a
storage solution containing (in mM): 92 NaCl, 2.5 KCl, 2 $CaCl_2$, 2 $MgSO_4$, 30
$NaHCO_3$, 1.25 NaH_2PO_4, 20 HEPES, 2 Thiourea, 5 Na-ascorbate, 3 Na-pyruvate
and 25 D-glucose (310 mOsm/kg, pH 7.3–7.4) and maintained at RT for up to 6 hrs.
After minimum 30 minutes a slice was transferred to a recording chamber mounted
on an upright microscope (BX51WI, Olympus) and perfused with ACSF containing
(in mM): 121 NaCl, 2.8 KCl, 1 NaH_2PO_4, 26 $NaHCO_3$, 2 $CaCl_2$, 2 $MgCl_2$ and 15 D-
glucose (310 mOsm/kg, pH 7.3–7.4) at a rate of ~6 mL/min. All solutions were
bubbled with a gas mixture of 95% O_2 and 5% CO_2. The recording ACSF was
supplemented with D-AP5 (50 μM, Tocris), CNQX (15 μM, Tocris) and picrotoxin
(100 μM, SigmaAldrich) to block fast glutamatergic and GABAergic synaptic
transmission. Whole-cell patch clamp recordings were performed using a Multiclamp
700b amplifier and Digidata 1440 digitizer together with the pCLAMP 10 recording
software (Molecular Devices). Borosilicate glass micropipettes were obtained using a

PC-100 puller (Narishige) and had resistances of 3–5 MΩ. Pipettes were filled with

intracellular solution containing (in mM): 130 K-gluconate, 10 KCl, 15 HEPES, 5 Na_2-

phosphocreatine, 4 Mg-ATP, 0.3 Na-GTP and 0.03 Alexa Fluor 594 (ThermoFisher)

(310 mOsm/kg, pH 7.3). L5 pyramidal neurons were visually targeted using

epifluorescence microscopy together with a CMOS camera (ORCAFusion,

Hamamatsu). A current/membrane potential response to blue light was confirmed for

each cell prior to recording. Excitation light (480 nm, 1 s, 30.8 mW/cm^2 or 540 nm, 1

s, 50 mW/cm^2) was delivered through a 40x water immersion objective using a 130

W metal halide light source (U-HGLGPS, Olympus) together with U-MNIBA3 and U-

MWIGA3 filter cubes (Olympus). An electronic shutter (Lambda SC, Sutter

Instruments) was used to control the time window of illumination. CTZ was prepared

and administered as in HEK cell recordings except at a higher final bath

concentration of 300 µM. The effect of CTZ on depolarization and spontaneous firing

was examined in continuous recordings from a starting Vm of -70 mV. Vehicle

consisted of an equal amount DMSO diluted in ACSF. In LMO7-expressing cells, the

effect of CTZ on neuronal input-output (f-I) was quantified using depolarizing square

current injections (800 ms, 0–300 pA., 25 pA increments) from -70 mV.

Bioluminescence was simultaneously recorded (0.2 Hz, 4x4 binning) using an ORCA

Fusion CMOS camera (Hamamatsu) in some cells (LMO3; n=2, LMO7; n=3). Data

was sampled at 10 kHz, filtered at 3 kHz and analyzed in Igor Pro (WaveMetrics). Rs

values were ≤15 MΩ in all cells after break-in and were not allowed to fluctuate more

than 25% from start to end of recording. The liquid junction potential was not

corrected for. Brain slices containing neurons filled with Alexa Fluor were fixed post

hoc in 4% paraformaldehyde overnight at 4°C (in PBS, pH 7.3–7.4). Prior to imaging,

slices were washed in PBS and tissue cleared using the ScaleSQ(5) protocol (Hama

et al 2015). Finally, slices were mounted on slides with ProLong Gold (ThermoFisher), coverslipped and imaged on an Axio Imager M2 microscope (Zeiss) using 10x, 20x and 40x air objectives.

In vivo electrophysiology and bioluminescence imaging

Electrophysiological data was acquired using an Open Ephys acquisition board (http://www.open-ephys.org/) connected via an SPI interface cable (Intan) to a 32-channel headstage (Intan). A 32-channel laminar probe (Neuronexus, A1x32-Poly2-5mm-50s-177) was connected to the headstage. The iridium electrode contacts on the probe covered a linear length of 790 μm and were arranged into two columns of 16 contacts spaced 50 μm apart. The data were acquired using the Open Ephys GUI software at a sampling rate of 30 kHz and referenced to a supra-dural silver wire inserted over the right frontal cortex. Imaging data were acquired using an Andor Ixon Ultra 888 EMCCD camera fitted with a Navitar Zoom 6000 lens system. The data were acquired using Andor Solis data acquisition software (Andor Solis 64 bit, v4.32). The field-of-view was centered over the craniotomy and adjusted to encompass the full 3 mm diameter of the craniotomy. Images (512 x 512 pixels, ~6 μm2/pixel) were acquired continuously at an exposure length of 2 s and an electron multiplication gain of 300. To test the optogenetic elements of the constructs, external 530 nm light was delivered by an LED (Mightex FCS-0530-0000) coupled to an optic fiber (105 μm diameter, 0.22 NA). The patch fiber was coupled to a fiber on the neuronexus probe (105 μm diameter, 0.22 NA), which terminated 200 μm above the top contact of the probe.

Experiments were conducted about seven weeks on average after surgery (LMO7: M = 6.98, SD = 1.60, median = 6.71, range = 5.00–9.71; LMO3: M = 7.18, SD = 1.94, median = 7.14, range = 4.85–10.14). On the day of the experiment, the animal

was anesthetized (1–2% isoflurane) and a dental drill was used to remove the

cement holding the window in place. The window was then lifted off to expose the

open craniotomy. For the duration of the experiment the craniotomy was kept

covered in saline. The animal remained anesthetized for the entirety of the

experiment and was transcardially perfused at its conclusion.

We identified the regions of strongest viral transfection within the craniotomy under

epifluorescent 530 nm illumination. For each animal, the center of this region was

selected for the location to insert the electrode, avoiding visible blood vessels. The

electrode was lowered into the cortex through the intact dura using a

micromanipulator (Siskiyou MD7700) at a rate of 1 µm/sec. The electrode was

lowered until the uppermost contact on the laminar probe disappeared from view into

the brain as viewed under a stereoscope. The probe was then left undisturbed for 30

minutes before beginning recording. Each animal was next tested for the presence of

an LED evoked optogenetic response. Fifty trials of two gaussian pulses (peak

intensity at fiber tip = 10.7 mW/mm^2) 200 ms in length with a 200 ms inter-pulse-

interval were delivered at an inter-trial-interval of 7 to 10 s (pseudorandom square

distribution). Multi-unit spiking activity (MUA) was monitored during stimulation. All

animals exhibited an increase in MUA with LED stimulation that was readily

observable in real-time on a trial to trial basis.

Either water soluble native coelenterazine (LMO3 animals, Nanolight CAT#3031) or

coelenterazine H (LMO7 animals, Nanolight CAT#3011) was dissolved in sterile

saline (4.55 µg/µL) to yield a final concentration of 11 mM. The solution was loaded

into a 100 µl glass syringe (Hamilton #80601) fitted with a ~1 cm length of 18 gauge

plastic tubing. The Hamilton syringe and tubing was placed in a motorized injector

(Stoelting Quintessential Stereotaxic Injector, QSI). The tip of the plastic tubing was

lowered into the pool of saline over the craniotomy using a micromanipulator until it

touched the surface of the skull. The tip of the tubing was further adjusted so that it

rested at a distance of ~7.5 mm from the opening edge of the craniotomy. The CTZ

was delivered by infusing 20 µL of the solution into the saline over the open

craniotomy at a rate of 200 µL/min. The volume of the saline well within the headpost

and surrounding the craniotomy was measured to be 200 µL in volume. The addition

of the 20 µL solution to the saline was chosen to yield a 1 mM final concentration in

the saline well.

A TTL pulse co-triggered the acquisition of electrophysiology and imaging data and a

minimum of 5 minutes of baseline activity was recorded for each animal. The CTZ

was then infused into the craniotomy and data were recorded for a minimum of 20

minutes.

Behavioral testing

Three female C57BL/6 mice were used. Several weeks after viral transduction the

behavior of mice in an open field chamber was assessed after intraperitoneal (i.p.)

injection of vehicle (saline) and fluorofurimazine (FFz; 175 µL, 1.53 µmoles) (Gaspar

et al 2021, Su et al 2020) in two recording sessions on separate days. Immediately

after the injection, mice were placed in the open field chamber (clear plexiglass,

10"x12"x15"). RaspberryPi 4 cameras were utilized to capture videos (1280 x 730 at

25 fps) from below with recordings starting 5–8 minutes after the i.p. injection. Motor

behaviors were tracked for 10 minutes. Videos were analyzed for overall turning

preferences (ipsi- vs contralateral of the transduced SNr) using DeepLabCut.

Number of rotations, grooming bout number and total time spent grooming were

manually scored by an experimenter blind to the treatment (vehicle, FFz) of each

mouse.

Surgery and viral injections

Brain slice electrophysiology

P2 Swiss Webster pups were cryo-anesthetized on ice for 5 minutes prior to injection. Upon cessation of movement pups were bilaterally injected with 1 uL AAV9-hsyn-LMO3-EYFP (3 females/3 males) or AAV9-hsyn-LMO7 (4 females/3 males) into the fronto-parietal cortex using a borosilicate glass pipette. Injections were performed manually by applying a small positive pressure to the pipette. Prior to, and following injections, pups were kept warm on a heating blanket and placed back with their dam after the procedure.

In vivo electrophysiology

A total of 15 C57BL/6J mice (7 female/8 male) were used. Seven of these animals (3 females) were injected with the LMO3 viral construct (AAV9-hsyn-LMO3-EYFP; age on day of surgery: M = 26.92 weeks, SD = 4.18, median = 30.14, range = 26.00–40.43). The remaining eight animals (4 females) were injected with the LMO7 viral construct (AAV9-hSyn-LMO7; age on day of surgery: M = 26.23 weeks, SD = 4.79, median = 30 , range = 26.28–40.28). Each animal was anesthetized (1–2% isoflurane), fitted with a steel headpost, and injected with viral constructs in a 3 mm craniotomy centered over left SI (-1.25 A/P, 3.5 M/L relative to bregma) made with a dental drill. Three injections of the constructs were made at locations equidistantly spaced around the central SI point at depths of 500 μm. Each injection was 400 nL in volume and delivered at a rate of 50 nL/min. Viral injections were performed through a glass pipette fitted in a motorized injector (Stoelting Quintessential Stereotaxic Injector, QSI). Finally a glass window was fitted over the open craniotomy and cemented in place with dental cement (C & B Metabond). Dexamethasone was given intraperitoneally (0.5 mg/kg) to reduce brain edema, and

sustained-release meloxicam (Meloxicam SR) was given subcutaneously (4 mg/kg)

for pain relief during recovery.

Behavior

Ten-week-old mice were stereotaxically injected into the substantia nigra pars

reticulata (SNr) with the LMO7 viral construct (AAV9-hSyn-LMO7) through a

unilateral craniotomy (from Bregma: A/P -3.4; M/L 1.25; D/V -4.2). Solution

containing virus at a concentration of 1×10^{13} vg/mL was injected at a volume of

500–800 nL and at a rate of 100 nL/min using a NanoFil syringe with a 35-gauge

beveled needle.

Data analysis

HEK cell electrophysiology

Photocurrent amplitudes were analyzed at steady state level as the average current

amplitude measured during the last 100 ms of illumination to circumvent

inconsistencies associated with light-adaptation and the peak current response. The

CTZ-induced current amplitude was quantified at the peak plateau level observed

within 5 minutes as a ± 10 s average around the peak amplitude. Only recordings

where initial holding current was below -100 pA and stable for minimum 2 minutes

following light stimulation were used in analysis. The CTZ-induced response was

normalized to photocurrent amplitude in each recorded cell to yield the coupling

efficiency (CTZ-induced current amplitude / maximum photocurrent amplitude). Only

cells displaying photocurrent amplitudes of minimum 500 pA were used for

evaluation of coupling efficiency. In non-expressing controls the effect of CTZ on

holding current was quantified 5 minutes post wash in and averaged over 20 s. All

data was analyzed in Clampfit (Molecular devices) and Igor Pro (WaveMetrics).

Brain slice electrophysiology

Cells displaying holding currents less than -150 pA and Rs ≤ 15 MΩ were included in

the analysis. CTZ-induced depolarization amplitudes were calculated as a ± 5 s

average around a manually identified peak amplitude observed at 50-80 s following

substrate delivery to bath. The effect of vehicle on Vm was calculated as a 10

second average 70-80 ms post substrate addition (no effect on Vm was observed

after an additional 60 s). In frequency-current (f-I) plots the number of action

potentials produced during 800 ms depolarizing square current injection was

calculated at baseline (pre), after 90 s post CTZ delivery to bath (CTZ) and following

5 minutes of wash out with ACSF (post). The rheobase current was defined as the

minimum stimulation magnitude required to evoke ≥ 1 action potentials. Action

potential threshold was defined as a dV/dt of ≥ 20 mV/ms and measured from the

first action potential generated by the depolarizing current protocol. Bioluminescence

traces were generated by plotting time-lapse data as mean intensity values from

somatic ROIs of LMO-expressing L5 pyramidal neurons in ImageJ, and time-

synchronized to membrane potential recordings. In simultaneous bioluminescence

and Vm recordings the peak intensity was normalized to the peak depolarization

amplitude. All data was analyzed in Igor Pro (WaveMetrics).

In vivo electrophysiology and imaging

Data were not used from one LMO3 mouse due to a bad reference connection

resulting in excessive noise in the data and no discernible spiking response to LED

stimulation, and one LMO7 mouse due to a hardware malfunction during the infusion

of CTZ. Offline analyses and statistical tests of both electrophysiological and imaging

data were performed in Matlab R2022a (Mathworks). For each recording, electrodes

with RMS values more than three times the interquartile range above the 3rd quartile

or less than three times the interquartile range below the 1st quartile of all 32

electrodes were marked as excessively noisy and removed from further analyses. (LMO7 number of bad electrodes: Mean = 2.28 , SD = 0.75 , median = 2 , range = 1–3; LMO3 number of bad electrodes: Mean = 2.67, SD = 1.21 , median = 2.5, range = 1–4). The remaining electrodes were re-referenced to the common average reference (Ludwig et al 2009). To isolate MUA, a bandpass filter (passband: 500 Hz to 5000 Hz, 3rd order Butterworth) was applied to the data in the forwards and backwards directions to avoid phase distortions. Spikes were defined as data points less than -3 times the standard deviation, where the standard deviation was estimated as the median divided by 0.6745 (Quiroga et al 2004). Each electrode's spike time series was then convolved with a 50 ms gaussian window (SD = 5 ms) to provide an estimate of the instantaneous firing rate.

In the case of LED stimulation, the MUA data were epoched into trials -2 to 2 s relative to stimulus onset. Next we identified responsive electrodes, defined as electrodes in which the lower bootstrapped 99% CI of the poststimulus response (mean MUA activity 0 to 600 ms after LED onset) was greater than the upper bootstrapped 99% CI of the prestimulus 2 s baseline, pooling all the trial data (number of responsive electrodes LMO7: M = 30.85, SD = 0.89, median = 31, range = 29–32; LMO3: M = 27.67, SD = 4.93, median = 29, range = 18–31). The data were then averaged across trials and responsive electrodes to produce a single LED evoked waveform for each animal and converted to percent change relative to baseline. LED response values for each animal are reported as the average percent change from baseline in the 0 to 600 ms poststimulus time period. Spike time series were binned at 1 s intervals in the case of CTZ administration. CTZ responsive electrodes were identified as those whose lower 99% bootstrapped CI of the MUA after CTZ delivery was greater than the upper 99% bootstrapped CI of the baseline

period baseline (Number CTZ responsive electrodes LMO7: M = 11.28, SD = 4.85, median = 10, range = 4–16; LMO3: M = 15.17, SD = 8.35, median = 14.5, range = 4–24). CTZ responsive electrodes were then averaged for each animal and converted to percent change relative to baseline. Among all animals, all electrodes identified as CTZ responsive were also identified as LED responsive. CTZ response values are reported as the maximum increase from baseline in a 60 s moving average across the entire post CTZ time period.

For the imaging data, each image was binned offline by taking the median value in an 8x8 pixel bin, resulting in a new image size of 64 x 64. For each animal, a circular region with a diameter of 8 pixels (~376 µm) was placed in the region directly adjacent to the electrode shank and in front of the surface with the exposed electrode contacts. The mean of these pixels was computed for each image to yield a time series of bioluminescent light output. Each animal's bioluminescence time series was then converted to percent change relative to baseline. Bioluminescence response values for each animal were calculated as maximum increase from baseline in a 60 s moving average across the entire post CTZ time period.

Unless otherwise stated all significance testing was performed using independent samples t-tests (Matlab ttest2).

Behavior

DeepLabCut 2.1 was run on a Lenovo Thinkstation P620 with a NVIDIA Quattro RTX 5000 graphics card. DeepLabCut was utilized for markerless quantification of body orientation by tracking the nose, four paws, and the base of the tail (Mathis et al 2018, Nath et al 2019). Three hundred frames were selected for labeling and used to train the network with a ResNet101 based neural network, utilizing 95% of the labeled frames with default parameters for a single training session. The network

was trained for 200,000 iterations until the loss reached a plateau. By validating with

three shuffles, the test error with a p-cutoff value of 0.9 was 4.39 pixels. We used a

p-cutoff of 0.9 to condition the X,Y coordinates for further analysis. Segments of

recordings with a likelihood of <0.9 were excluded from analysis. Custom scripts

were utilized to quantify cumulative instantaneous body angle after vehicle and FFz

i.p. injections. The body angle was calculated based on the tail base to nose vector

relative to the x-axis.

Statistics

Statistical significance was evaluated by paired or unpaired t-tests, or by a Mann–

Whitney U test when samples failed a test for normality, using SPSS (IBM) or Matlab

(Mathworks). Data is reported as mean ± SEM unless otherwise specified.

Acknowledgments

719 We would like to thank all members of Bioluminescence Hub

720 (http://www.bioluminescencehub.org/) laboratories for their feedback, discussions,

721 and thoughtful comments throughout the progression of this work.

Competing interests

724 The authors have no conflicts of interest to declare.

Author contributions

727 Conceptualization, AB, CIM, UH, NCS; Validation, AB, JM, ELC, CIM, UH, NCS;

728 Formal Analysis, AB, JM, ELC; Investigation, AB, JM, ELC, MP, NF, TB; Data

729 Curation, AB, JM, ELC; Writing—Original Draft, AB, JM, ELC; Writing—Review &

730 Editing, AB, NCS, CIM, UH; Visualization, AB, JM, ELC; Supervision, CIM, UH, NCS;

731 Project Administration, UH; Funding Acquisition, AB, CIM, UH, NCS.

Funding

734 This work was supported by the Swedish Research Council (AB; 2016-06760),

735 National Institutes of Health grants R21MH101525 and U01NS099709, and National

736 Science Foundation grants CBET-1464686 and DBI-1707352.

References

Berglund K, Birkner E, Augustine GJ, Hochgeschwender U. 2013. Light-emitting
channelrhodopsins for combined optogenetic and chemical-genetic control of
neurons. *PLoS One* 8: e59759
Berglund K, Clissold K, Li HE, Wen L, Park SY, et al. 2016. Luminopsins integrate opto-
and chemogenetics by using physical and biological light sources for opsin activation.
Proc. Natl. Acad. Sci. U. S. A. 113: E358-67
Boyden ES, Zhang F, Bamberg E, Nagel G, Deisseroth K. 2005. Millisecond-timescale,
genetically targeted optical control of neural activity. *Nat Neurosci* 8: 1263-8
Chou WC, Liao KW, Lo YC, Jiang SY, Yeh MY, Roffler SR. 1999. Expression of chimeric
monomer and dimer proteins on the plasma membrane of mammalian cells.
Biotechnol Bioeng 65: 160-9
Dixon AS, Schwinn MK, Hall MP, Zimmerman K, Otto P, et al. 2016. NanoLuc
Complementation Reporter Optimized for Accurate Measurement of Protein
Interactions in Cells. *ACS Chem Biol* 11: 400-8
Fenno L, Yizhar O, Deisseroth K. 2011. The development and application of optogenetics.
Annu Rev Neurosci 34: 389-412
Gaspar N, Walker JR, Zambito G, Marella-Panth K, Lowik C, et al. 2021. Evaluation of
NanoLuc substrates for bioluminescence imaging of transferred cells in mice. *J
Photochem Photobiol B* 216: 112128
Guillaumin A, Serra GP, Georges F, Wallen-Mackenzie A. 2021. Experimental investigation
into the role of the subthalamic nucleus (STN) in motor control using optogenetics in
mice. *Brain Res* 1755: 147226
Guo P, El-Gohary Y, Prasadan K, Shiota C, Xiao X, et al. 2012. Rapid and simplified
purification of recombinant adeno-associated virus. *J Virol Methods* 183: 139-46
Hall MP, Unch J, Binkowski BF, Valley MP, Butler BL, et al. 2012. Engineered luciferase
reporter from a deep sea shrimp utilizing a novel imidazopyrazinone substrate. *ACS
Chem Biol* 7: 1848-57
Hama H, Hioki H, Namiki K, Hoshida T, Kurokawa H, et al. 2015. ScaleS: an optical
clearing palette for biological imaging. *Nat Neurosci* 18: 1518-29
Inouye S, Watanabe K, Nakamura H, Shimomura O. 2000. Secretional luciferase of the
luminous shrimp Oplophorus gracilirostris: cDNA cloning of a novel

imidazopyrazinone luciferase(1). *FEBS Lett* 481: 19-25

Kim SB, Suzuki H, Sato M, Tao H. 2011. Superluminescent variants of marine luciferases for
bioassays. *Anal Chem* 83: 8732-40

Kim SB, Torimura M, Tao H. 2013. Creation of artificial luciferases for bioassays. *Bioconjug*
Chem 24: 2067-75

Lambert GG, Crespo EL, Boassa D, Luong S, Sprecher B, et al. 2023. CaBLAM: a high-
contrast bioluminescent calcium indicator derived from an engineered Oplophorus
gracilirostris luciferase. . *bioRxiv*

Lambert GG, Depernet H, Gotthard G, Schultz DT, Navizet I, et al. 2020. Aequorea's secrets
revealed: New fluorescent proteins with unique properties for bioimaging and
biosensing. *PLoS Biol* 18: e3000936

Ludwig KA, Miriani RM, Langhals NB, Joseph MD, Anderson DJ, Kipke DR. 2009. Using a
common average reference to improve cortical neuron recordings from
microelectrode arrays. *J Neurophysiol* 101: 1679-89

Marshel JH, Kim YS, Machado TA, Quirin S, Benson B, et al. 2019. Cortical layer-specific
critical dynamics triggering perception. *Science* 365

Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, et al. 2018. DeepLabCut: markerless
pose estimation of user-defined body parts with deep learning. *Nat Neurosci* 21:
1281-89

Moore CI, Berglund K. 2020. BL-OG: BioLuminescent-OptoGenetics. *J Neurosci Res* 98:
469-70

Nath T, Mathis A, Chen AC, Patel A, Bethge M, Mathis MW. 2019. Using DeepLabCut for
3D markerless pose estimation across species and behaviors. *Nat Protoc* 14: 2152-76

Park SY, Song SH, Palmateer B, Pal A, Petersen ED, et al. 2020. Novel luciferase-opsin
combinations for improved luminopsins. *J Neurosci Res* 98: 410-21

Quiroga RQ, Nadasdy Z, Ben-Shaul Y. 2004. Unsupervised spike detection and sorting with
wavelets and superparamagnetic clustering. *Neural Comput* 16: 1661-87

Saito K, Chang YF, Horikawa K, Hatsugai N, Higuchi Y, et al. 2012. Luminescent proteins
for high-speed single-cell and whole-body imaging. *Nat Commun* 3: 1262

Shaner NC, Lambert GG, Chammas A, Ni Y, Cranfill PJ, et al. 2013. A bright monomeric
green fluorescent protein derived from Branchiostoma lanceolatum. *Nat Methods* 10:
407-9

Sternson SM, Roth BL. 2014. Chemogenetic tools to interrogate brain functions. *Annu Rev*
Neurosci 37: 387-407

Su Y, Walker JR, Hall MP, Klein MA, Wu X, et al. 2023. An optimized bioluminescent
substrate for non-invasive imaging in the brain. *Nat Chem Biol*

Su Y, Walker JR, Park Y, Smith TP, Liu LX, et al. 2020. Novel NanoLuc substrates enable
bright two-population bioluminescence imaging in animals. *Nat Methods* 17: 852-60

Suzuki K, Kimura T, Shinoda H, Bai G, Daniels MJ, et al. 2016. Five colour variants of
bright luminescent protein for real-time multicolour bioimaging. *Nat Commun* 7:
13718

Ting JT, Lee BR, Chong P, Soler-Llavina G, Cobbs C, et al. 2018. Preparation of Acute
Brain Slices Using an Optimized N-Methyl-D-glucamine Protective Recovery
Method. *J Vis Exp*

Vierock J, Peter E, Grimm C, Rozenberg A, Chen IW, et al. 2022. WiChR, a highly
potassium-selective channelrhodopsin for low-light one- and two-photon inhibition of
excitable cells. *Sci Adv* 8: eadd7729

Welsh JP, Patel KG, Manthiram K, Swartz JR. 2009. Multiply mutated Gaussia luciferases
provide prolonged and intense bioluminescence. *Biochem Biophys Res Commun* 389:
563-8

Legends

FIGURE 1 – **Novel luciferase FRET pair characteristics. (A)** Bioluminescence emission spectra from the luciferases NanoLuc (Nluc) and eKL9h compared to the mNeonGreen-luciferase fusions NCS1 and NCS2. (**B**) Action spectrum of VChR1 compared with the peak emission wavelengths of NLuc, sbGluc, and NCS1/NCS2 illustrating the better alignment achievable using FRET. (**C**) Schematic of the LMOs compared for bioluminescence emission in HEK cell cultures. (**D**) Example image of bioluminescence recorded from HEK cell cultures in 4-well plates using IVIS imaging. (**E**) Summary bar graph showing the average radiance per copy of VChR1 recorded in the four groups. (**F**) Summary bar graph showing the average radiance of LMO3 and LMO7 relative to the radiance recorded from B7-tethered constructs (left).

FIGURE 2 – **Evaluation of the coupling efficiency of new LMOs in HEK cells.**
(**A**) Schematic of the four LMOs, with respective CTZ substrates, tested for coupling efficiency. (**B**) Effect of native coelenterazine (N-CTZ) on membrane current (Im) in a non-expressing HEK cell. (**C**) Left, EYFP fluorescence imaged in a stable HEK line expressing LMO3. Right, photocurrent response evoked by 480 nm light (1 s) and example trace showing the representative inward current amplitude recorded 5 min post N-CTZ application. (**D**) Left, EYFP fluorescence imaged in a stable HEK line expressing LMO5. Right, photocurrent response evoked by 480 nm light (1 s) and example trace showing the representative inward current amplitude recorded 5 min post H-CTZ application. (**E**) Effect of coelenterazine H (H-CTZ) on Im in a non-expressing HEK cell. (**F**) Left, mNeonGreen fluorescence imaged in a stable HEK line expressing LMO6. Right, photocurrent response evoked by 540 nm light (1 s) and example trace showing the representative inward current amplitude recorded 5 min post N-CTZ application. (**G**) Left, mNeonGreen fluorescence imaged in a stable HEK line expressing LMO7. Photocurrent evoked by 540 nm light for 1 s. Right, photocurrent response evoked by 540 nm light (1 s) and example trace showing the representative inward current amplitude recorded 5 min post H-CTZ application. (**H**) Example trace comparison of CTZ-induced inward current amplitude across stable HEK lines as indicated. (**I**) Summary bar graph of average CTZ-induced current amplitude generated with each construct. (**J**) Summary graph showing correlation

between CTZ-induced current and maximum photocurrent in all cells (**K**) Summary bar graph showing the coupling efficiency (CTZ-induced current amplitude / maximum photocurrent amplitude) achieved with each construct.

FIGURE 3 – **Comparison of LMO3 and LMO7 in acute mouse brain slices.**
(**A**) LMO3 expression in L5 neocortical neurons from fronto-parietal coronal slices (insets show a pipette-filled pyramidal cell visualized with AlexaFluor 594 and the EYFP tag post hoc). Right, firing response of the cell in response to 480 nm and 540 nm excitation. (**B**) Example trace showing the membrane potential response of the cell in *A* to N-CTZ (black trace, Vm) normalized to the recorded peak bioluminescence (blue, BL) and relative to control (vehicle, light grey trace). (**C**) Example trace showing a suprathreshold Vm response recorded from a LMO3-expressing L5 pyramidal neuron to CTZ. (**D**) LMO7 expression in L5 neocortical neurons from fronto-parietal coronal slices (insets show a pipette-filled pyramidal cell visualized with AlexaFluor 594 and mNeonGreen post hoc). Right, firing response of the cell in response to 480 nm and 540 nm excitation. (**E**) Example trace showing the membrane potential response of the cell in *D* to H-CTZ (black trace, Vm) normalized to the recorded peak bioluminescence (green, BL) and relative to control (vehicle, dark grey trace). (**F**) Example trace showing a suprathreshold Vm response recorded from a LMO7-expressing L5 pyramidal neuron to CTZ. (**G**) Representative example traces of CTZ-induced depolarization amplitude in LMO3 and LMO7-expressing cells relative to controls. (**H**) Summary graph showing the average CTZ-induced depolarization (circled data points represent recordings in *C* and *F*). (**I-L**) Effect of H-CTZ on the input-output function of LMO7-expressing L5 pyramidal neurons. (**I**) Example traces showing effect of CTZ on firing rate at indicated levels of square current injection. (**J**) Summary graph showing the frequency-current response of LMO7-expressing cells pre, during- and post CTZ application. (**K, L**) Summary bar graphs showing effect of CTZ on rheobase (**K**) and maximum number of action potentials (**L**) in LMO7-expressing L5 pyramidal neurons. Scale bars (A, D): 40 µm.

FIGURE 4 – **In vivo comparison of LMO3 and LMO7 (black and magenta throughout, respectively).** (*A*) Schematic of the in vivo experimental setup. The laminar probe was inserted perpendicularly to the primary somatosensory cortex. A fiber attached to the electrode delivered light stimuli. The luciferin (native or H-CTZ)

was infused over the top of the cortex, and bioluminescent light output was recorded
using an EMCCD camera. Example images of bioluminescence for each group are
depicted to the right of the schematic. The red lines indicate the location of the
electrode in each instance. (**B**) Average bioluminescent light output for the two
groups expressed as the percent change from baseline. Shaded error bars indicate
the bootstrapped 95% confidence interval. Inset: Dots are individual animal's peak
bioluminescent output, horizontal black bars indicate the median for each group. (**C**)
Average MUA response to the external LED stimulus for each group. Shaded gray
bars indicate the relative intensity of the LED stimulus through time. (**D**) The average
MUA response to luciferin delivery for the two groups. The onset of the 60 s luciferin
infusion began at time 0. (**E**) The linear fit of animals' LED evoked MUA to luciferin
evoked MUA. Dashed lines are the linear fits for the two groups and shaded error
bars are the 95% confidence intervals of the fitted lines. (**F**). The average luciferin
evoked MUA activity of the two LED evoked activity matched animal subsets.

FIGURE 5 – **LMO7-mediated bioluminescence-driven excitation of SNr neurons**
elicits motor behavioral changes. (**A**) Schematic illustration of the expression
strategy for unilateral photoactivation of substantia nigra pars reticulata (SNr)
neurons. AAV-hSyn LMO7 was injected in SNr of wildtype mice. (**B**) Instantaneous
time lapsed trajectory of the snout of a representative mouse in an open field treated
with either vehicle (left) or FFz (right). (**C**) Schematic illustration of body parts (snout
and base of the tail) extracted with DeepLabCut between consecutive video frames
for instantaneous rotational analysis. (**D**) Summary data showing the cumulative
rotation degrees in mice expressing LMO7 during a 160-second-long episode treated
with vehicle or FFz. Inset, calculation of rotation degree (θ) for each frame (40
millisecond bins). n = 3 animals. Data are presented as mean values ± SEM. (**E**)
Summary data showing the total degrees rotated in each experimental condition;
p=0.04. (**F**) Turning preference displayed as the ratio of ipsilateral over contralateral
revolutions during a 10-minute duration; p=0.01. (**G**) Total number of grooming bouts
(left, p=0.06) and total time spent grooming (right, p=0.006) between vehicle and FFz
treated mice.

SUPPLEMENTARY FIGURE 1 – **Average photocurrent amplitude and density**
across stable expression HEK lines. (**A**) Example traces of typical photocurrent

amplitudes evoked by 1 second illumination at maximum irradiance. (**B**) Summary graph of average VChR1 photocurrent amplitudes evoked at maximum irradiance across HEK lines. (**C**) Summary graph showing capacitance values of all HEK cells and across stable lines. (**D**) Summary graph showing average photocurrent density across HEK lines.

SUPPLEMENTARY FIGURE 2 – **Characterization of photocurrent responses to blue and green light in HEK cells.** (**A**) Example traces of photocurrent amplitudes evoked by 480 nm (left) and 540 nm (right) light in 1 second windows from -80 to +80 mV. (**B**) Summary graph of photocurrent-voltage relationship and reversal potential of VChR1 with blue and green light illumination (n = 4 cells from LMO3- and LMO7- expressing HEK lines). (**C**) Example traces of photocurrent amplitudes in response to increasing blue (left) and green (right) light intensity as indicated (**D**) Summary graph showing photocurrent amplitude relative to irradiance level with blue and green light (n = 3 LMO3 cells). Ee max; maximum irradiance level obtained, Ee sat; irradiance level sufficient to saturate photocurrent amplitude. Saturation occurs below maximum irradiance levels for both blue and green light.

SUPPLEMENTARY FIGURE 3 – **Expression of LMO7 in mouse neocortex after early postnatal bilateral AAV injection.** (**A**) Representative image showing the distribution of LMO7 (mNeonGreen) fluorescence at 4 weeks post AAV9-hSyn-LMO7 injection into the fronto-paritetal cortex of P2 Swiss Webster mice. (**B**, **C**) magnified images from 'A' showing LMO7-expressing neurons at 20x (**B**) and 40x (**C**). Scale bars; A: 2 mm, B: 200 µm, C: 100 µm.

SUPPLEMENTARY FIGURE 4 – **Average photocurrent amplitudes recorded from LMO3 and LMO7-expressing neurons.** (**A**) representative photocurrent traces (left) and average photocurrent amplitudes (right) recorded from LMO3 (black) and LMO7 (magenta) populations (480 nm excitation light, 1s). (**B**) scatter plot showing photocurrent amplitude relative to CTZ-mediated depolarization for each recorded cells (LMO3; n = 10, LMO7; n = 11).

Table 1. Properties and composition of tested LMOs

	LMO3	LMO5	LMO6	LMO7
Construct	sbGluc-VChR1	Nluc-VChR1	NCS1-VChR1	NCS2-VChR1
Light emitter	slowburn *Gaussia* luciferase	Nanoluc	sbAluc1–*d7N*mNeonGreen	mNeonGreen*d10C*–eKL9H
Peak em. λ (nm)	~ 470	~ 455	~ 518	~ 518
Substrate	Native coelente-razine (N-CTZ)	Coelenterazine H (H-CTZ) Fluorofurimazine (FFz)	Native coelente-razine (N-CTZ)	Coelenterazine H (H-CTZ) Fluorofurimazine (FFz)
Actuator	*Volvox* channel-rhodopsin 1	*Volvox* channel-rhodopsin 1	*Volvox* channel-rhodopsin 1	*Volvox* channel-rhodopsin 1

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure S1.

Figure S2.

Figure S3.

Figure S4.