1
2 **DNA hypomethylation activates Cdk4/6 and ATR to cause dormant origin firing**
3 **and cell cycle arrest that restricts liver outgrowth in zebrafish**
4

5 **Bhavani P. Madakashira, Elena Magnani, Shashi Ranjan, and Kirsten C. Sadler***
6
7 Program in Biology NYU Abu Dhabi, Abu Dhabi, UAE
8
9 *Corresponding author; Kirsten.edepli@nyu.edu

15 Running Title: Cdk4/6 causes DNA rereplication
16

Abstract

Coordinating epigenomic inheritance and cell cycle progression is essential for
organogenesis. UHRF1 connects these functions by facilitating maintenance of DNA
methylation and cell cycle progression. Here, we provide evidence resolving the
paradoxical phenotype of *uhrf1* mutant zebrafish embryos that have both activation
of pro-proliferative E2F target genes and increased number of hepatocytes in S-
phase, but the liver fails to grow. We find that Atr inhibition reduces DNA replication
and increases liver size in *uhrf1* mutants, suggesting that *uhrf1* mutant hepatocytes
have replication stress leading to Atr-mediated cell cycle inhibition and dormant
origin firing. We uncover persistent Cdk4/6 activation as the mechanism driving *uhrf1*
mutant hepatocytes into S-phase, activating Atr and restricting hepatic outgrowth.
Palbociclib treatment of *uhrf1* mutant embryos prevented aberrant S-phase entry,
and the DNA damage response. Palbociclib rescued most cellular and
developmental phenotypes in *uhrf1* mutants, except DNA hypomethylation,
transposon activation and the interferon response. Pro-proliferative genes were also
activated in a Cdk4/6 dependent fashion in the liver of *dnmt1* mutants, suggesting
DNA hypomethylation as a mechanism of Cdk4/6 activation. Thus, the
developmental defects caused by DNA hypomethylation are attributed to persistent
Cdk4/6 activation leading to DNA replication stress, dormant origin firing and cell
cycle inhibition, preventing hepatic outgrowth.

Introduction

Cell cycle progression during organogenesis requires precise coordination of the machinery that replicates the genome and those that pattern the epigenome. The chromatin landscape dictates the timing of regions of the genome get replicated early or late in S-phase and can influence when and where the origins of DNA replication fire [1]. Thus, the widespread changes to the epigenome that occurs during development when cell identity is established, in aging or in cancer can influence the pattern, timing or progression of DNA replication [2,3]. The crosstalk between the pathways that control cell cycle progression and epigenetic patterning during development has not been well studied.

DNA methylation is among the best studied epigenetic marks. The methylome is patterned during early vertebrate development and is maintained during cell division by a process that couples DNA replication to DNA methylation via interaction between proteins at the replication fork. Newly synthesized DNA is unmethylated and the hemimethylated structure of DNA generated at the replication fork is recognized by Ubiquitin-like, containing PHD and RING finger domains, 1 (UHRF1) which, through a base flipping mechanism [4-7], allows DNA methyltransferase 1 (DNMT1) to access and methylate the daughter strand [8-10]. Both proteins are anchored to the DNA replication fork [11,12] through interactions with PCNA [10], LIG [13] and modified histones [10,14,15]. Thus, the colocalization and coactivity of the methylation and replication factors assures faithful propagation of methylation patterns.

One way to assure that these factors act together is by co-regulating their expression. UHRF1, DNMT1 and all of the genes required for DNA replication are induced by the E2F family of transcription factors [16-20]. E2F is activated when the retinoblastoma (Rb1) protein becomes phosphorylated by the cyclin-dependent kinases 4/6 (CDK)-Cyclin D complex. Therefore, the downstream consequence of CDK4/6 activation in response to mitogen stimulation assures that the factors required for both DNA replication and DNA methylation are present as cells enter S-phase. This coupling is essential, as in both during development and in cancer cells, inactivation of either UHRF1 or DNMT1 blocks cell proliferation [21-29] by a mechanism that is not understood.

Just as these cell cycle regulatory pathways can regulate DNA methylation, the chromatin landscape influences the cell cycle. For instance, repressive epigenetic marks restrict E2F target gene accessibility, and removal of these repressive marks is essential for E2F function [30]. The p16^{INK4a} gene, which encodes a CDK4/6 inhibitor, is regulated by repressive epigenetic marks which can lead to suppression of p16^{INK4a}, causing inappropriate cell cycle entry [31]. Another study showed that with DNMT1/DNMT3B deficient cells, loss of 3D genome structure randomized replication timing [32]. These data demonstrate the complexity of the interplay between the ubiquitous cell cycle machinery and epigenetic modifiers to ensure fidelity of the epigenetic patterns as cells proliferate. There is a limited understanding of how these processes are regulated during the rapid cell cycles that take place during vertebrate development.

The developing liver provides an optimal model to study this relationship, as liver outgrowth is mediated by proliferation of differentiated hepatocytes. In zebrafish,

hepatocytes are specified and determined during the organogenesis stage of
development (between 24-48 hours post fertilization, hpf) [33,34] and the differentiated
hepatocytes proliferate while the liver undergoes morphogenesis during hepatic
outgrowth. This stage is largely complete by 120 hpf [35]. There are a few studies that
indicate that the mechanisms of cell cycle regulation during hepatic outgrowth utilize
the same signaling pathways that control proliferation in most other cell types. For
instance, in mice, genetic approaches showed that E2F regulates hepatocyte
proliferation during liver development and regeneration [36,37] and hepatocyte specific
knockout of Rb1 is sufficient to promote cell cycle entry, increased ploidy and
generated large nuclei in post-natal and adult mouse hepatocytes [38,39]. It is not
known if these mechanisms are conserved in zebrafish or how these and other
critical cell cycle regulators control organ growth.
128
We previously discovered a functional relationship between *uhrf1* and cell cycle
progression during hepatic outgrowth in zebrafish. *uhrf1* mutant embryos have
multiple developmental abnormalities, including a small liver [23,26,40]. These
phenotypes are first evident between 80-96 hpf and are exaggerated at 120 hpf,
resulting in lethality by 10 days post fertilization (dpf) [25]. The cellular phenotypes of
uhrf1 deficient hepatocytes are similar to those of *dnmt1* mutants: mutant larvae
have large, misshapen nuclei, near complete loss of nuclear structures [41], and
widespread cell death [23,25], which is dependent on Tnfa [42]. Most strikingly, at 120
hpf, when DNA replication is complete in wild-type larvae (WT), nearly half of all
mutant hepatocytes are undergoing DNA replication with a gene expression
signature typical for proliferating cells[25] despite the fact that the liver does not grow.
As in all other systems studied [8,9,26,43,44], *uhrf1* and *dnmt1* loss in zebrafish results in
DNA hypomethylation, which unleashes transposons and an interferon response to
these viral mimetics [42,45]. It is not known which cellular and molecular defects
contribute to the developmental phenotypes of these mutants.
144
Here, we resolving the paradoxical finding of the cellular phenotype and molecular
phenotypes associated with DNA replication and proliferation with the developmental
phenotype of hepatic outgrowth failure in *uhrf1* mutants. We identify Cdk4/6 activity
as the mechanism giving rise to most of the developmental and cellular defects in
uhrf1 zebrafish mutants, except for DNA hypomethylation, transposon activation and
induction of the interferon response. The Cdk4/6 inhibitor, Palbociclib, rescues all
embryonic morphological defects in *uhrf1* mutants, including liver size, DNA
rereplication, nuclear enlargement and euchromatinization and cell death. Strikingly,
embryo survival is improved by Palbociclib. Cdk4/6 signaling is also activated in
another model of DNA hypomethylation, in *dnmt1* mutants, suggesting that DNA
hypomethylation as the underlying mechanism. Since DNA replication does not
occur in *dnmt1* mutants and Dnmt1 depletion in *uhrf1* mutant hepatocytes blocks
DNA replication, we concluded that Dnmt1 is essential at the replication fork [11,12].
The Ataxia Telangiectasia And Rad3 Related (Atr) pathway is activated in *uhrf1*
mutants, signifying replication stress and we show that DNA replication in part
requires Atr signaling. These data provide new mechanistic insight into the
interaction between key cell cycle regulatory pathways and DNA methylation during
development, indicate the importance of the Cdk4/6 pathway in cellular
consequences of Uhrf1 depletion. This also suggests that the important
chemotherapeutic, Palbociclib, may modulate the cellular response to epigenetic
defects.

Results

DNA rereplication is a primary cellular response to *uhrf1* loss

We previously reported that *uhrf1* mutant livers have an increased number of cells that incorporate nucleotides during hepatic outgrowth, despite the livers being smaller sized [25]. To determine whether this was due to DNA replication or DNA repair, we investigated the pattern, timing and outcome of EdU incorporation in hepatocytes. FACS sorted hepatocytes from *uhrf1* and WT larvae at 120 hpf showed that 7% of *uhrf1* mutants had 4N DNA content compared to 2% of controls (Figure 1A), consistent with our previous finding of enlarged hepatocyte nuclei in *uhrf1* mutants [41], a phenotype that reflects increased ploidy [46]. We pulsed embryos at 80, 96 and 120 hpf for 30 minutes with EdU, sorted *uhrf1* mutants based on morphological phenotypes at 96 hpf and 120 hpf, and, since the phenotype is not apparent at 80 hpf, all embryos at this time point were individually genotyped and assessed for the number of cells with EdU incorporation. Nearly all hepatocytes in *uhrf1* mutants and their phenotypically normal sibling control larvae (heretofore referred to as controls) were EdU positive at 80 hpf. At 96, and 120 hours, this was reduced to 25% and 10%, respectively in controls. However, 55% and 43% of *uhrf1* mutant hepatocytes were EdU positive at these same time points (Figure 1B-C).

The stereotypical pattern of DNA replication has euchromatic DNA replicated in early S-phase, visualized as homogeneous EdU labeling in the nucleoplasm, followed by replication of DNA in the nuclear lamina during mid-S-phase and heterochromatin replicated last [47]. If the increase in EdU incorporation in *uhrf1* mutant hepatocytes reflected DNA rereplication, then these patterns should be apparent. Scoring S-phase stage based on the patterns of EdU incorporation showed that at 80 hpf, most hepatocytes in controls and *uhrf1* mutants were in early S-phase, while at 96 hpf and 120 hpf there was a significant increase in the number of *uhrf1*$^{-/-}$ hepatocytes in mid and late S-phase, respectively (Figure 1D). These data rule out the possibility that EdU incorporation in *uhrf1* mutant hepatocytes reflects DNA damage repair, as repair mediated nucleotide incorporation does not mirror the stereotypical stages of DNA replication [48,49]. Interestingly, we observed that *uhrf1*$^{-/-}$ hepatocytes incorporate EdU in a pattern that reflects a pattern of heterochromatin, even though these hepatocytes are completely devoid of the heterochromatin markers H3K9me3, H3K9me2, Hp1α (Figure 1E) and the nuclear lamina marker Lamin A [41]. We conclude that increased number of EdU positive hepatocytes in *uhrf1* mutants reflects DNA rereplication.

We hypothesized that the EdU incorporation phenotype in *uhrf1* mutant hepatocytes reflected either an acceleration of the fork progression rate or an increase in the number of active replication origins. To test this, we optimized the duration of the EdU pulse to be just below the limit of detection in control hepatocytes at 120 hpf (Figure S1). Following a 12 minute pulse with EdU, no hepatocytes were labeled at either 80 or 120 hpf in control larvae or in *uhrf1*$^{-/-}$ larvae at 80 hpf but nearly 50% of *uhrf1* mutant hepatocytes were labeled at 120 hpf (Figures 1F and Figure S1), and nearly all of these EdU positive *uhrf1*$^{-/-}$ nuclei were in early S-phase (Figure 1G).

216   This suggests that Uhrf1 deficiency either induces firing of more DNA replication
217   origins or increases replication fork speed.
218
219
220
CDK4/6 is activated in *uhrf1* mutant livers
To identify the basis of the DNA replication defect, we analyzed bulk RNAseq data
from pools of control and *uhrf1* mutant livers collected at 120 hpf [42], when the
hepatic developmental and cellular phenotypes are fully penetrant. Gene ontology
(GO) analysis using REVIGO revealed that downregulated genes were mainly
involved in hepatic functions, such as protein modification and lysis, signaling and
metabolic process, while the upregulated genes were enriched in pathways involved
in cell cycle regulation, DNA replication, anti-viral and immune response and cell
death (Figure 2A, Table S1). UPSET analysis of the differentially expressed genes
(DEGs) in all of the cell cycle pathways identified by REVIGO demonstrated that
many genes were common to multiple genesets, however a cluster of genes
involved in DNA replication and cell division were unique (Figure 2B; Table S1).
These data confirm and extend our previous findings [23,25,42] that *uhrf1* mutant livers
have aberrant cell cycle regulation.
235
We next used Ingenuity Pathway Analysis (IPA) of the unified set of all DEGs in the
GO term genesets related to the cell cycle to identify potential upstream regulators of
the cell cycle DEGs in *uhrf1* mutant livers. This uncovered a pattern whereby the
G1/S and G2/M checkpoint pathways were downregulated and cell cycle regulation,
proliferation and transition through the cell cycle were upregulated (Figure 2C, Table
S2). The key upstream regulators of the G1/S transition and S-phase entry and
progression, *Cdk4/6-Cyclin D, Cdk2-Cyclin E* and the DNA replication initiation
factor, *Cdc25a*, were upregulated in *uhrf1* mutant livers. Additionally, the cell cycle
inhibitor *cdkn1b* and the cell cycle regulated gene *rbl2,* which prevents access of
E2F1 and is a UHRF1 interacting protein [50], were downregulated (Figure 2D).
Together, these data suggest that loss of *uhrf1* results in repression of G1/S and
G2/M cell cycle checkpoints and activation of Cdk4/6 downstream targets that
promote G1/S transition and DNA replication.
249
These data establish a developmental timeline of the *uhrf1* mutant hepatic
phenotypes whereby DNA hypomethylation [42], and nuclear size defects are first
detected at 80 hpf [41], followed by DNA rereplication (Figure 1B-D,F) and cell death at
96 and 120 hpf. Since no cell cycle DEGs were detected at 80 hpf [25] and qPCR
analysis on RNA from pooled livers dissected at 96 hpf showed that several cell
cycle regulatory genes (*ccne1, ccna2)*, replication licensing factors (*mdm2, mcm3,*
orc3, cdc6) and processivity factors (*pola1, pold1* and *pcna)* were upregulated at this
time point (Figure 2E), we conclude that the DNA replication defect in uhrf1
hepatocytse occurs following loss of DNA methylation. Importantly, we found a
similar effect in the liver of neonatal mice at P4 that are deficient for *Uhrf1* in
hepatocytes ((*Alb:Cre^Tg/+^;Uhrf1^fl/f^ i.e. Uhrf1^HepKO^*) [44]. There was robust expression of
Cre (Figure 2F, inset) and downregulation of *Uhrf1*, indicating effective Cre-mediated
Uhrf1 depletion in these samples. All cell cycle genes analyzed showed increased
expression in *Uhrf1^HepKO^* livers (Figure 2F), suggesting that Uhrf1 plays a conserved
role in regulating the cell cycle during hepatic outgrowth by promoting S-phase entry
and DNA rereplication.

266
Cdk4/6 activation is required for the *uhrf1* mutant phenotype
We hypothesized that Cdk4/6 and E2F activation combined with loss of G1/S checkpoints could drive Uhrf1 deficient cells into S-phase. To test this, we used the Cdk4/6 inhibitor Palbociclib (PD in figures), which is an approved chemotherapeutic for breast cancer [51] that effectively suppresses DNA replication and cell proliferation in liver cancer cells [52]. We optimized the dose and timing of Palbociclib exposure to obtain the maximal tolerable concentration and administration timing to achieve no effects on control larvae based on survival, overall embryo phenotype, morphological criteria and liver size (Table S3). The final treatment scheme outlined in Figure 3A involved adding 20 µM Palbociclib in 0.5% DMSO or 0.5% DMSO as a control to 48 hpf embryos generated from an incross of *uhrf1*[+/-] adults which expressed nuclear localized signal targeted cherry transgene in hepatocytes (*Tg(fabp10a: nls-mcherry)*) to mark the livers. The resulting larvae were collected at 120 hpf, imaged to assess morphological features characteristic of *uhrf1* mutants and then all larvae were individually genotyped. In some experiments, larvae at 120 hpf were pulsed for 30 minutes with EdU to assess DNA replication.

At 120 hpf, *uhrf1* mutants are distinguished by smaller head, lower jaw deformity, microphthalmia, underdeveloped gut and small liver (Figure S2A; [23,25]). These phenotypes are fully penetrant at 120 hpf, so that on average, 25% of embryos resulting from an incross of *uhrf1*[+/-] adults display all of these phenotypes (Figure S2B), but only 7% of embryos produced from an incross of *uhrf1*[+/-] adults treated with 20 µM Palbociclib have all mutant phenotypes (Figure 3B). In treated clutches, 18% of embryos showed a partial mutant phenotype, with restoration of the lower jaw defect, increase in liver size, or both (Figure S2B-C). When these embryos were genotyped, we uncovered the expected genotype-phenotype relationship in 100% of embryos in DMSO treated clutches, so that all embryos displaying the mutant phenotypes were *uhrf1*[-/-] and all embryos that lacked these phenotypes were either *uhrf1*[+/+] or [+/-]. However, in Palbociclib treated clutches, embryos with a partial phenotype and even a few scored as normal were *uhrf1*[-/-] (Figure S2C-D).

Palbociclib had no significant effect on liver size in WT embryos, but significantly increased the size of the liver (Figure 3B-D) and eye (Figure S2E) in *uhrf1* mutants. The liver size of nearly all *uhrf1* mutants treated with Palbociclib exceeded the average liver size in DMSO treated mutants, and 20% of treated mutants had a liver size that was within the range of WT embryos (Figure 3C). Strikingly, Palbociclib also increased survival of *uhrf1* mutants at 9 dpf (Figure S2F).

To differentiate aspects of the *uhrf1* mutant phenotype that were a direct consequence of Uhrf1 loss compared to those that were mediated by Cdk4/6 activation, we assessed the effects of Palbociclib on the cellular aspects of hepatic phenotypes: EdU incorporation, nuclear size and shape and cell death. These parameters were unchanged by Palbociclib treatment of control embryos, but were significantly rescued in *uhrf1* mutants. The number of hepatocytes with EdU incorporation at 120 hpf was reduced from 40% to 28% in untreated compared to Palbociclib treated *uhrf1* mutants (Figure 3E). This was accompanied by a dramatic change in the nuclear morphology: DMSO and Palbociclib treated control hepatocyte nuclei displayed clear nuclear rims and distinct heterochromatin foci, with prominent H3K9me3 staining, whereas *uhrf1* mutant nuclei were large and oblong, with

significant nuclear debris (Figures 3F). The nuclear size, shape, DNA patterning and
H3K9me3 staining in *uhrf1* mutant hepatocytes were nearly completely restored to
the WT phenotype by Palbociclib treatment (Figure 3F-G). Finally, the amount of cell
death in *uhrf1* mutant livers was significantly reduced by Palbociclib (Figure 3H).
Therefore, Cdk4/6 activity is required for both the developmental and cellular
phenotypes caused by Uhrf1 loss.
322
Since Uhrf1 is a required component of the maintenance methylation machinery [8,9],
DNA hypomethylation is a direct consequence of Uhrf1 loss. Assessment of bulk
DNA methylation levels confirmed this, as both DMSO and Palbociclib treated *uhrf1*
mutants had significant DNA hypomethylation (Figure 3I). Since Palbociclib
treatment rescued most of the developmental and cellular defects in *uhrf1* mutants
despite persistent DNA hypomethylation, we conclude that the developmental and
cellular phenotypes caused by Uhrf1 loss is due to persistent activation of Cdk4/6
and is not a direct consequence of DNA hypomethylation.
331
Cdk4/6 inhibition blocks cell cycle and Tnfa target genes activation in *uhrf1*
mutant livers but does not suppress TE activation or the interferon response
We hypothesized that the cellular and developmental phenotypic rescue of *uhrf1*
mutants by Palbociclib was attributed to downregulation of the G1/S transition and S-
phase promoting genes. In contrast, we expected that activation of transposable
elements (TEs) and the ensuing interferon response [42,45] would not be affected by
Palbociclib, since these are directly mediated by DNA methylation loss in *uhrf1*
mutants.
340
We assessed this using bulk RNAseq analysis of the livers of 120 hpf *uhrf1* mutants
and their phenotypically normal siblings (control) treated with DMSO or Palbociclib.
Principal component analysis showed that controls cluster together regardless of
treatment, while mutants treated with Palbociclib were closer to control samples
(Figure 4A). Palbociclib treatment of control larvae resulted in modest
downregulation of genes that play roles in hepatocyte function and metabolism, but
had no effect on genes that regulate the cell cycle (Figure S3A-B; Table S4, S5). In
contrast, Palbociclib dramatically changed the hepatic transcriptome in *uhrf1*
mutants, reducing the number of DEGs from 2,272 in DMSO treated mutants to 650
in the liver of *uhrf1* mutants treated with Palbociclib, 560 DEGs common to both
(Figure 4B, Table S6). While the common DEGs showed a very high correlation in
their expression (r=0.974), the genes that were unique to each dataset genes were
less correlated (r= 0.666; Figure S3C). In contrast, the upregulation of transposable
elements (TEs) in *uhrf1* mutant livers was not suppressed by Palbociclib (Figure
S3D).
356
We predicted that Cdk4/6 inhibition in *uhrf1* mutants would reduce the induction of
E2F targets, but would have no effect on the interferon response as this is caused by
TE activation. Unsupervised clustering of the unified set of DEGs from both DMSO
and Palbociclib treated mutants showed that most of the upregulated genes in
DMSO treated *uhrf1* mutant livers were still upregulated, but to a lesser extent in
Palbociclib treated mutants (Figure 4C, cluster 5), however, a subset of genes
remain elevated in both samples (cluster 4 and 5) (Figure 4C, Table S7). GO
analysis of these two clusters shows that DNA replication related genesets were
predominant in cluster 5, and that immune pathways were predominant in cluster 4

(Figure 4D-E, Table S8). Specific analysis of key genes involved in the pre-replication phase of cell cycle, such as *ccna2*, the *orcs*, *mcms* and *cdc45* which is important for origin firing at the replication forks in association with *mcms* [53,54], showed that they were highly induced in DMSO treated mutants, but expressed at significantly lower levels in the liver of Palbociclib treated *uhrf1* mutants (Figure 4F, Figure S3E and Table S9).

Interestingly, the immune genes which were all upregulated in *uhrf1* mutants showed a nuanced response to Palbociclib treatment: interferon response genes remained highly expressed while genes downstream of Tnfa were downregulated in the liver of *uhrf1* mutants exposed to Palbociclib (Figure 4F, Table S10). Since CDK6 positively regulates TNFa [55], we conclude that Tnfa induction in *uhrf1* mutants is attributed to Cdk4/6 activation, in contrast to interferon related genes which are due to TE activation.

We further investigated the effect of Palbociclib on the expression of cell cycle regulatory genes in *uhrf1* mutants using IPA. We first analyzed the genes activated in *uhrf1* mutants compared to controls treated with DMSO and found significant activation of genes regulating G1/S *ccnd1*, *ccne* and *cdk4/6*, which lead to upregulation of transcription factors *e2fs, hdac, ftdp1*, as well as activation of DNA damage sensors *atm, atr* and *tp53* (Figure S4A). qPCR analysis of several of these genes confirmed that Palbociclib downregulated their expression the liver of *uhrf1* mutants (Figure S4B). We next compared the gene expression patterns in *uhrf1* mutants treated with Palbociclib to mutants treated with DMSO, and identified 280 DEGs (Figure S4C; Table S11). IPA analysis of these DEGs showed that genes involved in cell cycle checkpoints were upregulated and genes involved in cell cycle control and S-phase were downregulated in *uhrf1* mutant livers treated with Palbociclib (Figure 4G) but that Palbociclib had no effect on *atm/atr* and *tp53* induction in *uhrf1* mutants (Figure 4G, Figure S4D, Table S12), suggesting that DNA damage persists even when DNA replication is suppressed.

Together, these data indicate that Cdk4/6 activation does not affect DNA methylation, TE induction, interferon response and DNA damage caused by Uhrf1 loss, but is required for the downregulation of G1/S checkpoints, E2F targets, DNA rereplication and cell death. Based on our previous finding that blocking Tnfa reduced cell death in *uhrf1* mutant livers but has no effect on the mutant phenotype or liver size at 120 hpf [42], combined with the current finding that Palbociclib reduced Tnfa activation, we deduce Palbociclib prevents cell death in *uhrf1* mutants by inactivation of Tnfa.

Cdk4/6 is activated by *dnmt1* mutation
To understand if the cellular phenotypes that cause hepatic outgrowth failure in *uhrf1* mutants are attributed to DNA hypomethylation, we assessed larvae with a loss of function mutation in *dnmt1* [24]. These have a strikingly similar phenotype to *uhrf1* mutants, including DNA hypomethylation, large and disorganized hepatocyte nuclei and a small liver [25,26,41,42,56]. We asked whether the cell cycle related geneset that we found to be differentially expressed in *uhrf1* mutant livers (the unified set of all cell cycle genes in Figure 2B) were also differentially expressed in bulk RNAseq data from the liver of 120 hpf *dnmt1* mutants [42]. Similar to *uhrf1* mutants, many of the genes required for DNA replication (*pcna, orcs, mcms, cdc45* and *pold2*) were

upregulated in *dnmt1* mutant livers (Figure 5A). Surprisingly, however there were a
lower number of hepatocytes that incorporate EdU in *dnmt1* mutants at both 96 hpf
and 120 hpf (Figure 5B). Dnmt1 loss prevents DNA replication, consistent with a
requirement for Dnmt1 at the replication fork [11,12]. Previous analysis showed that
uhrf1 and *dnmt1* mutant livers have a similar pattern whereby genes required for
DNA replication were upregulated [25,42]. IPA provided a more detailed analysis, while
the Atm and Tp53 mediated DNA damage responses were upregulated in both
samples, some genes that regulate S-phase (*chek1, chek2, tlk1/2, cdc25),* M phase
progression (*cdk1, cyclin B*) and Atr mediated DNA damage response signaling were
upregulated only in *uhrf1* mutant livers (Figure S5). Thus, while both mutants have a
global upregulation of cell cycle promoting genes and the Atm mediated response to
double stranded breaks, *uhrf1* loss specifically causes the response to single
stranded breaks.

If Cdk4/6 is upstream of cell cycle activation in *uhrf1* and *dnmt1* mutants, treatment
of *dnmt1* mutants with Palbociclib should downregulate these genes. Palbociclib had
no significant effect on the developmental phenotype or liver size in *dnmt1* mutants
(Figure 5C-D). While there was variability in the induction of cell cycle genes in
dnmt1 mutant livers, treatment with Palbociclib decreased the expression of
assessed cell cycle genes, with *ccne1, cdc6* and *pold1* reaching significance. As in
uhrf1 mutants, Palbociclib had no effect on interferon response gene expression in
dnmt1 mutant livers (Figure 5E). Thus, while *dnmt1* and *uhrf1* loss activate Cdk4/6
and E2F target genes, derepress TEs and induce an interferon response, these
responses are independent. Induction of cell cycle genes requires Cdk4/6 signaling
while TE activation and the interferon response do not. Moreover, the finding that
dnmt1 mutants are not rescued by Palbociclib reflects the fact that Dnmt1 deficient
cells cannot undergo DNA replication, likely due to the requirement of Dnmt1 at the
replication fork [12,57,58].

DNA replication in *uhrf1* mutant hepatocytes requires *dnmt1*
We examined the relationship between Uhrf1 deficiency, DNA replication and
Dnmt1. Western blotting of whole embryos from control and *uhrf1* mutants [59] showed
a significant increase in Dnmt1 levels starting at 80 hpf in mutants (Figure 6A),
suggesting that Dnmt1 upregulation is an early molecular consequence of *uhrf1*
mutation. Cdk4/6 activity has been shown to stabilize Dnmt1 [60] and thus one
mechanism by which Palbociclib could reduce DNA rereplication in *uhrf1* mutants is
by reducing Dnmt1 levels. To test this, we assessed Dnmt1 protein levels in 120 hpf
hepatocytes using immunofluorescence. Dnmt1 was not detected in controls, but
there was strong expression in 70% of *uhrf1* mutant hepatocytes, and this was
depleted in *uhrf1* mutants treated with Palbociclib treatment (Figure 6B). Therefore,
Dnmt1 protein stabilization in *uhrf1* deficient hepatocytes requires Cdk4/6. This could
either be a direct effect of Cdk4/6 on Dnmt1 stability or could be indirect via induction
of dnmt1 transcriptional activation by E2F.

To determine if Dnmt1 was required for the EdU incorporation phenotype in *uhrf1*
mutant hepatocytes, we used CRISPR-Cas9 to target *dnmt1*. Efficacy was verified
using an *in vivo* assay for DNA methylation in which the UAS:GFP cassette is
silenced by DNA methylation, and GFP expression occurs in response to Gal4 when
methylation is lost on the UAS promoter (*Tg(c269off^{+/-}; 14X UAS:dsRed)* [42,61]. All
Tg(c269off^{+/-}; 14X UAS:dsRed^{+/-});dnmt1^{cr} larvae had GFP expression in the head at

120 hpf (Figure S6A), but had a mild phenotype compared to *dnmt1* mutants (Figure
5C). Crispr deletion of *dnmt1* did not change liver size of *uhrf1* mutants (Figure 6C-
D) or the volume or shape of the *uhrf1* mutant hepatocyte nuclei (Figure S6B-D).
dnmt1 deletion did not rescue aberrant expression of cell cycle genes in *uhrf1*
mutant livers (Figure 6E) but did significantly reduce EdU incorporation in
hepatocytes compared to uninjected *uhrf1*[-/-] (Figure 6F). We conclude that Dnmt1 is
required for DNA replication in *uhrf1* mutant hepatocytes.

Atr is required for DNA replication and hepatic outgrowth failure in *uhrf1*
mutants

ATR responds to DNA replication stress by stabilizing replication forks, delaying or
blocking the progress of the cell cycle, promoting DNA repair [62,63]. In response to
replication stress, ATR activates dormant origins while repressing firing within those
that are not yet activated [64]. We found Atr to be specifically activated in *uhrf1*
mutants, but not *dnmt1* mutants, suggesting that the DNA replication phenotype in
uhrf1 mutants, which is absent from *dnmt1* mutants, causes replication stress
mediated Atr activation. To test this, we established a treatment protocol using the
highly selective and potent ATR inhibitor VE- 821 [65] during zebrafish development.
While treatment from 48-120 hpf with 10 µm VE- 821 had no effect control larvae
and had minimal effects on the overall *uhrf1* mutant phenotype, it significantly
increased liver size (Figure 7A-B). Moreover, EdU incorporation in hepatocytes was
reduced from 36% to 24% in untreated and VE- 821 treated *uhrf1* mutants,
respectively (Figure 7C-D). This suggests that DNA replication in uhrf1 mutants is
dependent on Atr.

Discussion

We report that Cdk4/6 activation is the mechanism accounting for the unusual cell
cycle defect caused by *uhrf1* mutation in zebrafish and present a model that
illustrates the relationship between DNA methylation, cell cycle regulation, DNA
replication and development (Figure 7E). The striking rescue of the cellular and
developmental defects in Uhrf1 deficient zebrafish embryos by Cdk4/6 inhibition
suggests that S-phase entry causes all the downstream phenotypes. The data
presented here, combined with our finding that reducing cell death does not rescue
liver size [42], indicates that the cell cycle defect is the main mechanism of hepatic
outgrowth failure. Surprisingly, the cellular and developmental phenotypes resolve,
despite persistent DNA hypomethylation, TE activation and an interferon response.
This finding differentiates between the direct targets of Uhrf1 - i.e. DNA
hypomethylation – and those that are secondary consequences – i.e. DNA
rereplication. These data also show that loss of DNA methylation and TE activation
per se do not cause the developmental defects that occur in *uhrf1* mutants. Rather it
is the downstream consequences of DNA hypomethylation which are responsible.

The finding of similar Cdk4/6 dependent gene expression in *dnmt1* and *uhrf1*
mutants suggests that DNA hypomethylation is the mechanism for Cdk/4/6

activation, although the mechanism of Cdk4/6 activation is not clear. We previously
published that *uhrf1* mutant hepatocytes have both an abnormal DNA replication
pattern and widespread cell death [23,25]. Importantly, the role of Uhrf1 in regulating the
hepatic cell cycle is conserved in the developing mouse liver. We demonstrate that
both *uhrf1* and *dnmt1* mutants activate of cell cycle genes in a Cdk4/6 dependent
manner, but while *uhrf1* mutant hepatocytes undergo DNA rereplication, *dnmt1*
mutant hepatocytes do not. Since DNA replication in *uhrf1* mutant hepatocytes was
rescued by *dnmt1* depletion, we conclude that either the *uhrf1* mutant phenotype
could be partly attributed to elevated Dnmt1 levels or, more likely, it is due to the
requirement for Dnmt1 for a functional replication fork [57,58]. Hence, Dnmt1 is both
upstream and downstream of Cdk4/6 in this model. The finding that Atr activation is
required for DNA replication and small liver size in *uhrf1* mutant hepatocytes
suggests Atr signaling activates DNA replication and prevents cell cycle progression.
Taken together, these results provide mechanistic insight into cell cycle regulation
during development and point to a functional relationship between epigenetic
regulation and cell cycle control.

Uhrf1 loss has been implicated in the mis-regulation of both G1 and G2/M phases of
the cell cycle [18,28,66] and here we provide evidence for a functional role in regulating
DNA replication. There are several possible mechanisms underlying this phenotype.
The first, discussed above, is that DNA replication stress activates Atr to prevent cell
cycle progression while promoting dormant origin firing (Figure 7E). During normal
DNA replication, fork speed has to be moderated for the fidelity of normal cell cycle,
with an increased fork speed triggering replication stress and DNA damage response
[67,68]. However during DNA rereplication, DNA synthesis was reported to progress
slowly, and repeatedly utilize the same main origins [69]. In contrast, unscheduled
firing of dormant origins can lead to increased and uncoordinated DNA synthesis [3,70].
While the experiments required to assess fork speed is challenging in whole
embryos, the Atr inhibition data in *uhrf1* mutants supports the hypothesis that the
DNA rereplication in *uhrf1* deficient hepatocytes is caused by Atr dependent firing of
multiple dormant origins, not a repeated firing of the same origins. The question of
why origins of replication can be fired at all if there is a checkpoint activated by ATR
can be answered by one study where ATR/ Chk1 pathway activation in response to
replication stress caused dormant origin firing within active replication clusters [71-73].
Hence, in *uhrf1* mutants, Atr activation could be activated by two mechanisms: either
directly by Cdk4/6 and/or as a downstream effect of replicative stress brought on by
Cdk4/6 activation. Further investigation between the CDK4/6 pathway and the
players in the ATR pathway will provide better insight linking cell cycle checkpoint,
replication stress and dormant origin firing leading to DNA re replication in *uhrf1*
deficient hepatocytes.

Another possible mechanism of the DNA replication defect in *uhrf1* mutants could
involve a defect in regulating which origins fire due to a failure to degrade replication
licensing factor Cdt1, which is required to prevent reuse of replication origins. Since
the levels of Cdt1 protein are regulated by the Cul4–Ddb1–Cdt2 E3 ubiquitin ligase
complex [74], it is feasible that Cdt1 stabilization due to nuclear Cyclin D1/CDK4, as
shown in other models [75], could result in repeated origin use. Further, Cyclin
D1/CDK4 during S-phase specifically represses transcription of *CUL4,* resulting in
CDT1 stabilization [76]. Our finding that *uhrf1* loss induced overexpression of *ccnd1*
and *cdk4* and downregulation of *cul4* suggests that Ctd1 levels may persist and

promote DNA rereplication [77]. Further biochemical analysis is required to evaluate
this mechanism.

The most straightforward conclusion to explain how Palbociclib prevents DNA
replication in *uhrf1* mutants is through reinstatement of Rb mediated suppression of
E2f. Another possibility is that Cdk4/6 inhibition depletes Dnmt1, which we have
shown is required for DNA re replication. Uhrf1 targets Dnmt1 for degradation [78-82]
and CDK4 phosphorylates DNMT1, preventing its degradation and therefore
Palbociclib treatment reduces DNMT1 protein levels [83]. Our data suggest that in
uhrf1 mutant embryos, Dnmt1 accumulation is attributed to increased Cdk4/6
activity, as inhibition of Cdk4/6 caused significant reduction in Dnmt1 levels.
However, it is also possible that this effect was mediated through decreased E2f
mediated *dnmt1* transcription. We conclude that Dnmt1 levels are permissive to the
unscheduled replication phenotype in *uhrf1* mutant hepatocytes.
568
We were intrigued by the finding that Palbociclib blocked cell death in *uhrf1* mutant
hepatocytes. In other studies, Palbociclib has been shown to activate NF-κB and
TNF-induced gene expression in mammalian cells [84]. Given our previous finding that
Tnfa depletion in *uhrf1* mutants prevented hepatocyte cell death but did not rescue
liver size[42], we attribute the rescue of cell death in Palbociclib treated *uhrf1* mutants
to the downregulation of Tnfa signaling, and conclude that cell death is not caused
by cell cycle defects. These data also show that the small liver phenotype in this
context is not caused by cell death, but rather is due to failure of hepatocyte
proliferation.
578
Our data point to DNA hypomethylation as an upstream signal that leads to Cdk4/6
activation in hepatocytes. How this occurs is not known, but other studies suggest
that the p16^{INK4a} inhibitor of CDK4/6 could be suppressed through DNA methylation
[31,85], and inhibiting p16^{INK4a} leads to increased cell cycle progression and increased
DNMT1 expression [31]. While it would be surprising if *uhrf1* and *dnmt1* mutation
caused hypermethylation of the p16^{INK4a} regulatory sequences or any other Cdk4/6
inhibitor, it is possible that loss of methylation causes redistribution of another
repressive epigenetic marks to the p16^{INK4a} regulatory regions, as we described in
other contexts [44]. Alternatively, loss of DNA methylation could derepress Cyclin D,
leading to perpetual Cdk4/6 activation.
589
In summary, this work provides the first insight into two distinct functions of Uhrf1
during liver development. Here we show that Uhrf1 loss induces Cdk4/6, E2f and all
the genes that are required for S-phase. This is paradoxical, as UHRF1 is
overexpressed in nearly all types of solid tumors [86], including HCC [78,87], Palbociclib
effectively suppressed cell proliferation in tumor growth in preclinical models of HCC
by promoting a reversible cell cycle arrest [52]. Therefore, it is possible that in HCC,
UHRF1 overexpression can act as a dominant negative to activate CDK4/6 and
promote cell proliferation, and future studies may investigate whether Palbociclib
functions as a therapeutic by counteracting the oncogenic effects of UHRF1.
599

Materials and Methods

Zebrafish Husbandry and Genotyping
Adult zebrafish were maintained in a circulating aquaculture system on a 14:10 h light: dark cycle at 28 °C. The *uhrf1*[hi272] allele [88] was maintained by genotyping adults to identify heterozygous carriers as described [88]. Homozygous mutant embryos were generated by crossing heterozygous adults and were identified based on distinctive phenotypes as described [23,25,26] or by genotyping individual embryos as described [23]. *uhrf1*[hi272+/−] adults were genotyped by PCR as described [23]. *Tg(fabp10a: nls-mcherry)* in *uhrf1*[hi272+/−] and *Tg(fabp10a: CAAX-EGFP)* in *uhrf1*[hi272+/−] background was used for all experiments for ease of liver detection [78]. Homozygous *uhrf1*[hi272-/−] mutant larvae are hereafter called *uhrf1*[−/−]. Homozygous *dnmt1*[s904-/-] mutants (*dnmt1*[-/-]) were generated by crossing *dnmt1*[s904+/−] adults in *dnmt1*[s904+/−]; *Tg(c269*[off]*; 10XUAS:dsRed); (fabp10a: Gal4; cmlc2: GFP)* background to monitor for DNA methylation in the liver as described [42]. All protocols were approved by the NYU Abu Dhabi Institutional Animal Care and Use Committee (IACUC).

Mice Husbandry and Usage
Mice maintenance and experiments were approved by the New York University Abu Dhabi Animal Care and Use Committee. Temperature, humidity, and light:dark cycles were controlled and mice were fed food and water ad libitum. Mice with hepatocyte-specific deletion of the Uhrf1 gene (*Uhrf1*[fl/fl]*;Alb-Cre*; referred to as *Uhrf1*[HepKO] mice) were generated by crossing mice homozygous for floxed *Uhrf1* (*Uhrf1*[fl/fl]) with mice expressing the Cre recombinase under the liver-specific albumin promoter back- crossed onto the *Uhrf1*[fl/fl] background (*Alb-cre*[Tg/+]*; Uhrf1*[fl/+]) described previously [44]. P4 mouse embryos were sacrificed and their livers were collected, flash frozen in liquid nitrogen, and stored at -80 °C for subsequent analysis.

FACS Analysis of the Cell Cycle
For FACS, 20 *uhrf1*[-/-]*; Tg(fabp10a: nls-mcherry)* and 20 *Tg(fabp10a: nls-mcherry)* sibling larvae were collected and used for single cells suspension. Single cell suspension was prepared by incubating larvae at 37 °C for 20 minutes in Trypsin-EDTA 0.05% (Thermo Fisher). After 20 minutes 10% FBS was added to stop the activity of the trypsin and cells were centrifuged at 2000 g for 7 minutes at 4 °C followed by 2 washes in PBS. Cells were resuspended in PBS and fixed with ice cold ethanol overnight. Cells were staining in PBS containing 10 μg/ml Hoechst (Thermo Fisher) for 10 minutes in dark and washed twice with PBS. Cells were finally resuspended in PBS containing 1% BSA (Sigma Aldrich) and 10 μg/ml RNaseA (Invitrogen) and acquired at FACS Aria II (BD Bioscience). Hepatocytes were selected by gating for mCherry[+] population, and Hoechst levels recorded for analysis. Analysis was performed with FlowJo 10 and plotted in GraphPad Prism 9.

CRISPR/Cas9 Generation and T7 Endonuclease Assay
4 sgRNAs for *dnmt1 was* designed by using ChopChop (https:// chopchop.cbu.uib.no/). sgRNA for *slc45a2* (gene involved in pigmentation) were

previously designed and validated [40,89]. Genotyping primers were designed by
Primer3 (https://bioinfo.ut.ee/primer3- 0.4.0/) and validated in USCS Genome
Browser (https://genome.ucsc.edu/cgi-bin/hgPcr). sgRNAs were produced by sgRNA
IVT kit (Takara Bio) by following the manufacturer's instructions and RNA was
isolated by Trizol (Invitrogen). sgRNAs were quantified by Qubit RNA BR kit and
diluted at 50 ng/μl and stored as single use aliquots. The efficiency of the sgRNAs
were assessed by injecting all 4 *dnmt1* sgRNAs into WT embryos with equal volume
of previously diluted nls-Cas9 protein (IDT; 0.5 μl of nls-Cas9 added with 9.5 μl of 20
mM HEPES; 150 mM KCl, pH 7.5) and sgRNAs, incubated at 37 °C for 5 minutes
and then 1 nl was injected in 1–2 cells stage embryos. At 24–72 hpf, 12–16 sgRNA
injected embryos were individually collected and genomic DNA was extracted by
heat shock denaturation in 50 mM NaOH (95 °C for 20 minutes). For each embryo,
PCR was performed on genomic DNA by using Q5 High-Fidelity Taq Polymerase
(New England Biolabs) followed by T7 endonuclease I assay (New England Biolabs)
to detect indel mutations. For T7 endonuclease I assay, 10 μl of PCR product was
incubated with 0.5 μl of T7e1 enzyme (New England Biolabs) for 30 minutes at 37°C.
Digested and undigested fragments were run in parallel in 2% agarose gel to assess
the presence of indels. Efficiency was calculated as the number of embryos that
show a positive result based on T7e1 assay divided by the total number of embryos
assayed for the sgRNAs. Once established for efficiency, the pool of 4 *dnmt1*
sgRNAs were injected into the 1 cell embryos generated by an incross of *uhrf1*[+/-]
adults as previously described. Not injected embryos or *slc45a2* crispants were used
as control. The resulting F0 larvae were considered crispants. For each clutch, *uhrf1*[-/-]
mutants were divided from phenotypically WT siblings at 5 dpf based on
morphological differences and T7e1 assay. sgRNA sequence information and
genotyping primers are provided in Table S13.
673
Inhibitor Treatment
The CDK4/6 inhibitor Palbociclib (PD-0332991, Sigma Aldrich) [52] was dissolved in
DMSO (Sigma) to yield a 10 mM stock solution and stored at -80 °C. For the *in vivo*
experiments, Palbociclib was dissolved in 0.5% DMSO embryo water at a
concentration of 20 μM. 48 hpf dechorionated embryos in 6 well plates were treated
either with Palbociclib in 0.5% DMSO or 0.5% DMSO alone and collected at 120 hpf
for further analysis. The ATR inhibitor VE- 821 (ab219506, Abcam) [90] was dissolved
in DMSO (Sigma) to yield a 20 mM stock solution and stored at -80 °C. For the *in*
vivo experiments, VE- 821 was dissolved in embryo water at a concentration of 10
μM. 48 hpf dechorionated embryos were treated either with VE- 821 or DMSO
(1:2000 dilution) in embryo water and collected at 120 hpf for further analysis.
685
Protein Lysates and Western Blotting
For western blot, whole larvae from *uhrf1*[-/-] and control siblings were collected at 80,
96 and 120 hpf in RIPA buffer (10mM Tris-HCl pH 8.0, 1mM EDTA, 0.5mM EGTA,
1% Triton X-100, 0.1% Sodium Deoxycholate, 0.1% SDS, 140mM NaCl
supplemented with protease inhibitor cocktail (Machery-Nagel), sonicated using a
hand sonicator (Hielscher Ultrasonics), and centrifuged at 14000 g for 15 minutes at
4°C to remove cell debris. Supernatant was collected and SDS-PAGE loading
Leammli buffer (BioRad) was added and run on SDS-PAGE gel. Volume
corresponding to 4-5 larvae was loaded on 10% acrylamide gels, transferred onto
PVDF membranes, blocked with 5% w/v powdered skimmed milk (Sigma) in TBST

(20 mM Tris–HCl, 150 mM NaCl, 0.1% v/v Tween 20, pH 8.0) for 1 hour at room
temperature and incubated overnight in primary antibody (1:100 anti-Dnmt1
polyclonal, Santa Cruz, 1:1000 H3 polyclonal, Santa Cruz) diluted in blocking buffer
at 4°C. The following day membranes were washed 3 times n TBTS for 5 minutes
and incubated in secondary antibody (1:2500 anti-Rabbit HRP conjugated, Promega)
diluted in blocking buffer for 1 hour at room temperature and washed 3 times with
TBTS for 5 minutes at room temperature. Chemiluminescent signals using Pierce™
ECL Western Blotting Substrate (Thermo Fisher Scientific) or Clarity ECL substrate
(BioRad) and imaged using Bio-Rad ChemiDoc. Immunoblot bands are quantified by
densitometry using GelAnalyzer (http://www.gelanalyzer.com) and plotted using
GraphPad Prism 9. 3 independent clutches were used for quantification. Levels of
Dnmt1 are normalized on level of H3 for each sample and then for the level of Dnmt1
of 80 hpf wt looking larvae for that clutch.
709
RNA and DNA Extraction
For each clutch, 10 to 20 livers were microdissected from *dnmt1*[-/-] and their controls
or *uhrf1*[-/-] and their controls in Tg(fabp10a: CAAX-EGFP) background larvae to
facilitate liver identification according to published protocols [91]. RNA was extracted
using Trizol (Invitrogen) following the manufacturer's instructions with modifications.
Briefly, during precipitation in isopropanol, 10 µg of Glycoblue (Thermo Fisher
Scientific) was added, and precipitation performed overnight at –20 °C followed by 1
h centrifugation at 12,000× g at 4 °C. The resultant RNA pellet was resuspended in
water and quantified by Qubit RNA High Sensitivity kit (Thermo Fisher). For genomic
DNA extraction, each embryo or 10 livers were subjected to DNA extraction buffer
(10 mM Tris-HCl pH9, 10 mM EDTA, 200 mM NaCl, 0.5% SDS, 200 µg/mL
proteinase K) followed by DNA precipitation with isopropanol as previously described
[45]. DNA was resuspended in water and quantified by Qubit dsDNA High Sensitivity
kit (Thermo Fisher).

Slot Blot Analysis of 5-MeC
Slot blot was performed as previously described [25]. Briefly, 2 ng of genomic DNA
was denatured in 400 mM NaOH/10 mM EDTA and blotted onto nitrocellulose
membrane in duplicate for dsDNA and 5-MeC using a slot blot apparatus.
Membranes were incubated 1 h at 80 °C, blocked with 5% skim milk in TBST (37
mM NaCl, 20 mM Tris pH 7.5, 0.1% Tween 20), and incubated overnight at 4 °C in
either anti-dsDNA (Abcam, 1:8000 in 2% BSA in TBST) or anti-5-methyl-cytosine
(m5C– Aviva Biosystem clone 33D3, 1:2000 in 2% BSA in TBST). Membranes were
washed in TBST and probed with anti-mouse HRP secondary antibody (Promega;
1:5000 in 2% BSA in TBST) for 1 h at room temperature followed by development in
ECL (Thermo Fisher Scientific). ChemiDoc (BioRad) was used to detect and quantify
the chemiluminescent signal. Gel Analyzer was used to measure the signals and
ratio between 5-MeC and dsDNA was plotted for controls and mutants in each clutch
using GraphPad Prism 9.

cDNA Production and qPCR
RNA extracted from microdissected livers was retrotranscribed to cDNA using
Qscript cDNA synthesis kit (Quanta Bio) following the manufacturer's instructions.
cDNA was diluted to 1 ng/µL, and 5 µL was used per reaction for qPCR using
Maxima R SYBR green/ROX master mix (Thermo Fisher Scientific). *rplp0* was used
to normalize expression levels by using the calculations for delta-Ct, and control

siblings were used to calculate delta-delta-Ct (DDCt) as previously described [92]. To
determine changes in expression between control and experimental samples, the
fold change was calculated and expressed as the log 2 value (L2FC) for display. All
experiments were performed on samples from at least two independent clutches,
with the number of biological replicates indicated for each experiment. Primer
information is provided in Table S13.
750
Left Lobe Liver Size Measurements
Liver size measurements were carried out based on the 2-dimensional area of the
left liver lobe in 120 hpf larvae. Embryos from incross of *Tg(fabp10a: nls-mcherry)* in
uhrf1$^{hi272+/-}$ and *Tg(fabp10a: CAAX-EGFP)* in *uhrf1*$^{hi272+/-}$ background was used for
all experiments for ease of liver detection. The live embryos were mounted in 3%
methylcellulose on the right side and the left liver lobe was imaged using Nikon
Elements on a Nikon SMZ25 Stereoscope. Liver size measurements were calculated
in ImageJ using free hand selection of the EFGP/ cherry labeled area corresponding
to the left liver lobe, and analyzed using the shape descriptor sub-function to
determine the area of the left liver lobe, expressed in μM^2. For Palbociclib treatment,
uhrf1$^{-/-}$ liver size of 20,000 μM^2 and below was considered small liver phenotype and
uhrf1$^{-/-}$ liver size above 20,000 μM^2 considered as rescued.
763
Immunofluorescence
120 hpf control siblings and *uhrf1*$^{-/-}$ zebrafish embryos containing the *Tg(fabp10a:*
nls-mcherry) transgene to identify hepatocyte nuclei were fixed in 4%
paraformaldehyde for 4 h at room temperature, washed in PBS, and treated with 150
mM Tris-HCl at pH 9.0 for 5 minutes, followed by heating at 70 °C for 15 minutes
according to an established protocol [93] and following incubation at room temperature
to cool the embryos and washed in PBS. Livers were dissected as described,
permeabilized with 10 µg/mL Proteinase K (Macherey-Nagel) in PBS containing
0.1% tween (PBST) for 10 minutes, washed 3 times with PBS, and incubated in a
blocking solution containing 5% Fetal Bovine Serum (GIBCO) in PBS for 60 minutes
at room temperature. The blocking solution was removed, and the livers were then
incubated in 100 µL of H3k9me3 (39161, Activ Motif), H3k9me2 (39753, Activ Motif),
Heterochromatin protein (Hp1) (MAB3866, Millipore), or Dnmt1 (SantaCruz) antibody
in blocking solution (1:200 dilution) overnight at 4 °C. After 3 washes in PBST, the
livers were incubated in secondary antibody (Molecular Probes) in blocking solution
(1:400 dilution) in the dark for 2 h on a shaker. After 5 serial washes with PBST, the
nuclei were counterstained with Hoechst (Thermo Fisher Scientific) diluted 1:1000 in
PBS, washed twice with PBS to remove excess Hoechst, and mounted on a
microscope slide with Vectashield (Vector Laboratories) and covered with a 0.1 mM
coverslip for imaging using Leica SP8 confocal microscope.
784
Terminal Deoxynucleotidyl Transferase dUTP Nick end Labeling Assay
(TUNEL)
Larvae collected at 5 dpf were fixed in 4% Paraformaldehyde for 4 h at room
temperature, and gradually dehydrated through a graded series of methanol and
stored in 100% methanol at 4 ∘C overnight. Gradual rehydration to PBS through a
graded series of methanol/PBS dilutions was carried out at room temperature.
Larvae were permeabilized with 10 µg/ml Proteinase K (Macherey-Nagel) in PBS
containing 0.1% tween (PBST) and fixed in 4% Paraformaldehyde for 10 minutes at
room temperature. Livers were then dissected out of the larvae and subjected to

TUNEL assay according to manufacturer's instructions (*In Situ* Cell Death Detection
kit, Fluorescein; Roche). Nuclei were counterstained with Hoechst (Thermo Fisher
Scientific) diluted 1:1000 in PBS, mounted on a microscope slide with Vectashield
(Vector Laboratories) and covered with a 0.1 mM coverslip for imaging using Leica
SP8 confocal microscope.
799

EdU Incorporation for S-phase Analysis
10 live larvae at 80 hpf, 96 hpf or 120 hpf were transferred into 50 ml falcon tube with
250 uL solution 1 mM EdU (ThermoFisher) in 10% DMSO in embryo water, and
incubated on ice for 20 minutes. The tubes were moved to room temperature, filled
with embryo water to 50 ml, and incubated for either 12 minutes or 30 minutes at
28.5 °C. After removing the embryo water, the embryos were fixed in 4%
Paraformaldehyde (Electron Microscopy Sciences) overnight at 4 °C, then
dehydrated through a graded series of methanol and incubated in 100% methanol
overnight at 4°C. The embryos were then gradually rehydrated back to 100% PBS
by serial dilutions of methanol. The embryos were then permeabilized with PBS
containing 1% Triton-X for 15 minutes prior to the Click-iT reaction to attach the
Alexa azide (Invitrogen). The larvae were transferred to a solution of 172 µl PBS, 8
µl CuSO4 (100 mM), and 0.2 µl Alexa azide in a 0.6 µl microcentrifuge tube, and
rocked in the dark for 10 minutes. 20 µl of ascorbic acid (0.5 M) was added after,
and was rocked in the dark for additional 20 minutes. Then the mixture was removed
and a fresh Alexa azide solution was added, and the same steps were repeated
once more. 80 hpf embryos were individually genotyped by collecting tail samples.
The larvae were washed 4 times in PBST prior to dissection and mounted in
Vectashield mounting medium with DAPI before they were imaged with the Leica
SP8 confocal microscope.
820

Confocal Imaging, Image Processing, and Analysis
Confocal imaging was performed using Leica TCS SP8 microscope with a 40x oil
immersion at a scan speed of 100 Hz. For antibody immunofluorescence stains and
TUNEL experiments, images were acquired at different focal distances from each
liver and LAS X software (Leica software) was used for quantification from 3
separate optical sections per liver, which were then averaged from 3 livers per clutch
per condition, and a minimum of 2 clutches were analyzed. For EdU and nuclear
morphology measurements, Z-stacks were acquired using the galvo stage, with 2
µm intervals. Bit depth was 12, and to enhance image quality, field of view and laser
intensity were adjusted separately for each sample. The acquired images were
visualized using LASX software. 3D analysis of Z-stacks of Hoechst-stained nuclei
was performed using the interactive 3D measurement tool in LASX for volume,
surface area, sphericity, elliptical mean, and nuclei counting with Hoechst and EdU
stains. Z-stacks were compiled into a 3D image, were adjusted for threshold and
noise to define the nuclei clearly, set for minimum size to be measured as 1000
voxels, and measurements of the previously mentioned parameters obtained and
exported in excel spreadsheets. For each time point and parameter measured,
around 100 nuclei were sampled from at least 3 livers per condition. Results were
plotted in GraphPad Prism 9.
840

RNA-Seq
50 livers were dissected at 120 hpf from Palbociclib or DMSO treated control sibling
or *uhrf1*[-/-]; *Tg(fabp10a: CAAX-EGFP)* larvae as previously described. The livers

were individually collected and stored at -80 °C while the carcasses were genotyped
for *uhrf1* mutation [23]. Once genotyped, the *uhrf1*[-/-] livers and control livers from each
individual clutch were pooled and RNA extracted according to established protocols
[42]. 100 ng of RNA was used for library preparation according to the manufacturer's
instructions (Illumina, San Diego, CA, USA). Libraries were sequenced on NovaSeq
(Illumina) to obtain 150 bp paired-end reads. Raw Fastq files quality was assessed
by using FASTQC and the reads were quality trimmed using Trimmomatic [94] to
remove low quality reads and adapters. Qualified reads were mapped to the
reference genome Danio rerio GRCz10. To estimate and compare gene expression
in different data sets, raw reads were uploaded to http://tsar.abudhabi.nyu.edu/ for
pre-processing and DeSeq2 for Differential gene expression analysis. Datasets of
RNAseq performed on 120 hpf *uhrf1*[-/-] and *dnmt1*[-/-] and their controls is previously
published [42] and available at GEO (GSE160728). Dataset of RNAseq performed on
120 hpf *uhrf1*[-/-] and sibling controls treated with Palbociclib or DMSO will be available
on GEO.
859
Bioinformatic Analysis
For Gene Ontology and Ingenuity Pathways Analysis (IPA) zebrafish gene names
were converted into human gene names using Biomart. Differentially expressed
genes were determined by p-value adjusted 0.05 and log 2 fold change 0. GO
enrichment analysis of differential expressed genes was conducted using REVIGO [95]
to find enriched terms for different DEG data sets. An adjusted p-value < 0.05 was
considered significant for all analyses. UPSET plots were generated to show
intersections of enriched terms from REVIGO, their size, and unique terms [96].
Heatmaps were generated with Pheatmap on z-score normalized counts. For plotting
and statistical analysis, R package 'ggplot' and GraphPad Prism 9 software were
used. The R codes used are available at https://github.com/SadlerEdepli-NYUAD.
871
Statistical Analysis
All experiments were carried out on multiple larvae from at least 2 clutches, as
indicated in each figure legend and data represented as the median with range. The
unpaired student *t* test was employed to compare the medians between groups with
adjustment for multiple comparisons and all of the data were considered to be
significant at *$p < 0.05$, **$p < 0.005$, ***$p < 0.0005$. All plots along with statistical
analysis were generated in GraphPad Prism 9 (GraphPad Software). Bioinformatic
analysis and visualization of genomic data were performed and plotted in RStudio.
880

Author Contributions: Conceptualization, K.C.S., B.M.; methodology, B.M., E.M.,
S.R.; formal analysis, B.M., E.M., S.R.; investigation, K.C.S., B.M., E.M., S.R.;
writing—original draft preparation, K.C.S. and B.M., writing—review and editing,
K.C.S., B.M., E.M.; visualization, K.C.S., B.M., E.M., S.R.; supervision, K.C.S.;
project administration, K.C.S.; funding acquisition, K.C.S. All authors have read and
agreed to the published version of the manuscript.

Funding: This work is supported by the NYUAD Faculty Research Fund (to K.C.S.),
REF (RE188, to KSE) and by Tamkeen under the NYU Abu Dhabi Research Institute
Award to the NYUAD Center for Genomics and Systems Biology (ADHPG-CGSB)
and the NYUAD Center for Genomics and Systems Biology.

Data Availability Statement: All the datasets used in this paper are available as
GEO datasets under the number GSE160728. Code used in this manuscript is
available on https://github.com/SadlerEdepli-NYUAD.

Acknowledgments: FACS Cell sorting was performed in the NYUAD CGSB FACS
Core at New York University Abu Dhabi. Bioinformatics assistance was provided by
the NYUAD Bioinformatics Core. We are grateful to Mehar Sultana, Marc Arnoux,
Nizar Drou and Muhammad Arshad for assistance with sequencing and analysis
Mehar Sultana for assistance with technical support for the FACS experiments, and
Rachid Rezgui for assistance with NYUAD Light Microscopy Core Technology
Platform. All members of the Sadler group provided insightful discussion and help
throughout the project, in particular Patrice Delaney bioinformatics and editing
assistance.
903
904 **Conflicts of Interest:** The authors declare no conflicts of interest.
905

References

1. Petryk, N., Kahli, M., d'Aubenton-Carafa, Y., Jaszczyszyn, Y., Shen, Y., Silvain, M., Thermes, C., Chen, C.L., and Hyrien, O. (2016). Replication landscape of the human genome. Nat Commun *7*, 10208. 10.1038/ncomms10208.

2. Recolin, B., van der Laan, S., Tsanov, N., and Maiorano, D. (2014). Molecular mechanisms of DNA replication checkpoint activation. Genes (Basel) *5*, 147-175. 10.3390/genes5010147.

3. McIntosh, D., and Blow, J.J. (2012). Dormant origins, the licensing checkpoint, and the response to replicative stresses. Cold Spring Harb Perspect Biol *4*. 10.1101/cshperspect.a012955.

4. Arita, K., Ariyoshi, M., Tochio, H., Nakamura, Y., and Shirakawa, M. (2008). Recognition of hemi-methylated DNA by the SRA protein UHRF1 by a base-flipping mechanism. Nature *455*, 818-821.

5. Avvakumov, G.V., Walker, J.R., Xue, S., Li, Y., Duan, S., Bronner, C., Arrowsmith, C.H., and Dhe-Paganon, S. (2008). Structural basis for recognition of hemi-methylated DNA by the SRA domain of human UHRF1. Nature *455*, 822-825.

6. Hashimoto, H., Horton, J.R., Zhang, X., Bostick, M., Jacobsen, S.E., and Cheng, X. (2008). The SRA domain of UHRF1 flips 5-methylcytosine out of the DNA helix. Nature *455*, 826-829.

7. Qian, C., Li, S., Jakoncic, J., Zeng, L., Walsh, M.J., and Zhou, M.M. (2008). Structure and hemimethylated CpG binding of the SRA domain from human UHRF1. J Biol Chem *283*, 34490-34494.

8. Bostick, M., Kim, J.K., Esteve, P.O., Clark, A., Pradhan, S., and Jacobsen, S.E. (2007). UHRF1 plays a role in maintaining DNA methylation in mammalian cells. Science *317*, 1760-1764.

9. Sharif, J., Muto, M., Takebayashi, S., Suetake, I., Iwamatsu, A., Endo, T.A., Shinga, J., Mizutani-Koseki, Y., Toyoda, T., Okamura, K., et al. (2007). The SRA protein Np95 mediates epigenetic inheritance by recruiting Dnmt1 to methylated DNA. Nature *450*, 908-912.

10. Mancini, M., Magnani, E., Macchi, F., and Bonapace, I.M. (2021). The multi-functionality of UHRF1: epigenome maintenance and preservation of genome integrity. Nucleic Acids Res *49*, 6053-6068. 10.1093/nar/gkab293.

11. Lopez-Contreras, Andres J., Ruppen, I., Nieto-Soler, M., Murga, M., Rodriguez-Acebes, S., Remeseiro, S., Rodrigo-Perez, S., Rojas, Ana M., Mendez, J., Muñoz, J., and Fernandez-Capetillo, O. (2013). A Proteomic Characterization of Factors Enriched at Nascent DNA Molecules. Cell Reports *3*, 1105-1116. https://doi.org/10.1016/j.celrep.2013.03.009.

12. Sirbu, B.M., McDonald, W.H., Dungrawala, H., Badu-Nkansah, A., Kavanaugh, G.M., Chen, Y., Tabb, D.L., and Cortez, D. (2013). Identification of Proteins at Active, Stalled, and Collapsed Replication Forks Using Isolation of Proteins on Nascent DNA (iPOND) Coupled with Mass Spectrometry*. Journal of Biological Chemistry *288*, 31458-31467. https://doi.org/10.1074/jbc.M113.511337.

13. Ferry, L., Fournier, A., Tsusaka, T., Adelmant, G., Shimazu, T., Matano, S., Kirsh, O., Amouroux, R., Dohmae, N., Suzuki, T., et al. (2017). Methylation of DNA Ligase 1 by G9a/GLP Recruits UHRF1 to Replicating DNA and Regulates DNA Methylation. Mol Cell *67*, 550-565 e555. 10.1016/j.molcel.2017.07.012.

14. Kim, J.K., Esteve, P.O., Jacobsen, S.E., and Pradhan, S. (2009). UHRF1
binds G9a and participates in p21 transcriptional regulation in mammalian
cells. Nucleic Acids Res *37*, 493-505. gkn961 [pii]10.1093/nar/gkn961.
15. Nishiyama, A., Yamaguchi, L., Sharif, J., Johmura, Y., Kawamura, T.,
Nakanishi, K., Shimamura, S., Arita, K., Kodama, T., Ishikawa, F., et al.
(2013). Uhrf1-dependent H3K23 ubiquitylation couples maintenance DNA
methylation and replication. Nature *502*, 249-253. nature12488
[pii]10.1038/nature12488.
16. Magri, L., Swiss, V.A., Jablonska, B., Lei, L., Pedre, X., Walsh, M., Zhang, W.,
Gallo, V., Canoll, P., and Casaccia, P. (2014). E2F1 coregulates cell cycle
genes and chromatin components during the transition of oligodendrocyte
progenitors from proliferation to differentiation. J Neurosci *34*, 1481-1493.
34/4/1481 [pii]10.1523/JNEUROSCI.2840-13.2014.
17. Unoki, M., Nishidate, T., and Nakamura, Y. (2004). ICBP90, an E2F-1 target,
recruits HDAC1 and binds to methyl-CpG through its SRA domain. Oncogene
23, 7601-7610. 10.1038/sj.onc.12080531208053 [pii].
18. Bonapace, I.M., Latella, L., Papait, R., Nicassio, F., Sacco, A., Muto, M.,
Crescenzi, M., and Di Fiore, P.P. (2002). Np95 is regulated by E1A during
mitotic reactivation of terminally differentiated cells and is essential for S
phase entry. J Cell Biol *157*, 909-914.
19. McCabe, M.T., Davis, J.N., and Day, M.L. (2005). Regulation of DNA
methyltransferase 1 by the pRb/E2F1 pathway. Cancer Res *65*, 3624-3632.
10.1158/0008-5472.CAN-04-2158.
20. Kimura, H., Nakamura, T., Ogawa, T., Tanaka, S., and Shiota, K. (2003).
Transcription of mouse DNA methyltransferase 1 (Dnmt1) is regulated by both
E2F-Rb-HDAC-dependent and -independent pathways. Nucleic Acids Res *31*,
3101-3113. 10.1093/nar/gkg406.
21. Ge, T.T., Yang, M., Chen, Z., Lou, G., and Gu, T. (2016). UHRF1 gene
silencing inhibits cell proliferation and promotes cell apoptosis in human
cervical squamous cell carcinoma CaSki cells. J Ovarian Res *9*, 42.
10.1186/s13048-016-0253-8.
22. Zhang, J., Yang, C., Wu, C., Cui, W., and Wang, L. (2020). DNA
Methyltransferases in Cancer: Biology, Paradox, Aberrations, and Targeted
Therapy. Cancers (Basel) *12*. 10.3390/cancers12082123.
23. Sadler, K.C., Krahn, K.N., Gaur, N.A., and Ukomadu, C. (2007). Liver growth
in the embryo and during liver regeneration in zebrafish requires the cell cycle
regulator, uhrf1. Proc Natl Acad Sci U S A *104*, 1570-1575.
10.1073/pnas.0610774104.
24. Anderson, R.M., Bosch, J.A., Goll, M.G., Hesselson, D., Dong, P.D., Shin, D.,
Chi, N.C., Shin, C.H., Schlegel, A., Halpern, M., and Stainier, D.Y. (2009).
Loss of Dnmt1 catalytic activity reveals multiple roles for DNA methylation
during pancreas development and regeneration. Dev Biol *334*, 213-223.
10.1016/j.ydbio.2009.07.017.
25. Jacob, V., Chernyavskaya, Y., Chen, X., Tan, P.S., Kent, B., Hoshida, Y., and
Sadler, K.C. (2015). DNA hypomethylation induces a DNA replication-
associated cell cycle arrest to block hepatic outgrowth in uhrf1 mutant
zebrafish embryos. Development *142*, 510-521. 10.1242/dev.115980.
26. Tittle, R.K., Sze, R., Ng, A., Nuckels, R.J., Swartz, M.E., Anderson, R.M.,
Bosch, J., Stainier, D.Y., Eberhart, J.K., and Gross, J.M. (2011). Uhrf1 and
Dnmt1 are required for development and maintenance of the zebrafish lens.

Dev Biol *350*, 50-63. S0012-1606(10)01196-6 [pii]10.1016/j.ydbio.2010.11.009.

27. Qin, Y., Wang, J., Gong, W., Zhang, M., Tang, Z., Zhang, J., and Quan, Z. (2014). UHRF1 depletion suppresses growth of gallbladder cancer cells through induction of apoptosis and cell cycle arrest. Oncol Rep *31*, 2635-2643. 10.3892/or.2014.3145.

28. Jenkins, Y., Markovtsov, V., Lang, W., Sharma, P., Pearsall, D., Warner, J., Franci, C., Huang, B., Huang, J., Yam, G.C., et al. (2005). Critical Role of the Ubiquitin Ligase Activity of UHRF1, a Nuclear RING Finger Protein, in Tumor Cell Growth. Mol Biol Cell *16*, 5621-5629.

29. Chen, T., Hevi, S., Gay, F., Tsujimoto, N., He, T., Zhang, B., Ueda, Y., and Li, E. (2007). Complete inactivation of DNMT1 leads to mitotic catastrophe in human cancer cells. Nat Genet *39*, 391-396. ng1982 [pii]10.1038/ng1982.

30. Blais, A., and Dynlacht, B.D. (2007). E2F-associated chromatin modifiers and cell cycle control. Curr Opin Cell Biol *19*, 658-662. 10.1016/j.ceb.2007.10.003.

31. Jung, J.K., Arora, P., Pagano, J.S., and Jang, K.L. (2007). Expression of DNA methyltransferase 1 is activated by hepatitis B virus X protein via a regulatory circuit involving the p16INK4a-cyclin D1-CDK 4/6-pRb-E2F1 pathway. Cancer Res *67*, 5771-5778. 10.1158/0008-5472.CAN-07-0529.

32. Du, Q., Smith, G.C., Luu, P.L., Ferguson, J.M., Armstrong, N.J., Caldon, C.E., Campbell, E.M., Nair, S.S., Zotenko, E., Gould, C.M., et al. (2021). DNA methylation is required to maintain both DNA replication timing precision and 3D genome organization integrity. Cell Reports *36*. ARTN 10972210.1016/j.celrep.2021.109722.

33. Wang, S., Miller, S.R., Ober, E.A., and Sadler, K.C. (2017). Making It New Again: Insight Into Liver Development, Regeneration, and Disease From Zebrafish Research. Curr Top Dev Biol *124*, 161-195. 10.1016/bs.ctdb.2016.11.012.

34. Goessling, W., and Stainier, D.Y. (2016). Endoderm specification and liver development. Methods Cell Biol *134*, 463-483. 10.1016/bs.mcb.2016.03.042.

35. Chu, J., and Sadler, K.C. (2009). New school in liver development: lessons from zebrafish. Hepatology *50*, 1656-1663. 10.1002/hep.23157.

36. Sladky, V.C., Knapp, K., Soratroi, C., Heppke, J., Eichin, F., Rocamora-Reverte, L., Szabo, T.G., Bongiovanni, L., Westendorp, B., Moreno, E., et al. (2020). E2F-Family Members Engage the PIDDosome to Limit Hepatocyte Ploidy in Liver Development and Regeneration. Dev Cell *52*, 335-349 e337. 10.1016/j.devcel.2019.12.016.

37. Delgado, I., Fresnedo, O., Iglesias, A., Rueda, Y., Syn, W.K., Zubiaga, A.M., and Ochoa, B. (2011). A role for transcription factor E2F2 in hepatocyte proliferation and timely liver regeneration. Am J Physiol Gastrointest Liver Physiol *301*, G20-31. 10.1152/ajpgi.00481.2010.

38. Wu, S.C., Kim, A., Gu, Y.J., Martinez, D.I., Zocchi, L., Chen, C.C., Lopez, J., Salcido, K., Singh, S., Wu, J., et al. (2022). UHRF1 overexpression promotes osteosarcoma metastasis through altered exosome production and AMPK/SEMA3E suppression. Oncogenesis *11*. ARTN 5110.1038/s41389-022-00430-6.

39. Mayhew, C.N., Carter, S.L., Fox, S.R., Sexton, C.R., Reed, C.A., Srinivasan, S.V., Liu, X., Wikenheiser-Brokamp, K., Boivin, G.P., Lee, J.S., et al. (2007). RB loss abrogates cell cycle control and genome integrity to promote liver tumorigenesis. Gastroenterology *133*, 976-984. 10.1053/j.gastro.2007.06.025.

40. Marjoram, L., Alvers, A., Deerhake, M.E., Bagwell, J., Mankiewicz, J.,
Cocchiaro, J.L., Beerman, R.W., Willer, J., Sumigray, K.D., Katsanis, N., et al.
(2015). Epigenetic control of intestinal barrier function and inflammation in
zebrafish. Proc Natl Acad Sci U S A *112*, 2770-2775.
10.1073/pnas.1424089112.
41. Madakashira, B.P., Zhang, C., Macchi, F., Magnani, E., and Sadler, K.C.
(2021). Nuclear Organization during Hepatogenesis in Zebrafish Requires
Uhrf1. Genes (Basel) *12*. 10.3390/genes12071081.
42. Magnani, E., Macchi, F., Madakashira, B.P., Zhang, C., Alaydaroos, F., and
Sadler, K.C. (2021). uhrf1 and dnmt1 Loss Induces an Immune Response in
Zebrafish Livers Due to Viral Mimicry by Transposable Elements. Front
Immunol *12*, 627926. 10.3389/fimmu.2021.627926.
43. Feng, S., Cokus, S.J., Zhang, X., Chen, P.Y., Bostick, M., Goll, M.G., Hetzel,
J., Jain, J., Strauss, S.H., Halpern, M.E., et al. (2010). Conservation and
divergence of methylation patterning in plants and animals. Proc Natl Acad
Sci U S A *107*, 8689-8694. 10.1073/pnas.1002720107.
44. Wang, S., Zhang, C., Hasson, D., Desai, A., SenBanerjee, S., Magnani, E.,
Ukomadu, C., Lujambio, A., Bernstein, E., and Sadler, K.C. (2019). Epigenetic
Compensation Promotes Liver Regeneration. Dev Cell *50*, 43-56 e46.
10.1016/j.devcel.2019.05.034.
45. Chernyavskaya, Y., Mudbhary, R., Zhang, C., Tokarz, D., Jacob, V.,
Gopinath, S., Sun, X., Wang, S., Magnani, E., Madakashira, B.P., et al.
(2017). Loss of DNA methylation in zebrafish embryos activates
retrotransposons to trigger antiviral signaling. Development *144*, 2925-2939.
10.1242/dev.147629.
46. Knouse, K.A., Wu, J., Whittaker, C.A., and Amon, A. (2014). Single cell
sequencing reveals low levels of aneuploidy across mammalian tissues. Proc
Natl Acad Sci U S A *111*, 13409-13414. 10.1073/pnas.1415287111.
47. Zhang, J., Bellani, M.A., Huang, J., James, R.C., Pokharel, D., Gichimu, J.,
Gali, H., Stewart, G., and Seidman, M.M. (2021). Replication of the
Mammalian Genome by Replisomes Specific for Euchromatin and
Heterochromatin. Front Cell Dev Biol *9*, 729265. 10.3389/fcell.2021.729265.
48. Lee, S.H., Hao, E., Levine, F., and Itkin-Ansari, P. (2011). Id3 upregulates
BrdU incorporation associated with a DNA damage response, not replication,
in human pancreatic beta-cells. Islets *3*, 358-366. 10.4161/isl.3.6.17923.
49. Rieck, S., Zhang, J., Li, Z., Liu, C., Naji, A., Takane, K.K., Fiaschi-Taesch,
N.M., Stewart, A.F., Kushner, J.A., and Kaestner, K.H. (2012).
Overexpression of hepatocyte nuclear factor-4alpha initiates cell cycle entry,
but is not sufficient to promote beta-cell expansion in human islets. Mol
Endocrinol *26*, 1590-1602. 10.1210/me.2012-1019.
50. Macaluso, M., Montanari, M., Noto, P.B., Gregorio, V., Bronner, C., and
Giordano, A. (2007). Epigenetic modulation of estrogen receptor-alpha by
pRb family proteins: a novel mechanism in breast cancer. Cancer Res *67*,
7731-7737. 67/16/7731 [pii]10.1158/0008-5472.CAN-07-1476.
51. Laderian, B., and Fojo, T. (2017). CDK4/6 Inhibition as a therapeutic strategy
in breast cancer: palbociclib, ribociclib, and abemaciclib. Semin Oncol *44*,
395-403. 10.1053/j.seminoncol.2018.03.006.
52. Bollard, J., Miguela, V., Ruiz de Galarreta, M., Venkatesh, A., Bian, C.B.,
Roberto, M.P., Tovar, V., Sia, D., Molina-Sanchez, P., Nguyen, C.B., et al.
(2017). Palbociclib (PD-0332991), a selective CDK4/6 inhibitor, restricts

tumour growth in preclinical models of hepatocellular carcinoma. Gut *66*,
1286-1296. 10.1136/gutjnl-2016-312268.
53. Pacek, M., Tutter, A.V., Kubota, Y., Takisawa, H., and Walter, J.C. (2006).
Localization of MCM2-7, Cdc45, and GINS to the site of DNA unwinding
during eukaryotic DNA replication. Mol Cell *21*, 581-587.
10.1016/j.molcel.2006.01.030.
54. Tanaka, S., Tak, Y.S., and Araki, H. (2007). The role of CDK in the initiation
step of DNA replication in eukaryotes. Cell Div *2*, 16. 10.1186/1747-1028-2-
16.
55. Schmitz, M.L., and Kracht, M. (2016). Cyclin-Dependent Kinases as
Coregulators of Inflammatory Gene Expression. Trends in Pharmacological
Sciences *37*, 101-113. https://doi.org/10.1016/j.tips.2015.10.004.
56. Ganz, J., Melancon, E., Wilson, C., Amores, A., Batzel, P., Strader, M.,
Braasch, I., Diba, P., Kuhlman, J.A., Postlethwait, J.H., and Eisen, J.S.
(2019). Epigenetic factors Dnmt1 and Uhrf1 coordinate intestinal
development. Dev Biol *455*, 473-484. 10.1016/j.ydbio.2019.08.002.
57. Milutinovic, S., Zhuang, Q.L., Niveleau, A., and Szyf, M. (2003). Epigenomic
stress response - Knockdown of DNA methyltransferase 1 triggers an intra-S-
phase arrest of DNA replication and induction of stress response genes.
Journal of Biological Chemistry *278*, 14985-14995. 10.1074/jbc.M213219200.
58. Unterberger, A., Andrews, S.D., Weaver, I.C.G., and Szyf, M. (2006). DNA
methyltransferase 1 knockdown activates a replication stress checkpoint.
Molecular and Cellular Biology *26*, 7575-7586. 10.1128/Mcb.01887-05.
59. Zupkovitz, G., Kabiljo, J., Kothmayer, M., Schlick, K., Schöfer, C., Lagger, S.,
and Pusch, O. (2021). Analysis of Methylation Dynamics Reveals a Tissue-
Specific, Age-Dependent Decline in 5-Methylcytosine Within the Genome of
the Vertebrate Aging Model Nothobranchius furzeri. Front Mol Biosci *8*,
627143. 10.3389/fmolb.2021.627143.
60. Acevedo, M., Vernier, M., Mignacca, L., Lessard, F., Huot, G., Moiseeva, O.,
Bourdeau, V., and Ferbeyre, G. (2016). A CDK4/6-Dependent Epigenetic
Mechanism Protects Cancer Cells from PML-induced Senescence. Cancer
Res *76*, 3252-3264. 10.1158/0008-5472.CAN-15-2347.
61. Goll, M.G., Anderson, R., Stainier, D.Y., Spradling, A.C., and Halpern, M.E.
(2009). Transcriptional silencing and reactivation in transgenic zebrafish.
Genetics *182*, 747-755. genetics.109.102079
[pii]10.1534/genetics.109.102079.
62. Branzei, D., and Foiani, M. (2005). The DNA damage response during DNA
replication. Current Opinion in Cell Biology *17*, 568-575.
10.1016/j.ceb.2005.09.003.
63. Lambert, S., and Carr, A.M. (2005). Checkpoint responses to replication fork
barriers. Biochimie *87*, 591-602. 10.1016/j.biochi.2004.10.020.
64. Courtot, L., Hoffmann, J.S., and Bergoglio, V. (2018). The Protective Role of
Dormant Origins in Response to Replicative Stress. International Journal of
Molecular Sciences *19*. ARTN 356910.3390/ijms19113569.
65. Reaper, P.M., Griffiths, M.R., Long, J.M., Charrier, J.D., MacCormick, S.,
Charlton, P.A., Golec, J.M.C., and Pollard, J.R. (2011). Selective killing of
ATM- or p53-deficient cancer cells through inhibition of ATR. Nat Chem Biol *7*,
428-430. 10.1038/Nchembio.573.
66. Arima, Y., Hirota, T., Bronner, C., Mousli, M., Fujiwara, T., Niwa, S., Ishikawa,
H., and Saya, H. (2004). Down-regulation of nuclear protein ICBP90 by

p53/p21Cip1/WAF1-dependent DNA-damage checkpoint signals contributes
to cell cycle arrest at G1/S transition. Genes Cells *9*, 131-142.

67. Merchut-Maya, J.M., Bartek, J., and Maya-Mendoza, A. (2019). Regulation of
replication fork speed: Mechanisms and impact on genomic stability. DNA
Repair (Amst) *81*, 102654. 10.1016/j.dnarep.2019.102654.

68. Maya-Mendoza, A., Moudry, P., Merchut-Maya, J.M., Lee, M., Strauss, R.,
and Bartek, J. (2018). High speed of fork progression induces DNA replication
stress and genomic instability. Nature *559*, 279-284. 10.1038/s41586-018-
0261-5.

69. Fu, H., Redon, C.E., Thakur, B.L., Utani, K., Sebastian, R., Jang, S.-M.,
Gross, J.M., Mosavarpour, S., Marks, A.B., Zhuang, S.Z., et al. (2021).
Dynamics of replication origin over-activation. Nature Communications *12*,
3448. 10.1038/s41467-021-23835-0.

70. Bohly, N., Schmidt, A.K., Zhang, X., Slusarenko, B.O., Hennecke, M.,
Kschischo, M., and Bastians, H. (2022). Increased replication origin firing links
replication stress to whole chromosomal instability in human cancer. Cell Rep
41, 111836. 10.1016/j.celrep.2022.111836.

71. Koundrioukoff, S., Carignon, S., Techer, H., Letessier, A., Brison, O., and
Debatisse, M. (2013). Stepwise Activation of the ATR Signaling Pathway upon
Increasing Replication Stress Impacts Fragile Site Integrity. Plos Genetics *9*.
ARTN e100364310.1371/journal.pgen.1003643.

72. Shechter, D., and Gautier, J. (2004). MCM proteins and checkpoint kinases
get together at the fork. P Natl Acad Sci USA *101*, 10845-10846. DOI
10.1073/pnas.0404143101.

73. Ge, X.Q., and Blow, J.J. (2010). Chk1 inhibits replication factory activation but
allows dormant origin firing in existing factories. Journal of Cell Biology *191*,
1285-1297. 10.1083/jcb.201007074.

74. Truong, L.N., and Wu, X. (2011). Prevention of DNA re-replication in
eukaryotic cells. J Mol Cell Biol *3*, 13-22. mjq052 [pii]10.1093/jmcb/mjq052.

75. Aggarwal, P., Vaites, L.P., Kim, J.K., Mellert, H., Gurung, B., Nakagawa, H.,
Herlyn, M., Hua, X., Rustgi, A.K., McMahon, S.B., and Diehl, J.A. (2010).
Nuclear cyclin D1/CDK4 kinase regulates CUL4 expression and triggers
neoplastic growth via activation of the PRMT5 methyltransferase. Cancer Cell
18, 329-340. 10.1016/j.ccr.2010.08.012.

76. Aggarwal, P., Lessie, M.D., Lin, D.I., Pontano, L., Gladden, A.B., Nuskey, B.,
Goradia, A., Wasik, M.A., Klein-Szanto, A.J., Rustgi, A.K., et al. (2007).
Nuclear accumulation of cyclin D1 during S phase inhibits Cul4-dependent
Cdt1 proteolysis and triggers p53-dependent DNA rereplication. Genes Dev
21, 2908-2922. 10.1101/gad.1586007.

77. Kim, Y., and Kipreos, E.T. (2007). Cdt1 degradation to prevent DNA re-
replication: conserved and non-conserved pathways. Cell Div *2*, 18.
10.1186/1747-1028-2-18.

78. Mudbhary, R., Hoshida, Y., Chernyavskaya, Y., Jacob, V., Villanueva, A., Fiel,
M.I., Chen, X., Kojima, K., Thung, S., Bronson, R.T., et al. (2014). UHRF1
overexpression drives DNA hypomethylation and hepatocellular carcinoma.
Cancer Cell *25*, 196-209. 10.1016/j.ccr.2014.01.003.

79. Du, Z., Song, J., Wang, Y., Zhao, Y., Guda, K., Yang, S., Kao, H.Y., Xu, Y.,
Willis, J., Markowitz, S.D., et al. (2010). DNMT1 stability is regulated by
proteins coordinating deubiquitination and acetylation-driven ubiquitination.
Sci Signal *3*, ra80. 3/146/ra80 [pii]10.1126/scisignal.2001462.

80.   Felle, M., Joppien, S., Nemeth, A., Diermeier, S., Thalhammer, V., Dobner, T.,
Kremmer, E., Kappler, R., and Langst, G. (2011). The USP7/Dnmt1 complex
stimulates the DNA methylation activity of Dnmt1 and regulates the stability of
UHRF1. Nucleic Acids Res *39*, 8355-8365. gkr528 [pii]10.1093/nar/gkr528.

81.   Qin, W., Leonhardt, H., and Spada, F. (2011). Usp7 and Uhrf1 control
ubiquitination and stability of the maintenance DNA methyltransferase Dnmt1.
J Cell Biochem *112*, 439-444. 10.1002/jcb.22998.

82.   Zhang, H., Gao, Q., Tan, S., You, J., Lyu, C., Zhang, Y., Han, M., Chen, Z., Li,
J., Wang, H., et al. (2019). SET8 prevents excessive DNA methylation by
methylation-mediated degradation of UHRF1 and DNMT1. Nucleic Acids Res
47, 9053-9068. 10.1093/nar/gkz626.

83.   Bourdeau, V., and Ferbeyre, G. (2016). CDK4-CDK6 inhibitors induce
autophagy-mediated degradation of DNMT1 and facilitate the senescence
antitumor response. Autophagy *12*, 1965-1966.
10.1080/15548627.2016.1214779.

84.   Buss, H., Handschick, K., Jurrmann, N., Pekkonen, P., Beuerlein, K., Muller,
H., Wait, R., Saklatvala, J., Ojala, P.M., Schmitz, M.L., et al. (2012). Cyclin-
dependent kinase 6 phosphorylates NF-kappaB P65 at serine 536 and
contributes to the regulation of inflammatory gene expression. PLoS One *7*,
e51847. 10.1371/journal.pone.0051847.

85.   D'Arcangelo, D., Tinaburri, L., and Dellambra, E. (2017). The Role of
p16(INK4a) Pathway in Human Epidermal Stem Cell Self-Renewal, Aging and
Cancer. Int J Mol Sci *18*. 10.3390/ijms18071591.

86.   Ashraf, W., Ibrahim, A., Alhosin, M., Zaayter, L., Ouararhni, K., Papin, C.,
Ahmad, T., Hamiche, A., Mely, Y., Bronner, C., and Mousli, M. (2017). The
epigenetic integrator UHRF1: on the road to become a universal biomarker for
cancer. Oncotarget *8*, 51946-51962. 10.18632/oncotarget.17393.

87.   Liang, D., Xue, H., Yu, Y., Lv, F., You, W., and Zhang, B. (2015). Elevated
expression of UHRF1 predicts unfavorable prognosis for patients with
hepatocellular carcinoma. Int J Clin Exp Pathol *8*, 9416-9421.

88.   Amsterdam, A., Nissen, R.M., Sun, Z., Swindell, E.C., Farrington, S., and
Hopkins, N. (2004). Identification of 315 genes essential for early zebrafish
development. Proc Natl Acad Sci U S A *101*, 12792-12797.

89.   Irion, U., Krauss, J., and Nusslein-Volhard, C. (2014). Precise and efficient
genome editing in zebrafish using the CRISPR/Cas9 system. Development
141, 4827-4830. 10.1242/dev.115584.

90.   Prevo, R., Fokas, E., Reaper, P.M., Charlton, P.A., Pollard, J.R., McKenna,
W.G., Muschel, R.J., and Brunner, T.B. (2012). The novel ATR inhibitor VE-
821 increases sensitivity of pancreatic cancer cells to radiation and
chemotherapy. Cancer Biol Ther *13*, 1072-1081. 10.4161/cbt.21093.

91.   Evason, K.J., Francisco, M.T., Juric, V., Balakrishnan, S., Lopez Pazmino
Mdel, P., Gordan, J.D., Kakar, S., Spitsbergen, J., Goga, A., and Stainier,
D.Y. (2015). Identification of Chemical Inhibitors of beta-Catenin-Driven Liver
Tumorigenesis in Zebrafish. PLoS Genet *11*, e1005305.
10.1371/journal.pgen.1005305.

92.   Schmittgen, T.D., and Livak, K.J. (2008). Analyzing real-time PCR data by the
comparative C(T) method. Nat Protoc *3*, 1101-1108. 10.1038/nprot.2008.73.

93.   Inoue, D., and Wittbrodt, J. (2011). One for all--a highly efficient and versatile
method for fluorescent immunostaining in fish embryos. PLoS One *6*, e19713.
10.1371/journal.pone.0019713.

94.   Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible
trimmer for Illumina sequence data. Bioinformatics *30*, 2114-2120.
10.1093/bioinformatics/btu170.
95.   Supek, F., Bosnjak, M., Skunca, N., and Smuc, T. (2011). REVIGO
summarizes and visualizes long lists of gene ontology terms. PLoS One *6*,
e21800. 10.1371/journal.pone.0021800.
96.   Conway, J.R., Lex, A., and Gehlenborg, N. (2017). UpSetR: an R package for
the visualization of intersecting sets and their properties. Bioinformatics *33*,
2938-2940. 10.1093/bioinformatics/btx364.

1265

Figure legends

Figure 1. DNA rereplication leads to increased DNA content and heterochromatin loss in *uhrf1* mutant hepatocytes

(A) FACS sorting of 5 dpf *Tg(fabp10a:nls:mcherry)* hepatocytes with control or *uhrf1^-/-* at 120 hpf. (B) Representative Z-stack projection confocal images of EdU and Hoechst stained control and *uhrf1^-/-* livers. Whole control and *uhrf1^-/-* embryos were pulsed with EdU for 30 minutes at 80 hpf, 96 hpf and 120 hpf and processed for EdU fluorescence by Click-it. (C) Quantification of EdU incorporation in B. (D) 80 hpf, 96 hpf and 120 hpf hepatocytes were sorted into early, mid and late S phase based on EdU pattern and quantified. (E) Representative immunofluorescent images of 120 hpf *uhrf1^-/-* and control livers with heterochromatin markers H3K9me3, H3K9me2 and Heterochromatin and their quantification. (F) Quantification of EdU incorporation from a 12 minute EdU pulse at 80 hpf and 120 hpf control and *uhrf1^-/-* hepatocytes and (G) sorted based on their S phase stage. C indicates control siblings and M indicates *uhrf1^-/-*. Scale bar = 50 µm, the number of samples and clutches indicated for each condition and represented as median with range. p-value *< 0.05, **< 0.005, *** < 0.0005 by unpaired Students t-test with adjustment for multiple comparisons.

Figure 2. CDK4/6 is activated in *uhrf1* mutant livers

(A) Gene ontology (REVIGO) of the significant upregulated and downregulated genes in 120 hpf *uhrf1^-/-* liver RNA-Seq ($p_{adj} < 0.05$) for each category. The cell cycle GO terms are highlighted in bold. (B) UPSET plot of all the cell cycle GO category genes from A. (C) Ingenuity Pathway Analysis (IPA) of all the cell cycle genes differentially expressed in 120 hpf *uhrf1^-/-* liver RNA-Seq, pathways involved in cell cycle checkpoint regulation in bold, z- score in IPA indicates a predicted activation or inhibition of a pathway. (D) Genes from the G1/S checkpoint pathway generated by IPA plotted with log2fold change, important cell cycle genes highlighted in red. (E) qPCR on key cell cycle genes of 96 hpf *uhrf1^-/-* and control livers. (F) qPCR on key cell cycle genes in the livers of P4 neonatal mice deficient for *Uhrf1* in hepatocytes (*Uhrf1^HEPKO*) and controls, inset shows band for Cre transgene by RT-PCR. The number of clutches is indicated for each condition and represented as median with range. p-value *< 0.05, **< 0.005, *** < 0.0005 by unpaired Students t-test with adjustment for multiple comparisons.

Figure 3. Cdk4/6 inhibition by Palbociclib (PD) rescues the *uhrf1* mutant liver phenotype

(A) Treatment scheme for inhibitor Palbociclib (PD). (B) Representative images of 120 hpf *uhrf1^-/-* and control embryos after DMSO or PD treatment, liver indicated in red. (C) Quantification of the left liver lobe area of 120 hpf PD treated and untreated livers. (D) Liver phenotype rescue in 120 hpf *uhrf1^-/-* treated with PD. (E) quantification of 30 minute EdU incorporation at 120 hpf in DMSO and PD treated controls and *uhrf1^-/-*. (F) Representative confocal image of 120 hpf Hoechst stained *uhrf1* control and *uhrf1^-/-* hepatocytes from DMSO or PD treatment and quantification of the nuclear surface area. (G) Representative immunofluorescent images of 120 hpf livers with heterochromatin marker H3K9me3 and quantification of H3K9me3 positive hepatocytes. (H) TUNEL assay was performed to detect cell death in 120 hpf in DMSO and PD treated controls and *uhrf1^-/-* and quantified. (I) Representative image of 5mC levels in *uhrf1^-/-* embryos and controls as measured by slot blot and quantification of the methylation levels from three clutches. Scale: 1000 µm in B, 50

1315   µm in F,G, the number of samples and clutches indicated for each condition and
1316   represented as median with range. p-value *< 0.05, **< 0.005, *** < 0.0005 by
1317   unpaired Students t-test with adjustment for multiple comparisons.
1318
Figure 4. Cdk4/6 inhibition by Palbociclib (PD) rescues abnormal cell cycle
gene expression in *uhrf1* mutants
(A) PCA analysis of RNA Seq from 120 hpf Palbociclib (PD) treated and DMSO
treated *uhrf1*[-/-] and control livers. (B) Venn diagram of differential expressed genes
(DEGs) (cutoff p_{adj} < 0.05) between *uhrf1*[-/-] and their control siblings treated with
DMSO and PD. (C) Unsupervised clustering heatmap of the expression profile of
120 hpf DMSO and PD treated *uhrf1*[-/-] liver RNA-Seq with z-scores based on rows.
Rows are divided into 10 clusters calculated on the hierarchical clustering of
dendrogram (Euclidean distance). (D) REVIGO gene ontology of cluster 4 and (E)
and cluster 5. (F) Key cell cycle genes, immune and *tnfa* response gene expression
between PD and DMSO treated *uhrf1*[-/-] plotted as a heatmap with log2fold change.
(G) IPA of the cell cycle genes differentially expressed between 120 hpf *uhrf1*[-/-] PD
and DMSO livers, cell cycle checkpoint regulation pathways indicated in bold. p-
value *< 0.05, **< 0.005, *** < 0.0005 by unpaired Students t-test with adjustment for
multiple comparisons and represented as median with range.
1334
Figure 5. Loss of *uhrf1* leads to Cdk4/6 mediated Dnmt1 protein stabilization
and smaller liver size
(A) Unsupervised clustering heatmap of all the cell cycle genes of 120 hpf *dnmt1*[-/-]
liver RNA-Seq with z-scores based on rows, important S-phase genes are indicated.
(B) EdU quantification (pulse 30 minute) of 96 hpf and 120 hpf control and *dnmt1*[-/-]
hepatocytes. (C) Representative images of 120 hpf *dnmt1*[-/-] and controls with
c269off+/-; 14X UAS:dsRed+/-; Tg(fabp10a:Gal4;cmlc2:GFP)+/- treated with DMSO
or Palbociclib (PD), showing the liver (dsRed), the heart (GFP) as a marker of
transgenesis and in green, the liver with hypomethylation in mutants only. (D)
Quantification of the left liver lobe area of *dnmt1*[-/-] and controls, treated with PD or
DMSO. Scale: 50 µm, the number of samples and clutches indicated for each
condition. (E) QPCR of key cell cycle genes and immune genes in *dnmt1*[-/-] livers.
Scale: 1000 µm in C, the number of samples and clutches indicated for each
condition. p-value *< 0.05, **< 0.005, *** < 0.0005 by unpaired Students t-test with
adjustment for multiple comparisons and represented as median with range.
1350
1351   **Figure 6. *dnmt1* mutation rescues DNA replication defect in *uhrf1* mutant**
1352   **hepatocytes**
1353
(A) Western blot of 80, 96 and 120 hpf *uhrf1*[-/-] and control siblings, levels of Dnmt1
are normalized on level of H3 for each sample and plotted. (B) Representative
immunofluorescent images with Dnmt1 staining of 120 hpf control and *uhrf1*[-/-] livers
treated with DMSO or PD and Dnmt1 positive cell quantification. (C) Representative
images of 120 hpf *uhrf1*[-/-] and control embryos uninjected or injected with *dnmt1*[cr].
(D) Quantification of the left liver lobe area of 120 hpf *uhrf1*[-/-] and controls uninjected
or injected with *dnmt1*[cr]. (E) QPCR of key cell cycle genes in 120 hpf livers of *uhrf1*[-/-]
and controls uninjected or injected with *dnmt1*[cr]. (F) Representative images of 30
minute EdU incorporation at 120 hpf in uninjected and *dnmt1*[cr] injected controls and
uhrf1[-/-] livers. Scale: 50 µm in B, 1000 µm in C, the number of samples and clutches
        indicated for each condition p-value *< 0.05, **< 0.005, *** < 0.0005 by unpaired

1364 Students t-test with adjustment for multiple comparisons and represented as median
1365 with range.
1366

Figure 7. Atr inhibition rescues liver size and DNA rereplication in *uhrf1* mutant
hepatocytes
(A) Representative images of 120 hpf $uhrf1^{-/-}$ and control larvae after DMSO or VE-
821 treatment, liver indicated in red. (B) Quantification of the left liver lobe area of
120 hpf VE- 821 treated and untreated embryos from $uhrf1^{-/-}$ and control larvae. (C)
Representative Z-stack projection of 120 hpf EdU and Hoechst stained control and
$uhrf1^{-/-}$ livers from DMSO or VE- 821 treated larvae. (D) Quantification of 30 minute
EdU incorporation at 120 hpf in DMSO and VE- 821 treated controls and $uhrf1^{-/-}$
livers. Scale: 50 μm in A, 50 μm in C, the number of samples and clutches indicated
for each condition p-value *< 0.05, **< 0.005, *** < 0.0005 by unpaired Students t-
test with adjustment for multiple comparisons. Data represented as median with
range. (E) Model of the relationship between *uhrf1* mutation, DNA methylation, cell
cycle regulation, DNA replication and small liver. Uhrf1 and Dnmt1 loss cause DNA
hypomethylation, upregulation of TEs and inflammatory response. Cdk4/6 activation
upon DNA hypomethylation causes Tnfa/b activation, leading to cell death in *uhrf1*
deficient livers. Cdk4/6 dependent activation of DNA replication genes leads to DNA
rereplication, replication stress and Atr activation. Both TE activation and replication
stress trigger DNA damage response to inhibit cell division, restricting organ
outgrowth and causing lethality in *uhrf1* mutants.
1386
1387

Supplemental Materials

Supplementary materials and methods

Survival assay
Heterozygous *uhrf1*[hi272+/-]; *Tg(fabp10a: nls-mcherry)* [+/-] adults were incrossed and split into 40 embryos per 90 mm petriplate, with around 25% *uhrf1*[-/-] per plate. The embryos were dechorionated and treated with either 0.5% DMSO or 20 μm Palbociclib at 48 hpf. At 120 hpf, the embryos were transferred to fresh embryo water and monitored till 9 dpf when the live embryos were individually collected and genotyped for the *uhrf1* mutation.

Eye size measurement
Eye size measurements were carried out based on the 2-dimensional area of the left eye in 120 hpf larvae. Live larvae were mounted in 3% methylcellulose on the right side and the left eye was imaged using Nikon Elements on a Nikon SMZ25 Stereoscope. Eye size measurements were calculated in ImageJ using free hand selection of the eye area and analyzed using the shape descriptor sub-function to determine the area expressed in μM^2.

Supplement Figure Legends
Figure S1. Representative images of 5 minute, 12 minute and 30 minute EdU incorporation in 120 hpf *uhrf1* control and *uhrf1*[-/-] livers.

Figure S2. Phenotype of Palbociclib (PD) treated *uhrf1* control and mutants
(A) 120 hpf embryos from incross of *uhrf1*[+/-] *;Tg(fabp10a: nls-mcherry)* adults treated with DMSO and imaged, livers are in red. (B) Phenotypic assessment of the embryos at 120 hpf. (C) 120 hpf embryos from incross of *uhrf1*[+/-] *;Tg(fabp10a: nls-mcherry)* adults treated with PD and imaged, livers in red, embryos numbered and genotype on the right side of the image. (D) PCR based genotyping to ascertain the *uhrf1*[-/-] with PD treatment. (E) Eye size measurement of 120 hpf *uhrf1*[-/-] and control embryos with DMSO and PD treatment. (F) Survival of DMSO and PD treated embryos at 9 dpf. Scale: 50μm, the number of samples and clutches indicated for each condition. p-value *< 0.05, **< 0.005, *** < 0.0005 by two tailed Students t-test with adjustment for multiple comparisons.

Figure S3. Palbociclib (PD) treatment differentially regulates cell cycle gene expression but does not rescue transposon expression in *uhrf1* mutants
(A) Volcano plot of Differently Expressed Genes (DEGs) from RNA-Seq between 120 hpf *uhrf1* control PD treated and control DMSO treated livers. (B) IPA showing the pathways differentially expressed from RNA-Seq between 120 hpf *uhrf1* control PD treated and control DMSO treated livers. (C) Four quadrant cross-plot of the DEGs with log2fold change from RNA-Seq between 120 hpf *uhrf1*[-/-] PD treated and DMSO treated livers. (D) Correlation plot of repetitive elements in 120 hpf *uhrf1*[−/−] mutant livers treated with DMSO and PD. Upregulated TEs have log 2 fold change > 0; downregulated TEs have log 2 fold change < 0. Log 2 fold change is calculated between mutants and their own control siblings. (E) Gene ontology (REVIGO) of the upregulated and downregulated genes in 120 hpf DMSO and PD treated *uhrf1*[-/-] liver RNA-Seq using significant genes (p_{adj} < 0.05) for each category.

1438
Figure S4. Palbociclib (PD) treatment on *uhrf1* mutants rescues key cell cycle gene
overexpression
(A) IPA analysis of the G1-S pathway in DMSO *uhrf1*$^{-/-}$ compared to DMSO treated
controls. The color of the circles represents the observed results (red for activated/
upregulated and blue for repressed/ downregulated). (B) qPCR of key cell cycle
genes of 120 hpf *uhrf1*$^{-/-}$ and control livers treated with DMSO or PD, data from 2
clutches shown. (C) Volcano plot of Differently Expressed Genes (DEGs) from RNA-
Seq between 120 hpf *uhrf1*$^{-/-}$ PD treated and *uhrf1*$^{-/-}$ DMSO treated livers. (D) IPA
analysis of the G1-S pathway between PD and DMSO treated *uhrf1*$^{-/-}$. The color of
the circles represents the observed result (red for activated/ upregulated and blue for
repressed/ downregulated).
1450
1451 **Figure S5.** IPA analysis of ATM pathway
1452 IPA analysis of the ATM pathway in 120 *uhrf1*$^{-/-}$ liver and 120 *dnmt1*$^{-/-}$ liver
1453 transcriptomes. The color of the circles represents the observed results (orange for
1454 activated/ upregulated only in *uhrf1*$^{-/-}$, red for activated in both, light blue for
1455 repressed/ downregulated only in *uhrf1*$^{-/-}$ and dark blue for repressed in both.
1456
Figure S6. Testing the efficiency of *dnmt1*^{cr} *in c269off+/-; 14X UAS:dsRed+/-;*
Tg(fabp10a:Gal4;cmlc2:GFP)+/- and nuclear morphology of *dnmt1*^{cr} injected *uhrf1*
control and mutants
(A) Representative images of 120 hpf uninjected and *dnmt1*^{cr} embryos imaged for
dsRed and GFP in the head. GFP positivity indicates hypomethylation. (B) 120 hpf
embryos from incross of *uhrf1*$^{+/-}$;*Tg(fabp10a: nls-mcherry)* adults injected or
uninjected with *dnmt1*^{cr} were genotyped, PFA fixed, livers dissected, Hoechst
stained and imaged as a Z-stack which were then used to calculate (B) volume, (C)
Surface area and (D) Ellipsoid mean of the hepatocytes. The number of samples and
clutch indicated for each condition. p-value *< 0.05, **< 0.005, *** < 0.0005 by
ANOVA.
1468
Supplemental tables
Table S1. REVIGO analysis of RNA-Seq from 120 hpf *uhrf1* mutant livers
Table S2. Cell cycle pathways identified by IPA of RNA-Seq from 120 hpf *uhrf1*
mutant livers
Table S3. Optimization of PD exposure protocol on WT embryos
Table S4. DEGs in the liver of PD and DMSO treatment of 120 hpf control larvae
Table S5. IPA of DEGs from the liver of PD and DMSO treated 120 hpf control larvae
Table S6. DEGs from the liver of DMSO treated control and *uhrf1* mutant larvae and
from PD treated control and *uhrf1* mutants at 120 hpf
Table S7. ENSEMBL IDs of the heatmap clusters from DMSO and PD treated
control and *uhrf1* mutant larvae
Table S8. REVIGO analysis of clusters 4 and 5 from DMSO and PD treated control
and *uhrf1* mutant larvae
Table S9. REVIGO analysis of DEGs between *uhrf1* mutants treated with DMSO or
PD (from Table S6)
Table S10. DNA replication DEGs, immune DEGs and TNFa DEGs between *uhrf1*
mutants treated with DMSO or PD
Table S11. DEGs from the liver of DMSO vs. PD treated *uhrf1* mutants at 120 hpf
Table S12. IPA of DEGs from DMSO vs. PD treated *uhrf1* mutants at 120 hpf

Table S13. Primers for qPCR and CRISPR sgRNA

A.

UP DOWN

Term	UP	DOWN
mitotic cell cycle process	133	
DNA replication	68	
nuclear division	83	
organelle fission	85	
cell cycle process regulation	82	
cell division	97	
response to biotic stimulus	100	
response to other organism	99	
regeneration	59	
neg.regulation of chr. organization	15	
chromosome condensation	18	
tissue regeneration	43	
regulation of immune system process	79	
regulation of DNA metabolic process	36	
protein−DNA complex assembly	33	
response to cytokine	51	
immune effector process	38	
chromosome localization	12	
intrinsic apoptotic signaling pathway	23	
carboxylic acid catabolism		87
precursor metabolites generation		101
oxidation of organic compounds		76
purine metabolism		103
lipid oxidation		32
mitochondrion organization		76
hydroxy compound metabolism		76
sulfur compound metabolism		70
cytoplasmic translation		31
lipid homeostasis		27
modified amino acid metabolism		42
carboydrate metabolic process		93
ketone metabolism		28
intracellular protein transmembrane		20
ribosome assembly		17
macromolecule biosynthesis regulation		50
vitamin metabolic process		17
aldehyde metabolic process		17
small molecule metabolism regulation		25
autophagy of nucleus		9

Adjusted p−value
2e−04
4e−04
6e−04

Ratio of Genes in Gene List
0.02
0.04

B.

45 32 19 15 13 12 12 10 9 8 8 8 7 7 6 5 5 4 3 3 3 3 3 1 1 1 1 1 1 1 1 1

100 50 0

negative regulation of chromosome organization
chromosome condensation
DNA replication
cell cycle process regulation
nuclear division
cell division
mitotic cell cycle process

C.

G2/M DNA Damage Checkpoint Regulation
G1/S Checkpoint Regulation
GADD45 Signaling
CDK5 Signaling
DNA damage-induced 14-3-3 σ Signaling
ATM Signaling
Cyclins and Cell Cycle Regulation
Integrin Signaling
Mitotic Roles of Polo-Like Kinase
Cell Cycle Regulation by BTG Family Proteins
Cell Cycle Control of Chromosomal Replication

-2   0   2   4
z-score

D.

Log 2 Fold change ($uhrf1^{-/-}$/ Control)

rbl2, cdkn1b, rpl11, gsk3b, hdac10, mdm2, cul1, max, atr, ccnd1, cdk4, alb1, cdk2, rb1, tgfb2, skp2, tfdp1, tgfb1, ccne1, cdk6, cdc25a, suv39h1, tp53, bmi1, e2f5, nrg1, cdkn2c

E. Zebrafish liver; 96 hpf

● Control
● $uhrf1^{-/-}$

Log 2 Fold change ($uhrf1^{-/-}$/ Control)

ccne1, ccna2, mdm2, mcm3, orc3, cdc6, pola1, pold1, pcna

F. Mouse liver; P4

Cre+ Cre-

● Control (Alb;Cre$^{+/+}$)
● $Uhrf1^{HepKO}$ (Alb;Cre$^{Tg/+}$;Uhrf1$^{fl/fl}$)

Log 2 Fold change ($Uhrf1^{HepKO}$/ Control)

Ccna2, Mcm3, Orc1, Cdc6, Pcna, Uhrf1

*

Figure 2

Figure 2

Figure 3

Figure 4

A.

Control_1 Control_2 Control_3 dnmt1⁻/⁻_1 dnmt1⁻/⁻_2 dnmt1⁻/⁻_3

cdc45
mcm2
orc6
pold1
mcm5
cdc6
orc5
pola1
orc 1
mdm2
pcna

atr

pold2

Z-score

B.

% EdU/ Hoechst

	Control	dnmt1⁻/⁻	Control	dnmt1⁻/⁻
	96 hpf		120 hpf	
No: of cells =	240/800	30/500	98/750	14/700
livers =	6	6	9	9
clutch=	2	2	3	3

C.

120 hpf

c269off⁺/⁻;14X UAS:dsRed⁺/⁻;Tg(fabp10a:Gal4;cmlc2:GFP)⁺/⁻

DMSO | PD

Control
ds:Red — liver
GFP — heart

dnmt1⁻/⁻
ds:Red — liver
GFP — liver, heart

D.

Left liver area (µm²)

	DMSO	PD	DMSO	PD
	Control		dnmt1⁻/⁻	
livers =	12	16	13	20
clutch =	2	2	2	2

E.

● DMSO ● Palbociclib

Log 2 Fold change (dnmt1⁻/⁻/ Control)

ccne1 ccna2 mdm2 mcm3 orc3 cdc6 pola1 pold1 pcna isg15 nfkb2

Figure 5

A.

WT sibling   *uhrf1⁻/⁻*

hpf 80 96 120   80 96 120

Dnmt1   ─250 KDa
       ─130 KDa

H3   ─15 KDa

● Control   ● *uhrf1⁻/⁻*

Dnmt1 (normalized to H3)

80 hpf   96 hpf   120 hpf

B.

120 hpf; hepatocytes

Control	*uhrf1⁻/⁻*	
DMSO	PD	PD

Hoechst/Dnmt1

Dnmt1

% Dnmt1 positive nuclei

DMSO   PD   DMSO   PD
Control   *uhrf1⁻/⁻*

Dnmt1/ Hoechst =	5/200	5/200	152/250	38/200
livers =	3	4	4	4
clutch =	2	2	2	2

C.

120 hpf; *Tg(fabp10a:nls-mCherry)*

	Uninjected	*dnmt1ᶜʳ*
Control		
uhrf1⁻/⁻		

D.

Left liver area (µm²)

Uninj   *dnmt1ᶜʳ*   Uninj   *dnmt1ᶜʳ*
Control   *uhrf1⁻/⁻*

| livers = | 12 | 16 | 13 | 20 |
| clutch = | 2 | 2 | 2 | 2 |

E.

● Uninjected   ● *dnmt1ᶜʳ*

Log 2 Fold change (*uhrf1⁻/⁻*/Control)

***

ccna2 mcm3 mdm2 orc3 cdc6 pcna pola1 pold1 dnmt1

F.

% of cells with EdU incorporation

ns   **** **   ****

Uninj   *dnmt1ᶜʳ*   Uninj   *dnmt1*
Control   *uhrf1⁻/⁻*

No: of cells =	80/4000	65/3750	690/1450	120/1150
livers =	8	7	7	7
clutch =	2	2	2	2

Figure 6

A.

120 hpf; *Tg(fabp10a:nls-mCherry)*

DMSO | VE-821

Control

uhrf1⁻/⁻

B.

Left liver area (μm²)

Control		*uhrf1⁻/⁻*	
DMSO	VE-821	DMSO	VE-821
livers = 26	29	20	19
clutch = 3	3	3	3

C.

120 hpf; hepatocytes

DMSO | VE-821

EdU/ Hoechst

Control

uhrf1⁻/⁻

D.

% EdU incorporation

Control		*uhrf1⁻/⁻*	
DMSO	VE-821	DMSO	VE-821
No: of cells = 187/2144	162/2022	868/2500	334/1441
livers = 6	6	10	7
clutch = 3	3	3	3

E.

↓Uhrf1 ↓Dnmt1

Palbociclib ⊣ ↓DNA Methylation

↑Cdk4/6

↑Tnfa/b → Cell death

E2F

E2F site ON → DNA replication genes

dnmt1^cr ⊣ Re replication

Dnmt1 Pcna Mcms Orcs

↑TE → Interferon response

DNA Damage Response

Replication Stress

↑ATR

VE-821 ⊣

Dormant Origin Firing

Cell division

Small liver

Death

Figure 7