CRISPR/Cas9 DNA synthesis disruption as a tool to control cell proliferation in vitro and in vivo

Rocio Sartori Maldonado[1], Hossam Montaser[1], Inkeri Soppa[1], Melanie Balaz[1], Henri

Puttonen[4], Solja Eurola[1], Timo Otonkoski[1,2], Jonna Saarimäki-Vire[1], Kirmo Wartiovaara[1,3].

Affiliations:

1- Stem Cells and Metabolism Research Program, Faculty of Medicine, University of Helsinki,

Helsinki, Finland

2- Children's Hospital, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

3- Clinical Genetics, Helsinki University Hospital, Helsinki, Finland

4- Pathology, Helsinki University Hospital, Helsinki, Finland

Abstract

Cell therapies hold a lot of promise for regenerative medicine and gene therapy. However,

many living cell products entail a fundamental biological risk of unwanted growth. Here, we

describe a novel metabolic safety system to control cell proliferation without exogenous genetic

elements. The method allows robust cell culture and manufacture while mitigating the risk of

uncontrolled growth of transplanted cells.

Using CRISPR/Cas9, we inactivated the enzyme in charge of the *de novo* thymidylate

synthesis (*TYMS*) in several cell lines, including human induced pluripotent stem cells. This

resulted in cells that proliferate when supplemented with exogenous dTMP but fail to grow in

its absence. Under dTMP supplementation, $TYMS^{-/-}$ hiPSCs produce mature teratomas *in vivo*

and differentiate normally into potentially therapeutic cell types, such us pancreatic beta cells.

After terminal differentiation, the cells no longer require dTMP to remain functional, as seen by

prolonged *in vivo* production of human insulin in mice.

Introduction

Somatic cell therapies offer new possibilities for previously untreatable or challenging medical conditions. Among the cell types used in such therapies, induced pluripotent stem cells (hiPSCs) have garnered significant attention due to their availability from patient-specific somatic cells[1,2] and their capability to differentiate into potentially therapeutic cell lineages, such as glucose-responsive insulin secreting pancreatic islets[3]. Furthermore, the improved technologies for gene transfer and genome editing enable the development of genetically modified cell-based therapies with enhanced therapeutic potential[4–6].

The use of hiPSC offers several advantages over mesenchymal and embryonic stem cells (MSC and ESC, respectively). hiPSCs circumvent ethical concerns associated with the embryo derived ESCs[7,8]. Unlike donor-derived MSCs and ESCs, hiPSCs' patient-specificity minimizes the risk of immune rejection and graft-versus-host disease[9,10]. Moreover, they can be further genetically modified by silencing or modifying genes involved in, for instance, immune recognition[11]. These features broaden their applications in disease modeling and therapy development, and may enhance their compatibility for allogeneic cell therapies, reducing the need for immunosuppression compared to other cell sources. However, although the universal off-the-shelf cell products decrease the risk for treatment rejection, this diminished immunogenicity can also lead to problematic immune evasion in potential oncogenic events.

On top of creating a biologically useful cells for a desired clinical purpose, a successful commercial therapy product development needs to meet the criteria for safety and manufacture scalability. These two goals, however, often compromise each other, since manufacturing benefits from robust cell growth, but intensive or unlimited proliferation poses a threat when the cells are transferred to patients[12,13].

Attempts to create safer cells for therapy have yielded different strategies[14,15]. These safety systems aim to prevent uncontrolled proliferation or eliminate the transplanted cells if necessary[16,17]. Some of the existing safety strategies include the use of inducible apoptotic systems, suicide genes, antibody-mediated cell depletion, and gene editing-based switches (Table 1). Often, these switches work by reversibly activating or inactivating genes upon exposure to a small molecule (ON- and OFF-switches, respectively). However, most of them are transgenic, could be silenced or genetically altered, take long time to work, derive from viruses, cause neurotoxicity and/or present rapid remission[15,18].

In this context, we envisioned the generation of an *endogenous* safety strategy, by

disrupting a key metabolic gene, whose function can be compensated with a dispensable

compound. Using CRISPR/Cas9, we genetically inactivated thymidylate synthase (*TYMS*), the

only known enzyme in charge of the *de novo* thymidylate (dTMP) synthesis[19]. This knock-out

generates rescuable auxothrophy towards thymidine and restricts the synthesis of DNA,

without altering normal RNA or protein production (Fig. 1a). This means that dividing cells

depend on external supplementation of thymidine to proliferate and achieve robust

manufacturing of the cells of interest. However, once the cells have exited the cell cycle and

terminally differentiated, they do not require supplementation to synthesize nucleic acids.

While preventing dividing cells from proliferating uncontrollably, this approach does not

require the use of small molecules nor the insertion of large transgenic elements, hence it

cannot be silenced; and it does not result leaky or immunogenic.

Our results provide evidence that the culture and proliferation of *TYMS*[-/-] cells can be

regulated with exogenous dTMP *in vitro* and *in vivo*. Furthermore, the characterization and

differentiation of *TYMS*[-/-] hiPSC show that they provide fully functional source for cell therapy

but are unable to sustain unwanted proliferation.

Results

Generation of thymidine-dependent cell proliferation with genome editing.

The human thymidylate synthase (PDB: 5X5D) is evolutionarily highly conserved. The

functional enzymatic ligand pocket is encoded by exons 4 (amino acid R175), 5 (P193, C195,

Q214, S216, D218, R225) and 6 (H256, Y258)[20]. Therefore, to interrupt TYMS activity, we deleted

the exon 5 by targeting both intron 4 and intron 5. (Fig. 1.b, Supplementary Fig. 1a). This

depleted the enzyme expression at both mRNA and protein levels (Fig. 1.C, Supplementary Fig.

1a-b) in several cell lines.

We hypothesized that *TYMS*[-/-] cells could be cultured and manufactured normally with

external dTMP supplementation, but they would cease to proliferate in its absence. We

followed the growth of *TYMS*[-/-]-iPS (Hel24.3[21]) and immortal cell lines without or with different

concentrations of dTMP (from 0.3 μM to 40 μM). The untreated (wild-type, WT) cell lines were

used as controls. The results suggest that *TYMS*[-/-] cells proliferate at a rate comparable to WT

when supplemented with thymidine but undergo apoptosis early upon dTMP withdrawal (Fig.

1d). Moreover, the *TYMS*[-/-] -hiPSCs used from here onwards could not sustain growth at

physiological concentrations of thymidine (Fig. 1e, 1 µM; reference: 0.5-1.4 µM[22]). Contrarily,

WT *TYMS*[+/+] lines maintain equal growth rate in all tested dTMP concentrations (Supplementary

Fig. 1d). Cell cycle analysis by flow cytometry of Hel24.3 *TYMS*[-/-] cells reveal that in the first 24h

of dTMP withdrawal, the cells accumulate in G0-G1 and S-phase, and then, as they should

continue to G2, accumulate in the subG0-G1 phase (Supplementary Fig. 1e). In accordance with

the reports by Diehl et al[23], this suggests that this nucleotide starvation causes a replicative

stress sensed in the S-G2 checkpoint.

Knock-out hiPSCs pluripotency and genomic integrity.

The *TYMS*[-/-]-iPS cell lines showed no detrimental effects on the identity or functions of the

stem cells, tested by standard pluripotency analysis. The undifferentiated *TYMS*[-/-]-Hel24.3 lines

showed positive staining for pluripotency markers Nanog, OCT4, Sox2 and SSEA4 and

comparable mRNA expression of Sox2, Nanog and OCT4 to wild-type Hel24.3 (Fig. 2a-b). Their

chromosomal integrity remained unaltered by the editing process in prolonged culture, tested

in several time points for up to 25 passages (Fig. 2c and Supplementary Fig. 1e). The

differentiation into the three germ layers with or without dTMP supplementation revealed that

the cells successfully differentiate into ectoderm, mesoderm and endoderm when

supplemented with dTMP at the early stages of differentiation (Fig. 2d). Supplementation

withdrawal during the first stage produced significantly less embryoid bodies at d14 (Fig. 1e).

By day 28, however, all attached cells expressed characteristic markers of the three germ layers

(Fig 2.f, Supplementary Fig. 1f). The successful survival in 3D aggregates and differentiation of

non-supplemented cells may suggest the recycling of thymidine from apoptotic cells until cell

cycle exit.

TYMS[-/-]*-hiPSCs produce teratomas when supplemented in vivo.*

To test the behavior of our Hel24.3 *in vivo*, we implanted undifferentiated *TYMS*[+/+] and *TYMS*[-/-]

*-hiPSC aggregates subcutaneously in mice, with 100 mg/kg/day or without dTMP

supplementation in the drinking water (Fig. 3a). After 6 weeks, we analyzed the size,

composition, and proliferation of the resulting teratomas (Fig 3b-f).

TYMS[+/+]-hiPSC produced teratoma-like growth regardless of the thymidine supplementation,

while *TYMS*[-/-]-hiPSC produced visible teratomas only under dTMP supplementation (Fig 3B).

Without supplementation, only one of the mice presented a small tumor. Histological analysis

and immunocytochemistry showed that *TYMS*[+/+]-hiPSC generated dense teratomas with high

cellularity (Fig. 3c-d). Contrarily, *TYMS*[-/-]-hiPSC overall produced smaller teratomas. Under the

chosen concentration of dTMP, the teratomas were bigger and, although they presented overall

lower cellularity than their WT counterparts, we could identify denser areas with differentiated

structures, consisting mostly of neural tubules, and epithelial cells. Absence of thymidine

supplementation resulted in very cystic tumors with low cellularity, significantly less

proliferative cells (Fig. 3.f), and composed mainly of primitive glial, epithelial, and mesenchymal

tissue.

Non-supplemented knock-out beta cells produce insulin in vitro.

One potential and intensively studied cell-therapy target is diabetes, in which stem cell-

derived insulin-producing beta cells act as the therapeutic source of insulin. We evaluated the

safety, quality, and function of *TYMS*[-/-]-hiPSC-derived pancreatic islets using well-defined, multi-

step protocol for beta cell differentiation (Fig. 4a). We differentiated *TYMS*[-/-] hiPSCs (Hel24.3)

in vitro, providing thymidine supplementation (40 µM) until maturation. Non-supplemented

wild-type Hel24.3 were used as controls, while non-supplemented *TYMS*[-/-] cells died by day 2,

making further characterization of this condition impossible.

We evaluated the progression of the differentiation protocol by flow cytometry, qPCR, and

immunostaining of characteristic markers of several maturation stages. Wild-type and *TYMS*[-/-]

Hel24.3 cells showed comparable *in vitro* expression of markers at the definitive endoderm

stage (CXCR4), S4 (PDX1, NKX6.1) and S7 (insulin – INS, glucagon – GCG, NKX6.1, PDX1, C-

peptide) by flow cytometry (Fig. 4b and Supplementary Fig. 2a-b). In transcriptomic analysis by

qPCR, we found that both cell lines expressed characteristic markers at stages DE and S4, and

markers of mature beta-cells at week 4 of stage 7 (S7W4) (Fig. 4c). Moreover, at stage 7, the

TYMS[-/-]- derived islets presented significantly lower *TYMS* mRNA levels. By the last stage before

implantation (S7W4), our *TYMS*[-/-]-Hel24.3 islets showed normal morphology and comparable

percentage of insulin and glucagon expressing cells (Fig. 4D and Supplementary Fig 2C).

Accordingly, $TYMS^{-/-}$ and $TYMS^{+/+}$ Hel24.3 showed similar *in vitro* levels of glucose and KCl

induced insulin secretion (Fig. 4e-f).

We did not observe differences between $TYMS^{+/+}$-hiPSC differentiated with or without dTMP

supplementation, suggesting that the addition of dTMP at our working concentrations (up to

40 µM) does not affect $TYMS^{+/+}$-cells.

In vivo function of non-supplemented $TYMS^{-/-}$-hiPSC derived differentiated cells.

We next evaluated the function of mature $TYMS^{-/-}$ beta cells (S7) *in vivo* by engrafting them

under the kidney capsule in immunosuppressed mice without dTMP supplementation. We

implanted seven mice with $TYMS^{+/+}$-Hel24.3-derived control islets and six with $TYMS^{-/-}$-Hel24.3-

derived islets, and monitored their body weight, human C-peptide production, and glucose

levels for three months (Fig. 5a-c).

Both $TYMS^{-/-}$ and control islet successfully mature in the first month, produce comparable

levels of human C-peptide, and lower the mouse blood glucose closer to human physiological

values (Fig. 5a, b). All mice maintained their normal body weights (Fig 5b-c) and showed no

signs of other health concerns.

Immunohistochemistry of the $TYMS^{-/-}$-implanted kidneys showed vascularization in the graft

and high percentage of insulin producing cells (Fig. 5d).

Additionally, to monitor the proliferation of the cells in the grafts *in vivo* through time (1 and

3 months), we supplemented the drinking water of the mice with EdU for 5 days before

sacrificing them. Interestingly, while we see reduction on active proliferation in knock-out cells,

the uptake of EdU does not seem significantly different. (Fig 5e-f and Supplementary Fig. 3d).

This might point to the uptake of EdU as a source of thymidine.

Discussion

Here, we have developed a method to generate controllable cells without inserting any

external genetic elements. By disrupting the only known *de novo* DNA synthesis reaction, we

have obtained cells auxotrophic to thymidine. This approach worked in several cell lines and

successfully generated cells that require the external administration of thymidine to

proliferate. With dTMP supplementation, the cells grow exponentially for undefined periods

in culture showing no signs of exhaustion, rendering them useful in commercial biological

product development, for example.

The *TYMS^-/-* hiPSCs retain their pluripotency and differentiate successfully into different

cell types. Non-supplemented *TYMS^-/-*-hiPSC generate small or no teratomas *in vivo* with a

significantly lower proliferative index (KI67), pointing to the loss of proliferative capacity of

these cells *in vivo* under physiological concentrations of thymidine.

Importantly, TYMS activity is required for DNA, but not RNA, synthesis pathways. Hence,

we see no functional alterations in the terminally differentiated cells without dTMP

supplementations *in vitro* or *in vivo*. The islets survived implantation and proved to

successfully secrete human insulin and lower blood glucose up to three months after

implantation, without affecting the weight or overall wellbeing of the mice. This suggests that

in differentiated postmitotic cells, *TYMS* gene is dispensable.

The advances in basic science and technological developments simultaneously pressure

and aid pre-clinical studies in their translation to clinical applications. This remains particularly

true for the so-called *advanced therapy medicinal products* (ATMP), which include somatic-

cell and gene therapies, and tissue-engineered medicines. The regulatory agencies categorize

CRISPR-based products as ATMP. Although some of these products have reached the clinical

trial stage, the safety concerns arising from altering DNA challenge further applications and

the continuity of ongoing trials. In this context, attempts to control the activity and

proliferation of the therapeutic cells have proven relatively efficient. These systems, however,

have not yet led to the progression of clinical trials to further regulatory phases, possibly due

to their complexity or partial/remissive function.

We hope that future differentiations of *TYMS^-/-* hiPSC into other therapeutic tissues will

further confirm the utility of this approach for this form of cell-based therapies. In addition,

it is possible to expand this strategy to other therapeutic cell types. Immunotherapies, for

instance, currently present the challenge of cells overactivation and overexpansion and are

restricted to the use of senescent cells to limit risks. We expect them to benefit from our

controllable safety approach, as it would enable and regulate the activity of the therapeutic

cells according to supplementation.

Materials and methods

Culture of HEK293 FT, A431, A549, HSC3

We cultured HEK293, epidermoid carcinoma cell line A431, adenocarcinoma human alveolar basal epithelial cells A549, and human oral squamous carcinoma cell line HSC-3 cells in Dulbecco's modified Eagle's medium (DMEM) containing 10% fetal bovine serum (FBS), 100 µg/ml penicillin-streptomycin and 2 mM GlutaMAX (all ThermoFisher Scientific). After *TYMS* knock-out, cell media were additionally supplemented with 5-20µM dTMP (ThermoFisher Scientific). We kept the cell lines at 37 °C and 5% CO2, with media change every other day until splitting with TrypLE select. All cell lines tested negative for mycoplasma.

Induced Pluripotent Stem cells culture and characterization

Human induced pluripotent stem cell (hiPSC) line Hel24.3 were cultured on Matrigel (Corning) coated plates in E8 medium (ThermoFisher Scientific) and split using 0.5 mM EDTA. We kept the cell lines at 37 °C and 5% CO2, with media change every other day. All cell lines tested negative for mycoplasma. After *TYMS* knock-out, cell media were additionally supplemented with 5-20µM dTMP. The karyotyping was carried out by Ambar Anàlisis Mèdiques (Barcelona, Spain) by G-banding. We treated edited (passage 20 and 45) and non-edited (passage 20) hiPSC with colcemid for 4 h, 37°C, 5% CO2 and proceeded to prepare the cells as recommended by the service provider.

Genome editing

To knock-out exon 5, we designed guide RNA targeting intron 4 (I4) and intron 5 (I5) using online tools (https://benchling.com, CRISPOR,[24], supplementary table 1). We generated the sgRNA by incubating 5 min at 95 °C our gRNA (customized Alt-R CRISPR-Cas9 gRNA, iDT) with Alt-R® CRISPR-Cas9 tracrRNA, ATTO™ 550 (iDT).

For each stem cell electroporation experiment, we dissociated 2x10^6 cells to single-cells with StemPro Accutase (ThermoFisher Scientific). We complexed the sgAlt-R® S.p. HiFi Cas9 Nuclease V3 and and both sgRNA to form the functional ribonucleoprotein (RNP) and delivered this RNP into the cells, along the Alt-R® Cas9 electroporation enhancer (all from integrated DNA Technologies – iDT), by electroporation with Neon transfection systems (1100V, 20ms, 2

pulses). Cells were plated onto Matrigel-coated plates containing E8 with 5 µM ROCK inhibitor

(Y-27632, Selleckchem) and 20-40 µM dTMP, and incubated at 37 ˚C, 5% CO2.

24-48 h after electroporation, we sorted ATTO550 positive cells for monoclonal expansion in

E8 with 5 µM ROCK inhibitor, 20-40 µM dTMP and 10% CloneR (StemCell Technologies). The

media was refreshed every 72 h until splitting. We individually screened monoclonal colonies

by PCR. All PCR products of the selected edited clones were validated by Sanger sequencing,

along with the integrity of the top-7 off target sequences (Supplementary table 2).

Embryoid body differentiation

To test the pluripotency for embryonal lineage differentiation, we grew the cells until 90-

100% confluency and performed embryoid body (EB) differentiation assay as previously

described [2]. Final plated EBs were fixed with 4%PFA-PBS and immunostained for characteristic

markers of the three germ-layers (see list below).

Teratoma assay

Knock-out and wild-type Hel24.3 cells were cultured in ultra-low attachment plates with E8

supplemented with 5 µM ROCK inhibitor for 24-48h to form aggregates. The medium for knock-

out cells was additionally supplemented with 40 µM dTMP until 2h before implantation. Groups

of approximately 500 spheres were collected in syringe cannulas for subcutaneous implantation

in the legs of the mice. For half of the mice, the drinking water was supplemented with 100

mg/kg/day dTMP. 6-8 weeks post implantation the teratomas were extracted and fixed for

immunocytochemistry and hematoxilin-eosin stainings as described below.

dTMP auxotrophy

Wild-type and knock-out clones from each cell line were split onto Matrigel coated 6-well

plates at a reason of a twenty-five thousand cells per well. Each well contained the

corresponding medium for the cell line (see *cell culture*) supplemented with 0 to 2mM dTMP.

Cells were washed and media were refreshed daily to clear away dead cells debris and keep the

thymidine concentration as constant as possible.

Beta-cell differentiation

The *in vitro* beta cell differentiation was carried out as previously described[3]. Briefly, *TYMS* knock-out and control Hel24.3 iPS cells were seeded onto Matrigel coated 10 cm plates as previously stated. Differentiation protocol was started 24 h post-seeding, when the medium was changed to D0 medium. Both cell lines were supplemented with 0 or 40 µM dTMP until maturation. At stage 7, dTMP was withdrawn or kept until analysis (week 1 to 4 from first day of S7). The media was refreshed every 2-3 days until analysis.

Flow cytometry

Stage 4 cells and stage 7 stem cell derived islets (SC-islets) were dissociated with TrypLE for 10 min at 37 °C and resuspended in 5% FBS-containing PBS. Fixation and permeabilization were done using Cytofix/Cytoperm (BD Biosciences, catalog no. 554714) for 20 min at RT. Then, samples were incubated overnight with primary antibodies at 4 °C, followed by secondary antibodies for 30 min in RT in Perm/Wash buffer (BD Biosciences, catalog no. 554714) supplemented with 5% FBS. The cells were run on FACSCalibur cytometer (BD Biosciences); data were collected with CellQuest Pro v.4.0.2 (BD Biosciences) and analyzed with FlowJo™ v10.8 Software (BD Life Sciences). Antibodies are listed in Supplementary Table 3.

For cell cycle analysis, wild type and knock out hiPSC were plated overnight on 6 well plates containing E8 supplemented with ROCK inhibitor and 20 µM dTMP. Then, the media were refreshed and for some plates the dTMP supplementation was stopped. Cells from individual wells were collected at 0, 2, 4, 10, 16 and 24h after dTMP withdrawal and fixed using 4% PFA in in PBS. The same timepoint were collected for the supplemented plates. After fixation, samples were washed and incubated 30 min with 10 µg/ml DAPI diluted in 1% Triton-X (Sigma-Aldrich) in PBS. Then, they were immediately analyzed using the NovoCyte Quanteon 4025.

Insulin secretion analysis *in vitro*

For static analysis of insulin secretion, a total of 30–50 SC-islets were picked and incubated for 90 min in a 12 well plate containing 2.8 mM glucose (G2.8) in Krebs-Ringer buffer (KRB) for equilibration. This was followed by sequential 30-min incubations of G2.8, 16.8 mM glucose (G16.8) and G2.8+ 30 mM KCl in KRB. After each incubation, 200uL samples were taken for further insulin analysis. Once all samples were taken, the islets were collected for DNA quantification. Dynamic insulin secretion test was carried out using a perifusion apparatus

(Brandel Suprafusion SF-06) at a 0.25 ml min–1 flow rate, sampling every 4 min. 50 SC-islets

were handpicked and perfused with KRB; sample collection started after 90 min of equilibration

in G2.8. Insulin content of secretion fractions and SC-islet lysates was analyzed with enzyme-

linked immunosorbent assay (ELISA) (Mercodia).

Islet *in vivo* characterization

Animal care and experiments were approved by National Animal Experiment Board in

Finland (ESAVI/14852/2018). NOD-SCID-Gamma (NSG, Jackson Laboratories, catalog no.

0055577) mice were housed in the Biomedicum Helsinki conventional facility in 12 h light/dark

cycle and fed standard chow. SC-islet implantations were done following a stablished previously

described protocol[3]. Non-fasted blood samples were collected monthly from the saphenous

vein for glucose measurement and C-peptide secretion analysis. For further characterization of

the SC-islet graft, the engrafted kidney was removed after 1 or 3 months and processed as

described below.

Immunocytochemistry, immunohistochemistry, and image analysis

Samples from hiPSC were fixed using 4%PFA-PBS for 20 min and permeabilized using 1%

Triton-X in PBS. Samples from S4 and S7 SC-islets were fixed with 4% PFA in PBS for 2h, and

explanted SC-islet grafts and teratoma-like growths were fixed overnight. After fixation, these

samples were embedded in paraffin and sectioned. 5 μm sections were deparaffinized and

subjected to HIER in 0.1 mmol l–1 citrate buffer. The cells and tissue slides were blocked with

UV-block (Thermo Scientific, catalog no. TA-125-PBQ), and incubated with primary antibodies

in 0.1% Tween-20 overnight at 4 °C with the given dilutions (Supplementary Table 3). After

washing, secondary antibodies diluted in a similar manner were incubated at RT for 1 h in the

presence of Hoechst33342 for nuclear staining.

The cells and slides were imaged using Apotome II from a Zeiss AxioImager with the same

exposure and export setting used on all slides of each immunostaining. Images were processed

in Zen2 Blue Edition v.2 (Zeiss) and analyzed using FiJi [25] or CellProfiler v.4.0[26] with pipelines

adapted from a previous report[27]. The same pipeline settings were used on all images of each

immunostaining.

Protein expression analysis

mRNA expression levels were analyzed by qPCR (supplementary table 4-6). TYMS protein

expression was assessed by standard Western blot. Briefly, protein was extracted by using

RIPA lysis buffer (20-188, Millipore) supplemented with cOmplete Mini protease and

phosphatase inhibitors tablets (Roche). Samples were sonicated (30 sec, %50 duty cycle pulse)

and centrifuged for soluble protein extraction. Protein quantification was done using a Pierce

BCA protein assay kit (ThermoFisher) and 20-30 µg total protein were run in Mini-PROTEAN

TGX Precast gels (#4561033, Bio-Rad). Transfer to a nitrocellulose membrane was done using

25V for 6 min in the iBlot2 Gel Transfer device (IB21001, ThermoFisher). Then, the membrane

was probed with a rabbit polyclonal primary antibody against TYMS (1:2500, Proteintech;

Supplementary table 3). For detection, we used an anti-rabbit HRP-coupled secondary

antibody (1:5000, Cell Signalling) and Western ECL substrate (#1705061, Bio-Rad. As a loading

control, we probed the membrane with a b-actin-HRP coupled antibody (sc-47778, Santa

Cruz).

355
356

References

1.   Takahashi, K. & Yamanaka, S. Induction of Pluripotent Stem Cells from Mouse
Embryonic and Adult Fibroblast Cultures by Defined Factors. *Cell* **126**, 663–676
(2006).

2.   Weltner, J. *et al.* Human pluripotent reprogramming with CRISPR activators. *Nat*
Commun (2018) doi:10.1038/s41467-018-05067-x.

3.   Balboa, D., Barsby, T. & Lithovius, V. ✉. Functional, metabolic and transcriptional
maturation of human pancreatic islets derived from stem cells. doi:10.1038/s41587-
022-01219-z.

4.   Maldonado, R., Jalil, S. & Wartiovaara, K. Curative gene therapies for rare diseases. *J*
Community Genet (2020) doi:10.1007/s12687-020-00480-6.

5.   Maeder, M. L. & Gersbach, C. A. Genome-editing Technologies for Gene and Cell
Therapy. (2016) doi:10.1038/mt.2016.10.

6.   Hsu, P. D., Lander, E. S. & Zhang, F. Development and applications of CRISPR-Cas9 for
genome engineering. *Cell* **157**, 1262–1278 (2014).

7.   Picken, A. *et al.* A Monte Carlo framework for managing biological variability in
manufacture of autologous cell therapy from mesenchymal stromal cells therapies.
Cytotherapy **22**, 227–238 (2020).

8.   Aboul-Soud, M. A. M., Alzahrani, A. J. & Mahmoud, A. cells Induced Pluripotent Stem
Cells (iPSCs)-Roles in Regenerative Therapies, Disease Modelling and Drug Screening.
(2021) doi:10.3390/cells10092319.

9.   Zhao, T., Zhang, Z.-N., Rong, Z. & Xu, Y. Immunogenicity of induced pluripotent stem
cells. (2011) doi:10.1038/nature10135.

10.   Themeli, M. *et al.* Generation of tumor-targeted human T lymphocytes from induced
382   pluripotent stem cells for cancer therapy. (2013) doi:10.1038/nbt.2678.

383   11.   Xu, H. *et al.* Targeted Disruption of HLA Genes via CRISPR-Cas9 Generates iPSCs with
384   Enhanced Immune Compatibility. *Cell Stem Cell* **24**, 566-578.e7 (2019).

385   12.   Lee, A. S. *et al.* Effects of cell number on teratoma formation by human embryonic
386   stem cells. *Cell Cycle* **8**, 2608–2612 (2009).

387   13.   Hentze, H. *et al.* Teratoma formation by human embryonic stem cells: Evaluation of
388   essential parameters for future safety studies. (2009) doi:10.1016/j.scr.2009.02.002.

389   14.   Wiebking, V. *et al.* Metabolic engineering generates a transgene-free safety switch for
390   cell therapy. doi:10.1038/s41587-020-0580-6.

391   15.   Bonifant, C. L., Jackson, H. J., Brentjens, R. J. & Curran, K. J. Toxicity and management
392   in CAR T-cell therapy. *Mol Ther Oncolytics* **3**, 16011 (2016).

393   16.   Lim, T.-T., Geisen, C., Hesse, M., Fleischmann, B. K. & Zimmermann, K. Lentiviral
394   Vector Mediated Thymidine Kinase Expression in Pluripotent Stem Cells Enables
395   Removal of Tumorigenic Cells) Lentiviral Vector Mediated Thymidine Kinase
396   Expression in Pluripotent Stem Cells Enables Removal of Tumorigenic Cells. *PLoS One*
397   **8**, 70543 (2013).

398   17.   Chen, F. *et al.* Suicide gene-mediated ablation of tumor-initiating mouse pluripotent
399   stem cells. doi:10.1016/j.biomaterials.2012.11.018.

400   18.   Volarevic, V. *et al.* Ethical and Safety Issues of Stem Cell-Based Therapy. *Int. J. Med.*
401   *Sci* **15**, 36–45 (2018).

402   19.   Andersona, D. D., Quintero, C. M. & Stovera, P. J. Identification of a de novo
403   thymidylate biosynthesis pathway in mammalian mitochondria. *Proc Natl Acad Sci U S*
404   *A* **108**, 15163–15168 (2011).

405   20.   Chen, D., Jansson, A., Sim, D., Larsson, A. & Nordlund, P. Structural analyses of human
406   thymidylate synthase reveal a site that may control conformational switching
407   between active and inactive states. *J Biol Chem* **292**, 13449–13458 (2017).

408   21.   Trokovic, R., Weltner, J. & Otonkoski, T. Generation of iPSC line HEL24.3 from human
409   neonatal foreskin fibroblasts. *Stem Cell Res* **15**, 266–268 (2015).

410   22.   Traut, T. W. Physiological concentrations of purines and pyrimidines. *Mol Cell*
411   *Biochem* **140**, 1–22 (1994).

412   23.   Diehl, F. F. *et al.* Nucleotide imbalance decouples cell growth from cell proliferation.
413   doi:10.1038/s41556-022-00965-1.

414   24.   Concordet, J. P. & Haeussler, M. CRISPOR: Intuitive guide selection for CRISPR/Cas9
415   genome editing experiments and screens. *Nucleic Acids Res* (2018)
416   doi:10.1093/nar/gky354.

417   25.   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nature*
418   *Methods 2012 9:7* **9**, 676–682 (2012).

419   26.   Stirling, D. R. *et al.* CellProfiler 4: improvements in speed, utility and usability. *BMC*
420   *Bioinformatics* **22**, 1–11 (2021).

421   27.   Lithovius, V. *et al.* SUR1-mutant iPS cell-derived islets recapitulate the
422   pathophysiology of congenital hyperinsulinism. *Diabetologia* **64**, 630–640 (2021).

423   28.   Karjoo, Z., Chen, X. & Hatefi, A. Progress and Problems with the Use of Suicide Genes
424   for Targeted Cancer Therapy. *Adv Drug Deliv Rev* **99**, 113 (2016).

425   29.   Berger, C., Flowers, M. E., Warren, E. H. & Riddell, S. R. Analysis of transgene-specific
426   immune responses that limit the in vivo persistence of adoptively transferred HSV-

TK–modified donor T cells after allogeneic hematopoietic cell transplantation. *Blood*
107, 2294 (2006).

30.   Sheikh, S., Ernst, D. & Keating, A. Prodrugs and prodrug-activated systems in gene
therapy. doi:10.1016/j.ymthe.2021.04.006.

31.   Sahillioglu, A. C. & Schumacher, T. N. Safety switches for adoptive cell therapy. *Curr*
Opin Immunol **74**, 190–198 (2022).

32.   Buil, B. A., Tackenberg, C. & Rust, R. Editing a gateway for cell therapy across the
blood–brain barrier. *Brain* **146**, 823 (2023).

33.   Pellegrini, S., Zamarian, V. & Sordi, V. Strategies to Improve the Safety of iPSC-Derived
β Cells for β Cell Replacement in Diabetes. doi:10.3389/ti.2022.10575.

34.   Tiberghien, P. *et al.* Ganciclovir treatment of herpes simplex thymidine kinase-
transduced primary T lymphocytes: an approach for specific in vivo donor T-cell
depletion after bone marrow transplantation? *Blood* **84**, 1333–1341 (1994).

35.   Austin, E. A. & Huber, B. E. A first step in the development of gene therapy for
colorectal carcinoma: cloning, sequencing, and expression of Escherichia coli cytosine
deaminase. *Mol Pharmacol* **43**, (1993).

36.   Felmer, R. N. & Clark, J. A. The gene suicide system NTR/CB1954 causes ablation of
differentiated 3T3L1 adipocytes by apoptosis. *Biol Res* **37**, 449–460 (2004).

37.   Gadi, V. K. *et al.* In vivo sensitization of ovarian tumors to chemotherapy by
expression of E. coli purine nucleoside phosphorylase in a small fraction of cells. *Gene*
Ther **7**, 1738–1743 (2000).

38.   Sato, T. *et al.* Engineered Human tmpk/AZT As a Novel Enzyme/Prodrug Axis for
Suicide Gene Therapy. *Molecular Therapy* **15**, 962–970 (2007).

39.   Serafini, M. *et al.* Characterization of CD20-transduced T lymphocytes as an
alternative suicide gene therapy approach for the treatment of graft-versus-host
disease. *Hum Gene Ther* **15**, 63–76 (2004).

40.   Wang, X. *et al.* A transgene-encoded cell surface polypeptide for selection, in vivo
tracking, and ablation of engineered cells. *Blood* **118**, 1255–1263 (2011).

41.   Kieback, E., Charo, J., Sommermeyer, D., Blankenstein, T. & Uckert, W. A safeguard
eliminates T cell receptor gene-modified autoreactive T cells after adoptive transfer.
Proc Natl Acad Sci U S A **105**, 623–628 (2008).

42.   Birrer, M. J., Moore, K. N., Betella, I. & Bates, R. C. Antibody-Drug Conjugate-Based
Therapeutics: State of the Science. *J Natl Cancer Inst* **111**, 538–549 (2019).

43.   Straathof, K. C. *et al.* An inducible caspase 9 safety switch for T-cell therapy. *Blood*
105, 4247–4254 (2005).

44.   Liu, Y. *et al.* Inducible caspase-9 suicide gene under control of endogenous oct4 to
safeguard mouse and human pluripotent stem cell therapy. *Mol Ther Methods Clin*
Dev **24**, 332–341 (2022).

45.   Wunderlich, S. *et al.* Targeted biallelic integration of an inducible Caspase 9 suicide
gene in iPSCs for safer therapies. *Mol Ther Methods Clin Dev* **26**, 84–94 (2022).

Conflict of interest

RSM and KW are marked as inventors in a patent application by the University of Helsinki

Aknowledgements

We thank MD Vaino Lithovius for his assistance in image analysis, MSc Sami Jalil for scientific discussion and support, MSc Hazem Ibrahim and MSc Eliisa Vahankangas for assistance with animal care. We thank the Flow cytometry Unit, the Functional Genetics Unit (FuGU) and the Animal facility at Biomedicum Helsinki for their possible contribution to this study. Lastly, we thank Helsinki University Hospital research funding, Paulo Foundation, Signe and Ane Gyllenberg Foundation, Magnus Ehrnrooth Foundation, K. Albin Johansson Foundation, Ida Montin Foundation, Finnish Red Cross Blood Service Research Fund for financial support.

517 Table 1 – Summary table of selected "safety" systems, and potential advantages and
518 disadvantages.
519

Type	Function	Example	Advantages	Disadvantages
Enzyme/Prodrug [28–33]	Transgenes transform prodrug into toxic metabolite	HSV-TK + Ganciclovir[34] CD+5-FC[35] NTR+CB1954[36] PNP+MEP[37] mTMPK+AZT[38]	Some have the ability to eliminate the whole graft Could be combined with other strategies Safe Generally efficient	Transgene derive from viruses, bacteria or yeast Immuno-rejection of the therapeutic cells Toxicity of the prodrug Bystander effect Slow activation time Silencing of transgene
MAb mediated [32,33]	Engineer a gene targeted by an Ab into the transplanted cells	CD20t+Rituximab hEGFRt+Cetuximab[40] c-myc tag + Anti-c-myc tag[41] X+ADC anti-X[42]	Avoid GVHD[39] Efficient Different payloads for ADC	CD20 present in endogenous B cells
Inducible Dimerization[32]	Insertion of a modified Caspase-9. The recruitment sequence is replaced by a dimerization domain of FK506 binding protein	iCas9[43]+CID	Low immunogenicity Inert small molecule Fast action Tested to eliminate iPSC [44]	Silencing of transgene CID resistance[45]
Metabolic	Endogenous disruption of pyrimidine de novo synthesis	UMPS KO[14]	Transgene-free	May affect RNA syntesis
CID: Chemical inducer of dimerization; ADC: antibody–drug conjugate; GVHD: Graft-versus-host disease; CD: cytosine deaminase; NTR: Nitroreductase; HSV-TK: herpes simplex virus thymidine kinase; PNP: purine nucleoside phosphorylase; mTMPK: mutated thymidylate monophosphate kinase; OxPD: Oxazaphorine drugs; AZT: azidothymidine				

520
521

Figure 1. **A.** Diagram showing the nucleotide pathways for DNA (orange) or RNA (blue) synthesis. **B.** Schematic representation of the gene of interest (in TYMS), showing the two gRNAs for our CRISPR/Cas9 approach, and the amino acids that form the active site of the protein over the corresponding exon that code for them. The lower line represents the change of frame upon DNA repair. **C.** TYMS expression in wild-type versus knock-out hiPSC (Hel 24.3), HEK293, A431, A549 and HSC3 (n=3/cell line). Results shown as ratio of TYMS mRNA over GAPDH mRNA contents. **D.** Growth curve of knock-out hiPSC under different concentrations of dTMP. Wild-type without dTMP supplementation (blue) used as control. **E.** Growth curve of wild-type and knock-out hiPSC supplemented with 0 or 25 μM dTMP. Results shown as average confluency per image ⍰ SD. Results shown as average confluency per image ⍰ SD. Statistical significance based on Wilcoxon test; $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***), $p < 0.0001$ (****).

Figure 2. A. hiPSC immunostaining against pluripotency markers NANOG, OCT4, Sox2 and SSEA4. **B.** hiPSC qPCR for expression analysis of pluripotency markers NANOG, OCT4 and Sox2. Results shown as mRNA over Cyclog mRNA. **C.** Karyotype of TYMS-knock-out hiPSC, passage 45. **D.** Graphical representation of trilineage differentiation protocol, dTMP supplementation shown below. **E.** Number of resulting 300 μm aggregates from wild-type and knock-out hiPSC with (+dTMP) and without (-dTMP) supplementation during the first stage of differentiation (day 0 to 14). **F.** Staining against markers for endoderm (Sox17), mesoderm (SMA), and ectoderm (beta-3-tubulin) in cells derived from wild-type and knock-out hiPSC under dTMP supplementation during the first stage of differentiation. Statistical significance in panels B and E is based on Wilcoxon test; $p > 0.05$ (ns), $p < 0.05$ (*), $p < 0.01$ (**).

Figure 3. A. Graphical representation of teratoma formation experiment. **B.** Knock-out and wild-type hiPSC-derived teratomas with and without dTMP supplementation as extracted from the mice, before processing for molecular analysis. **C.** Nuclear count of sliced teratomas. **D.** Macro (1.25x) and micro (10x) images of haematoxylin-eosin staining show dense and cystic structures within the tumours, along with structures of the three germ layers (arrows – not there, we need to spot them). **E.** Immunostainings against KI67 (proliferation markers, green) and total nuclei (Hoechst, blue). **F.** Quantification of KI67 positive nuclei from immunostainings. Statistical significance based on Wilcoxon test; $p>0.05$ (ns), $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***), $p < 0.0001$ (****).

Figure 4. A. Graphical representation of in vitro beta cell differentiation from hiPSC. dTMP supplementation was kept until stage 7 (S7). Following tests were carried out four weeks from stage 7 (S7W4). **B.** Flow cytometry analysis for markers for different stages of differentiation: CXCR4 for definite endoderm stage; NKX6.1 and PDX1 for stage 4; NKX6.1, C-peptide, PDX1, insulin (INS) and glucagon (GCG) for stage 7. **C.** mRNA expression analysis of maturation markers at different stages. Results are shown as each tested mRNA over GAPDH mRNA. **D.** Immunostaining against insulin (INS, magenta) and glucagon (GCG, yellow) in hiPSC-derive islets at stage 7, after 5 days of dTMP supplementation withdrawal. **E.** Dynamic insulin secretion in response to different stimuli (explain). **F.** Insulin secretion in response to low (2.8) and high (16.8) glucose concentration and glucose plus KCl (2.8 KCl). Statistical significance based on Wilcoxon test; p>0.05 (ns), p < 0.05 (*), p < 0.01 (**).

Figure 5. A-C. Follow up of human C-peptide secretion (to serum), blood glucose and mice body weight after 1-, 2- and 3-months from implantation (month 0). **D.** Immunostaining of engrafted kidney 3-months post-implantation. INS (magenta), GCG (yellow). **E.** Immunostaining against EdU and KI67 in grafts after 1 and 3 months. Statistical significance based on Wilcoxon test; p>0.05 (ns).