1 **No sex differences in oxygen uptake or extraction kinetics in the moderate or heavy**
2 **exercise intensity domains**
3
4

5 Maria Solleiro Pons[1], Lina Bernert[1,2], Emily Hume[1], Luke Hughes[1], Zander Williams[3], Mark
6 Burnley[4], Paul Ansdell[1]
7
8
[1] Department of Sport, Exercise and Rehabilitation, Faculty of Health and Life Sciences,
Northumbria University, UK
[2] Institute of Sport and Exercise Sciences, University of Muenster, Germany
[3] Department of Respiratory Medicine, Royal Brompton Hospital, Royal Brompton and
Harefield NHS Trust, UK
[4] School of Sport, Health and Exercise Sciences, Loughborough University, UK

15
16
17
18 **Running Title:** The influence of sex on oxygen uptake and extraction
19
20
21

Corresponding Author:
Paul Ansdell, PhD
Department of Sport, Exercise and Rehabilitation
Faculty of Health and Life Sciences
Northumbria University
Newcastle upon Tyne
NE1 8ST
UK
p.ansdell@northumbria.ac.uk

Abstract

The integrative response to exercise differs between sexes, with oxidative energy provision purported as a potential mechanism. However, there is a lack of systematic investigation into the kinetics of oxygen uptake ($\dot{V}O_2$) and extraction (HHb+Mb) during exercise in both sexes.

Sixteen young adults (8 females, age: 27±5 years) completed three experimental visits. Incremental exercise testing was performed to obtain lactate threshold and $\dot{V}O_{2peak}$. Subsequent visits involved three six-minute cycling bouts at 80% of lactate threshold and one 30-minute bout at a work rate 30% between the lactate threshold and maximal ramp test power. Pulmonary gas exchange and near-infrared spectroscopy continuously sampled $\dot{V}O_2$ and HHb+Mb, respectively. The mean of six moderate and two heavy intensity transitions were modelled with mono-exponential curves to quantify the phase II response to exercise. Slow component amplitudes were also quantified for the heavy intensity domain.

Males and females demonstrated similar relative $\dot{V}O_{2peak}$ values (46.2±6.6 vs 40.5±6.7 ml.kg^{-1}.min^{-1}, p=0.111), with males achieving ~30% greater power outputs (p=0.002). In both the moderate and heavy intensity domains, the relative amplitude of the phase II transition was similar between sexes for $\dot{V}O_2$ ($p \geq 0.179$) and HHb+Mb ($p \geq 0.193$). Similarly, there were no sex differences in the time constants for $\dot{V}O_2$ ($p \geq 0.385$) or HHb+Mb ($p \geq 0.274$). In the heavy intensity domain, neither $\dot{V}O_2$ ($p \geq 0.686$) or HHb+Mb ($p \geq 0.432$) slow component amplitudes were different between sexes.

The oxidative response to moderate and heavy intensity exercise did not differ between males and females, suggesting both sexes experience similar degrees of bioenergetic stress during intensity-matched exercise.

New and Noteworthy

This study demonstrated no sex differences in the bioenergetic response to moderate and heavy intensity cycling exercise. The change in oxygen uptake and deoxyhaemoglobin were modelled with mono-exponential curve fitting, which revealed that both sexes increased the rate of oxidative energy provision similarly. This provides insight into previously reported sex differences (e.g., fatigability) suggesting that the mechanisms are contractile rather than metabolic.

Introduction

The transition from rest to exercise involves an integrated response from the pulmonary, cardiovascular, and muscular systems to rapidly increase the supply and utilisation of oxygen for oxidative adenosine triphosphate (ATP) provision (Poole & Jones, 2012). The speed at which this process can occur can be quantified using pulmonary oxygen uptake ($\dot{V}O_2$) kinetics, and is thought to be an influencing factor in high-intensity exercise performance (Burnley & Jones, 2007). The $\dot{V}O_2$ response can be broken down into three phases, beginning with the initial cardio-dynamic phase (phase I, 10-20 s) which represents an increased venous return via the muscle pump effect, as well as increased pulmonary blood flow (Grassi *et al.*, 1996). Thereafter, increases in pulmonary $\dot{V}O_2$ are considered to reflect increased muscle oxygen uptake in response to exercise (phase II), until the energy demand of exercise is met by oxidative phosphorylation and $\dot{V}O_2$ reaches a steady state (Hughson *et al.*, 2001). A steady state response is attainable within the moderate intensity domain, whereas in either heavy or severe intensity domains, a further rise in $\dot{V}O_2$ is observed, termed the slow component. This three-phase response is ubiquitous in exercising humans; however, the biological characteristics of the individual can influence the rates at which they occur.

The time constant of phase II kinetics are considered to be a crucial determinant of the decrease in contractile function experienced by the exercising individual (Goulding *et al.*, 2021). As oxidative phosphorylation does not immediately meet the demand for ATP, substrate-level phosphorylation is required (Burnley & Jones, 2007). Intuitively, the rate at which oxidative metabolism can be upregulated at the onset of exercise is inversely linked with the accumulation of deleterious metabolites such as hydrogen ions [H^+] and inorganic phosphate [P_i], as well as the depletion of phosphocreatine [PCr] stores, which all interfere with excitation-contraction coupling (Allen *et al.*, 2008). Accordingly, Temesi *et al.* (2017) demonstrated a positive correlation between the time constant ($\tau\dot{V}O_2$) of the phase II response and the decrease in quadriceps potentiated twitch force. Similarly, elite endurance athletes demonstrate faster $\tau\dot{V}O_2$ (Koppo *et al.*, 2004) and lesser declines in contractile function (Ducrocq *et al.*, 2021) compared to untrained individuals when exercising at similar relative intensities. Collectively, this evidence suggests that those individuals able to meet the ATP demand with oxidative metabolism quicker, experience less contractile impairment during exercise.

A consistent finding in studies comparing males and females exercising at the same metabolic intensity is that females experience a lesser degree of contractile impairment of the knee-extensors (Ansdell *et al.*, 2019; Ansdell *et al.*, 2020a; Azevedo *et al.*, 2021). Previously, this has been suggested to be a result of sex differences in skeletal muscle composition, whereby

females consistently demonstrate a greater proportional area of type I fibres (Staron *et al.*,
2000; Roepstorff *et al.*, 2006). The consequences of this fibre type difference are multi-
factorial; for-instance, it is well established that type I fibres are more fatigue-resistant
(Schiaffino & Reggiani, 2011). Additionally, female *vastus lateralis* capillary density is ~23%
greater in females compared to males (Roepstorff *et al.*, 2006), while females also
demonstrate greater mitochondrial oxidative function and intrinsic respiratory rates than males
of equivalent training status (Cardinale *et al.*, 2018). One factor that remains unexplored is
whether these physiological sex differences result in differences in the metabolic response to
exercise. Conceivably, the superior aerobic phenotype of female skeletal muscle could imply
that females might be able to meet the ATP demand of exercise through oxidative means
faster than males, however this is a hypothesis that remains untested.

When the slow component is considered, the increase in $\dot{V}O_2$ during constant-load exercise
implies an impairment of efficiency and is likely an amalgamation of several concurrent
physiological changes. Within skeletal muscle, the accumulation of metabolites and
associated contractile dysfunction is linked with the loss of efficiency (Grassi *et al.*, 2015). This
dysfunction then leads to a compensatory recruitment of larger motor units that express
predominantly type II muscle fibres (Krustrup *et al.*, 2004). Of relevance here, is that female
skeletal muscle has consistently been demonstrated to be more fatigue-resistant (Ansdell *et*
al., 2019; Ansdell *et al.*, 2020a), and require less additional motor unit recruitment in states of
fatigue (Ansdell *et al.*, 2017; Ansdell *et al.*, 2019). This could represent a lesser requirement
for additional oxygen uptake during constant load exercise in females, however, this remains
unexplored.

Despite more aerobically-suited skeletal muscle, females have lower levels of haemoglobin
(Murphy, 2014), which is thought to impair O_2 carrying capacity during exercise (Harms *et al.*,
1998). During exercise where O_2 delivery and utilisation are both limiting factors (e.g., cycling),
these factors are thought to counteract each other to enable comparable relative metabolic
thresholds between the sexes (Ansdell *et al.*, 2020b). To date, the only investigation to
systematically investigate the oxidative adjustment at the onset of exercise between sexes did
so during low intensity treadmill walking (Beltrame *et al.*, 2017). Data from this study
suggested faster O_2 extraction in females, fitting with the notion that phase II $\dot{V}O_2$ kinetics are
influenced by intramuscular factors in healthy humans (Poole & Jones, 2012). Despite this,
treadmill walking is not considered to be limited by O_2 delivery, therefore the findings do not
necessarily translate to high-intensity locomotor exercise, during which all levels of the O_2
cascade are considered to be influential in determining metabolic responses to exercise
(Goulding & Marwood, 2023).

Accordingly, the present study aimed to compare the kinetics of pulmonary $\dot{V}O_2$ as well as

muscle oxygen extraction, inferred from deoxyhaemoglobin and myoglobin (HHb+Mb)

kinetics, in both sexes during moderate and heavy intensity exercise. It was hypothesised that

females would demonstrate a smaller value for the phase II time constant (i.e., faster kinetics)

for $\dot{V}O_2$ and HHb+Mb at the onset of exercise, and a smaller slow component amplitude in the

heavy intensity domain.

Methods

Ethical Approval

This study received institutional ethical approval from the Northumbria University Health and

Life Sciences Research Ethics Committee (submission reference: 49189) and was conducted

according to all aspects of the Declaration of Helsinki, apart from pre-registration in a

database. Participants volunteered for the study and provided written informed consent.

Participants

Using the effect size for the sex difference in *vastus lateralis* tissue oxygenation during heavy

intensity exercise ($\eta p^2 = 0.509$) from (Ansdell *et al.*, 2020a), an *a priori* sample size calculation

determined a minimum of 14 participants (seven females and seven males) were required to

detect an effect ($\alpha = 0.05$, power = 0.95). Therefore, eight males (mean ± SD age: 27 ± 3

years; stature: 182 ± 5 cm; body mass: 75.3 ± 10.2 kg) and eight females (mean ± SD age:

27 ± 7; stature: 163 ± 4 cm; body mass: 61.8 ± 5.9 kg) volunteered to take part in the study.

Hormonal status was not an exclusion criterion or controlled for in this study. Female

participants were tested in any phase of their menstrual cycle and there were no restrictions

on hormonal contraceptive usage. This decision was based on evidence from (Mattu *et al.*,

2020) who demonstrated no hormonal effects on $\dot{V}O_2$ kinetics during cycling exercise. All

participants were free from cardiovascular, respiratory, and neurological disease as well as

musculoskeletal injury.

Experimental Design

All participants visited the laboratory on three occasions. During the first visit, participants were

familiarised with the experimental procedures and completed two incremental exercise tests

to quantify lactate threshold and peak oxygen uptake ($\dot{V}O_{2peak}$). The second and third visits

were identical and involved three six-minute bouts of moderate intensity exercise (80% of

lactate threshold, LT), separated by six minutes of unloaded pedalling. Thereafter, a single

bout of heavy intensity exercise (30% ΔLT – $\dot{V}O_{2peak}$) was performed for 30 minutes.

 Visit 1: Familiarisation & Incremental Testing

The first visit began with participants completing a screening questionnaire to ensure inclusion

criteria were met. Thereafter, participants moved onto the cycle ergometer (Velotron, SRAM,

Chicago, IL, USA) which was set up with the seat height aligned with the hip, and handlebar

height set according to the participants' comfort, these measurements were recorded and

replicated for subsequent trials. The breath-by-breath gas exchange mask was then placed

over the participant's mouth and nose, and an air-tight seal was ensured before resting data

was recorded. Following resting measures, participants completed five minutes of warm-up

cycling at a light intensity (60 W) at a self-selected cadence between 70-100 rpm, before

commencing an incremental exercise test. The first incremental exercise test began at 75 W

and increased by 25 W every five minutes. At the end of each stage, a capillary blood sample

was drawn from the participants' fingertip and immediately analysed to determine whole blood

lactate concentration (mmol.L^{-1}, Biosen C-Line, EKF Diagnostics, Germany). The test was

terminated once LT was identified as the first work rate at which a non-linear increase in blood

lactate concentration was observed, after which, participants were provided 20 minutes of

passive rest.

Next, participants began the second incremental test with five minutes of warm-up cycling at

a light intensity (60 W) at the same self-selected cadence as before. Thereafter, a ramp test

beginning at 75 W commenced, with power output increasing 1 W every 2.4 seconds. This

test was terminated at volitional exhaustion, defined as cadence falling >10 rpm for five

seconds. Strong verbal encouragement was provided to participants throughout. A final blood

lactate sample was drawn immediately after volitional exertion. The greatest 30 second

average $\dot{V}O_2$ value was used to quantify $\dot{V}O_{2peak}$, whilst the final power output was used to

quantify maximal ramp test power (Pmax).

Visits 2 & 3: Square-Wave Exercise Bouts

Visits two and three were identical, and performed with a minimum of 24 h between visits. The

visits involved continuous sampling of pulmonary gas exchange and near infrared

spectroscopy (NIRS) of the *vastus lateralis*. Trials commenced with participants performing

three minutes of unloaded pedalling on the cycle ergometer. Thereafter, participants

performed three repetitions of six-minute cycling bouts at 80% of LT (moderate intensity

exercise), interspersed with six minutes of unloaded pedalling. Following this, participants

cycled for 30 minutes at a work rate 30% between the LT and $\dot{V}O_{2peak}$ (30%Δ, heavy intensity

exercise). Exercise intensity was altered abruptly in a 'square-wave' fashion for each

repetition.

Immediately following all exercise, participants were laid supine on a physiotherapy table, and

an automatic personalised tourniquet system for blood flow restriction (Delfi Medical

Innovations Inc., Vancouver BC, Canada) was placed around the thigh, proximal to the NIRS

optode, and inflated for five minutes at 120% of limb occlusion pressure, in order to occlude

blood flow. This system automatically measures limb occlusion pressure, defined as the

minimum pressure required for complete restriction of arterial blood flow in a limb, and

maintains the pressure during inflation to ensure consistent occlusion. Following this, pressure

in the cuff was released and the hyperaemic response measured (see *Near Infrared*

Spectroscopy). This protocol is termed 'physiological calibration' (Barstow, 2019), and allows

all NIRS data to be expressed as a % of an individual's physiological minimum and maximum

values. Physiological calibration negates any potential influence of adipose tissue thickness

on NIRS data (Ryan *et al.*, 2012).

Pulmonary Gas Exchange

During all visits, expired gas was analysed breath-by-breath using an online system (Vyntus

CPX, Jaeger, CareFusion, Germany). Oxygen (O_2) and carbon dioxide (CO_2) concentrations

were quantified via a paramagnetic chemical fuel cell and non-dispersive infrared cell

respectively. Before each test, the analysers were calibrated using ambient air and a gas of

known O_2 (14.00%) and CO_2 (4.97%) concentrations. Ventilatory volumes were inferred from

measurement of gas flow using a digital turbine transducer (volume 0 to 10 L, resolution 3 mL,

flow 0 to 15 $L \cdot s^{-1}$) and calibrated prior to each test (Hans Rudolph Inc. Kansas City, USA).

Near Infrared Spectroscopy

A multi-distance, continuous-wave, single channel NIRS (NIRO-200NX, Hamamatsu) was

used to evaluate changes in vastus lateralis muscle deoxyhaemoglobin and myoglobin

(HHb+Mb) concentrations, sampled at a rate of 5 Hz. The light-emitting probe comprises of

light emitting diodes operating at three wavelengths (735, 810, and 850 nm). The probe was

placed on the *vastus lateralis*, 20 cm above the fibular head. Optodes were held in place by

an elasticised bandage and covered by an opaque, dark material to avoid motion and ambient

light influences.

Data Analysis

$\dot{V}O_2$ *Kinetics*

The breath by breath data was manually filtered to remove outlying breaths, defined as breaths

deviating more than 500 $ml.min^{-1}$ from the mean value from the preceding five breaths.

Thereafter, breath by breath data was linearly interpolated to provide second-by-second

values. The multiple repetitions of the square-wave exercise bouts were then averaged, and

246 $\dot{V}O_2$ responses were time aligned to the onset of exercise. Data from the onset of the transition

247 to 20 s was removed, then the resultant data was modelled with a monoexponential curve,

248 including data from -60 to 360 seconds (moderate) or -60 to 120 seconds (heavy), with the

249 following equation:

$$\dot{V}O_2(t) = \dot{V}O_2(b) + Ap(1 - e^{-(\tau-TDp)/\tau p})$$

252 Where $\dot{V}O_2(t)$ is the $\dot{V}O_2$ at time t; $\dot{V}O_2(b)$ is the baseline $\dot{V}O_2$ measured in the 60 s preceding

253 the transition in work rate; and A_p, TD_p, and τ_p are the amplitude, time delay, and the time

254 constant of the phase II response, respectively (Rossiter *et al.*, 2001). For exercise in the

255 heavy intensity domain, the amplitude of the $\dot{V}O_2$ slow component was determined by

256 subtracting the phase II amplitude from the highest 30 s average of $\dot{V}O_2$ during the 30 min

257 bout. To facilitate comparisons between sexes, amplitudes were also normalised to each

258 individual's $\dot{V}O_{2peak}$ as well as being presented in L.min^{-1}.

Figure 1: Visualisation of the monoexponential curve fitting procedures for a representative participant's data in the moderate intensity domain. Panel A describes the $\dot{V}O_2$ data (1 Hz) and Panel B describes the HHb+Mb data (0.2 Hz).

Deoxyhaemoglobin Kinetics

Prior to curve fitting, data from NIRS were normalised to the minimum values during, and the

maximum values following the five minute arterial occlusion (Ryan *et al.*, 2012), then averaged

into 1 s and 5 s bins. The multiple repetitions of the square-wave exercise bouts were then

averaged, and HHb+Mb responses were time aligned to the onset of exercise. The TD for the

HHb+Mb response was determined using the 1 s averaged data as the time between exercise

onset and the first point at which HHb+Mb signal started to systematically increase. This was

performed for each transition individually, with all TDs averaged to provide a single value. The

5 s averaged data was then modelled with a monoexponential curve in the same manner as

$\dot{V}O_2$ data, including data up to 90 s after the transition (Murias *et al.*, 2010).

Statistical Analysis

Data are presented as mean ± SD within the text and figures. Normal distribution of data was

confirmed with the Shapiro-Wilk test. As all variables had normally distributed data, males and

females were compared with independent samples t tests. The significance level for all

statistical tests was set at $p < 0.05$.

 Results

Incremental Exercise Testing

Anthropometric data and outcome variables from the two incremental exercise tests
performed in the first visit are presented in Table 1. As expected, males had a greater stature
and body mass than females ($p \leq 0.006$) as well as a greater absolute $\dot{V}O_{2peak}$ (mean
difference: 39%, $p = 0.002$). However, when $\dot{V}O_{2peak}$ was expressed relative to body mass, no
sex difference was observed (mean difference: 14%, $p = 0.111$). Males also exercised at
greater power outputs than females, with Pmax and LT being ~30% greater in males ($p \leq$
0.023), however when LT was expressed as a % of Pmax, no sex difference was observed (p
= 0.373). This resulted in the power outputs for the moderate and heavy intensity bouts being
greater in males compared to females ($p \leq 0.023$).

Table 1: Anthropometric data and outcome variables from incremental exercise testing.

	Males (n = 8)	Females (n = 8)	P value
Age (years)	27 ± 3	27 ± 7	0.734
Stature (cm)	182 ± 5	163 ± 4	**< 0.001**
Body Mass (kg)	75.3 ± 10.2	61.8 ± 5.9	**0.006**
$\dot{V}O_{2peak}$ (L.min^{-1})	3.47 ± 0.58	2.50 ± 4.42	**0.002**
Relative $\dot{V}O_{2peak}$ (ml.kg^{-1}.min^{-1})	46.2 ± 6.6	40.5 ± 6.7	0.111
Pmax (W)	328 ± 54	236 ± 43	**0.002**
Power at LT (W)	153 ± 25	119 ± 29	**0.023**
LT (% Pmax)	47 ± 6	50 ± 7	0.373
80% LT (W)	123 ± 20	95 ± 23	**0.023**
30% Δ (W)	206 ± 30	154 ± 32	**0.005**

LT: lactate threshold, Pmax: Maximal ramp test power output, $\dot{V}O_{2peak}$: Maximal rate of oxygen consumption.

$\dot{V}O_2$ Kinetics

The transition from unloaded pedalling to moderate and heavy intensity cycling elicited an

increase in $\dot{V}O_2$ (see Figure 2), and the monoexponential curve used to describe the

increase in $\dot{V}O_2$ in males and females demonstrated excellent r^2 values (see Table 2).

Table 2: Data from the monoexponential modelling of $\dot{V}O_2$ kinetics during moderate and heavy
intensity transitions.

	Males (n = 8)	Females (n = 8)	P value
Moderate Intensity Domain			
TD (s)	16.6 ± 4.1	17.9 ± 4.9	0.575
Baseline $\dot{V}O_2$ (L.min^{-1})	0.97 ± 0.05	0.77 ± 0.09	**< 0.001**
Amplitude (L.min^{-1})	0.83 ± 0.19	0.60 ± 0.18	**0.024**
Amplitude (% $\dot{V}O_{2peak}$)	24 ± 3	24 ± 5	0.949
τ (s)	27.9 ± 7.5	24.8 ± 6.6	0.385
r^2	0.965 ± 0.032	0.947 ± 0.034	
Heavy Intensity Domain			
TD (s)	15.0 ± 4.8	14.5 ± 4.5	0.858
Baseline $\dot{V}O_2$ (L.min^{-1})	1.00 ± 0.06	0.85 ± 0.11	**0.005**
Amplitude (L.min^{-1})	1.50 ± 0.38	0.96 ± 0.25	**0.005**
Amplitude (% $\dot{V}O_{2peak}$)	43 ± 5	38 ± 7	0.179
τ (s)	28.8 ± 7.9	27.2 ± 4.4	0.633
r^2	0.961 ± 0.031	0.927 ± 0.030	
SC Amplitude (L.min^{-1})	0.38 ± 0.16	0.28 ± 0.12	0.158
SC Amplitude (% $\dot{V}O_{2peak}$)	11.9 ± 6.9	10.8 ± 3.2	0.686

SC: slow component, τ: time constant, TD: time delay, $\dot{V}O_2$: rate of oxygen consumption

Figure 2: Group mean V̇O₂ data from males (blue) and females (red) during moderate (Panel A) and
heavy (Panel B) intensity transitions. The bold lines represent group means and the thin lines represent
standard deviation.

In absolute units (L.min^{-1}) males experienced greater rises in V̇O$_2$ during the phase II kinetics
($p \leq 0.024$), however when this was made relative to the individual (% V̇O$_{2peak}$) no sex
differences were observed ($p \geq 0.179$). Similarly, the V̇O$_2$ slow component amplitude was
similar between sexes in relative units ($p = 0.686$). As visualised in Figure 1, there were no
sex differences in τV̇O$_2$ in either the moderate ($p = 0.385$) or heavy ($p = 0.633$) intensity
domains.

Deoxyhaemoglobin Kinetics

The transition from unloaded pedalling to moderate and heavy intensity cycling elicited an
increase in HHb+Mb concentration (Figure 3), and the monoexponential curve used to
describe the increase in HHb+Mb in males and females demonstrated excellent r^2 values
(Table 3). One female's data had to be removed due to issues with the NIRS signal, resulting
in n = 7 females being used for NIRS analyses.

	Males (n = 8)	**Females (n = 7)**	*P value*
Moderate Intensity Domain			
TD (s)	8.7 ± 1.5	9.6 ± 2.6	0.447
Baseline HHb+Mb (% ischemia)	23 ± 8	29 ± 7	0.186
Amplitude (% ischemia)	21 ± 7	17 ± 5	0.225
τ (s)	8.1 ± 2.8	10.0 ± 3.8	0.274
r^2	0.974 ± 0.022	0.956 ± 0.037	
Heavy Intensity Domain			
TD (s)	6.2 ± 2.8	7.0 ± 2.3	0.582
Baseline HHb+Mb (% ischemia)	23 ± 6	17 ± 7	0.146
Amplitude (% ischemia)	36 ± 9	30 ± 8	0.193
τ (s)	11.3 ± 3.7	12.2 ± 4.0	0.665
r^2	0.983 ± 0.011	0.979 ± 0.023	
SC Amplitude (% ischemia)	14 ± 5	16 ± 4	0.432

HHb+Mb: deoxyhaemoglobin, SC: slow component, τ: time constant, TD: time delay consumption

As shown in Figure 3, the phase II amplitude of HHb+Mb increase in both intensity domains
was similar between sexes ($p \geq 0.193$). Similarly, there was no sex difference in the amplitude
of the slow component in the heavy intensity domain ($p = 0.432$). The time constant for phase
II HHb+Mb kinetics (τHHb+Mb) was also similar in males and females in both intensity
domains ($p \geq 0.274$).

Figure 3: Group mean HHb+Mb data from males (blue) and females (red) during moderate (Panel A)
and heavy (Panel B) intensity transitions. The bold lines represent group means and the thin lines
represent standard deviation.

Discussion

This study aimed to compare the kinetics of $\dot{V}O_2$ and HHb+Mb during moderate and heavy intensity exercise in males and females. In contrast to the hypothesis, at the onset of exercise the phase II time constants (τ) for $\dot{V}O_2$ and HHb+Mb were not different between the sexes, implying that both males and females were able to increase oxidative phosphorylation at comparable rates. In absolute units, males had larger amplitude increases than females, however when normalised to the individuals' maximum values, the rise in $\dot{V}O_2$ and HHb+Mb were also equivalent between the sexes. Combined, these data demonstrate that the bioenergetic response to exercise is similar between sexes, which has implications for exercise prescription and provides mechanistic insight into previously observed sex differences in the integrative response to exercise.

Previous literature investigating sex differences in the onset kinetics of oxygen transport and utilisation conflicts with the present data, with Beltrame *et al.* (2017) demonstrating quicker $\tau\dot{V}O_2$ and τHHb+Mb in females compared to males. One potential explanation for this discrepancy could be that Beltrame *et al.* utilised a treadmill walking task, compared to cycling. In tasks where O_2 delivery is not a limiting factor, females often outperform males. For instance, Ansdell *et al.* (2019) showed female knee-extensors had a greater relative critical torque than males during single-limb exercise. Whereas during cycling, where O_2 delivery is a determinant of critical power (Goulding & Marwood, 2023), this metabolic threshold was not different between sexes (Ansdell *et al.*, 2020a). Whilst consensus on whether O_2 delivery does (Hughson *et al.*, 2001) or does not (Grassi, 2001) limit $\tau\dot{V}O_2$ has not been reached, it is conceivable that during tasks where O_2 delivery and utilisation are both determinants in the metabolic response to exercise, the superior female skeletal muscle oxidative capacity (Cardinale *et al.*, 2018) is counteracted by an inferior O_2 carrying capacity (Murphy, 2014). Within the present data, this balance manifests as a comparable $\tau\dot{V}O_2$ in males and females, which agrees with data from do Nascimento Salvador *et al.* (2019), who demonstrated no sex difference in $\tau\dot{V}O_2$ during a transition from unloaded pedalling to 'very heavy' (60% Δ) cycling exercise.

Data from incremental exercise suggests that the poorer O_2 delivery in females results in a greater degree of O_2 extraction to compensate (Murias *et al.*, 2013). The present data contradicts this notion, as the amplitude of phase II HHb+Mb kinetics was similar between sexes (see Table 3). However, it is important to note that Murias *et al.* noted that this sex difference only occurred once incremental exercise exceeded the respiratory compensation point (i.e., the severe intensity domain), whereas the present study compared sexes in the

moderate and heavy intensity domains. The lack of a sex difference in the phase II amplitude

for HHb+Mb kinetics contradicts previously published NIRS data that demonstrated a smaller

rise in HHb+Mb in females compared to males during constant-load exercise (Ansdell *et al.*,

2020a). The crucial difference in methodologies employed between the previous study and

the present study is the application of a 'physiological calibration' to negate the influence of

adipose tissue thickness on NIRS signals (Ryan *et al.*, 2012). Previously, the sex difference

in the rise in HHb+Mb was suggested to reflect a lower oxygen cost of muscle contraction in

female knee-extensors, however the present data, with more rigorous methodologies

employed, refutes this.

The sex difference in muscle fibre type, whereby females demonstrate a greater proportional

area of type I fibres, appears to have not influenced either $\dot{V}O_2$ or HHb+Mb onset kinetics in

the present study. This would concur with data from Barstow *et al.* (1996), who demonstrated

no relationship between type I fibre percentage of the *vastus lateralis* and the time constant of

phase II $\dot{V}O_2$ kinetics. In contrast, Pringle *et al.* (2003) observed a negative correlation

between type I fibre percentage and the phase II time constant in the heavy intensity domain

only. Of note is that Pringle *et al.* included a wide range of participants with ~27 – 85% type I

fibres, and when groups were split into discrete groups of low and high fibre type percentages

(mean difference: 25%), the high percentage group had faster phase II kinetics. The present

study was not able to quantify the sex difference in muscle fibre typology, however, previous

literature has observed a 5-13% difference in type I fibre percentage of the *vastus lateralis*

(Simoneau & Bouchard, 1989; Staron *et al.*, 2000; Roepstorff *et al.*, 2006). Therefore, it could

be the case that the sex difference in muscle fibre typology is not large enough to affect the

phase II $\dot{V}O_2$ or HHb+Mb kinetics. Muscle fibre typology has previously been demonstrated to

affect the amplitude of the slow component within the heavy intensity domain, as individuals

with a lower type I fibre percentage experience larger rises in $\dot{V}O_2$ during constant-load

exercise (Barstow *et al.*, 1996; Pringle *et al.*, 2003). It is suggested that the slow component

is mechanistically underpinned by factors such as additional motor unit recruitment (Poole *et*

al., 1994; Burnley *et al.*, 2002) to compensate for fatigue-related changes in muscle

metabolism. For instance, muscle PCr stores demonstrate a similar slow component in

depletion during heavy intensity exercise (Rossiter *et al.*, 2002). Given that female knee-

extensors appears more fatigue-resistant and demonstrate lesser rises in the amplitude of

surface electromyography during constant-load exercise (Ansdell *et al.*, 2017; Ansdell *et al.*,

2019; Ansdell *et al.*, 2020a), we hypothesised that the relative amplitude of the slow

component would be greater in males to reflect a greater rate of metabolic disturbance.

However, as is evident in Tables 2 and 3, no sex difference was observed in the relative slow

component amplitude, implying that there was no difference in the metabolic response to constant-load exercise.

The lack of sex differences in either the phase II kinetics or slow component amplitude collectively suggest that the bioenergetic response to exercise was similar in males and females. Data on this topic is sparse, and due to the nature of methods such as magnetic resonance spectroscopy (MRS), limited to single-joint, isometric muscle contractions. Previous literature using this technique to study muscle metabolic changes during a 60 s contraction of the dorsiflexors showed no sex difference in changes in PCr, Pi, or pH (Russ *et al.*, 2005). Data from muscle biopsies of the *vastus lateralis* taken before and after repeated 30s cycling sprints suggested a greater preservation of ATP concentrations in females across a ~60 minute protocol (Esbjörnsson-Liljedahl *et al.*, 2002); however the authors suggested that this was likely a result of sex differences in the 20 minute recovery periods, rather that metabolic differences during exercise. Accordingly, the same group observed no sex differences in the metabolic response to a single 30-second cycling sprint (Esbjörnsson-Liljedahl *et al.*, 1999). Collectively, across multiple tasks and methodologies (MRS, biopsy and $\dot{V}O_2$ kinetics), the data suggest that there is no sex difference in the bioenergetic response to high-intensity exercise. This information provides mechanistic insight into the sex differences in the integrative response to exercise. For instance, sex differences in fatigability have partly been attributed to a lesser accumulation of fatiguing metabolites (Hunter, 2014; Ansdell *et al.*, 2020b). It is perhaps more accurate to suggest that previously observed sex differences in fatigue during intensity-matched exercise (Ansdell *et al.*, 2020a; Azevedo *et al.*, 2021) are more likely due to a greater fatigue-resistance of female muscle contractile apparatus, which experience similar degrees of metabolic stress as males. It is established that males and females differ in contractile properties such as calcium (Ca^{2+}) kinetics of the sarcoplasmic reticulum (Harmer *et al.*, 2014), with lower Ca^{2+}ATPase activity thought to permit a more fatigue-resistant skeletal muscle profile during equivalent exercise tasks (Hunter, 2014). Therefore, the present study advances the contemporary understanding of sex differences in the integrative response to exercise and provides mechanistic insight into previously observed phenomena.

These data have applications across the spectrum of health and disease. For example, those prescribing steady state exercise to improve skeletal muscle performance in athletes or patients might not need to account for the sex of their participants (Gloeckl *et al.*, 2022; Furrer *et al.*, 2023). This statement should however, be caveated by the fact that evidence regarding the influence of sex on long-term adaptation to exercise is sparse (Ansdell *et al.*, 2020b). Indeed, one area for further exploration is the bioenergetic response to exercise within the

441   severe intensity domain, where sex differences in fatigability have previously been observed
442   (Ansdell *et al.*, 2020a; Azevedo *et al.*, 2021). The employment of complementary techniques
443   to quantify O_2 delivery and muscle fibre typology could also provide greater insight into the
444   influence of sex on the O_2 cascade in a variety of tasks.
445

Conclusion
The present study aimed to compare the oxygen extraction and uptake kinetics during
moderate and heavy intensity cycling exercise. Contrary to our hypotheses, no sex differences
were observed in either the phase II or slow component kinetics for $\dot{V}O_2$ or HHb+Mb. The lack
of sex difference implies that males and females experience similar bioenergetic responses to
exercise, which provides mechanistic insight into previously observed phenomena such as
the sex difference in fatigability. Furthermore, based on these data and others demonstrating
no hormonal influences (Mattu *et al.*, 2020), we suggest that there is no rationale for the
exclusion of female participants in research investigating cardiopulmonary responses to
exercise.
456

Acknowledgements

The authors wish to thank the participants for their time and effort, as well as the technical staff within the Department of Sport, Exercise and Rehabilitation for their support.

Funding

This project was supported by a Physiological Society Research Springboard Studentship awarded to MSP, as well as Erasmus+ funding awarded to LB (2020-1-DE01-KA103-005569).

Disclosures

There are no conflicts of interest, financial or otherwise.

Author Contributions

MSP, ZW, MB, and PA conceived and designed the research. MSP, LB, EH, LH, and PA performed the experiments. LB, EH, LH, ZW, and PA analysed data. MSP, LB, EH, LH, ZW, MB, and PA interpreted results of the experiments. PA drafted the manuscript. MSP, LB, EH, LH, ZW, MB, and PA edited and revised the manuscript. All authors approved the final version of the manuscript.

Reference List

Allen DG, Lamb GD & Westerblad H. (2008). Skeletal muscle fatigue: cellular mechanisms. *Physiol Rev* **88,** 287-332.

Ansdell P, Brownstein CG, Škarabot J, Hicks KM, Howatson G, Thomas K, Hunter SK & Goodall S. (2019). Sex differences in fatigability and recovery relative to the intensity–duration relationship. *The Journal of Physiology* **597,** 5577-5595.

Ansdell P, Škarabot J, Atkinson E, Corden S, Tygart A, Hicks KM, Thomas K, Hunter SK, Howatson G & Goodall S. (2020a). Sex differences in fatigability following exercise normalised to the power–duration relationship. *The Journal of Physiology* **598,** 5717-5737.

Ansdell P, Thomas K, Hicks KM, Hunter SK, Howatson G & Goodall S. (2020b). Physiological sex differences affect the integrative response to exercise: acute and chronic implications. *Experimental Physiology* **105,** 2007-2021.

Ansdell P, Thomas K, Howatson G, Hunter S & Goodall S. (2017). Contraction intensity and sex differences in knee-extensor fatigability. *Journal of Electromyography and Kinesiology* **37,** 68-74.

Azevedo RdA, Forot J, Iannetta D, MacInnis MJ, Millet GY & Murias JM. (2021). Slight power output manipulations around the maximal lactate steady state have a similar impact on fatigue in females and males. *Journal of Applied Physiology* **130,** 1879-1892.

Barstow TJ. (2019). Understanding near infrared spectroscopy and its application to skeletal muscle research. *Journal of Applied Physiology* **126,** 1360-1376.

Barstow TJ, Jones AM, Nguyen PH & Casaburi R. (1996). Influence of muscle fiber type and pedal frequency on oxygen uptake kinetics of heavy exercise. *Journal of Applied Physiology* **81,** 1642-1650.

Beltrame T, Villar R & Hughson RL. (2017). Sex differences in the oxygen delivery, extraction, and uptake during moderate-walking exercise transition. *Applied Physiology, Nutrition, and Metabolism* **42,** 994-1000.

Burnley M, Doust JH, Ball D & Jones AM. (2002). Effects of prior heavy exercise on $\dot{V}O_2$ kinetics during heavy exercise are related to changes in muscle activity. *Journal of Applied Physiology* **93,** 167-174.

Burnley M & Jones AM. (2007). Oxygen uptake kinetics as a determinant of sports performance. *European Journal of Sport Science* **7,** 63-79.

Cardinale DA, Larsen FJ, Schiffer TA, Morales-Alamo D, Ekblom B, Calbet JAL, Holmberg HC & Boushel R. (2018). Superior Intrinsic Mitochondrial Respiration in Women Than in Men. *Front Physiol* **9,** 1133.

do Nascimento Salvador PC, Schäfer L, Grassi B, Guglielmo LGA & Denadai BS. (2019). Changes in VO2 Kinetics After Elevated Baseline Do Not Necessarily Reflect Alterations in Muscle Force Production in Both Sexes. *Frontiers in Physiology* **10**.

Ducrocq GP, Hureau TJ, Bøgseth T, Meste O & Blain GM. (2021). Recovery from Fatigue after Cycling Time Trials in Elite Endurance Athletes. *Med Sci Sports Exerc* **53,** 904-917.

Esbjörnsson-Liljedahl M, Bodin K & Jansson E. (2002). Smaller muscle ATP reduction in women than in men by repeated bouts of sprint exercise. *Journal of Applied Physiology* **93,** 1075-1083.

Esbjörnsson-Liljedahl M, Sundberg CJ, Norman B & Jansson E. (1999). Metabolic response in type I and type II muscle fibers during a 30-s cycle sprint in men and women. *Journal of Applied Physiology* **87,** 1326-1332.

Furrer R, Hawley JA & Handschin C. (2023). The molecular athlete: exercise physiology from mechanisms to medals. *Physiological Reviews* **103,** 1693-1787.

Gloeckl R, Zwick RH, Fürlinger U, Jarosch I, Schneeberger T, Leitl D, Koczulla AR, Vonbank K, Alexiou C, Vogiatzis I & Spruit MA. (2022). Prescribing and adjusting exercise

training in chronic respiratory diseases – Expert-based practical recommendations. *Pulmonology*.

Goulding RP & Marwood S. (2023). Interaction of Factors Determining Critical Power. *Sports Medicine* **53,** 595-613.

Goulding RP, Rossiter HB, Marwood S & Ferguson C. (2021). Bioenergetic Mechanisms Linking V̇O2 Kinetics and Exercise Tolerance. *Exercise and Sport Sciences Reviews* **49**.

Grassi B. (2001). Regulation of Oxygen Consumption at Exercise Onset: Is It Really Controversial? *Exercise and Sport Sciences Reviews* **29,** 134-138.

Grassi B, Poole DC, Richardson RS, Knight DR, Erickson BK & Wagner PD. (1996). Muscle O2 uptake kinetics in humans: implications for metabolic control. *Journal of Applied Physiology* **80,** 988-998.

Grassi B, Rossiter HB & Zoladz JA. (2015). Skeletal muscle fatigue and decreased efficiency: two sides of the same coin? *Exerc Sport Sci Rev* **43,** 75-83.

Harmer AR, Ruell PA, Hunter SK, McKenna MJ, Thom JM, Chisholm DJ & Flack JR. (2014). Effects of type 1 diabetes, sprint training and sex on skeletal muscle sarcoplasmic reticulum Ca2+ uptake and Ca2+-ATPase activity. *The Journal of Physiology* **592,** 523-535.

Harms CA, McClaran SR, Nickele GA, Pegelow DF, Nelson WB & Dempsey JA. (1998). Exercise-induced arterial hypoxaemia in healthy young women. *J Physiol* **507 (Pt 2),** 619-628.

Hughson RL, Tschakovsky ME & Houston ME. (2001). Regulation of Oxygen Consumption at the Onset of Exercise. *Exercise and Sport Sciences Reviews* **29,** 129-133.

Hunter SK. (2014). Sex differences in human fatigability: mechanisms and insight to physiological responses. *Acta Physiologica* **210,** 768-789.

Koppo K, Bouckaert J & Jones AM. (2004). Effects of Training Status and Exercise Intensity on Phase II V̇O2 Kinetics. *Medicine & Science in Sports & Exercise* **36,** 225-232.

Krustrup P, Söderlund K, Mohr M, González-Alonso J & Bangsbo J. (2004). Recruitment of fibre types and quadriceps muscle portions during repeated, intense knee-extensor exercise in humans. *Pflugers Arch* **449,** 56-65.

Mattu AT, Iannetta D, MacInnis MJ, Doyle-Baker PK & Murias JM. (2020). Menstrual and oral contraceptive cycle phases do not affect submaximal and maximal exercise responses. *Scandinavian Journal of Medicine & Science in Sports* **30,** 472-484.

Murias JM, Keir DA, Spencer MD & Paterson DH. (2013). Sex-related differences in muscle
deoxygenation during ramp incremental exercise. *Respiratory Physiology &*
Neurobiology **189,** 530-536.

Murias JM, Kowalchuk JM & Paterson DH. (2010). Speeding of V̇o2 kinetics with endurance
training in old and young men is associated with improved matching of local O2
delivery to muscle O2 utilization. *Journal of Applied Physiology* **108,** 913-922.

Murphy WG. (2014). The sex difference in haemoglobin levels in adults — Mechanisms,
causes, and consequences. *Blood Reviews* **28,** 41-47.

Poole DC, Barstow TJ, Gaesser GA, Willis WT & Whipp BJ. (1994). VO2 slow component:
physiological and functional significance. *Med Sci Sports Exerc* **26,** 1354-1358.

Poole DC & Jones AM. (2012). Oxygen Uptake Kinetics. *Compr Physiol* **2,** 933-996.

Pringle JSM, Doust JH, Carter H, Tolfrey K, Campbell IT & Jones AM. (2003). Oxygen uptake
kinetics during moderate, heavy and severe intensity 'submaximal' exercise in
humans: the influence of muscle fibre type and capillarisation. *European Journal of*
Applied Physiology **89,** 289-300.

Roepstorff C, Thiele M, Hillig T, Pilegaard H, Richter EA, Wojtaszewski JF & Kiens B. (2006).
Higher skeletal muscle alpha2AMPK activation and lower energy charge and fat
oxidation in men than in women during submaximal exercise. *J Physiol* **574,** 125-138.

Rossiter HB, Ward SA, Howe FA, Kowalchuk JM, Griffiths JR & Whipp BJ. (2002). Dynamics
of intramuscular 31P-MRS Pi peak splitting and the slow components of PCr and O2
uptake during exercise. *Journal of Applied Physiology* **93,** 2059-2069.

Rossiter HB, Ward SA, Kowalchuk JM, Howe FA, Griffiths JR & Whipp BJ. (2001). Effects of
prior exercise on oxygen uptake and phosphocreatine kinetics during high-intensity
knee-extension exercise in humans. *J Physiol* **537,** 291-303.

Russ DW, Lanza IR, Rothman D & Kent-Braun JA. (2005). Sex differences in glycolysis during
brief, intense isometric contractions. *Muscle & Nerve* **32,** 647-655.

Ryan TE, Erickson ML, Brizendine JT, Young H-J & McCully KK. (2012). Noninvasive
evaluation of skeletal muscle mitochondrial capacity with near-infrared spectroscopy:
correcting for blood volume changes. *Journal of Applied Physiology* **113,** 175-183.

Schiaffino S & Reggiani C. (2011). Fiber types in mammalian skeletal muscles. *Physiol Rev*
91, 1447-1531.

Simoneau JA & Bouchard C. (1989). Human variation in skeletal muscle fiber-type proportion
and enzyme activities. *American Journal of Physiology-Endocrinology and Metabolism*
257, E567-E572.

636
637 Staron RS, Hagerman FC, Hikida RS, Murray TF, Hostler DP, Crill MT, Ragg KE & Toma K.
638   (2000). Fiber type composition of the vastus lateralis muscle of young men and
639   women. *J Histochem Cytochem* **48,** 623-629.

640
641 Temesi J, Mattioni Maturana F, Peyrard A, Piucco T, Murias JM & Millet GY. (2017). The
642   relationship between oxygen uptake kinetics and neuromuscular fatigue in high-
643   intensity cycling exercise. *European Journal of Applied Physiology* **117,** 969-978.

644
645