No sex differences in oxygen uptake or extraction kinetics in the moderate or heavy exercise intensity domains

Maria Solleiro Pons[1], Lina Bernert[1,2], Emily Hume[1], Luke Hughes[1], Zander Williams[3], Mark Burnley[4], Paul Ansdell[1]

[1] Department of Sport, Exercise and Rehabilitation, Faculty of Health and Life Sciences, Northumbria University, UK

[2] Institute of Sport and Exercise Sciences, University of Münster, Germany

[3] Department of Respiratory Medicine, Royal Brompton Hospital, UK

[4] School of Sport, Health and Exercise Sciences, Loughborough University, UK

Running Title: The influence of sex on oxygen uptake and extraction

Corresponding Author:

Paul Ansdell, PhD

Department of Sport, Exercise and Rehabilitation

Faculty of Health and Life Sciences

Northumbria University

Newcastle upon Tyne

NE1 8ST

UK

p.ansdell@northumbria.ac.uk

Abstract

The integrative response to exercise differs between sexes, with oxidative energy contribution purported as a potential mechanism. The present study investigated whether this difference was evident in the kinetics of oxygen uptake ($\dot{V}O_2$) and extraction (HHb+Mb) during exercise.

Sixteen adults (8 males, 8 females, age: 27±5 years) completed three experimental visits. Incremental exercise testing was performed to obtain lactate threshold and $\dot{V}O_{2peak}$. Subsequent visits involved three six-minute cycling bouts at 80% of lactate threshold and one 30-minute bout at a work rate 30% between the lactate threshold and power at $\dot{V}O_{2peak}$. Pulmonary gas exchange and near-infrared spectroscopy of the vastus lateralis were used to continuously sample $\dot{V}O_2$ and HHb+Mb, respectively. The phase II $\dot{V}O_2$ kinetics were quantified using mono-exponential curves during moderate and heavy exercise. Slow component amplitudes were also quantified for the heavy intensity domain.

Relative $\dot{V}O_{2peak}$ values were not different between sexes ($p=0.111$). Males achieved ~30% greater power outputs ($p=0.002$). In the moderate and heavy intensity domains, the relative amplitude of the phase II transition was not different between sexes for $\dot{V}O_2$ (~24 and ~40% $\dot{V}O_{2peak}$, $p \geq 0.179$) and HHb+Mb (~20 and ~32% ischemia, $p \geq 0.193$). Similarly, there were no sex differences in the time constants for $\dot{V}O_2$ (~28 s, $p \geq 0.385$) or HHb+Mb (~10s, $p \geq 0.274$). In the heavy intensity domain, neither $\dot{V}O_2$ ($p \geq 0.686$) or HHb+Mb ($p \geq 0.432$) slow component amplitudes were different between sexes.

The oxidative response to moderate and heavy intensity exercise did not differ between males and females, suggesting similar dynamic responses of oxidative metabolism during intensity-matched exercise.

New and Noteworthy

This study demonstrated no sex differences in the oxidative response to moderate and heavy intensity cycling exercise. The change in oxygen uptake and deoxyhaemoglobin were modelled with mono-exponential curve fitting, which revealed no differences in the rate of oxidative energy provision between sexes. This provides insight into previously reported sex differences in the integrative response to exercise.

Introduction

The transition from rest to exercise involves an integrated response from the pulmonary, cardiovascular, and muscular systems to rapidly increase the supply and utilisation of oxygen for oxidative adenosine triphosphate (ATP) provision (Poole & Jones, 2012). The speed at which this process can occur can be quantified using pulmonary oxygen uptake ($\dot{V}O_2$) kinetics, and is thought to determine metabolic stability and exercise tolerance across the spectrum of athletic performance and disease (Burnley & Jones, 2007; Grassi *et al.*, 2011). The $\dot{V}O_2$ response can be broken down into three phases, beginning with the initial cardio-dynamic phase (phase I, 10-20 s) which represents an increased venous return via the muscle pump effect, as well as increased pulmonary blood flow (Grassi *et al.*, 1996). Thereafter, increases in pulmonary $\dot{V}O_2$ are considered to reflect increased muscle oxygen uptake in response to exercise (phase II), until the energy demand of exercise is met by oxidative phosphorylation and $\dot{V}O_2$ reaches a steady state (Hughson *et al.*, 2001). A steady state response is attainable quickly within the moderate intensity domain, whereas in either heavy or severe intensity domains, a further rise in $\dot{V}O_2$ is observed before the steady state is attained (heavy) or $\dot{V}O_{2max}$ is reached (severe), termed the slow component. This three-phase response is ubiquitous in exercising humans; however, the biological characteristics of the individual can influence the rates at which they occur (for review see Poole & Jones, 2012).

The time constant of phase II kinetics are considered to be a crucial determinant of the decrease in contractile function experienced by the exercising individual (Temesi *et al.*, 2017; Goulding *et al.*, 2021), and could explain previously observed sex differences in the integrative response to exercise (Ansdell *et al.*, 2020b). As oxidative phosphorylation does not immediately meet the demand for ATP, substrate-level phosphorylation is required (Burnley & Jones, 2007). Intuitively, the rate at which oxidative metabolism can be upregulated at the onset of exercise is inversely linked with the accumulation of deleterious metabolites such as hydrogen ions [H^+] and inorganic phosphate [P_i], as well as the depletion of phosphocreatine [PCr] stores, which all interfere with excitation-contraction coupling (Allen *et al.*, 2008). However, the relationship between the aforementioned metabolites and the upregulation of oxidative phosphorylation is multi-faceted, with the progressive change in the phosphate energy state (i.e., the [ATP]/[ADP][P_i] balance) driving the rate at which mitochondrial respiration increases, whilst [H^+] accumulation concurrently inhibits anaerobic glycolysis at the onset of exercise (Tschakovsky & Hughson, 1999; Korzeniewski & Rossiter, 2015). Accordingly, Temesi *et al.* (2017) demonstrated a positive correlation between the time constant ($\tau\dot{V}O_2$) of the phase II response and the decrease in quadriceps potentiated twitch force. Similarly, elite endurance athletes demonstrate faster

$\tau \dot{V}O_2$ (Koppo *et al.*, 2004) and lesser declines in contractile function (Ducrocq *et al.*, 2021)
compared to untrained individuals when exercising at similar relative intensities.

A consistent finding in studies comparing males and females exercising at the same
metabolic intensity is that females experience a lesser degree of contractile impairment of
the knee-extensors (Ansdell *et al.*, 2019; Ansdell *et al.*, 2020a; Azevedo *et al.*, 2021).
Previously, this has been suggested to be a result of sex differences in skeletal muscle
composition, whereby females consistently demonstrate a greater proportional area of type I
fibres (Staron *et al.*, 2000; Roepstorff *et al.*, 2006). The consequences of this fibre type
difference are multi-factorial; for-instance, it is well established that type I fibres are more
fatigue-resistant (Schiaffino & Reggiani, 2011). Additionally, female *vastus lateralis* capillary
density is ~23% greater in females compared to males (Roepstorff *et al.*, 2006), while
females also demonstrate greater mitochondrial oxidative function and intrinsic respiratory
rates than males of equivalent training status (Cardinale *et al.*, 2018). One factor that
remains unexplored is whether these physiological sex differences result in differences in the
metabolic response to exercise. Conceivably, the superior aerobic phenotype of female
skeletal muscle could imply that females might be able to meet the ATP demand of exercise
through oxidative means faster than males. Despite this, *ex vivo* evidence suggests that
female skeletal muscle fibres have a lower ADP sensitivity of mitochondrial respiration
(Miotto *et al.*, 2018), which could result in a slower rate of oxidative phosphorylation at the
onset of exercise. Therefore, *in vivo* assessment of $\dot{V}O_2$ kinetics would provide insight into
the balance between these morphological and cellular sex differences.

The $\dot{V}O_2$ slow component is underpinned by different mechanisms to the phase II kinetics
and describes the increase in $\dot{V}O_2$ during constant-load exercise. This increase in $\dot{V}O_2$
implies an impairment of efficiency and is likely an amalgamation of several concurrent
physiological changes. Within skeletal muscle, the accumulation of metabolites (e.g., [Pi])
and associated contractile dysfunction is linked with the loss of efficiency (Grassi *et al.*,
2015). Of relevance here, is that female skeletal muscle has consistently been demonstrated
to be more fatigue-resistant (Ansdell *et al.*, 2019; Ansdell *et al.*, 2020a), and shows lesser
increases in the surface electromyogram in states of fatigue (Ansdell *et al.*, 2017; Ansdell *et*
al., 2019). Furthermore, it is well-established that females have a greater reliance on lipid
metabolism during sustained exercise at similar relative work rates (Cano *et al.*, 2022),
which could be related to the more oxidative phenotype of skeletal muscle, as described
above. Combined, these physiological sex differences could represent a slower loss of
efficiency during constant load exercise in females, however, this remains unexplored.

Despite more aerobically-suited skeletal muscle, females have lower levels of haemoglobin
(Murphy, 2014), which is thought to impair O_2 carrying capacity during exercise (Harms *et
al.*, 1998; Diaz-Canestro *et al.*, 2022). During exercise where O_2 delivery and utilisation are
both limiting factors (e.g., cycling, Goulding & Marwood, 2023), these factors are thought to
counteract each other to enable comparable relative metabolic thresholds (i.e., critical
power) between the sexes (Ansdell *et al.*, 2020b). To date, the only investigation to
systematically investigate the oxidative adjustment at the onset of exercise between sexes
did so during low intensity treadmill walking (Beltrame *et al.*, 2017). Data from this study
suggested faster O_2 extraction in females, quantified as the change in de-oxyhaemoglobin
and myoglobin (HHb+Mb) signal in near-infrared spectroscopy (NIRS), fitting with the notion
that phase II $V\dot{}O_2$ kinetics are influenced by intramuscular factors in healthy humans (Poole
& Jones, 2012). However, the demands of treadmill walking differ to those of high-intensity
cycling exercise, where it has recently been argued that all levels of the O_2 cascade are
considered to be influential in determining metabolic responses to exercise (Goulding &
Marwood, 2023). Thus, given the lack of evidence regarding physiological responses to
exercise in females (James *et al.*, 2023), it remains to be determined how sex differences in
convective and diffusive contributions to O_2 delivery mediate the rate of $V\dot{}O_2$ adjustment to
exercise.

Accordingly, the present study employed a multi-method approach of measuring pulmonary
gas exchange and near-infrared spectroscopy simultaneously to compare the kinetics of
pulmonary $V\dot{}O_2$ as well as muscle oxygen extraction in both sexes during moderate and
heavy intensity exercise. Previously this experimental approach has been used to obtain
information about O_2 delivery and utilisation to gain insights into the integrative response to
exercise (DeLorey *et al.*, 2003). It was hypothesised that females would demonstrate a
smaller value for the phase II time constant (i.e., faster kinetics) for $V\dot{}O_2$ and HHb+Mb at
the onset of exercise, and a smaller slow component amplitude in the heavy intensity
domain.

Methods

Ethical Approval

This study received institutional ethical approval from the Northumbria University Health and Life Sciences Research Ethics Committee (submission reference: 49189) and was conducted according to all aspects of the Declaration of Helsinki, apart from pre-registration in a database. Participants volunteered for the study and provided written informed consent.

Participants

Using the effect size for the sex difference in *vastus lateralis* tissue oxygenation during heavy intensity exercise ($\eta p^2 = 0.509$) from (Ansdell *et al.*, 2020a), an *a priori* sample size calculation determined a minimum of 14 participants (seven females and seven males) were required to detect an effect ($\alpha = 0.05$, power = 0.95). Therefore, eight males (mean ± SD age: 27 ± 3 years; stature: 182 ± 5 cm; body mass: 75.3 ± 10.2 kg) and eight females (mean ± SD age: 27 ± 7; stature: 163 ± 4 cm; body mass: 61.8 ± 5.9 kg) volunteered to take part in the study. Hormonal status was not an exclusion criterion or controlled for in this study. Female participants were tested in any phase of their menstrual cycle and there were no restrictions on hormonal contraceptive usage. This decision was based on evidence from (Mattu *et al.*, 2020) who demonstrated no hormonal effects on $\dot{V}O_2$ kinetics during cycling exercise. Of the eight females, two were using combined oral contraceptive pills (Lucette and Rigevidon) and six were naturally menstruating. All participants were free from cardiovascular, respiratory, and neurological disease as well as musculoskeletal injury.

Experimental Design

All participants visited the laboratory on three occasions across an average of 10 ± 4 days (range: 5 – 21 days). During the first visit, participants were familiarised with the experimental procedures and completed two incremental exercise tests to quantify lactate threshold and peak oxygen uptake ($\dot{V}O_{2peak}$). The second and third visits were identical and involved three six-minute bouts of moderate intensity exercise (80% of lactate threshold, LT), separated by six minutes of unloaded pedalling. Thereafter, a single bout of heavy intensity exercise (30% ΔLT – $\dot{V}O_{2peak}$) was performed for 30 minutes.

Visit 1: Familiarisation & Incremental Testing

The first visit began with participants completing a screening questionnaire to ensure inclusion criteria were met. Thereafter, participants moved onto the cycle ergometer (Velotron, SRAM, Chicago, IL, USA) which was set up with the seat height aligned with the hip, and handlebar height set according to the participants' comfort, these measurements

were recorded and replicated for subsequent trials. The breath-by-breath gas exchange mask was then placed over the participant's mouth and nose, and an air-tight seal was ensured before resting data was recorded. Following resting measures of pulmonary gas exchange and muscle oxygenation, participants completed five minutes of warm-up cycling at a light intensity (60 W) at a self-selected cadence between 70-100 rpm, before commencing an incremental exercise test. The first incremental exercise test began at 75 W and increased by 25 W every five minutes. At the end of each stage, a capillary blood sample was drawn from the participants' fingertip and immediately analysed to determine whole blood lactate concentration (mmol.L^{-1}, Biosen C-Line, EKF Diagnostics, Germany). The test was terminated once LT was identified as the first work rate at which a non-linear increase in blood lactate concentration was observed (Faude *et al.*, 2009), after which, participants were provided 20 minutes of passive rest.

Next, participants began the second incremental test with five minutes of warm-up cycling at a light intensity (60 W) at the same self-selected cadence as before. Thereafter, a ramp test beginning at 75 W commenced, with power output increasing 1 W every 2.4 seconds (25 W.min^{-1}). This test was terminated at volitional exhaustion, defined as cadence falling >10 rpm for five seconds. Strong verbal encouragement was provided to participants throughout. A final blood lactate sample was drawn immediately after volitional exertion. The greatest 30 second average V̇O$_2$ value was used to quantify V̇O$_{2peak}$, whilst the final power output was used to quantify maximal ramp test power (Pmax).

Visits 2 & 3: Square-Wave Exercise Bouts

Visits two and three were identical, and performed with a minimum of 24 h between visits. The visits involved continuous sampling of pulmonary gas exchange and near infrared spectroscopy (NIRS) of the *vastus lateralis*. Trials commenced with participants performing three minutes of unloaded pedalling on the cycle ergometer. Thereafter, participants performed three repetitions of six-minute cycling bouts at 80% of the work rate associated with LT (moderate intensity exercise), interspersed with six minutes of unloaded pedalling. Following this, participants cycled for 30 minutes at a work rate 30% between the LT and V̇O$_{2peak}$ (30%Δ, heavy intensity exercise). Throughout this visit, participants were asked to replicate their self-selected cadence from visit 1, which was monitored by an experimenter throughout. Exercise intensity was altered abruptly in a 'square-wave' fashion for each repetition.

Immediately following all exercise, a 'physiological calibration' of NIRS signals was performed as per recommendations from Barstow (2019). Participants were laid supine on a

physiotherapy table, with their leg placed horizontal, and an automatic personalised

tourniquet system for blood flow restriction (Delfi Medical Innovations Inc., Vancouver BC,

Canada) was placed around the thigh via a nylon cuff (11.5 cm☐×☐86 cm, 5 mm thick),

proximal to the NIRS optode. The cuff was inflated for five minutes at 120% of limb occlusion

pressure in order to occlude blood flow, and the mean pressure was not different between

males and females (120% limb occlusion pressure: 248 ± 20 vs. 249 ± 30 mmHg, p =

0.780). This system automatically measures limb occlusion pressure, defined as the

minimum pressure required for complete restriction of arterial blood flow in a limb, and

maintains the pressure during inflation to ensure consistent occlusion. Following this,

pressure in the cuff was released and the hyperaemic response measured (see *Near*

Infrared Spectroscopy). This protocol allowed all NIRS data to be expressed as a % of an

individual's physiological minimum and maximum values. Physiological calibration negates

any potential influence of adipose tissue thickness on NIRS data (Barstow, 2019).

Pulmonary Gas Exchange

During all visits, expired gas was analysed breath-by-breath using an online system (Vyntus

CPX, Jaeger, CareFusion, Germany). Oxygen (O_2) and carbon dioxide (CO_2) concentrations

were quantified via a paramagnetic chemical fuel cell and non-dispersive infrared cell

respectively. Before each test, the analysers were calibrated using ambient air and a gas of

known O_2 (15.00%) and CO_2 (4.97%) concentrations. Ventilatory volumes were inferred from

measurement of gas flow using a digital turbine transducer (volume 0 to 10 L, resolution 3

mL, flow 0 to 15 L·s^{-1}) and calibrated prior to each test (Hans Rudolph Inc. Kansas City,

USA).

Near Infrared Spectroscopy

A multi-distance, continuous-wave, single channel NIRS (NIRO-200NX, Hamamatsu) was

used to evaluate changes in vastus lateralis muscle deoxyhaemoglobin and myoglobin

(HHb+Mb), as well as oxyhaemoglobin and myoglobin (HbO_2+MbO_2) concentrations,

sampled at a rate of 5 Hz. Tissue oxygenation index (TOI) was calculated as HbO_2+MbO_2 ÷

[HbO_2+MbO2 + HHb+Mb] × 100. The light-emitting probe comprised of light emitting diodes

operating at three wavelengths (735, 810, and 850 nm). The probe was placed on the *vastus*

lateralis, 20 cm above the fibular head. Optodes were held in place by an elasticised

bandage and covered by an opaque, dark material to avoid motion and ambient light

influences.

Data Analysis

V☐O$_2$ Kinetics

The breath by breath data was manually filtered to remove outlying breaths, defined as
breaths deviating more than 500 ml.min^{-1} from the mean value from the preceding five
breaths. Thereafter, breath by breath data was linearly interpolated to provide second-by-
second values. The multiple repetitions of the square-wave exercise bouts were then
averaged, and $V\dot{}O_2$ responses were time aligned to the onset of exercise. Data from the
onset of the transition to 20 s was removed, then the resultant data was modelled with a
monoexponential curve, including data from -60 to 360 seconds (moderate) or -60 to 120
seconds (heavy), with the following equation:

$$V\dot{}O_2(t) = V\dot{}O_2(b) + Ap(1 - e^{-(\tau-TDp)/\tau p})$$

Where $V\dot{}O_2(t)$ is the $V\dot{}O_2$ at time t, $V\dot{}O_2(b)$ is the baseline $V\dot{}O_2$ measured in the 60 s
preceding the transition in work rate; and A_p, TD_p, and τ_p are the amplitude, time delay, and
the time constant of the phase II response, respectively. We chose to constrain the
modelling of heavy intensity domain onset kinetics to 120 seconds in order to minimise the
influence of the slow component, however this cannot be guaranteed (Burnley *et al.*, 2006).
For exercise in the heavy intensity domain, the amplitude of the $V\dot{}O_2$ slow component was
determined by subtracting the phase II amplitude from the highest 30 s average of $V\dot{}O_2$
during the 30 min bout (Rossiter *et al.*, 2001). To facilitate comparisons between sexes,
amplitudes were also normalised to each individual's $V\dot{}O_{2peak}$ and end-exercise $V\dot{}O_2$ as
well as being presented in L.min^{-1}. The O_2 cost of the transitions was estimated by
calculating the $V\dot{}O_2$ gain (Porcelli *et al.*, 2016), where the amplitude of the phase II
response was divided by work rate (ml.min^{-1}.W^{-1}).

Figure 1: Visualisation of the monoexponential curve fitting procedures for a representative participant's data in the moderate intensity domain. Panel O_2 data (1 Hz) and Panel B describes the HHb+Mb data (0.2 Hz).

Deoxyhaemoglobin Kinetics

Prior to curve fitting, data from NIRS were normalised to the minimum values during, and the

maximum values following the five minute arterial occlusion (Ryan *et al.*, 2012), then

averaged into 1 s and 5 s bins. The multiple repetitions of the square-wave exercise bouts

were then averaged, and HHb+Mb responses were time aligned to the onset of exercise.

The TD for the HHb+Mb response was determined using the 1 s averaged data as the time

between exercise onset and the first point at which HHb+Mb signal started to systematically

increase. This was performed for each transition individually, with all TDs averaged to

provide a single value. The 5 s averaged data was then modelled

309   O_2 data, including data up to 90 s after the transition

310 (Murias *et al.*, 2010). Other NIRS-derived variables (TOI and HbO_2+MbO_2) were quantified

311 as 30 s averages at the following time points: the 30s of unloaded pedalling immediately

312 prior to the transition, 90 s following the transition (heavy intensity domain only), and the final

313 30s of the square wave bout of exercise.

Statistical Analysis

Data are presented as mean ± SD within the text and figures. Normal distribution of data was confirmed with the Shapiro-Wilk test. As all variables had normally distributed data, males and females were compared with independent samples t tests for variables with a single value or time point. For repeated measures variables during exercise, two-way (sex × time) repeated measures ANOVA were performed, followed by Bonferroni-corrected post hoc tests if significant main effects were observed. Effect sizes for comparisons were calculated as Cohen's *d.* The significance level for all statistical tests was set at $p < 0.05$.

Results

Incremental Exercise Testing

Anthropometric data and outcome variables from the two incremental exercise tests

performed in the first visit are presented in Table 1. As expected, males had a greater

stature and body mass than females ($p \leq 0.006$) as well as a greater absolute V̇O$_{2peak}$

(mean difference: 39%, $p = 0.002$). However, when V̇O$_{2peak}$ was expressed relative to body

mass, no sex difference was observed (mean difference: 14%, $p = 0.111$). Males also

exercised at greater power outputs than females, with Pmax and LT being ~30% greater in

males ($p \leq 0.023$), however when LT was expressed as a % of Pmax, no sex difference was

observed ($p = 0.373$). This resulted in the power outputs for the moderate and heavy

intensity bouts being greater in males compared to females ($p \leq 0.023$).

Table 1: Anthropometric data and outcome variables from incremental exercise testing.

	Males (n = 8)	Females (n = 8)	P value	Cohen's d
Age (years)	27 ± 3	27 ± 7	0.734	0.117
Stature (cm)	182 ± 5	163 ± 4	**< 0.001**	**1.383**
Body Mass (kg)	75.3 ± 10.2	61.8 ± 5.9	**0.006**	**0.501**
V̇O$_{2peak}$ (L.min^{-1})	3.47 ± 0.58	2.50 ± 4.42	**0.002**	**0.627**
Relative V̇O$_{2peak}$ (ml.kg^{-1}.min^{-1})	46.2 ± 6.6	40.5 ± 6.7	0.111	0.326
Pmax (W)	328 ± 54	236 ± 43	**0.002**	**0.646**
Pmax (W.kg^{-1})	4.4 ± 0.8	3.8 ± 0.5	0.118	0.253
Power at LT (W)	153 ± 25	119 ± 29	**0.023**	**0.528**
LT (% Pmax)	47 ± 6	50 ± 7	0.373	0.183
LT (W.kg^{-1})	2.0 ± 0.2	1.9 ± 0.4	0.440	0.127
80% LT (W)	123 ± 20	95 ± 23	**0.023**	**0.528**
80% LT (W.kg^{-1})	1.6 ± 0.2	1.5 ± 0.3	0.440	0.110
30% Δ (W)	206 ± 30	154 ± 32	**0.005**	**0.653**
30% Δ (W.kg^{-1})	2.7 ± 0.4	2.5 ± 0.4	0.198	0.213

LT: lactate threshold, Pmax: Maximal ramp test power output, V̇O$_{2peak}$: Maximal rate of oxygen consumption.

$\dot{V}O_2$ *Kinetics*

The transition from unloaded pedalling to moderate and heavy intensity cycling elicited an

increase in $\dot{V}O_2$ (see Figure 2), and the monoexponential curve used to describe the

increase in $\dot{V}O_2$ in males and females demonstrated excellent r^2 values (see Table 2).

Table 2: Data from the monoexponential modelling of $\dot{V}O_2$ kinetics during moderate and heavy
intensity transitions.

	Males (n = 8)	Females (n = 8)	P value	Cohen's d
Moderate Intensity Domain				
TD (s)	16.6 ± 4.1	17.9 ± 4.9	0.575	0.121
Baseline $\dot{V}O_2$ (L.min^{-1})	0.97 ± 0.05	0.77 ± 0.09	**< 0.001**	**1.516**
Amplitude (L.min^{-1})	0.83 ± 0.19	0.60 ± 0.18	**0.024**	**0.477**
Amplitude (ml.kg^{-1}.min^{-1})	11.0 ± 1.6	9.7 ± 2.6	0.230	0.323
Amplitude (% $\dot{V}O_{2peak}$)	24 ± 3	24 ± 5	0.949	0.018
$\dot{V}O_2$ gain (ml.min^{-1}.W^{-1})	10.2 ± 0.7	10.5 ± 1.1	0.587	0.130
τ (s)	27.9 ± 7.5	24.8 ± 6.6	0.385	0.160
r^2	0.965 ± 0.032	0.947 ± 0.034		
Heavy Intensity Domain				
TD (s)	15.0 ± 4.8	14.5 ± 4.5	0.858	0.033
Baseline $\dot{V}O_2$ (L.min^{-1})	1.00 ± 0.06	0.85 ± 0.11	**0.005**	**0.894**
Amplitude (L.min^{-1})	1.50 ± 0.38	0.96 ± 0.25	**0.005**	**0.541**
Amplitude (ml.kg^{-1}.min^{-1})	19.9 ± 4.7	15.6 ± 3.7	0.060	0.351
Amplitude (% $\dot{V}O_{2peak}$)	43 ± 5	38 ± 7	0.179	0.355
$\dot{V}O_2$ gain (ml.min^{-1}.W^{-1})	9.4 ± 0.8	9.3 ± 0.9	0.766	0.059
τ (s)	28.8 ± 7.9	27.2 ± 4.4	0.633	0.075
r^2	0.961 ± 0.031	0.927 ± 0.030		
SC Amplitude (L.min^{-1})	0.38 ± 0.16	0.28 ± 0.12	0.158	0.250
SC Amplitude (% $\dot{V}O_{2peak}$)	11.9 ± 6.9	10.8 ± 3.2	0.686	0.061
SC Amplitude (% end exercise $\dot{V}O_2$)	13.6 ± 6.5	12.9 ± 3.9	0.822	0.035

SC: slow component, τ: time constant, TD: time delay, $\dot{V}O_2$: rate of oxygen consumption

Figure 2: Group mean V̇O₂ data from males (blue, n=8) and females (red, n=8) during moderate
(Panel A) and heavy (Panel B) intensity transitions. The bold lines represent group means and the
thin lines represent standard deviation.

In absolute units (L.min⁻¹) males experienced greater amplitudes of V̇O₂ during the phase II
kinetics ($p \leq 0.024$), however when this was made relative to the individual (V̇O₂ gain and
% V̇O₂peak) no sex differences were observed ($p \geq 0.179$). Similarly, the V̇O₂ slow
component amplitude was not different between sexes in relative units ($p = 0.686$). As
visualised in Figure 1, there were no sex differences in τV̇O₂ in either the moderate ($p = $
0.385) or heavy ($p = 0.633$) intensity domains. Baseline V̇O₂ was slightly elevated at the
onset of heavy intensity exercise compared to the onset of moderate intensity exercise for
females (mean difference: 0.08 L.min⁻¹, $p = 0.028$), whereas male baseline V̇O₂ was not
different ($p = 0.312$).

Oxygen Extraction Kinetics

The transition from unloaded pedalling to moderate and heavy intensity cycling elicited an
increase in HHb+Mb concentration (Figure 3), and the monoexponential curve used to
describe the increase in HHb+Mb in males and females demonstrated excellent r^2 values
(Table 3). One female's data had to be removed due to issues with the NIRS signal,
resulting in n = 7 females being used for NIRS analyses.

Table 3: Data from the monoexponential modelling of deoxyhaemoglobin kinetics during moderate
and heavy intensity transitions.

	Males (n = 8)	Females (n = 7)	P value	Cohen's d
Moderate Intensity Domain				
TD (s)	8.7 ± 1.5	9.6 ± 2.6	0.447	0.212
Baseline HHb+Mb (% ischemia)	23 ± 8	29 ± 7	0.186	0.251
Amplitude (% ischemia)	21 ± 7	17 ± 5	0.225	0.219
τ (s)	8.1 ± 2.8	10.0 ± 3.8	0.274	0.260
r^2	0.974 ± 0.022	0.956 ± 0.037		
Heavy Intensity Domain				
TD (s)	6.2 ± 2.8	7.0 ± 2.3	0.582	0.101
Baseline HHb+Mb (% ischemia)	23 ± 6	17 ± 7	0.146	0.332
Amplitude (% ischemia)	36 ± 9	30 ± 8	0.193	0.252
τ (s)	11.3 ± 3.7	12.2 ± 4.0	0.665	0.090
r^2	0.983 ± 0.011	0.979 ± 0.023		
SC Amplitude (% ischemia)	14 ± 5	16 ± 4	0.432	0.142

HHb+Mb: deoxyhaemoglobin and myoglobin, SC: slow component, τ: time
constant, TD: time delay consumption

As shown in Figure 3, the phase II amplitude of HHb+Mb increase in both intensity domains
was not different between sexes ($p \geq 0.193$). Similarly, there was no sex difference in the
amplitude of the slow component in the heavy intensity domain ($p = 0.432$). The time
constant for phase II HHb+Mb kinetics (τHHb+Mb) was also not different between males and
females in both intensity domains ($p \geq 0.274$). Baseline HHb+Mb was slightly lower at the
onset of heavy intensity exercise for females compared to the onset of moderate exercise
(mean difference: 12% ischemia, $p = 0.002$), but not for males ($p = 0.583$).

Figure 3: Group mean HHb+Mb data from males (blue, n=8) and females (red, n=7) during moderate
(Panel A) and heavy (Panel B) intensity transitions. The bold lines represent group means and the
thin lines represent standard deviation.

Near-infrared spectroscopy

During the moderate intensity cycling, a significant effect of sex was observed for
HbO_2+MbO_2 ($F_{1,13} = 10.85$, $p = 0.006$), but no time ($p = 0.804$) or sex × time interaction ($p =$
0.054) effects were observed. For TOI, significant time ($F_{1,13} = 48.59$, $p < 0.001$) and sex
($F_{1,13} = 12.57$, $p = 0.004$) effects were observed, but no sex × time interaction effect ($p =$
0.095). Post-hoc tests revealed that females had greater values before (6%, $p = 0.002$) and
during the stage (10%, $p = 0.007$). However, when TOI values were normalised as %
ischemia, there was a main effect of time ($F_{1,6} = 115.09$, $p < 0.001$), but neither the main
effect of sex ($p = 0.181$) or the sex × time interaction effect ($p = 0.381$) were evident.

Table 4: Data from near-infrared spectroscopy before and during the moderate and heavy intensity
exercise transitions in males (n = 8) and females (n = 7).

		Moderate Intensity		Heavy Intensity		
		Unloaded Pedalling	End Stage	Unloaded Pedalling	90 secs	End Stage
$HbO_2 + MbO_2$	Male	57 ± 6	53 ± 7	68 ± 11*	54 ± 10*[#]	47 ± 10[#]
(%ischemia)	Female	41 ± 10	44 ± 9	51 ± 7	41 ± 4[#]	53 ± 8
TOI	Male	69 ± 3	61 ± 5[#]	71 ± 4	57 ± 6[#]	50 ± 8[#$]
(%)	Female	75 ± 3*	71 ± 6*[#]	78 ± 4*	70 ± 8*[#]	69 ± 9*[#]
TOI	Male	72 ± 4	60 ± 7[#]	76 ± 6	53 ± 9[#]	39 ± 10[#$]
(% ischemia)	Female	67 ± 9	53 ± 10[#]	77 ± 6	52 ± 9[#]	51 ± 17[#]

HbO_2+MbO_2: oxygenated haemoglobin and myoglobin; TOI: tissue oxygenation index; * = greater than the opposite sex ($p < 0.05$); [#] = lower than unloaded pedalling ($p < 0.05$); [$] = lower than 90 secs.

During heavy intensity cycling, a main effect of time was observed for HbO_2+MbO_2 ($F_{2,26}$ =

17.39, $p < 0.001$), as well as sex × time interaction effect ($F_{2,26} = 15.67$, $p < 0.001$), but no

main effect of sex ($p = 0.052$). Post-hoc tests revealed that females had lower values before

(-17% ischemia, $p = 0.004$) and 90 seconds after the transition (-13% ischemia, $p = 0.005$),

but not at the end of the stage ($p = 0.216$). Furthermore, HbO_2+MbO_2 decreased in both

sexes from unloaded pedalling to 90 seconds into the transition ($p < 0.001$), but for females,

this returned to baseline by the end of the stage ($p = 0.056$) whereas males remained

decreased ($p = 0.002$). For TOI, main effects of time ($F_{2,26} = 70.21$, $p < 0.001$) and a sex ×

time interaction effect ($F_{2,26} = 15.67$, $p < 0.001$) were observed, but no main effect of sex ($p =$

0.052). Post-hoc tests revealed that females had greater values than males at all timepoints

($p \leq 0.005$). Additionally, while males demonstrated a progressive decrease in TOI at each of

the three time points ($p \leq 0.001$), females only decreased from unloaded pedalling to 90

seconds ($p < 0.001$), then no further decrease was observed at 30 minutes ($p = 1.000$).

When TOI was normalised to % ischemia, a main effect of time ($F_{1.37,17.78} = 49.50$ $p < 0.001$)

remained, however the sex ($p = 0.266$) and sex × time interaction effect ($p = 0.112$) were not

observed.

Discussion

This study aimed to compare the kinetics of $\dot{V}O_2$ and HHb+Mb during moderate and heavy

intensity exercise in males and females. In contrast to the hypothesis, at the onset of

exercise the phase II time constants (τ) for $\dot{V}O_2$ and HHb+Mb were not different between

the sexes, implying that both males and females were able to increase oxidative

phosphorylation at comparable rates. In absolute units, males had larger amplitude

increases than females, however when normalised to the individuals' maximum values, the

rise in $\dot{V}O_2$ and HHb+Mb was not different. Combined, these data demonstrate that the

oxidative response to exercise is not different between sexes, which provides mechanistic

insight into previously observed sex differences in the integrative response to exercise.

Previous literature investigating sex differences in the onset kinetics of oxygen transport and

utilisation conflicts with the present data, with Beltrame *et al.* (2017) demonstrating quicker

$\tau\dot{V}O_2$ and τHHb+Mb in females compared to males. One potential explanation for this

discrepancy could be that Beltrame *et al.* utilised a treadmill walking task, compared to

cycling. In tasks where O_2 delivery is not a limiting factor, females often outperform males.

For instance, Ansdell *et al.* (2019) showed female knee-extensors had a greater relative

critical torque than males during single-limb exercise. Whereas during cycling, where O_2

delivery is a determinant of critical power (Goulding & Marwood, 2023), this metabolic

threshold was not different between sexes (Ansdell *et al.*, 2020a). Whilst consensus on

whether O_2 delivery does (Hughson *et al.*, 2001) or does not (Grassi, 2001) limit $\tau \dot{V}O_2$ has

not been reached, it is conceivable that during tasks where O_2 delivery and utilisation are

both determinants in the metabolic response to exercise, the superior female skeletal muscle

oxidative capacity (Cardinale *et al.*, 2018) and vasodilatory response to exercise (Parker *et*

al., 2007) is counteracted by an inferior O_2 carrying capacity (Murphy, 2014). Within the

present data, this balance manifests as a comparable $\tau \dot{V}O_2$ in males and females, which

agrees with data from do Nascimento Salvador *et al.* (2019), who demonstrated no sex

difference in $\tau \dot{V}O_2$ during a transition from unloaded pedalling to 'very heavy' (60% Δ)

cycling exercise.

Data from incremental exercise suggests that the poorer O_2 delivery in females results in a

greater degree of O_2 extraction to compensate (Murias *et al.*, 2013). The present data

contradicts this notion, as the amplitude of phase II HHb+Mb kinetics was not different

between sexes (see Table 3). However, it is important to note that Murias *et al.* noted that

this sex difference only occurred once incremental exercise exceeded the respiratory

compensation point (i.e., the severe intensity domain), whereas the present study compared

sexes in the moderate and heavy intensity domains. The lack of a sex difference in the

phase II amplitude for HHb+Mb kinetics contradicts previously published NIRS data that

demonstrated a smaller rise in HHb+Mb and lesser decrease in TOI in females compared to

males during constant-load exercise (Ansdell *et al.*, 2020a). The crucial difference in

methodologies employed between the previous study and the present study is the

application of a 'physiological calibration' to negate the influence of adipose tissue thickness

on NIRS signals (Ryan *et al.*, 2012). Previously, the sex difference in the rise in HHb+Mb

was suggested to reflect a lower oxygen cost of muscle contraction in female knee-

extensors, however the present data, with more rigorous methodologies employed, refutes

this. One aspect of the modelling that did demonstrate a sex difference was the reduction in

HHb+Mb (and concomitant increase in $\dot{V}O_2$) baseline for the heavy intensity transition in

females, but not males. This could reflect a sex difference in how O_2 utilisation is altered by

prolonged or intermittent exercise (i.e., the preceding three bouts of moderate intensity

cycling); however, the present study was not configured in a manner appropriate to answer

that research question. Females did demonstrate lower HbO_2+MbO_2 levels during heavy

intensity cycling, perhaps indicating a lesser O_2 availability. This would fit with the notion that

O_2 carrying capacity is inferior in females during high-intensity exercise (Diaz-Canestro *et*

al., 2022); however, the lack of difference in the speed and amplitude of HHb+Mb onset

kinetics (and pulmonary $\dot{V}O_2$) implies that oxygen extraction is not negatively affected by

this in the moderate and heavy intensity domains.

Although not measured in the present study, the sex difference in muscle fibre type, whereby

females demonstrate a greater proportional area of type I fibres (Staron *et al.*, 2000;

Roepstorff *et al.*, 2006), appears to have not influenced either $\dot{V}O_2$ or HHb+Mb onset

kinetics in the present study. This would concur with data from Barstow *et al.* (1996), who

demonstrated no relationship between type I fibre percentage of the *vastus lateralis* and the

time constant of phase II $\dot{V}O_2$ kinetics. In contrast, Pringle *et al.* (2003) observed a

negative correlation between type I fibre percentage and the phase II time constant in the

heavy intensity domain only. Of note is that Pringle *et al.* included a wide range of

participants with ~27 – 85% type I fibres, and when groups were split into discrete groups of

low and high fibre type percentages (mean difference: 25%), the high percentage group had

faster phase II kinetics. The present study was not able to quantify the sex difference in

muscle fibre typology, however, previous literature has observed a 5-13% difference in type I

fibre percentage of the *vastus lateralis* (Simoneau & Bouchard, 1989; Staron *et al.*, 2000;

Roepstorff *et al.*, 2006). Therefore, it could be the case that the sex difference in muscle

fibre typology is not large enough to affect the phase II $\dot{V}O_2$ or HHb+Mb kinetics. Indeed,

recent evidence suggests that during exercise normalised to metabolic thresholds, sex

differences in muscle fibre typology do not influence fatigability (McDougall *et al.*, 2023).

Muscle fibre typology has previously been demonstrated to affect the amplitude of the slow

component within the heavy intensity domain, as individuals with a lower type I fibre

percentage experience larger rises in $\dot{V}O_2$ during constant-load exercise (Barstow *et al.*,

1996; Pringle *et al.*, 2003). It is suggested that the slow component is mechanistically

underpinned by factors such as additional motor unit recruitment (Poole *et al.*, 1994; Burnley

et al., 2002) to compensate for fatigue-related changes in muscle metabolism. For instance,

muscle PCr stores demonstrate a similar slow component in depletion during heavy intensity

exercise (Rossiter *et al.*, 2002). Given that female knee-extensors appears more fatigue-

resistant and demonstrate lesser rises in the amplitude of surface electromyography during

constant-load exercise (Ansdell *et al.*, 2017; Ansdell *et al.*, 2019; Ansdell *et al.*, 2020a), we

hypothesised that the relative amplitude of the slow component would be greater in males to

reflect a greater rate of metabolic disturbance. However, as is evident in Tables 2 and 3, no

sex difference was observed in the relative slow component amplitude, implying that there

was no difference in the metabolic response to constant-load exercise.

The lack of sex differences in either the phase II kinetics or slow component amplitude

collectively suggest that the oxidative response to exercise was not different between males

and females. Data on this topic is sparse, and due to the nature of methods such as

magnetic resonance spectroscopy (MRS), limited to single-joint, isometric muscle
contractions. Previous literature using this technique to study muscle metabolic changes
during a 60 s contraction of the dorsiflexors showed no sex difference in changes in PCr, Pi,
or pH (Russ *et al.*, 2005). Data from muscle biopsies of the *vastus lateralis* taken before and
after repeated 30s cycling sprints suggested a greater preservation of ATP concentrations in
females across a ~60 minute protocol (Esbjörnsson-Liljedahl *et al.*, 2002); however the
authors suggested that this was likely a result of sex differences in the 20 minute recovery
periods, rather that metabolic differences during exercise. Accordingly, the same group
observed no sex differences in the metabolic response to a single 30-second cycling sprint
(Esbjörnsson-Liljedahl *et al.*, 1999). Collectively, across multiple tasks and methodologies
(MRS, biopsy and $V O_2$ kinetics), the data suggest that there is no sex difference in the
bioenergetic response to high-intensity exercise. This information provides mechanistic
insight into the sex differences in the integrative response to exercise. For instance, sex
differences in fatigability have partly been attributed to a lesser accumulation of fatiguing
metabolites (Hunter, 2014; Ansdell *et al.*, 2020b). It is perhaps more accurate to suggest that
previously observed sex differences in fatigue during intensity-matched exercise (Ansdell *et*
al., 2020a; Azevedo *et al.*, 2021) are more likely due to a greater fatigue-resistance of
female muscle contractile apparatus, which experience similar degrees of metabolic stress
as males. It is established that males and females differ in contractile properties such as
calcium (Ca^{2+}) kinetics of the sarcoplasmic reticulum (Harmer *et al.*, 2014), with lower
Ca^{2+}ATPase activity thought to permit a more fatigue-resistant skeletal muscle profile during
equivalent exercise tasks (Hunter, 2014). Therefore, the present study advances the
contemporary understanding of sex differences in the integrative response to exercise and
provides mechanistic insight into previously observed phenomena.

These data have applications across the spectrum of health and disease. For example,
those prescribing steady state exercise to improve skeletal muscle performance in athletes
or patients might not need to account for the sex of their participants (Gloeckl *et al.*, 2022;
Furrer *et al.*, 2023). This statement should however, be caveated by the fact that evidence
regarding the influence of sex on long-term adaptation to exercise is sparse (Ansdell *et al.*,
2020b). Indeed, one area for further exploration is the bioenergetic response to exercise
within the severe intensity domain, where sex differences in fatigability have previously been
observed (Ansdell *et al.*, 2020a; Azevedo *et al.*, 2021). The employment of complementary
techniques to quantify O_2 delivery (for instance, the present study did not quantify total
Hb+Mb) and muscle fibre typology could also provide greater insight into the influence of sex
on the O_2 cascade in a variety of tasks.

Previous evidence from pre-pubertal children and adolescents suggests that boys have a
lower $\tau\dot{V}O_2$ and $\dot{V}O_2$ slow component than girls (Armstrong & Barker, 2009), which
contradicts the present findings in adults. Previous studies comparing children and adults
also found that intramuscular PCr kinetics were similar in males and females regardless of
age (Willcocks *et al.*, 2010), implying similar oxidative capacity. Interestingly, the same study
found a greater 'PCr cost' ($mM.W^{-1}$) in females compared to males suggesting a greater
inefficiency in females, which contradicts the present findings. Caution is urged when
comparing data from Willcocks *et al.* (2010) and the present study, however, given the
nature of exercise (single joint vs. whole body), the lack of matching for aerobic fitness, and
the low sample size (6 males vs. 5 females).

Finally, hormonal status was not an exclusion criterion or controlled for within female
participants in the present study. We based this decision on evidence from Mattu *et al.*
(2020) who demonstrated that $\dot{V}O_2$ kinetics did not differ between the follicular and luteal
phases of the eumenorrheic menstrual cycle, or with oral contraceptive usage. However, we
do acknowledge that the aforementioned study only investigated the moderate intensity
domain, and therefore might not apply to $\dot{V}O_2$ kinetics in the heavy intensity domain. In the
heavy intensity domain, contributing factors to the $\dot{V}O_2$ slow component (e.g., substrate
utilisation) might be affected by hormonal status, although the available evidence is
conflicting (Oosthuyse *et al.*, 2023). Therefore, further research is required to determine
whether endogenous and exogenous hormones influence the oxidative response to high-
intensity exercise.

Conclusion
The present study aimed to compare the oxygen extraction and uptake kinetics during
moderate and heavy intensity cycling exercise. Contrary to our hypotheses, no sex
differences were observed in either the phase II or slow component kinetics for $\dot{V}O_2$ or
HHb+Mb. The lack of sex difference implies that males and females do not experience
different oxidative responses to exercise, which provides mechanistic insight into previously
observed phenomena such as the sex difference in fatigability. Furthermore, based on these
data and others demonstrating no hormonal influences (Mattu *et al.*, 2020), we suggest that
there is no rationale for the exclusion of female participants in research investigating
cardiopulmonary responses to exercise.

Acknowledgements

The authors wish to thank the participants for their time and effort, as well as the technical

staff within the Department of Sport, Exercise and Rehabilitation for their support.

Funding

This project was supported by a Physiological Society Research Springboard Studentship

awarded to MSP, as well as Erasmus+ funding awarded to LB (2020-1-DE01-KA103-

005569). PA is supported by the UK Office for Veteran's Affairs (G2-SCH-2022-11-12245).

Disclosures

There are no conflicts of interest, financial or otherwise.

Author Contributions

MSP, ZW, MB, and PA conceived and designed the research. MSP, LB, EH, LH, and PA

performed the experiments. LB, EH, LH, ZW, and PA analysed data. MSP, LB, EH, LH, ZW,

MB, and PA interpreted results of the experiments. PA drafted the manuscript. MSP, LB, EH,

LH, ZW, MB, and PA edited and revised the manuscript. All authors approved the final

version of the manuscript.

Reference List

Allen DG, Lamb GD & Westerblad H. (2008). Skeletal muscle fatigue: cellular mechanisms. *Physiol*
Rev **88,** 287-332.

Ansdell P, Brownstein CG, Škarabot J, Hicks KM, Howatson G, Thomas K, Hunter SK & Goodall S.
(2019). Sex differences in fatigability and recovery relative to the intensity–duration
relationship. *The Journal of Physiology* **597,** 5577-5595.

Ansdell P, Škarabot J, Atkinson E, Corden S, Tygart A, Hicks KM, Thomas K, Hunter SK, Howatson G &
Goodall S. (2020a). Sex differences in fatigability following exercise normalised to the
power–duration relationship. *The Journal of Physiology* **598,** 5717-5737.

Ansdell P, Thomas K, Hicks KM, Hunter SK, Howatson G & Goodall S. (2020b). Physiological sex
differences affect the integrative response to exercise: acute and chronic implications.
Experimental Physiology **105,** 2007-2021.

Ansdell P, Thomas K, Howatson G, Hunter S & Goodall S. (2017). Contraction intensity and sex
differences in knee-extensor fatigability. *Journal of Electromyography and Kinesiology* **37,**
68-74.

Armstrong N & Barker AR. (2009). Oxygen uptake kinetics in children and adolescents: a review.
Pediatr Exerc Sci **21,** 130-147.

Azevedo RdA, Forot J, Iannetta D, MacInnis MJ, Millet GY & Murias JM. (2021). Slight power output
manipulations around the maximal lactate steady state have a similar impact on fatigue in
females and males. *Journal of Applied Physiology* **130,** 1879-1892.

Barstow TJ. (2019). Understanding near infrared spectroscopy and its application to skeletal muscle
research. *Journal of Applied Physiology* **126,** 1360-1376.

Barstow TJ, Jones AM, Nguyen PH & Casaburi R. (1996). Influence of muscle fiber type and pedal
frequency on oxygen uptake kinetics of heavy exercise. *Journal of Applied Physiology* **81,**
1642-1650.

Beltrame T, Villar R & Hughson RL. (2017). Sex differences in the oxygen delivery, extraction, and
uptake during moderate-walking exercise transition. *Applied Physiology, Nutrition, and*
Metabolism **42,** 994-1000.

Burnley M, Doust JH, Ball D & Jones AM. (2002). Effects of prior heavy exercise on$\dot{V}o_2$ kinetics
during heavy exercise are related to changes in muscle activity. *Journal of Applied Physiology*
93, 167-174.

Burnley M, Doust JH & Jones AM. (2006). Time required for the restoration of normal heavy exercise
$\dot{V}o_2$ kinetics following prior heavy exercise. *Journal of Applied Physiology* **101,** 1320-1327.

Burnley M & Jones AM. (2007). Oxygen uptake kinetics as a determinant of sports performance.
European Journal of Sport Science **7,** 63-79.

Cano A, Ventura L, Martinez G, Cugusi L, Caria M, Deriu F & Manca A. (2022). Analysis of sex-based
differences in energy substrate utilization during moderate-intensity aerobic exercise. *Eur J*
Appl Physiol **122,** 29-70.

Cardinale DA, Larsen FJ, Schiffer TA, Morales-Alamo D, Ekblom B, Calbet JAL, Holmberg HC & Boushel
R. (2018). Superior Intrinsic Mitochondrial Respiration in Women Than in Men. *Front Physiol*
9, 1133.

DeLorey DS, Kowalchuk JM & Paterson DH. (2003). Relationship between pulmonary O2 uptake
kinetics and muscle deoxygenation during moderate-intensity exercise. *Journal of Applied*
Physiology **95,** 113-120.

Diaz-Canestro C, Pentz B, Sehgal A & Montero D. (2022). Differences in Cardiac Output and Aerobic
Capacity Between Sexes Are Explained by Blood Volume and Oxygen Carrying Capacity. *Front*
Physiol **13,** 747903.

do Nascimento Salvador PC, Schäfer L, Grassi B, Guglielmo LGA & Denadai BS. (2019). Changes in
VO2 Kinetics After Elevated Baseline Do Not Necessarily Reflect Alterations in Muscle Force
Production in Both Sexes. *Frontiers in Physiology* **10**.

Ducrocq GP, Hureau TJ, Bøgseth T, Meste O & Blain GM. (2021). Recovery from Fatigue after Cycling
Time Trials in Elite Endurance Athletes. *Med Sci Sports Exerc* **53**, 904-917.

Esbjörnsson-Liljedahl M, Bodin K & Jansson E. (2002). Smaller muscle ATP reduction in women than
in men by repeated bouts of sprint exercise. *Journal of Applied Physiology* **93**, 1075-1083.

Esbjörnsson-Liljedahl M, Sundberg CJ, Norman B & Jansson E. (1999). Metabolic response in type I
and type II muscle fibers during a 30-s cycle sprint in men and women. *Journal of Applied*
Physiology **87**, 1326-1332.

Faude O, Kindermann W & Meyer T. (2009). Lactate Threshold Concepts. *Sports Medicine* **39**, 469-
490.

Furrer R, Hawley JA & Handschin C. (2023). The molecular athlete: exercise physiology from
mechanisms to medals. *Physiological Reviews* **103**, 1693-1787.

Gloeckl R, Zwick RH, Fürlinger U, Jarosch I, Schneeberger T, Leitl D, Koczulla AR, Vonbank K, Alexiou
C, Vogiatzis I & Spruit MA. (2022). Prescribing and adjusting exercise training in chronic
respiratory diseases – Expert-based practical recommendations. *Pulmonology*.

Goulding RP & Marwood S. (2023). Interaction of Factors Determining Critical Power. *Sports*
Medicine **53**, 595-613.

Goulding RP, Rossiter HB, Marwood S & Ferguson C. (2021). Bioenergetic Mechanisms Linking $\dot{V}O_2$
Kinetics and Exercise Tolerance. *Exercise and Sport Sciences Reviews* **49**.

Grassi B. (2001). Regulation of Oxygen Consumption at Exercise Onset: Is It Really Controversial?
Exercise and Sport Sciences Reviews **29**, 134-138.

Grassi B, Poole DC, Richardson RS, Knight DR, Erickson BK & Wagner PD. (1996). Muscle O2 uptake
kinetics in humans: implications for metabolic control. *Journal of Applied Physiology* **80**, 988-
998.

Grassi B, Porcelli S, Salvadego D & Zoladz JA. (2011). Slow VO_2 kinetics during moderate-intensity
exercise as markers of lower metabolic stability and lower exercise tolerance. *Eur J Appl*
Physiol **111**, 345-355.

Grassi B, Rossiter HB & Zoladz JA. (2015). Skeletal muscle fatigue and decreased efficiency: two sides
of the same coin? *Exerc Sport Sci Rev* **43**, 75-83.

Harmer AR, Ruell PA, Hunter SK, McKenna MJ, Thom JM, Chisholm DJ & Flack JR. (2014). Effects of
type 1 diabetes, sprint training and sex on skeletal muscle sarcoplasmic reticulum Ca2+
uptake and Ca2+-ATPase activity. *The Journal of Physiology* **592,** 523-535.

Harms CA, McClaran SR, Nickele GA, Pegelow DF, Nelson WB & Dempsey JA. (1998). Exercise-
induced arterial hypoxaemia in healthy young women. *J Physiol* **507 (Pt 2),** 619-628.

Hughson RL, Tschakovsky ME & Houston ME. (2001). Regulation of Oxygen Consumption at the
Onset of Exercise. *Exercise and Sport Sciences Reviews* **29,** 129-133.

Hunter SK. (2014). Sex differences in human fatigability: mechanisms and insight to physiological
responses. *Acta Physiologica* **210,** 768-789.

James JJ, Klevenow EA, Atkinson MA, Vosters EE, Bueckers EP, Quinn ME, Kindy SL, Mason AP,
Nelson SK, Rainwater KAH, Taylor PV, Zippel EP & Hunter SK. (2023). Underrepresentation of
women in exercise science and physiology research is associated with authorship gender.
Journal of Applied Physiology **135,** 932-942.

Koppo K, Bouckaert J & Jones AM. (2004). Effects of Training Status and Exercise Intensity on Phase II
$\dot{V}O2$ Kinetics. *Medicine & Science in Sports & Exercise* **36,** 225-232.

Korzeniewski B & Rossiter HB. (2015). Each-step activation of oxidative phosphorylation is necessary
to explain muscle metabolic kinetic responses to exercise and recovery in humans. *The
Journal of Physiology* **593,** 5255-5268.

Mattu AT, Iannetta D, MacInnis MJ, Doyle-Baker PK & Murias JM. (2020). Menstrual and oral
contraceptive cycle phases do not affect submaximal and maximal exercise responses.
Scandinavian Journal of Medicine & Science in Sports **30,** 472-484.

McDougall RM, Tripp TR, Frankish BP, Doyle-Baker PK, Lun V, Wiley JP, Aboodarda SJ & MacInnis MJ.
(2023). The influence of skeletal muscle mitochondria and sex on critical torque and
performance fatiguability in humans. *The Journal of Physiology* **601,** 5295-5316.

Miotto PM, McGlory C, Holloway TM, Phillips SM & Holloway GP. (2018). Sex differences in
mitochondrial respiratory function in human skeletal muscle. *American Journal of
Physiology-Regulatory, Integrative and Comparative Physiology* **314,** R909-R915.

Murias JM, Keir DA, Spencer MD & Paterson DH. (2013). Sex-related differences in muscle
deoxygenation during ramp incremental exercise. *Respiratory Physiology & Neurobiology*
189, 530-536.

Murias JM, Kowalchuk JM & Paterson DH. (2010). Speeding of $\dot{V}o2$ kinetics with endurance training
in old and young men is associated with improved matching of local O2 delivery to muscle
O2 utilization. *Journal of Applied Physiology* **108,** 913-922.

Murphy WG. (2014). The sex difference in haemoglobin levels in adults — Mechanisms, causes, and
753    consequences. *Blood Reviews* **28,** 41-47.

755 Oosthuyse T, Strauss JA & Hackney AC. (2023). Understanding the female athlete: molecular
756    mechanisms underpinning menstrual phase differences in exercise metabolism. *Eur J Appl*
757    *Physiol* **123,** 423-450.

759 Parker BA, Smithmyer SL, Pelberg JA, Mishkin AD, Herr MD & Proctor DN. (2007). Sex differences in
760    leg vasodilation during graded knee extensor exercise in young adults. *J Appl Physiol (1985)*
761    **103,** 1583-1591.

763 Poole DC, Barstow TJ, Gaesser GA, Willis WT & Whipp BJ. (1994). VO2 slow component: physiological
764    and functional significance. *Med Sci Sports Exerc* **26,** 1354-1358.

766 Poole DC & Jones AM. (2012). Oxygen Uptake Kinetics. *Compr Physiol* **2,** 933-996.

768 Porcelli S, Marzorati M, Morandi L & Grassi B. (2016). Home-based aerobic exercise training
769    improves skeletal muscle oxidative metabolism in patients with metabolic myopathies.
770    *Journal of Applied Physiology* **121,** 699-708.

772 Pringle JSM, Doust JH, Carter H, Tolfrey K, Campbell IT & Jones AM. (2003). Oxygen uptake kinetics
773    during moderate, heavy and severe intensity 'submaximal' exercise in humans: the influence
774    of muscle fibre type and capillarisation. *European Journal of Applied Physiology* **89,** 289-300.

776 Roepstorff C, Thiele M, Hillig T, Pilegaard H, Richter EA, Wojtaszewski JF & Kiens B. (2006). Higher
777    skeletal muscle alpha2AMPK activation and lower energy charge and fat oxidation in men
778    than in women during submaximal exercise. *J Physiol* **574,** 125-138.

780 Rossiter HB, Ward SA, Howe FA, Kowalchuk JM, Griffiths JR & Whipp BJ. (2002). Dynamics of
781    intramuscular 31P-MRS Pi peak splitting and the slow components of PCr and O2 uptake
782    during exercise. *Journal of Applied Physiology* **93,** 2059-2069.

784 Rossiter HB, Ward SA, Kowalchuk JM, Howe FA, Griffiths JR & Whipp BJ. (2001). Effects of prior
785    exercise on oxygen uptake and phosphocreatine kinetics during high-intensity knee-
786    extension exercise in humans. *J Physiol* **537,** 291-303.

788 Russ DW, Lanza IR, Rothman D & Kent-Braun JA. (2005). Sex differences in glycolysis during brief,
789    intense isometric contractions. *Muscle & Nerve* **32,** 647-655.

791 Ryan TE, Erickson ML, Brizendine JT, Young H-J & McCully KK. (2012). Noninvasive evaluation of
792    skeletal muscle mitochondrial capacity with near-infrared spectroscopy: correcting for blood
793    volume changes. *Journal of Applied Physiology* **113,** 175-183.

Schiaffino S & Reggiani C. (2011). Fiber types in mammalian skeletal muscles. *Physiol Rev* **91,** 1447-
1531.

Simoneau JA & Bouchard C. (1989). Human variation in skeletal muscle fiber-type proportion and
enzyme activities. *American Journal of Physiology-Endocrinology and Metabolism* **257,** E567-
E572.

Staron RS, Hagerman FC, Hikida RS, Murray TF, Hostler DP, Crill MT, Ragg KE & Toma K. (2000). Fiber
type composition of the vastus lateralis muscle of young men and women. *J Histochem*
Cytochem **48,** 623-629.

Temesi J, Mattioni Maturana F, Peyrard A, Piucco T, Murias JM & Millet GY. (2017). The relationship
between oxygen uptake kinetics and neuromuscular fatigue in high-intensity cycling
exercise. *European Journal of Applied Physiology* **117,** 969-978.

Tschakovsky ME & Hughson RL. (1999). Interaction of factors determining oxygen uptake at the
onset of exercise. *Journal of Applied Physiology* **86,** 1101-1113.

Willcocks RJ, Williams CA, Barker AR, Fulford J & Armstrong N. (2010). Age- and sex-related
differences in muscle phosphocreatine and oxygenation kinetics during high-intensity
exercise in adolescents and adults. *NMR in Biomedicine* **23,** 569-577.

Figure Legends

Figure 1: Visualisation of the monoexponential curve fitting procedures for a representative

participant's data in the moderate intensity domain. Panel A describes the $\dot{V}O_2$ data (1 Hz)

and Panel B describes the HHb+Mb data (0.2 Hz).

Figure 2: Group mean $\dot{V}O_2$ data from males (blue, n=8) and females (red, n=8) during

moderate (Panel A) and heavy (Panel B) intensity transitions. The bold lines represent group

means and the thin lines represent standard deviation.

Figure 3: Group mean HHb+Mb data from males (blue, n=8) and females (red, n=7) during

moderate (Panel A) and heavy (Panel B) intensity transitions. The bold lines represent group

means and the thin lines represent standard deviation.