Beyond venomous fangs: Uloboridae spiders have lost their venom but not their toxicity

Xiaojing Peng[1], Ludwig Dersch[3,4], Josephine Dresler[3,4], Tim Lüddecke[3,4], Tim Dederichs[2], Peter Michalik[2], Steve Peigneur[5], Jan Tytgat[5], Afrah Hassan[1], Antonio Mucciolo[6], Marc Robinson-Rechavi[1,7], Giulia Zancolli[1,7,*]

[1] Department of Ecology and Evolution, University of Lausanne, 1015, Lausanne, Switzerland
[2] Zoological Institute and Museum, University of Greifswald, Loitzer Straße 26, 17489, Greifswald, Germany
[3] Animal Venomics Lab, Fraunhofer Institute for Molecular Biology and Applied Ecology (IME), Ohlebergsweg 12, 35392, Gießen, Germany
[4] LOEWE-Centre for Translational Biodiversity Genomics (LOEWE-TBG), Senckenberganlage 25, 60325 Frankfurt am Main, Germany
[5] Toxicology and Pharmacology, University of Leuven (KU Leuven), P.O. Box 922, Herestraat 49, 3000 Leuven, Belgium
[6] Electron Microscopy Facility, University of Lausanne, 1015, Lausanne, Switzerland
[7] Swiss Institute of Bioinformatics, 1015, Lausanne, Switzerland

* Corresponding author: Giulia Zancolli
Email: giulia.zancolli@gmail.com

25   **Keywords**: toxins, evolution, secondary loss, Uloboridae, digestive fluids, defensin
26

Abstract

Venom, one of nature's most potent secretions, has played a crucial role in the evolutionary
success of many animal groups, including spiders. However, Uloboridae spiders appear to
lack venom and capture their prey, unlike venomous spiders, by extensive silk-wrapping and
regurgitation of digestive fluids onto the entire prey. A prevailing hypothesis posits that
toxins may have been reallocated from the venom to alternative secretions, like silk or
digestive fluids. Yet, whether uloborids have retained venom toxins and the mechanisms
underlying prey immobilisation remain unresolved. Here, we employed a multi-disciplinary
approach to assess the absence of venom glands in *Uluborus plumipes*, toxin gene
expression and toxicity of digestive proteins. Our findings confirm that *U. plumipes* lacks a
venom apparatus, while neurotoxin-like transcripts were highly expressed in the digestive
system. Midgut gland extract had comparable toxicity levels to that of the venomous
Parasteatoda tepidariorum. However, no inhibitory effects on sodium nor potassium
channels were observed, indicating a different toxic mechanism. These findings support the
hypothesis that Uloboridae spiders have lost their venom apparatus while retaining toxin-
like genes. The potent toxicity of their digestive fluids, a trait conserved across spiders, likely
compensate for the absence of venom, ensuring effective prey immobilisation and
digestion.

Background

The arms race between predators and their prey is a fundamental driving force of animal
adaptation. Over time, animals have evolved a range of strategies to succeed in this ongoing
battle for survival and to gain advantages over their opponents. Among nature's most
effective weapons is venom - a potent cocktail of proteins and peptides capable of
immobilising prey or deterring predators. Venomous animals are widespread across most
branches of the animal tree of life [1,2]. This success can be attributed to the unique ability
of venom to shift the battleground from the physical to the chemical level, enabling smaller
and slower creatures to prevail over larger and faster ones. Moreover, venom has been
linked to adaptive radiations in several animal clades, such as cone snails [3] and spiders [4–
6], with diversification rates twice as high in venomous compared to non-venomous animal
families [7].

The loss of venom might seem counterintuitive given its advantages, but it can be explained
by its high metabolical cost due to the protein-rich nature of venom. Producing venom
demands significant resources, creating competition with other physiological needs [6,8–
10]. In species where venom use becomes less critical for survival, natural selection may
favour its reduction or complete loss. Relaxed purifying selection can allow loss-of-function
mutations to accumulate overtime through genetic drift, ultimately leading to the
degradation of venom systems [11]. For example, in some species of catfish that rely on
body size for predator defence, or sea snakes feeding exclusively on eggs [12], their venoms
have become unnecessary [13]. In spiders, a similar phenomenon may have occurred in the
Uloboridae family, which appears to lack functional venom glands [14]. In these spiders, the
frontal part of their prosoma contains only muscle bundles where venom glands are

typically located in araneomorph spiders. Whether uloborids are effectively venomless and
how they subdue their prey without venom remains unclear.
Uloboridae spiders employ a unique hunting strategy characterised by extensive wrapping
of prey in silk (up to hundreds of metres per prey), followed by the application of digestive
fluids over the prey's entire surface. This is in contrast to other spiders which typically
regurgitated the fluids only on the area around their own mouthparts [15–18]. The broad
application of the uloborid digestive fluids onto the prey appears to be lethal; Weng and
colleagues [19] found that most ants in silk packages exposed to digestive fluids of the
uloborid *Philoponella vicina* were dead, while most of the unwetted ants, or ants wetted
with water, survived. This observation suggests that uloborids might have shifted the
secretion of toxins from venom to digestive fluids, raising the question of whether they have
retained venom components in other body secretions.
82
In this study, we tested the hypothesis that uloborids have replaced venom with toxic
molecules in their digestive fluids, rendering venom glands redundant and leading to their
evolutionary loss. We confirmed the absence of specialised venom-secreting glands in the
prosoma of *Uloborus plumipes* and detected the expression of venom toxin homologs in
various body tissues, particularly the midgut gland. Functional assays of midgut gland
extracts showed no neurotoxicity against voltage-gated sodium or potassium channels.
Nevertheless, they rapidly killed *Drosophila* flies, demonstrating the innately potency of
uloborid digestive fluids. Similar results were observed for digestive fluids of the venomous
Theridiidae common house spider, *Parasteatoda tepidariorum*, suggesting that the
innovation in Uloboridae spiders lies not in the repurposing of toxins but rather in the
strategic use of pre-existing secretions.
94
Results
Morphological evidence for the lack of venom glands
The objective of this study was to examine the structure of the chelicerae of *U. plumipes*
and compare them with those of *P. tepidariorum*. Based on visual examination of dissected
individuals and histological transverse sections across the chelicerae and anterior part of the
prosoma, we confirmed the absence of venom glands in *U. plumipes* (Fig. 1). The chelicerae
have a prominent epithelium and large muscles (Fig. 1A-E) but lack any kind of duct of the
kind observed in *P. tepidariorum* (Fig. 1F, G).
103

Figure 1: **Histological transversal sections of the chelicerae of *Uloborus plumipes* and, for comparison, the venomous *Parasteatoda tepidariorum*.** A-E: Cross-sections of *U. plumipes* (red), F-G: Cross-sections from *P. tepidariorum* (purple). A, B: Distal end of the chelicera and fang. C: Section through the area just above the fang base. Note that the fang does not contain any kind of canal. D, E: The proximal end of the chelicera, close to the transition to the prosoma, is filled with very large flexor muscles, but no duct nor venom gland base is visible. F: Venom duct and opening visible in the fang. G: Venom duct clearly visible in the chelicera. H: Image of *U. plumipes* by Olei under CC BY-SA 2.5 and I: Image of *P. tepidariorum* by J. Gallagher under CC-BY-2.0 via Wikimedia Commons.

Electron microscopy revealed the presence of three small pores at the tip of the uloborid fangs (Fig. 2A); however, compared to the much larger pore size of the theridiid fang (Fig. 2B), it is unlikely that the uloborid pores represent the opening of a venom duct. Similar small cuticular pits have been observed at the fang's tip of the orb-web spider *Nephila clavata*, although their function remains unknown [20].

121
Figure 2: **Fangs of *Uloborus plumipes* and *Parasteatoda tepidariorum*.** A: *U. plumipes*
possesses small pits at the tip of the fangs (black arrows in the zoomed insert). B: As a
comparison, *P. tepidariorum* possesses a needle-like opening of the venom duct at the tip of
the fangs.

RNA-seq reveals the expression of putative toxins in the midgut gland of *U. plumipes*
De novo transcriptome assembly statistics
Although our morphological analysis confirmed the absence of a specialised venom-
producing and delivery system in the anterior part of the spider's body, it is possible that
uloborids relocated the production of venom components to other organs that produce
secretions that come into contact with the prey. To investigate this further, we performed
RNA sequencing of 13 libraries from different tissues of *U. plumipes* including midgut gland,
silk glands, chelicerae, prosoma, and gonads. The libraries ranged in size from 53 to 107
million reads (Supplementary Table S1).
The resulting *de novo* transcriptome contained 302'504 assembled transcripts, of which
20'243 (6.7%) passed our annotation pipeline (Table S2, Dataset 1). Our final assembly was
83.5% complete according to OMArk [21] (Fig. S1). OrthoVenn3 [22] identified 18'258
transcripts (90% of the total) as homologous to the *Uloborus diversus* genome annotation,
and these were then clustered into 10'518 orthogroups. Among these orthogroups, 9'817
were shared with *U. diversus*.
The distribution of transcript counts was similar among libraries, except the midgut sample
"UM1", which had a smaller median value than the other libraries (Fig S2, S3). Principal
component analysis showed that samples clustered by tissue type, with the first two
components explaining 60% of the total variance (Fig S4).

Identification of toxin-like transcripts in the midgut gland
Manual screening of the final assembly identified 124 translated sequences matching
venom-related proteins in the UniProt-ToxProt database [23] with biased expression across
tissues (Dataset 2), and with 27 predicted to have neurotoxic activity using the NT-
estimation model. The most represented gene families included astacin-like
metalloprotease with 32 sequences (26%), serine proteases (13%), and cysteine-rich
peptides such as atracotoxins, prokineticins, kunitz-type serine protease inhibitors, and
thyroglobulin type-1 (Fig. 3A). Notably, most toxin-like transcripts were significantly

upregulated in the midgut gland, supporting the hypothesis of venom components repurposed into digestive secretion (Fig. 3B).

Figure 3: Classification and tissue distribution of *U. plumipes* toxin-like transcripts. A: Distribution of toxin-like transcripts classified based on their InterProScan predicted domains. The colour corresponds to the body tissue where their expression was significantly upregulated. 'No prediction' includes sequences without InterProScan annotation but with a positive BlastP hit. B) Expression levels of the 124 toxin-like transcripts across the tissues investigated. The transcripts are sorted by their InterPro domain as in (A), and neurotoxicity prediction is also reported.

Endopeptidases, such as metallo and serine proteases, constituted 43% of the upregulated transcripts in the midgut gland, consistent with their role in proteolysis and findings from prior proteomic studies [24–26]. While diverse, these transcripts were not the most highly expressed (Fig. 3b). Instead, the topmost expressed transcript was a cysteine-rich peptide orthologous to XP_054706256.1 in *U. diversus* (86% identity) with a prokineticin domain and a BlastP hit to U3-aranetoxin-Ce1a. Other highly expressed cysteine-rich peptides were predicted to be neurotoxic, with also BlastP hits to U3-araneotoxin as well as U24-ctenitoxins, the latter previously identified at the proteomic level in digestive fluids of *Uloborus* [26] and other spider species [24,25].

A notable toxin homolog expressed in the midgut gland was a Phospholipase D-like protein known as Sphingomyelinase D, a potent dermonecrotic component of *Loxosceles* venoms. Valladão et al. [26] confirmed the enzyme's presence in the midgut gland, but as a β-type variant, which lack the severe necrotic activity observed in the α-type. However, the insecticidal potential of this β variant remains untested [26].

Defensin expression: an enigmatic toxin-like component

The second most highly expressed toxin-like transcript of *U. plumipes* was a defensin with a BlastP hit to the scorpion venom peptide BmKDfsin3 from *Mesobuthus martensii* (Fig. 4). Interestingly, this defensin was not annotated in the *U. diversus* genome. BlastN searches revealed the ortholog defensin gene on chromosome 4, while searches against the recently available *U. plumipes* genome identified four hits on four different contigs, one of which was 100% identical to our transcript.

Similarly, in another Uloboridae species, *Octonoba sinensis* [27], three defensin genes were found: the ortholog of *U. plumipes* on chromosome 4, a second gene on the same

chromosome with less identity, and another on chromosome 3. The ortholog on
chromosome 4 is embedded within the gene "g1332" which seems to have been
misannotated since its coding sequence does not match any spider genes except for a
retrotransposon in *Araneus ventricosus* with 32% identity. Mapping available SRA libraries
from multiple *O. sinensis* tissues to the added defensin annotation showed high expression
in the 'abdomen' sample (Fig. S5). It remains uncertain if the midgut gland was dissected
within the abdomen sample, although the expression pattern suggests that it likely was.
Beyond Uloboridae, a defensin gene is present on chromosome 3 in *P. tepidariorum*, as well
as in other spider genomes. Notably, defensins were found in the venom glands of
Cupiennius species, distinct from those expressed in the haemolymph [28]. Phylogenetic
analysis and Alphafold 3 [29] predictions of the mature region of the defensin sequence
suggest two copies (Fig. 4A): one likely expressed in haemolymph on chromosome 3 and
another more derived on chromosome 4, highly expressed in the midgut gland of
Uloboridae spiders, resembling the venom gland defensin of *Cupiennius* (Fig. 4B, C).
205
206

Figure 4. Phylogenetic analysis and 3D structure prediction of defensin. A: Neighbour-
joining tree with branch bootstrap values and multiple sequence alignment of the mature
region of the defensin found strongly upregulated in the midgut gland of *U. plumipes*. B:
Alphafold 3 structure prediction of the defensins. Note the similarity between the copy in *U.*
plumipes and that found in the venom glands of *C. salei*. C: Overlaid 3D structure predictions
of defensins from *U. plumipes* and that from other species. The colour-code for the species
is the same as in (B).
215

Toxin-like transcripts in other body parts
Outside the midgut gland, toxin-like transcripts were less highly expressed, although
significantly upregulated to a specific body part. For instance, a U8-agatoxin-Ao1a-like
sequence with a knottin domain, which is one of the most abundant domains in spider

venoms [6], was highly expressed in the prosoma. However, it presented a disruption in the
canonical cysteine framework (C-CXC-CC-CXC-CXC-C), with the absence of the second and
tenth cysteine residues (Fig. S6).
In the silk glands, we detected various acetylcholinesterase genes. Acetylcholinesterase is an
enzyme family typically involved in neurotransmitter breakdown at postsynaptic
neuromuscular junctions. While its presence in the venoms of spiders, centipedes, and
other organisms has been noted, its role in spider venom, as well as in the silk, remains
uncertain. Nevertheless, it has been hypothesized to serve a trophic function [30].
Finally, several ferritin transcripts ranked among the most highly expressed in the whole
transcriptome. Ferritin, commonly reported in cone snail venom glands [31,32], has also
been detected in spider digestive fluids [24], suggesting a broader physiological role.
231
Midgut gland extracts are insecticidal
The upregulation of toxin-like transcripts in the midgut gland, particularly of peptides with
predicted neurotoxic activity, supports the hypothesis that Uloboridae spiders have shifted
venom-like components into their digestive fluids to aid in prey immobilisation. Since
Uloboridae do not inject venom, these potent digestive fluids may play a central role in
incapacitating prey. To test this hypothesis, we compared the midgut gland extracts of *U.*
plumipes and *P. tepidariorum* by injecting them into *Drosophila suzukii*.
Both extracts were highly insecticidal, with 230 ng killing over 50% of flies within one-hour
post-injection. Even as little as 46 ng killed approximately 40% of flies within 24h, with *U.*
plumipes extracts slightly more potent, though not statistically significant (t = -0.25, df = 3.9,
p = 0.4) (Fig. 4A).
243
Midgut gland extract does not block sodium and potassium channels
Considering the high expression of transcripts with predicted neurotoxic activity, we tested
whether the midgut extract's insecticidal effect was due to neurotoxins targeting sodium
and potassium channels, which are among the most common voltage-gated ion channel
targets of spider venoms [6]. Using a two-electrode voltage clamp assay, we examined
whether 10 µg of extract inhibited sodium channels from *Blatella germanica* (BgNav1) or
potassium channels from *D. melanogaster* (Shaker IR).
The extracts showed no inhibitory effects on either channel (Fig. 4B). Furthermore, even at
concentrations up to 30 µg, the extracts did not induce cytolysis in oocytes, indicating that
the midgut secretions do not cause membrane damage or cell integrity disruption, as pore-
forming toxins do.
255

Figure 5. Bioassays of spider midgut gland extract. A: Survival rates of *D. melanogaster* injected with midgut gland extracts from *U. plumipes* and *P. tepidariorum* using 46nl of 5mg/ml (230ng) and 1mg/ml (46ng) of protein extract. The negative control is represented by the untreated group and flies injected with PBS. The positive control is represented by injections of 70% ethanol. B: Electrophysiological experiments with *U. plumipes* (red) and *P. tepidariorum* (purple) midgut gland extracts as representative traces of currents through the potassium (Shaker IR) and sodium (BgNav1) ion channels after application of 10 μg of extract.

Discussion

Spiders are efficient predators, relying on foraging webs and potent venoms to capture and subdue prey. Nearly all spiders inject venom, with spiders of the Uloboridae family being a remarkable exception. For nearly a century, uloborids were believed to lack venom glands, a notion originating from an early anatomical sketch [14]. However, this assumption had remained largely unexamined until now. In this work, we investigated whether uloborids have indeed lost their venom apparatus and whether venom-associated toxins were repurposed in other secretions.

Uloboridae spiders have effectively lost their venom apparatus but not their toxins

Our histological analyses confirmed the absence of venom glands in the prosoma of *U. plumipes*. Furthermore, the venom duct opening at the fang tip was absent, indicating complete loss of the venom system. Interestingly, we observed small pores on the fangs, a

feature also noted in other spiders [20], though their function remains unclear. While histology confirmed the absence of venom structures, our transcriptomic analysis revealed a more complex story: venom-like components are still be present and expressed, specifically in the midgut gland, the organ secreting digestive fluids. A proteomic study on a related species confirmed the presence of similar toxins [26], and our functional assays demonstrated the lethal effects of midgut gland extracts on *Drosophila* flies. Even at low concentrations, these extracts were highly insecticidal, underscoring their potency. This discovery suggests that uloborids may have repurposed venom toxins, integrating them into their digestive system for prey immobilisation.

Toxins in spider digestive fluids: an evolutionary adaptation?

The presence of venom components in digestive fluids is not unique to Uloboridae. Studies on venomous species of different spider taxa, such as the golden orb-weaver *Nephila cruentata* [24], or the Brazilian white-knee tarantula *Acanthoscurria geniculata* and the velvet spider *Stegodyphus mimosarum* [25], have detected similar proteins in their digestive secretions, albeit in varying proportions. For instance, 66% of venom components in *S. mimosarum* were also found in digestive fluids, compared to only 9% in *A. geniculata* [25]. Conversely, digestive enzymes have been detected in venoms, possibly starting the extra-oral digestion [25]. Why toxins exist in digestive fluids remains unclear, with no definitive hypotheses proposed thus far.

We suggest that this dual presence serves as an evolutionary safety mechanism, ensuring prey immobilisation when venom is inadequate. In large spiders such as *A. geniculata*, which has strong jaws and can kill prey just with the strength of their bite alone, fewer toxins are found in the digestive fluids. While their 'safety mechanism' is brute force, smaller spiders may rely on a dual chemical weapon. Our functional assay demonstrated that also the midgut extract of the venomous *P. tepidariorum* was equally potent to that of *U. plumipes* in killing *Drosophila* flies.

How uloborids kill their prey

We propose that Uloboridae spiders have not evolved novel toxic secretions but rather altered how existing secretions are deployed. Unlike typical spiders, which apply digestive fluids after venom injection and only near their mouthparts, uloborids exhibit a unique feeding behaviour. They wrap their prey extensively in silk and cover the entire body with digestive fluids [15–18]. It is only after the application of digestive fluids that prey death occurs [19]. We can think of this external application of toxin secretion as mimicking the internal diffusion of venom. Proteolytic enzymes, such as astacin-like metalloproteases and serine proteases, degrade intersegmental membranes, acting as spreading factors and facilitating toxin entry or potentially causing death through enzymatic activity alone [33].

A key question remains: do uloborids digestive secretion contain neurotoxins capable of paralysing prey? And if so, are these toxins unique to Uloboridae, or shared with venomous species?

While our transcriptomic analysis of *U. plumipes* revealed upregulation of genes with predicted neurotoxic activity in the midgut gland, their exact role remains speculative. Functional assays on sodium (Shaker IR) and potassium (BgNav1) channels showed no inhibition of these commonly targeted voltage-gated ion channels. However, many potential targets, such as calcium channels, acid-sensing ion channels, or glutamate receptors, remain

327 untested [6]. Furthermore, toxin efficacy often depends on isoform specificity. For example,
328 scorpion toxin MeuTXKα1 effectively blocks rKv1.1, hKv1.3, and *Shaker* IR channels but not
329 other isoforms such as rKv1.2, rKv1.4, rKv1.5, rKv1.6, or hERG [34]. This suggests that
330 uloborid neurotoxins may act on yet unidentified targets, necessitating further exploration.
331

A promising candidate: defensin
Among the identified candidate toxins, defensin stood out due to its high expression,
particularly in the midgut gland. Defensins are small peptides with antimicrobial properties
found in various organisms, including plants, arthropods, and vertebrates. In scorpions,
defensins have dual roles: they exhibit antimicrobial activity and block potassium channels
(Kv1.1, Kv1.2, Kv1.3 and SK3), functioning similarly to neurotoxins like OSK1 and ScyTx [35].
Notably, slight modifications can transform an insect defensin into a potent scorpion-like
potassium channel inhibitor [36]. Despite the translated sequence of *U. plumipes* defensin
being predicted neurotoxic, our electrophysiology experiment showed no inhibition on the
BgNav1 channel. However, it is possible that other ion channels are targeted, as discussed
earlier.
Beyond scorpions, defensins have also been detected in the venom glands of the bromeliad
spider species *C. salei* and *C. getazi* [28]. Our genomic and phylogenetic analyses show that,
after duplication, the copy on chromosome 4 underwent upregulation in the midgut gland
of Uloboridae and the venom glands of *Cupiennius*, while the copy on chromosome 3
appears to maintain the original immunity role. Further genomic studies using the newly
available spider genomes could provide more clarity on the evolutionary history of this
intriguing gene family. Our BlastN searches also highlight the challenges in annotating these
small peptides, which may hinder proteomic analyses. Defensins have not been reported in
the digestive fluids of various spiders [25], including *Uloborus* sp. [26], and this absence may
reflect technical limitations due to the lack of the defensin sequence in the databases used
for peptide identification rather than a true absence in secretion.
354

Conclusions
This study investigated the evolutionary loss of venom in the spider *U. plumipes*, confirming
the absence of a venom apparatus in Uloboridae while revealing the high expression of
venom toxins in the midgut gland. These findings demonstrate that spider toxins are not
exclusively confined to specialised venom-secreting glands but are also play a role in the
digestive system. This supports an evolutionary link between the two systems, suggesting
that toxins may have initially served digestive functions before being co-opted for venom
use. In venomous spiders, toxins in the digestive fluids likely act as a secondary mechanism
for effective prey immobilisation, while Uloboridae have adapted this trait as their primary
strategy by dispersing toxic fluids across the prey's body. This study highlights the
complexity of spider biology and emphasises the importance of investigating toxin
expression across tissues to fully understand their evolutionary and functional roles.
367

Methods
Specimens
Individuals of *Uloborus plumipes* were collected from greenhouses in Viernheim and Merzig
in Germany, and in Lausanne, Switzerland. *Parasteatoda tepidariorum* spiders are from a
laboratory colony which is kept at 27°C with 70% humidity on a gradual light/dark cycle of
16/8 hours and fed twice a week with *Musca domestica* flies.

374
Morphological and histological analyses
To verify the absence of venom-producing glands in *U. plumipes* we performed a histological analysis of the chelicerae and anterior part of the prosoma. We dissected five individuals of *U. plumipes* directly in the fixative, Karnovsky's solution [37], and fixed the chelicerae with part of the prosoma overnight. After washing in 0.1M phosphate buffer, the samples were post-fixed in 2% osmium tetroxide solution for two hours and dehydrated using a graded series of ethanol. Embedding was carried out using Embed812 resin embedding kit (Science Services GmbH, München, Germany). During the final step, samples were transferred into a "VacuTherm" vacuum heating cabinet (Thermo Fisher Scientific, Waltham, Massachusetts, USA) and incubated at 40 °C and 100 mbar for 3 × 30 min. Polymerization of the resin blocks was carried out in a heating cabinet at 60 °C for a minimum of 24 h. Semi-thin sections were obtained with a Leica UC6 ultra-microtome (Leica Microsystems GmbH, Wetzlar, Germany), with a DiATOME histo Jumbo diamond knife (Diatome Ltd., Nidau, Switzerland) at a thickness of 700 nm. Staining was done with toluidine blue at 70 °C, and images were obtained with a Fritz Slide Scanner (PreciPoint GmbH, Freising, Germany).

Individuals of *P. tepidariorum* were fixed in 2.5% glutaraldehyde for 1 hour at room temperature and subsequently fixed overnight at 4°C. After several washes with 0.1 M phosphate buffer, the samples were postfixed in 2% osmium tetroxide solution for 1 hour at room temperature, then washed with water, dehydrated through a graded series of acetone and subsequently a graded series of resin, and finally embedded using Spurr Low-Viscosity Embedding kit (SIGMA). The embedded spiders were incubated in resin blocks at 60°C for 48 hours. Semi-thin sections were obtained with a Leica EM UC7 Ultramicrotome (Leica Microsystem), with a DiATOME diamond knife (Diatome Ltd., Switzerland) at 700 nm thickness. Sections were stained with 0.5% toluidine blue, and images were acquired using Zeiss Axio Imager Z2 (Leica Microsystem).

Scanning electron microscopy
We used three individuals to examine the fang morphology. Specifically, we aimed at verifying whether the opening of the venom duct at the tip of the fangs were present, or whether they disappeared as a consequence of the loss of the venom-secreting apparatus.
The spiders were fixed in liquid nitrogen, placed on an aluminium holder with double-sided carbon adhesive, and examined with a scanning electron microscope (Quanta FEG 250, TFS) using the environmental mode (partial pressure 80Pa). The detectors used were the Large Field Detector and the Backscattered electron detector (BSED) at 10kV spot 4, working distance between 7.9 and 6.6mm.

RNA-seq
Tissue samples of chelicerae, prosoma, midgut gland, silk gland and gonads (ovaries) were dissected from approximately 12 adult individuals. To obtain enough RNA, for each tissue, we pooled multiple samples for a total of three replicates except the ovaries for which we had only one. Total RNA was isolated using the TRIzol[TM] Plus RNA Purification kit (ThermoFisher) following manufacturer's instructions, with an additional on-column DNA purification step. Thirteen cDNA libraries were generated with the TruSeq RNA Sample Preparation kit (Illumina) with 150 read length, followed by pair-end sequencing on an Illumina NovaSeq at the Genomic Technologies Facility of the University of Lausanne, Switzerland.

Assembly and annotation

Raw reads were assessed with FastQC v0.11.9 [38] and quality-filtered with Fastp v0.22.0 [39]. Reads shorter than 30bp were discarded. As no genome for Uloboridae spiders was available at time of the project, all reads from all tissues were concatenated and used for *de novo* assembly with SPAdes v3.15.3 [40]. We adopted a multi-step quality filtering approach to prune the raw transcriptome assembly. First, we used the program Borf v1.2 [41] to predict open reading frames (ORFs) and retained only sequences with a minimum length of 20 amino acids and a complete ORF. Next, we annotated the filtered amino acid sequences with BlastP [42] against multiple databases, including the NCBI non-redundant pre-formatted Refseq database, UniProt/SwissProt [43], UniProt-ToxProt [23], Arachnoserver [44], and a customised database consisting of 11 spider genomes (Supplementary Table S3). All databases were downloaded on July 4[th], 2022. Protein families and domains were predicted with InterProScan v5.51.85.0 [45], and signal peptides were detected using SignalP-6.0 [46]. Only transcripts with a BlastP hit to at least one database and e-value < 1e-5 were retained. We further reduced redundancy by clustering all nucleotide sequences with > 99% identity using CD-HIT v4.8.1 (Fu et al. 2012). Orthologous clusters with *Uloborus diversus* (GCF_026930045) genes [47] were identified using the web platform OrthoVenn3 [22]. The completeness of the final assembled transcriptome was assessed by OMArk [21] by calculating the overlap of the non-redundant annotated genes and conserved ancestral gene set of the phylum Arthropod.

Toxin-like transcripts identification

We assigned a transcript as toxin-like if its best hit, defined as having the highest bitscore, was a protein listed in the UniProt-ToxProt database [23]. Furthermore, based on the assumption that toxins should have biased expression specifically to the gland that secretes them, we further minimise the chances of false positives by running a differential expression analysis and keeping only those toxin-like transcripts which were significantly upregulated (see below). Presence of a knottin domain was tested with the tool Knotter 1D from the Knottin database website [48], and neurotoxic activity was predicted using NT_estimation, a deep learning approach which uses a peptide data augmentation method to improves the recognition of spider neurotoxic peptides via a convolutional neural network model (Lee et al. 2021). The putative toxins were classified into their corresponding protein families based on their predicted InterPro protein domain [45]

Expression level quantification

Transcript abundances were quantified using Kallisto v0.48.0 (Bray et al., 2016) with default parameters for paired-end reads. Downstream analyses were conducted in R version 4.1.3 [49]. Count distribution across libraries was inspected with the package *vioplot* v 0.3.7 (Adler et al., 2021) and genes with TPM value >= 1 in at least one library were kept for further analyses. Consistency of expression patterns between samples from the same tissue was assessed by means of principal component analysis, and the percentage of explained variance by each component was calculated using the function fviz_eig in *Ade4* v1.7.19 (Dray and Dufour, 2007).

Differential expression analysis was performed in *Sleuth* v0.30.0 (Pimentel et al., 2017) and the fold change (FC) of transcripts was calculated as the ratio between the highest and the

468 second highest expression values. Only putative toxin transcripts with fold change >= 2 and
469 q-val < 0.05 were retained.
470 Because *Sleuth* automatically filter out genes which are exclusively expressed in one tissue
471 type, we also checked those which did not pass *Sleuth* filter and kept all the sequences with
472 FC >= 2.
473
Genomic investigation of defensin
Considering the high expression levels of defensin in the midgut gland, and the lack of
ortholog with *U. diversus*, we thought to investigate further this putative toxin in other
spider species, specifically other Uloboridae whose genome became available during the
course of the study. First, we performed BlastN [42] searches against the whole genome of
the Ulobordae *U. plumipes*, *U. diversus,* and *Octonoba sinensis*, as well as *P. tepidariorum*
and the NCBI non-redundant database. We used SeaView v5.0.4 [50] for multisequence
alignment with MUSCLE v3.8.31 [51] and to obtain a Neighbour-Joining tree with bootstrap
from 100 replicates. Three-dimensional (3D) structures were predicted using Alphafold v3
[29]. Expression levels were examined for those species with multi-tissue RNA-Seq data, i.e.
O. sinensis and *P. tepidariorum*. For the latter, we used the expression quantification from
Hassan et al. [52], while for *O. sinensis* we aligned reads from a subset of SRA (see Table S4)
using STAR v2.7.10b [53] and quantified transcript abundances with Kallisto v0.48.0 [54]
after adding the defensin transcripts to the annotation. As the defensin was embedded into
gene "g1332" which seem to be spurious annotation, we performed the quantification also
after removing this gene.
490
Midgut gland protein extraction and *in vivo* experiments
Dissected midgut gland tissues from *U. plumipes* and *P. tepidariorum* were physically
disrupted with a Potter-Elvehjem homogeniser as previously described by Valladão et al.
[26] Briefly, samples were homogenized in a falcon tube containing 500μL of water and then
centrifuged at 16.000g for 30 min at 4°C. The supernatant containing the digestive fluids
was collected and lyophilised using a Beta 2-8 LSCplus (Christ) freeze drier. Lyophilised
midgut extracts were used for downstream experiments. Protein content was quantified
using a BCA-Protein-Assay kit (Merck).
Insecticidal activity was assessed *in vivo* as described earlier [55]. For the injections,
concentrations of 5 mg/mL and 1 mg/mL of the extracts were prepared in PBS. *Drosophila*
suzukii flies from a laboratory stock, originally sourced from Ontario, Canada, were reared in
a climate chamber at 26 °C with 60% humidity and a 12 h photoperiod. They were fed on
10.8% soybean and cornmeal mix, 0.8% agar, 8% malt, 2.2% molasses, 1% nipagin and
0.625% propionic acid media. For the insecticidal assay, we used 4–6-day old adult flies.
Flies were anaesthetized on a CO_2-Pad (Inject+Matic). Volumes of 46 nl were injected
intrathoracically using glass capillaries pulled on a P-2000 Laser-Based Micropipette Puller
(Sutter Instrument) held on a Nanoject II device (Drummond Scientific). Injections were
performed under a Stemi 508 Stereomicroscope (Zeiss). Negative controls consisted of
untreaded flies and flies injected with 46 nl of PBS, while positive controls were injected
with 46 nl of 70% ethanol. Batches of 10 flies then incubated in Ø 29 × 95 mm vials with
foam stoppers (Nerbe Plus) filled 1/16 with food media. Survival rates and signs of paralysis
were assessed 1h, 3h and 24h post injection.
513
514 **Two-electrode voltage clamp electrophysiology**

We investigated neurotoxicity of the digestive fluids by examining whether 10 μg of midgut extract would inhibit potassium and the sodium channels. Recordings were performed at room temperature (18–22 °C) using a Geneclamp 500 amplifier (Molecular Devices, San Jose, CA, USA) controlled by a pClamp data acquisition system (Axon Instruments, San Jose, CA USA). Whole-cell currents from *Xenopus laevis* oocytes were recorded 1–4 days after cRNA injection. The bath solution composition was ND96. Voltage and current electrodes were filled with 3 M KCl. Resistances of both electrodes were kept at 0.7–1.5 MΩ. Elicited currents were sampled at 1 kHz and filtered at 0.5 kHz (for potassium currents) or sampled at 20 kHz and filtered at 2 kHz (for sodium currents) using a four-pole low-pass Bessel filter. Leak subtraction was performed using a −P/4 protocol.

For the electrophysiological measurements, we used a 2 s ramp protocol from −120 to +80 mV, with cells clamped at −120 mV holding potential. Shaker IR currents were evoked by 500 ms step depolarizations to 0 mV, followed by a 500 ms pulse to −50 mV from a holding potential of −90 mV. Sodium current traces were evoked by a 100 ms depolarization to 0 mV. All data were obtained in at least five independent experiments (n ≥ 5). Non-injected oocytes without expressing any type of ion channel were used to check whether the midgut extracts were cytotoxic to oocytes. Animal experiments using *X. laevis* were approved by the Animal Ethics Committee of KU Leuven in accordance with EU Council Directive 2010/63/EU.

Expression of ion channels in *X. Laevis* oocytes

The following genes encoding ion channel subunits were expressed in *X. laevis* oocytes: the voltage-gated potassium channel, Shaker-IR from *Drosophila melanogaster* and the voltage-gated sodium channel BgNav1 from *Blattella germanica*. Linearised plasmids bearing the ion channel genes were transcribed using the mMESSAGE mMACHINE T7 transcription kits (Ambion, Austin, TX, USA) to prepare the respective cRNA. The harvesting of stage V–VI oocytes from anesthetized female *X. laevis* frogs was described previously [56]. Oocytes were injected with 50 nl of cRNA at a concentration of 1 ng/nl using a micro-injector (Drummond Scientific, Broomall, PA, USA). The oocytes were incubated at 16°C in ND96 solution containing (in mM): NaCl, 96; KCl, 2; CaCl2, 1.8; MgCl2, 2; and HEPES, 5 (pH 7.4), supplemented with 50 mg/l gentamicin sulfate.

Acknowledgements

This project has received funding from the European Union's Horizon 2020 research and innovation programme under Marie Skłodowska-Curie Grant Agreement 845674 to G.Z. and the Etat de Vaud to M.R-R. We are grateful to Daniel Gritz and Vanessa Oehmig from the German Arachnological Society (DeArGe) who provided specimens of *U. plumipes* to our study. We thank J. Marquis from GTF for support with library preparation and sequencing.

Author contributions

X.P. performed the RNA extraction and the transcriptome annotation and analysis. L.D., J.D, and T.L. performed the *in vivo* experiments. T.D., P.M., and A.H. performed the morphological and histological analyses. S.P. and J.T. performed the electrophysiology experiments. A.M. performed the electron microscopy. M.R-R. contributed to the transcriptomic analysis. G.Z. and T.L. conceived the study. G.Z. contributed to the transcriptomic analysis and wrote the manuscript, with input from all authors.

References

561
562
563
564
1. Schendel V, Rash LD, Jenner RA, Undheim EAB. 2019 The diversity of venom: The
importance of behavior and venom system morphology in understanding its ecology and
evolution. *Toxins* **11**, 666. (doi:10.3390/toxins11110666)

2. Modica MV *et al.* 2021 The new COST Action European Venom Network (EUVEN)—
synergy and future perspectives of modern venomics. *GigaScience* **10**, giab019.
(doi:10.1093/gigascience/giab019)

3. Puillandre N, Fedosov AE, Kantor YI. 2017 Systematics and evolution of the Conoidea. In
Evolution of Venomous Animals and Their Toxins (ed A Malhotra), pp. 367–398.
Dordrecht: Springer Netherlands. (doi:10.1007/978-94-007-6458-3_19)

4. Vassilevski AA, Kozlov SA, Grishin EV. 2009 Molecular diversity of spider venom.
Biochem. Mosc. **74**, 1505–1534. (doi:10.1134/S0006297909130069)

5. King GF, Hardy MC. 2013 Spider-venom peptides: Structure, pharmacology, and
potential for control of insect pests. *Annu. Rev. Entomol.* **58**, 475–496.
(doi:10.1146/annurev-ento-120811-153650)

6. Lüddecke T, Herzig V, von Reumont BM, Vilcinskas A. 2021 The biology and evolution of
spider venoms. *Biol. Rev.* **97**, 163-178. (doi:10.1111/brv.12793)

7. Arbuckle K, Harris RJ. 2021 Radiating pain: Venom has contributed to the diversification
of the largest radiations of vertebrate and invertebrate animals. *BMC Ecol. Evol.* **21**, 150.
(doi:10.1186/s12862-021-01880-z)

8. Nisani Z, Hayes WK. 2011 Defensive stinging by *Parabuthus transvaalicus* scorpions: Risk
assessment and venom metering. *Anim. Behav.* **81**, 627–633.
(doi:10.1016/j.anbehav.2010.12.010)

9. Nelsen DR, Kelln W, Hayes WK. 2014 Poke but don't pinch: Risk assessment and venom
metering in the western black widow spider, *Latrodectus hesperus*. *Anim. Behav.* **89**,
107–114. (doi:10.1016/j.anbehav.2013.12.019)

10. Cooper AM, Nelsen DR, Hayes WK. 2017 The Strategic use of venom by spiders. In
Evolution Of Venomous Animals And Their Toxins (ed A Malhotra), pp. 145–166.
Dordrecht: Springer Netherlands. (doi:10.1007/978-94-007-6458-3_13)

11. Morgenstern D, King GF. 2013 The venom optimization hypothesis revisited. *Toxicon* **63**,
120–128. (doi:10.1016/j.toxicon.2012.11.022)

12. Li M, Fry BG, Kini RM. 2005 Eggs-only diet: Its implications for the toxin profile changes
and ecology of the marbled sea snake (Aipysurus eydouxii). *J. Mol. Evol.* **60**, 81–89.
(doi:10.1007/s00239-004-0138-0)

13. Wright JJ. 2017 Evolutionary history of venom glands in the Siluriformes. In *Evolution of*
Venomous Animals and Their Toxins (ed A Malhotra), pp. 279–301. Dordrecht: Springer
Netherlands. (doi:10.1007/978-94-007-6458-3_9)

14. Millot J. 1931 Les glandes venimeuses des Araneides. *Ann. Sci. Nat. Zool.* **14**, 113–147.

15. Lubin YD. 1986 Web building and prey capture in the Uloboridae. *SpidersWebs Behav.*
Evol., 132–171.

16. Opell BD. 1988 Prey handling and food extraction by the triangle-web spider *Hyptiotes*
cavatus (Uloboridae). *J. Arachnol.* **16**, 272–274.

17. Eberhard WG, Barrantes G, Weng J-L. 2006 The mystery of how spiders extract food
without masticating prey. *Bull.-Br. Arachnol. Soc.* **13**, 372.

18. Eberhard WG, Barrantes G, Weng J-L. 2006 Tie them up tight: wrapping by *Philoponella*
vicina spiders breaks, compresses and sometimes kills their prey. *Naturwissenschaften*
93, 251–254. (doi:10.1007/s00114-006-0094-1)

19. Weng J-L, Barrantes G, Eberhard WG. 2006 Feeding by *Philoponella vicina* (Araneae,
Uloboridae) and how uloborid spiders lost their venom glands. *Can. J. Zool.* **84**, 1752–
1762. (doi:10.1139/z06-149)

20. Moon M-J, Yu M-H. 2007 Fine structure of the chelicera in the spider *Nephila clavata*.
Entomol. Res. **37**, 167–172. (doi:10.1111/j.1748-5967.2007.00108.x)

21. Nevers Y, Rossier V, Train CM, Altenhoff A, Dessimoz C, Glover N. 2022 Multifaceted
quality assessment of gene repertoire annotation with OMArk. *Nat. Biotechnol.*,
2022.11.25.517970. (doi:10.1101/2022.11.25.517970)

22. Sun J, Lu F, Luo Y, Bie L, Xu L, Wang Y. 2023 OrthoVenn3: an integrated platform for
exploring and visualizing orthologous data across genomes. *Nucleic Acids Res.* **51**,
W397–W403. (doi:10.1093/nar/gkad313)

23. Jungo F, Bougueleret L, Xenarios I, Poux S. 2012 The UniProtKB/Swiss-Prot Tox-Prot
program: A central hub of integrated venom protein data. *Toxicon* **60**, 551–557.
(doi:10.1016/j.toxicon.2012.03.010)

24. Fuzita FJ, Pinkse MWH, Patane JSL, Verhaert PDEM, Lopes AR. 2016 High throughput
techniques to reveal the molecular physiology and evolution of digestion in spiders.
BMC Genomics **17**, 716. (doi:10.1186/s12864-016-3048-9)

25. Walter A, Bechsgaard J, Scavenius C, Dyrlund TS, Sanggaard KW, Enghild JJ, Bilde T. 2017
Characterisation of protein families in spider digestive fluids and their role in extra-oral
digestion. *BMC Genomics* **18**, 600. (doi:10.1186/s12864-017-3987-9)

26. Valladão R, Neto OBS, de Oliveira Gonzaga M, Pimenta DC, Lopes AR. 2023 Digestive
enzymes and sphingomyelinase D in spiders without venom (Uloboridae). *Sci. Rep.* **13**,
2661. (doi:10.1038/s41598-023-29828-x)

27. Zhang Y *et al.* 2024 A trade-off in evolution: the adaptive landscape of spiders without
venom glands. *GigaScience* **13**, giae048. (doi:10.1093/gigascience/giae048)

28. Kuhn-Nentwig L, Langenegger N, Heller M, Koua D, Nentwig W. 2019 The dual prey-
inactivation strategy of spiders—in-depth venomic analysis of Cupiennius salei. *Toxins*
11, 167. (doi:10.3390/toxins11030167)

29. Abramson J *et al.* 2024 Accurate structure prediction of biomolecular interactions with
AlphaFold 3. *Nature* **630**, 493–500. (doi:10.1038/s41586-024-07487-w)

30. Dresler J, Avella I, Damm M, Dersch L, Krämer J, Vilcinskas A, Lüddecke T. 2024 A
roadmap to the enzymes from spider venom: biochemical ecology, molecular diversity,
and value for the bioeconomy. *Front. Arachn. Sci.* **3**. (doi:10.3389/frchs.2024.1445500)

31. Hu H, Bandyopadhyay PK, Olivera BM, Yandell M. 2011 Characterization of the *Conus*
bullatus genome and its venom-duct transcriptome. *BMC Genomics* **12**, 60.
(doi:10.1186/1471-2164-12-60)

32. Abalde S, Tenorio MJ, Afonso CML, Zardoya R. 2018 Conotoxin diversity in *Chelyconus*
ermineus (Born, 1778) and the convergent origin of piscivory in the Atlantic and Indo-
Pacific cones. *Genome Biol. Evol.* **10**, 2643–2662. (doi:10.1093/gbe/evy150)

33. Dresler J, Avella I, Damm M, Dersch L, Krämer J, Vilcinskas A, Lüddecke T. 2024 A
roadmap to the enzymes from spider venom: biochemical ecology, molecular diversity,
and value for the bioeconomy. *Front. Arachn. Sci.* **3**. (doi:10.3389/frchs.2024.1445500)

34. Zhu S, Peigneur S, Gao B, Luo L, Jin D, Zhao Y, Tytgat J. 2011 Molecular diversity and
functional evolution of scorpion potassium channel toxins. *Mol. Cell. Proteomics* **10**, S1–
S11. (doi:10.1074/mcp.M110.002832)

35. Meng L *et al.* 2020 Ion channel modulation by scorpion hemolymph and its defensin
ingredients highlights origin of neurotoxins in telson formed in Paleozoic scorpions. *Int.*
J. Biol. Macromol. **148**, 351–363. (doi:10.1016/j.ijbiomac.2020.01.133)

36. Zhu S, Peigneur S, Gao B, Umetsu Y, Ohki S, Tytgat J. 2014 Experimental conversion of a
defensin into a neurotoxin: implications for origin of toxic function. *Mol. Biol. Evol.* **31**,
546–559. (doi:10.1093/molbev/msu038)

37. Karnovsky MJ. 1965 A formaldehyde glutaraldehyde fixative of high osmolality for use in
electron microscopy. *J Cell Biol* **27**, 1A-149A.

38. Andrews S. 2010 FastQC: A quality control tool for high throughput sequence data. See
https://www.bioinformatics.babraham.ac.uk/projects/fastqc/ (accessed on 4 June
2021).

39. Chen S, Zhou Y, Chen Y, Gu J. 2018 fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**, i884–i890. (doi:10.1093/bioinformatics/bty560)

40. Bushmanova E, Antipov D, Lapidus A, Prjibelski AD. 2019 rnaSPAdes: a de novo
transcriptome assembler and its application to RNA-Seq data. *GigaScience* **8**, giz100.
(doi:10.1093/gigascience/giz100)

41. Signal B, Kahlke T. 2021 Borf: Improved ORF prediction in de-novo assembled
transcriptome annotation. *BioRxiv*, 2021.04.12.439551.
(doi:10.1101/2021.04.12.439551)

42. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009
BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421. (doi:10.1186/1471-
2105-10-421)

43. The UniProt Consortium. 2023 UniProt: the universal protein knowledgebase in 2023.
Nucleic Acids Res. **51**, D523–D531. (doi:10.1093/nar/gkac1052)

44. Pineda SS *et al.* 2018 ArachnoServer 3.0: an online resource for automated discovery,
analysis and annotation of spider toxins. *Bioinforma. Oxf. Engl.* **34**, 1074–1076.
(doi:10.1093/bioinformatics/btx661)

45. Quevillon E, Silventoinen V, Pillai S, Harte N, Mulder N, Apweiler R, Lopez R. 2005
InterProScan: protein domains identifier. *Nucleic Acids Res.* **33**, W116–W120.
(doi:10.1093/nar/gki442)

46. Teufel F *et al.* 2022 SignalP 6.0 predicts all five types of signal peptides using protein
language models. *Nat. Biotechnol.* **40**, 1023–1025. (doi:10.1038/s41587-021-01156-3)

47. Miller J, Zimin AV, Gordus A. 2023 Chromosome-level genome and the identification of
sex chromosomes in *Uloborus diversus*. *GigaScience* **12**, giad002.
(doi:10.1093/gigascience/giad002)

48. Postic G, Gracy J, Périn C, Chiche L, Gelly J-C. 2018 KNOTTIN: the database of inhibitor
cystine knot scaffold after 10 years, toward a systematic structure modeling. *Nucleic
Acids Res.* **46**, D454–D458. (doi:10.1093/nar/gkx1084)

49. R Core Team. 2019 R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. URL http://www.R-project.org/.
(doi:URL http://www.R-project.org/)

50. Gouy M, Guindon S, Gascuel O. 2010 SeaView Version 4: A Multiplatform graphical user
interface for sequence alignment and phylogenetic tree building. *Mol. Biol. Evol.* **27**,
221–224. (doi:10.1093/molbev/msp259)

51. Edgar RC. 2004 MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res.* **32**, 1792–1797. (doi:10.1093/nar/gkh340)

52. Hassan A, Blakeley G, McGregor AP, Zancolli G. 2024 Venom gland organogenesis in the
common house spider. *Sci. Rep.* **14**, 15379. (doi:10.1038/s41598-024-65336-2)

53. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
Gingeras TR. 2013 STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21.
(doi:10.1093/bioinformatics/bts635)

54. Soneson C, Love MI, Robinson MD. 2015 Differential analyses for RNA-seq: transcript-
level estimates improve gene-level inferences. *F1000Research* **4**, 1521.
(doi:10.12688/f1000research.7563.1)

55. Fischer ML *et al.* 2024 Divergent venom effectors correlate with ecological niche in
neuropteran predators. *Commun. Biol.* **7**, 981. (doi:10.1038/s42003-024-06666-9)

56. Peigneur S *et al.* 2021 Small cyclic sodium channel inhibitors. *Biochem. Pharmacol.* **183**,
114291. (doi:10.1016/j.bcp.2020.114291)

714