Title: A whole organism map of X-inactivation reveals tissue-specific escape

as an exceptionally rare event in humans

Authors: Björn Gylemo[1] and Colm E. Nestor[1*]

Email: bjorn.gylemo@liu.se and colm.nestor@liu.se

Affiliations:

[1]Crown Princess Victoria Children´s Hospital, Department of Biomedical and Clinical

Sciences (BKV), Linköping University, Linköping, Sweden

*Corresponding author. Email: Colm.Nestor@liu.se

Abstract

As females are mosaic for X-inactivation, direct determination of X-linked allelic expression

in bulk tissues is typically unfeasible. Using females that are non-mosaic for X-inactivation

(nmXCI) as a powerful and natural genetic system we directly determined X-inactivation status

of 401 X-linked genes across 32 normal tissues. We reveal that nmXCI is common in the female

population and that contrary to previous reports, variable escape is an exceedingly rare event in

humans. As the most complete atlas of XCI in humans to date, our findings represent a major

advance in our understanding of human X-inactivation and will serve as a reference for

dissecting the genetic origin of sex-bias in human traits.

Keywords

X-chromosome inactivation, allele-specific expression, non-mosaic XCI

Background

Human females are typically 'mosaic' for X-inactivation; female tissues contain a mixture of

cells that have inactivated either the maternal (Xm) or paternal X-chromosome (Xp) (Fig. 1A).

Mosaicism renders analysis of X-inactivation (XCI) in primary human tissues highly complex

or unfeasible. Consequently, our knowledge of XCI across human tissues is surprisingly sparse

and typically based on indirect and imprecise measures of XCI, such as DNA methylation, sex-

biased expression or from observations in animal models (1,2). Here, we establish a unique set

of three unrelated females in the Genotype-Tissue Expression (GTEx) project in which the same

parental X-chromosome was inactivated in all cells, thereby removing the confounding effect

of mosaicism (3). Such non-mosaic (nmXCI) females enable direct determination of the XCI

status even from bulk tissue samples. Using these powerful and natural human genetic models

we could directly determine the XCI status of 401 genes across 32 normal human tissues,

representing a major advance in our understanding of human X-inactivation.

Results and Discussion

Non-mosaic X-inactivation (nmXCI) is common in human females

Females exhibiting non-mosaic XCI (nmXCI) are a powerful system in which to dissect the

XCI status of human X-linked genes, but their frequency in the general population is unknown

with current estimates ranging from 1:50 to 1:500 individuals (*4-7*). To identify nmXCI females

we first screened allele-specific expression of genes known to undergo X-inactivation (**Table**

S1) in a single tissue across all 285 female donors (**Table S2**) in the GTEx database (v8 release)

(*7*), including a single nmXCI female previously identified in GTEx (donor UPIC) as a control

(*7*). Tissues used for the screen were selected based on abundance in the entire data set. Six

female samples showed complete mono-allelic expression of inactive genes (allelic expression

> 0.4) in the selected tissue, consistent with expression from a single parental chromosome,

nmXCI (**Fig. 1B**). Extending allele-specific expression analysis to all available tissues (**Table**

S3) across the six potential nmXCI females confirmed the complete nmXCI status of three

individuals (including the nmXCI control, UPIC) (**Fig. 1C**). The remaining three potential

nmXCI females showed a mixture of mosaic and highly skewed XCI across tissues (**Fig. 1C**).

We found no obvious detrimental mutations in the coding or regulatory regions of the *XIST*

gene as previously described in a report of familial nmXCI (**see methods**) (*8*), thus the genetic

basis of nmXCI in these three individuals is unknown.

Fig. 1. Identification of females with non-mosaic X-inactivation. (**A**), The patterns of X-

chromosome inactivation (XCI) in women resulting in mosaic or non-mosaic XCI. (**B**), Single-

tissue allelic expression (AE) of 237 inactive genes (*7*) (see **Table S1**) in all 285 women in

GTEx. Tissue abbreviations can be found in **Table S9**. Each dot represents mean AE across all

inactive genes in that sample, grey ribbon represents standard error and the red box highlights

women with a mean AE > 0.4 (UPIC, 13PLJ, ZZPU, 131XW, 14PJM and 1GF9X). (**C),**

Multiple-tissue AE of the 237 inactive genes (7) (see **Table S1**) in the six women with a mean

AE > 0.4 and 6 randomly selected females (14BIN, 1211K, Y8LW, 14JG6, 1129S and 1122O).

Boxes indicate mean ± 25th and 75th inter quartile range (IQR).

A landscape of X-inactivation across normal human tissues

The ability to detect allele-specific expression is dependent on the presence of a heterozygous

variant in the gene of interest. Using both whole exome sequencing and whole genome

sequencing for all three nmXCI individuals we identified 756 heterozygous SNPs (**Fig. 2A and**

Table S4) across 401 X-linked genes (**Fig 2B, Fig. S1A and Table S4**), representing 43% of

all protein coding X-linked genes (**Fig. S1B**). For 177 of these X-linked genes this is the first

direct determination of their XCI status across multiple tissues in humans (**Fig. S1A and Table**

S5). The tissue-specific allelic expression status of X-linked genes was highly similar across all

three individuals and independent of the SNP used in each comparison, supporting the

robustness of our approach (**Fig. S1C and Fig. S1D**). As expected, genes located in the pseudo-

autosomal regions (PARs) of the X-chromosome and well-characterised nonPAR escape genes

showed clear bi-allelic expression across tissues (**Fig. 2D and Fig. 2E**), genes undergoing X-

inactivation showed mono-allelic expression (**Fig. 2F**) whereas 66 genes exhibited varying

degrees of mono- and bi-allelic expression across tissues, 'variable' escape genes (**Fig. 2G**).

Tissue-specific escape from XCI is exceptionally rare in humans

Unexpectedly, whereas PAR, escape and inactive genes identified here showed clear overlap

with previous classifications of XCI status, only ~7% of variable escape genes were similarly

classified in previous studies of XCI (**Fig. 2H**). We noted that genes previously classified as

'variable' escape were significantly less expressed than all other classes of X-linked genes

across 50 tissues in 103 randomly selected females in the GTEx, regardless of whether the

current or previous classification of XCI was used (**Fig. 2I, Fig. S1E, Fig. S1F and Table S6**).

This suggests that classification of many genes as variable escape may have reflected a lower

expression level and consequently fewer sequencing reads, which increased risk for inaccurate

determination of allelic balance. To explore this phenomenon further we manually examined

the SNP quality and read depth for all 66 variable genes across all tissues and individuals.

Whereas some known (e.g., *ANOS1*, *CLIC2, NAA10*) and novel (*ARSD, CA5B, TRAPPC2*)

genes show clear and reproducible variability in escape across tissues and individuals, most

genes previously classified as variable have expression at the limit of detection and have most

likely been miss-classified as a result (**Fig. 2J, Fig. 2K, Fig. S2A and Table S7**).

Fig. 2. An extended landscape of X-inactivation in humans. (A,B), Overlap of heterozygous

SNPs (hetSNPs) (A) and genes (B) across the three females with non-mosaic X-inactivation

(nmXCI). (**C**), Genomic location and allele-specific classification count of covered X-linked

genes. PAR = genes located in the pseudoautosomal region, biallelic = genes with an allelic

expression (AE) < 0.4 in at least 85% of individual sample and tissue, monoallelic = genes with

an AE > 0.4 in at least 85% of individual sample and tissue, variable = genes with AE < 0.4 in

more than 15% and less than 85 % of individual sample and tissues. (**D-G**), AE across tissues

in the three nmXCI females for two example genes per category: PAR genes (D), escape genes

(E), inactive genes (F), variable genes (G). Data points missing in (E, left panel) and (G, left

panel) reflect the lack of a hSNPs present in that gene in ZZPU. (**H**), Number of genes per XCI

classification and their overlap with a previous assessment (*7*). (**I**), Mean transcript-per-million

(TPM) expression of each gene categories previously described (*7*) across 103 randomly

selected females (see **Fig. S1F** for number of samples per tissue). Each dot indicates the mean

expression of all genes per tissue within each gene category. Tissues are merged on tissue

category, see **Fig. S1E** for all tissues included in GTEx. ****: p-value <= 0.0001, Benjamini-

Hochberg adjusted unpaired two-sided Wilcoxon test. (**J**), Reference and alternative allele read

counts of example novel and known (*7*) variable genes and variable genes reclassified as

inactive genes. (**K**), Reclassification of previous assessments (*7*) of XCI status of genes. (D-I),

tissue abbreviations can be found in **Table S9.**

Conclusions

Here we describe the most comprehensive map of human X-inactivation to date. This data

represents a doubling of the number of X-linked genes for which XCI status has been directly

determined, across multiple normal tissues and individuals. Our results suggest that XCI is

highly stable both within and across individuals, and that previously suggested levels of tissue-

specific escape from X-inactivation may have been vastly overestimated. The classification of

many genes as variable escape may be a consequence of a low read count, which increases the

risk for inaccurate determination of XCI status. Variability in X-linked gene expression is

important to characterise, as XCI variability would directly modify expressivity of X-linked

pathogenic variants and provide a mechanism for sex-biased expression differences in certain

biological contexts, which in turn can modify disease-risk. Moreover, our identification of

additional nmXCI females among the 285 females in the GTEx database further substantiates

recent claims of extreme skewing of XCI as a frequent and major modifier of trait penetrance

and expressivity in the human population (5,6).

Materials and Methods

<u>Allele-specific expression analysis</u>

For initial screen for identifying nmXCI females in GTEx, pre-processed single-nucleotide

polymorphisms (SNP) calling performed from whole-exome sequencing (WES) data from

GTEx was downloaded (GTEx_Analysis_2017-06-

05_v8_WholeExomeSeq_979Indiv_VEP_annot.vcf.gz, see

https://gtexportal.org/home/methods for specifics on how GTEx processed the data), lifted over

from hg19 to hg38 with LiftoverVcf (GATK v.4.1.9.0) (*9*), bcftools (v1.10.2) (*10*) was utilized

to include only heterozygous SNPs (hetSNPs) on the X-chromosome. One RNA-seq sample

per participant (see **Table S2**) was downloaded from AnVIL using the Gen3 client, as aligned

BAM files (see https://gtexportal.org/home/methods for specifics on how GTEx processed the

data). Tissues used for the screen were selected based on abundance in the entire data set

(muscle (n=239), skin-lleg (n=21), thyroid (n=14), adipose-subc (n=2), artery-tibi (n=2),

esophagus-muco (n=2), nerve (n=2), adipose-visc (n=1), ovary (n=1), uterus (n=1), (see **Table**

S9 for tissue abbreviations)). Following data download, allelic expression (AE) analysis of

allele counts for the hetSNPs was retrieved from the RNA-seq data using GATK

ASEReadCounter (v.4.1.9.0), requiring a minimum base quality of 20 in the RNA-seq data. To

remove potentially spurious sites, conservative filters were applied to the data (variant call read

depth of at least 20 per allele, minor allele read count > 10% of the total read count and RNA-

seq read depth of more than 10). If a gene carried multiple hetSNPs, only the one with the

highest RNA-seq read count was included. To assess the XCI status of each individual

participant, the mean AE (allelic expression $= |0.5 -$ reference reads/total reads$|$) of genes

previously shown to be inactive (*7*) was calculated (see **Table S2**). A mean AE of inactive genes

> 0.4 was used as a cut-off between mosaic XCI (mXCI) and non-mosaic XCI (nmXCI), with

values > 0.4 indicating all or most cells has inactivated the same parental X-chromosome. As a

control, UPIC was included. UPIC is a female which has previously been shown to have a

nmXCI (*7*). As UPIC is not included in the GTEx_Analysis_2017-06-

05_v8_WholeExomeSeq_979Indiv_VEP_annot.vcf.gz, SNP calling was performed using

WES data and ASE analysis was performed (see next section for detailed pipeline), ultimately

leveraging her ASE data as a positive control. For further assessing XCI status of the 6 samples

with a high mean AE, the remaining RNA-seq tissue samples available for those participants

were downloaded from GTEx and processed in the same manner as above (see **Table S3**).

Following the initial screen, we sought to refine the resolution of the XCI map. To this end, we

downloaded WES and whole-genome sequencing (WGS) data available for the three nmXCI

females (UPIC, 13PLJ and ZZPU) and performed SNP calling. Briefly, WES BAM and WGS

CRAM files for the participants were downloaded from AnVIL using the Gen3 client. The files

were sorted and converted into fastq files using Samtools (v1.9) (*10*). Raw fastq reads were

quality trimmed using FastP (v.0.20.0) (*11*) with default settings. BWA-MEM (*12*) was used for

mapping reads to the human genome build 38 (hg38,

GCA_000001405.15_GRCh38_no_alt_analysis_set.fna) using default settings. Sambamba

v0.8.2 (*13*) was used to mark duplicate reads. The GATK best practice germline short variant

discovery pipeline was used to process the aligned reads, using base quality score recalibration

and local realignment at known indels (GATK v.4.2.6.1). Indels (insertions and deletions) and

SNPs were called jointly across all samples, for both WGS and WES data. Default filters were

applied to indel and SNP calls using the variant quality score recalibration (VQSR) approach

of GATK. All RNA-seq sample available for each participant (**Table S8**) was downloaded from

AnVIL using the Gen3 client as aligned BAM files, using Samtools (v1.9) (*10*) to sort and

convert the BAM files to fastq files and quality trimming with FastP (v.0.20.0) (*11*) with default

settings. RNA-seq reads were aligned with STAR 2-pass mode (v2.7.10a) (*14*) with a

GCA_000001405.15_GRCh38_no_alt_analysis_set.fna   and   --sjdbGTFfile

gencode.v40.annotation.gtf index. WASP filtering was performed to reduce the issue of

reference bias (*15*). Briefly, GATK SelectVariants was used to extract hetSNPs (that passed the

VQSR filtering) for each individual, from either the WGS or WES VCFs, and subsequently

passed to STAR via --varVCFfile and–waspOutputMode. Reads failing WASP filtering were

removed. Following data pre-processing, AE analysis of the allele counts for hetSNPs were

retrieved from RNA-seq data using GATK ASEReadCounter. Heterozygous variants that

passed VQSR filtering were first extracted for each sample from WES and WGS VCFs using

GATK SelectVariants. Sample-specific VCFs and RNA-seq BAMs were inputted to

ASEReadCounter requiring minimum base quality of 20 in the RNA-seq data.

ASEReadCounter outputs from WGS and WES pipelines were merged, if overlapping sites

were identified the variant read counts were merged. To remove potentially spurious sites,

conservative filters were applied to the data (variant call read depth >= 8 per allele, minor allele

read count > 10% of the total read count and RNA-seq read depth > 8). If a gene carried multiple

hetSNPs, only the one with the highest RNA-seq read count was included. For comparisons to

previous gene coverage of UPIC (**Fig. S1A**), supplementary table 5 in Tukiainen *et al* (*7*) was

utilized as a prior.

<u>X-inactivation status categorization</u>

All 401 genes were first classified into PAR or nonPAR depending on their chromosomal

location, and further categorized into 'biallelic', 'monoallelic' and 'variable' depending on their

AE in each individual and tissue. Genes with an AE < 0.4 in at least 85% of tissues and

individuals was classified as 'biallelic', Genes with an AE > 0.4 in at least 85% of tissues and

individuals was 'monoallelic' and genes that varied between in their 'monoallelic' or 'biallelic'

classification across tissues was classified as 'variable' (i.e., AE > 0.4 in more than 15% but

less than 85% of tissues. The genes categorized as 'variable' were subject to manual curation,

as it was difficult to accurately define a computational approach to robustly call XCI on

hetSNPs with low read coverage. Briefly, genes (i) with a low read count per allele and (ii) for

which changes in mono-allelic/bi-allelic classification across tissue were derived from single-

digit read counts were classified as 'undeterminable'. Genes for which variation in XCI status

across tissues were supported by higher read counts were considered more high-confidence

variable genes and thus classified as 'curated variable'.

GTEx RNA-seq data

GTEx RNA-seq data for was downloaded from AnVIL using the Gen3 client (as aligned BAM

files), sorted and converted into fastq files with Samtools (v1.9) (*10*) and quality trimmed with

FastP (v.0.20.0) using default settings. Transcriptome mapping was performed using Salmon

v1.9.0 (*16*) (--gcBias --seqBias --numBootstraps 100 -l A) using known Refseq transcripts

(NM_* and NR_*, excluding all Y-PAR transcripts) from the GRCh38.p12 assembly

(GCF_000001405.38_GRCh38.p12_rna.fna.gz) as a reference. Abundance estimates were

converted to h5 format using Wasabi (https://github.com/COMBINE-lab/wasabi) and data

normalization and gene-level transcript per million (TPM) was calculated using the gene_mode

function implemented in Sleuth (v0.30.0) (*17-18*).

Statistical analysis

For statistical testing between groups a Benjamini-Hochberg adjusted unpaired two-sided

Wilcoxon test was used. P-values < 0.05 was deemed significant.

<u>Abbreviations</u>

XCI: X-chromosome inactivation

Xm: Maternal X-chromosome

Xp: Paternal X-chromosome

GTEx: Genotype-Tissue Expression

nmXCI: non-mosaic X-chromosome inactivation

PAR: pseudo-autosomal region

SNP: single-nucleotide polymorphisms

WES: whole-exome sequencing

WGS: whole-genome sequencing

hetSNP: heterozygous single-nucleotide polymorphisms

AE: allelic expression

Declarations

<u>Ethics approval and consent to participate</u>

Not applicable.

<u>Consent for publication</u>

Not applicable.

<u>Availability of data and materials</u>

No raw data was generated as part of this study. All GTEx data are available from 'The database

of Genotypes and Phenotypes (dbGaP)' under accession number phs000424. All code for

figure   generation   have   been   made   freely   available   on   GitHub

(https://github.com/bgylemo/EpiLiu0987).

<u>Competing interests</u>

Authors declare that they have no competing interests.

Funding

European Research Council grant 101045171 XX-Health (CEN)

Swedish Research Council gran 2020-01277_VR (CEN)

Authors' contributions

Conceptualization: CEN

Methodology: CEN, BG

Investigation: BG

Visualization: BG

Funding acquisition: CEN

Project administration: CEN

Supervision: CEN

Writing – original draft: CEN, BG

Acknowledgements

-

Authors' information (optional)

-

References

1. Carrel, L. & Willard, H.F. X-inactivation profile reveals extensive variability in X-
linked gene expression in females. *Nature* **434**, 400-4 (2005).

2. Cotton, A.M. *et al.* Analysis of expressed SNPs identifies variable extents of expression
from the human inactive X chromosome. *Genome Biol* **14**, R122 (2013).

3. Consortium, G.T. The Genotype-Tissue Expression (GTEx) project. *Nat Genet* **45**, 580-
5 (2013).

4.   Amos-Landgraf, J.M. *et al.* X chromosome-inactivation patterns of 1,005

phenotypically unaffected females. *Am J Hum Genet* **79**, 493-9 (2006).

5.   Roberts, A.L. *et al.* Age acquired skewed X chromosome inactivation is associated with

adverse health outcomes in humans. *Elife* **11**(2022).

6.   Shvetsova, E. *et al.* Skewed X-inactivation is common in the general female population.

Eur J Hum Genet **27**, 455-465 (2019).

7.   Tukiainen, T. *et al.* Landscape of X chromosome inactivation across human tissues.

Nature **550**, 244-248 (2017).

8.   Plenge, R.M. *et al.* A promoter mutation in the XIST gene in two unrelated families

with skewed X-chromosome inactivation. *Nat Genet* **17**, 353-6 (1997).

9.   Castel, S.E., Levy-Moonshine, A., Mohammadi, P., Banks, E. & Lappalainen, T. Tools

and best practices for data processing in allelic expression analysis. *Genome Biol* **16**,

195 (2015).

10.   Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**(2021).

11.   Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ

preprocessor. *Bioinformatics* **34**, i884-i890 (2018).

12.   Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows-Wheeler

transform. *Bioinformatics* **26**, 589-95 (2010).

13.   Tarasov, A., Vilella, A.J., Cuppen, E., Nijman, I.J. & Prins, P. Sambamba: fast

processing of NGS alignment formats. *Bioinformatics* **31**, 2032-4 (2015).

14.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21

(2013).

15.   van de Geijn, B., McVicker, G., Gilad, Y. & Pritchard, J.K. WASP: allele-specific

software for robust molecular quantitative trait locus discovery. *Nat Methods* **12**, 1061-

3 (2015).

16.   Patro, R., Duggal, G., Love, M.I., Irizarry, R.A. & Kingsford, C. Salmon provides fast

297   and bias-aware quantification of transcript expression. *Nat Methods* **14**, 417-419 (2017).

298   17.   Pimentel, H., Bray, N.L., Puente, S., Melsted, P. & Pachter, L. Differential analysis of

299   RNA-seq incorporating quantification uncertainty. *Nat Methods* **14**, 687-690 (2017).

300   18.   Yi, L., Pimentel, H., Bray, N.L. & Pachter, L. Gene-level differential analysis at

301   transcript-level resolution. *Genome Biol* **19**, 53 (2018).