1   **Title:** Transcriptomics Reveals Tumorigenesis and Pathogen Response in Scombrid Puffy Snout
2   Syndrome

3   **Authors:** Savanah L. Leidholt[1], Kalia Bistolas,[1] Mark Dasenko[2], Emily Miller[3], Kyle Van Houtan[3,4], Andre
4   Boustany[3],[4], Tatiana Galivn[1], Rebecca Vega Thurber[1]
5
6   **Affiliations**:
7   1 Oregon State University, Corvallis, OR, USA
8   2 OSU Center for Quantitative Life Sciences, Oregon State University, Corvallis, OR, USA
9   3 Monterey Bay Aquarium, 886 Cannery Row, Monterey, California, 93940, USA.
10 4 Nicholas School of the Environment, Duke University, Durham, North Carolina, 27708, USA.
11
12 **Corresponding Author Information**: Email: sleidholt1994@gmail.com; Mailing Address: 2820 SW
13 Campus Way, Corvallis, OR 97331; Phone: (541) 737-4441
14

Abstract:
Scombrids represent some of the most economically important fisheries globally. However, increased
interest in creating aquaculture systems for these fish has increased the risk of disease emergence. One such
disease, Puffy Snout Syndrome (PSS), causes collagenous tissue growths on the face, in numerous scombrid
taxa. PSS has only been documented in captive-held fish populations and can lead to high mortality rates.
Despite this, little is known about PSS's causative agent(s) and the immune response they elicit. Therefore,
we leveraged metatranscriptomic data of symptomatic-captive, asymptomatic-captive, and healthy-wild
Pacific Mackerel (*Scomber japonicus*) to evaluate the physiological characteristics of PSS infections and
identify a mechanism of disease. PSS symptomatic-captive and asymptomatic-captive mackerel showed
distinct gene expression patterns from their wild counterparts. Genes involved in tumorigenesis, immune
response, and pathophysiology were over-expressed in captive-held fish. In particular, the *Wnt9* pathway
was increased by more than 2 orders of magnitude in captive fish, potentially representing a diagnostic
feature of this disease. Together, these host physiological data and our past visual identification of RNA
virus-like particles in afflicted tissues suggest that viral-mediated oncogenesis may be driving PSS in
captive mackerel.
30

Introduction:
The Scombridae fishes (tunas, bonitos, mackerel) comprise one of the largest global finfish economies,
projected to generate ~$50-80 billion USD annually by 2027 [1]. Efforts have been undertaken in recent
years to increase the efficiency of tuna ranching and farming, which involves keeping tunas captive for
extended periods of time, potentially years, at high densities. Tunas and mackerels are robust species in
aquaculture with negligible economic loss from diseases [2]–[4]. However, high fish-stock densities, low-
quality food sources, and increasing temperatures in mariculture operations have contributed to large-scale
disease outbreaks [4], [5]. Mass mortality outbreaks in tuna aquaculture are often associated with parasitic
blood flukes (*Cordicola* spp.) ranging from 50-80% of an enclosure [6]. Incidences of mass mortality events
from bacterial and viral diseases are less common than parasitic pathogens, but significant die-offs occur
in scombrid aquaculture, mostly associated with juveniles. An outbreak of *Photobacterium damselae*
contributed to over 2500 Atlantic bluefin tuna (*Thunnus thynnus*) fish deaths in a mariculture operation
over a six-month period in 2004 [7]. This outbreak was directly correlated with increased water
temperatures and compounded by the long-term feeding of low-quality baitfish [8]. Other bacterial

outbreaks reported in hatchery-reared tuna larvae include *Mycobacterium, Vibrio*, and *Tenacibaculum* sp.
[10],[11]. Both nervous necrosis virus (NNV) and red seabream iridovirus (RSIV) have had high mortality
rates in larval and juvenile Pacific Bluefin tunas (*Thunnus orientalis*)but have yet to be described in adults
[12]–[14]. However, a study did describe that juvenile *T. orientalis* were highly susceptible to RSIV when
experimentally challenged orally, with nearly 90% mortality in the infected group [15].

Puffy Snout Syndrome (PSS) is a disease of captive scombrids characterized by excessive collagen
growths on the fish head. The first signs are a darkening and wrinkling of the skin, though this phenotype
is more pronounced in tunas. As the disease progresses, a gelatinous substance overgrows on the dorsal
area of the head. Work using Masson trichrome staining found that normal facial tissue was composed
primarily of striated muscle fiber; however, in PSS-afflicted fish, this tissue broke down and was replaced
by collagen [16]. Further, mucus-producing goblet cells which create a healthy slime coat on the fish, were
noticeably decreased in PSS-affected tunas [16]. Severe degradation at the cellular level renders
mitochondria one of the only recognizable structures [17]. A survey of 11 rearing facilities that held tunas,
mackerels, or both found that of the total 22 rearing units in facilities, PSS developed in 68% of these units
[16]. Fish with severe PSS cannot feed and, as a result, are either euthanized or starve to death. Further,
PSS may be exclusively associated with fish held in captivity. None of the facilities surveyed reported
seeing PSS in wild-caught fish, with PSS developing in the first 6 months of captivity [16].

Despite its unique disease characteristics, density-dependent prevalence, and wide host range, PSS
is an understudied syndrome with mortality rates ranging from 20-100% of an enclosure [16], [18]–[20].
Only 9 published reports exist on the topic, and only two of those attempts to identify etiological agents
[16], [17]. Recent work used electron microscopy techniques and identified viral-like particles (VLPs)
within PSS-afflicted tissues of *S. japonicus* [17]. These VLPs shared morphology with small RNA viruses
being amorphous in shape and ~100 nm in diameter. Along with VLPs, malformed mitochondria had a high
abundance in PSS-afflicted tissues. The VLPs and disease prevalence in increased fish stock densities [16]
strongly suggest that PSS onset is associated with a viral infectious agent.

As a next step to identify the underlying mechanisms of PSS, we leveraged high throughput
sequencing methods to further characterize this disease at the physiological and genomic levels. Our aims
were to 1) analyze comparative host gene expression dynamics in PSS of apparently healthy mackerel,
asymptomatic-captive, and symptomatic-captive mackerel; 2) identify expression pathways indicative of
specific diseases (i.e. viral, bacterial, cancerous). Here we present the results of differential gene expression
analysis for *Scomber japonicus* afflicted with PSS and, for the first time, evidence of tumorigenesis in this
species.

Methods and Materials:
Sample Collection:
Sample collection and euthanasia of captive-held and wild-caught mackerel were performed as described
previously [17]. Briefly, captive *S. japonicus* were obtained from the Monterey Bay Aquarium (MBA) and
euthanized with an overdose of MS-222 (i.e. tricaine mesylate). Wild-caught fish were captured from the
coast of Monterey Bay, California, USA, using a hook and line and placed in a holding tank for 2-12 days
before euthanasia via decapitation. All fish were placed in separate bags and placed on ice or at -20°C until
facial necropsy. Epithelial and muscle tissue were taken from behind the nares and anterior to the eyes. A
total of 3 symptomatic-captive, 3 asymptomatic-captive, and 3 healthy wild-caught mackerel were used for
this analysis.

RNA Extraction, Sequencing, and Trimming:
At Oregon State University (OSU), total RNA was isolated from frozen *S. japonicus* tissue. All surfaces
were sterilized with 95% ethanol. Frozen mackerel samples were removed from the -80°C freezer and held
in liquid nitrogen before RNA extraction to ensure RNA stability within tissues. Approximately 0.5 mg of
tissue was removed with a sterile scalpel and forceps, homogenized in liquid nitrogen using an autoclaved
mortar and pestle, and extracted using the E.Z.N.A DNA/RNA Isolation kit from Omega Bio-tek per
manufacturer instructions [21]. RNA was eluted in 50 μl of nuclease-free water and stored at -80 °C until
library preparation.
97
RNAseq library preparation and sequencing were performed at OSU's Center for Quantitative Life
Sciences. Briefly, ribosomal RNA was removed from the total RNA extractions using an Illumina
mouse/rat/human Ribo-Zero kit for host rRNA removal and then a second Ribo-Zero kit for bacterial rRNA
removal. After rRNA removal, a stranded RNA-seq library was prepared using the Takara PrepX DNA
library prep. Then samples were sequenced on one lane of a 2x150bp PE Illumina HiSeq 3000 run. Illumina
HiSeq libraries ranged from 28,745,252 to 49,799,840 paired-end reads. The sequence data was quality-
filtered by removing reads with a Phred score lower than 30. Adapters removal, poly(A) tail removal, and
read normalized to 30x coverage were all done using the BBtools suite [22]. Library read counts after each
quality control step can be found in Supplementary Table 1. The resulting 593,829,134 reads were merged
with BBmerge [23].
108
Informatic pipeline:
Trinity v3.9.1 [24] was employed to build the transcriptome using default settings. Quality-controlled
transcripts of individual libraries (i.e. samples) were mapped to the de novo assembled transcriptome using
the Salmon Estimation Method packaged with Trinity (v.1.8.0 [25]). Compared to two additional reference-
guided assemblies (*Scomber colias* genome and *Scomber scombrus* transcriptome), the de novo
transcriptome recruited the most reads among libraries and was used in this study. All three assemblies'
metrics and recruitment statistics can be found in Supplemental Material.
116
117 DESeq2 (v.3.16) [26] was employed with R statistical programming software (v.4.1.2) [27] to test for
118 differential expression between treatment group pairs to determine whether each model coefficient differed
119 significantly from zero. A two-way ANOVA was used for significance testing. The resulting *p*-values were
120 adjusted for multiple comparisons using Tukey's honest significant difference test in PRISM by GraphPad
121 (v.9.4.1) [28].
122
Results:
A total of 697 differentially expressed genes (DEGs) were identified with distinct differences between
captive and wild-caught fish from the DeSeq2 package (Figure 1c). Genes with adjusted P-value less than
0.01 and an absolute value of log2 more than 1 were defined as differentially expressed. A two-way
ANOVA to analyze the effect of captivity and disease status on the transcriptome of *Scomber japonicus*
revealed a statistically significant difference between wild and captive status groups (p < 0.0001). This
result was consistent for all pairwise comparisons between all DGEs in captive and wild groups (Tukey's
multiple comparisons test p < 0.0001, n=697). No significant difference was seen between the
asymptomatic and symptomatic-captive groups. Tumor suppression and oncogenesis genes were highly
expressed in captive compared to wild-caught animals (Figure 2a). Immune-related genes and

pathophysiological markers were also significantly over-expressed (Figure 2b-c) (p-value < 0.01 n = 100).
Wild-caught *S. japonicus* were assumed to be healthy in all cases as there were no visible signs of disease
externally or internally.
136
Elevated Tumorigenesis Markers Are Characteristic of PSS
Wnt9 was identified as the most highly expressed DEG within captive-held *S. japonicus* tissue. *Wnt9*
showed a two-fold difference in expression between captive and wild-caught animals (Figure 1d). This gene
is the starting component of the canonical Wnt-signaling pathway responsible for cell proliferation and
growth. Other genes identified as differentially expressed in this pathway were *LDLR*, *CCTNB1*, and
CNNA2 (Figure 2A).
143
Immune-Related Gene Expression Altered in Captive Fish
All immune genes identified as differentially expressed are innate immune system components with some
of the most important in fish being *TRIM16* and *TLR8* (Figure 2b). Several of these genes trigger the
inflammatory response such as: *HIF-PHI*, *il-11* and *HSP90A1*. All of the aforementioned genes had a 1-1.5
fold increase in expression in the captive-held mackerel. Only *USP25* appears significantly underexpressed
in this system by a 1-1.5 fold. *USP25* has been implicated as an innate immune response to pathogens and
some cancers.
151
Pathophysiology are overexpressed in Captive Fish
Genes related to tissue regeneration were overexpressed within captive-held mackerel (Figure 2C). The
most abundant of these genes induces collagen production, which appears to be a hallmark of PSS. Genes
associated with ossification (*NPPC*), glucose transport and tissue regeneration (*ADIPOQ*), fibrosis
(*ENTPD2*), and vascular repair (*B2R*) were also overexpressed in captive-held mackerel. No genes
associated with muscle regeneration were found to be differentially expressed.

Figure 1: Transcriptomic data shows distinct differences between captive and wild mackerel A) Adapted from Miller et al., 2021[17]; From left to right images of healthy wild fish, asymptomatic-captive fish, and PSS symptomatic-captive fish. B) Principal component analysis plot of all transcripts showing the top 2 axes which explain the most variance. C) Top 100 differentially expressed genes in a heatmap showing clear differentiation by fish origin. D) Raw counts of the gene, *Wnt-9B* in each group. Black boxes and symbols represent healthy-wild mackerel, green represents symptomatic-captive mackerel, and yellow represents asymptomatic-captive mackerel.

Discussion

This work characterizes Puffy Snout Syndrome (PSS) at the transcriptomic level, showing facial tumorigenesis evidence through the pan-cancer marker overexpression within damaged tissues. However, PSS remains an enigmatic syndrome with multiple focal areas for future study. Transcriptomics offers insight into variations in animal physiology by highlighting differences in gene expression amongst healthy, asymptomatic, and symptomatic fish. Here, we found that gene expression in tumorigenesis, immune response, and pathophysiology were dramatically altered in captive fish compared to their wild counterparts. Many genes can be associated with PSS's gross pathology (Figure 2d).

PSS tissues are highly enriched in genes involved in tumorigenesis.
Transcriptomic data suggest that the primary cause of PSS is dysregulation of cell proliferation &
differentiation. This could lead to the observed collagenous growths and progressive evolution through
hyperplasia and dysplasia in the facial tissue, often characteristic of cancers. This is reflected in the largest
detected shift in gene expression in *Wnt9* between wild and captive groups. *Wnt9* is a morphogen in adult
animals' early development, cell differentiation/migration, and tissue homeostasis [29]. This gene was
elevated ~2-fold in captive fish and may be diagnostic of PSS. The gene for beta-catenin, *CTNNB1*, the
other major gene in the *Wnt9* pathway, was also overexpressed in captive fish. Genes in the canonical
WNT/beta-catenin pathway (Figure 3) are often diagnostic for many human and non-human animal cancers
[30]–[32].
Despite its role in tumor suppression, retinoblastoma protein (*Rb1)* was found to be significantly
overexpressed in captive fish. *Rb1* regulates mitosis and gene expression by binding chromatin and limiting
gene expression of proteins involved in moving cells from the G1 to the S stage [33]. However, *Rb1*
phosphorylation by the cyclin-mediated pathway alters its ability to bind these genes [34]. *CCNA2*, which
encodes cyclin-A2, which regulates this transition, is also highly expressed ~1.5 fold, suggesting that tumor
suppression activity of Rb1 may be inhibited upstream by cyclin over-expression.
203
Pathogens may explain the differences in the immune gene expression.
We hypothesize that the immune-specific transcriptomic profile results from prolonged
physiological stress in captivity or an underlying pathogenic infection such as a virus that captivity may
exacerbate. Essential immune genes expressed in response to PSS are indicative of the viral pathogen
response in fish and mammals. For example, *USP25* is a form of post-translational modification and has
been shown to positively regulate RNA virus-induced signaling by the innate immune system, therefore
playing a critical role in antiviral immunity [35]. Studies with *USP25* deficient mice showed they were
more susceptible to infection to both herpes simplex virus 1 and influenza A than their wild-type
counterparts [36]. *USP25* successfully activated the antiviral cascades of IFN3 and NF-κβ by stabilizing
Traf3 and *Traf6* from viral degradation [31].
TRIM16 is part of the fish-specific TRIM proteins and has been associated with viral infections in
fish through negatively regulating interferon production [37]–[40]; higher expression of TRIM16
expression is associated with higher viral loads and poor survival outcomes of fish. Further, in fish, TLR8
primarily identifies single-stranded RNA molecules and is presumed to be a sensor for single-stranded RNA
viral infection [41]. Further investigation is required to determine if USP*25, TRIM16, and TLR8* expression
patterns are due to a stress response or virus.
220
Pathophysiology corresponds to the histological examination of tissues.
Often pathophysiological markers show an organism's response to tissue damage but do not
necessarily provide information on the cause of that damage. Voorhees, 2015 reported an initial
inflammatory response in PSS-afflicted tissue with extensive fibrogenesis followed by collagen's slow
replacement of muscle tissue. The overexpressed pathophysiological genes in captive-held animals support
this finding. A total of four different collagen types were reported to be over-expressed in captive-held
animals. Likewise, no genes associated with muscle regeneration were found to be overexpressed in captive
mackerel. Thus, the observed collagen gene expression profiles could be affiliated with the morphology of
PSS increased tissue growth on the head and overall tissue inflammation.

We propose that PSS is a condition induced by an infectious oncogenic virus that induces tumor
formation, physiological cancer induced by alternate factors and not previously described in wild fish, or
transmissible cancer. Given that transmissible cancers are extremely rare in wildlife, with few exceptions,
such as in marine molluscs, terrestrial canines, and rodents, we can mostly but not completely rule this out.
Further, no transmissible cancer has been described in teleosts, so it remains an unlikely explanation.
However, our previous microscopy work showed the presence of small viral particles and large numbers of
deformed mitochondria in affiliated fish. Given these visual observations, the increased expression of
tumorigenesis genes, and genes involved in the innate immune response, viral infection-inducing
tumorigenesis seems most likely. Further characterization of the presence and abundance of viral-specific
genes and genomes and additional viral-induced cytopathology will be required to confirm these
hypotheses.
241
242
243

244
245   Figure 2: Selected DGEs: A) tumor-related genes; B) immune genes; C) and pathophysiological genes.
246   D) Venn diagram showing the overlap of the top 100 genes within these three groups. Black boxes represent
247   healthy-wild mackerel, yellow represent asymptomatic mackerel, green represents symptomatic mackerel.
248

249   Data Accessibility:
250   Raw reads and transcriptome were deposited to the figshare data repository under DOI:
251   10.6084/m9.figshare.22779725. Raw reads were also uploaded to NCBI under BioProject number
252   PRJNA967399.
253

Authors' contributions:
All authors contributed to the writing in this manuscript. E.M performed all necropsies. S.L., M.D., and
T.G. performed DNA/RNA extractions and library preparation. S.L and K.V created all figures. R.V.T.
provided the grants for this work.
symptomatic-captive mackerel, and yellow represents asymptomatic mackerel.

Acknowledgements: This research was funded through the Pernot Endowment fund and Tartar Award
through Oregon State University. We would like to thank Sherrie Bain and Manny Ezcurra for their
expertise in animal care in handling during the deaccession process.

Citations:

[1]   Research and Markets ltd, "Global Tuna Market Report and Forecast 2022-2027."
264   https://www.researchandmarkets.com/reports/5558096/global-tuna-market-report-and-forecast-2022-
265   2027 (accessed Mar. 21, 2023).
266   [2]   G. Jelić Mrčelić, V. Nerlović, M. Slišković, and I. Zubak Čižmek, "An Overview of Atlantic Bluefin
267   Tuna Farming Sustainability in the Mediterranean with Special Regards to the Republic of Croatia,"
268   *Sustain. Sci. Pract. Policy*, vol. 15, no. 4, p. 2976, Feb. 2023.
269   [3]   B. Nowak, I. Mladineo, H. Aiken, N. Bott, and C. Hayward, "farmed tuna: southern bluefin tuna
270   (Thunnus maccoyii) in Australia and northern bluefin tuna (Thunnus thynnus) in the Mediterranean."
271   https://eafp.org/download/2006-Volume26/Issue%201/26_038.pdf (accessed Mar. 21, 2023).
272   [4]   J. Balli, I. Mladineo, S. Shirakashi, and B. F. Nowak, "Chapter 11 - Diseases in Tuna Aquaculture,"
273   in *Advances in Tuna Aquaculture*, D. D. Benetti, G. J. Partridge, and A. Buentello, Eds. San Diego:
274   Academic Press, 2016, pp. 253–272.
275   [5]   I. Mladineo, J. Zilic, and M. Cankovic, "Health survey of Atlantic bluefin tuna, thunnusthynnus
276   (Linnaeus, 1758), reared in Adriatic cages from 2003 to 2006."
277   https://d1wqtxts1xzle7.cloudfront.net/67831536/j.1749-7345.2008.00156.x20210701-2244-n4jmzo-
278   libre.pdf?1625140226=&response-content-
279   disposition=inline%3B+filename%3DHealth_survey_of_Atlantic_Bluefin_Tuna_T.pdf&Expires=16
280   79434350&Signature=cxGfkXeFjT~UBwQvMg0CD3IBZFL5WOb7Yw6lQlYBnqh4Mf1N4XmeW
281   9ZomQuUK1C2c7quK-
282   GMTJv6qItrd5ptaDSVPsmtgjNLmDEkX9AB7OaXcURKUXyFqhMc9Dwl3K6Jjmp-
283   45LBojuLie9AZoGofT9nLoRwN2plRpau9-wfuV8jBoxCb60TH~UuaZ0pyubMPJVmw-
284   eDgjQdfbOmpK4SdbIU6OTHJc0i-r83mR7VBCiYC1gdCb~cw0oR2ePYymPRY5iyyLtz5FS0ZPa-
285   S8J9yMeAfwT23zs3tmxccHzmNrC6XjQkYiWaw6~LQVdOvMbw3wCtLRMXJYD6c4Pvs4PD-
286   w__&Key-Pair-Id=APKAJLOHF5GGSLRBV4ZA (accessed Mar. 21, 2023).
287   [6]   J. J. Minich *et al.*, "The Southern Bluefin Tuna Mucosal Microbiome Is Influenced by Husbandry
288   Method, Net Pen Location, and Anti-parasite Treatment," *Front. Microbiol.*, vol. 11, p. 2015, Aug.
289   2020.
290   [7]   I. Mladineo, I. Miletić, and I. Bočina, "Photobacterium damselae subsp. piscicida Outbreak in Cage-
291   Reared Atlantic Bluefin Tuna Thunnus thynnus," *J. Aquat. Anim. Health*, vol. 18, no. 1, pp. 51–54,
292   Mar. 2006.
293   [8]   V. Å imat, J. MarÅ¡iÄ-LuÄiÄ, M. Tudor, and I. Mladineo, "Long-term storage influence on volatile
294   amines (TVB-N and TMA-N) in sardines and herring utilized as food for tuna fattening," *J. Appl.
295   Ichthyol.*, vol. 25, no. 6, pp. 766–770, Dec. 2009.
296   [9]   B. L. Munday, Y. Sawada, T. Cribb, and C. J. Hayward, "Diseases of tunas, Thunnus spp," *J. Fish
297   Dis.*, vol. 26, no. 4, pp. 187–206, Apr. 2003.
298   [10] N. Sandlund, C. Skår, and E. Karlsbakk, "First identification of mycobacteriosis in Atlantic mackerel
299   (Scomber scombrus)," *J. Fish Dis.*, Feb. 2023, doi: 10.1111/jfd.13765.
300   [11] Gustinelli, A., Campana, M., Caggiano, M., Florio, D., Caffara, M., Quaglio, F., Corriero, A. and
301   Fioravanti, M.L., "Observations on health problems in hatchery reared tuna (Thunnus thynnus) in
302   Ital," presented at the 15th EAFP International Conference on Diseases of Fish and Shellfish,
303   Croatia, 2011, p. 482.
304   [12] K. N. Hidemasa Kawakami, "Marine cultured fish species with red sea bream iridovirus disease
305   confirmed from 1996 to 2000," *Fish Disease Research*, vol. 37, no. 1, pp. 45–47, 2002.
306   [13] Sawada, Y., S. Miyoshita, O. Murata, and H. Kumai, "Problems in the seedling production of
307   northern BFT, Thunnus orientalis," in *Proceeding of European Aquaculture Society "'Seafarming
308   today & tomorrow,'"* Trieste, Itally, 2002, pp. 464–465 i.
309   [14] T. Sugaya *et al.*, "Genetic heterogeneity of betanodaviruses in juvenile production trials of Pacific
310   bluefin tuna, Thunnus orientalis (Temminck & Schlegel)," *J. Fish Dis.*, vol. 32, no. 10, pp. 815–823,
311   Oct. 2009.

[15] Y. Matsuura *et al.*, "Development of a method for experimental infection of Pacific bluefin tuna with red seabream iridoviral disease," *Aquaculture*, vol. 539, p. 736627, Jun. 2021.

[16] T. M. Voorhees, "Epidemiology of Puffy Snout Syndrome in Tuna," University of Rhode Island, 2015. doi: 10.23860/thesis-voorhees-taylor-2015.

[17] E. A. Miller *et al.*, "Electron microscopy reveals viral-like particles and mitochondrial degradation in scombrid puffy snout syndrome," *Dis. Aquat. Organ.*, vol. 147, pp. 25–31, Nov. 2021.

[18] A. Tester, "Establishing Tunas and Other Pelagic Species in Ponds and Tanks," United States Department of the Interior; Fish and Wildlife Services, 71, 1952. [Online]. Available: https://spo.nmfs.noaa.gov/sites/default/files/legacy-pdfs/SSRF71.pdf

[19] E. L. Nakamura, "Observations on the Behavior of Skipjack Tuna, Euthynnus pelamis, in Captivity," *Copeia*, vol. 1962, no. 3, pp. 499–505, 1962.

[20] E. L. Nakamura, "Development and Uses of Facilities for Studying Tuna Behavior," in *Behavior of Marine Animals: Current Perspectives in Research Volume 2: Vertebrates*, H. E. Winn and B. L. Olla, Eds. Boston, MA: Springer US, 1972, pp. 245–277.

[21] "E.Z.N.A.®DNA/RNA Kit," OMEGA Biotek, 2018. [Online]. Available: www.omegabiotek.com

[22] "BBMap," *SourceForge*, Jul. 15, 2022. https://sourceforge.net/projects/bbmap/ (accessed Mar. 23, 2023).

[23] B. Bushnell, J. Rood, and E. Singer, "BBMerge - Accurate paired shotgun read merging via overlap," *PLoS One*, vol. 12, no. 10, p. e0185056, Oct. 2017.

[24] M. G. Grabherr *et al.*, "Full-length transcriptome assembly from RNA-Seq data without a reference genome," *Nat. Biotechnol.*, vol. 29, no. 7, pp. 644–652, May 2011.

[25] R. Patro, G. Duggal, M. I. Love, R. A. Irizarry, and C. Kingsford, "Salmon provides fast and bias-aware quantification of transcript expression," *Nat. Methods*, vol. 14, no. 4, pp. 417–419, Apr. 2017.

[26] M. I. Love, W. Huber, and S. Anders, "Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2," *Genome Biol.*, vol. 15, no. 12, p. 550, 2014.

[27] R Core Team, "R: A language and environment for statistical ## computing. R Foundation for Statistical Computing, Vienna, Austria," 2021. Accessed: Mar. 23, 2021. [Online]. Available: https://www.R-project.org/

[28] "Prism - GraphPad." https://www.graphpad.com/features (accessed Apr. 03, 2023).

[29] J. C. Croce and D. R. McClay, "Evolution of the Wnt Pathways," in *Wnt Signaling*, E. Vincan, Ed. Totowa, NJ: Humana Press, 2009, pp. 3–18.

[30] M. Ramos-Solano, M. Álvarez-Zavala, B. García-Castro, L. F. Jave-Suárez, and A. Aguilar-Lemarroy, "Wnt signalling pathway and cervical cancer," *Rev. Med. Inst. Mex. Seguro Soc.*, vol. 53, no. S2, pp. 218–224, May 2015.

[31] M. Crumbaker *et al.*, "The Impact of Whole Genome Data on Therapeutic Decision-Making in Metastatic Prostate Cancer: A Retrospective Analysis," *Cancers*, vol. 12, no. 5, May 2020, doi: 10.3390/cancers12051178.

[32] H. M. Penrose *et al.*, "High-fat diet induced leptin and Wnt expression: RNA-sequencing and pathway analysis of mouse colonic tissue and tumors," *Carcinogenesis*, vol. 38, no. 3, pp. 302–311, Mar. 2017.

[33] N. J. Dyson, "RB1: a prototype tumor suppressor and an enigma," *Genes Dev.*, vol. 30, no. 13, pp. 1492–1502, Jul. 2016.

[34] K. Engeland, "Cell cycle regulation: p53-p21-RB signaling," *Cell Death Differ.*, vol. 29, no. 5, pp. 946–960, May 2022.

[35] D. Lin *et al.*, "Induction of USP25 by viral infection promotes innate antiviral responses by mediating the stabilization of TRAF3 and TRAF6," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 112, no. 36, pp. 11324–11329, Sep. 2015.

[36] W. Zhu, D. Zheng, D. Wang, L. Yang, C. Zhao, and X. Huang, "Emerging Roles of Ubiquitin-Specific Protease 25 in Diseases," *Front Cell Dev Biol*, vol. 9, p. 698751, Jun. 2021.

[37] L. M. van der Aa *et al.*, "A large new subset of TRIM genes highly diversified by duplication and positive selection in teleost fish," *BMC Biol.*, vol. 7, p. 7, Feb. 2009.

[38] Y. Yu *et al.*, "Fish TRIM16L exerts negative regulation on antiviral immune response against grouper iridoviruses," *Fish Shellfish Immunol.*, vol. 59, pp. 256–267, Dec. 2016.

[39] C. Langevin, J.-P. Levraud, and P. Boudinot, "Fish antiviral tripartite motif (TRIM) proteins," *Fish Shellfish Immunol.*, vol. 86, pp. 724–733, Mar. 2019.

[40] E. Misk, P. Huber, J. I. MacInnes, S. M. Sherif, M. Abo-Ismail, and J. S. Lumsden, "Innate response of rainbow trout gill epithelial (RTgill-W1) cell line to ultraviolet-inactivated VHSV and FliC and rhabdovirus infection," *Fish Shellfish Immunol Rep*, vol. 3, p. 100043, Dec. 2022.

[41] D. Pietretti and G. F. Wiegertjes, "Ligand specificities of Toll-like receptors in fish: indications from infection studies," *Dev. Comp. Immunol.*, vol. 43, no. 2, pp. 205–222, Apr. 2014.

[42] M. Kanehisa and S. Goto, "KEGG: kyoto encyclopedia of genes and genomes," *Nucleic Acids Res.*, vol. 28, no. 1, pp. 27–30, Jan. 2000.