Title:

Dual-specific autophosphorylation of kinase IKK2 enables phosphorylation of

substrate IκBα through a phosphoenzyme intermediate

Prateeka Borar[1], Tapan Biswas[2,#], Ankur Chaudhuri[3,#], Tom Huxford[4], Saikat Chakrabarti[3]

Gourisankar Ghosh[2] & Smarajit Polley[1,*]

[1]Department of Biological Sciences, Bose Institute, Kolkata, India; [2]Department of Chemistry

and Biochemistry, University of California San Diego, La Jolla, USA; [3]Structural Biology and

Bioinformatics Division, CSIR-Indian Institute of Chemical Biology, Kolkata, India;

[4]Department of Chemistry and Biochemistry, San Diego State University, San Diego, USA.

[#] These two authors have contributed equally to this work.

*Correspondence to be addressed to: spolley@jcbose.ac.in, smarajit.polley@gmail.com

Abstract:

Rapid and high-fidelity phosphorylation of two serines (S32 and S36) of IκBα by a prototype

Ser/Thr kinase IKK2 is critical for fruitful canonical NF-κB activation. Here, we report that

IKK2 is a dual specificity Ser/Thr kinase that autophosphorylates itself at tyrosine residues in

addition to its activation loop serines. Mutation of one such tyrosine, Y169, located in

proximity to the active site, to phenylalanine, renders IKK2 inactive for phosphorylation of

S32 of IκBα. Surprisingly, auto-phosphorylated IKK2 relayed phosphate group(s) to IκBα

without ATP when ADP is present. We also observed that mutation of K44, an ATP-binding

lysine conserved in all protein kinases, to methionine renders IKK2 inactive towards specific

phosphorylation of S32 or S36 of IκBα, but not non-specific substrates. These observations

highlight an unusual evolution of IKK2, in which autophosphorylation of tyrosine(s) in the

activation loop and the invariant ATP-binding K44 residue define its signal-responsive

substrate specificity ensuring the fidelity of NF-κB activation.

Introduction:
42
Protein kinases bestow novel identity to their substrates by adding phosphate

moieties at the site of modification that often play key regulatory roles in a diverse array of

signaling events. For example, inflammatory or pathogenic signals trigger precisely

regulated induction of the transcription factor NF-κB in metazoans(Hoffmann and Baltimore,

2006) that depends upon the upstream activation of the Inhibitor of κB Kinase-complex (IKK-

complex), comprising two catalytic subunits, IKK1 (also known as IKKα) and IKK2 (or IKKβ),

and an essential regulatory scaffolding protein, NEMO (IKKγ)(DiDonato et al., 1997;

Rothwarf et al., 1998; Zandi et al., 1997). IKK2 activity is maintained at a very low basal level

in resting cells(Ghosh and Karin, 2002; Hacker and Karin, 2006; Hinz and Scheidereit, 2014;

Liu et al., 2012). Upon activation, IKK2 phosphorylates the N-terminus of IκBα (Inhibitor of

κBα) protein specifically at two Serine residues (S32 and S36) marking it for ubiquitinylation-

mediated proteasomal degradation (Figure 1A). Signal-responsive phosphorylation of IκBα

refers to the phosphorylation of S32 and S36 by the IKK-activity. Degradation of IκBα

liberates NF-κB to execute its gene expression program (Hayden and Ghosh, 2008; Hinz

and Scheidereit, 2014; Karin and Ben-Neriah, 2000; Scheidereit, 2006). Mutation of both

S32 and S36 to alanine residues converts IκBα into a non-phosphorylatable super-repressor

of NF-κB(Brown et al., 1995; Lin et al., 1995). Furthermore, IKK possesses exquisite

specificity for these two serines, and it fails to phosphorylate IκBα when S32 and S36 are

mutated even to another phosphorylatable residue, threonine(DiDonato et al., 1997).

IKK2 is an essential Ser/Thr kinase that ensures the activation of a particular

signaling cascade - NF-κB, with high fidelity despite its reported pleiotropy in other

contexts(Antonia et al., 2021; Schröfelbauer et al., 2012; Schrofelbauer and Hoffmann,

2011). IKK2 is a multidomain protein consisting of a kinase domain (KD) followed by a

Ubiquitin-like domain (ULD) and the scaffold dimerization domain (SDD) (Figure 1B). IKK2

toggles between inactive and active states based on the phosphorylation status of two

serine residues (S177 and S181; S176 and S180 in the case of IKK1) within its activation

loop (activation loop) in the KD(Huse and Kuriyan, 2002; Liu et al., 2013; Polley et al., 2013;

Xu et al., 2011). Replacement of these two serines with phospho-mimetic glutamate

residues (S177E, S181E; henceforth EE) renders the kinase constitutively active. IKK2 is

reported to be primed through phosphorylation at S177 by the upstream kinase TAK1 that

leads to auto-phosphorylation at S181, thus rendering the kinase fully active in cells(Zhang

et al., 2014). However, oligomerization upon association with NEMO, in the presence of

linear or Lys-63 linked poly-ubiquitin (Ub) chains or at a high concentration of IKK2, also

enables *trans* auto-phosphorylation at S177 and S181(Chen, 2012; Du et al., 2022; Ea et al.,

2006; Polley et al., 2013). It was also shown that NEMO not only helps in the activation of

the IKK-complex, but also ensures that the IKK-activity is well directed towards IκBα in the

inhibitor bound transcriptionally inactive NF-κB (NF-κB: IκBα complex)(Schröfelbauer et al.,

2012). Activated IKK2 phosphorylates IκBα specifically at residues S32/S36, for which the

C-terminal SDD of IKK2 (residue range 645-756) is critical – a shorter construct of IKK2

lacking the NBD and SDD displayed CK2-like activity that did not retain the exquisite

specificity of IKK2 to phosphorylate the S32/36 of IκBα (Shaul et al., 2008). Other sites of

IκBα e.g., the residues in the C-terminal PEST region that contains proline (P), glutamic acid

(E), serine (S) and threonine (T) residues, but not S32/S36, can be phosphorylated by

various kinases(Barroga et al., 1995; Tergaonkar et al., 2003). The role of IκBα-

phosphorylation at sites other than the signal responsive S32/S36 phosphorylation in NF-κB

activation and signaling is elusive. These findings underscore the ability of IKK2 to function

specifically in order to guarantee rapid NF-κB activation in canonical signaling. It is however,

still not fully understood how IKK2 achieves its exquisite specificity towards S32/S36 of IκBα

to mark it for proteasomal degradation that liberates active NF-κB dimers.

Here we report an in-depth analysis of the catalytic features of IKK2 that revealed

that IKK2 undergoes multisite autophosphorylation and can phosphorylate its own tyrosine

residue(s) in addition to its own serines or the serines of its substrate IκBα, i.e., IKK2

possesses dual specificity. This phosphorylation at tyrosine residue(s) is autocatalytic. This

phosphorylation at tyrosine residues is dependent upon the prior activation of IKK2 via

activation loop S177/S181 phosphorylation, and is critical to phosphorylation of signal

responsive S32/S36 of IκBα. Mutation of Y169 of IKK2 to phospho-ablative phenylalanine

severely compromised phosphorylation of IκBα at S32. Mutation of Y188 to phenylalanine

rendered the kinase practically inactive *in vitro*. IKK2 activity towards signal responsive

S32/S36 of IκBα was severely compromised in MEF cells reconstituted with these two

mutant IKK2s in response to treatment with TNFα. We also observed that mutation of the

conserved ATP-interacting residue K44 in IKK2 to methionine leads to a loss of tyrosine as

well as S177,S181 auto-phosphorylation activity of the kinase along with its ability to

phosphorylate S32/S36 of IκBα, thereby clearly suggesting a critical role of this Lysine

residue in regulating IKK2's kinase activity for tyrosine as well as serine phosphorylations.

Interestingly, IKK2 K44M retained non-specific kinase activity referring to phosphorylation of

IκBα at residues other than S32 and S36. Furthermore, we observed that, after auto-

phosphorylation at tyrosine, IKK2 can phosphorylate IκBα in the absence of exogenous

supply of ATP, as long as ADP is present in the reaction. Overall, our results reveal unique,

and likely transient, autophosphorylated intermediates of IKK2 that appear to be critical in

relaying phosphoryl group(s) specifically to S32/S36 of IκBα rather than following the

conventional mode of phosphate group transfer directly from ATP to the substrate seen in

eukaryotic protein kinases. IKK2 thus ensures fidelity of NF-κB activation. Recent reports in

other kinases indicate that multisite (auto)phosphorylation is a major determinant in

regulating function of respective kinases and regulation exerted by those kinases. Our study

adds another layer of regulation and critical nature of such autophosphorylation that plays a

significant biological role in metazoans.

Results:
120
121   ***IKK2 undergoes auto-phosphorylation at uncharacterized sites:***

122   Signal-induced phosphorylation of activation loop Ser177/Ser181 marks the general

activation of IKK2 (Figure 1B) from an unphosphorylated inactive state. Phosphorylation of

several other serines located within the flexible C-terminal region of IKK2, spanning amino

acid 701 to the end, has also been reported (Figure 1B)(Delhase et al., 1999).

Hyperphosphorylation of these C-terminal sites down-regulates IKK2 activity in cells. It was

also reported earlier that, ectopically overexpressed IKK2 is capable of autophosphorylating

these two serines in the activation loop in *trans* even in the absence of an activating

physiological signal. Indeed, recombinantly purified FL IKK2WT is capable of

autophosphorylation *in vitro* when incubated with γ^{32}P-ATP that increases with the time of

incubation (Figure 1C). The spread-out of slower migrating phospho-IKK2 band likely

indicates phosphorylation of the protein at multiple sites. We also observed that

autophosphorylation of IKK2 occurred irrespective of the presence of its cognate binding

partner NEMO and *bonafide* substrate IκBα (Figure 1D). To confirm that this

phosphorylation of IKK2 was indeed autophosphorylation and not due to an unrelated kinase

copurified with IKK2, we generated a K44M mutant of full-length IKK2 where the ATP-

binding invariant lysine residue conserved in Ser/Thr/Tyr kinases was mutated to

methionine. This lysine residue anchors ATP in the respective binding pocket of the kinase

by forming hydrogen bonds with the α- and β- phosphate of ATP that is critical for productive

phosphorylation reaction in protein kinases. Mutating this K to M is expected to greatly

reduce or eliminate the kinase activity. An *in vitro* kinase assay using this mutant along with

the WT kinase using γ^{32}P-ATP as the phosphor-donor confirmed that the IKK2 K44M lacked

both IκBα (substrate) and autophosphorylation activities (Figure S1A). That these

phosphorylations were indeed autophosphorylation was further supported by another *in vitro*

radioactive kinase assay performed in the presence or absence of an IKK-specific ATP-

competitive inhibitor, Calbiochem Inhibitor VII (Figure 1E) where the kinase and kinase-

substrate reactions were incubated for 30 minutes with the inhibitor prior to adding the

phosphor-donor, γ^{32}P-ATP. A similar assay was performed with cold ATP and specific

phosphorylation of S32/36 on IκBα was monitored by immunoblotting in presence and

absence of the same inhibitor (Figure S1B). Purification status of recombinant FL IKK2 WT

and FL IKK2 K44M proteins used in this study are shown in Figure S1C.

Next, we used different versions of the IKK2 protein where all previously known

phosphorylation sites were lacking: Ser177 and 181 were mutated to phosphor-mimetic Glu

(EE; this version of IKK2 is known to be constitutively active (Zandi et al., 1998)), and the C-

terminal serine-rich region was truncated to different extents. The shortest construct,

Δ664EE that entirely lacked the flexible C-terminus (670-756) was further subjected to

limited proteolysis using trypsin to eliminate residues, if any, from the flexible ends of the

enzymes and this trypsin-treated IKK2 was purified prior to *in vitro* kinase assays. Each of

these constructs was active in phosphorylating itself and the substrate IκBα (Figure 1F). The

occurrence of autophosphorylation despite the absence of activation loop serines and

serines within the C-terminus strongly suggested the presence of hitherto uncharacterized

sites of autophosphorylation in IKK2.

Auto-phosphorylation of IKK2 reveals its dual specificity:

IKK2 is known as a *bonafide* Ser/Thr kinase that undergoes autophosphorylation. In
the previous section, we found that IKK2's autophosphorylation activity might extend beyond
the previously characterized site(s). We asked if it is a dual specificity kinase. To test if Tyr
residues can undergo auto-phosphorylation in addition to Ser, IKK2 incubated with Mg^{2+}-
ATP for various time periods was probed with antibodies that specifically recognize
activation loop serines of IKK2 (Figure 2A, upper panel) or phospho-tyrosine residues
(Figure 2A, middle panel). Unexpectedly, we observed *de novo* auto-phosphorylation of
IKK2 on tyrosine residues along with the anticipated auto-phosphorylation of activation loop
serines. It may be noted that autophosphorylation activation loop-S177/181 also increased
with time, indicating inhomogeneous and incomplete phosphorylation of activation loop-
S177/181 in the recombinant IKK2 when purified from Sf9 cells. We tested another phospho-
tyrosine specific antibody to further substantiate this *de novo* tyrosine autophosphorylation of
IKK2. It confirmed that detection of *de novo* phosphotyrosine on IKK2 was independent of
the source of the antibody used (Figure 2B).   The tyrosine auto-phosphorylation was
abolished in the presence of an IKK-specific inhibitor, Inhibitor VII (Figure 2C). A similar
assay was performed to monitor the phospho-tyrosine level in IKK2 as a function of urea
concentration. This revealed the detrimental effects of urea on tyrosine autophosphorylation
of IKK2 at concentrations beyond 1M (Figure S2A). A more sensitive assay with γ^{32}P-ATP as
the phospho-donor also indicated that urea concentrations beyond 1M inhibited both auto-
phosphorylation (pan-phosphorylation) and substrate (IκBα) phosphorylation (Figure S2B).
Furthermore, the IKK2-K44M mutant (kinase inactivating) did not display any tyrosine
autophosphorylation when incubated with ATP unlike WT IKK2 (Figure 2D). These results
confirm the novel tyrosine phosphorylation ability of IKK2, demonstrating its dual specificity
to both its own Ser and Tyr residues. We also observed that activity of IKK2 towards S32
and S36 of IκBα and its own tyrosine residue(s) were affected similarly when the kinase
assays were performed in the presence of various kinase inhibitors, both general (AMPPNP
and Staurosporine) and IKK-specific (MLN-120B, TPCA, Inhibitor VII) (Figure S2C). These

observations confirm that canonical activity of IKK2 towards S32 and S36 of IκBα, and

towards its own tyrosine residues are mediated by the same active site.

Next, we checked the requirement of the C-terminal <u>NE</u>MO <u>b</u>inding <u>d</u>omain (NBD) for

autocatalytic dual specificity. Constructs of various truncations (lacking NBD) were capable

of phosphorylating itself on tyrosine residue(s), and phosphorylating IκBα at signal-

responsive S32 and S36 (Figure 2E), indicating that the NBD domain is not required for

either its ability to phosphorylate S32 and S36, as well as autophosphorylating itself on

tyrosine. Next, we assessed the autocatalysis at Ser and Tyr by IKK2 using a constitutively

active NBD-truncated version of IKK2 (IKK2$^{\Delta664}$EE). As anticipated, the protein was

recognized by monoclonal phospho-serine antibody with or without fresh ATP-treatment;

however, phospho-tyrosine antibody recognized it strongly only upon fresh ATP-treatment

(Figure 2F) further confirming dispensability of the NBD in *de novo* tyrosine

autophosphorylation of IKK2.

Next, we asked if phosphorylation of activation loop S177/S181 was necessary for

this dual specificity of IKK2. To this end, we purified the S177A, S181A (AA; non-

phosphorylatable activation loop) phospho-ablative mutant of IKK2, and checked its tyrosine

autophosphorylation activity. We found that the IKK2AA mutant lacked tyrosine

autophosphorylation activity (Figure 2G) in an *in vitro* kinase assay. It suggests that

activation loop S177/S181 serine phosphorylation is essential in conducting dual specificity

in IKK2. Indeed, the IKK2 AA mutant was incapable of phosphorylating S32/36 of IκBα as

well (Figure S2D) in an *in vitro* kinase assay.

IKK2 autophosphorylation is critical to specificity of IκBα phosphorylation:

As mentioned earlier, it was long known that activation loop-serine phosphorylation

was necessary for IKK-activation; we have now established that IKK2 phosphorylates itself

on tyrosine residue(s) in addition to activation loop-serines. We wanted to further understand

the significance of IKK2 autophosphorylation in the context of IκBα S32/S36

phosphorylation. It is known that disruption of a universal salt bridge mediated by the ATP-

binding Lys (K44 in IKK2) is detrimental for kinase activity. However, it is not known if disruption of this salt bridge has any bearing on substrate specificity of eukaryotic protein kinases (EPKs). During our investigation, we noted unusual behavior in the IKK2-K44M mutant. We had noticed earlier that IKK2-K44M was incapable of *de novo* autophosphorylation on tyrosine residues (Figure 2D). It also failed to phosphorylate itself on the activation loop Ser177 and 181 (Figure 3A). Interestingly, IKK2-K44M phosphorylated full-length WT IκBα robustly albeit much less efficiency than WT IKK2 (Figure 3B) in a radioactive *in vitro* kinase assay. IKK2-K44M also showed a minor auto-phosphorylation activity compared to WT IKK2 in the absence of substrate (Figure 3B). We performed a similar kinase assay using full-length WT IκBα where cold ATP was used as the phosphate donor, and phosphorylation status of IκBα at S32, S36 was monitored by immunoblotting using an antibody that specifically recognizes IκBα when phosphorylated at those two N-terminal serines. It revealed that IKK2-K44M is unable to phosphorylate S32/S36 of IκBα (Figure 3C). Since radioactive kinase assays report pan-phosphorylation of a substrate (Figure 3B), it possibly reflects phosphorylation of the PEST region of FL IκBα. We additionally compared IKK2-K44M with WT IKK2 using radioactive kinase assay with both FL IκBα WT and FL IκBαAA substrates in the absence or presence of Inhibitor VII. As anticipated, FL IκBα was phosphorylated much more efficiently than FL IκBαAA by WT IKK2, and phosphorylation was practically abolished in the presence of 10 μM Inhibitor VII (Figure 3D). In contrast, IKK2-K44M phosphorylated FL IκBαAA more efficiently than FL IκBα, and reduction in the presence of inhibitor VII was strikingly less (Figure 3D). It indicates that IKK2-K44M, despite retaining non-specific kinase activity towards the PEST region, lost phosphorylation activity towards S32, S36 of the *bona fide* substrate IκBα. We subjected 100ug of FL IKK2 K44M protein used in the above studies to LC-MS/MS analysis, which did not detect any contaminating kinase from Sf9 cells. It is noteworthy that an equivalent K to M mutation in Erk2 also did not abolish the basal activity, but rather retained ~5 % of it(Robbins et al., 1993). The above set of experiments confirm that K44 residue is

not only crucial for both IKK2's well-established serine phosphorylation as well as newly-

found tyrosine autophosphorylation activities, as well as its specificity for phosphorylating its

bona fide substrate IκBα at S32/36. Hence, these results strongly suggest that IKK2-

autophosphorylation is critical for its substrate specificity.

Tyrosine phosphorylation is critically important for IKK2 activity:

A number of earlier studies indicated that phosphorylation of IKK2 on Y169, Y188

and/or Y199 to be critical for its function(Darwech et al., 2010; Meyer et al., 2013). Our mass

spectrometric analyses identified multiple tyrosine residues to be phosphorylated, e.g.,

Y169, Y188, Y199, Y325 (data not shown). We compared available structural models of

human IKK2 (PDB ID: 4KIK, 4E3C) to investigate whether tyrosine phosphorylation within

the activation loop could be supported based on its structural features(Liu et al., 2013; Polley

et al., 2013). The structure of IKK2 dimer is asymmetric in that the activation loop of one

monomer is largely disordered while that of the other monomer is mostly ordered (PDB ID:

4KIK)(Liu et al., 2013). In the active monomers S177 and S181 are clearly observed. In

these active state conformers, the position and conformation of Tyr169 appears to be well-

poised for accepting a phosphate from ATP (Figure S4A) intramolecularly. A superposition

of IKK2 with a pseudo-substrate-bound PKA shows that the hydroxyl of Tyr169 in IKK2

projects toward the γ-phosphate of ATP similarly to the serine hydroxyl on the PKA inhibitor

pseudo-substrate(Knighton et al., 1991; Nolen et al., 2004) (Figure S4A & B). Interestingly,

Y169 is the DFG+1 (DLG in case of IKK2 and IKK1) residue while Y188 (equivalent to Y204

of PKA) is located at the end of activation loop prior to the conserved APE motif (Figure 4A

and 4B). It was previously established that the DFG+1 position is crucial for defining the

substrate specificity of a kinase(Chen et al., 2014). Y188 counterpart Y204 of PKA is

reported to allosterically modulate PKA activity by altering its conformational entropy(Ahuja

et al., 2017; Yang et al., 2005). We further focused our attention to Y169 and Y188 as they

are located in the activation loop of the kinase.

To test the effect of phosphorylation at Y169 and Y188, we generated Y169F and

Y188F mutants of IKK2. Radioactive *in vitro* kinase assays using recombinant purified

IKK2^{FL}-Y188F and IKK2^{FL}-Y169F mutants revealed a drastic reduction in the activity of IKK2

(Figure 4C). Since the DFG+1 residue in kinases (Y169 in IKK2) is known to dictate its

specificity, we created different mutants of the IκBα substrate where S32 and S36 were

singly and independently altered to either alanine (phosphoablative) or glutamic acid

(phosphomimetic) residues. Kinase activity of IKK2-Y169F was monitored in an *in vitro*

radioactive kinase assay using WT as well as mutant IκBα. Mutation of S36 to either alanine

or glutamate drastically reduced the phosphorylation signal of IκBα while that for the S32A/E

mutants were minimally affected (Figure 4D). This suggests that the presence of Tyr (and

possibly its phosphorylation) is critically important for signal-responsive phosphorylation of

IκBα.

Next, we tested the consequence of Y169F and Y188F mutants on the pro-

inflammatory signal-induced IKK activity in cells. We reconstituted IKK2 knock-out mouse

embryonic fibroblast (*ikk2$^{-/-}$* MEF) cells with WT, Y169F and Y188F mutants of IKK2, and

measured signal-induced activation of IKK2 upon TNF-α treatment. IKK2 activity is

significantly reduced in Y169F and nearly abolished in Y188F (Figure 4E**)**. Interestingly,

phosphorylation of activation loop serines of the less active Y169F mutant was only

marginally defective (Fig. 4F). This suggests that phosphorylation of the activation loop

serines at 177 and 181 is necessary but not sufficient for IKK2 to become fully active.

Together, these results suggest an unreported communication between the phosphorylation

of Y169 of IKK2 and that of S32 of IκBα.

Structural analyses of IKK2 autophosphorylation:

To get a more comprehensive picture, it is necessary to analyze at least three

phosphorylated states of IKK2: unphosphorylated (UnP-IKK2), phosphorylated at S177 and

S181 (p-IKK2), and phosphorylated at Y169, S177 and S181 (P-IKK2). We took advantage

of computational approaches using molecular dynamic (MD) simulations and flexible

molecular docking (see methods).[43–50] MD simulation was performed for 200ns for each of

these three states of IKK2, unphosphorylated and phosphorylated. Phosphorylation at S177,

S181 increased the stability of the kinase, which was further increased by the

phosphorylation at Y169 as evidenced by a gradual decrease in the total energy of system

(Figure 5A). A striking enhancement in stability of IKK2 upon this auto-phosphorylation is

reflected in the change of Tm from ~ 40 °C to ~ 50 °C in differential scanning calorimetry

(DSC) scans in the presence of ADP and ATP (Figure S4A). This behavior supports our

observation in MD simulation, and tallies with increased solubility of IKK2 observed after

auto-phosphorylation that was essential for its successful crystallization (Polley et al., 2013)

Interestingly, RMSD values increased with more phosphorylation (Figure 5B) despite

phosphorylation-induced stabilization, P-IKK2 showed the highest RMSD followed by p-IKK2

and UnP-IKK2, indicating increased internal motion in the kinase as a result of

phosphorylation. It may be noted that, the changes in total energy and RMSD are consistent

throughout the course of simulation though the magnitude is small. We found that,

phosphorylation at S177, S181 and S177, S181, Y169 induced distinct structural alteration

in the KD when compared with the unphosphorylated version (Figure 5C, models shown

separately in Figure S4B). Careful analyses of the trajectory of RMSF values for these three

states revealed that, different regions of the kinase domain had different RMSF values in

different phosphorylated states (Figure S4C). We observed that, while the glycine-rich loop

and the αC-helix in p-IKK2 were situated in a manner reminiscent of an active kinase (Figure

S4D), those in UnP and P-IKK2 were placed very differently. In fact, helix αC was found to

be distorted in P-IKK2, that is unlikely to support canonical phosphotransfer from ATP.

Activation segment/loop (activation loop) in these three differently phosphorylated states

adopted conformations distinct from each other. activation loop in the UnP model was

displayed out and moved closer to the tip of the C-lobe, whereas activation loop in p-IKK2

and P-IKK2 moved inwards and adopted conformations closer to the N-lobe. The formation

of the proper R- and C-spines in p-IKK2 confirmed its active form (Figure 5D). p-IKK2 KD

also exhibited other canonical features of an active kinase, viz., a salt bridge between K44

and E61 (K72, E91 in PKA, respectively), 'DFG in' (DLG here) conformation. Next, we

docked ATP onto the p-IKK2 structure (ΔG = -10.64 kcal/mol) using a flexible docking

protocol (see methods). The ATP and its phosphate groups exhibited a pose (Figure S4D)

very similar to that in the PKA structure (PDB ID: 1ATP), and the terminal phosphate was in

close proximity to Y169-OH making autophosphorylation of Y169 highly plausible (Figure

5E). Further analyses of the ATP-bound p-IKK2 structure confirm that presence of ATP

helped the R and C-spine form the desired continuum seen in the active form of a kinase

(Figure 5F). Superposition of the ATP from the p-IKK2 on P-IKK2 indicated that the

narrowing of ATP-binding pocket in P-IKK2 may lead to rejection of ATP emanating from

severe clashes between the glycine-rich loop and ATP (Figure 5G). Furthermore, the R-

spine in P-IKK2 was not continuous (Figure 5H), wherein M65 (L95 in PKA) of the four

conserved residues were located at a position far away from the rest of three residues that

constitute the R-spine, and from the glycine rich loop.

In addition, dynamic cross-correlation matrix (DCCM) or contact map of each structure

suggests that specific phosphorylation events render the kinase display distinct allosteric

changes at locations far from the phosphorylation sites themselves (Figure S4E).

ATP binding to p-IKK2 leads to autophosphorylation at Y169 generating P-IKK2 in an

ADP bound state. Interestingly, the superposition of ADP (from the docked complex of p-

IKK2 and ADP, ΔG = -9.88 kCal/mol, Figure 5I) onto the P-IKK2 structure confirmed that

while P-IKK2 was unable to accommodate ATP in a manner similar to that in other known

experimentally derived kinase structures, it could comfortably accommodate ADP into the

ATP-binding cleft (Figure 5J). It may be noted that, p-IKK2 has higher affinity for ATP than

ADP as indicated by their respective binding energy values. We speculate that ADP-bound

P-IKK2 is an intermediate step where presence of as many phosphate groups in the

activation loop and ADP in such a close proximity persuades the kinase to adopt a

conformation that can withstand this high density of negative charges albeit risking the

integrity of an active fold, and making some of those *de novo* phosphate groups transient in

nature to return back to the p-IKK2 state to readily bind to ATP that is abound in the

reaction.

These structural observations suggest that it is possible for the residues within the

flexible activation segment to undergo auto-phosphorylation and possibly render the kinase

with a novel phospho-transfer activity distinct from the conventional transfer of γ-phosphate

from ATP to the substrate.

Freshly auto-phosphorylated IKK2 phosphorylates IκBα even in absence of ATP:

We observed phosphorylation of tyrosine residue(s) only upon fresh ATP-treatment,

indicating that tyrosines of purified IKK2 loses its phosphate group during the course of

purification (Eckhart et al., 1979). It may be noted that, we did not observe any density of

phosphates on any of these residues in the human IKK2 crystal structure derived from ATP-

treated protein sample (Polley et al., 2013). Of note, human IKK2 crystals grown with

autophosphorylated IKK2 (PDB ID 4E3C) took about a month at 18-20°C before they were

subjected to X-ray diffraction. This suggested an intriguing possibility that the phosphate

linked to Tyr residue(s) may serve as a temporary phosphate-sink and could eventually be

transferred to substrate IκBα. To test this possibility, we first allowed purified IKK2 to

undergo auto-phosphorylation by incubating the excess kinase with limiting amounts of

radiolabeled γ-^{32}P-ATP followed by two passes through desalting spin columns (40 kDa MW

cut-off) to remove unreacted ATP. Nucleotide-free auto-phosphorylated IKK2 (P-IKK2) was

incubated with IκBα either in absence of nucleotide or with newly added unlabeled

nucleotides, and phosphorylation status of the substrate was subsequently monitored by

autoradiography after resolving the reaction mixture on an SDS-PAGE (Figure 6A). Transfer

of radiolabeled phosphate to IκBα was observed and the transfer was significantly enhanced

in the presence of cold ADP (Figure 6B). Phosphorylation of IκBα did not significantly

decrease in the presence of a high concentration of cold ATP. That addition of excess cold

ATP did not reduce the extent of IκBα phosphorylation-signal on the autoradiograph strongly

suggests the transfer of ^{32}P-phosphate from ^{32}P-IKK2 to IκBα (Figure 6B) by a relay

mechanism. This transfer perhaps occurs through a phosphoenzyme intermediate with or

without involving ADP.

We undertook a different strategy to further confirm transfer of phosphate from the

kinase (and not ATP) to the substrate. We auto-phosphorylated IKK2^{FL}-WT with excess cold

ATP and removed the unreacted ATP from the reaction mixture using size-exclusion

chromatography on a Superdex200 column (Figure 6C). P-IKK2 eluted at fractions

corresponding to MW between 670 and 158 kDa, i.e., much larger than free ATP (Figure

6D). This P-IKK2 protein was incubated with FL IκBα either in presence or absence of ADP

for different time periods. A similar reaction in presence of ATP instead of ADP was

performed as a positive control. Phosphorylation of S32 and S36 (detected by the phospho-

serine antibody specific to S32 and S36 of IκBα) was efficient in presence of 10 and 50 μM

ADP but absent when ADP was absent in the reaction mixture. Phosphorylation of IκBα in

the presence of 50 μM ATP was significantly more robust as anticipated (Figure 6E). The

minor detection of IκBα phosphorylation in absence of ADP in the radioactive assay and lack

of it in the cold immunoblotting assay (compare lanes marked with vertical arrows in Figures

6B and 6F) may indicate higher sensitivity of the radioactive assay. The need of ADP in

phosphotransfer prompted us to consider whether the ADP was contaminated with a trace

amount of ATP. ESI-MS analysis of 50 μM ADP did not detect any peak corresponding to

ATP (Figure S5A). Dependence of IκBα phosphorylation on ADP could raise another

possibility if microscopic reversibility generated ATP from ADP in the active site of the

kinase. To this end, we employed TLC-based analysis to detect generation/presence of ATP

in a similar setting as described in Fig. 5A-B which revealed that no detectable ATP was

present in the system, i.e., no detectable ATP was retained in the protein after passing it

through the spin column twice, nor did any γP^{32}-ATP was generated when incubated with

cold ADP (Figure S5B).

__Phosphotransfer from P-IKK2 is critical to fidelity and specificity of IκBα phosphorylation:__

We next tested if phosphotransfer is restricted only to the N-terminal signal responsive S32/S36 or to serines/threonines of the PEST domain as well. Radioactive kinase assay as described earlier was performed with full-length WT and mutant S32A, S36A (AA) IκBα as the phosphate recipient. Phosphate group(s) of P-IKK2 was not transferred to full-length IκBα AA which retains the serines and threonines within its PEST domain (Figure 7A), indicating phosphotransfer only to the N-terminal serines of IκBα that are S32 and S36. We also tested the effect of AMPPNP, an unhydrolyzable analogue of ATP, on the phosphotransfer, and found no inhibitory effect irrespective of the presence of ADP. This suggests adenine nucleotides poising the kinase for phosphotransfer (Figure 7A). A recent report proposed that IKK2 phosphorylates IκBα at S32 followed by S36 following a single binding event and that phosphorylation of S32 increases the phosphorylation rate of S36(Stephenson et al., 2022). Based on our results and other evidence, we propose that IKK2 phosphorylated at S177/S181 undergoes autophosphorylation on tyrosine residues in its activation loop area that serves as a phosphoenzyme intermediate to transfer phosphate group from IKK2 to S32 that enhances the rate of second site (S36) phosphorylation (Figure 7B). For the second site phosphorylation ATP serves as the phosphate donor.

Discussion:

Phosphorylation of substrates by protein kinases is regulated in various ways(Cullati et al., 2022; Sang et al., 2022). IKK2, a kinase central to inflammation, phosphorylates serine residues of its primary substrate IκBα with γ-phosphate of ATP. Here, we report multiple intriguing properties of IKK2 that provide specificity to its substrate phosphorylation. These include: 1) dual specificity that entails autophosphorylation of tyrosine residues of IKK2, in addition to known auto-phosphorylation at activation loop serines; 2) relay of phosphate to substrate IκBα from auto-phosphorylated IKK2 (P-IKK2) and 3) loss of

specificity of the kinase due to disruption of a universally conserved salt bridge (mediated by

K44).

Ser/Thr/Tyr family of eukaryotic protein kinases transfer γ-phosphate of ATP directly

to substrate, in general to S, T and Y residues(Cohen, 2002; Fischer and Krebs, 1955;

Hunter, 1991; Krebs and Fischer, 1955; Nolen et al., 2004). Histidine kinases (in prokaryotes

and in some lower eukaryotes) however shuttles the γ-phosphate to Asp residue on a

response regulator (RR) substrate through a phospho-His intermediate(Borkovich and

Simon, 1990; Gao and Stock, 2009; Hunter, 2022; Kalagiri and Hunter, 2021; Laub et al.,

2007; Robbins et al., 1993). Our present analyses of eukaryotic protein kinase IKK2

revealed its surprising property in which IKK2 phosphorylated at activation loop-serines

(S177/S181) could be further auto-phosphorylated by ATP at multiple other residues yielding

another form (P-IKK2) (Figure 7B). This marks IKK2 as an autocatalytic dual-specificity

kinase rather than a prototype S/T kinase. This multisite phosphorylated form of IKK2 (P-

IKK2) could transfer phosphate(s) to substrate IκBα in presence of ADP without requiring

exogenous supply of fresh ATP, hinting at an unusual property of IKK2 acting possibly as a

phosphate-sink. It is possible that phospho-tyrosines residues of IKK2 act as this sink. The

Y169 located at the DFG+1 (DLG in IKK2) position is unique only to IKK1 and IKK2, and is

absent even in close mammalian homologues such as IKKε and TBK1. Interestingly, neither

IKKε nor TBK1 can phosphorylate IκBα at S32 and S36, *in vitro* or *in vivo*(Larabi et al., 2013;

Tu et al., 2013). A recent report also indicated the role of DFG+1 residue in differentiating

Ser versus Thr residues(Chen et al., 2014). These observations hinted to an involvement of

Y169 in substrate selectivity. Indeed, our experimental data suggests a pivotal role of Y169

in phosphorylating S32 of IκBα both *in vitro* and *ex vivo* (Figure 4D & 4E). It is worthwhile to

note that, a phosphoablative mutation at S32 in IκBα to Ile (S32I) is associated with

ectodermal dysplasia and T-cell immunodeficiency in patients(Courtois et al., 2003; Mooster

et al., 2015). More interestingly, the placement of Y169 in the kinase active site makes it

amenable to a phosphorylation-dephosphorylation cycle, suggesting its possible role both in

phosphotransfer and substrate specificity. Mutation of Y188, a conserved tyrosine, to non-

phosphorylatable phenylalanine abolished the kinase activity of IKK2 *in vitro* as well as in

MEFs upon TNF-treatment (Figure 4C & 4E). The placement of Y188 indicates that its

mutation will be detrimental to the structural fold of kinase, thus it is unclear if or how Y188

plays a role in the phosphotransfer. Since phosphotransfer to S32, S36 of IκBα is observed

with auto-phosphorylated IKK2 EE, transferred phosphates are not from S177 or S181.

Kinase domain autophosphorylation is known to regulate function of other kinases as well,

e.g., autophosphorylation at T220 greatly influenced both the activity and substrate

specificity of CK1(Cullati et al., 2022).

The strength of input signal, level of IκBα degradation, and NF-κB activation

correlate with the amplitude, duration, and kinetics of IKK activity(Behar and Hoffmann,

2013; Cheong et al., 2006). This digital (all-or-none) activation profile of NF-κB appears to

be due to rapid activation and inactivation of IKK2. IKK targets multiple substrates *in vivo*;

however, the rapid activation of NF-κB is regulated primarily by IκBα. How IKK specifically

selects IκBα for phosphorylation is unclear, although NEMO is reported to direct the kinase

activity of IKK2 specifically towards IκBα(Schröfelbauer et al., 2012). We observe that P-

IKK2 could transfer phosphate(s) specifically to the S32/S36 of IκBα, and not to other sites

(such as the PEST sequence of IκBα) (Figure 6B, 6E and 7A). We surmise if transiency of

multisite phosphorylated IKK2 (P-IKK2) underlie this activation-inactivation events during

signal response. It is possible that regulatory system of IKK2 is somewhat analogous to the

two-component histidine kinase-effector systems in prokaryotes that show a threshold

activation response(Bhate et al., 2015; Lamarche et al., 2008; Laub and Goulian, 2007). It is

noteworthy that, transfer of a phosphate moiety directly from pHis is energetically much

more probable than that from a pTyr, unless it is an enzymatic process which is likely the

case for IKK2:IκBα system.

Tyrosines in activation loop of IKK2 are previously reported to undergo signal-

induced phosphorylation in cells. These modifications were proposed to be mediated by a

tyrosine kinase although there is no evidence of a specific tyrosine kinase(Meyer et al., 2013; Otero et al., 2008; Rieke et al., 2011). Several members of the Ser/Thr kinase family undergo tyrosine phosphorylation and these kinases employ diverse mechanisms for auto-phosphorylation(Bhattacharyya et al., 2006; Ge et al., 2002; Lochhead, 2009; Lochhead et al., 2006, 2005; Tigno-Aranjuez et al., 2010).

The phosphotransfer from P-IKK2 to IκBα monitored here is detection of a single turnover event in absence of the phosphate donor, ATP in the bulk (Figure 6B). Phosphorylation on IKK2 occurs at multiple sites, and not every phosphorylated residue in P-IKK2 expected to transfer its phosphate to IκBα. There is more than one type of phosphorylated residues in P-IKK2 distributed over the length of the kinase, arguably with differential stability wherein only a fraction of the total pool can be relayed. In addition, it is likely that the phosphorylated IKK2 in the bulk harbors a non-homogenous pool where differently phosphorylated species of the kinase are present, and are indistinguishable in our assay schemes. Another intriguing feature of phosphorelay in IKK2 is IKK2's dependence on ADP (Figure 6B & 6E). It was also observed that, the generic kinase inhibitor AMPPNP (unhydrolyzable ATP analogue) failed to inhibit phosphor-transfer from IKK2 (Figure 7A) and IκBα phosphorylation (Figure S2C). Taken together, we surmise that presence of ADP or AMP-PNP help IKK2 attain a conformational state poised for phosphorylating S32/36 of IκBα even in absence of fresh exogenous ATP. Similar phenomenon was reported for IRE1, where ADP was found to be a more potent stimulator of its ribonuclease activity and that AMPPNP could also potentiate this activity in IRE1(Lee et al., 2008; Sidrauski and Walter, 1997).The list of Ser/Thr kinase family proteins that undergo tyrosine phosphorylation is growing that employ diverse mechanisms for auto-phosphorylation(Bhattacharyya et al., 2006; Ge et al., 2002; Lochhead, 2009; Lochhead et al., 2006, 2005; Tigno-Aranjuez et al., 2010). The most notable example of this dual specificity are DYRK and GSK3β. However, the role of Tyr-phosphorylation in DYRK/GSK3β is not related to that in IKK2, nor does it render DYRKs/ GSK3β capable of relaying phosphates to the substrates. To the best of our

knowledge the only other reported example of an EPK, though atypical, transferring

phosphate through a phosphoenzyme intermediate is MHCK of *Dictyostelium*

discoideum(Ye et al., 2010).

Our study reveals tyrosine auto-phosphorylation in IKK2 and indicates that

consequences of Ser and Tyr autophosphorylations in dual-specific IKK2 are distinct.

Moreover, the crucial tyrosine residues are positioned in such a manner that some that are

amenable to auto-phosphorylation are essential for substrate phosphorylation. Most

surprisingly, the tyrosine-linked phosphate groups can be transferred by a relay mechanism

to specific serines in IκBα, an event reminiscing phospho-relay in two-component systems

prevalent in prokaryotes and some lower eukaryotes. Our study also reveals the role of K44

of IKK2 in auto-phosphorylation and thus in specific phosphorylation of S32, S36 of IκBα,

although the mutant kinase retains its non-specific kinase activity. It may be noted that, while

a monomeric version of IKK2 (devoid of NBD and large portion of the dimeric interface in the

SDD) still retained its specificity for the N-terminal serines of IκBα(Hauenstein et al., 2014),

while a shorter construct containing only the KD and ULD lacked this specificity that

phosphorylated the PEST domain like CK2(Shaul et al., 2008). It requires further

investigation to determine if a structural state of IKK2, perhaps through a different

modification or through binding to another protein, allow it to phosphorylate substrates

relatively non-specifically.

In all, it is intriguing how the gateway to the activation of NF-κB is regulated through

unique phosphorylation events in IKK2 (Figure 7C). It appears that a multi-layered fail-safe

mechanism is devised to tightly control NF-κB activation. Firstly, in absence of upstream

signals IKK-complex is unable to phosphorylate Ser 32,36 of IκBα efficiently thereby keeping

check on the aberrant and untimely activation of NF-κB. Upon encountering proinflammatory

cues, cells need to activate NF-κB immediately and efficiently, i.e., IKK needs to be

activated and phosphorylate Ser 32,36 of IκBα. NEMO warrants IKK-activation and ensures

that the IKK-complex specifically chooses IκBα from a large pool of substrates of IKK2. IKK2

is designed in such a manner that it phosphorylates itself at activation loop-tyrosines when

activated, such that phosphate group(s) can be relayed directly to Ser32,36 of IκBα with

great fidelity, thus leaving little chance of a misconstrued signaling event thereby confirming

NF-κB activation (Figure 7C). Our discovery might also present the beginning of a new

aspect in eukaryotic cell-signaling by EPKs and provide a foundation for mechanistic studies

of an intriguing phospho-transfer reaction that helps accomplish the desired specificity. To

the best of our knowledge, it is the first report, apart from MHCK of *Dictyostelium discoideum*

(an atypical EPK), to suggest that a metazoan EPK confirms its specificity through the

formation of a phosphoenzyme intermediate.

Methods summary:

Protein purification, auto-phosphorylation assay, immunoprecipitation coupled kinase

assay, phosphotransfer assay and other experiments have been described in details in the

supplementary document.

Acknowledgement:

We thank Tony Hunter (Salk Institute) and Kaushik Biswas (BI) for commenting on

the work. Authors also thank members of Polley lab (BI) and Ghosh Lab (UCSD) for their

continuous support. This work was supported by an Intermediate Fellowship from the DBT

Wellcome Trust India Alliance (IA/I/15/1/501852) & intramural funding from Bose Institute to

SP, NIH grant to GG (AI163327) and American Cancer Society grant RSG-08-287-01-GMC

to TH. Biochemistry research at SDSU is supported in part by the California Metabolic

Research Foundation. PB was supported by the graduate research fellowship from Bose

Institute. SC acknowledges CSIR-IICB for infrastructure support. AC acknowledges ICMR

for post-doctoral Research Associateship [BMI/11(55)2022].

562
Author contribution:

SP, GG and TB designed experiments; PB, SP, TB & GG performed experiments.

Computational analyses were performed by AC & SC. SP, GG, TB, PB, TH, SC and AC

analyzed the results. SP wrote the manuscript with the help of PB, TB and AC in preparing

figures and images. GG, TB, PB and TH edited/modified the manuscript. All authors

commented on the manuscript.

Ethics declarations:

Competing interests: The authors declare no competing financial interests.

References:

Ahuja LG, Kornev AP, McClendon CL, Veglia G, Taylor SS. 2017. Mutation of a kinase
allosteric node uncouples dynamics linked to phosphotransfer. *Proc Natl Acad Sci U*
S A **114**:E931–E940. doi:10.1073/pnas.1620667114
Antonia RJ, Hagan RS, Baldwin AS. 2021. Expanding the View of IKK: New Substrates and
New Biology. *Trends Cell Biol* **31**:166–178. doi:10.1016/j.tcb.2020.12.003
Barroga CF, Stevenson JK, Schwarz EM, Verma IM. 1995. Constitutive phosphorylation of I
kappa B alpha by casein kinase II. *Proceedings of the National Academy of Sciences*
92:7637–7641. doi:10.1073/pnas.92.17.7637
Behar M, Hoffmann A. 2013. Tunable Signal Processing through a Kinase Control Cycle: the
IKK Signaling Node. *Biophys J* **105**:231–241. doi:10.1016/j.bpj.2013.05.013
Bhate MP, Molnar KS, Goulian M, DeGrado WF. 2015. Signal Transduction in Histidine
Kinases: Insights from New Structures. *Structure* **23**:981–994.
doi:10.1016/j.str.2015.04.002
Bhattacharyya RP, Reményi A, Good MC, Bashor CJ, Falick AM, Lim WA. 2006. The Ste5
scaffold allosterically modulates signaling output of the yeast mating pathway.
Science **311**:822–826. doi:10.1126/science.1120941
Borkovich KA, Simon MI. 1990. The dynamics of protein phosphorylation in bacterial
chemotaxis. *Cell* **63**:1339–1348. doi:10.1016/0092-8674(90)90429-i
Brown K, Gerstberger S, Carlson L, Franzoso G, Siebenlist U. 1995. Control of I kappa B-
alpha proteolysis by site-specific, signal-induced phosphorylation. *Science*
267:1485–1488. doi:10.1126/science.7878466
Chen C, Ha BH, Thevenin AF, Lou HJ, Zhang R, Yip KY, Peterson JR, Gerstein M, Kim PM,
Filippakopoulos P, Knapp S, Boggon TJ, Turk BE. 2014. Identification of a major
determinant for serine-threonine kinase phosphoacceptor specificity. *Mol Cell*
53:140–7. doi:10.1016/j.molcel.2013.11.013
Chen ZJ. 2012. Ubiquitination in Signaling to and Activation of IKK. *Immunol Rev* **246**:95–
106. doi:10.1111/j.1600-065X.2012.01108.x
Cheong R, Bergmann A, Werner SL, Regal J, Hoffmann A, Levchenko A. 2006. Transient
IkappaB kinase activity mediates temporal NF-kappaB dynamics in response to a
wide range of tumor necrosis factor-alpha doses. *J Biol Chem* **281**:2945–50.
doi:10.1074/jbc.M510085200
Cohen P. 2002. The origins of protein phosphorylation. *Nat Cell Biol* **4**:E127–E130.
doi:10.1038/ncb0502-e127
Courtois G, Smahi A, Reichenbach J, Döffinger R, Cancrini C, Bonnet M, Puel A, Chable-
Bessia C, Yamaoka S, Feinberg J, Dupuis-Girod S, Bodemer C, Livadiotti S, Novelli
F, Rossi P, Fischer A, Israël A, Munnich A, Le Deist F, Casanova J-L. 2003. A
hypermorphic IkappaBalpha mutation is associated with autosomal dominant
anhidrotic ectodermal dysplasia and T cell immunodeficiency. *J Clin Invest*
112:1108–1115. doi:10.1172/JCI18714
Cullati SN, Chaikuad A, Chen J-S, Gebel J, Tesmer L, Zhubi R, Navarrete-Perea J, Guillen
RX, Gygi SP, Hummer G, Dötsch V, Knapp S, Gould KL. 2022. Kinase domain

autophosphorylation rewires the activity and substrate specificity of CK1 enzymes.
Mol Cell **82**:2006-2020.e8. doi:10.1016/j.molcel.2022.03.005
Darwech I, Otero JE, Alhawagri MA, Abu-Amer Y. 2010. Tyrosine phosphorylation is
required for IkappaB kinase-beta (IKKbeta) activation and function in
osteoclastogenesis. *J Biol Chem* **285**:25522–25530. doi:10.1074/jbc.M110.121533
Delhase M, Hayakawa M, Chen Y, Karin M. 1999. Positive and Negative Regulation of IκB
Kinase Activity Through IKKβ Subunit Phosphorylation. *Science* **284**:309–313.
doi:10.1126/science.284.5412.309
DiDonato JA, Hayakawa M, Rothwarf DM, Zandi E, Karin M. 1997. A cytokine-responsive
IκB kinase that activates the transcription factor NF-κB. *Nature* **388**:548–554.
doi:10.1038/41493
Du M, Ea CK, Fang Y, Chen ZJ. 2022. Liquid phase separation of NEMO induced by
polyubiquitin chains activates NF-kappaB. *Mol Cell* **82**:2415-2426 e5.
doi:10.1016/j.molcel.2022.03.037
Ea CK, Deng L, Xia ZP, Pineda G, Chen ZJ. 2006. Activation of IKK by TNFalpha requires
site-specific ubiquitination of RIP1 and polyubiquitin binding by NEMO. *Mol Cell*
22:245–57. doi:10.1016/j.molcel.2006.03.026
Eckhart W, Hutchinson MA, Hunter T. 1979. An activity phosphorylating tyrosine in polyoma
T antigen immunoprecipitates. *Cell* **18**:925–933. doi:10.1016/0092-8674(79)90205-8
Fischer EH, Krebs EG. 1955. Conversion of phosphorylase b to phosphorylase a in muscle
extracts. *J Biol Chem* **216**:121–32.
Gao R, Stock AM. 2009. Biological Insights from Structures of Two-Component Proteins.
Annu Rev Microbiol **63**:133–154. doi:10.1146/annurev.micro.091208.073214
Ge B, Gram H, Di Padova F, Huang B, New L, Ulevitch RJ, Luo Y, Han J. 2002. MAPKK-
independent activation of p38alpha mediated by TAB1-dependent
autophosphorylation of p38alpha. *Science* **295**:1291–1294.
doi:10.1126/science.1067289
Ghosh S, Karin M. 2002. Missing pieces in the NF-kappaB puzzle. *Cell* **109 Suppl**:S81-96.
Hacker H, Karin M. 2006. Regulation and function of IKK and IKK-related kinases. *Sci STKE*
2006:re13. doi:10.1126/stke.3572006re13
Hauenstein AV, Rogers WE, Shaul JD, Huang DB, Ghosh G, Huxford T. 2014. Probing
kinase activation and substrate specificity with an engineered monomeric IKK2.
Biochemistry **53**:2064–73. doi:10.1021/bi401551r
Hayden MS, Ghosh S. 2008. Shared principles in NF-kappaB signaling. *Cell* **132**:344–62.
doi:10.1016/j.cell.2008.01.020
Hinz M, Scheidereit C. 2014. The IκB kinase complex in NF-κB regulation and beyond.
EMBO Rep **15**:46–61. doi:10.1002/embr.201337983
Hoffmann A, Baltimore D. 2006. Circuitry of nuclear factor kappaB signaling. *Immunol Rev*
210:171–86. doi:10.1111/j.0105-2896.2006.00375.x
Hunter T. 2022. A journey from phosphotyrosine to phosphohistidine and beyond. *Mol Cell*
82:2190–2200. doi:10.1016/j.molcel.2022.05.007
Hunter T. 1991. Protein kinase classification. *Methods Enzymol* **200**:3–37.
doi:10.1016/0076-6879(91)00125-g
Huse M, Kuriyan J. 2002. The conformational plasticity of protein kinases. *Cell* **109**:275–282.
doi:10.1016/s0092-8674(02)00741-9
Kalagiri R, Hunter T. 2021. The many ways that nature has exploited the unusual structural
and chemical properties of phosphohistidine for use in proteins. *Biochem J*
478:3575–3596. doi:10.1042/BCJ20210533
Karin M, Ben-Neriah Y. 2000. Phosphorylation meets ubiquitination: the control of NF-
[kappa]B activity. *Annu Rev Immunol* **18**:621–63.
doi:10.1146/annurev.immunol.18.1.621
Knighton DR, Zheng JH, Ten Eyck LF, Xuong NH, Taylor SS, Sowadski JM. 1991. Structure
of a peptide inhibitor bound to the catalytic subunit of cyclic adenosine
monophosphate-dependent protein kinase. *Science* **253**:414–420.
doi:10.1126/science.1862343

Krebs EG, Fischer EH. 1955. Phosphorylase activity of skeletal muscle extracts. *J Biol*
Chem **216**:113–20.
Lamarche MG, Wanner BL, Crépin S, Harel J. 2008. The phosphate regulon and bacterial
virulence: a regulatory network connecting phosphate homeostasis and
pathogenesis. *FEMS Microbiol Rev* **32**:461–473. doi:10.1111/j.1574-
6976.2008.00101.x
Larabi A, Devos JM, Ng SL, Nanao MH, Round A, Maniatis T, Panne D. 2013. Crystal
structure and mechanism of activation of TANK-binding kinase 1. *Cell Rep* **3**:734–46.
doi:10.1016/j.celrep.2013.01.034
Laub MT, Biondi EG, Skerker JM. 2007. Phosphotransfer profiling: systematic mapping of
two-component signal transduction pathways and phosphorelays. *Methods Enzymol*
423:531–48. doi:10.1016/S0076-6879(07)23026-5
Laub MT, Goulian M. 2007. Specificity in two-component signal transduction pathways.
Annu Rev Genet **41**:121–45. doi:10.1146/annurev.genet.41.042007.170548
Lee KPK, Dey M, Neculai D, Cao C, Dever TE, Sicheri F. 2008. Structure of the dual
enzyme Ire1 reveals the basis for catalysis and regulation in non-conventional RNA
splicing. *Cell* **132**:89–100. doi:10.1016/j.cell.2007.10.057
Lin YC, Brown K, Siebenlist U. 1995. Activation of NF-kappa B requires proteolysis of the
inhibitor I kappa B-alpha: signal-induced phosphorylation of I kappa B-alpha alone
does not release active NF-kappa B. *Proceedings of the National Academy of*
Sciences **92**:552–556. doi:10.1073/pnas.92.2.552
Liu F, Xia Y, Parker AS, Verma IM. 2012. IKK biology. *Immunol Rev* **246**:239–53.
doi:10.1111/j.1600-065X.2012.01107.x
Liu S, Misquitta YR, Olland A, Johnson MA, Kelleher KS, Kriz R, Lin LL, Stahl M, Mosyak L.
2013. Crystal Structure of a Human IκB Kinase β Asymmetric Dimer. *J Biol Chem*
288:22758–22767. doi:10.1074/jbc.M113.482596
Lochhead PA. 2009. Protein kinase activation loop autophosphorylation in cis: overcoming a
Catch-22 situation. *Sci Signal* **2**:pe4. doi:10.1126/scisignal.254pe4
Lochhead PA, Kinstrie R, Sibbet G, Rawjee T, Morrice N, Cleghon V. 2006. A chaperone-
dependent GSK3beta transitional intermediate mediates activation-loop
autophosphorylation. *Mol Cell* **24**:627–633. doi:10.1016/j.molcel.2006.10.009
Lochhead PA, Sibbet G, Morrice N, Cleghon V. 2005. Activation-loop autophosphorylation is
mediated by a novel transitional intermediate form of DYRKs. *Cell* **121**:925–936.
doi:10.1016/j.cell.2005.03.034
Meyer AN, Drafahl KA, McAndrew CW, Gilda JE, Gallo LH, Haas M, Brill LM, Donoghue DJ.
2013. Tyrosine Phosphorylation Allows Integration of Multiple Signaling Inputs by
IKKβ. *PLoS One* **8**:e84497. doi:10.1371/journal.pone.0084497
Mooster JL, Le Bras S, Massaad MJ, Jabara H, Yoon J, Galand C, Heesters BA, Burton OT,
Mattoo H, Manis J, Geha RS. 2015. Defective lymphoid organogenesis underlies the
immune deficiency caused by a heterozygous S32I mutation in IκBα. *J Exp Med*
212:185–202. doi:10.1084/jem.20140979
Nolen B, Taylor S, Ghosh G. 2004. Regulation of Protein Kinases: Controlling Activity
through Activation Segment Conformation. *Molecular Cell* **15**:661–675.
doi:10.1016/j.molcel.2004.08.024
Otero JE, Dai S, Foglia D, Alhawagri M, Vacher J, Pasparakis M, Abu-Amer Y. 2008.
Defective osteoclastogenesis by IKKbeta-null precursors is a result of receptor
activator of NF-kappaB ligand (RANKL)-induced JNK-dependent apoptosis and
impaired differentiation. *J Biol Chem* **283**:24546–24553.
doi:10.1074/jbc.M800434200
Polley S, Huang D-B, Hauenstein AV, Fusco AJ, Zhong X, Vu D, Schröfelbauer B, Kim Y,
Hoffmann A, Verma IM, Ghosh G, Huxford T. 2013. A structural basis for IκB kinase
2 activation via oligomerization-dependent trans auto-phosphorylation. *PLoS Biol*
11:e1001581. doi:10.1371/journal.pbio.1001581

Rieke C, Papendieck A, Sokolova O, Naumann M. 2011. Helicobacter pylori-induced
tyrosine phosphorylation of IKKβ contributes to NF-κB activation. *Biol Chem*
392:387–393. doi:10.1515/BC.2011.029
Robbins DJ, Zhen E, Owaki H, Vanderbilt CA, Ebert D, Geppert TD, Cobb MH. 1993.
Regulation and properties of extracellular signal-regulated protein kinases 1 and 2 in
vitro. *J Biol Chem* **268**:5097–106.
Rothwarf DM, Zandi E, Natoli G, Karin M. 1998. IKK-gamma is an essential regulatory
subunit of the IkappaB kinase complex. *Nature* **395**:297–300. doi:10.1038/26261
Sang D, Shu T, Pantoja CF, Ibáñez de Opakua A, Zweckstetter M, Holt LJ. 2022.
Condensed-phase signaling can expand kinase specificity and respond to
macromolecular crowding. *Mol Cell* **82**:3693-3711.e10.
doi:10.1016/j.molcel.2022.08.016
Scheidereit C. 2006. IkappaB kinase complexes: gateways to NF-kappaB activation and
transcription. *Oncogene* **25**:6685–705. doi:10.1038/sj.onc.1209934
Schrofelbauer B, Hoffmann A. 2011. How do pleiotropic kinase hubs mediate specific
signaling by TNFR superfamily members? *Immunol Rev* **244**:29–43.
doi:10.1111/j.1600-065X.2011.01060.x
Schröfelbauer B, Polley S, Behar M, Ghosh G, Hoffmann A. 2012. NEMO ensures signaling
specificity of the pleiotropic IKKβ by directing its kinase activity toward IκBα. *Mol Cell*
47:111–121. doi:10.1016/j.molcel.2012.04.020
Shaul JD, Farina A, Huxford T. 2008. The human IKKbeta subunit kinase domain displays
CK2-like phosphorylation specificity. *Biochem Biophys Res Commun* **374**:592–7.
doi:10.1016/j.bbrc.2008.07.082
Sidrauski C, Walter P. 1997. The transmembrane kinase Ire1p is a site-specific
endonuclease that initiates mRNA splicing in the unfolded protein response. *Cell*
90:1031–1039. doi:10.1016/s0092-8674(00)80369-4
Stephenson AA, Taggart DJ, Xu G, Fowler JD, Wu H, Suo Z. 2022. The inhibitor of κB
kinase β (IKKβ) phosphorylates IκBα twice in a single binding event through a
sequential mechanism. *J Biol Chem* **299**:102796. doi:10.1016/j.jbc.2022.102796
Tergaonkar V, Bottero V, Ikawa M, Li Q, Verma IM. 2003. IkappaB kinase-independent
IkappaBalpha degradation pathway: functional NF-kappaB activity and implications
for cancer therapy. *Mol Cell Biol* **23**:8070–83.
Tigno-Aranjuez JT, Asara JM, Abbott DW. 2010. Inhibition of RIP2's tyrosine kinase activity
limits NOD2-driven cytokine responses. *Genes Dev* **24**:2666–2677.
doi:10.1101/gad.1964410
Tu D, Zhu Z, Zhou AY, Yun CH, Lee KE, Toms AV, Li Y, Dunn GP, Chan E, Thai T, Yang S,
Ficarro SB, Marto JA, Jeon H, Hahn WC, Barbie DA, Eck MJ. 2013. Structure and
ubiquitination-dependent activation of TANK-binding kinase 1. *Cell Rep* **3**:747–58.
doi:10.1016/j.celrep.2013.01.033
Xu G, Lo Y-C, Li Q, Napolitano G, Wu X, Jiang X, Dreano M, Karin M, Wu H. 2011. Crystal
structure of inhibitor of κB kinase β (IKKβ). *Nature* **472**:325–330.
doi:10.1038/nature09853
Yang J, Garrod SM, Deal MS, Anand GS, Woods VL Jr, Taylor S. 2005. Allosteric network of
cAMP-dependent protein kinase revealed by mutation of Tyr204 in the P+1 loop. *J*
Mol Biol **346**:191–201. doi:10.1016/j.jmb.2004.11.030
Ye Q, Crawley SW, Yang Y, Côté GP, Jia Z. 2010. Crystal structure of the alpha-kinase
domain of Dictyostelium myosin heavy chain kinase A. *Sci Signal* **3**:ra17.
doi:10.1126/scisignal.2000525
Zandi E, Chen Y, Karin M. 1998. Direct phosphorylation of IkappaB by IKKalpha and
IKKbeta: discrimination between free and NF-kappaB-bound substrate. *Science*
281:1360–1363. doi:10.1126/science.281.5381.1360
Zandi E, Rothwarf DM, Delhase M, Hayakawa M, Karin M. 1997. The IkappaB kinase
complex (IKK) contains two kinase subunits, IKKalpha and IKKbeta, necessary for
IkappaB phosphorylation and NF-kappaB activation. *Cell* **91**:243–252.
doi:10.1016/s0092-8674(00)80406-7

Zhang J, Clark K, Lawrence T, Peggie MW, Cohen P. 2014. An unexpected twist to the
activation of IKKbeta: TAK1 primes IKKbeta for activation by autophosphorylation.
Biochem J **461**:531–7. doi:10.1042/BJ20140444

Figure legends:

Figure 1: Autophosphorylation of IKK2 at hitherto uncharacterized sites. (A) NEMO is known to remain bound to the IKK-complex pre- and post-stimulation, helps in activation as well as substrate recognition to channelize IKK-activity towards IκBα. However, it is not clear how IKK specifically phosphorylates the Ser32&36 of IκBα. **(B)** Domain organisation of FL IKK2 depicting functional kinase domain (KD), ubiquitin-like domain (ULD), and scaffold dimerization domain (SDD) based on the X-ray structures. Known sites of Ser phosphorylation and NEMO interacting regions at the extreme C-termini are highlighted. Serine and tyrosine residues in the activation loop are shown, mutation of which to Glu renders IKK2 constitutively active. The conserved ATP-interacting Lys (at position 44) is also shown. **(C)** Autophosphorylation of FL IKK2WT upon incubation with radioactive ATP for different time periods in an *in vitro* kinase assay. **(D)** *In vitro* radioactive kinase assays performed to check the effect of NEMO on autophosphorylation of FL IKK2WT (left panel), and that of NEMO and IκBα on autophosphorylation and substrate phosphorylation (right panel) activity of FL IKK2WT. **(E)** Effect of Inhibitor VII on FL IKK2WT autophosphorylation and IκBα substrate phosphorylation at different inhibitor concentrations in a radioactive *in vitro* kinase assay. A schematic representation of the assay procedure is shown in Figure S1B. This assay was performed twice. **(F)** Radioactive kinase assay conducted to check auto- and substrate phosphorylation of full length and deletion constructs of constitutively active form of IKK2 harbouring phosphomimetic Ser177Glu and Ser181Glu mutations.

Figure 2: IKK2 possesses autocatalytic dual specificity. (A) *In vitro* kinase assay, followed by immunoblotting with FL IKK2WT using antibodies against phosphor-Ser (177/181) and phospho-Tyr. **(B)** A different commercial source of phospho-Tyr antibody was used to detect the same. **(C)** Effect of Inhibitor VII on tyrosine autophosphorylation of FL IKK2WT. **(D)** Immunoblotting performed with phospho-Tyr antibody to check autophosphorylation of IKK2 mutant K44M at different time points compared to that of IKK2 WT. **(E)** Tyrosine autophosphorylation and substrate phosphorylation of GST (1-54) IκBα in presence of full length and deletion mutants of IKK2 harbouring phosphomimetic Ser177Glu and Ser181Glu mutations. **(F)** *De novo* auto-phosphorylation of IKK2 on tyrosine monitored by phospho-Ser and phospho-Tyr specific monoclonal antibodies. **(G)** Autophosphorylation and subsequent immunoblot analysis of FLIKK2 mutant S177A, S181A using antibodies against phospho-IKK2-Ser(177/181) and phospho-Tyr.

Figure 3: Autophosphorylation of IKK2 is critical to the specificity of IκBα phosphorylation. (A) Immunoblotting performed with phospho-IKK2 Ser(177/181) antibody

to check autophosphorylation of IKK2 mutant K44M at different time points compared to that

of IKK2 WT. **(B)** Comparison of substrate and autophosphorylation activities of FL IKK2 WT

and K44M mutant in an *in vitro* radioactive kinase assay performed with or without FL IκBα

WT as the substrate. **(C)** Residue level specificity of phosphorylation by the FL IKK2 K44M

was analysed using an antibody that specifically recognizes phospho-S32/36 of IκBα. **(D)**

Differential sensitivity of specific vs. non-specific phosphorylation by IKK2 as assessed by

the effect of Inhibitor VII on FL IKK2 WT and IKK2 K44M using IκBα WT and AA as

substrates.

Figure 4: Tyrosine autophosphorylation is critical for IKK2 activity. (A) Position of

canonically important residues in the AL (purple ribbon) of IKK2-KD are shown in the context

of its structure (PDB ID 4E3C; surface representation; KD is shown light green with respect

to the SDD in teal). Autophosphorylated tyrosine residues identified in this study are shown

in green. **(B)** Alignment of the activation segment sequences of different IKK family kinases.

Conservation of Tyr at the DFG+1 (DLG in case of IKK1 and IKK2) is retained only in a

stress response related plant kinase SnRK2, but not in other IKK homologues or any known

dual specificity kinases, e.g., DYRK and GSK3β (both contain Ser at that position). Tyr188

(204 in PKA) is conserved.   **(C)** Radioactive *in vitro* kinase assay performed with IKK2

Y169F and Y188F mutants using GST (1-54) IκBα WT and AA as substrates. **(D)**

Radioactive *in vitro* kinase assay performed with IKK2 Y169F mutant with a range of GST

(1-54) IκBα mutants where the substrate phosphorylation signal is abolished in case of S36

A/E IκBα. **(E)** Severe reduction in TNFα-induced IKK activity in *ikk2-/-* MEF-3T3 cells

reconstituted with mutant (Y169F and Y188F) IKK2s compared to wild-type. Kinase assay

was performed with IKK immunoprecipitated (IP-ed) with anti-NEMO antibody from whole

cell extract. **(F)** AL-serine phosphorylation status of IKK2 WT and Y169F was assessed in

reconstituted MEFs post-stimulation with TNF-α after immunoprecipitating the kinase with

monoclonal anti-HA antibody.

Figure 5: Structural analyses of IKK2 autophosphorylation. (A) Molecular dynamics

(MD) simulations of all three differently phosphorylated states of IKK2 was performed. 200ns

trajectory of energy for each state is shown in golden yellow, blue and copper red

representing UnP-IKK2, p-IKK2 and P-IKK2, respectively. Same color coding has been

followed throughout this figure and the corresponding supplementary figure. **(B)** 200ns

trajectory of RMSD for each state is shown**. (C)** Residues forming canonical R- (labelled in

blue) & C-spines (labelled in dark gray) representative of an active conformation in p-IKK2

are shown.   **(D)** Superposition of structures representing the differently phosphorylated

states post 200ns MD simulation. Relevant sections, e.g., activation loop, αC-helix, Gly-rich

loop areas are highlighted in the right panel. **(E)** ATP-bound p-IKK2 structure is depicted. In

the right panel, relevant residues are highlighted. Y169 is found in a position poised for

autophosphorylation. **(F)** ATP completes the desired continuum of R- & C-spine observed in

active kinase conformations. **(G)** P-IKK2 cannot accommodate ATP in its cleft, and αC-helix

is displaced. **(H)** M is moved away from the R-spine, failing to form canonically active

conformation of R- & C-spines. **(I & J)** ADP-bound p-IKK2 and P-IKK2 states are shown,

respectively. Both p-IKK2 and P-IKK2 states can accommodate ADP in their respective

clefts.

Figure 6: Freshly autophosphorylated IKK2 relays phosphates to IκBα. (A) Schematic

representation of the experiment performed to monitor the path traversed by phosphate(s)

from phospho-IKK2 to substrate by autoradiography detection method. **(B)** Purified

autophosphorylated (radiolabelled) IKK2 transfers its phosphate to IκBα substrate in

absence of any nucleotide, and transfer efficiency is enhanced upon addition of ADP or

ATP. **(C)** Schematic representation of the experiment performed to monitor the path

traversed by phosphate(s) from phospho-IKK2 to substrate by immunoblotting detection

method. **(D)** Elution profile of the size exclusion chromatography run of phospho-IKK2 to get

rid of excess unlabelled ATP performed in Superdex200 10/30 increase. **(E)** Purified

autophosphorylated (cold) IKK2 transfers its phosphate to IκBα substrate with enhanced

transfer efficiency upon addition of ADP or ATP monitored by immunoblotting using specific

antibodies as indicated in the figure.

Figure 7: Specificity and fidelity of phosphotransfer by IKK2 to IκBα. (**A**) AMP-PNP

can support efficient phosphotransfer. Also, phosphotransfer is observed in IκBα WT but not

in S32,36Ala double mutant. Domain organization and position of relevant S/T/Y residues of

IκBα are shown above the autoradiograph. **(B)** A detailed scheme of reactions showing

IKK2-activation and subsequent phosphorylation of the signal responsive S32,36 on IκBα by

phosphotransfer in presence of ADP. **(C)** A cartoon representation of the proposed model of

NF-κB activation by IKK2 employing phosphotransfer to IκBα.

Figure Supplement 1: (A) Autoradiograph of *in vitro* kinase assay reactions carried out with

FL IKK2 WT and K44M using GST (1-54) IκBα as the substrate as a function of time. **(B)**

Effect of Inhibitor VII on FL IKK2WT autophosphorylation and IκBα substrate

phosphorylation specifically on S32/36 at different inhibitor concentrations in a non-

radioactive *in vitro* kinase assay monitored using a monoclonal antibody that specifically

recognizes IκBα phosphorylated at those two serines. A scheme of kinase assays done to
check the effect of the inhibitors is shown in the box below the gel image. **(C)** Coomassie
stained SDS-PAGE showing purity of some of the IKK2 proteins used in this study.
113
114
Figure Supplement 2: (A) Effect of increasing concentration of urea on tyrosine
autophosphorylation of FL IKK2WT assessed in an *in vitro* kinase assay. **(B)** Radioactive *in*
vitro kinase assay performed with increasing concentration of urea up to 6M to check auto-
and substrate phosphorylation. **(C)** Inhibition of both substrate phosphorylation and tyrosine
autophosphorylation activities of IKK2 checked in presence of common kinase inhibitors
such as AMPPNP and Staurosporine, highly specific IKK2 inhibitors TPCA, Calbiochem
Inhibitor VII and MLN120B. **(D)** Mutation of AL-serines to non-phosphorylatable alanine (FL
IKK2 S177A,181A double mutant) debilitates its kinase activity towards S32,36 of IκBα
compared to WT or EE version of IKK2 as observed in an *in vitro* kinase assay.
124
125   **Figure Supplement 3: (A)** A close-up view of the active site in active form in the context of
126   the entire kinase domain of IKK2. Three tyrosines (Y169, Y188 and Y199) in immediate
127   vicinity of active sites are highlighted. Position of ATP is modelled based on the structure of
128   PKA. **(B)** Substrate bound PKA in its active form; phosphorylatable serine of the substrate
129   peptide (in magenta) is highlighted.
130
Figure Supplement 4: (A) A differential scanning calorimetry (DSC) thermogram showing a
striking enhancement in folding stability of IKK2 upon ATP-treatment. **(B)** Ribbon
representation of different phosphorylated versions of IKK2 are shown: UnP-IKK2 in golden
yellow, p-IKK2 in blue and P-IKK2 in copper red. **(C)** Trajectories of RMSF for these
structures are shown (left panel). For clarity, different regions of the KD are shown in the
right. **(D)** ATP-bound structure of p-IKK2 is shown. **(E)** Dynamic cross-correlation matrix
(DCCM) or contact map of each structure reveals phosphorylation-induced local as well as
allosteric changes.
139
140   **Figure Supplement 5: (A)** ESI-MS scan of the ADP used in this assay system showing
141   negligible trace, if any, of ATP at 50μM ADP concentration. **(B)** Regeneration of ATP, if any,
142   through microscopic reversibility was checked by TLC in a variety of conditions. γ-P^{32} was
143   used as the control/standard.

Figure 1

Figure 2

Figure 3

A

	ATP				
FL IKK2:	K44M				WT
Time (mins):	0	15	30	60	5

IB: (P)-Ser

IB: IKK2

C

	FL IκBα +ATP				
FL IKK2:	K44M				WT
Time (mins):	0	15	30	60	5

In vitro kinase assay

IB: IKK2

IB: (P)-IκBα

IB: IκBα

B

	ATP*					FL IκBα WT				
FL IKK2:	K44M			WT		K44M				WT
Time:	0	15	30	60	15	0	15	30	60	15	(mins)

In vitro Kinase Assay

Autoradiograph

Autophosphorylation

◄ FL IκBα WT

Coomassie

◄ FL IκBα WT

◄ FL IKK2 K44M/WT

D

	ATP*, 30mins							
FL IκBα:	WT				SS32,36AA			
FL IKK2:	WT		K44M		WT		K44M	
Inhibitor VII (µM):	0	10	0	10	0	10	0	10

In vitro Kinase Assay

Autophosphorylation

FL IκBα ►

Autoradiograph

His-FL IκBα AA ►
FL IκBα WT ►

Coomassie

FL IKK2

Figure 4

Figure 5

A Trajectory of energy

B Trajectory of RMSD

C p-IKK2 (no ATP): R & C Spines

D UnP-IKK2 P-IKK2 p-IKK2

E p-IKK2 with ATP

F p-IKK2 (with ATP): Spines

G P-IKK2:ATP

H P-IKK2 (with ATP): Spines

I p-IKK2:ADP

J P-IKK2:ADP

Figure 6

Figure 7

A

B

C