Title:

Dual-specific autophosphorylation of kinase IKK2 enables phosphorylation of

substrate IκBα without requiring ATP

Prateeka Borar[1], Tapan Biswas[2,#], Ankur Chaudhuri[3,#], Tom Huxford[4], Saikat Chakrabarti[3]

Gourisankar Ghosh[2] & Smarajit Polley[1,5,*]

[1]Department of Biological Sciences, Bose Institute, Kolkata, India; [2]Department of Chemistry

and Biochemistry, University of California San Diego, La Jolla, USA; [3]Structural Biology and

Bioinformatics Division, CSIR-Indian Institute of Chemical Biology, Kolkata, India;

[4]Department of Chemistry and Biochemistry, San Diego State University, San Diego, USA.

[5] Lead contact

[#] These two authors have contributed equally to this work.

*Correspondence to be addressed to: spolley@jcbose.ac.in, smarajit.polley@gmail.com

Abstract:

Rapid and high-fidelity phosphorylation of two serines (S32 and S36) of IκBα by a prototype S/T kinase IKK2 is critical for fruitful canonical NF-κB activation. Here, we report that IKK2 is a dual specificity kinase that autophosphorylates itself at tyrosine residues in *cis* in addition to its activation loop serines. Mutation of one such tyrosine, Y169, located in proximity to the active site, to phenylalanine, renders IKK2 inactive for phosphorylation of S32 of IκBα. Surprisingly, auto-phosphorylated IKK2 relayed phosphate group(s) to IκBα without ATP when ADP is present. We also observed that mutation of K44, an ATP-binding lysine conserved in all protein kinases, to methionine renders IKK2 inactive towards specific phosphorylation of S32 or S36 of IκBα, but not non-specific substrates. These observations highlight an unusual evolution of IKK2, in which autophosphorylation of tyrosine(s) in the activation loop and the invariant ATP-binding K44 residue define its signal-responsive substrate specificity ensuring the fidelity of NF-κB activation.

Introduction:

Inflammatory or pathogenic signals trigger precisely regulated induction of the transcription factor NF-κB in metazoans[1]. This process depends upon the upstream activation of the Inhibitor of κB Kinase-complex (IKK-complex), comprising two catalytic subunits, IKK1 (also known as IKKα) and IKK2 (or IKK2β), and an essential regulatory scaffolding protein, NEMO (IKKγ)[2–4]. IKK2 activity is maintained at a very low basal level in resting cells[5–8]. Upon activation, IKK2 phosphorylates the N-terminus of IκBα (Inhibitor of κBα) protein specifically at two Serine residues (S32 and S36) marking it for ubiquitinylation-mediated proteasomal degradation (Figure 1A). This liberates NF-κB to execute its gene expression program[6,9–11]. Mutation of both S32 and S36 to alanine residues converts IκBα into a non-phosphorylatable super-repressor of NF-κB[12,13]. Furthermore, IKK possesses exquisite specificity for these two serines, and it fails to phosphorylate IκBα when S32 and S36 are mutated even to another phosphorylatable residue, threonine[2].

A majority of the eukaryotic protein kinases are Ser/Thr/Tyr kinases (STYKs) that phosphorylate specific namesake residues of their cognate substrates[14,15]. STYKs could specifically phosphorylate S/T or Y or S/T and Y residues[16–18]. STYKs are rather promiscuous but can display a high substrate specificity in an appropriate context[19–21]. IKK2 is an essential kinase that ensures the activation of a particular signaling cascade - NF-κB, with high fidelity[22,23] despite its reported pleiotropy in other contexts. IKK2 is a multidomain protein consisting of a kinase domain (KD) followed by a Ubiquitin-like domain (ULD) and the scaffold dimerization domain (SDD) (Figure 1B). IKK2 toggles between inactive and active states based on the phosphorylation status of two serine residues (S177 and S181; S176 and S180 in the case of IKK1) within its activation loop (AL) in the KD[24–27]. Replacement of these two serines with phospho-mimetic glutamate residues (S177E, S181E; henceforth EE) renders the kinase constitutively active. IKK2 is reported to be primed through phosphorylation at S177 by the upstream kinase TAK1 that leads to auto-phosphorylation at S181, thus rendering the kinase fully active in cells[28]. However, oligomerization upon association with NEMO, in the presence of linear or Lys-63 linked poly-ubiquitin (Ub) chains or at a high concentration, also enables *trans* auto-phosphorylation at S177 and S181[26,29–31]. It was also shown that NEMO not only helps in the activation of the IKK-complex, but also ensures that the IKK-activity is well directed towards IκBα in the inhibitor bound transcriptionally inactive NF-κB (NF-κB: IκBα complex)[32]. Activated IKK2 phosphorylates IκBα specifically at residues S32/S36 and the C-terminal SDD of IKK2 (residue range 645-756) is necessary for this[33]. Other sites of IκBα, but not S32/S36, can be phosphorylated by various kinases[34,35]. The role of IκBα-phosphorylation at sites other than the signal responsive S32/S36 phosphorylation in NF-κB activation and signaling is elusive. These findings underscore the ability of IKK2 to function specifically in guaranteeing rapid NF-κB activation in canonical signaling. It is however, still not fully understood how IKK2 achieves its exquisite specificity towards S32/S36 of IκBα to induce NF-κB transcriptional activity.

An in-depth analysis of the catalytic features of IKK2 revealed that IKK2 undergoes

multisite autophosphorylation and can phosphorylate its own tyrosine residue(s) in addition to

its own serines or the serines of its substrate IκBα, i.e., IKK2 possesses dual specificity. This

phosphorylation at tyrosine residue(s) is autocatalytic. Furthermore, we observed that, after

auto-phosphorylation at tyrosine, IKK2 can phosphorylate IκBα in the absence of ATP, as long

as ADP is present in the reaction. Mass spectrometry of IKK2 identified that Y169 and Y188,

both present within proximity of the enzyme active site, can undergo auto-phosphorylation in

cis. This phosphorylation at tyrosine residues is dependent upon the prior activation of IKK2

via AL S177/S181 phosphorylation, and critical to phosphorylation of signal responsive

S32/S36 of IκBα. Mutation of Y169 of IKK2 to phenylalanine severely compromised

phosphorylation of IκBα at S32 (which has long been associated with pathological conditions

such as ectodermal dysplasia). We also found that mutation of the conserved ATP-interacting

residue K44 in IKK2 to methionine leads to a loss of Tyr-auto-phosphorylation activity of the

kinase and its ability to phosphorylate S32/S36 of IκBα. Interestingly, K44M retained non-

specific kinase activity. Overall, our results reveal unique, and likely transient, tyrosine-

autophosphorylated intermediates of IKK2 that appear to be critical in relaying phosphoryl

group(s) specifically to S32/S36 of IκBα in presence of ADP, thus ensuring fidelity of NF-κB

activation. Recent reports in other kinases indicate that multisite (auto)phosphorylation is a

major determinant in regulating function of respective kinases and regulation exerted by those

kinases. Our study adds another layer of regulation and critical nature of such

autophosphorylation that plays a significant biological role in metazoan.

Results:

IKK2 undergoes auto-phosphorylation at uncharacterized sites:

Signal-induced phosphorylation of AL Ser177/Ser181 marks the general activation of IKK2

(Figure 1B). Phosphorylation of several other serines located within the flexible C-terminal

region of IKK2, spanning amino acid 701 to the end, has also been reported (Figure 1B)[36].

Hyperphosphorylation of these C-terminal sites down-regulates IKK2 activity in cells. It was

also reported earlier that, ectopically overexpressed IKK2 is capable of autophosphorylating

these two AL serines in *trans* even in the absence of an activating physiological signal. Indeed,

recombinantly purified FL IKK2WT is capable of autophosphorylation *in vitro* when incubated

with γ^{32}P-ATP that increases with the time of incubation (Figure 1C). The spread-out of slower

migrating phospho-IKK2 band likely indicates phosphorylation of the protein at multiple sites.

We also observed that autophosphorylation of IKK2 occurred irrespective of the presence of

its cognate binding partner NEMO and *bonafide* substrate IκBα (Figure 1D). To confirm that

this phosphorylation of IKK2 was indeed autophosphorylation and not due to an unrelated

kinase copurified with IKK2, we generated a K44M mutant of full-length IKK2 where the ATP-

binding invariant lysine residue conserved in STYKs was mutated to methionine, that is

expected to greatly reduce or eliminate the kinase activity. An *in vitro* kinase assay using this

mutant along with the WT kinase using γ^{32}P-ATP as the phosphor-donor confirmed that the

IKK2 K44M lacked both IκBα (substrate) and autophosphorylation activities (Figure S1). That

these phosphorylations were indeed autophosphorylation was further supported by another *in*

vitro radioactive kinase assay performed in the presence or absence of an IKK-specific ATP-

competitive inhibitor, Calbiochem Inhibitor VII (Figure 1E).

Next, we used different versions of the IKK2 protein where all previously known

phosphorylation sites were lacking: Ser177 and 181 were mutated to phosphor-mimetic Glu

(EE; this version of IKK2 is known to be constitutively active [37]), and the C-terminal serine-rich

region was truncated to different extents. The shortest construct, Δ664EE that entirely lacked

the flexible C-terminus (670-756) was further subjected to limited proteolysis using trypsin to

eliminate residues, if any, from the flexible ends of the enzymes and this trypsin-treated IKK2

was purified prior to *in vitro* kinase assays. Each of these constructs was active in

phosphorylating itself and the substrate IκBα (Figure 1F). The occurrence of

autophosphorylation despite the absence of AL serines and serines within the C-terminus

strongly suggested the presence of hitherto uncharacterized sites of autophosphorylation in

IKK2.

Auto-phosphorylation of IKK2 reveals dual specificity: IKK2 is known as a *bonafide* S/T

kinase that undergoes autophosphorylation. In the previous section, we found that IKK2's

autophosphorylation activity might extend beyond the previously characterized site(s). We

asked if it is a dual specificity kinase. To test if Tyr residues can undergo auto-phosphorylation

in addition to Ser (and Thr), IKK2 incubated with Mg^{2+}-ATP for various time periods was

probed with antibodies that specifically recognize AL serines of IKK2 (Figure 2A, upper panel)

or phospho-tyrosine residues (Figure 2A, middle panel). Unexpectedly, we observed *de novo*

auto-phosphorylation of IKK2 on tyrosine residues along with the anticipated auto-

phosphorylation of AL serines. We tested another phospho-tyrosine specific antibody to

further substantiate this *de novo* tyrosine autophosphorylation of IKK2. It confirmed that

detection of *de novo* phosphotyrosine on IKK2 was independent of the source of antibody

used for the purpose (Figure 2B). The tyrosine auto-phosphorylation was abolished in the

presence of an IKK-specific inhibitor, Inhibitor VII (Figure 2C). A similar assay was performed

to monitor the phospho-tyrosine level in IKK2 as a function of urea concentration. This

revealed the detrimental effects of urea on tyrosine autophosphorylation of IKK2 at

concentrations beyond 1M (Figure S2A). A more sensitive assay with γ^{32}P-ATP as the

phospho-donor also indicated that urea concentrations beyond 1M inhibited both auto-

phosphorylation (pan-phosphorylation) and substrate (IκBα) phosphorylation (Figure S2B).

Furthermore, the IKK2-K44M mutant (kinase inactivating) did not display any tyrosine

autophosphorylation when incubated with ATP unlike WT IKK2 (Figure 2D). These results

confirm the novel tyrosine phosphorylation ability of IKK2, demonstrating its dual specificity to

both its own Ser and Tyr residues. We also observed that activity of IKK2 towards S32 and

S36 of IκBα and its own tyrosine residue(s) were affected similarly when the kinase assays

were performed in the presence of various kinase inhibitors, both general (AMPPNP and

Staurosporine) and IKK-specific (MLN-120B, TPCA, Inhibitor VII) (Figure S2C). These

observations confirm that canonical activity of IKK2 towards S32 and S36 of IκBα, and towards

its own tyrosine residues are mediated by the same active site pocket.

Next, we checked the requirement of the C-terminal <u>NE</u>MO <u>b</u>inding <u>d</u>omain (NBD) for

autocatalytic dual specificity. Constructs of various truncations (lacking NBD) were capable of

phosphorylating itself on tyrosine residue(s), and phosphorylating IκBα at signal-responsive

S32 and S36 (Figure 2E). Next, we assessed the autocatalysis at Ser and Tyr by IKK2 using

a constitutively active NBD-truncated version of IKK2 (IKK2$^{\Delta664}$EE). As anticipated, the protein

was recognized by monoclonal phospho-serine antibody with or without fresh ATP-treatment;

however, phospho-tyrosine antibody recognized it strongly only upon fresh ATP-treatment

(Figure 2F).

Next, we asked if phosphorylation of AL S177/S181 was necessary for this dual

specificity of IKK2. To this end, we purified the S177A, S181A (AA; non-phosphorylatable AL)

mutant of IKK2, and checked its tyrosine autophosphorylation activity. We found that the IKK2

AA mutant lacked tyrosine autophosphorylation activity (Figure 2G) in an *in vitro* kinase assay.

It suggests that AL S177/S181 serine phosphorylation is essential in conducting dual

specificity in IKK2. Indeed, the IKK2 AA mutant was incapable of phosphorylating IκBα as well

(Figure S2D) in an *in vitro* kinase assay.

To investigate if tyrosine phosphorylation occurs in *cis* (within the same kinase

molecule, intramolecular) or *trans* (by a different kinase molecule, intermolecular), we

incubated full-length IKK2-K44M with ATP in the absence or presence of a constitutively active

IKK2$^{\Delta743}$-EE lacking the extreme C-terminal end. Prior to this assay, IKK2$^{\Delta743}$-EE was pre-

treated with ATP and purified through a Superdex200 size-exclusion column to remove

unutilized ATP. The absence of a phospho-tyrosine signal on IKK2^{FL}-K44M irrespective of the

presence of IKK2$^{\Delta743}$-EE, and the presence of the phospho-tyrosine signal on the IKK2$^{\Delta743}$-EE

strongly suggested that tyrosine auto-phosphorylation occurs in *cis*, and not in *trans* (Figure

2H, upper panel). In contrast, strong phosphorylation of AL S177/S181 on the IKK2^{FL}-K44M

by IKK2$^{\Delta743}$-EE suggests that AL serine auto-phosphorylation of IKK2 could occur in *trans*,

agreeing with our previous findings (Figure 2H, middle panel) [26].

Tyrosine phosphorylation is critically important for IKK2 activity: Next, we performed

LC/MS/MS analyses to identify tyrosine residues that are autophosphorylated *de novo* on

IKK2 upon the addition of fresh ATP. We identified at least two residues, Y169 and Y188, to

be phosphorylated upon fresh ATP treatment[38,39] (Figure 3A, and Figure S3A & S3B).

Interestingly, both are located within the AL (Figure 3B); Y169 is the DFG+1 (DLG in case of

IKK2 and IKK1) residue while Y188 (equivalent to Y204 of PKA) is located at the end of AL

prior to the conserved APE motif (Figure 3A). It was previously established that the DFG+1

position is crucial for defining the substrate specificity of a kinase[40]. Y188 counterpart Y204 of

PKA is reported to allosterically modulate PKA activity by altering its conformational

entropy[41,42].

To test the effect of phosphorylation at Y169 and Y188, we generated Y169F and

Y188F mutants of IKK2. Radioactive *in vitro* kinase assays using recombinant purified IKK2^{FL}-

Y188F and IKK2^{FL}-Y169F mutants revealed a drastic reduction in the activity of IKK2 (Figure

3C). Since the DFG+1 residue in kinases (Y169 in IKK2) is known to dictate its specificity, we

created different mutants of the IκBα substrate where S32 and S36 were singly and

independently altered to either alanine (non-phosphorylatable) or glutamic acid

(phosphomimetic) residues. Mutation of S36 to either Ala or Glu drastically reduced the

phosphorylation signal of IκBα while that for the S32A/E mutants were minimally affected

(Figure 3D). This suggests that the presence of Tyr (and possibly its phosphorylation) is

critically important for signal-responsive phosphorylation of IκBα. Since pTyr mimic residue is

not available, testing the roles of modified tyrosine residues is not feasible.

Next, we tested the consequence of Y169F and Y188F mutants on the pro-

inflammatory signal-induced IKK activity in cells. We reconstituted IKK2 knock-out mouse

embryonic fibroblast (*ikk2$^{-/-}$* MEF) cells with WT, Y169F and Y188F mutants of IKK2, and

measured signal-induced activation of IKK2 upon TNF-α treatment. IKK2 activity is

significantly reduced in Y169F and nearly abolished in Y188F (Figure 3E**)**. Together, these

results suggest communication between the phosphorylation of Y169 of IKK2 and that of S32

of IκBα.

Structural analyses of IKK2 autophosphorylation: We compared available structural

models of human IKK2 (PDB ID: 4KIK, 4E3C) to investigate whether tyrosine phosphorylation

within the activation loop could be supported based on its structural features[23],[24]. We observed

distinct conformations of the activation loop containing S177/S181. The structure of IKK2

dimer is asymmetric in that the activation loop of one monomer is largely disordered while that

of the other monomer is mostly ordered (PDB ID: 4KIK)[23]. In the active monomers S177 and

S181 are clearly observed. In these active state conformers, the position and conformation of

Tyr169 appears to be well-poised for accepting a phosphate from ATP (Figure S4A). A

superposition of IKK2 with a pseudo-substrate-bound PKA shows that the hydroxyl of Tyr169

in IKK2 projects toward the γ-phosphate of ATP similarly to the serine hydroxyl on the PKA

inhibitor pseudo-substrate[40],[41] (Figure S4A & B).

To get a more comprehensive picture, it is necessary to analyze at least three phosphorylated

states of IKK2: unphosphorylated (UnP-IKK2), phosphorylated at S177 and S181 (p-IKK2),

and phosphorylated at Y169, S177 and S181 (P-IKK2). We took advantage of computational

approaches using molecular dynamic (MD) simulations and flexible molecular docking (see

methods).[43–50] MD simulation was performed for 200ns for each of these three states of IKK2,

unphosphorylated and phosphorylated. Phosphorylation at S177, S181 increased the stability

of the kinase, which was further increased by the phosphorylation at Y169 as evidenced by a

gradual decrease in the total energy (Figure 4A). A striking enhancement in stability of IKK2

upon this auto-phosphorylation is reflected in the change of Tm from ~ 40 °C to ~ 50 °C in

differential scanning calorimetry (DSC) scans in the presence of ADP and ATP (Figure S4C).

This behavior supports our observation in MD simulation, and tallies with increased solubility

of IKK2 observed after auto-phosphorylation that was essential for its successful

crystallization.[26]

Interestingly, RMSD values increased with more phosphorylation (Figure 4B) despite

phosphorylation-induced stabilization, P-IKK2 showed the highest RMSD followed by p-IKK2

and UnP-IKK2. We found that, phosphorylation at S177, S181 and S177, S181, Y169 induced

distinct structural alteration in the KD when compared with the unphosphorylated version

(Figure 4C, models shown separately in Figure S4D; see Movies S1, S2 & S3 for better clarity

of the structural changes involved). Careful analyses of the trajectory of RMSF values for

these three states revealed that, different regions of the kinase domain had different RMSF

values in different phosphorylated states (Figure S4E). We observed that, while the glycine-

rich loop and the αC-helix in p-IKK2 were situated in a manner reminiscent of an active kinase,

those in UnP and P-IKK2 were placed very differently. In fact, helix αC was found to be

distorted in P-IKK2 (Figure 4C and S4D), that is unlikely to support canonical phosphotransfer

from ATP. Activation segment/loop (AL) in these three differently phosphorylated states

adopted conformations distinct from each other (Movies S1, S2 & S3). AL in the UnP model

was displayed out and moved closer to the tip of the C-lobe, whereas AL in p-IKK2 and P-

IKK2 moved inwards and adopted conformations closer to the N-lobe. The formation of the

proper R- and C-spines in p-IKK2 confirmed its active form (Figure 4D). p-IKK2 KD also

exhibited other canonical features of an active kinase, viz., a salt bridge between K44 and E61

(K72, E91 in PKA, respectively), 'DFG in' (DLG here) conformation. Next, we docked ATP

onto the p-IKK2 structure (ΔG = -10.64 kcal/mol) using a flexible docking protocol (see

methods). The ATP and its phosphate groups exhibited a pose (Figure S4F) very similar to

that in the PKA structure (PDB ID: 1ATP), and the terminal phosphate was in close proximity

to Y169-OH making *Cis*-autophosphorylation of Y169 highly plausible (Figure 4E). Further

analyses of the ATP-bound p-IKK2 structure confirm that presence of ATP helped the R and

C-spine form the desired continuum of an active kinase (Figure 4F). Superposition of the ATP

from the p-IKK2 on P-IKK2 indicated that the narrowing of ATP-binding pocket in P-IKK2 may

lead to rejection of ATP emanating from severe clashes between the glycine-rich loop and

ATP (Figure 4G). Furthermore, the R-spine in P-IKK2 was not continuous (Figure 4H), wherein

M65 (L95 in PKA) of the four conserved residues were located at a position far away from the

rest of three residues that constitute the R-spine, and from the glycine rich loop.

In addition, dynamic cross-correlation matrix (DCCM) or contact map of each structure

suggests that specific phosphorylation events render the kinase display distinct allosteric

changes at locations far from the phosphorylation sites themselves (Figure S4G).

ATP binding to p-IKK2 leads to autophosphorylation at Y169 generating P-IKK2 in an ADP

bound state. Interestingly, the superposition of ADP (from the docked complex of p-IKK2 and

ADP, ΔG = -9.88 kCal/mol, Figure 4I) onto the P-IKK2 structure confirmed that while P-IKK2

was unable to accommodate ATP in a manner similar to that in other known experimentally

derived kinase structures, it could comfortably accommodate ADP into the ATP-binding cleft

(Figure 4J). It may be noted that, p-IKK2 has higher affinity for ATP than ADP as indicated by

their respective binding energy values. We speculate that ADP-bound P-IKK2 is an

intermediate step where presence of as many phosphate groups in the AL and ADP in such

a close proximity persuades the kinase to adopt a conformation that can withstand this high

density of negative charges albeit risking the integrity of an active fold, and making some of

those *de novo* phosphate groups transient in nature to return back to the p-IKK2 state to

readily bind to ATP.

These structural observations suggest that it is possible for the tyrosine residues within the

flexible activation segment to undergo auto-phosphorylation in *cis* and possibly render the

kinase with a novel phospho-transfer activity without requiring ATP.

Freshly auto-phosphorylated IKK2 phosphorylates IκBα in absence of ATP: We

observed phosphorylation of tyrosine residue(s) by WB or mass spectrometry only upon fresh

ATP-treatment, indicating that tyrosines of purified IKK2 loses its phosphate group [51]. It may

be noted that, we did not observe any density of phosphates on any of these residues in the

human IKK2 crystal structure derived from ATP-treated protein sample [26]. This suggested an

intriguing possibility that the phosphate linked to Tyr residue(s) may serve as a temporary

phosphate-sink and could eventually be transferred to substrate IκBα. To test this possibility,

we first allowed purified IKK2 to undergo auto-phosphorylation by incubating the excess kinase with limiting amounts of radiolabeled γ-^{32}P-ATP followed by two passes through desalting spin columns (40 kDa MW cut-off) to remove unreacted ATP. Nucleotide-free auto-phosphorylated IKK2 (P-IKK2) was incubated with IκBα either in absence of nucleotide or with newly added unlabeled nucleotides, and phosphorylation status of the substrate was subsequently monitored by autoradiography after resolving the reaction mixture on an SDS-PAGE (Figure 5A). Transfer of radiolabeled phosphate to IκBα was observed and the transfer was significantly enhanced in the presence of cold ADP (Figure 5B). Phosphorylation of IκBα did not significantly decrease in the presence of a high concentration of cold ATP. That addition of excess cold ATP did not reduce the extent of IκBα phosphorylation strongly suggests the transfer of ^{32}P-phosphate from ^{32}P-IKK2 to IκBα (Figure 5B) by a relay mechanism with or without involving ADP.

We undertook a different strategy to further confirm transfer of phosphate from the kinase (and not ATP) to the substrate. We auto-phosphorylated IKK2^{FL}-WT with excess cold ATP and removed the unreacted ATP from the reaction mixture using size-exclusion chromatography on a Superdex200 column (Figure 5C). P-IKK2 eluted at fractions corresponding to MW between 670 and 158 kDa, i.e., much larger than free ATP (Figure 5D). This P-IKK2 protein was incubated with FL IκBα either in presence or absence of ADP for different time periods. A similar reaction in presence of ATP instead of ADP was performed as a positive control. Phosphorylation of S32 and S36 was efficient in presence of 10 and 50 μM ADP but absent when ADP was absent in the reaction mixture. Phosphorylation of IκBα in the presence of 50 μM ATP was significantly more robust as anticipated (Figure 5E). The minor detection of IκBα phosphorylation in absence of ADP in the radioactive assay and lack of it in the cold immunoblotting assay (compare lanes marked with vertical arrows in Figs. 5B and 5F) may indicate higher sensitivity of the radioactive assay. The need of ADP in phosphotransfer prompted us to consider whether the ADP was contaminated with a trace

amount of ATP. ESI-MS analysis of 50 μM ADP did not detect any peak corresponding to ATP

(Figure S5A).

Phosphotransfer from P-IKK2 is critical to fidelity and specificity of IκBα

phosphorylation:

We next tested if phosphotransfer is restricted only to the N-terminal signal responsive

S32/S36 or to serines/threonines of the PEST domain as well. Radioactive kinase assay as

described earlier was performed with full-length WT and mutant S32A, S36A (AA) IκBα as the

phosphate recipient. Phosphate group(s) of P-IKK2 was not transferred to full-length IκBα AA

which retains the serines and threonines within its PEST domain (Figure 6A), indicating

phosphotransfer only to the N-terminal serines of IκBα. We also tested the effect of AMPPNP,

an unhydrolyzable analogue of ATP, on the phosphotransfer, and found no inhibitory effect

irrespective of the presence of ADP. This suggests adenine nucleotides poising the kinase for

phosphotransfer (Figure 6A).

During our investigation, we noted unusual behavior in the IKK2-K44M mutant. IKK2-

K44M phosphorylated full-length WT IκBα robustly but with much less efficiency than WT

IKK2. IKK2-K44M also showed a minor auto-phosphorylation activity compared to WT IKK2

in the absence of substrate (Figure 6B). We performed a similar kinase assay using full-length

WT IκBα where cold ATP was used as the phosphate donor, and phosphorylation status of

IκBα at S32, S36 was monitored by immunoblotting. It revealed that IKK2-K44M is unable to

phosphorylate S32/S36 (Figure 6C). Since in radioactive kinase assay, we observed pan-

phosphorylation of the substrate (Figure 6B), it possibly reflects phosphorylation of the PEST

region of FL IκBα. We additionally compared IKK2-K44M with WT IKK2 using radioactive

kinase assay with both FL IκBα and FL IκBαAA substrates in the absence or presence of

Inhibitor VII. As anticipated, FL IκBα was phosphorylated much more efficiently than FL

IκBαAA by WT IKK2, and phosphorylation was practically abolished in the presence of 10 μM

Inhibitor VII (Figure 6D). In contrast, IKK2-K44M phosphorylated FL IκBαAA more efficiently

than FL IκBα, and reduction in the presence of inhibitor VII was strikingly less (Figure 6D). It

indicates that IKK2-K44M, despite retaining non-specific kinase activity, lost phosphorylation

activity towards S32, S36 of the *bona fide* substrate IκBα. It is noteworthy that an equivalent

K to M mutation in Erk2 also did not abolish the basal activity, but rather retained ~5 % of it[52].

Discussion:

Phosphorylation of substrates by protein kinases is regulated in various ways[20,21]. IKK2, a

kinase central to inflammation, phosphorylates serine residues of its primary substrate IκBα

with γ-phosphate of ATP. Here, we report multiple intriguing properties of IKK2 that provide

specificity to its substrate phosphorylation. These include: 1) dual specificity that entails

autophosphorylation of tyrosine residues of IKK2 in *cis*, in addition to known auto-

phosphorylation at AL serines; 2) relay of phosphate to substrate IκBα from auto-

phosphorylated IKK2 (P-IKK2) likely through ADP; and 3) loss of specificity of the kinase due

to disruption of a universally conserved salt bridge (mediated by K44).

STYK family of eukaryotic protein kinases transfer γ-phosphate of ATP directly to

substrate, in general to S, T and Y residues[53–57]. Histidine kinases (in prokaryotes and in some

lower eukaryotes) however shuttles the γ-phosphate to Asp residue on a response regulator

(RR) substrate through a phospho-His intermediate[52,58–62]. Phospho-His functions as a

phosphoenzyme intermediate in several metabolic enzymes as well[61]. Our present analyses

of eukaryotic protein kinase IKK2 revealed its surprising property in which IKK2

phosphorylated at AL-serines (S177/S181) could be further auto-phosphorylated by ATP at

two (Y169 & Y188) or more tyrosine residues (and perhaps other residues) yielding another

form (P-IKK2) (Figure 6E). This marks IKK2 as an autocatalytic dual-specificity kinase rather

than a prototype S/T kinase. Interestingly, while auto-phosphorylation at AL-serines (S177

and S181) occurs in *trans*[63], auto-phosphorylation of tyrosine residues occurs in *cis*. This

multisite phosphorylated form of IKK2 (P-IKK2) could transfer some of its phosphate(s) to

substrate IκBα in presence of ADP without requiring ATP, hinting at an unusual property of

IKK2 acting possibly as a phosphate-sink. It is possible that phospho-tyrosines residues of

IKK2 act as this sink. The Y169 located at the DFG+1 (DLG in IKK2) position is unique only

to IKK1 and IKK2, and is absent even in close mammalian homologues such as IKKε and

TBK1. Interestingly, neither IKKε nor TBK1 can phosphorylate IκBα at S32 and S36, *in vitro*

or *in vivo*[64,65]. A recent report also indicated the role of DFG+1 residue in differentiating Ser

versus Thr residues[40]. These observations hinted to an involvement of Y169 in substrate

selectivity. Indeed, our experimental data suggests a pivotal role of Y169 in phosphorylating

S32 of IκBα both *in vitro* and *ex vivo* (Figure 3D-E). It is worthwhile to note that, a un-

phosphorylatable mutation at S32 in IκBα to Ile (S32I) is associated with ectodermal dysplasia

and T-cell immunodeficiency in patients[66,67]. More interestingly, the placement of this residue

in the kinase active site makes it amenable to a phosphorylation-dephosphorylation cycle,

suggesting its possible role both in phosphotransfer and substrate specificity. Mutation of

Y188, a conserved tyrosine, to non-phosphorylatable phenylalanine abolished the kinase

activity of IKK2 *in vitro* as well as in MEFs upon TNF-treatment (Figure 3C & 3E). The

placement of Y188 indicates that its mutation will be detrimental to the structural fold of kinase,

thus it is unclear if or how Y188 plays a role in the phosphotransfer. Since phosphotransfer to

S32, S36 of IκBα is observed with auto-phosphorylated IKK2 EE, transferred phosphates are

not from S177 or S181. Kinase domain autophosphorylation is known to regulate function of

other kinases as well, e.g., autophosphorylation at T220 greatly influenced both the activity

and substrate specificity of CK1[21].

The strength of input signal, level of IκBα degradation, and NF-κB activation correlate

with the amplitude, duration, and kinetics of IKK activity[68,69]. This digital (all-or-none) activation

profile of NF-κB appears to be due to rapid activation and inactivation of IKK2. IKK targets

multiple substrates *in vivo*; however, the rapid activation of NF-κB is regulated primarily by

IκBα. How IKK specifically selects IκBα for phosphorylation is unclear, although NEMO is

reported to direct the kinase activity of IKK2 specifically towards IκBα[32]. We observe that P-

IKK2 could transfer phosphate(s) specifically to the S32/S36 of IκBα, and not to other sites

(such as the PEST sequence of IκBα) (Figure 5B, 5E and 6A). We surmise if transiency of

multisite phosphorylated IKK2 (P-IKK2) underlie this rapid activation-inactivation events

during signal response. It is possible that regulatory system of IKK2 is somewhat analogous

the two-component histidine kinase-effector systems in prokaryotes that show a threshold

activation response[70–72].

All three tyrosines in AL of IKK2 are previously reported to undergo signal-induced

phosphorylation in cells. These modifications were proposed to be mediated by a tyrosine

kinase although there is no evidence of a specific tyrosine kinase[39,73,74]. Several members of

the Ser/Thr kinase family undergo tyrosine phosphorylation and these kinases employ diverse

mechanisms for auto-phosphorylation[75–80].

The phosphotransfer from P-IKK2 to IκBα appears inefficient as no apparent decrease

in phospho-label in P-IKK2 is observed post transfer (Figure 5B). This is in contrast to HKs

where a single phosphate from the phosphorylated HK is stoichiometrically transferred to the

cognate RR. There are also other prominent differences, such as phosphorylation on IKK2

occurs at multiple sites unlike at single site for the HKs, and not every phosphorylated residue

in P-IKK2 expected to transfer its phosphate to IκBα. It is likely that, there are more than one

type of phosphorylated residues in P-IKK2 with respect to their stability wherein only a fraction

of the total pool can be relayed. Another notable difference between phosphorelay in IKK2

from that in HK is IKK2's dependence on ADP (Figure 5B & 5E). It was also observed that,

the generic kinase inhibitor AMPPNP (unhydrolyzable ATP analogue) failed to inhibit

phosphor-transfer from IKK2 (Figure 6A) and IκBα phosphorylation (Figure S2C). Taken

together, we surmise that presence of ADP or AMP-PNP help IKK2 attain a conformational

state poised for phosphorylating S32/36 of IκBα even in absence of ATP. Similar

phenomenon was reported for IRE1, where ADP was found to be a more potent stimulator of

its ribonuclease activity and that AMPPNP could also potentiate this activity in IRE1[81,82].The

list of Ser/Thr kinase family proteins that undergo tyrosine phosphorylation is growing that

employ diverse mechanisms for auto-phosphorylation[75–80]. The most notable example of this

dual specificity is the DYRK. However, the role of Tyr-phosphorylation in DYRK is not related

to that in IKK2, nor does it render DYRKs capable of relaying phosphates to the substrates. It

will be interesting to see whether phosphotransfer by relay (with or without ADP) occurs in any

other kinase-substrate system other than IKK2: IκBα system described here.

Our study reveals tyrosine auto-phosphorylation in IKK2 and indicates that

consequences of Ser and Tyr autophosphorylations in dual-specific IKK2 are distinct.

Moreover, positions of the critical tyrosine residues are unique, such that some that are

amenable to *cis* auto-phosphorylation are essential for substrate phosphorylation. Most

surprisingly, the tyrosine-linked phosphate groups can be transferred by a relay mechanism

to specific serines in IκBα, an event reminiscing phospho-relay in two-component systems

prevalent in prokaryotes and some lower eukaryotes. Our study also reveals the role of K44

of IKK2 in auto-phosphorylation and thus in specific phosphorylation of S32, S36 of IκBα,

although it does not affect its non-specific kinase activity. It requires further investigation to

determine if a structural state of IKK2, perhaps through a different modification or through

binding to another protein, allow it to phosphorylate substrates relatively non-specifically.

In all, it is intriguing how the gateway to the activation of NF-κB is regulated through

unique phosphorylation events in IKK2 (Figure 6F). It appears that a multi-layered fail-safe

mechanism is devised to tightly control NF-κB activation. Firstly, in absence of upstream

signals IKK-complex is unable to phosphorylate Ser 32,36 of IκBα efficiently thereby keeping

check on the aberrant and untimely activation of NF-κB. Upon proinflammatory cues, cells

need to activate NF-κB immediately and efficiently, i.e., IKK needs to be activated and

phosphorylate Ser 32,36 of IκBα. NEMO warrants IKK-activation and ensures that the IKK-

complex specifically chooses IκBα from a large pool of substrates of IKK2. IKK2 is designed

in such a manner that it phosphorylates itself at AL-tyrosines when activated, such that

phosphate group(s) can be relayed directly or through ADP to Ser32,36 of IκBα with great

fidelity, thus leaving little chance of a misconstrued signaling event thereby confirming NF-κB

activation. Our discovery might also present the beginning of a new aspect in eukaryotic cell-

signaling by EPKs and provide a foundation for mechanistic studies of an intriguing phospho-

transfer reaction that helps accomplish the desired specificity. To the best of our knowledge,

it is the first report to strongly suggest that a metazoan EPK harbors HK-like phosphotransfer

activity through relay mechanism to its substrate.

Methods summary:

Protein purification, auto-phosphorylation assay, immunoprecipitation coupled kinase assay,

phosphotransfer assay and other experiments have been described in details in the

supplementary document.

Acknowledgement: We thank Joe Adams and Simpson Joseph (UCSD), Tony Hunter (Salk

Institute) and Alex Hoffmann (UCLA) for commenting on the work. Authors also thank

members of Polley lab (BI) and Ghosh Lab (UCSD) for their continuous support. We also

thank Majid Ghassemian (Biomolecular Mass spectrometry facility, UCSD) and Souvik Roy

(Mass spectrometry facility, BI) for the mass spectrometry experiments and analyses. This

work was supported by an Intermediate Fellowship from the DBT Wellcome Trust India

Alliance (IA/I/15/1/501852) & intramural funding from Bose Institute to SP, NIH grant to GG

(AI163327) and American Cancer Society grant RSG-08-287-01-GMC to TH. Biochemistry

research at SDSU is supported in part by the California Metabolic Research Foundation. PB

was supported by the graduate research fellowship from Bose Institute. SC acknowledges

CSIR-IICB for infrastructure support. AC acknowledges ICMR for post-doctoral Research

Associateship [BMI/11(55)2022].

Author contribution: SP, GG and TB designed experiments; PB, SP, TB & GG performed

experiments. Computational analyses were performed by AC & SC. SP, GG, TB, PB, TH, SC

and AC analyzed the results. SP wrote the manuscript with the help of PB, TB and AC in

preparing figures and images. GG, TB, PB and TH edited/modified the manuscript. All authors

commented on the manuscript.

Declarations:

Competing interests: The authors declare no competing financial interests.

Generative AI: Authors did not use any generative AI tool in preparing this manuscript.

Figure 1

Figure 1: Autophosphorylation of IKK2 at hitherto uncharacterized sites. A, NEMO is known to remain bound to the IKK-complex pre- and post-stimulation, helps in activation as well as substrate recognition to channelize IKK-activity towards IκBα. However, it is not clear how IKK specifically phosphorylates the Ser32&36 of IκBα. **B,** Domain organisation of FL IKK2 depicting functional kinase domain (KD), ubiquitin-like domain (ULD), and scaffold dimerization domain (SDD) based on the X-ray structures. Known sites of Ser phosphorylation and NEMO interacting regions at the extreme C-termini are highlighted. Serine and tyrosine residues in the activation loop are shown, mutation of which to Glu renders IKK2 constitutively active. The conserved ATP-interacting Lys (at position 44) is also shown. **C,** Autophosphorylation of FL IKK2WT upon incubation with radioactive ATP for different time periods in an *in vitro* kinase assay. **D,** *In vitro* radioactive kinase assays performed to check the effect of NEMO on autophosphorylation of FL IKK2WT (left panel), and that of NEMO and IκBα on autophosphorylation and substrate phosphorylation (right panel) activity of FL IKK2WT. **E,** Effect of Inhibitor VII on FL IKK2WT autophosphorylation and IκBα substrate phosphorylation at different inhibitor concentrations in a radioactive *in vitro* kinase assay. **F,** Radioactive kinase assay conducted to check auto- and substrate phosphorylation of full length and deletion constructs of constitutively active form of IKK2 harbouring phosphomimetic Ser177Glu and Ser181Glu mutations.

Figure 2

Figure 2: IKK2 possesses autocatalytic dual specificity. A, *In vitro* kinase assay, followed by immunoblotting with FL IKK2WT using antibodies against phosphor-Ser (177/181) and phospho-Tyr. **B,** A different commercial source of phospho-Tyr antibody was used to detect the same. **C,** Effect of Inhibitor VII on tyrosine autophosphorylation of FL IKK2WT. **D,** Immunoblotting performed with phospho-Tyr antibody to check autophosphorylation of IKK2 mutant K44M at different time points compared to that of IKK2 WT. **E,** Tyrosine autophosphorylation and substrate phosphorylation of GST (1-54) IκBα in presence of full length and deletion mutants of IKK2 harbouring phosphomimetic Ser177Glu and Ser181Glu mutations. **F,** *De novo* Auto-phosphorylation of IKK2 on tyrosine monitored by phospho-Ser and phospho-Tyr specific monoclonal antibodies. **G,** Autophosphorylation and subsequent immunoblot analysis of FLIKK2 mutant S177A, S181A using antibodies against phospho-Ser (177/181) and phospho-Tyr. **H,** *In vitro* kinase assay followed by immunoblotting done showing active IKK2 cannot phosphorylate tyrosines of an inactive (K44M) IKK2 mutant; however, it can efficiently phosphorylate AL serines.

Figure 3: Tyrosine autophosphorylation is critical for IKK2 activity. A, Position of canonically important residues in the AL (purple ribbon) of IKK2-KD are shown in the context of its structure (PDB ID 4E3C; surface representation; KD is shown light green with respect to the SDD in teal). Autophosphorylated tyrosine residues identified in this study are shown in green. **B,** Alignment of the activation segment sequences of different IKK family kinases. Conservation of Tyr at the DFG+1 (DLG in case of IKK1 and IKK2) is retained only in a stress response related plant kinase SnRK2, but not in other IKK homologues or any known dual specificity kinases, e.g., DYRK and GSK3β (both contain Ser at that position). Tyr188 (204 in PKA) is conserved. **C,** Radioactive *in vitro* kinase assay performed with IKK2 Y169F and Y188F mutants using GST (1-54) IκBα WT and AA as substrates. **D,** Radioactive *in vitro* kinase assay performed with IKK2 Y169F mutant with a range of GST (1-54) IκBα mutants where the substrate phosphorylation signal is abolished in case of S36 A/E IκBα. **E,** Severe reduction in TNFα-induced IKK activity in *ikk2-/-* MEF-3T3 cells reconstituted with mutant (Y169F and Y188F) IKK2s compared to wild-type. Kinase assay was performed with IKK immunoprecipitated (IP-ed) with anti-NEMO antibody from whole cell extract.

Figure 4

Figure 4: Structural analyses of IKK2 autophosphorylation. A, Molecular dynamics (MD) simulations of all three differently phosphorylated states of IKK2 was performed. 200ns trajectory of energy for each state is shown in golden yellow, blue and copper red representing UnP-IKK2, p-IKK2 and P-IKK2, respectively. Same color coding has been followed throughout this figure and the corresponding supplementary figure. **B,** 200ns trajectory of RMSD for each state is shown. **C,** Residues forming canonical R- (labelled in blue) & C-spines (labelled in dark gray) representative of an active conformation in p-IKK2 are shown. **D,** Superposition of structures representing the differently phosphorylated states post 200ns MD simulation. Relevant sections, e.g., activation loop, αC-helix, Gly-rich loop areas are highlighted in the right panel. **E,** ATP-bound p-IKK2 structure is depicted. In the right panel, relevant residues are highlighted. Y169 is found in a position poised for *Cis*-autophosphorylation. **F,** ATP completes the desired continuum of R- & C-spine observed in active kinase conformations. **G,** P-IKK2 cannot accommodate ATP in its cleft, and αC-helix is displaced. **H,** M is moved away from the R-spine, failing to form canonically active conformation of R- & C-spines. **I & J,** ADP-bound p-IKK2 and P-IKK2 states are shown, respectively. Both p-IKK2 and P-IKK2 states can accommodate ADP in their respective clefts.

Figure 5

Figure 5: Freshly autophosphorylated IKK2 relays phosphates to IκBα. A, Schematic representation of the experiment performed to monitor the path traversed by phosphate(s) from p-IKK2 to substrate by autoradiography detection method. **B,** Purified autophosphorylated (radiolabelled) IKK2 transfers its phosphate to IκBα substrate in absence of any nucleotide, and transfer efficiency is enhanced upon addition of ADP or ATP. **C,** Schematic representation of the experiment performed to monitor the path traversed by phosphate(s) from p-IKK2 to substrate by immunoblotting detection method. **D,** Elution profile of the size exclusion chromatography run of p-IKK2 to get rid of excess unlabelled ATP performed in Superdex200 10/30 increase. **E,** Purified autophosphorylated (cold) IKK2 transfers its phosphate to IκBα substrate with enhanced transfer efficiency upon addition of ADP or ATP.

Figure 6

Figure 6: Specificity and fidelity of phosphorelay by IKK2 to IκBα. A, AMP-PNP can support efficient phosphorelay. Also, phospho-relay is observed in IκBα WT but not in S32,36Ala double mutant. Domain organization and position of relevant S/T/Y residues of IκBα are shown above the autoradiograph. **B,** Comparison of substrate and autophosphorylation activities of FL IKK2 WT and K44M mutant in an *in vitro* radioactive kinase assay performed with or without FL IκBα WT as the substrate. **C,** Residue level specificity of phosphorylation by the FL IKK2 K44M was analysed using an antibody that specifically recognizes phosphor-S32/36 of IκBα. **D,** Differential sensitivity of specific vs. non-specific phosphorylation by IKK2 as assessed by the effect of Inhibitor VII on FL IKK2 WT and IKK2 K44M using IκBα WT and AA as substrates. **E,** A detailed scheme of reactions showing IKK2-activation and subsequent phosphorylation of the signal responsive S32,36 on IκBα by phosphorelay in presence of ADP. **F,** A cartoon representation of the proposed model of NF-κB activation by IKK2 employing phosphorelay to IκBα.

References:

1. Hoffmann, A., and Baltimore, D. (2006). Circuitry of nuclear factor kappaB signaling. Immunol Rev *210*, 171–186. 10.1111/j.0105-2896.2006.00375.x.

2. DiDonato, J.A., Hayakawa, M., Rothwarf, D.M., Zandi, E., and Karin, M. (1997). A cytokine-responsive IκB kinase that activates the transcription factor NF-κB. Nature *388*, 548–554. 10.1038/41493.

3. Zandi, E., Rothwarf, D.M., Delhase, M., Hayakawa, M., and Karin, M. (1997). The IkappaB kinase complex (IKK) contains two kinase subunits, IKKalpha and IKKbeta, necessary for IkappaB phosphorylation and NF-kappaB activation. Cell *91*, 243–252. 10.1016/s0092-8674(00)80406-7.

4. Rothwarf, D.M., Zandi, E., Natoli, G., and Karin, M. (1998). IKK-gamma is an essential regulatory subunit of the IkappaB kinase complex. Nature *395*, 297–300. 10.1038/26261.

5. Hacker, H., and Karin, M. (2006). Regulation and function of IKK and IKK-related kinases. Sci STKE *2006*, re13. 10.1126/stke.3572006re13.

6. Hinz, M., and Scheidereit, C. (2014). The IκB kinase complex in NF-κB regulation and beyond. EMBO Rep. *15*, 46–61. 10.1002/embr.201337983.

7. Ghosh, S., and Karin, M. (2002). Missing pieces in the NF-kappaB puzzle. Cell *109 Suppl*, S81-96.

8. Liu, F., Xia, Y., Parker, A.S., and Verma, I.M. (2012). IKK biology. Immunol Rev *246*, 239–253. 10.1111/j.1600-065X.2012.01107.x.

9. Scheidereit, C. (2006). IkappaB kinase complexes: gateways to NF-kappaB activation and transcription. Oncogene *25*, 6685–6705. 10.1038/sj.onc.1209934.

10. Karin, M., and Ben-Neriah, Y. (2000). Phosphorylation meets ubiquitination: the control of NF-[kappa]B activity. Annu Rev Immunol *18*, 621–663. 10.1146/annurev.immunol.18.1.621.

11. Hayden, M.S., and Ghosh, S. (2008). Shared principles in NF-kappaB signaling. Cell *132*, 344–362. 10.1016/j.cell.2008.01.020.

12. Brown, K., Gerstberger, S., Carlson, L., Franzoso, G., and Siebenlist, U. (1995). Control of I kappa B-alpha proteolysis by site-specific, signal-induced phosphorylation. Science *267*, 1485–1488. 10.1126/science.7878466.

13. Lin, Y.C., Brown, K., and Siebenlist, U. (1995). Activation of NF-kappa B requires proteolysis of the inhibitor I kappa B-alpha: signal-induced phosphorylation of I kappa B-alpha alone does not release active NF-kappa B. Proceedings of the National Academy of Sciences *92*, 552–556. 10.1073/pnas.92.2.552.

14. Manning, G., Whyte, D.B., Martinez, R., Hunter, T., and Sudarsanam, S. (2002). The protein kinase complement of the human genome. Science *298*, 1912–1934. 10.1126/science.1075762.

15. Manning, G., Plowman, G.D., Hunter, T., and Sudarsanam, S. (2002). Evolution of
protein kinase signaling from yeast to man. Trends Biochem Sci *27*, 514–520.
10.1016/s0968-0004(02)02179-5.

16. Dhanasekaran, N., and Premkumar Reddy, E. (1998). Signaling by dual specificity
kinases. Oncogene *17*, 1447–1455. 10.1038/sj.onc.1202251.

17. He, X., Jones, M.H., Winey, M., and Sazer, S. (1998). Mph1, a member of the Mps1-like
family of dual specificity protein kinases, is required for the spindle checkpoint in S.
pombe. J Cell Sci *111 (Pt 12)*, 1635–1647. 10.1242/jcs.111.12.1635.

18. Lauze, E., Stoelcker, B., Luca, F.C., Weiss, E., Schutz, A.R., and Winey, M. (1995).
Yeast spindle pole body duplication gene MPS1 encodes an essential dual specificity
protein kinase. EMBO J *14*, 1655–1663. 10.1002/j.1460-2075.1995.tb07154.x.

19. Ubersax, J.A., and Ferrell, J.E., Jr. (2007). Mechanisms of specificity in protein
phosphorylation. Nat Rev Mol Cell Biol *8*, 530–541. 10.1038/nrm2203.

20. Sang, D., Shu, T., Pantoja, C.F., Ibáñez de Opakua, A., Zweckstetter, M., and Holt, L.J.
(2022). Condensed-phase signaling can expand kinase specificity and respond to
macromolecular crowding. Mol Cell *82*, 3693-3711.e10. 10.1016/j.molcel.2022.08.016.

21. Cullati, S.N., Chaikuad, A., Chen, J.-S., Gebel, J., Tesmer, L., Zhubi, R., Navarrete-
Perea, J., Guillen, R.X., Gygi, S.P., Hummer, G., et al. (2022). Kinase domain
autophosphorylation rewires the activity and substrate specificity of CK1 enzymes. Mol
Cell *82*, 2006-2020.e8. 10.1016/j.molcel.2022.03.005.

22. Schrofelbauer, B., and Hoffmann, A. (2011). How do pleiotropic kinase hubs mediate
specific signaling by TNFR superfamily members? Immunol Rev *244*, 29–43.
10.1111/j.1600-065X.2011.01060.x.

23. Antonia, R.J., Hagan, R.S., and Baldwin, A.S. (2021). Expanding the View of IKK: New
Substrates and New Biology. Trends Cell Biol *31*, 166–178. 10.1016/j.tcb.2020.12.003.

24. Huse, M., and Kuriyan, J. (2002). The conformational plasticity of protein kinases. Cell
109, 275–282. 10.1016/s0092-8674(02)00741-9.

25. Liu, S., Misquitta, Y.R., Olland, A., Johnson, M.A., Kelleher, K.S., Kriz, R., Lin, L.L.,
Stahl, M., and Mosyak, L. (2013). Crystal Structure of a Human IκB Kinase β
Asymmetric Dimer. J Biol Chem *288*, 22758–22767. 10.1074/jbc.M113.482596.

26. Polley, S., Huang, D.-B., Hauenstein, A.V., Fusco, A.J., Zhong, X., Vu, D.,
Schröfelbauer, B., Kim, Y., Hoffmann, A., Verma, I.M., et al. (2013). A structural basis for
IκB kinase 2 activation via oligomerization-dependent trans auto-phosphorylation. PLoS
Biol. *11*, e1001581. 10.1371/journal.pbio.1001581.

27. Xu, G., Lo, Y.-C., Li, Q., Napolitano, G., Wu, X., Jiang, X., Dreano, M., Karin, M., and
Wu, H. (2011). Crystal structure of inhibitor of κB kinase β (IKKβ). Nature *472*, 325–330.
10.1038/nature09853.

28. Zhang, J., Clark, K., Lawrence, T., Peggie, M.W., and Cohen, P. (2014). An unexpected
twist to the activation of IKKbeta: TAK1 primes IKKbeta for activation by
autophosphorylation. Biochem J *461*, 531–537. 10.1042/BJ20140444.

29. Du, M., Ea, C.K., Fang, Y., and Chen, Z.J. (2022). Liquid phase separation of NEMO
induced by polyubiquitin chains activates NF-kappaB. Mol Cell *82*, 2415-2426 e5.
10.1016/j.molcel.2022.03.037.

30. Ea, C.K., Deng, L., Xia, Z.P., Pineda, G., and Chen, Z.J. (2006). Activation of IKK by
TNFalpha requires site-specific ubiquitination of RIP1 and polyubiquitin binding by
NEMO. Mol Cell *22*, 245–257. 10.1016/j.molcel.2006.03.026.

31. Chen, Z.J. (2012). Ubiquitination in Signaling to and Activation of IKK. Immunol Rev *246*,
95–106. 10.1111/j.1600-065X.2012.01108.x.

32. Schröfelbauer, B., Polley, S., Behar, M., Ghosh, G., and Hoffmann, A. (2012). NEMO
ensures signaling specificity of the pleiotropic IKKβ by directing its kinase activity toward
IκBα. Mol. Cell *47*, 111–121. 10.1016/j.molcel.2012.04.020.

33. Shaul, J.D., Farina, A., and Huxford, T. (2008). The human IKKbeta subunit kinase
domain displays CK2-like phosphorylation specificity. Biochem Biophys Res Commun
374, 592–597. 10.1016/j.bbrc.2008.07.082.

34. Barroga, C.F., Stevenson, J.K., Schwarz, E.M., and Verma, I.M. (1995). Constitutive
phosphorylation of I kappa B alpha by casein kinase II. Proceedings of the National
Academy of Sciences *92*, 7637–7641. 10.1073/pnas.92.17.7637.

35. Tergaonkar, V., Bottero, V., Ikawa, M., Li, Q., and Verma, I.M. (2003). IkappaB kinase-
independent IkappaBalpha degradation pathway: functional NF-kappaB activity and
implications for cancer therapy. Mol Cell Biol *23*, 8070–8083.

36. Delhase, M., Hayakawa, M., Chen, Y., and Karin, M. (1999). Positive and Negative
Regulation of IκB Kinase Activity Through IKKβ Subunit Phosphorylation. Science *284*,
309–313. 10.1126/science.284.5412.309.

37. Zandi, E., Chen, Y., and Karin, M. (1998). Direct phosphorylation of IkappaB by
IKKalpha and IKKbeta: discrimination between free and NF-kappaB-bound substrate.
Science *281*, 1360–1363. 10.1126/science.281.5381.1360.

38. Darwech, I., Otero, J.E., Alhawagri, M.A., and Abu-Amer, Y. (2010). Tyrosine
phosphorylation is required for IkappaB kinase-beta (IKKbeta) activation and function in
osteoclastogenesis. J Biol Chem *285*, 25522–25530. 10.1074/jbc.M110.121533.

39. Otero, J.E., Dai, S., Foglia, D., Alhawagri, M., Vacher, J., Pasparakis, M., and Abu-
Amer, Y. (2008). Defective osteoclastogenesis by IKKbeta-null precursors is a result of
receptor activator of NF-kappaB ligand (RANKL)-induced JNK-dependent apoptosis and
impaired differentiation. J Biol Chem *283*, 24546–24553. 10.1074/jbc.M800434200.

40. Chen, C., Ha, B.H., Thevenin, A.F., Lou, H.J., Zhang, R., Yip, K.Y., Peterson, J.R.,
Gerstein, M., Kim, P.M., Filippakopoulos, P., et al. (2014). Identification of a major
determinant for serine-threonine kinase phosphoacceptor specificity. Mol Cell *53*, 140–
147. 10.1016/j.molcel.2013.11.013.

41. Ahuja, L.G., Kornev, A.P., McClendon, C.L., Veglia, G., and Taylor, S.S. (2017).
Mutation of a kinase allosteric node uncouples dynamics linked to phosphotransfer. Proc
Natl Acad Sci U S A *114*, E931–E940. 10.1073/pnas.1620667114.

42. Yang, J., Garrod, S.M., Deal, M.S., Anand, G.S., Woods, V.L., Jr., and Taylor, S. (2005).
Allosteric network of cAMP-dependent protein kinase revealed by mutation of Tyr204 in
the P+1 loop. J Mol Biol *346*, 191–201. 10.1016/j.jmb.2004.11.030.

43. Berman, H.M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T.N., Weissig, H., Shindyalov,
I.N., and Bourne, P.E. (2000). The Protein Data Bank. Nucleic Acids Res *28*, 235–242.
10.1093/nar/28.1.235.

44. Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng,
E.C., and Ferrin, T.E. (2004). UCSF Chimera--a visualization system for exploratory
research and analysis. J Comput Chem *25*, 1605–1612. 10.1002/jcc.20084.

45. Abraham, M.J., Murtola, T., Schulz, R., Páll, S., Smith, J.C., Hess, B., and Lindahl, E.
(2015). GROMACS: High performance molecular simulations through multi-level
parallelism from laptops to supercomputers. SoftwareX *1–2*, 19–25.
10.1016/j.softx.2015.06.001.

46. Huang, J., and MacKerell, A.D. (2013). CHARMM36 all-atom additive protein force field:
validation based on comparison to NMR data. J Comput Chem *34*, 2135–2145.
10.1002/jcc.23354.

47. Jorgensen, W.L., Chandrasekhar, J., Madura, J.D., Impey, R.W., and Klein, M.L. (1983).
Comparison of simple potential functions for simulating liquid water. The Journal of
Chemical Physics *79*, 926–935. 10.1063/1.445869.

48. Nocedal, J., and Wright, S.J. eds. (1999). Numerical Optimization (Springer-Verlag)
10.1007/b98874.

49. On the Extension of the Davidon-Broyden Class of Rank One, Quasi-Newton
Minimization Methods to an Infinite Dimensional Hilbert Space with Applications to
Optimal Control Problems - NASA Technical Reports Server (NTRS)
https://ntrs.nasa.gov/citations/19710026200.

50. Liu, N., and Xu, Z. (2019). Using LeDock as a docking tool for computational drug
design. *218*, 012143. 10.1088/1755-1315/218/1/012143.

51. Eckhart, W., Hutchinson, M.A., and Hunter, T. (1979). An activity phosphorylating
tyrosine in polyoma T antigen immunoprecipitates. Cell *18*, 925–933. 10.1016/0092-
8674(79)90205-8.

52. Robbins, D.J., Zhen, E., Owaki, H., Vanderbilt, C.A., Ebert, D., Geppert, T.D., and Cobb,
M.H. (1993). Regulation and properties of extracellular signal-regulated protein kinases
1 and 2 in vitro. J Biol Chem *268*, 5097–5106.

53. Nolen, B., Taylor, S., and Ghosh, G. (2004). Regulation of Protein Kinases: Controlling
Activity through Activation Segment Conformation. Molecular Cell *15*, 661–675.
10.1016/j.molcel.2004.08.024.

54. Hunter, T. (1991). Protein kinase classification. Methods Enzymol *200*, 3–37.
10.1016/0076-6879(91)00125-g.

55. Cohen, P. (2002). The origins of protein phosphorylation. Nat Cell Biol *4*, E127–E130.
10.1038/ncb0502-e127.

56. Fischer, E.H., and Krebs, E.G. (1955). Conversion of phosphorylase b to phosphorylase
a in muscle extracts. J Biol Chem *216*, 121–132.

57. Krebs, E.G., and Fischer, E.H. (1955). Phosphorylase activity of skeletal muscle
extracts. J Biol Chem *216*, 113–120.

58. Borkovich, K.A., and Simon, M.I. (1990). The dynamics of protein phosphorylation in
bacterial chemotaxis. Cell *63*, 1339–1348. 10.1016/0092-8674(90)90429-i.

59. Gao, R., and Stock, A.M. (2009). Biological Insights from Structures of Two-Component
Proteins. Annu Rev Microbiol *63*, 133–154. 10.1146/annurev.micro.091208.073214.

60. Hunter, T. (2022). A journey from phosphotyrosine to phosphohistidine and beyond. Mol
Cell *82*, 2190–2200. 10.1016/j.molcel.2022.05.007.

61. Kalagiri, R., and Hunter, T. (2021). The many ways that nature has exploited the unusual
structural and chemical properties of phosphohistidine for use in proteins. Biochem J
478, 3575–3596. 10.1042/BCJ20210533.

62. Laub, M.T., Biondi, E.G., and Skerker, J.M. (2007). Phosphotransfer profiling: systematic
mapping of two-component signal transduction pathways and phosphorelays. Methods
Enzymol *423*, 531–548. 10.1016/S0076-6879(07)23026-5.

63. Pike, A.C., Rellos, P., Niesen, F.H., Turnbull, A., Oliver, A.W., Parker, S.A., Turk, B.E.,
Pearl, L.H., and Knapp, S. (2008). Activation segment dimerization: a mechanism for
kinase autophosphorylation of non-consensus sites. EMBO J *27*, 704–714.
10.1038/emboj.2008.8.

64. Larabi, A., Devos, J.M., Ng, S.L., Nanao, M.H., Round, A., Maniatis, T., and Panne, D.
(2013). Crystal structure and mechanism of activation of TANK-binding kinase 1. Cell
Rep *3*, 734–746. 10.1016/j.celrep.2013.01.034.

65. Tu, D., Zhu, Z., Zhou, A.Y., Yun, C.H., Lee, K.E., Toms, A.V., Li, Y., Dunn, G.P., Chan,
E., Thai, T., et al. (2013). Structure and ubiquitination-dependent activation of TANK-
binding kinase 1. Cell Rep *3*, 747–758. 10.1016/j.celrep.2013.01.033.

66. Mooster, J.L., Le Bras, S., Massaad, M.J., Jabara, H., Yoon, J., Galand, C., Heesters,
B.A., Burton, O.T., Mattoo, H., Manis, J., et al. (2015). Defective lymphoid
organogenesis underlies the immune deficiency caused by a heterozygous S32I
mutation in IκBα. J Exp Med *212*, 185–202. 10.1084/jem.20140979.

67. Courtois, G., Smahi, A., Reichenbach, J., Döffinger, R., Cancrini, C., Bonnet, M., Puel,
A., Chable-Bessia, C., Yamaoka, S., Feinberg, J., et al. (2003). A hypermorphic
IkappaBalpha mutation is associated with autosomal dominant anhidrotic ectodermal
dysplasia and T cell immunodeficiency. J Clin Invest *112*, 1108–1115.
10.1172/JCI18714.

68. Cheong, R., Bergmann, A., Werner, S.L., Regal, J., Hoffmann, A., and Levchenko, A.
(2006). Transient IkappaB kinase activity mediates temporal NF-kappaB dynamics in
response to a wide range of tumor necrosis factor-alpha doses. J Biol Chem *281*, 2945–
2950. 10.1074/jbc.M510085200.

69. Behar, M., and Hoffmann, A. (2013). Tunable Signal Processing through a Kinase
Control Cycle: the IKK Signaling Node. Biophys J *105*, 231–241.
10.1016/j.bpj.2013.05.013.

70. Lamarche, M.G., Wanner, B.L., Crépin, S., and Harel, J. (2008). The phosphate regulon
and bacterial virulence: a regulatory network connecting phosphate homeostasis and
pathogenesis. FEMS Microbiol Rev *32*, 461–473. 10.1111/j.1574-6976.2008.00101.x.

71. Bhate, M.P., Molnar, K.S., Goulian, M., and DeGrado, W.F. (2015). Signal Transduction
in Histidine Kinases: Insights from New Structures. Structure *23*, 981–994.
10.1016/j.str.2015.04.002.

72. Laub, M.T., and Goulian, M. (2007). Specificity in two-component signal transduction
pathways. Annu Rev Genet *41*, 121–145. 10.1146/annurev.genet.41.042007.170548.

73. Rieke, C., Papendieck, A., Sokolova, O., and Naumann, M. (2011). Helicobacter pylori-
induced tyrosine phosphorylation of IKKβ contributes to NF-κB activation. Biol Chem
392, 387–393. 10.1515/BC.2011.029.

74. Meyer, A.N., Drafahl, K.A., McAndrew, C.W., Gilda, J.E., Gallo, L.H., Haas, M., Brill,
L.M., and Donoghue, D.J. (2013). Tyrosine Phosphorylation Allows Integration of
Multiple Signaling Inputs by IKKβ. PLoS One *8*, e84497. 10.1371/journal.pone.0084497.

75. Lochhead, P.A., Sibbet, G., Morrice, N., and Cleghon, V. (2005). Activation-loop
autophosphorylation is mediated by a novel transitional intermediate form of DYRKs.
Cell *121*, 925–936. 10.1016/j.cell.2005.03.034.

76. Lochhead, P.A., Kinstrie, R., Sibbet, G., Rawjee, T., Morrice, N., and Cleghon, V. (2006).
A chaperone-dependent GSK3beta transitional intermediate mediates activation-loop
autophosphorylation. Mol Cell *24*, 627–633. 10.1016/j.molcel.2006.10.009.

77. Lochhead, P.A. (2009). Protein kinase activation loop autophosphorylation in cis:
overcoming a Catch-22 situation. Sci Signal *2*, pe4. 10.1126/scisignal.254pe4.

78. Bhattacharyya, R.P., Reményi, A., Good, M.C., Bashor, C.J., Falick, A.M., and Lim, W.A.
(2006). The Ste5 scaffold allosterically modulates signaling output of the yeast mating
pathway. Science *311*, 822–826. 10.1126/science.1120941.

79. Tigno-Aranjuez, J.T., Asara, J.M., and Abbott, D.W. (2010). Inhibition of RIP2's tyrosine
kinase activity limits NOD2-driven cytokine responses. Genes Dev *24*, 2666–2677.
10.1101/gad.1964410.

80. Ge, B., Gram, H., Di Padova, F., Huang, B., New, L., Ulevitch, R.J., Luo, Y., and Han, J.
(2002). MAPKK-independent activation of p38alpha mediated by TAB1-dependent
autophosphorylation of p38alpha. Science *295*, 1291–1294. 10.1126/science.1067289.

81. Lee, K.P.K., Dey, M., Neculai, D., Cao, C., Dever, T.E., and Sicheri, F. (2008). Structure
of the dual enzyme Ire1 reveals the basis for catalysis and regulation in non-
conventional RNA splicing. Cell *132*, 89–100. 10.1016/j.cell.2007.10.057.

82. Sidrauski, C., and Walter, P. (1997). The transmembrane kinase Ire1p is a site-specific
endonuclease that initiates mRNA splicing in the unfolded protein response. Cell *90*,
1031–1039. 10.1016/s0092-8674(00)80369-4.

Supplemental information:

Figure S1

732 **Figure S1:** Autoradiograph of *in vitro* kinase assay reactions carried out with FL IKK2 WT and
733 K44M using GST (1-54) IκBα as the substrate as a function of time.

Figure S2

Figure S2: A, Effect of increasing concentration of urea on tyrosine autophosphorylation of FL IKK2WT assessed in an *in vitro* kinase assay. **B,** Radioactive *in vitro* kinase assay performed with increasing concentration of urea up to 6M to check auto- and substrate phosphorylation. **C,** Inhibition of both substrate phosphorylation and tyrosine autophosphorylation activities of IKK2 checked in presence of common kinase inhibitors such as AMPPNP and Staurosporine, highly specific IKK2 inhibitors TPCA, Calbiochem Inhibitor VII and MLN120B. **D,** Mutation of AL-serines to nonphosphorylatable alanine (FL IKK2 S177A,181A double mutant) debilitates its kinase activity towards S32,36 of IκBα compared to WT or EE version of IKK2 as observed in an *in vitro* kinase assay.

Figure S3

Figure S3: A& B, Mass spectrometry analysis of ATP-treated IKK2 showing phosphorylation of Tyr188 and Tyr169, respectively.

Figure S4: A, A close-up view of the active site in active form in the context of the entire kinase domain of IKK2. Three tyrosines (Y169, Y188 and Y199) in immediate vicinity of active sites are highlighted. Position of ATP is modelled based on the structure of PKA. **B,** Substrate bound PKA in its active form; phosphorylatable serine of the substrate peptide (in magenta) is highlighted. **C,** A differential scanning calorimetry (DSC) thermogram showing a striking enhancement in folding stability of IKK2 upon ATP-treatment. **D,** Ribbon representation of different phosphorylated versions of IKK2 are shown: UnP-IKK2 in golden yellow, p-IKK2 in blue and P-IKK2 in copper red. **E,** Trajectories of RMSF for these structures are shown (left panel). For clarity, different regions of the KD are shown in the right. **F,** ATP-bound structure of p-IKK2 is shown. **G,** Dynamic cross-correlation matrix (DCCM) or contact map of each structure reveals phosphorylation-induced local as well as allosteric changes.

Figure S5

Figure S5

857
858
859
860 **Figure S5:** ESI-MS scan of the ADP used in this assay system showing negligible trace, if
861 any, of ATP at 50µM ADP concentration.

METHOD DETAILS

Reagents and Cell culture

Monoclonal mouse anti-IκBα (L35A5) (catalogue number: 4814), mouse anti-Phospho-IκBα (Ser$^{32/36}$) (5A5) (catalogue number: 9246) and rabbit anti-Phospho-IKKα/β (Ser$^{176/180}$) (16A6) (catalogue number: 2697) were from Cell Signaling Technology (CST). Monoclonal mouse anti-IKKβ (10AG2) (catalogue number: NB100-56509) was from Novus Biologicals. Polyclonal rabbit anti-6XHis (catalogue number: BB-AB0010) was from BioBharti Life Science. Monoclonal mouse anti-Phospho-Tyrosine (catalogue number: 610000) was from BD Biosciences. Polyclonal rabbit anti-GST (catalogue number: 924801), monoclonal mouse anti-HA 11 epitope (catalogue number: 901502) and mouse anti-Tubulinβ3 (catalogue number: 657402) were from BioLegend. Polyclonal rabbit anti-IKKα/β (H-470): sc7607 was from Santa Cruz Biotechnology (has been discontinued), and mouse anti-Phospho-Serine Q5 (catalogue number: 37430) was from Qiagen. Horseradish peroxidase-conjugated secondary antibodies against rabbit (catalogue number: 31460) and mouse IgG (catalogue number: 31430) were Pierce antibodies from Thermo Scientific. Adenosine-5'-Triphosphate (ATP), Disodium trihydrate (catalogue number: A-081-25) was from Gold Biotechnology and Adenosine Diphosphate (ADP) (Reference number: V916A) provided in ADP-Glo^{TM} Kinase assay (catalogue number: V6930) from Promega. The recombinant murine TNF-α used was from Roche.

Sf9 cells used for expressing proteins with baculovirus system were procured from Gibco™ Sf9 cells in Sf-900™ III SFM, Thermo Scientific (catalogue number: 12659017) and maintained in Gibco™Sf-900™ III SFM, Thermo Scientific (catalogue number: 12658019).

Protein expression and purification

IKK2 (WT and other constructs): Wild-type and mutant IKK2s were purified following a published protocol (Polley et al, 2013). Sf9 cells (Gibco™ Sf9 cells in Sf-900™ III SFM) at a density of 1.5-2 million/ml were infected with high titre viral stock (generally after the P3 stage) at a ratio of 1:100 or 1:50 as optimized for higher expression level as judged by a pilot experiment performed previously, and grown for 48 hours upon infection. Cells were harvested and lysed by sonication in a buffer containing 25 mM Tris-HCl pH 8.0, 200 mM NaCl, 10% Glycerol, 5 mM β-Mercaptoethanol (β-ME), 10 mM Imidazole, 20 mM β-glycerophosphate, 10 mM NaF, and 1 mM sodium orthovanadate and 1X Protease Inhibitor Cocktail (PIC, Sigma). The lysate was clarified by centrifugation at 20,000 x g for 45 minutes at 4˚C and incubated with Ni-NTA agarose resin (Qiagen, catalogue number: 30230) pre-equilibrated with lysis buffer for 2 hours at 4˚C. The resin was subsequently washed with lysis buffer containing 30 mM Imidazole. Protein was eluted with lysis buffer containing 250 mM Imidazole. Protein fractions were analysed on SDS-PAGE, pooled together and either flash frozen or subjected to size-exclusion chromatography on Superdex 200 Increase 10/300 GL (Cytiva) column equilibrated with 20 mM Tris-HCl pH 8.0, 200 mM NaCl, 10% Glycerol and 2 mM DTT. To make untagged IKK2, protein fractions eluted from the Ni-NTA resin of desired purity were treated with TEV-protease at a ratio of 50:1 (weight by weight, IKK2: TEV) in the cold room for more than 10 hours after the addition of 1 mM of EDTA, the resulting protein mix was then loaded on to the SEC as described above. To prepare autophosphorylated IKK2, the fractions with the desired purity level were treated with 0.5-1 mM of ATP, 10 mM of $MgCl_2$ and phosphatase inhibitors and the autophosphorylation reaction was carried out at 27˚C for 1 hour. The resulting protein solution was then concentrated in a YM30 concentrator and loaded onto Superdex 200 Increase 10/300 GL column. Elution fractions from the size-

exclusion step were analysed on SDS-PAGE and peak fractions of desired quality

(Cytiva) were pooled together and concentrated using a YM30 Centriprep or Centricon

devices (Millipore) depending upon the volume of the sample and flash frozen

immediately.

IκBα (WT, mutant and other constructs): His-tagged full-length and GST-tagged

deletion (1-54) constructs of IκBα (both the wild-type and Ser32Ala, Ser36Ala,

Ser32,36Ala single and double-mutant versions) were expressed in *E. coli* Rosetta2

(DE3). The cells were grown overnight at 16˚C after 0.3 mM IPTG induction. The cell

pellet was resuspended in 40 ml of Lysis Buffer (25 mM Tris pH 7.5, 200 mM NaCl,

10 mM imidazole, 10% Glycerol, 5 mM β-mercaptoethanol, and 1 mM PMSF). The

cells were lysed by using a sonicator keeping the cell suspension chilled on ice. The

lysate was clarified by centrifugation at 15,000 x g for one hour at 4°C. About 3 ml of

Ni-NTA Agarose resin was equilibrated with lysis buffer, and the clarified lysate

(supernatant) was mixed with it later. The protein was allowed to bind the resin by

incubating for 2 hours on a rotary mixer in a 4°C room. The resin was washed with the

lysis buffer containing 500 mM NaCl (high salt wash) for about 10 CV to clear out non-

specific proteins from binding. Then the resin was thoroughly washed with the lysis

buffer containing 20 mM imidazole. His-tagged IκBα protein was eluted under gravity

flow, using elution buffer (the lysis buffer containing 250 mM imidazole). ~1 ml volume

of each fraction was collected. [For GST-tagged proteins, Glutathione Sepharose resin

(Cytiva, catalogue number: 17-5279-01] was used and elution was performed with the

help of reduced glutathione). The main elution fractions were subjected to size-

exclusion chromatography (SEC) on a HiLoad 16/600 pg Superdex 200 column

(Cytiva) and the gel run revealed cleaner protein fractions. The peak fractions were

saved and stored at -80˚C.

NEMO (WT, mutant and other constructs): NEMO FL, NEMO double mutant C76A

C95S (site-directed mutagenesis done in pET24d vector) and NEMO (44-196)

(truncates in pET24d) were expressed in *E. coli* Rosetta2 DE3 cells. The cells were

grown overnight at ~21˚C after 0.25 mM IPTG induction (at 0.3-0.4 O.D.) and 50-100

µM Zn^{2+}. [OR The cells were grown overnight (~17 hours) at 16˚C after 0.1 mM IPTG

induction (at 0.2-0.3 O.D.) and 50-100 µM Zn^{2+}]. The cell pellet was resuspended in

40 ml of Lysis Buffer (25 mM Tris pH 7.5, 200 mM NaCl, 10 mM imidazole, 10%

Glycerol, 5 mM β-mercaptoethanol, 20 µM $ZnCl_2$, 0.5 M urea (acting like an osmolyte).

The rest of the steps performed were typical of a Ni-NTA purification as described

above. The main elution fractions were subjected to size-exclusion chromatography

on a HiLoad 16/600 pg Superdex 200 column (Cytiva) and the cleaner peak fractions

were stored at -80˚C.

Kinase Assay with purified proteins

Typically, 100-200 ng (unless mentioned otherwise) of purified kinase was used in

each *in vitro* kinase reaction. 0.5-1 µg of IκBα substrate (full-length or GST-tagged 1-

54) was incubated with kinase for 30 mins or indicated time periods at 27˚C in

presence of 20 µM ATP with 0.1 µCi of γ-P32-ATP in case of radioactive assay or 50-

100 µM ATP in case of Western blot assay in a reaction buffer containing 20mM

HEPES pH 7.5, 100 mM NaCl, 15 mM $MgCl_2$, 2 mM DTT, 0.2 mM Na_3VO_4, 10 mM

NaF, 20 mM β-Glycerophosphate. The reaction was stopped by heating the samples

at 95˚C for 5 mins after the addition of 1X-Laemmli buffer. Samples were run on 8 or

10% SDS-PAGE. For radioactive assays, the gel was then dried and analysed either

by film or phosphor-imaging. For Western blot assays, proteins from the gel were

electro-transferred onto nitrocellulose/PVDF membrane and probed with desired

antibodies. Phosphorylated tyrosine and serine residues were detected using

monoclonal antibodies against phospho-tyrosine and phospho-serine.

IP-Kinase assay

Immortalized *ikk2⁻/⁻* mouse embryonic fibroblast 3T3 cells were reconstituted by

retroviral transduction with HA-tagged WT or mutant pBABE-IKK2 constructs. Stable

expression of IKK2 was first confirmed by immunoblotting. Cells were then stimulated

with recombinant murine TNF-α (Roche) at 70-80% cell confluency. IKK complex was

immunoprecipitated from whole cell extracts using a NEMO-specific antibody (BD) and

Protein A agarose (Upstate) for 2 hrs in the cold room. Whole-cell extract used was

pre-cleared by incubation with Protein A agarose bead for an hour. The

immunoprecipitated complex was washed extensively, including a final washing step

with the kinase assay buffer devoid of ATP and incubated with GST-IκBα (1-54) and

20 μM ATP and 5 μCi of γ-^{32}P-ATP for 30 mins at 30˚C in kinase assays. Reactions

were stopped by the addition of 2X-Laemmli SDS buffer to samples and heating at

95˚C for 5 mins. Reaction products were resolved on a 12% SDS PAGE. The extent

of substrate phosphorylation was measured by cutting out the appropriate area of the

gel containing the substrate and subjecting it to phosphor-imaging (Typhoon, Cytiva)

after drying. As a control, the amount of IKK2 in each reaction was estimated by

transferring the IKK2-containing gel area onto a nitrocellulose membrane and probing

it with anti-IKK1/2 antibody.

Sample preparation for LC-MS/MS analyses for Tyrosine phosphorylation

detection

Protein samples were diluted in TNE (50 mM Tris pH 8.0, 100 mM NaCl, 1 mM EDTA)

buffer. RapiGest SF reagent (Waters Corp.) was added to the mix to a final

concentration of 0.1% and samples were boiled for 5 min. Tris (2-carboxyethyl)

phosphine (TCEP) was added to a final concentration of 1mM and the samples were

incubated at 37˚C for 30 min. Subsequently, the samples were carboxymethylated

with 0.5 mg/ml of iodoacetamide for 30 min at 37˚C followed by neutralization with

2mM final concentration of TCEP. Protein samples prepared as above were digested

with trypsin (trypsin: protein ratio - 1:50) overnight at 37˚C. RapiGest was degraded

and removed by treating the samples with 250 mM HCl at 37˚C for 1 h followed by

centrifugation at 14000 rpm for 30 min at 4˚C. The soluble fraction was then added to

a new tube and the peptides were extracted and desalted using C18 desalting columns

(Thermo Scientific).

LC-MS/MS analysis

Trypsin-digested peptides were analysed by ultra-high-pressure liquid

chromatography (UPLC) coupled with tandem mass spectroscopy (LC-MS/MS) using

nano-spray ionization. The nano-spray ionization experiments were performed using

a TripleTOF 5600 hybrid mass spectrometer (ABSCIEX) interfaced with nano-scale

reversed-phase UPLC (Waters corporation nano ACQUITY) using a 20 cm-75-micron

ID glass capillary packed with 2.5 µm C18 (130) CSHTM beads (Waters corporation).

Peptides were eluted from the C18 column into the mass spectrometer using a linear

gradient (5–80%) of Acetonitrile (ACN) at a flow rate of 250 µl/min for 1h. The buffers

used to create the ACN gradient were: Buffer A (98% H2O, 2% ACN, 0.1% formic

acid, and 0.005% TFA) and Buffer B (100% ACN, 0.1% formic acid, and 0.005% TFA).

MS/MS data were acquired in a data-dependent manner in which the MS1 data was

acquired for 250ms at m/z of 400 to 1250 Da and the MS/MS data was acquired from

m/z of 50 to 2,000Da. The Independent data acquisition (IDA) parameters were as

follows; MS1-TOF acquisition time of 250 milliseconds, followed by 50 MS2 events of

48 milliseconds acquisition time for each event. The threshold to trigger MS2 event

was set to 150 counts when the ion had the charge state +2, +3 and +4. The ion

exclusion time was set to 4 seconds. Finally, the collected data were analysed using

Protein Pilot 4.5 (ABSCIEX) for peptide identifications.

ESI-MS analyses of ADP

50 µM of ADP solution was prepared to be analysed by Electrospray Ionisation Mass

Spectrometry (ESI-MS). It was performed in a Xevo G2-XS QToF mass spectrometer

(Waters Corporation) using capillary at 3 kV, with voltage parameters of sampling cone

at 40V and source offset at 80V. The source temperature was set at 100˚C and the

desolvation temperature at 40˚C. The gas flow rates were as follows: Cone gas flow

rate at 50 litre/hr and desolvation gas flow rate at 400 litre/hr. The sample was passed

at a 5 µl/min flow rate. The MS data was acquired for a mass range of 50-2000 m/z

and the acquisition time was 1 minute.

Radioactive Phosphotransfer assay

Phosphotransfer assays were performed in a wide range of concentration of IKK2,

IκBα and ATP. Typically, 10 µM purified IKK2 protein was first autophosphorylated in

the presence of 100-200 nM of γ32P-ATP (PerkinElmer) in kinase assay buffer for 1

hour at 30˚C. 70 µl of the resulting reaction mixture was first passed through a Micro

Bio-Spin Bio-Gel P-30 spin column (Bio-Rad; 40 kDa MWCO) equilibrated with 20 mM

HEPES pH 7.9, 100 mM NaCl, 0.2 mM Na_3VO_4, 10 mM NaF, 20 mM β-

Glycerophosphate, 2 mM DTT, 2 mM EDTA and 1X protease inhibitor cocktail. Eluted

protein was subjected to another round of buffer exchange on a Bio-Gel P-30 column

equilibrated with the same buffer mentioned above devoid of EDTA and containing 15

mM $MgCl_2$. Phosphorylated IKK2 thus obtained was used in the phosphotransfer

reaction. 2-3 µM of Phosphorylated IKK2 was incubated with 10-20 µM of WT or

mutant IκBα protein at 30˚C in kinase assay buffer in a total volume of 20 µl for different

time periods as indicated in Figure 5B and 6A. The reactions were stopped by the

addition of 10 µl of 4X Laemmli SDS buffer containing 100 mM EDTA to samples and

heating at 95˚C for 5 mins. Different concentration of ADP or ATP was added to the

assay to assess the role of these nucleotides in phosphotransfer efficiency. Routinely,

10-15µl of reaction products were resolved on 10% SDS PAGE and radiolabelled

samples were monitored by phosphor imaging of the dried gel on a Typhoon scanner.

For the phosphotransfer reaction described in Figure 4a, 10 µM purified IKK2 was

autophosphorylated with 100 nM of γ^{32}P-ATP in the presence of 40 µM of cold ATP.

Phospho-transfer assay with cold ATP

About 10 µM of purified FL IKK2WT was treated with 100 µM of cold ATP in kinase

assay buffer (containing 1X protease inhibitor cocktail) for one hour at 30˚C. The

resulting reaction mixture was then loaded onto a Superdex 200 increase 10/300 GL

(24ml bed volume, Cytiva) SEC column to purify autophosphorylated FL IKK2WT

separated from excess ATP. SEC was performed in a buffer containing 20 mM

HEPES, 100 mM NaCl, 20 mM β-glycerophosphate, 10 mM NaF, 0.2 mM Na_3VO_4 and

2 mM DTT. The elution fractions containing phosphorylated IKK2 were incubated with

FL IκBαWT in the presence of different concentrations of ATP or ADP at different time

points (as mentioned in Figure 3f) at 30˚C. The reactions were stopped by adding 4X

Laemmli SDS dye, heated at 95˚C for 5 minutes and were run on a 10% SDS-PAGE.

This was followed by immunoblotting and detection was done using monoclonal

antibodies against phospho-tyrosine and phospho Ser[32/36] IκBα.

Molecular Dynamics Simulation

The kinase domain structure of IKK2 (PDB ID: 4E3C), spanning residues 16 to 310,

was obtained from the PDB database. We created unphosphorylated forms (at Ser177

& Ser181) and two phosphorylated forms (phosphorylated at Ser177 & Ser181 and at

Tyr169 & Ser177 & Ser181) of IKK2 using UCSF CHIMERA software. These

structures underwent molecular dynamics (MD) simulations using the

GROMACSv2023.1 simulation package, enabling a comparative analysis of their

structural and energetic stability. The Charmm36 force field was applied to generate

coordinates and topology files for the IKK2 structures. For simulation, a cubic box was

defined and filled with TIP3 water molecules. Sodium and chloride ions were added to

neutralize the three systems. The system underwent two-stage minimization using

steepest-descent and conjugate-gradient algorithms. Equilibration was carried out

under NVT (constant particle number, volume, and temperature) and NPT (constant

particle number, pressure, and temperature) conditions for 200 ps at 300 K and 1 atm

pressure. MD simulations were conducted for 200 ns for each protein. To assess

global structural changes, Root Mean Square Deviation (RMSD) analysis was

employed, while Root Mean Square Fluctuation (RMSF) analysis revealed the overall

and local motion/flexibility of the proteins. The *gmx_energy* module was utilized to

compute energy during the simulation. To understand the correlations of the atomic

motions in different regions, a dynamic cross-correlation analysis was carried out. The

initial step of dynamic cross-correlation analysis was to construct the covariance

matrix which examines the linear relationship of atomic fluctuations for individual

atomic pairs. The covariance matrix was calculated using the *gmx_covar* tool.

Dynamic cross-correlation matrices (DCCM) were calculated from the C_α-trace

covariance matrix principal components. Results are displayed as colour-coded with

a value of −1 indicating completely anti-correlated motions and a value of +1 indicating

completely correlated motions.

Molecular Docking Analysis

The terminal structures from the simulations (200th ns) doubly phosphorylated IKK2

at S177, S181 (p-IKK2) form were subjected to molecular docking analysis using ATP

and ADP as ligands, allowing for comparison of binding affinities. LeDock software

was employed for ligand-protein docking. The top 100 docking solutions were

extracted. The p-IKK2: ATP poses were compared with available kinase: ATP

complexes, e.g., PKA: ATP complex (PDB ID: 1ATP). The pose that resembled most

with the pose found in PDB ID 1ATP was chosen for further analysis. p-IKK2:ADP

poses were also compared likewise, where IRE1: ADP (PDB ID: 2RIO) complex was

chosen for comparison and the pose closest to that in PDB ID: 2RIO was chosen for

further analysis.

For assessing the ability of P-IKK2 to accommodate ATP or ADP, the MD-simulation-

derived P-IKK2 structure was superimposed on the p-IKK2 structure bound to

ATP/ADP, and compared as described.

1129