Title:

Dual-specific autophosphorylation of kinase IKK2 enables phosphorylation of substrate IκBα without requiring ATP

Prateeka Borar[1], Tapan Biswas[2], Tom Huxford[3], Gourisankar Ghosh[2] & Smarajit Polley[1,*]

[1]Department of Biological Sciences, Bose Institute, Kolkata, India, [2]Department of Chemistry and Biochemistry, University of California San Diego, La Jolla, USA, [3]Department of Chemistry and Biochemistry, San Diego State University, San Diego, USA.

*Correspondence to be addressed to: spolley@jcbose.ac.in, smarajit.polley@gmail.com

Abstract:

IKK2, a Ser/Thr kinase, acts as a gateway to canonical activation of NF-κB underlying stress

signalling. The immediate targets of IKK2 are two serines (S32 and S36) of IκBα, which

undergo phosphorylation within minutes of a specific stimuli. Here, we report that IKK2

autophosphorylates itself at tyrosine residues in *cis* in addition to its activation loop serines.

Mutation of one such tyrosine, Y169, located in proximity to the active site, to phenylalanine

renders IKK2 inactive for phosphorylation of S32 of IκBα. Surprisingly, we observe that auto-

phosphorylated IKK2 relayed phosphate group(s) to IκBα even in the absence of ATP if ADP

is present. We also observed that mutation of K44, an ATP-binding lysine conserved in all

protein kinases, to methionine renders IKK2 inactive towards specific phosphorylation of S32

or S36 of IκBα, but not non-specific substrates. These observations highlight an unusual

evolution of IKK2, in which autophosphorylation of tyrosine(s) in the activation loop and the

invariant ATP-binding K44 residue are required in defining its substrate specificity for S32,S36

of IκBα. We speculate that this dual specificity may provide the desired specificity to substrate

phosphorylation ensuring the fidelity of NF-κB activation.

Introduction:

44
Inflammatory or pathogenic signals trigger precisely regulated induction of the transcription

factor NF-κB in metazoans [1]. This process depends upon the upstream activation of Inhibitor

of κB Kinase-complex (IKK-complex), comprising two catalytic subunits, IKK1 (also known as

IKKα) and IKK2 (or IKK2β), and an essential regulatory scaffolding protein, NEMO (IKKγ)[2-4].

IKK2 activity is maintained at a very low basal level in resting cells [5-8]. Upon activation, IKK2

phosphorylates the N-terminus of IκBα (Inhibitor of κBα) protein specifically at two Serine

residues (S32 and S36) marking it for ubiquitinylation-mediated proteasomal degradation. This

liberates NF-κB to execute its gene expression program [6,9-11]. Mutation of both S32 and S36

to alanine residues converts IκBα into a nonphosphorylatable super-repressor of NF-κB[12,13].

Furthermore, IKK possesses exquisite specificity for these two serines, it fails to phosphorylate

IκBα when S32 and S36 are mutated even to another phosphorylatable residue, threonine[2].

A majority of the eukaryotic protein kinases are Ser/Thr/Tyr kinases (STYKs) that phosphorylate specific namesake residues of their cognate substrates[14,15]. STYKs could specifically phosphorylate S/T or Y[14,16] or S/T as well as Y residues[17,18]. STYKs are rather promiscuous but can display a high substrate specificity in an appropriate context [19]. IKK2 is one such pleotropic kinase that also ensures activation of a particular signaling cascade with fidelity [20,21]. IKK2 toggles between inactive and active states based on phosphorylation status of two serine residues (S177 and S181; S176 and S180 in IKK1) within its activation loop (AL) [22], [23-25]. Replacement of these two serines with phospho-mimetic glutamate residues (S177E,S181E; henceforth EE) renders the kinase constitutively active. IKK2 is reported to be primed through phosphorylation at S177 by the upstream kinase TAK1 that leads to auto-phosphorylation at S181, thus rendering the kinase fully active in cells[26]. However, oligomerization upon association with NEMO, in presence of linear or Lys-63 linked poly-ubiquitin (Ub) chains or at a high concentration, also enables *trans* auto-phosphorylation at S177 and S181 [24,27-29]. Activated IKK2 phosphorylates IκBα specifically at residues S32/S36 and the C-terminal SDD (residue range 645-756) is necessary for this [30]. Other sites of IκBα, but not S32/S36, can be phosphorylated by various kinases [31,32]. These findings underscore the ability of IKK2 to function specifically in ensuring rapid NF-κB activation in canonical-signaling. It is, however, still not fully understood how IKK2 achieves its exquisite specificity towards S32/S36 of IκBα and induces NF-κB transcriptional activity.

An in-depth analysis of the catalytic features of IKK2 revealed that IKK2 can phosphorylate its own tyrosine residue(s) in addition to its own serines or the serines of its substrate IκBα, i.e., IKK2 possesses dual specificity. This phosphorylation at tyrosine residue(s) is autocatalytic. Furthermore, we observed that, after auto-phosphorylation at tyrosine, IKK2 can phosphorylate IκBα in the absence of ATP, as long as ADP is present in the reaction. Mass spectrometry of IKK2 identified that Y169 and Y188, both present within proximity of the enzyme active site, can undergo auto-phosphorylation in *cis*. This phosphorylation at tyrosine residues is dependent upon the prior activation of IKK2 via AL

S177/S181 phosphorylation, and critical to phosphorylation of signal responsive S32/S36 of

IκBα. Mutation of Y169 of IKK2 (which has long been associated with pathological conditions

such as ectodermal dysplasia) to phenylalanine severely compromised phosphorylation of

IκBα at S32. We also found that mutation of the conserved ATP-interacting residue K44 in

IKK2 to methionine leads to a loss of Tyr-auto-phosphorylation activity of the kinase and its

ability to phosphorylate S32/S36 of IκBα. Interestingly, K44M retained non-specific kinase

activity. Overall, our results reveal unique and transient tyrosine phosphoester intermediates

of IKK2 that appear to be critical in relaying phosphoryl group(s) specifically to S32/S36 of

IκBα, thus ensuring fidelity of NF-κB activation.

Results:
93
IKK2 undergoes auto-phosphorylation at uncharacterized sites: During our study of IKK2

activation, we found increasingly higher levels of IKK2 phosphorylation upon incubation of a

recombinant pure IKK2 with radioactive ATP. We surmised that one or multiple residues

besides the known phosphorylation sites (S177/181) might also be phosphorylated (**Fig. 1a**

and **Fig. 1b**). This phosphorylation of IKK2 did not display any reliance upon the presence of

substrate IκBα or NEMO (**Fig. 1c**). To confirm if this phosphorylation of IKK2 is indeed auto-

phosphorylation, and not dependent upon another kinase co-purified with IKK2, we generated

a K44M mutant of full-length IKK2, which is largely inactive (K44 is a conserved ATP-binding

residue in STYK kinases). An *in vitro* kinase assay of IKK2-K44M in absence or presence of

wild type (WT) IKK2 using γ^{32}P-ATP as the phospho-donor confirmed that the IKK2-K44M

lacked both substrate (IκBα) phosphorylation and auto-phosphorylation activities

(**Supplementary Data Fig. 1a**). The auto-phosphorylation is further supported by *in vitro*

radioactive kinase assay in the presence or absence of an IKK-specific ATP-competitive

inhibitor, Calbiochem Inhibitor VII (**Fig. 1d**).

Next, we used various C-terminal (serine-rich region) truncation mutants of the IKK2-

EE lacking known phosphorylation sites S177 and S181 [33]. The shortest construct, Δ664EE,

lacked the whole flexible C-terminus (665-756); and this was further treated with trypsin to

eliminate residues, if any, from the flexible ends of the enzymes prior to *in vitro* kinase assays.

All these mutants are found active in auto- and IκBα substrate phosphorylation (**Fig. 1e**). Auto-

phosphorylation despite the absence of AL serines and serines within the C-terminus strongly

suggested the presence of hitherto uncharacterized phosphorylation sites in IKK2. A striking

enhancement in stability of IKK2 upon this auto-phosphorylation is reflected in the change of

Tm from ~ 40 °C to ~ 50 °C in differential scanning calorimetry (DSC) scans in the presence

of ADP and ATP (**Supplementary Data Fig. 1b**). This behavior tallies with increased solubility

of IKK2 observed after auto-phosphorylation that was essential for its successful

crystallization[24].

Auto-phosphorylation of IKK2 reveals dual specificity: To test if Tyr residues can undergo

auto-phosphorylation in addition to Ser (and Thr), IKK2 incubated with Mg^{2+}-ATP for various

time periods was probed with antibodies that specifically recognize AL serines of IKK2 (**Fig.**

2a, upper panel) or phospho-tyrosine residues (**Fig. 2a**, middle panel). Unexpectedly, we

observed *de novo* auto-phosphorylation of IKK2 on tyrosine residues along with the

anticipated auto-phosphorylation of AL serines. The tyrosine auto-phosphorylation was

abolished in the presence of an IKK-specific inhibitor, Inhibitor VII (**Fig. 2b**). A similar assay

was performed to monitor the phospho-tyrosine level in IKK2 as a function of urea

concentration. This revealed the detrimental effects of urea at concentrations beyond 1M

(**Supplementary Data Fig. 2a**). A more sensitive assay with γ^{32}P-ATP as the phospho-donor

also indicated that urea concentrations beyond 1M inhibited both auto-phosphorylation (pan-

phosphorylation) and IκBα phosphorylation (**Supplementary Data Fig. 2b**). Furthermore, the

IKK2-K44M mutant (kinase inactivating) did not display any tyrosine phosphorylation when

incubated with ATP unlike WT IKK2 (**Fig. 2c**). These results confirm novel tyrosine

phosphorylation ability of IKK2, demonstrating its dual specificity to both its own Ser and Tyr

residues.

Next, we checked the requirement of the C-terminal <u>NE</u>MO <u>b</u>inding <u>d</u>omain (NBD) for

autocatalytic dual specificity. Constructs of various truncations (lacking NBD) were capable of

phosphorylating IκBα at signal-responsive S32 and S36, and phosphorylating itself on tyrosine

residue(s) (**Fig. 2d**). Next, we assessed the autocatalysis at Ser/Thr and Tyr by IKK2 using a

constitutively active NBD-truncated version of IKK2 (IKK2$^{\Delta664}$EE). The protein was recognized

by monoclonal anti-p-Ser antibody with or without fresh ATP-treatment as anticipated;

however, p-Tyr antibody recognized it strongly only upon fresh ATP-treatment (**Fig. 2e**).

To identify tyrosine residues that are phosphorylated *de novo* on IKK2 upon addition

of fresh ATP, we performed LC/MS/MS analyses. We identified at least two residues, Y169

and Y188, to be phosphorylated upon fresh ATP treatment[34,35] (**Supplementary Data Fig.**

2c,d). Interestingly, both are located within the AL (**Supplementary Data Fig. 2e**); Y169 is

the DFG+1 (DLG in case of IKK2 and IKK1) residue while Y188 (equivalent to Y204 of PKA)

is located at the end of AL. It was previously established that DFG+1 position is crucial for

defining the substrate specificity of a kinase[36]. Y188 counterpart Y204 of PKA is reported to

allosterically modulate PKA activity by altering its conformational entropy[37,38]. We also

observed that activity of IKK2 towards S32 and S36 of IκBα and its own tyrosine residue(s)

were affected similarly when the kinase assays were performed in the presence of various

kinase inhibitors, both general (AMPPNP and Staurosporine) and IKK-specific (MLN-120B,

TPCA, Inhibitor VII) (**Supplementary Data Fig. 2f**). These observations confirm that canonical

activity of IKK2 towards S32 and S36 of IκBα, and towards its own tyrosine residues are

mediated by the same active site pocket.

To investigate if Tyr phosphorylation occurs in *cis* (within the same kinase molecule,

intramolecular) or *trans* (by a different kinase molecule, intermolecular), we incubated full-

length IKK2-K44M with ATP in absence or presence of a constitutively active IKK2$^{\Delta743}$-EE

lacking the extreme C-terminal end. IKK2$^{\Delta743}$-EE was pre-treated with ATP, and purified

through a Superdex200 size-exclusion column to remove unutilized ATP. Absence of p-Tyr

signal on IKK2^{FL}-K44M irrespective of the presence of IKK2$^{\Delta743}$-EE, and presence of the p-Tyr

signal on the IKK2$^{\Delta743}$-EE strongly suggested that tyrosine auto-phosphorylation occurs in *cis*,

and not in *trans* (**Fig. 2i**, **upper panel**). In contrast, strong phosphorylation of AL S177/S181

on the IKK2^{FL}-K44M by IKK2$^{\Delta 743}$-EE suggests that AL serine auto-phosphorylation of IKK2

could occur in *trans*, agreeing with our previous findings (**Fig. 2f, middle panel**)[24].

167   To test the effect of phosphorylation at Y169 and Y188, we generated Y169F and

Y188F mutants. Radioactive *in vitro* kinase assays using recombinant purified IKK2^{FL}-Y188F

and IKK2^{FL}-Y169F mutants revealed drastic reduction in activity of IKK2 (**Fig. 2g**). Since the

DFG+1 residue in kinases (Y169 in IKK2) is known to dictate its specificity, we created

different mutants of the IκBα substrate where S32 and S36 were singly and independently

altered to either alanine (unphosphorylatable) or glutamic acid (phosphomimetic) residues.

Mutation of S36 to either Ala or Glu drastically reduced the phosphorylation signal of IκBα

while that for the S32A/E mutants were minimally affected (**Fig. 2h**). This suggests that

presence of Tyr (and possibly its phosphorylation) is critically important for signal responsive

phosphorylation of IκBα. Since pTyr mimic residue is not available, testing roles of modified

tyrosine residues is not easily amenable.

178   Next, we tested the consequence of Y169F and Y188F mutants on NF-κB signaling.

We reconstituted IKK2 knock-out mouse embryonic fibroblast cells with WT, Y169F and

Y188F mutants of IKK2, and measured signal induced activation of IKK2 upon TNF-α

treatment. IKK2 activity is significantly reduced in Y169F and nearly abolished in Y188F (**Fig.**

2i). Together these results suggest a communication between the phosphorylation of Y169 of

IKK2 and that of S32 of IκBα.

Freshly auto-phosphorylated IKK2 phosphorylates IκBα in absence of ATP: We

observed phosphorylation of tyrosine residue(s) by WB or mass spectrometry only upon fresh

ATP-treatment, indicating that tyrosines of purified IKK2 loses its phosphate group[39]. It may

be noted that, we did not observe any density of phosphates on any of these residues in the

human IKK2 crystal structure derived from ATP-treated protein sample[24]. This suggested an

intriguing possibility that the phosphate linked to Tyr residue(s) may serve as a temporary

phosphate-sink and could eventually be transferred to substrate IκBα. To test this possibility,

we first allowed purified IKK2 to undergo auto-phosphorylation by incubating the excess

kinase with limiting amounts of radiolabeled γ-[32]P-ATP followed by two passes through

desalting spin columns (40 kDa MW cut off) to remove unreacted ATP. Nucleotide-free auto-

phosphorylated IKK2 (P-IKK2) was incubated with IκBα either in absence of nucleotide or with

newly added unlabeled nucleotides, and phosphorylation status of the substrate was

subsequently monitored by autoradiography after resolving the reaction mixture on an SDS-

PAGE (**Fig. 3a**). Transfer of radiolabeled phosphate to IκBα was observed and the transfer

was significantly enhanced in the presence of cold ADP (**Fig. 3b**). This phosphotransfer did

not show a dependence on ADP concentrations nor did it significantly decrease in the

presence of a high concentration of cold ATP. That addition of excess cold ATP did not reduce

the extent of IκBα phosphorylation strongly supports transfer of [32]P-phosphate from [32]P-IKK2

to IκBα (**Fig. 3b**) by a relay mechanism with or without involving ADP.

We compared available structural models of human IKK2 to investigate whether

tyrosine phosphorylation within the activation loop could be supported based on its structural

features[23],[24]. We observe distinct conformations of the activation loop containing S177/S181.

The structure of IKK2 dimer is asymmetric in which the activation loop of one monomer is

largely disordered while that of the other monomer is mostly ordered (PDB id 4KIK)[23]. In the

active monomers S177 and S181 are clearly observed. In these active state conformers, the

position and conformation of Tyr169 appears to be well-poised for accepting a phosphate from

ATP (**Fig. 3c**). A superposition of IKK2 with a pseudo-substrate-bound PKA shows that the

hydroxyl of Tyr169 in IKK2 projects toward the γ-phosphate of ATP similarly to the serine

hydroxyl on the PKA inhibitor pseudo-substrate[40,41] (**Fig. 3c, Supplementary Data Fig. 3a**).

Strikingly, Tyr1238 within the activation loop of a constitutively active oncogenic mutant of

RON tyrosine kinase, which is thought to be activated through *trans* auto-phosphorylation

under normal conditions, also appears well-poised for auto-phosphorylation in *cis* as the

Tyr169 of IKK2[42] (**Supplementary Data Fig. 3b**). These structural observations suggest that

it is possible for tyrosine residues within the flexible activation segment to undergo auto-

phosphorylation in *cis* and possibly render the kinase with a novel phospho-transfer activity.

We undertook a different strategy to further confirm transfer of phosphate from the

kinase (and not ATP) to the substrate. We auto-phosphorylated IKK2^{FL}-WT with excess cold

ATP and removed the unreacted ATP from the reaction mixture using size-exclusion

chromatography on a Superdex200 column (**Fig. 3d**). P-IKK2 eluted at fractions

corresponding to MW between 670 and 158 kDa, i.e., much larger than free ATP (**Fig. 3e**).

This P-IKK2 protein was incubated with FL IκBα either in presence or absence of ADP for

different time periods. A similar reaction in presence of ATP instead of ADP was performed

as a positive control. Phosphorylation of S32 and S36 was efficient in presence of 10 and 50

μM ADP but absent when ADP was absent in the reaction mixture. Phosphorylation of IκBα

in the presence of 50 μM ATP was significantly more robust as anticipated (**Fig. 3f**). The minor

detection of IκBα phosphorylation in absence of ADP in the radioactive assay and lack of it in

the cold immunoblotting assay (compare lanes marked with arrows in Figs. 3b and 3f) may

indicate higher sensitivity of the radio assay. The need of ADP in phosphotransfer prompted

us to consider whether the ADP was contaminated with a trace amount of ATP. ESI-MS

analysis of 50 μM ADP indicated no peak corresponding to ATP (**Supplementary Data Fig.**

3c).

We performed another phospho-transfer assay that bypasses the need for purification

of P-IKK2 from unreacted ATP. In this assay condition ATP should be totally depleted within

10-20 min as no detectable enhancement of Tyr phosphorylation level was observed after that

(**Supplementary Data Fig. 3d**). Substrate was added to the reaction post-20, 30, and 40 min

ATP treatment at 30°C. We observed phosphorylation of the substrate, indicating transfer of

phosphate from P-IKK2. Additionally, substrate phosphorylation reduced significantly at 40

min preincubation set perhaps hinting at transient nature of transferable phosphates, and its

loss prior to addition of the substrate (**Supplementary Data Fig. 3d**). The results of these

experiments allowed us to conclude that transiently auto-phosphorylated tyrosine residues are

capable of transferring phosphate to the substrate.

Phosphotransfer from P-IKK2 is critical to fidelity and specificity of IκBα phosphorylation:

We next tested if phosphotransfer is restricted only to the N-terminal signal responsive S32/S36 or to serines of PEST domain as well. Radioactive kinase assay as described earlier was performed with full-length WT and mutant S32A,S36A (AA) IκBα as the phosphate recipient. Phosphate group(s) of P-IKK2 was not transferred to full-length IκBα AA which retains the serines and threonines within its PEST domain (**Fig. 4a**), indicating phosphotransfer only to the N-terminal serines of IκBα. We also tested the effect of AMPPNP, an unhydrolyzable analogue of ATP, on the phosphotransfer, and found no inhibitory effect irrespective of the presence of ADP. This suggests adenine nucleotides poising the kinase for phosphotransfer (**Fig. 4a**).

During our investigation, we noted unusual behavior in the IKK2-K44M mutant. IKK2-K44M phosphorylated full-length WT IκBα robustly but with much less efficiency than WT IKK2. IKK2-K44M also showed a minor auto-phosphorylation activity compared to WT IKK2 in the absence of substrate (**Fig. 4b**). We performed a similar kinase assay using full-length WT IκBα where cold ATP was used as the phosphate donor, and phosphorylation status of IκBα at S32,S36 was monitored by immunoblotting. It revealed that IKK2-K44M is unable to phosphorylate S32/S36 (**Fig. 4c**). Since in radioactive kinase assay we observed pan-phosphorylation of the substrate (**Fig. 4b**), it possibly reflects phosphorylation of the PEST region of FL IκBα. We additionally compared IKK2-K44M with WT IKK2 using radioactive kinase assay with both FL IκBα and FL IκBαAA substrates in the absence or presence of Inhibitor VII. As anticipated, FL IκBα was phosphorylated much more efficiently than FL IκBαAA by WT IKK2, and phosphorylation was practically abolished in the presence of 10 μM Inhibitor VII (**Fig. 4d**). In contrast, IKK2-K44M phosphorylated FL IκBαAA more efficiently than FL IκBα, and reduction in the presence of inhibitor VII was strikingly less (**Fig. 4d**). It indicates that IKK2-K44M, despite retaining non-specific kinase activity, lost phosphorylation activity towards S32,S36 of the *bona fide* substrate IκBα. It is noteworthy, an equivalent K to M

mutation in Erk2 also didn't abolish the basal activity, but rather retained ~5 % of it[43].

Discussion:

Phosphorylation of substrates by protein kinases is regulated in various ways. IKK2, a kinase

central to inflammation, phosphorylates serine residues of its primary substrate IκBα with γ-

phosphate ATP. Here, we report multiple intriguing properties of IKK2 that provide specificity

to its substrate phosphorylation. These include: 1) autophosphorylation of tyrosine residues

of IKK2 in cis, in addition to known auto-phosphorylation at AL serines *i.e.,* dual-specificity; 2)

relay of phosphate to substrate IκBα from auto-phosphorylated IKK2 (P-IKK2) either directly

or through ADP; and 3) disruption of a universally conserved salt bridge (mediated by K44)

renders the kinase non-specific.

282   STYK family of eukaryotic protein kinases transfer γ-phosphate of ATP directly to

substrate, in general to S, T and Y residues [44] [41,45-47]. Histidine kinases (in prokaryotes and in

some lower eukaryotes) however shuttles the γ-phosphate to Asp residue on a response

regulator (RR) substrate through a phospho-His intermediate [48,49] [43,50-52]. Phospho-His

functions as a phosphoenzyme intermediate in several metabolic enzymes as well [51]. Our

present analyses of eukaryotic protein kinase IKK2 revealed its surprising property in which

IKK2 phosphorylated at AL-serines (S177/S181) could be further auto-phosphorylated by ATP

at two (Y169 & Y188) or more tyrosine residues (and perhaps other residues) yielding another

active form (P-IKK2) (**Fig. 4e**). This marks IKK2 as an autocatalytic dual-specificity kinase

rather than a prototype S/T kinase. Interestingly, while auto-phosphorylation at AL-serines

(S177 and S181) occurs in *trans* [53], auto-phosphorylation of tyrosine residues occurs in *cis*.

This multisite phosphorylated form of IKK2 (P-IKK2) could transfer some of its phosphate(s)

to substrate IκBα in presence of ADP without requiring ATP, hinting at an unusual property of

IKK2 acting possibly as a phosphate-sink. It is possible that phospho-tyrosines residues of

IKK2 act as this sink. The Y169 located at the DFG+1 (DLG in IKK2) position is unique only

to IKK1 and IKK2, and is absent even in close mammalian homologues such as IKKε and

TBK1. Interestingly, neither IKKε nor TBK1 can phosphorylate IκBα at S32 and S36, *in vitro*

or *in vivo*[54,55]. A recent report also indicated the role of DFG+1 residue in differentiating Ser

versus Thr residues [36]. These observations hinted to an involvement of Y169 in substrate

selectivity. Indeed, our experimental data suggests a pivotal role of Y169 in phosphorylating

S32 of IκBα both *in vitro* and ex-vivo (**Fig. 2g-i**). It is worthwhile to note that, an

unphosphorylatable mutation at S32 in IκBα to Ile (S32I) is associated with ectodermal

dysplasia and T-cell immunodeficiency in patients[56,57]. More interestingly, the placement of

this residue in the kinase active site makes it amenable to a phosphorylation-

dephosphorylation cycle, suggesting its possible role both in phosphotransfer and substrate

specificity. Mutation of Y188, a conserved tyrosine, to unphosphorylatable phenylalanine

abolished the kinase activity of IKK2 *in vitro* as well as in MEFs upon TNF-treatment (**Fig. 2g**

and i). Placement of Y188 indicates that its mutation will be detrimental to the structural fold

of kinase, thus it is unclear if or how Y188 plays a role in the phosphor transfer. Since

phosphotransfer to S32,S36 of IκBα is observed with auto-phosphorylated IKK2 EE,

transferred phosphates are not from S177 or S181.

The strength of input signal, level of IκBα degradation, and NF-κB activation correlate

with the amplitude, duration, and kinetics of IKK activity [58,59]. This digital (all-or-none) activation

profile of NF-κB appears to be due to rapid activation and inactivation of IKK2. IKK targets

multiple substrates *in vivo*; however, the rapid activation of NF-κB is regulated primarily by

IκBα. How IKK specifically selects IκBα for phosphorylation is unclear, although NEMO is

reported to direct the kinase activity of IKK2 specifically towards IκBα [60]. We observe that P-

IKK2 could transfer phosphate(s) specifically to the S32/S36 of IκBα, and not to other sites

(such as PEST sequence of IκBα) (**Fig. 3b, f and 4a**). We surmise if transiency of multisite

phosphorylated IKK2 (P-IKK2) underlie this rapid activation-inactivation events during signal

response. It is possible that regulatory system of IKK2 is somewhat analogous the two-

component histidine kinase-effector systems in prokaryotes that show a threshold activation

response[61 62,63].

All three tyrosines in AL of IKK2 are previously reported to undergo signal-induced

phosphorylation in cells. These modifications were proposed to be mediated by a tyrosine

kinase although there is no evidence of a specific tyrosine kinase [35,64,65]. Several members of

the Ser/Thr kinase family undergo tyrosine phosphorylation and these kinases employ diverse

mechanisms for auto-phosphorylation[66-71].

The phosphotransfer from P-IKK2 to IκBα appears inefficient as no apparent decrease

in phospho-label in P-IKK2 is observed post transfer (**Fig. 3b**). This is in contrast to HKs where

a single phosphate from the phosphorylated HK is stoichiometrically transferred to the cognate

RR. There are also other prominent differences, such as phosphorylation on IKK2 occurs at

multiple sites unlike at single site for the HKs, and not every phosphorylated residue in P-IKK2

expected to transfer its phosphate to IκBα. It is likely that, there are more than one type of

phosphorylated residues in P-IKK2 with respect to their stability wherein only a fraction of the

total pool can be relayed. Another notable difference between phosphorelay in IKK2 from that

in HK is IKK2's dependence on ADP (**Fig. 3b and f**). It was also observed that, the generic

kinase inhibitor AMPPNP (unhydrolyzable ATP analogue) failed to inhibit phosphor-transfer

from IKK2 (**Fig. 4a**) and IκBα phosphorylation (**Supplementary Data Fig. 2f**). Taken together,

we surmise that presence of ADP or AMPPNP help IKK2 attain conformational state poised

for phosphorylating S32/36 of IκBα even in absence of ATP. The list of Ser/Thr kinase family

proteins that undergo tyrosine phosphorylation is growing that employ diverse mechanisms

for auto-phosphorylation[66-71]. The most notable example of this dual specificity is the DYRK.

However, the role of Tyr-phosphorylation in DYRK is not related to that in IKK2, nor does it

render DYRKs capable of relaying phosphates to the substrates. It will be interesting to see

whether phosphotransfer by relay (with or without ADP) occurs in any other kinase-substrate

system other than IKK2: IκBα system described here.

Our study reveals tyrosine auto-phosphorylation in IKK2 and indicates that

consequences of Ser and Tyr autophosphorylations in dual-specific IKK2 are distinct.

Moreover, positions of the critical tyrosine residues are unique, such that some that are

amenable to *cis* auto-phosphorylation are essential for substrate phosphorylation. Most

surprisingly, the tyrosine-linked phosphate groups can be transferred by a relay-mechanism

to specific serines in IκBα, an event reminiscing phospho-relay in two component systems

prevalent in prokaryotes and some lower eukaryotes. Our study also reveals the role of K44

of IKK2 in auto-phosphorylation and thus in specific phosphorylation of S32,S36 of IκBα,

although it does not affect its non-specific kinase activity. It requires further investigation to

determine if a structural state of IKK2, perhaps through a different modification or through

binding to another protein, allow it to phosphorylate substrates relatively non-specifically.

In all, it is intriguing how the gateway to the activation of NF-κB is regulated through

unique phosphorylation events in IKK2 (**Fig. 4f**). It appears that a multi-layered fail-safe

mechanism is devised to tightly control NF-κB activation. Firstly, in absence of upstream

signals IKK-complex is unable to phosphorylate Ser 32,36 of IκBα efficiently thereby keeping

check on the aberrant and untimely activation of NF-κB. Upon proinflammatory cues cells

need to activate NF-κB immediately and efficiently, i.e., IKK needs to be activated and

phosphorylate Ser 32,36 of IκBα. NEMO warrants IKK-activation and ensures that the IKK-

complex specifically chooses IκBα from a large pool of substrates of IKK2. IKK2 is designed

in such a manner that it phosphorylates itself at AL-tyrosines when activated, such that

phosphate group(s) can be relayed directly or through ADP to Ser32,36 of IκBα with great

fidelity, thus leaving little chance of a misconstrued signaling event thereby confirming NF-κB

activation. Our discovery might also present the beginning of a new aspect in eukaryotic cell-

signaling by EPKs and provide a foundation for mechanistic studies of an intriguing phospho-

transfer reaction that helps accomplish the desired specificity. To the best of our knowledge,

it is the first report to strongly suggest that a metazoan EPK harbors HK-like phosphotransfer

activity through relay mechanism to its substrate.

Acknowledgement: We thank Joe Adams and Simpson Joseph (UCSD), Tony Hunter (Salk

Institute) and Alex Hoffmann (UCLA) for commenting on the work. Authors also thank

members of Polley lab (BI) and Ghosh Lab (UCSD) for their continuous support. We also

thank Majid Ghassemian (Biomolecular Mass spectrometry facility, UCSD) and Souvik Roy

(Mass spectrometry facility, BI) for the mass spectrometry experiments and analyses. This

work was supported by an Intermediate Fellowship from the DBT Wellcome Trust India

Alliance (IA/I/15/1/501852) & intramural funding from Bose Institute to SP, NIH grant to GG

and American Cancer Society grant RSG-08-287-01-GMC to TH. Biochemistry research at

SDSU is supported in part by the California Metabolic Research Foundation. PB was

supported by the graduate research fellowship from Bose Institute.

386
387 **Author contribution:** SP, GG and TB designed experiments; PB, SP, TB & GG performed

388 experiments. SP, GG, TB, TH and PB analyzed the results. SP wrote the manuscript with the

389 help of PB and TB in preparing figures and images, GG, TB and TH modified the manuscript.

391 **Ethics declarations:**

392 Competing interests: The authors declare no competing financial interests.

References

Hoffmann, A. & Baltimore, D. Circuitry of nuclear factor kappaB signaling. *Immunol Rev* **210**, 171-186, doi:10.1111/j.0105-2896.2006.00375.x (2006).

DiDonato, J. A., Hayakawa, M., Rothwarf, D. M., Zandi, E. & Karin, M. A cytokine-responsive IkappaB kinase that activates the transcription factor NF-kappaB. *Nature* **388**, 548-554, doi:10.1038/41493 (1997).

Zandi, E., Rothwarf, D. M., Delhase, M., Hayakawa, M. & Karin, M. The IkappaB kinase complex (IKK) contains two kinase subunits, IKKalpha and IKKbeta, necessary for IkappaB phosphorylation and NF-kappaB activation. *Cell* **91**, 243-252 (1997).

Rothwarf, D. M., Zandi, E., Natoli, G. & Karin, M. IKK-gamma is an essential regulatory subunit of the IkappaB kinase complex. *Nature* **395**, 297-300, doi:10.1038/26261 (1998).

Hacker, H. & Karin, M. Regulation and function of IKK and IKK-related kinases. *Sci STKE* **2006**, re13, doi:10.1126/stke.3572006re13 (2006).

Hinz, M. & Scheidereit, C. The IkappaB kinase complex in NF-kappaB regulation and beyond. *EMBO Rep* **15**, 46-61, doi:10.1002/embr.201337983 (2014).

Ghosh, S. & Karin, M. Missing pieces in the NF-kappaB puzzle. *Cell* **109 Suppl**, S81-96 (2002).

Liu, F., Xia, Y., Parker, A. S. & Verma, I. M. IKK biology. *Immunol Rev* **246**, 239-253, doi:10.1111/j.1600-065X.2012.01107.x (2012).

Scheidereit, C. IkappaB kinase complexes: gateways to NF-kappaB activation and transcription. *Oncogene* **25**, 6685-6705, doi:10.1038/sj.onc.1209934 (2006).

Karin, M. & Ben-Neriah, Y. Phosphorylation meets ubiquitination: the control of NF-[kappa]B activity. *Annu Rev Immunol* **18**, 621-663, doi:10.1146/annurev.immunol.18.1.621 (2000).

Hayden, M. S. & Ghosh, S. Shared principles in NF-kappaB signaling. *Cell* **132**, 344-362, doi:10.1016/j.cell.2008.01.020 (2008).

Brown, K., Gerstberger, S., Carlson, L., Franzoso, G. & Siebenlist, U. Control of I kappa B-alpha proteolysis by site-specific, signal-induced phosphorylation. *Science* **267**, 1485-1488, doi:10.1126/science.7878466 (1995).

Lin, Y. C., Brown, K. & Siebenlist, U. Activation of NF-kappa B requires proteolysis of the inhibitor I kappa B-alpha: signal-induced phosphorylation of I kappa B-alpha alone does not release active NF-kappa B. *Proc Natl Acad Sci U S A* **92**, 552-556, doi:10.1073/pnas.92.2.552 (1995).

Manning, G., Whyte, D. B., Martinez, R., Hunter, T. & Sudarsanam, S. The protein kinase complement of the human genome. *Science* **298**, 1912-1934, doi:10.1126/science.1075762 (2002).

Manning, G., Plowman, G. D., Hunter, T. & Sudarsanam, S. Evolution of protein kinase signaling from yeast to man. *Trends Biochem Sci* **27**, 514-520, doi:10.1016/s0968-0004(02)02179-5 (2002).

16   Dhanasekaran, N. & Premkumar Reddy, E. Signaling by dual specificity kinases.
Oncogene **17**, 1447-1455, doi:10.1038/sj.onc.1202251 (1998).

17   He, X., Jones, M. H., Winey, M. & Sazer, S. Mph1, a member of the Mps1-like family
of dual specificity protein kinases, is required for the spindle checkpoint in S. pombe.
J Cell Sci **111 (Pt 12)**, 1635-1647, doi:10.1242/jcs.111.12.1635 (1998).

18   Lauze, E. *et al.* Yeast spindle pole body duplication gene MPS1 encodes an essential
dual specificity protein kinase. *EMBO J* **14**, 1655-1663, doi:10.1002/j.1460-
2075.1995.tb07154.x (1995).

19   Ubersax, J. A. & Ferrell, J. E., Jr. Mechanisms of specificity in protein phosphorylation.
Nat Rev Mol Cell Biol **8**, 530-541, doi:10.1038/nrm2203 (2007).

20   Schrofelbauer, B. & Hoffmann, A. How do pleiotropic kinase hubs mediate specific
signaling by TNFR superfamily members? *Immunol Rev* **244**, 29-43,
doi:10.1111/j.1600-065X.2011.01060.x (2011).

21   Antonia, R. J., Hagan, R. S. & Baldwin, A. S. Expanding the View of IKK: New
Substrates and New Biology. *Trends Cell Biol* **31**, 166-178,
doi:10.1016/j.tcb.2020.12.003 (2021).

22   Huse, M. & Kuriyan, J. The conformational plasticity of protein kinases. *Cell* **109**, 275-
282, doi:10.1016/s0092-8674(02)00741-9 (2002).

23   Liu, S. *et al.* Crystal structure of a human IkappaB kinase beta asymmetric dimer. *J
Biol Chem* **288**, 22758-22767, doi:10.1074/jbc.M113.482596 (2013).

24   Polley, S. *et al.* A structural basis for IkappaB kinase 2 activation via oligomerization-
dependent trans auto-phosphorylation. *PLoS Biol* **11**, e1001581,
doi:10.1371/journal.pbio.1001581 (2013).

25   Xu, G. *et al.* Crystal structure of inhibitor of kappaB kinase beta. *Nature* **472**, 325-330,
doi:10.1038/nature09853 (2011).

26   Zhang, J., Clark, K., Lawrence, T., Peggie, M. W. & Cohen, P. An unexpected twist to
the activation of IKKbeta: TAK1 primes IKKbeta for activation by autophosphorylation.
Biochem J **461**, 531-537, doi:10.1042/BJ20140444 (2014).

27   Du, M., Ea, C. K., Fang, Y. & Chen, Z. J. Liquid phase separation of NEMO induced
by polyubiquitin chains activates NF-kappaB. *Mol Cell* **82**, 2415-2426 e2415,
doi:10.1016/j.molcel.2022.03.037 (2022).

28   Ea, C. K., Deng, L., Xia, Z. P., Pineda, G. & Chen, Z. J. Activation of IKK by TNFalpha
requires site-specific ubiquitination of RIP1 and polyubiquitin binding by NEMO. *Mol
Cell* **22**, 245-257, doi:10.1016/j.molcel.2006.03.026 (2006).

29   Chen, Z. J. Ubiquitination in signaling to and activation of IKK. *Immunol Rev* **246**, 95-
106, doi:10.1111/j.1600-065X.2012.01108.x (2012).

30   Shaul, J. D., Farina, A. & Huxford, T. The human IKKbeta subunit kinase domain
displays CK2-like phosphorylation specificity. *Biochem Biophys Res Commun* **374**,
592-597, doi:10.1016/j.bbrc.2008.07.082 (2008).

31 Barroga, C. F., Stevenson, J. K., Schwarz, E. M. & Verma, I. M. Constitutive
phosphorylation of I kappa B alpha by casein kinase II. *Proc Natl Acad Sci U S A* **92**,
7637-7641, doi:10.1073/pnas.92.17.7637 (1995).

32 Tergaonkar, V., Bottero, V., Ikawa, M., Li, Q. & Verma, I. M. IkappaB kinase-
independent IkappaBalpha degradation pathway: functional NF-kappaB activity and
implications for cancer therapy. *Mol Cell Biol* **23**, 8070-8083 (2003).

33 Zandi, E., Chen, Y. & Karin, M. Direct phosphorylation of IkappaB by IKKalpha and
IKKbeta: discrimination between free and NF-kappaB-bound substrate. *Science* **281**,
1360-1363, doi:10.1126/science.281.5381.1360 (1998).

34 Darwech, I., Otero, J. E., Alhawagri, M. A. & Abu-Amer, Y. Tyrosine phosphorylation
is required for IkappaB kinase-beta (IKKbeta) activation and function in
osteoclastogenesis. *J Biol Chem* **285**, 25522-25530, doi:10.1074/jbc.M110.121533
(2010).

35 Otero, J. E. *et al.* Defective osteoclastogenesis by IKKbeta-null precursors is a result
of receptor activator of NF-kappaB ligand (RANKL)-induced JNK-dependent apoptosis
and impaired differentiation. *J Biol Chem* **283**, 24546-24553,
doi:10.1074/jbc.M800434200 (2008).

36 Chen, C. *et al.* Identification of a major determinant for serine-threonine kinase
phosphoacceptor specificity. *Mol Cell* **53**, 140-147, doi:10.1016/j.molcel.2013.11.013
(2014).

37 Ahuja, L. G., Kornev, A. P., McClendon, C. L., Veglia, G. & Taylor, S. S. Mutation of a
kinase allosteric node uncouples dynamics linked to phosphotransfer. *Proc Natl Acad
Sci U S A* **114**, E931-E940, doi:10.1073/pnas.1620667114 (2017).

38 Yang, J. *et al.* Allosteric network of cAMP-dependent protein kinase revealed by
mutation of Tyr204 in the P+1 loop. *J Mol Biol* **346**, 191-201,
doi:10.1016/j.jmb.2004.11.030 (2005).

39 Eckhart, W., Hutchinson, M. A. & Hunter, T. An activity phosphorylating tyrosine in
polyoma T antigen immunoprecipitates. *Cell* **18**, 925-933, doi:10.1016/0092-
8674(79)90205-8 (1979).

40 Knighton, D. R. *et al.* Structure of a peptide inhibitor bound to the catalytic subunit of
cyclic adenosine monophosphate-dependent protein kinase. *Science* **253**, 414-420,
doi:10.1126/science.1862343 (1991).

41 Nolen, B., Taylor, S. & Ghosh, G. Regulation of protein kinases; controlling activity
through activation segment conformation. *Mol Cell* **15**, 661-675,
doi:10.1016/j.molcel.2004.08.024 (2004).

42 Wang, J. *et al.* The crystal structure of a constitutively active mutant RON kinase
suggests an intramolecular autophosphorylation hypothesis. *Biochemistry* **49**, 7972-
7974, doi:10.1021/bi100409w (2010).

43 Robbins, D. J. *et al.* Regulation and properties of extracellular signal-regulated protein
kinases 1 and 2 in vitro. *J Biol Chem* **268**, 5097-5106 (1993).

44 Hunter, T. Protein kinase classification. *Methods Enzymol* **200**, 3-37,
doi:10.1016/0076-6879(91)00125-g (1991).

45   Cohen, P. The origins of protein phosphorylation. *Nat Cell Biol* **4**, E127-130,
515   doi:10.1038/ncb0502-e127 (2002).

516   46   Fischer, E. H. & Krebs, E. G. Conversion of phosphorylase b to phosphorylase a in
517   muscle extracts. *J Biol Chem* **216**, 121-132 (1955).

518   47   Krebs, E. G. & Fischer, E. H. Phosphorylase activity of skeletal muscle extracts. *J Biol*
519   *Chem* **216**, 113-120 (1955).

520   48   Borkovich, K. A. & Simon, M. I. The dynamics of protein phosphorylation in bacterial
521   chemotaxis. *Cell* **63**, 1339-1348, doi:10.1016/0092-8674(90)90429-i (1990).

522   49   Gao, R. & Stock, A. M. Biological insights from structures of two-component proteins.
523   *Annu Rev Microbiol* **63**, 133-154, doi:10.1146/annurev.micro.091208.073214 (2009).

524   50   Hunter, T. A journey from phosphotyrosine to phosphohistidine and beyond. *Mol Cell*
525   **82**, 2190-2200, doi:10.1016/j.molcel.2022.05.007 (2022).

526   51   Kalagiri, R. & Hunter, T. The many ways that nature has exploited the unusual
527   structural and chemical properties of phosphohistidine for use in proteins. *Biochem J*
528   **478**, 3575-3596, doi:10.1042/BCJ20210533 (2021).

529   52   Laub, M. T., Biondi, E. G. & Skerker, J. M. Phosphotransfer profiling: systematic
530   mapping of two-component signal transduction pathways and phosphorelays.
531   *Methods Enzymol* **423**, 531-548, doi:10.1016/S0076-6879(07)23026-5 (2007).

532   53   Pike, A. C. *et al.* Activation segment dimerization: a mechanism for kinase
533   autophosphorylation of non-consensus sites. *EMBO J* **27**, 704-714,
534   doi:10.1038/emboj.2008.8 (2008).

535   54   Larabi, A. *et al.* Crystal structure and mechanism of activation of TANK-binding kinase
536   1. *Cell Rep* **3**, 734-746, doi:10.1016/j.celrep.2013.01.034 (2013).

537   55   Tu, D. *et al.* Structure and ubiquitination-dependent activation of TANK-binding kinase
538   1. *Cell Rep* **3**, 747-758, doi:10.1016/j.celrep.2013.01.033 (2013).

539   56   Mooster, J. L. *et al.* Defective lymphoid organogenesis underlies the immune
540   deficiency caused by a heterozygous S32I mutation in IkappaBalpha. *J Exp Med* **212**,
541   185-202, doi:10.1084/jem.20140979 (2015).

542   57   Courtois, G. *et al.* A hypermorphic IkappaBalpha mutation is associated with
543   autosomal dominant anhidrotic ectodermal dysplasia and T cell immunodeficiency. *J*
544   *Clin Invest* **112**, 1108-1115, doi:10.1172/JCI18714 (2003).

545   58   Cheong, R. *et al.* Transient IkappaB kinase activity mediates temporal NF-kappaB
546   dynamics in response to a wide range of tumor necrosis factor-alpha doses. *J Biol*
547   *Chem* **281**, 2945-2950, doi:10.1074/jbc.M510085200 (2006).

548   59   Behar, M. & Hoffmann, A. Tunable signal processing through a kinase control cycle:
549   the IKK signaling node. *Biophys J* **105**, 231-241, doi:10.1016/j.bpj.2013.05.013 (2013).

550   60   Schrofelbauer, B., Polley, S., Behar, M., Ghosh, G. & Hoffmann, A. NEMO ensures
551   signaling specificity of the pleiotropic IKKbeta by directing its kinase activity toward
552   IkappaBalpha. *Mol Cell* **47**, 111-121, doi:10.1016/j.molcel.2012.04.020 (2012).

61 Lamarche, M. G., Wanner, B. L., Crepin, S. & Harel, J. The phosphate regulon and
bacterial virulence: a regulatory network connecting phosphate homeostasis and
pathogenesis. *FEMS Microbiol Rev* **32**, 461-473, doi:10.1111/j.1574-
6976.2008.00101.x (2008).

62 Bhate, M. P., Molnar, K. S., Goulian, M. & DeGrado, W. F. Signal transduction in
histidine kinases: insights from new structures. *Structure* **23**, 981-994,
doi:10.1016/j.str.2015.04.002 (2015).

63 Laub, M. T. & Goulian, M. Specificity in two-component signal transduction pathways.
Annu Rev Genet **41**, 121-145, doi:10.1146/annurev.genet.41.042007.170548 (2007).

64 Rieke, C., Papendieck, A., Sokolova, O. & Naumann, M. Helicobacter pylori-induced
tyrosine phosphorylation of IKKbeta contributes to NF-kappaB activation. *Biol Chem*
392, 387-393, doi:10.1515/BC.2011.029 (2011).

65 Meyer, A. N. *et al.* Tyrosine phosphorylation allows integration of multiple signaling
inputs by IKKbeta. *PLoS One* **8**, e84497, doi:10.1371/journal.pone.0084497 (2013).

66 Lochhead, P. A. Protein kinase activation loop autophosphorylation in cis: overcoming
a Catch-22 situation. *Sci Signal* **2**, pe4, doi:10.1126/scisignal.254pe4 (2009).

67 Lochhead, P. A. *et al.* A chaperone-dependent GSK3beta transitional intermediate
mediates activation-loop autophosphorylation. *Mol Cell* **24**, 627-633,
doi:10.1016/j.molcel.2006.10.009 (2006).

68 Lochhead, P. A., Sibbet, G., Morrice, N. & Cleghon, V. Activation-loop
autophosphorylation is mediated by a novel transitional intermediate form of DYRKs.
Cell **121**, 925-936, doi:10.1016/j.cell.2005.03.034 (2005).

69 Bhattacharyya, R. P. *et al.* The Ste5 scaffold allosterically modulates signaling output
of the yeast mating pathway. *Science* **311**, 822-826, doi:10.1126/science.1120941
(2006).

70 Tigno-Aranjuez, J. T., Asara, J. M. & Abbott, D. W. Inhibition of RIP2's tyrosine kinase
activity limits NOD2-driven cytokine responses. *Genes Dev* **24**, 2666-2677,
doi:10.1101/gad.1964410 (2010).

71 Ge, B. *et al.* MAPKK-independent activation of p38alpha mediated by TAB1-
dependent autophosphorylation of p38alpha. *Science* **295**, 1291-1294,
doi:10.1126/science.1067289 (2002).
584
585

Figure 1

Figure 1: Autophosphorylation of IKK2 at hitherto uncharacterized sites. a) Domain organisation of FL IKK2 depicting functional kinase domain (KD), ubiquitin-like domain (ULD), and scaffold dimerization domain (SDD) based on the X-ray structures. Known sites of Ser phosphorylation and NEMO interacting regions at the extreme C-termini are highlighted. Serine and tyrosine residues in the activation loop are shown, mutation of which to Glu renders IKK2 constitutively active. The conserved ATP-interacting Lys (at position 44) is also shown. **b)** Autophosphorylation of FL IKK2WT upon incubation with radioactive ATP for different time periods in an *in vitro* kinase assay. **c)** *In vitro* radioactive kinase assays performed to check the effect of NEMO on autophosphorylation of FL IKK2WT (left panel), and that of NEMO and IκBα on autophosphorylation and substrate phosphorylation (right panel) activity of FL IKK2WT. **d)** Effect of Inhibitor VII on FL IKK2WT autophosphorylation and IκBα substrate phosphorylation at different inhibitor concentrations by in a radioactive *in vitro* kinase assay. **e)** Radioactive kinase assay conducted to check auto- and substrate phosphorylation of full length and deletion constructs of constitutively active form of IKK2 harbouring

Figure 2

Figure 2: IKK2 possesses autocatalytic dual specificity. a) *In vitro* kinase assay followed by immunoblotting with FL IKK2WT against phospho Ser (177/181) and phospho-Tyr antibodies. **b)** Effect of Inhibitor VII on tyrosine autophosphorylation of FL IKK2WT. **c)** Immunoblotting performed with phospho-Tyr antibody to check autophosphorylation of IKK2 mutant K44M at different time points compared to that of IKK2 WT. **d)** Tyrosine auto phosphorylation and substrate phosphorylation of GST (1-54) IκBα in presence of full length and deletion mutants of IKK2 harbouring phosphomimetic Ser177Glu and Ser181Glu mutations. **e)** *De novo* Auto-phosphorylation of IKK2 on tyrosine monitored by phospho-Ser and phospho-Tyr specific monoclonal antibodies. **f)** *In vitro* kinase assay followed by immunoblotting done showing active IKK2 cannot phosphorylate tyrosines of an inactive (K44M) IKK2 mutant; however, can efficiently phosphorylate AL serines. **g)** Radioactive *in vitro* kinase assay performed with IKK2 Y169F and Y188F mutants using GST (1-54) IκBα WT and AA as substrates. **h)** Radioactive *in vitro* kinase assay performed with IKK2 Y169F mutant with a range of GST (1-54) IκBα mutants where the substrate phosphorylation signal is abolished in case of S36 A/E IκBα. **i)** Severe reduction in TNFα-induced IKK activity in *ikk2-/-* MEF-3T3 cells reconstituted with mutant (Y169F and Y188F) IKK2s compared to wild-type. Kinase assay was performed with IKK immunoprecipitated (IP-ed) with anti-NEMO antibody from whole cell extract.

Figure 3

Figure 3: Freshly autophosphorylated IKK2 relays phosphates to IκBα. a) Schematic representation of the experiment performed to monitor the path traversed by phosphate(s) from p-IKK2 to substrate by autoradiography detection method. **b)** Purified autophosphorylated (radiolabelled) IKK2 transfers its phosphate to IκBα substrate in absence of any nucleotide, and transfer efficiency is enhanced upon addition of ADP or ATP. **c)** A close-up view of the active site in active form in the context of the entire kinase domain. Three tyrosines (Y169, Y188 and Y199) in immediate vicinity of active sites are highlighted. Position of ATP is modelled based on the structure of PKA. **d)** Schematic representation of the experiment performed to monitor the path traversed by phosphate(s) from p-IKK2 to substrate by immunoblotting detection method. **e)** Elution profile of the size exclusion chromatography run of p-IKK2 to get rid of excess unlabelled ATP performed in Superdex200 10/30 increase. **f)** Purified autophosphorylated (cold) IKK2 transfers its phosphate to IκBα substrate with enhanced transfer efficiency upon addition of ADP or ATP.

Figure 4

Figure 4: Specificity and fidelity of phosphorelay by IKK2 to IκBα. a) AMP-PNP can support efficient phosphorelay. Phospho-relay reaction is observed in IκBα WT but not in S32,36Ala double mutant. Domain organization and position of relevant S/T/Y residues of IκBα are shown above the autoradiograph. **b)** Comparison of kinase activity of FL IKK2WT and K44M mutant phosphorylation signal compared to that of IKK2 WT in radioactive kinase assay performed with FL IκBα WT as the substrate. **c)** Residue level specificity of phosphorylation by the FL IKK2K44M was analysed using an antibody that specifically phosphor-S32/36 of IκBα. **d)** Differential sensitivity of specific vs. non-specific phosphorylation by IKK2 as assessed by the effect of Inhibitor VII on FL IKK2 WT and IKK2 K44M using IκBα WT and AA as substrates. **e)** A detailed scheme of reactions showing IKK2-activation and subsequent phosphorylation of the signal responsive S32,36 on IκBα by phosphorelay. **f)** A cartoon representation of the proposed model of NF-kB activation by IKK2 employing phosphorelay to IκBα.

Figure S1

a

Sample	T_m (°C)	ΔH (Kcal/mol)
IKK2+ADP	40.1	164.0
IKK2+ATP	50.8	251.9

Supplementary Figure 1: a, Autoradiograph of *in vitro* kinase assay reactions carried out with FL IKK2WT and K44M using GST (1-54) IκBα as the substrate as a function of time. **b,** A differential scanning calorimetry (DSC) thermogram showing a striking enhancement in folding stability of IKK2 upon ATP-treatment.

Supplementary Figure 2: a, Effect of increasing concentration of urea on tyrosine autophosphorylation of FL IKK2WT assessed in an *in vitro* kinase assay. **b,** Radioactive *in vitro* kinase assay performed with increasing concentration of urea up to 6M to check auto- and substrate phosphorylation. **c,** Mass spectrometry analysis of ATP-treated IKK2 showing phosphorylation of Tyr169. **d,** Mass spectrometry analysis of ATP-treated IKK2 showing phosphorylation of Tyr188. **e,** Alignment of the activation segment sequences of different IKK family kinases. Conservation of Tyr at the DFG+1 (DLG in case of IKK1 and IKK2) is retained only in a stress response related plant kinase SnRK2, but not in other IKK homologues or any known dual specificity kinases, e.g., DYRK and GSK3β. **f,** Inhibition of both substrate phosphorylation and tyrosine autophosphorylation activities of IKK2 checked in presence of common kinase inhibitors such as AMPPNP and Staurosporine, highly specific IKK2 inhibitors TPCA, Calbiochem Inhibitor VII and MLN120B.

Figure S3

Supplementary Figure 3: a, Substrate bound PKA in its active form; phosphorylatable serine of the substrate peptide (in magenta) is highlighted. **b,** Hydroxyl of Tyr1238 in the activation loop of RON kinase, which is activated through trans auto-phosphorylation, projects towards ATP similarly to that of Tyr169 -OH in IKK2. **c,** ESI-MS scan of the ADP used in this assay system showing negligible trace, if any, of ATP at 50µM ADP concentration. **d,** WB analysis showing auto-phosphorylation of full length IKK2 and phospho-transfer to IκBα.

Supplementary information:

Methods:

Reagents and Cell culture: Monoclonal mouse anti-IκBα (L35A5) (catalogue number: 4814),

mouse anti-Phospho-IκBα (Ser$^{32/36}$) (5A5) (catalogue number:9246) and rabbit anti-Phospho-

IKKα/β (Ser$^{176/180}$) (16A6) (catalogue number: 2697) were from Cell Signaling Technology

(CST). Monoclonal mouse anti-IKKβ (10AG2) (catalogue number: NB100-56509) was from

Novus Biologicals. Polyclonal rabbit anti-6XHis (catalogue number: BB-AB0010) was from

BioBharti Life Science. Monoclonal mouse anti-Phospho-Tyrosine (catalogue number:

610000) was from BD Biosciences. Polyclonal rabbit anti-GST (catalogue number: 924801),

monoclonal mouse anti-HA 11 epitope (catalogue number: 901502) and mouse anti-Tubulinβ3

(catalogue number: 657402) were from BioLegend. Polyclonal rabbit anti-IKKα/β (H-470):

sc7607 was from Santa Cruz Biotechnology (has been discontinued) and mouse anti-

Phospho-Serine Q5 (catalogue number: 37430) was from Qiagen. Horseradish peroxidase-

conjugated secondary antibodies against rabbit (catalogue number: 31460) and mouse IgG

(catalogue number: 31430) were Pierce antibodies from Thermo Scientific. Adenosine-5'-

Triphosphate (ATP), Disodium trihydrate (catalogue number: A-081-25) was from Gold

Biotechnology and Adenosine Diphosphate (ADP) (Reference number: V916A) provided in

ADP-Glo^{TM} Kinase assay (catalogue number: V6930) from Promega. Recombinant murine

TNF-α used was from Roche.

Sf9 cells used for expressing proteins with baculovirus system was procured from Gibco™

Sf9 cells in Sf-900™ III SFM, Thermo Scientific (catalogue number: 12659017) and

maintained in Gibco™Sf-900™ III SFM, Thermo Scientific (catalogue number: 12658019).

Protein expression and purification:

IKK2 (WT and other constructs): Wild-type and mutant IKK2s were purified following a

published protocol (Polley et al, 2013). Sf9 cells (Gibco™ Sf9 cells in Sf-900™ III SFM) at a

density of 1.5-2 millions/ml were infected with high titre viral stock (generally after P3 stage)

at a ratio of 1:100 or 1:50 as optimized for higher expression level as judged by a pilot

experiment performed previously, and grown for 48 hours upon infection. Cells were harvested

and lysed by sonication in a buffer containing 25 mM Tris-HCl pH 8.0, 200 mM NaCl, 10%

Glycerol, 5 mM β-Mercaptoethanol (β-ME), 10 mM Imidazole, 20 mM β-glycerophosphate, 10

mM NaF, and 1 mM sodium orthovanadate and 1X Protease Inhibitor Cocktail (PIC, Sigma).

The lysate was clarified by centrifugation at 20,000 x g for 45 minutes at 4°C and incubated

with Ni-NTA agarose resin (Qiagen, catalogue number: 30230) pre-equilibrated with lysis

buffer for 2 hours at 4°C. The resin was subsequently washed with lysis buffer containing

30mM Imidazole. Protein was eluted with lysis buffer containing 250mM Imidazole. Protein

fractions were analysed on SDS-PAGE, pooled together and either flash frozen or subjected

to size-exclusion chromatography on Superdex 200 Increase 10/300 GL (GE Healthcare)

column equilibrated with 20mM Tris-HCl pH 8.0, 200mM NaCl, 10% Glycerol and 2 mM DTT.

To make untagged IKK2, protein fractions eluted from the Ni-NTA resin of desired purity was

treated with TEV-protease at a ratio of 50:1 (weight by weight, IKK2: TEV) in the cold room for

more than 10 hours after addition of 1 mM of EDTA, the resulting protein mix was then loaded

on to the SEC as described above. To prepare autophosphorylated IKK2, the fractions with

desired purity level were treated with 0.5-1 mM of ATP,10 mM of $MgCl_2$ and phosphatase

inhibitors and the autophosphorylation reaction was carried out at 27°C for 1 hour. Resulting

protein solution was then concentrated in a YM30-concentrator and loaded onto Superdex

200 Increase 10/300 GL column. Elution fractions from the size-exclusion step were analysed

on SDS-PAGE and peak fractions of desired quality (GE Healthcare) were pooled together

and concentrated using a YM30 Centriprep or Centricon devices depending upon the volume

of the sample and flash frozen immediately.

IκBα (WT, mutant and other constructs): His-tagged full length and GST-tagged deletion

(1-54) constructs of IκBα (both the wild-type and Ser32Ala, Ser36Ala, Ser32,36Ala single and

double-mutant versions) were expressed in *E. coli* Rosetta2 (DE3). The cells were grown

overnight at 16°C after 0.3 mM IPTG induction. The cell pellet was resuspended in 40 ml of

Lysis Buffer (25 mM Tris pH 7.5, 200 mM NaCl, 10 mM imidazole, 10% Glycerol, 5 mM β-

mercaptoethanol, and 1 mM PMSF). The cells were lysed by using sonicator keeping the cell

suspension chilled on ice. The lysate was clarified by centrifugation at 15,000 x g for one hour

at 4°C. About 3 ml of Ni-NTA Agarose resin was equilibrated with lysis buffer, and the clarified

lysate (supernatant) was mixed with it later. The protein was allowed to bind the resin by

incubating for 2 hours on a rotary mixer in a 4°C room. The resin was washed with the lysis

buffer containing 500 mM NaCl (high salt wash) for about 10 CV to clear out non-specific

proteins from binding. Then the resin was thoroughly washed with the lysis buffer containing

20 mM imidazole. His-tagged IκBα protein was eluted under gravity flow, using elution buffer

(the lysis buffer containing 250 mM imidazole). ~1 ml volume of each fraction was collected.

(For GST tagged proteins, Glutathione Sepharose resin (GE Healthcare, catalogue number:

17-5279-01) was used and elution was performed with the help of reduced glutathione). The

main elution fractions were subjected to size-exclusion chromatography (SEC) on a HiLoad

16/600 pg Superdex 200 column (GE Healthcare) and the gel run revealed cleaner protein

fractions. The peak fractions were saved and stored at -80°C.

NEMO (WT, mutant and other constructs): NEMO FL, NEMO double mutant C76A C95S

(site directed mutagenesis done in pET24d vector) and NEMO (44-196) (truncates in pET24d)

were expressed in *E. coli* Rosetta2 DE3 cells. The cells were grown overnight at ~21°C after

0.25 mM IPTG induction (at 0.3-0.4 O.D.) and 50-100 μM Zn^{2+}. [OR The cells were grown

overnight (~17 hours) at 16°C after 0.1 mM IPTG induction (at 0.2-0.3 O.D.) and 50-100 μM

Zn^{2+}]. The cell pellet was resuspended in 40 ml of Lysis Buffer (25 mM Tris pH 7.5, 200 mM

NaCl, 10 mM imidazole, 10% Glycerol, 5 mM β-mercaptoethanol, 20μM $ZnCl_2$, 0.5 M urea

(acting like an osmolyte). Rest of the steps performed were typical to a Ni-NTA purification as

described above. The main elution fractions were subjected to size-exclusion chromatography

on a HiLoad 16/600 pg Superdex 200 column (GE Healthcare) and the cleaner peak fractions

were stored at -80°C.

Kinase Assay with purified proteins: Typically, 100-200 ng (unless mentioned otherwise)

of purified kinase was used in each *in vitro* kinase reaction. 0.5-1µg of IκBα substrate (full-

length or GST-tagged 1-54) was incubated with kinase for 30 mins or indicated time periods

at 27˚C in presence of 20µM ATP with 0.1µCi of γ-P32-ATP in case of radioactive assay or

50-100 µM ATP in case of Western blot assay in a reaction buffer containing 20mM HEPES

pH 7.5, 100mM NaCl, 15mM $MgCl_2$, 2mM DTT, 0.2mM Na_3VO_4, 10mM NaF, 20mM β-

Glycerophosphate. Reaction was stopped by heating the samples at 95˚C for 5 mins after the

addition of 1X-Laemmli buffer. Samples were run on 8 or 10% SDS-PAGE. For radioactive

assays, the gel was then dried and analysed either by film or phosphor-imaging. For Western

blot assays, proteins from gel were electro-transferred onto nitrocellulose/PVDF membrane

and probed with desired antibodies. Phosphorylated tyrosine and serine residues were

detected using monoclonal antibodies against phospho-tyrosine and phospho-serine.

IP-Kinase assay: Immortalized $ikk2^{-/-}$ mouse embryonic fibroblast 3T3 cells were

reconstituted by retroviral transduction with HA-tagged WT or mutant pBABE-IKK2 constructs.

Stable expression of IKK2 was first confirmed by immunoblotting. Cells were then stimulated

with recombinant murine TNF-α (Roche) at 70-80% cell confluency. IKK complex was

immunoprecipitated from whole cell extracts using a NEMO-specific antibody (BD) and Protein

A agarose (Upstate) for 2 hrs in the cold room. Whole cell extract used was pre-cleared by

incubation with Protein A agarose bead for an hour. Immunoprecipitated complex was washed

extensively, including a final washing step with the kinase assay buffer devoid of ATP and

incubated with GST-IκBα (1-54) and 20 µM ATP and 5 µCi of γ32P-ATP for 30 mins at 30˚C

in kinase assays. Reactions were stopped by addition of 2X-Laemmli SDS buffer to samples

and heating at 95˚C for 5 mins. Reaction products were resolved on a 12% SDS PAGE. Extent

of substrate phosphorylation was measured by cutting out the appropriate area of the gel

containing the substrate and subjecting it to phosphor-imaging (Typhoon, GE Life Sciences)

after drying. As a control, amount of IKK2 in each reaction was estimated by transferring IKK2

containing gel area onto nitrocellulose membrane and probing it with anti-IKK1/2 antibody.

Sample preparation for LC-MS/MS analyses for Tyrosine phosphorylation detection:

Protein samples were diluted in TNE (50mM Tris pH 8.0, 100mM NaCl, 1mM EDTA) buffer. RapiGest SF reagent (Waters Corp.) was added to the mix to a final concentration of 0.1% and samples were boiled for 5 min. Tris (2-carboxyethyl) phosphine (TCEP) was added to a final concentration of 1mM and the samples were incubated at 37˚C for 30 min. Subsequently, the samples were carboxymethylated with 0.5 mg/ml of iodoacetamide for 30 min at 37˚C followed by neutralization with 2mM final concentration of TCEP. Proteins samples prepared as above were digested with trypsin (trypsin: protein ratio - 1:50) overnight at 37˚C. RapiGest was degraded and removed by treating the samples with 250 mM HCl at 37˚C for 1 h followed by centrifugation at 14000 rpm for 30 min at 4˚C. The soluble fraction was then added to a new tube and the peptides were extracted and desalted using C18 desalting columns (Thermo Scientific).

LC-MS/MS analysis: Trypsin-digested peptides were analysed by ultra-high-pressure liquid chromatography (UPLC) coupled with tandem mass spectroscopy (LC-MS/MS) using nano-spray ionization. The nano-spray ionization experiments were performed using a TripleTOF 5600 hybrid mass spectrometer (ABSCIEX) interfaced with nano-scale reversed-phase UPLC (Waters corporation nano ACQUITY) using a 20 cm-75-micron ID glass capillary packed with 2.5 µm C18 (130) CSHTM beads (Waters corporation). Peptides were eluted from the C18 column into the mass spectrometer using a linear gradient (5–80%) of Acetonitrile (ACN) at a flow rate of 250 µl/min for 1h. The buffers used to create the ACN gradient were: Buffer A (98% H_2O, 2% ACN, 0.1% formic acid, and 0.005% TFA) and Buffer B (100% ACN, 0.1% formic acid, and 0.005% TFA). MS/MS data were acquired in a data-dependent manner in which the MS1 data was acquired for 250ms at m/z of 400 to 1250 Da and the MS/MS data was acquired from m/z of 50 to 2,000Da. The Independent data acquisition (IDA) parameters were as follows; MS1-TOF acquisition time of 250 milliseconds, followed by 50 MS2 events of 48 milliseconds acquisition time for each event. The threshold to trigger MS2 event was set to 150 counts when the ion had the charge state +2, +3 and +4. The ion exclusion time was set

to 4 seconds. Finally, the collected data were analysed using Protein Pilot 4.5 (ABSCIEX) for

peptide identifications.

ESI-MS analyses of ADP: 50 µM of ADP solution was prepared to be analysed by

Electrospray Ionisation Mass Spectrometry (ESI-MS). It was performed in a Xevo G2-XS QToF

mass spectrometer (Waters Corporation) using capillary at 3kV, with voltage parameters of

sampling cone at 40V and source offset at 80V. The source temperature was set at 100˚C and

desolvation temperature at 40˚C. The gas flow rates were as follows: Cone gas flow rate at 50

litre/hr and desolvation gas flow rate at 400 litre/hr. The sample was passed at 5 µl/min flow

rate. The MS data was acquired for a mass range of 50-2000 m/z and the acquisition time was

1 minute.

Radioactive Phosphotransfer assay: Phosphotransfer assays were performed in a wide

range of concentration of IKK2, IκBα and ATP. Typically, 10 µM purified IKK2 protein was first

autophosphorylated in presence of 100-200 nM of γ32P-ATP (PerkinElmer) in kinase assay

buffer for 1 hour at 30˚C. 70 µl of the resulting reaction mixture was first passed through a

Micro Bio-Spin Bio-Gel P-30 spin column (Bio-Rad; 40 kDa MWCO) equilibrated with 20 mM

HEPES pH 7.9, 100 mM NaCl, 0.2 mM Na_3VO_4, 10 mM NaF, 20 mM β-Glycerophosphate, 2

mM DTT, 2 mM EDTA and 1X protease inhibitor cocktail. Eluted protein was subjected to

another round of buffer exchange on a Bio-Gel P-30 column equilibrated with the same buffer

mentioned above devoid of EDTA and containing 15 mM $MgCl_2$. Phosphorylated IKK2 thus

obtained was used in phosphotransfer reaction. 2-3 µM of Phosphorylated IKK2 was

incubated with 10-20 µM of WT or mutant IκBα protein at 30˚C in kinase assay buffer in a total

volume of 20 µl for different time periods as indicated in Figure 3b and 4a. The reactions were

stopped by addition of 10 µl of 4X Laemmli SDS-buffer containing 100 mM EDTA to samples

and heating at 95˚C for 5 mins. Different concentration of ADP or ATP was added to the assay

to assess the role of these nucleotides in phosphotransfer efficiency. Routinely, 10-15µl of

reaction products were resolved on 10% SDS PAGE and radiolabelled samples were

monitored by phosphor imaging of the dried gel on a Typhoon scanner. For the

phosphotransfer reaction described in Figure 4a, 10 µM purified IKK2 was autophosphorylated

with of 100 nM of γ^{32}P-ATP in presence of 40 µM of cold ATP.

Phospho-transfer assay with cold ATP: About 10 µM of purified FL IKK2WT was treated

with 100 µM of cold ATP in kinase assay buffer (containing 1X protease inhibitor cocktail) for

one hour at 30°C. The resulting reaction mixture was then loaded onto a Superdex 200

increase 10/300 GL (24ml bed volume, Cytiva) SEC column to purify autophosphorylated FL

IKK2WT separated from excess ATP. SEC was performed in buffer containing 20 mM HEPES,

100 mM NaCl, 20 mM β-glycerophosphate, 10 mM NaF, 0.2 mM Na3VO4 and 2 mM DTT.

The elution fractions containing phosphorylated IKK2 were incubated with FL IκBαWT in

presence of different concentrations of ATP or ADP at different time points (as mentioned in

Figure 3f) at 30°C. The reactions were stopped by adding 4X Laemmli SDS dye, heated at

95°C for 5 minutes and were run on a 10% SDS-PAGE. This was followed by immunoblotting

and detection was done using monoclonal antibodies against phospho-tyrosine and phospho

Ser$^{32/36}$ IκBα.

184