Phylogenetic placement of *Ceratophyllum submersum* based on a complete plastome sequence derived from nanopore long read sequencing data

Samuel Nestor Meckoni[1,†], Benneth Nass[1,2,†] & Boas Pucker[1,3]*

[1] Plant Biotechnology and Bioinformatics, Institute of Plant Biology, TU Braunschweig, 38106 Braunschweig, Germany

[2] Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Kamýcká 129, Praha 6 - Suchdol, CZ-165 21, Czech Republic

[3] Braunschweig Integrated Centre of Systems Biology (BRICS), TU Braunschweig, 38106 Braunschweig, Germany

[†] Even contribution

* Corresponding author

Emails: s.meckoni@tu-bs.de, nassb@fzp.czu.cz, b.pucker@tu-braunschweig.de

Abstract

Objective

18 Eutrophication poses a mounting concern in today's world. *Ceratophyllum submersum* is a
19 plant capable of living in eutrophic conditions, therefore it could play a critical role in addressing
20 the problem of eutrophication. This study aimed to take a look at the genetic basis underlying
21 *C. submersum*.

Results

23 Sequencing of gDNA from *C. submersum* yielded enough reads to assemble a plastome.
24 Subsequent annotation and phylogenetic analysis validated existing information regarding
25 angiosperm relationships and the positioning of Ceratophylalles in a wider phylogenetic
26 context.

27
28 Keywords: ONT, angiosperms, Ceratophyllaceae, Ceratophyllales plastome, plastomics,
29 chloroplast-genome phylogenomics, phylogenetic pipeline, supermatrix tree

Introduction

Ceratophyllum submersum, commonly known as soft hornwort, is a subaquatic plant whose
genus is the only extant member in the order of Ceratophyllales, placed as a sister clade to
the eudicots [1,2]. It is native to Europe, Africa and Asia and grows in stagnant freshwater
bodies [3]. Morphological features include long, branching stems that can reach up to several
metres in length and leaves that are forked into narrow, filament-like segments that grow in
multiple whorls around the stem (Figure 1) [4]. The plant is often green, but can vary in colour
from brownish to reddish depending on environmental conditions. Anthocyanins contribute to
the colouration of many plant species, and metabolic analyses have detected several

derivatives in *C. submersum* [5–7]. It thrives in eutrophic conditions, characterised by low light

intensity and high nutrient levels [8,9]. Eutrophication of aquatic environments is indicated by

the accumulation of nutrients albeit other parameters and multiple classification systems exist

[10]. Due to anthropogenic effects the occurrence of eutrophic environments is rising which

poses a problem e.g. for greenhouse gas emissions [11]. In aquatic systems, eutrophication

is inducing harmful algal blooms (HABs), which are responsible for environmental hazards like

the Oder ecological disaster in 2022 [12]. *C. submersum* has the ability to enhance the water

quality by absorbing nutrients (especially nitrogen and phosphor) and is therefore popular in

amateur aquacultures. It can also prevent the formation of HABs, even though it suffers from

them [7].

Figure 1: *Ceratophyllum submersum* and its habitat. A: The pond (Braunschweig,

52.28062 N / 10.54896 E) where *C. submersum* was collected, **B**: Whole *C. submersum* plant,

C: Close up view of *C. submersum*.

Further research of *C. submersum* can help to develop ecosystem engineering solutions to

combat human-induced eutrophication and HABs in freshwaters. Here, we used the potential

of nanopore long read sequencing to unravel the plastome sequence of *C. submersum* and

provide a comprehensive annotation.

57 Main Text

58 Results and Discussion

A total of 1,544 long reads representing the plastome were extracted from the raw read

sequencing data (Additional file 1, PRJEB62706). From that, 752 reads were passed to the

Flye assembler by ptGAUL to reach a maximum coverage depth of 50x. The plastome

assembly of *C. submersum* resulted in a 155,767 bp long sequence with a GC content of

38.26%. In total, 75 unique protein coding genes were annotated (Additional file 2). The

C. submersum plastome sequence is 485 bp shorter than the reference plastome sequence

of *C. demersum* that has a length of 156,252 bp [13]. While the GC content is similar

(*C. demersum*: 38.22, *C. submersum*: 38.26) the amount of unique annotated genes in

C. submersum is smaller than in *C. demersum* (75 against 79). After the annotation some

predicted coding sequences were flawed (e.g. containing multiple stop codons). Manual

evaluation revealed four homopolymeric regions in which a frameshift would correct the

annotation. Since homopolymers are a frequent error type in ONT reads, manual correction in

those regions is appropriate [14]. The comparison of those regions to all plastome reads

suggested a correction of two of these regions, namely the adenine at position 3,094 and the

thymines at position 3,143, at position 86,261, and at position 86,262 were inserted.

To validate the plastome assembly from *C. submersum*, a phylogenetic tree was calculated

(Figure 2).

Figure 2: Phylogenetic tree of selected spermatophytes based on plastome protein sequences. The asterisks (*) indicate plastome assemblies generated in this study. Within the angiosperms, the orders are differently coloured. The eudicotyledons are the only exception as they are represented by several orders. All nodes received full bootstrapping support (100%), except those displaying the actual value. Please see the material and methods section for further description of the tree calculation. Visualisation was done in iTOL 6.7.6 [15]. The full tree including the outgroup *Chlamydomonas reinhardtii* can be found in the additional files (Additional file 4). GS = gymnosperms. The sketches are licensed by adobe and depositphotos (Standard License: https://wwwimages2.adobe.com/content/dam/cc/en/legal/servicetou/Stock-Additional-Terms_en_US_20221205.pdf, https://depositphotos.com/license.html).

Multiple plastome reference sequences were chosen to represent the angiosperm clade as well as *Chlamydomonas reinhardtii* as outgroup (for full list and references see Additional file

3). Additionally, our own plastome assembly of *Digitalis purpurea* was included as a control,

because gDNA from this plant was also sequenced on the same flow cell as the gDNA from

C. submersum (*D. purpurea* data: PRJEB62706). The phylogenetic tree (figure 2, for full tree

see Additional file 4) classifies *C. submersum* close to their reference *C. demersum* and our

D. purpurea plastome assembly close to the RefSeq *D. purpurea* plastome from Zhao *et al.*,

2023 [16]. The angiosperm clade is represented in accordance with the current APG IV

classification except for the exact separation/placement of the Magnoliids and the

Chloranthales, which is still controversial [2]. This underlines the significance of plastome

sequences for modern plant phylogenetics [17].

The plastome assembly and phylogenetic analysis presented in this study provide initial

insights into the genetic basis of *Ceratophyllum submersum*. Further research is needed to

unravel its nuclear genome and to explore potential human applications of this plant, such as

its use as a valuable resource or as an agent to mitigate environmental hazards.

105 Material and Methods

106 Plant Material, gDNA extraction and sequencing

Ceratophyllum submersum was collected from a small pond in Braunschweig (52.28062 N /

10.54896 E) and kept in cultivation. The artificial pond needs occasional water replenishment.

Foliage from surrounding shrubs and dead aquatic plants lead to a high eutrophy (see figure

1A). *C. submersum* shoot tips (figure 1C) were harvested for gDNA extraction conducted with

a CTAB method [18–20]. Short DNA fragments were depleted with the Short Read Eliminator

kit (Pacific Biosciences). Library preparation for ONT sequencing was started with 1µg of DNA

that was first repaired with the NEBNext□ Companion Module and then processed according

to the SQK-LSK109 protocol (Oxford Nanopore Technologies). For ONT sequencing, a R9.4.1

flow cell was used with a MinION. Prior to *C. submersum* sequencing, the flow cell was already

utilised for *Digitalis purpurea* gDNA sequencing. Processing of raw data was performed with

guppy v6.4.6+ae70e8f (https://community.nanoporetech.com) and minimap2 v2.24-r1122 [21]

on a graphical processor unit (GPU) in the de.NBI cloud to generate FASTQ files.

Plastome Assembly and Annotation

The FASTQ files were subjected to a plastome assembly with ptGAUL using standard

parameters [21–24]. As reference the *C. demersum* plastome from the NCBI RefSeq database

(release 216) was used [13,25]. Since the ptGAUL results consist of two assemblies (one per

path) the OGDRAW plastome maps were compared between the reference and the

assemblies to decide which one is closer to the reference. The *C. submersum* assembly

needed to be reverse complemented and the sequence start was adjusted to the *C. demersum*

sequence. GC contents of the assembly were calculated using the contig_stats.py script [26].

For annotation GeSeq was applied (see Additional file 5 for parameters) [27–36]. Protein

coding genes were extracted from the resulting GBSON.json file using our own script

JSON_2_peptide_fasta_v1.0.py [37]. Manual evaluation of wrong annotated CDSs was

conducted with the help of EMBOSS Sixpack to have an overview of all reading frames which

could then be compared to the results from NCBI blastx to find possible wrong frameshifts

[38,39]. Further evaluation of these frameshifts was conducted with the help of a read

mapping. All plastome reads were mapped against the plastome sequence with minimap2

v2.24-r1122 [21]. Subfurther generation of a bam file and its indexing with samtools 1.10 [40]

was conducted to be able to analyse the mapping in the integrative genomics viewer (IGV)

2.16.1 [41]. Assembly corrections were only then conducted, if a significant number of reads

support it.

Data preparation prior to submission was performed by extracting all the read IDs from the

'new_filter_gt3000.fa', generated by ptGAUL, which contains all the identified plastome reads.

The plastome reads were extracted from the FAST5 and FASTQ raw read datasets via the

ont_fast5_api tool and a custom script [37,42].

Phylogenetic analysis

Reference data retrieval and phylogenetic supermatrix tree construction was performed by our newly developed Python pipeline PAPAplastomes (Pipeline for the Automatic Phylogenetic Analysis of plastomes) which integrates established external tools for complex steps (Additional file 6) [43]. Reference species, closely related species of interest, and outgroup species can be specified via a config file. First, NCBI RefSeq plastome data is downloaded and the reference plastome peptide sequences are extracted from this collection. These peptide sequences are further combined with the peptide sequences derived from the assemblies representing plastomes of interest. The pre-OrthoFinder trimming step removes potential paralogs with the exact same sequence, sequences which contain asterisks and sequences that are shorter than 10 amino acids (this threshold value is adjustable). Next, OrthoFinder v2.5.4 [44–47] is applied. Post-Orthofinder processing includes four steps: (1) removal of outlier sequences, (2) deletion of orthogroups missing the species of interest or their references, (3) removal of orthogroups harbouring fewer species than the pre-defined outgroup species, (4) paralog cleaning. (1) Since the OrthoFinder results include phylogenetic trees of each orthogroup, outlier identification was conducted with the help of the Python module Dendropy v4.5.2 [48]. Per orthogroup the edge length for each taxon is accessed except for outgroup species. Based on these lengths, outliers are identified by the 1.5*IQR method (inter quartile range) i.e. sequences with a distance larger than 1.5 times the variation are excluded. (2) The species of an orthogroup are listed and if pre-defined species are all either present or absent the orthogroup will be kept. Otherwise, the orthogroup will be discarded. This is suggested for closely related species i.e. from the same genus and can be specified by the user in the config file. (3) Orthogroups consisting solely of the outgroup species are exempt from this criterion. (4) From remaining paralogs within one species, only the longest sequence is kept.

The cleaned orthogroups are then aligned with MAFFT v7.453 (--maxiterate 1000 --localpair) [49,50] and alignments are concatenated. Phylogenetic supermatrix tree calculation is

169   performed by IQ-TREE (multicore version 1.6.12 for Linux 64-bit built Aug 15 2019; with the '-

170   nt AUTO -bb 1000' options, seed: 291752) based on the concatenated alignment of all

171   orthogroups [51–53].

172   Limitations

173   Before sequencing *C. submersum*, the flow cell had already been utilised for sequencing

174   gDNA of *D. purpurea*. Insufficient DNA availability hindered the complete sequencing and

175   assembly of *C. submersum*'s nuclear genome. Future optimisations in DNA extraction

176   methods dedicated to small aquatic plants could overcome this limitation.

177   Abbreviations

CDS - coding sequence

gDNA - genomic DNA

GPU - graphical processing unit

GS - gymnosperm

HAB - harmful algal bloom

RefSeq - NCBI Reference Sequence Database

IQR - interquartile range

ONT - Oxford Nanopore Technologies

186   References

1.   Les DH. The Origin and Affinities of the Ceratophyllaceae. Taxon. 1988

May;37(2):326–45.

2.   The Angiosperm Phylogeny Group, Chase MW, Christenhusz MJM, Fay MF, Byng JW,

Judd WS, et al. An update of the Angiosperm Phylogeny Group classification for the

orders and families of flowering plants: APG IV. Bot J Linn Soc. 2016 May 1;181(1):1–

20.

3. POWO. Plants of the World Online. Facilitated by the Royal Botanic Gardens, Kew.

[Internet]. Plants of the World Online. 2023 [cited 2023 Apr 16]. Available from:

http://powo.science.kew.org/taxon/urn:lsid:ipni.org:names:163088-1

4. Düll R, Kutzelnigg H. Taschenlexikon der Pflanzen Deutschlands und angrenzender

Länder: die häufigsten mitteleuropäischen Arten im Porträt. 7th ed. Wiebelsheim:

Quelle & Meyer; 2011. 932 p. (Quelle & Meyer Taschenlexikon).

5. Les DH. Systematics and Evolution of *Ceratophyllum* L. (Ceratophyllaceae): A

Monograph (Taxonomy, Flavonoids, Aquatic plants, Phytogeography, Variation).

[Columbus]: The Ohio State University; 1986.

6. Les DH. Ceratophyllaceae. In: Kubitzki K, Rohwer JG, Bittrich V, editors. Flowering

Plants · Dicotyledons [Internet]. Berlin, Heidelberg: Springer Berlin Heidelberg; 1993

[cited 2023 Apr 16]. p. 246–50. Available from: http://link.springer.com/10.1007/978-3-

662-02899-5_24

7. Ujvárosi AZ, Riba M, Garda T, Gyémánt G, Vereb G, M-Hamvas M, et al. Attack of

Microcystis aeruginosa bloom on a *Ceratophyllum submersum* field: Ecotoxicological

measurements in real environment with real microcystin exposure. Sci Total Environ.

2019 Apr;662:735–45.

8. Zartes Hornblatt (*Ceratophyllum submersum*) [Internet]. [cited 2023 Apr 5]. Available

from: https://www.oekologie-seite.de/index.php?id=24&pid=642

9. Ellenberg H. Vegetation Mitteleuropas mit den Alpen. 6., vollständig neu bearbeitete

und stark erweiterte Auflage. Leuschner C, Dierschke H, editors. Verlag Eugen Ulmer;

2010.

10. Ayele HS, Atlabachew M. Review of characterization, factors, impacts, and solutions of

Lake eutrophication: lesson for lake Tana, Ethiopia. Environ Sci Pollut Res. 2021

Mar;28(12):14233–52.

11. Li Y, Shang J, Zhang C, Zhang W, Niu L, Wang L, et al. The role of freshwater

eutrophication in greenhouse gas emissions: A review. Sci Total Environ. 2021

May;768:144582.

12. European Commission: Joint Research Centre. An EU analysis of the ecological

disaster in the Oder River of 2022: lessons learned and research based

recommendations to avoid future ecological damage in EU rivers, a joint analysis from

DG ENV, JRC and the EEA. [Internet]. Publications Office of the European Union; 2023

[cited 2023 Apr 16]. Available from: https://data.europa.eu/doi/10.2760/067386

13. Moore MJ, Bell CD, Soltis PS, Soltis DE. Using plastid genome-scale data to resolve

enigmatic relationships among basal angiosperms. Proc Natl Acad Sci. 2007 Dec

4;104(49):19363–8.

14. Delahaye C, Nicolas J. Sequencing DNA with nanopores: Troubles and biases. PLOS

ONE. 2021 Oct 1;16(10):e0257521.

15. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree

display and annotation. Nucleic Acids Res. 2021 Jul 2;49(W1):W293–6.

16. Zhao F, Liu B, Liu S, Min DZ, Zhang T, Cai J, et al. Disentangling a 40-year-old

taxonomic puzzle: the phylogenetic position of *Mimulicalyx* (Lamiales). Bot J Linn Soc.

2023 Jan 25;201(2):135–53.

17. Gitzendanner MA, Soltis PS, Yi TS, Li DZ, Soltis DE. Plastome Phylogenetics: 30

Years of Inferences Into Plant Evolution. In: Advances in Botanical Research [Internet].

Elsevier; 2018 [cited 2022 Aug 9]. p. 293–313. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0065229617300885

18. Pucker B. Plant DNA extraction and preparation for ONT sequencing. 2020 Mar 21

[cited 2023 Feb 14]; Available from: https://dx.doi.org/10.17504/protocols.io.bcvyiw7w

19. Siadjeu C, Pucker B, Viehöver P, Albach DC, Weisshaar B. High Contiguity de novo

Genome Sequence Assembly of Trifoliate Yam (*Dioscorea dumetorum*) Using Long

Read Sequencing. Genes. 2020 Mar;11(3):274.

20. Pucker B, Rückert C, Stracke R, Viehöver P, Kalinowski J, Weisshaar B. Twenty-Five

Years of Propagation in Suspension Cell Culture Results in Substantial Alterations of

the Arabidopsis Thaliana Genome. Genes. 2019 Sep;10(9):671.

21. Li H. Minimap2: pairwise alignment for nucleotide sequences. Birol I, editor.

Bioinformatics. 2018 Sep 15;34(18):3094–100.

22. Zhou W, Armijos C, Lee C, Lu R, Wang J, Ruhlman T, et al. Plastid Genome Assembly

Using Long-read Data (ptGAUL) [Internet]. bioRxiv; 2022 [cited 2022 Nov 23]. p.

2022.11.19.517194. Available from:

https://www.biorxiv.org/content/10.1101/2022.11.19.517194v1

23. Shen W, Le S, Li Y, Hu F. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q

File Manipulation. Zou Q, editor. PLOS ONE. 2016 Oct 5;11(10):e0163962.

24. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using

repeat graphs. Nat Biotechnol. 2019 May;37(5):540–6.

25. O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, et al. Reference

sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and

functional annotation. Nucleic Acids Res. 2016 Jan 4;44(D1):D733–45.

26. Pucker B, Holtgräwe D, Rosleff Sörensen T, Stracke R, Viehöver P, Weisshaar B. A De

Novo Genome Sequence Assembly of the *Arabidopsis thaliana* Accession Niederzenz-

1 Displays Presence/Absence Variation and Strong Synteny. Vandepoele K, editor.

PLOS ONE. 2016 Oct 6;11(10):e0164321.

27. Tillich M, Lehwark P, Pellizzer T, Ulbricht-Jones ES, Fischer A, Bock R, et al. GeSeq –

versatile and accurate annotation of organelle genomes. Nucleic Acids Res. 2017 Jul

3;45(W1):W6–11.

28. Laslett D. ARAGORN, a program to detect tRNA genes and tmRNA genes in

nucleotide sequences. Nucleic Acids Res. 2004 Jan 2;32(1):11–6.

29. Kent WJ. BLAT —The BLAST -Like Alignment Tool. Genome Res. 2002 Apr

1;12(4):656–64.

30. Ian Small, Ian Castleden. Chloë: Organelle Annotator [Internet]. 2020. Available from:

https://github.com/ian-small/chloe

31. Nawrocki EP, Eddy SR. Infernal 1.1: 100-fold faster RNA homology searches.

Bioinformatics. 2013 Nov 15;29(22):2933–5.

32. Wheeler TJ, Eddy SR. nhmmer: DNA homology search with profile HMMs.

Bioinformatics. 2013 Oct 1;29(19):2487–9.

33. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res. 2004 Mar 1;32(5):1792–7.

34. Greiner S, Lehwark P, Bock R. OrganellarGenomeDRAW (OGDRAW) version 1.3.1:

expanded toolkit for the graphical visualization of organellar genomes. Nucleic Acids

Res. 2019 Jul 2;47(W1):W59–64.

35. Abascal F, Zardoya R, Telford MJ. TranslatorX: multiple alignment of nucleotide

sequences guided by amino acid translations. Nucleic Acids Res. 2010 Jul

1;38(suppl_2):W7–13.

36. Chan PP, Lin BY, Mak AJ, Lowe TM. tRNAscan-SE 2.0: improved detection and

functional classification of transfer RNA genes. Nucleic Acids Res. 2021 Sep

20;49(16):9077–96.

37. Meckoni SN. Small scripts codeberg repository [Internet]. 2023 [cited 2023 May 30].

Available from: https://codeberg.org/snmeckoni/scripts

38. Madeira F, Pearce M, Tivey ARN, Basutkar P, Lee J, Edbali O, et al. Search and

sequence analysis tools services from EMBL-EBI in 2022. Nucleic Acids Res. 2022 Jul

1;50(W1):W276–9.

39. NCBI blastx [Internet]. Bethesda (MD): National Library of Medicine (US), National

Center for Biotechnology Information; [cited 2023 Jun 8]. Available from:

https://blast.ncbi.nlm.nih.gov/Blast.cgi?PROGRAM=blastx&PAGE_TYPE=BlastSearch

&LINK_LOC=blasthome

40. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics. 2009 Aug 15;25(16):2078–9.

41. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al.

Integrative genomics viewer. Nat Biotechnol. 2011 Jan;29(1):24–6.

42. ont_fast5_api github repository [Internet]. nanoporetech; 2022 [cited 2023 May 30].

Available from: https://github.com/nanoporetech/ont_fast5_api

43.   Meckoni SN. PAPAplastomes [Internet]. 2023 [cited 2023 Jun 26]. Available from:

https://codeberg.org/snmeckoni/PAPAplastomes

44.   Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative

genomics. Genome Biol. 2019 Dec;20(1):238.

45.   Emms DM, Kelly S. OrthoFinder: solving fundamental biases in whole genome

comparisons dramatically improves orthogroup inference accuracy. Genome Biol. 2015

Dec;16(1):157.

46.   Emms DM, Kelly S. STRIDE: Species Tree Root Inference from Gene Duplication

Events. Mol Biol Evol. 2017 Dec 1;34(12):3267–78.

47.   Emms DM, Kelly S. STAG: Species Tree Inference from All Genes [Internet].

Evolutionary Biology; 2018 Feb [cited 2023 Mar 20]. Available from:

http://biorxiv.org/lookup/doi/10.1101/267914

48.   Sukumaran J, Holder MT. DendroPy: a Python library for phylogenetic computing.

Bioinformatics. 2010 Jun 15;26(12):1569–71.

49.   Katoh K. MAFFT: a novel method for rapid multiple sequence alignment based on fast

Fourier transform. Nucleic Acids Res. 2002 Jul 15;30(14):3059–66.

50.   Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. Mol Biol Evol. 2013 Apr 1;30(4):772–80.

51.   Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and Effective

Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol Biol Evol.

2015 Jan;32(1):268–74.

52.   Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder:

fast model selection for accurate phylogenetic estimates. Nat Methods. 2017

Jun;14(6):587–9.

53.   Minh BQ, Nguyen MAT, von Haeseler A. Ultrafast Approximation for Phylogenetic

Bootstrap. Mol Biol Evol. 2013 May 1;30(5):1188–95.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and material

The datasets generated and analysed during the current study are available under

PRJEB62706. Scripts developed for the data analysis are available from Codeberg

(https://codeberg.org/snmeckoni/scripts, https://codeberg.org/snmeckoni/PAPAplastomes)

and GitHub (https://github.com/bpucker/GenomeAssembly).

Competing interests

The authors declare that they have no conflict of interest.

Funding

This work was supported by the BMBF-funded de.NBI Cloud within the German Network for

Bioinformatics Infrastructure (de.NBI) (031A532B, 031A533A, 031A533B, 031A534A,

031A535A, 031A537A, 031A537B, 031A537C, 031A537D, 031A538A).

Author's contributions

All authors contributed to the concept and design of the studies, wet and dry lab work and manuscript writing.

Acknowledgments

We thank Dr. Christiane Evers from the Institute for Plant Biology at the TU Braunschweig for excellent support in taxonomically classifying several specimens.

Legends

Additional file 1: FASTQ stats of the *Ceratophyllum submersum* plastome reads. The stats were calculated with the help of the script FASTQ_stats3.py (https://github.com/bpucker/GenomeAssembly).

Additional file 2: *Ceratophyllum submersum* plastome map derived from OGDRAW.

Additional file 3: List of references of the reference plastomes. The data was extracted by PAPAplastomes and originates from the RefSeq database.

Additional file 4: Full phylogenetic tree with outgroup species *Chlamydomonas reinhardtii*.

Additional file 5: Exact parameters for the GeSeq run. Some parameters from the online tool GeSeq consist of a checkbox that has to be checked. Therefore, the value "check" is stated.

Additional file 6: PAPAplastome workflow chart.