Eukaryotic antiviral immune proteins arose via convergence, horizontal transfer, and ancient inheritance

Edward M. Culbertson[a] and Tera C. Levin*[a]

[a] University of Pittsburgh, Department of Biological Sciences

* Address correspondence to Tera C. Levin: teralevin@pitt.edu

Abstract

Animals use a variety of cell-autonomous innate immune proteins to detect viral infections and prevent replication. Recent studies have discovered that a subset of mammalian antiviral proteins have homology to anti-phage defense proteins in bacteria, implying that there are aspects of innate immunity that are shared across the Tree of Life. While the majority of these studies have focused on characterizing the diversity and biochemical functions of the bacterial proteins, the evolutionary relationships between animal and bacterial proteins are less clear. This ambiguity is partly due to the long evolutionary distances separating animal and bacterial proteins, which obscures their relationships. Here, we tackle this problem for three innate immune families (CD-NTases [including cGAS], STINGs, and Viperins) by deeply sampling protein diversity across eukaryotes. We find that Viperins and OAS family CD-NTases are truly ancient immune proteins, likely inherited since the last eukaryotic common ancestor and possibly longer. In contrast, we find other immune proteins that arose via at least four independent events of horizontal gene transfer (HGT) from bacteria. Two of these events allowed algae to acquire new bacterial viperins, while two more HGT events gave rise to distinct superfamilies of eukaryotic CD-NTases: the Mab21 superfamily (containing cGAS) which has diversified via a series of animal-specific duplications, and a previously undefined eSMODS superfamily, which more closely resembles bacterial CD-NTases. Finally, we found that cGAS and STING proteins have substantially different histories, with STINGs arising via convergent domain shuffling in bacteria and eukaryotes. Overall, our findings paint a picture of eukaryotic innate immunity as highly dynamic, where eukaryotes build upon their ancient antiviral repertoires through the reuse of protein domains and by repeatedly sampling a rich reservoir of bacterial anti-phage genes.

Introduction

As the first line of defense against pathogens, all forms of life rely on cell-autonomous innate immunity to recognize threats and respond with countermeasures. Until recently, many components of innate immunity were thought to be lineage-specific[1]. However, new studies have revealed that an ever-growing number of proteins used in mammalian antiviral immunity are homologous to bacterial immune proteins used to fight off bacteriophage infections. This list includes Argonaute, CARD domains, cGAS and other CD-NTases, Death-like domains, Gasdermin, NACHT domains, STING, SamHD1, TRADD-N domains, TIR domains, and Viperin, among others[2–13]. This discovery has been surprising and exciting, as it implies that some cellular defenses have deep commonalities spanning across the entire Tree of Life. But despite significant homology, these bacterial and animal immune proteins are often distinct in their molecular functions and operate within dramatically different signaling pathways (reviewed here[5]). *How, then, have animals and other eukaryotes acquired these immune proteins?*

One common hypothesis in the field is that these immune proteins are ancient, and have been inherited since the last common ancestor of bacteria and eukaryotes[5]. In other cases, horizontal gene transfer (HGT) between bacteria and eukaryotes has been invoked to explain the similarities[6,14]. However, because most papers in this field have focused on searching

genomic databases for new bacterial immune genes and biochemically characterizing them, the
evolution of these proteins in eukaryotes has not been as thoroughly investigated.
To address this knowledge gap, we turned to the EukProt database, which has been
specifically developed to reflect the true scope of eukaryotic diversity through the genomes and
transcriptomes of nearly 1,000 species, specifically selected to span the eukaryotic tree [15].
EukProt contains sequences from NCBI and Ensemble, plus many diverged eukaryotic species
not found in any other database, making it a unique resource for eukaryotic diversity[15]. While
it can be challenging to acquire diverse eukaryotic sequences from traditional databases due to
an overrepresentation of metazoan data[16], EukProt ameliorates this bias by downsampling
traditionally overrepresented taxa.
Using this database, we investigated the ancestry of three gene families that are shared
between animal and bacterial immunity: Stimulator of Interferon Gamma (STING), cyclic GMP-
AMP synthase (cGAS) and its broader family of cGAS-DncV-like nucleotidyltransferases (CD-
NTases), and Viperin. STING, CD-NTases, and Viperin are all interferon-stimulated genes that
function as antiviral immune modules, disrupting the viral life cycle by activating downstream
immune genes, sensing viral infection, or disrupting viral processes, respectively[17]. We found
eukaryotic CD-NTases arose following multiple HGT events between bacteria and eukaryotes.
cGAS falls within a unique, mainly metazoan clade. In contrast, OAS-like proteins were
independently acquired and are the predominant type of CD-NTase found across most
eukaryotes. Separately, we have discovered diverged eukaryotic STING proteins that bridge the
evolutionary gap between metazoan and bacterial STINGs, as well as two separate instances
where bacteria and eukaryotes have acquired similar proteins via convergent domain shuffling.
Finally, we find that Viperin is likely to be truly ancient, with both broad representation across
the eukaryotic tree of life and evidence of two additional HGT events where eukaryotes recently
acquired new bacterial viperins. Overall, our results demonstrate that immune proteins shared
between bacteria and eukaryotes are evolutionarily dynamic, with eukaryotes taking multiple
routes to acquire and deploy these ancient immune modules.

74 Results

Discovering immune homologs across the eukaryotic tree of life
The first step to understanding the evolution of CD-NTases, STINGs, and viperins was
to acquire sequences for these proteins from across the eukaryotic tree. To search for diverse
immune homologs, we employed a hidden Markov model (HMM) strategy, which has high
sensitivity, a low number of false positives, and the ability to separately analyze multiple
(potentially independently evolving) domains in the same protein[18–20]. To broaden our
searches from initial animal homologs to eukaryotic sequences more generally, we used
iterative HMM searches of the EukProt database, incorporating the hits from each search into
the subsequent HMM. After using this approach to create pan-eukaryotic HMMs for each protein
family, we then added in bacterial homologs to generate universal HMMs (Fig. 1A and Supp.
Fig. 1), continuing our iterative searches until we either failed to find any new protein sequences
or began finding proteins outside of the family of interest (Supp. Fig. 1).
Our searches for CD-NTases, STINGs, and viperins recovered hundreds of eukaryotic
proteins from each family, including a particularly large number of metazoan sequences (red

bars, Fig. 1B). It is not surprising that we found so many metazoan homologs, as each of these
proteins was discovered and characterized in metazoans and these animal genomes tend to be
of higher quality than other taxa (Supp. Fig 2). We also recovered homologs from other species
spread across the Eukaryotic tree, demonstrating that our approach could successfully identify
deeply diverged homologs (Fig. 1B). However, outside of Metazoa, these homologs were
sparsely distributed, such that for most species in our dataset (711/993), we did not recover
proteins from the three immune families examined (white space, lack of colored bars, Fig. 1B).
We believe this pattern reflects a pattern of ongoing, repeated gene losses across eukaryotes,
as has been found for other innate immune proteins[21–23] and other types of gene families
surveyed across eukaryotes[22,24–26]. We found that BUSCO completeness scores and data
type (genomes vs. transcriptomes) were insufficient to explain the pattern of gene loss (Supp.
Fig. 2). Thus, although it is always possible that our approach has missed some homologs, we
believe the resulting data represents a fair assessment of the true diversity across eukaryotes.
To analyze these genes, we aligned the homologs with MAFFT and MUSCLE and then
generated phylogenetic trees with IQtree, FastTree, and RaxML-ng (see Materials and
Methods). We considered our results to be robust if they were concordant across the majority of
six trees generated per gene.
106
107   **Eukaryotes acquired CD-NTases from bacteria through at least three independent HGT**
108   **events**
109   We next studied the evolution of the innate immune proteins, beginning with cGAS and
its broader family of CD-NTase enzymes, which generate diverse oligonucleotides. In addition
to the well-studied cGAS, a number of other eukaryotic CD-NTases have been previously
described: 2'-5'-Oligoadenylate Synthetase 1/2/3 (OAS1/2/3), Male abnormal 21-Like 1/2/3/4
(MAB21L1/2/3/4), Mab-21 domain containing protein 2 (MB21D2), Mitochondrial dynamics
protein 49/51(MID49/51), and Inositol 1,4,5 triphosphate receptor-interacting protein 1/2
(ITPRILP/1/2)[27]. Of these, cGAS and OAS1 are the best characterized and both play roles in
immune signaling. cGAS, Mab21L1, and MB21D2 are all cGAS-like receptors (cGLRs), and
recent work has shown that cGLRs are present in nearly all metazoan taxa and generate
diverse cyclic dinucleotide signals[28]. However, the immune functions of Mab21L1 and
MB21D2 remain unclear, although they have been shown to be important for development[29–
31].
121   Following infections or cellular damage, cGAS binds cytosolic DNA and generates cyclic
GMP-AMP (cGAMP)[32–35], which then activates downstream immune responses via STING
[34,36–38]. OAS1 synthesizes 2',5'-oligoadenylates which bind and activate Ribonuclease L
(RNase L)[39]. Activated RNase L is a potent endoribonuclease that degrades both host and
viral RNA species, reducing viral replication (reviewed here[40,41]). Some bacterial CD-NTases
such as *DncV* behave similar to animal cGAS; they are activated by phage infection and
produce cGAMP[8,42,43]. Other bacterial CD-NTases generate a wide variety of dinucleotides,
cyclic trinucleotides, and cyclic oligonucleotides[11]. These CD-NTases are commonly found
within cyclic oligonucleotide-based anti-phage signaling systems (CBASS) across many
bacterial phyla and even archaea[8,27,43].
131   To understand the evolutionary history of CD-NTases we used the Pfam domain
PF03281 as a eukaryotic starting point. As representative bacterial CD-NTases, we used 6,132

bacterial sequences, representing a wide swath of CD-NTase diversity[43]. Following our
iterative HMM searches, we recovered 313 sequences from 109 eukaryotes, of which 34 were
metazoans (Supplemental Data and Fig. 1B). Most eukaryotic sequences clustered into one of
two distinct superfamilies, which we name here for their highest scoring PFAM domain: Mab21
(Mab21: PF03281) or OAS (OAS1-C: PF10421) (Fig. 2A). Bacterial CD-NTases typically had
sequences matching the HMM for the Second Messenger Oligonucleotide or Dinucleotide
Synthetase domain (SMODS: PF18144).
140   The Mab21 superfamily is composed almost entirely of metazoan sequences, with only a
few homologs from Amoebozoa, choanoflagellates, and other eukaryotes (Fig. 2A). Indeed, the
majority of animal CD-NTases (cGAS, Mid51, Mab21, Mab21L1/2/3/4, Mb21d2, ITPRI) are
paralogs within the Mab21 superfamily, which arose from repeated animal-specific
duplications[44] (Supp. Fig 6). In contrast, unlike the animal-dominated Mab21 superfamily, the
OAS superfamily spans a broad group of eukaryotic taxa, with OAS-like homologs present in
8/12 eukaryotic supergroups. This distribution makes OAS proteins the most common CD-
NTases found across eukaryotes and implies that they arose very early in eukaryotic history,
possibly within the last eukaryotic common ancestor (LECA).
149   Given the connections between cGAS and STING in both animals and some
bacteria[3,43,45], we asked whether species that encode STING also have Mab21 and/or OAS
proteins. Because the Mab21 superfamily is largely animal-specific, we performed this analysis
separately in either Metazoa or with all non-metazoan eukaryotes (Fig. 2B). In animal species
where we found a STING homolog, we also typically found Mab21 superfamily sequence
(32/34), and a cGAS homolog in (26/34) species (Fig. 2B), consistent with the consensus that
these proteins are functionally linked. We also observed 19 metazoan species that had a
Mab21-like sequence with no detectable STING homolog. Almost half of these species (10/19)
were arthropods, agreeing with prior findings of STING sparseness among arthropods[45].
Outside of animals, we found that species with a STING homolog typically did not have a
detectable CD-NTase protein from either superfamily (22/34). While it remains possible that
these STING proteins function together with a to-be-discovered CD-NTase that was absent from
our dataset, we therefore hypothesize that many eukaryotes outside of metazoans and their
close relatives[46] use STING and CD-NTase homologs independently of each other.
163   What was the evolutionary origin of eukaryotic CD-NTases? Interestingly, the Mab21
and OAS superfamilies are only distantly related to one another. Each lies nested within a
different, previously defined, bacterial CD-NTase clade (Fig. 2 C and D). The OAS superfamily
falls within bacterial Clade C (with the closest related bacterial CD-NTases being those of
subclade C02-C03, Fig. 2C), while the metazoan Mab21 superfamily lies within bacterial Clade
D (subclade D12) (Fig. 2D). We note that in this tree (Fig. 2D), Clade D does not form a single
coherent clade, as was also true in the phylogeny that originally defined the bacterial CD-NTase
clades [11].
171   We also observed a number of eukaryotic sequences scattered across different bacterial
CD-NTase clades (Fig. 2A, colored branches within gray clades). While some of these may
reflect additional HGT events, others may come from technical artifacts such as bacterial
contamination of eukaryotic sequences. To minimize such false positive HGT calls, we took a
conservative approach in our analyses, considering potential bacteria-eukaryote HGT events to
be trustworthy only if: 1) eukaryotic and bacterial sequences branched near one another with

strong support (bootstrap values >70); 2) the eukaryotic sequences formed a distinct subclade,
represented by at least 2 species from the same eukaryotic supergroup; 3) the eukaryotic
sequences were produced by at least 2 different studies; and 4) the position of the horizontally
transferred sequences was robust across all alignment and phylogenetic reconstruction
methods used (Supp. Fig. 3). While these restrictions limit our attention to relatively old HGT
events, they also give us confidence these events are likely to be real.
The Mab21 superfamilies passed all four of these HGT thresholds, as did another
eukaryotic clade of CD-NTases that were all previously undescribed. We name this clade the
eukaryotic SMODS (eSMODS) superfamily, because the top scoring domain from hmmscan for
each sequence in this superfamily was the SMODS domain (PF18144), which is typically found
only in bacterial CD-NTases (Supplementary Data). This sequence similarity suggests that
eSMODS arose following a recent HGT from bacteria and/or that these CD-NTases have
diverged from their bacterial predecessors less than the eukaryotic OAS and Mab21 families
have. Additionally, all of these sequences were predicted to have a Nucleotidyltransferase
domain (PF01909), and (8/12) had a Polymerase Beta domain (PF18765), which are features
shared with many bacterial CD-NTases in Clades D,E, and F (Supplementary Data). The
eSMODS superfamily is made up of sequences from Amoebozoa, choanoflagellates,
Ancryomonadida, and one animal (the sponge *Oscarella pearsei*), which clustered robustly and
with high support within bacterial Clade D(e.g. subclade D04, CD-NTase 22 from *Myxococcus*
xanthus) (Supp. Fig 4). The eSMODS placement on the tree, which was robust to all alignment
and phylogenetic algorithms used (Supp. Fig. 3), suggesting that eSMODS represent an
additional, independent acquisition of CD-NTases from bacteria.
CD-NTases from bacterial Clade C and Clade D are the only CD-NTases to produce
cyclic trinucleotides, producing cyclic tri-Adenylate and cAAG, respectively[11,47–49].
Interestingly, OAS produces linear adenylates, which is one step away from the cAAA product
made by Class C CD-NTases, and similarly cGAMP (made by cGAS) is one adenylate away
from the class D product cAAG. As of this writing, the Clade D CD-NTases closest to the
eSMODS and Mab21 superfamilies (D04 and D12, respectively), have not been well
characterized. Therefore we argue that these CD-NTases should be a focus of future studies,
as they may hint at the evolutionary stepping stones that allow eukaryotes to acquire bacterial
immune proteins.
208
209 Diverged eukaryotic STINGs bridge the gap between bacteria and animals
We next turned to analyze Stimulator of Interferon Gamma (STING) proteins. In animals,
STING is a critical cyclic dinucleotide sensor, important during viral, bacterial, and parasitic
infections (reviewed here[50]). Structurally, most metazoan STINGs consist of an N-terminal
transmembrane domain (TM), made of 4 alpha helices fused to a C-terminal STING domain[51].
Canonical animal STINGs show distant homology with STING effectors from the bacterial cyclic
oligonucleotide-based antiphage signaling system (CBASS), with major differences in protein
structure and pathway function between these animal and bacterial defenses. For example, in
bacteria, the majority of STING proteins are fusions of a STING domain to a TIR
(Toll/interleukin-1 receptor) domain (Fig. 3A). Bacterial STING proteins recognize cyclic di-GMP
and oligomerize upon activation, which promotes TIR enzymatic activity[3,52,53]. Some
bacteria, such as *Flavobacteriaceae,* encode proteins that fuse a STING domain to a

transmembrane domain, although it is unclear how these bacterial TM-STINGs function[3].
Other bacteria have STING domain fusions with deoxyribohydrolase, α/β- hydrolase, or trypsin
peptidase domains[14]. In addition to eukaryotic TM-STINGs, a few eukaryotes such as the
oyster *Crassostrea gigas* have TIR-STING fusion proteins, although the exact role of their TIR
domain remains unclear[3,54,55].
Given these major differences in domain architectures, ligands, and downstream
immune responses, how have animals and bacteria evolved their STING-based defenses, and
what are the relationships between them? Prior to this work, the phylogenetic relationship
between animal and bacterial STINGs has been difficult to characterize with high support[14].
Indeed, when we made a tree of previously known animal and bacterial STING domains, we
found that the metazoan sequences were separated from the bacterial sequences by one very
long branch, along which many changes had occurred (Fig. 3B).
To improve the phylogeny through the inclusion of a greater diversity of eukaryotic
STING sequences, we began by carefully identifying the region of STING that was homologous
between bacterial and animal STINGs, as we expected this region to be best conserved across
diverse eukaryotes. Although Pfam domain PF15009 (TMEM173) is commonly used to define
animal STING domains, this HMM includes a portion of STING's transmembrane domain which
is not shared by bacterial STINGs. Therefore, we compared the crystal structures of HsSTING
(6NT5), *Flavobacteriaceae sp.* STING (6WT4) and *Crassostrea gigas* STING (6WT7) to define
a core "STING" domain. We used the region corresponding to residues 145-353 of 6NT5 as an
initial HMM seed alignment of 15 STING sequences from PF1500915 ("Reviewed" sequences
on InterPro). Our searches yielded 146 eukaryotic sequences from 64 species, which included
STING homologs from 34 metazoans (Supplemental Data and Fig. 1). Using maximum
likelihood phylogenetic reconstruction, we identified STING-like sequences from 26 diverse
microeukaryotes that clustered in between bacterial and metazoan sequences, breaking up the
long branch. We name these sequences the bacteria-like STINGs (blSTINGs) because they
were the only eukaryotic group of STINGs with a bacteria-like Prok_STING domain (PF20300)
and due to the short branch length (0.86 vs. 1.8) separating them from bacterial STINGs on the
tree (Fig. 3C). While a previous study reported STING domains in two eukaryotic species (one
in Stramenopiles and one in Haptista) [14], we were able to expand this set to additional species
and also recover blSTINGs from Amoebozoa, Rhizaria and choanoflagellates. This diversity
allowed us to place the sequences on the tree with high confidence, recovering a substantially
different tree than previous work[14]. As for CD-NTases, the tree topology we recovered was
robust across multiple different alignment and phylogenetic tree construction algorithms (Supp.
Fig. 3).
Given the similarities between the STING domains of the blSTINGs and bacterial
STINGs, we next asked whether the domain architectures of these proteins were similar using
Hmmscan and AlphaFold. The majority of the new eukaryotic blSTINGs were predicted to have
four N-terminal alpha helices (Fig. 3A, and Supplementary Data), similar to human STING.
While bacterial TM-STINGs were superficially similar with N-terminal transmembrane domains,
these proteins were predicted to have only two alpha helices and in 5/6 phylogenetic trees
bacterial TM-STINGs were more similar to other bacterial STINGs than to eukaryotic homologs
(Supp. Fig. 3). These results suggest that eukaryotes and bacteria independently converged on
a common TM-STING domain architecture through domain shuffling.

Interestingly, a similar pattern of convergent domain shuffling appears to have occurred
a second time with the TIR-STING proteins. It was previously known that some eukaryotes such
as the oyster *C. gigas,* have a TIR-STING fusion protein[3,54,55]. The STING domain of these
TIR-STINGs clustered closely to other metazoan STINGs, suggesting an animal origin (Fig. 3B).
We also investigated the possibility that *C. gigas* acquired the TIR-domain of its TIR-STING
protein via HGT from bacteria, however this analysis also suggested an animal origin for the TIR
domain (Supp. Fig. 7). Eukaryotic TIR-STINGs are rare, further supporting the hypothesis that
this protein resulted from recent convergence, where animals independently fused STING and
TIR domains to make a protein resembling bacterial TIR-STINGs, consistent with previous
reports[14]. Overall, we find that the TM-STING and TIR-STING proteins represent at least two
independent examples of convergent evolution, where bacteria and eukaryotes have created
similar proteins through the reuse of ancient protein domains. Our work also identified a number
of non-metazoan STINGs (the blSTINGs) that have a domain architecture similar to animal
STINGs but a STING domain more similar to bacterial STINGs.
279
Viperin is an ancient and widespread immune family
Viperins are innate immune proteins that restrict the replication of a diverse array of
viruses by conversion of nucleotides into 3'-deoxy-3',4'didehdro- (ddh) nucleotides[4,56–58].
Incorporation of these ddh nucleotides into a nascent RNA molecule leads to chain termination,
blocking RNA synthesis and inhibiting viral replication[56,59]. While metazoan viperin
specifically catalyzes CTP to ddhCTP[56], homologs from archaea and bacteria can generate
ddhCTP, ddhGTP, and ddhUTP[4,60]. Previous structural and phylogenetic analysis showed
that eukaryotic viperins are highly conserved at both the sequence and structural level and that,
phylogenetically, animal and fungal viperins form a distinct monophyletic clade compared to
bacterial viperins[4,57,60].
As viperin proteins consist of a single Radical SAM protein domain, we iteratively
searched EukProt beginning with domain PF04055 (Radical_SAM). The 194 viperin-like
proteins we recovered came from 158 species spanning the full range of eukaryotic diversity,
including organisms from all of the major eukaryotic supergroups, as well as some orphan taxa
whose taxonomy remains open to debate (Fig. 1). When we constructed phylogenetic trees
from these sequences, we found that the large majority of the eukaryotic viperins cluster
together in a single, monophyletic clade, separate from bacterial or archaeal viperins (Fig. 4).
Within the eukaryotic viperin clade, sequences from more closely related eukaryotes often
clustered together (Fig. 4, colored blocks), as would be expected if viperins were present and
vertically inherited within eukaryotes for an extended period of time. The vast species diversity
and tree topology both strongly support the inference that viperins are a truly ancient immune
module and have been present within the eukaryotic lineage likely dating back to the last
eukaryotic common ancestor (LECA).
In addition to this deep eukaryotic ancestry, we also uncovered two examples of
bacteria-eukaryote HGT that have occurred much more recently, both in Chloroplastida, a group
within Archaeplastida. The first of these consists of a small clade of Archaeplastida (Clade A)
consisting of marine algae such as *Chloroclados australicus* and *Nemeris dumetosa.* These
algal viperins cluster closely with the marine cyanobacteria *Anabaena cylindrica* and
Plankthriodies (Fig. 4 and Supp. Fig. 6). The second clade (Clade B) includes four other

Archaeplastida green algal species, mostly *Chlamydomonas spp.* In some of our trees the
Clade B viperins branched near to eukaryotic sequences from other eukaryotic supergroups,
however the placement of the neighboring eukaryotic sequences varied depending on the
algorithms we used; only the Archaeplastida placement was consistent. (Fig. 4 and Supp. Fig. 3
& 6). Taken together, we conclude that viperins represent a class of ancient immune proteins
that have likely been present in eukaryotes since the LECA. Yet, we also find ongoing
evolutionary innovation in viperins via HGT, both among eukaryotes and between eukaryotes
and bacteria.

Discussion

The recent discoveries that bacteria and mammals share mechanisms of innate
immunity have been surprising, because they imply that there are similarities in immunity that
span the Tree of Life. But how did these similarities come to exist? Here we uncover several
evolutionary trajectories that have led animals and bacteria to share homologous immune
proteins (summarized in Fig. 5). We found that Viperin is truly ancestral, dating back to at least
the Last Eukaryotic Common Ancestor (LECA), and likely further. We also uncovered examples
of convergence, as in STING, where the shuffling of ancient domains has led animals and
bacteria to independently arrive at similar protein architectures. Finally, we found evidence of
multiple examples of bacteria-eukaryote HGTs that have given rise to immune protein families.
An essential part of our ability to make these discoveries was the analysis of data from nearly
1000 diverse eukaryotic taxa. These organisms allowed us to distinguish between proteins
found across eukaryotes vs. animal-specific innovations, to document both recent and ancient
HGT events from bacteria that gave rise to eukaryotic immune protein families (Fig. 2 & 4), and
to identify STING proteins with eukaryotic domain architectures but more bacteria-like domains
(bISTINGs, Fig. 3). Because these diverged eukaryotic STINGs were found in organisms where
we typically did not find any CD-NTase proteins, we hypothesize that bISTINGs may detect and
respond to exogenous cyclic nucleotides, such as those generated by pathogens. In contrast to
the STINGs, the eukaryotic CD-NTases had substantially different evolutionary histories, with
multiple major CD-NTase superfamilies each emerging from within larger bacterial clades. While
these analyses cannot definitively determine the directionality of the transfer, we favor the most
parsimonious explanation that these components came into the eukaryotic lineage from
bacterial origins.
While not as prevalent as in bacteria, HGT in eukaryotes represents a significant force in
eukaryotic evolution, especially for unicellular eukaryotes[61–64]. In this study, our criteria for
'calling' HGT events was relatively strict, meaning that our estimate of HGT events is almost
certainly an underestimate. Importantly, this pattern suggests that the bacterial pan-genome has
been a rich reservoir that eukaryotes have repeatedly sampled to acquire novel innate immune
components. Some of these HGT events have given rise to new eukaryotic superfamilies (e.g.
eSMODS) that have never been characterized and could represent novel types of eukaryotic
immune proteins. We speculate that the eSMODS superfamily CD-NTases and the bISTINGs
may function more similarly to their bacterial homologs, potentially producing and responding to
a variety of cyclic di- or tri-nucleotides[11] Similarly, bacterial viperins have been shown to
generate ddhCTP, ddhGTP, and ddhUTP, whereas animal viperins only make

ddhCTP[4,56,60]. Thus, the two algal viperin clades arising from HGT may have expanded
functional capabilities as well. A caveat of this work is that such strictly bioinformatic
investigations are insufficient to reveal protein biochemical functions, nor can they determine
whether diverse homologs have been co-opted for non-immune functions. We therefore urge
future, functional studies to focus on these proteins to resolve the questions of 1) whether/how
bISTINGs operate in the absence of CD-NTases, 2) whether/how the functions of algal viperins
and eSMODS changed following their acquisition from bacteria, and 3) whether the homologs
truly function in immune defense.
In addition to these instances of gene gain, eukaryotic gene repertoires have been
dramatically shaped by losses. Even for viperins, which likely date back to the eukaryotic last
common ancestor, these proteins were sparsely distributed across eukaryotes and were absent
from the majority of species we surveyed. While some of this finding may be due to technical
limitations, such as dataset incompleteness or inability of the HMMs to recover distant
homologs, we believe this explanation is insufficient to fully explain the sparseness, as many
plant, fungal, and amoebozoan species are represented by well-assembled genomes where
these proteins are certifiably absent (Supp. Fig. 2). Instead, we propose that the sparse
distribution likely arises from ongoing and repeated gene loss, as has been previously
documented for other gene families across the Tree of Life[22,24–26].
Overall, our results yield a highly dynamic picture of immune protein evolution across
eukaryotes, wherein multiple mechanisms of gene gain are offset by ongoing losses.
Interestingly, this pattern mirrors the sparse distributions of many of these immune homologs
across bacteria[65–67], as anti-phage proteins tend to be rapidly gained and lost from genomic
defense islands[68,69]. It will be interesting to see if some eukaryotes evolve their immune
genes in similarly dynamic islands, particularly in unicellular eukaryotes that undergo more
frequent HGT[70].
We expect that our examination of STING, CD-NTases, and Viperin represents just the
tip of the iceberg when it comes to the evolution of eukaryotic innate immunity. New links
between bacterial and animal immunity continue to be discovered and other immune families
and domains such as Argonaute, Gasdermins, NACHT domains, CARD domains, TIR domains,
and SamHD1 have been shown to have bacterial roots[2,6,7,9,10]. To date, the majority of
studies have focused on proteins specifically shared between metazoans and bacteria. We
speculate that there are probably many other immune components shared between bacteria
and eukaryotes outside of animals. Further studies of immune defenses in microeukaryotes are
likely to uncover new mechanisms of cellular defense and to better illustrate the origins and
evolution of eukaryotic innate immunity.
386

Methods

Iterative HMM Search

The goal of this work was to search the breadth of EukProt v3 for immune proteins from
the CD-NTase, STING, and viperin families that span the gap between metazoan and bacterial
immunity. Our overall strategy was to first search with eukaryotes alone (starting from mainly
Metazoa). Then we added in bacterial sequences and searched with a mixed bacterial-

eukaryotic HMM search until we either found no new hits, or until we began getting hits from an
outgroup gene family. In parallel, we also performed bacteria-only and eukaryote-only searches,
to ensure that we found as many homologs as possible (schematized in Fig 1A, and further in
Supp. Fig. 1A).
397   Phase 1: Eukaryotic searches To begin, hidden Markov model (HMM) profiles from Pfam
(for CD-NTases and Viperin) or an HMM profile generated from a multiple sequence alignment
(for STING) were used to search EukProt V3[15], for diverse eukaryotic sequences. For CD-
NTases and Viperin, HMM profiles of Pfams PF03281 and PF04055 were used respectively.
For STING, where the Pfam profile includes regions of the protein outside of the STING
domain, we generated a new HMM for the initial search. First, we aligned crystal structures of
HsSTING (6NT5), *Flavobacteriaceae sp.* STING (6WT4) and *Crassostrea gigas* STING (6WT7)
to define a core "STING" domain. Then we aligned 15 eukaryotic sequences from PF15009
("Reviewed" sequences on InterPro) with MAFFT(v7.4.71)[71] and manually trimmed the
sequences down to the boundaries defined by our crystal alignment (residues 145-353 of
6NT5). We then trimmed the alignment with TrimAl (v1.2)[72] with options -gt 0.2. The trimmed
MSA was then used to generate an HMM profile with hmmbuild from the hmmer (v3.2.1)
package (hmmer.org).
HMM profiles were used to search EukProt via hmmsearch (also from hmmer v3.2.1)
with a statistical cutoff value of 1e-3 and -hit parameter set to 10 (i.e. the contribution of a single
species to the output list is capped at 10 sequences). The resulting hits from this search were
then aligned with hmmalign (included within hmmer) and used to generate a new HMM profile
with hmmbuild. This profile was used to search EukProt v3 again and the process was repeated
for a total of 3-4 eukaryotic searches.
Phase 2: combining eukaryotic and bacterial sequences into an HMM After the
eukaryotic searches reached saturation (i.e. no additional eukaryotic sequences were recovered
after additional searches), bacterial sequences were acquired from previous literature (Viperins
from[4], CD-NTases from[11], and STINGs from[3,8,43]). To ensure the combined HMM did not
have an overrepresentation of either bacterial or eukaryotic sequences, we downsampled the
bacterial sequences and eukaryotic sequences to obtain 50 phylogenetically diverse sequences
of each, and then combined the two downsampled lists. To do this, eukaryotic and bacterial
sequences were each separately aligned with MAFFT, phylogenetic trees were built with
FastTree (v2.1.10)[73], and the Phylogenetic Diversity Analyzer (pda/1.0.3)[74] software was
used to downsample the sequences while maximizing remaining sequence diversity.
The combined bacterial-eukaryotic sequence list was then aligned with hmmalign and
used to construct a new HMM profile with hmmbuild. This HMM profile was used to search
EukProt v3. The eukaryotic hits from this search were then aligned with MAFFT, and a tree was
constructed with FastTree. From this tree the sequences were then downsampled with PDA and
once again combined with the bacterial list, aligned, used to generate a new HMM, and a new
search. This process was iterated 3-5 times until saturation or until the resulting sequence hits
included other gene families that branched outside of the sequence diversity defined by the
metazoan and bacterial homologs.
Phase 3: Searching with a bacteria-only or existing eukaryote-only HMM profiles
To search EukProt v3 with a bacteria-only HMM for each protein family, we aligned the
full set of published bacterial sequences with MAFFT, trimmed with TrimAl, and hmmbuild was

used to generate an HMM profile which was used to search EukProt v3. As a point of
comparison, we also searched the database with only the starting, animal-dominated Pfam
(PF15009) for STING.
<u>Phase 4: Combining all hits into a single list and scanning for domains</u>
Sequences from all iterative searches were combined to generate a total hits FASTA file
for STING, CD-NTase, and Viperin. First, duplicate sequences were removed, then the fasta
files were scanned using hmmscan (also from hmmer v3.2.1) against the Pfam database (Pfam-
A.hmm) and all predicted domains with an E-value <1e-3 were considered. Next, we generated
phylogenetic trees (first by aligning with MAFFT and then building a tree with FastTree), and
used these trees along with the hmmscan domains to determine in-group and out-group
sequences. Out-group sequences were removed from the fasta file. We determined outgroup
sequences by these criteria: 1) if the sequence clustered outside of known outgroup sequences
(e.g. Poly(A) RNA polymerase (PAP) sequences for the CD-NTases, and molybdenum cofactor
biosynthetic enzyme (MoaA) for Viperin), or 2) if sequence did not have at least one of the
relevant domains (Mab21/OAS1-C/SMODS for CD-NTases, TMEM173/Prok_STING for STING,
and Radical_SAM for Viperin). These three FASTA files were used for the final alignments and
phylogenetic trees. To identify protein domains in each sequence, the FASTA files were
scanned using hmmscan (also from hmmer v3.2.1) against the Pfam database (Pfam-A.hmm)
and all predicted domains with an E-value <1e-3 were considered. See Supplemental Data for
the hmmscan results of all included homologs.
457

Final Alignment and Tree Building
To generate final phylogenetic trees, all eukaryotic search hits and bacterial sequences
were aligned using MAFFT. We downsampled the CD-NTase bacterial sequences from ~6000
down to 500 as described above, to facilitate more manageable computation times on
alignments and tree construction. For the STING and Viperin trees we included all bacterial
sequences. These initial alignments were first trimmed manually in Geneious (v2023.1.2) to
remove unaligned N- and C-terminal regions, and then realigned with MAFFT or
MUSCLE(v5.1)[75] and trimmed with TrimAl (v1.2)[72]. MUSCLE was used with the "-super5"
option, and otherwise default parameters. These alignments were used to generate
phylogenetic trees using three tree inference softwares: FastTree (v2.1.10)[73], IQtree
(2.2.2.7)[76] and RaxML-ng (v0.9.0)[77]. FastTree was utilized with default settings. IQtree was
used to determine the appropriate evolutionary model, and was run with 1000 ultrafast
bootstraps (IQtree settings: -s, -bb 1000, -m TEST, -nt AUTO). RaxML-ng trees were produced
with 100 bootstraps using the molecular model specified from the IQtree analysis (Raxml-ng
settings: --all, --model [specified by IQtree], --tree pars{10} --bs-trees 100). Phylogenetic trees
were visualized with iTOL[78].
474

TIR Domain Alignment and Tree
We used hmmscan to identify the coordinates of TIR domains in a list of 203 TIR domain
containing-sequences from InterPro (Family: IPR015032) and 104 bacterial TIR-STING proteins
(the same TIR-STING proteins used in Fig. 3)[3]. Next, we trimmed the sequences down to the
TIR coordinates and aligned the TIR domains with MUSCLE. We trimmed the alignments with
TrimAL and built a phylogenetic tree with IQtree.

481
Venn Diagrams

483 Venn diagrams were generated via DeepVenn[79] using presence/absence information for Mab21, OAS, and STING from each eukaryotic species that encoded at least one of these proteins.

Protein Structure Modeling

488 To model 3-D protein structures for STING homologs without a published crystal structure, we ran AlphaFold (v2.1.1)[80,81]. We generated 5 ranked models for STINGs from *Flavobacteriaceae* (IMG ID: 2624319773), *Nitzschia sp.* (EukProt ID: P007051), and *Caveostelium apophsatum* (EukProt ID: P019191). Fig. 2C shows highest ranked models only.

Acknowledgements

We thank Daniel Richter for his feedback, encouragement, and scientific guidance. Maureen Stolzer, Kevin Forsberg, Patrick Mitchell, and members of the Levin lab also provided helpful input on the project and manuscript. Thanks to Andrew VanDemark for helpful discussions about 3-D modeling and to Jacob Durrant for help running AlphaFold. This research was supported in part by the University of Pittsburgh Center for Research Computing, RRID:SCR_022735, through the resources provided. Specifically, this work used the HTC cluster, which is supported by NIH award number S10OD028483. Ed Culbertson was supported by NSF Postdoctoral fellowship 2208971 and Tera Levin was supported by NIH R00AI139344 and R35GM150681.

References:

1. Litman GW, Cannon JP, Dishaw LJ. Reconstructing immune phylogeny: new perspectives. Nat Rev Immunol. 2005;5: 866–879.

2. Johnson AG, Wein T, Mayer ML, Duncan-Lowey B, Yirmiya E, Oppenheimer-Shaanan Y, et al. Bacterial gasdermins reveal an ancient mechanism of cell death. bioRxiv. 2021. p. 2021.06.07.447441. doi:10.1101/2021.06.07.447441

3. Morehouse BR, Govande AA, Millman A, Keszei AFA, Lowey B, Ofir G, et al. STING cyclic dinucleotide sensing originated in bacteria. Nature. 2020;586: 429–433.

4. Bernheim A, Millman A, Ofir G, Meitav G, Avraham C, Shomar H, et al. Prokaryotic viperins produce diverse antiviral molecules. Nature. 2021;589: 120–124.

5.   Wein T, Sorek R. Bacterial origins of human cell-autonomous innate immune
mechanisms. Nat Rev Immunol. 2022. doi:10.1038/s41577-022-00705-4

6.   Kibby EM, Conte AN, Burroughs AM, Nagy TA, Vargas JA, Whalen LA, et al.
Bacterial NLR-related proteins protect against phage. Cell. 2023.
doi:10.1016/j.cell.2023.04.015

7.   Swarts DC, Makarova K, Wang Y, Nakanishi K, Ketting RF, Koonin EV, et al.
The evolutionary journey of Argonaute proteins. Nat Struct Mol Biol.
2014;21: 743–753.

8.   Cohen D, Melamed S, Millman A, Shulman G, Oppenheimer-Shaanan Y,
Kacen A, et al. Cyclic GMP–AMP signalling protects bacteria against viral
infection. Nature. 2019;574: 691–695.

9.   Ofir G, Herbst E, Baroz M, Cohen D, Millman A, Doron S, et al. Antiviral
activity of bacterial TIR domains via immune signalling molecules. Nature.
2021;600: 116–120.

10.   Tal N, Millman A, Stokar-Avihail A, Fedorenko T, Leavitt A, Melamed S, et
al. Bacteria deplete deoxynucleotides to defend against bacteriophage
infection. Nat Microbiol. 2022;7: 1200–1209.

11.   Whiteley AT, Eaglesham JB, de Oliveira Mann CC, Morehouse BR, Lowey
B, Nieminen EA, et al. Bacterial cGAS-like enzymes synthesize diverse
nucleotide signals. Nature. 2019;567: 194–199.

12.   Kaur G, Burroughs AM, Iyer LM, Aravind L. Highly regulated, diversifying
NTP-dependent biological conflict systems with implications for the
emergence of multicellularity. Elife. 2020;9. doi:10.7554/eLife.52696

13.   Wein T, Johnson AG, Millman A, Lange K, Yirmiya E, Hadary R, et al.
CARD-like domains mediate anti-phage defense in bacterial gasdermin
systems. bioRxiv. 2023. p. 2023.05.28.542683.
doi:10.1101/2023.05.28.542683

14.   Burroughs AM, Aravind L. Identification of Uncharacterized Components of
Prokaryotic Immune Systems and Their Diverse Eukaryotic Reformulations.
J Bacteriol. 2020;202. doi:10.1128/JB.00365-20

15.   Richter DJ, Berney C, Strassert JFH, Poh Y-P, Herman EK, Muñoz-Gómez
SA, et al. EukProt: a database of genome-scale predicted proteins across
the diversity of eukaryotes. bioRxiv. bioRxiv; 2020.
doi:10.1101/2020.06.30.180687

16. del Campo J, Sieracki ME, Molestina R, Keeling P, Massana R, Ruiz-Trillo I.
The others: our biased perspective of eukaryotic genomes. Trends Ecol
Evol. 2014;29: 252–259.

17. Shaw AE, Hughes J, Gu Q, Behdenna A, Singer JB, Dennis T, et al.
Fundamental properties of the mammalian innate immune system revealed
by multispecies comparison of type I interferon responses. PLoS Biol.
2017;15: e2004086.

18. Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR, et al.
Pfam: the protein families database. Nucleic Acids Res. 2014;42: D222–30.

19. Eddy SR. Accelerated Profile HMM Searches. PLoS Comput Biol. 2011;7:
e1002195.

20. Eddy SR. Profile hidden Markov models. Bioinformatics. 1998;14: 755–763.

21. Demuth JP, De Bie T, Stajich JE, Cristianini N, Hahn MW. The evolution of
mammalian gene families. PLoS One. 2006;1: e85.

22. Albalat R, Cañestro C. Evolution by gene loss. Nat Rev Genet. 2016;17:
379–391.

23. Wang X, Grus WE, Zhang J. Gene losses during human origins. PLoS Biol.
2006;4: e52.

24. Koumandou VL, Wickstead B, Ginger ML, van der Giezen M, Dacks JB,
Field MC. Molecular paleontology and complexity in the last eukaryotic
common ancestor. Crit Rev Biochem Mol Biol. 2013;48: 373–396.

25. Prokopchuk G, Butenko A, Dacks JB, Speijer D, Field MC, Lukeš J. Lessons
from the deep: mechanisms behind diversification of eukaryotic protein
complexes. Biol Rev Camb Philos Soc. 2023. doi:10.1111/brv.12988

26. Farkas Z, Kovács K, Sarkadi Z, Kalapis D, Fekete G, Birtyik F, et al. Gene
loss and compensatory evolution promotes the emergence of morphological
novelties in budding yeast. Nat Ecol Evol. 2022;6: 763–773.

27. Kranzusch PJ. cGAS and CD-NTase enzymes: structure, mechanism, and
evolution. Curr Opin Struct Biol. 2019;59: 178–187.

28. Li Y, Slavik KM, Toyoda HC, Morehouse BR, de Oliveira Mann CC, Elek A,
et al. cGLRs are a diverse family of pattern recognition receptors in innate
immunity. Cell. 2023. doi:10.1016/j.cell.2023.05.038

29. de Oliveira Mann CC, Kiefersauer R, Witte G, Hopfner K-P. Structural and biochemical characterization of the cell fate determining nucleotidyltransferase fold protein MAB21L1. Sci Rep. 2016;6: 27498.

30. Chow KL, Hall DH, Emmons SW. The mab-21 gene of Caenorhabditis elegans encodes a novel protein required for choice of alternate cell fates. Development. 1995;121: 3615–3626.

31. Yamada R, Mizutani-Koseki Y, Hasegawa T, Osumi N, Koseki H, Takahashi N. Cell-autonomous involvement of Mab21l1 is essential for lens placode development. Development. 2003;130: 1759–1770.

32. Keating SE, Baran M, Bowie AG. Cytosolic DNA sensors regulating type I interferon induction. Trends Immunol. 2011;32: 574–581.

33. Hornung V, Latz E. Intracellular DNA recognition. Nat Rev Immunol. 2010;10: 123–130.

34. Ablasser A, Goldeck M, Cavlar T, Deimling T, Witte G, Röhl I, et al. cGAS produces a 2'-5'-linked cyclic dinucleotide second messenger that activates STING. Nature. 2013;498: 380–384.

35. Gao P, Ascano M, Wu Y, Barchet W, Gaffney BL, Zillinger T, et al. Cyclic [G(2′,5′)pA(3′,5′)p] Is the Metazoan Second Messenger Produced by DNA-Activated Cyclic GMP-AMP Synthase. Cell. 2013. pp. 1094–1107. doi:10.1016/j.cell.2013.04.046

36. Sun L, Wu J, Du F, Chen X, Chen ZJ. Cyclic GMP-AMP synthase is a cytosolic DNA sensor that activates the type I interferon pathway. Science. 2013;339: 786–791.

37. Burdette DL, Monroe KM, Sotelo-Troha K, Iwig JS, Eckert B, Hyodo M, et al. STING is a direct innate immune sensor of cyclic di-GMP. Nature. 2011;478: 515–518.

38. Gui X, Yang H, Li T, Tan X, Shi P, Li M, et al. Autophagy induction via STING trafficking is a primordial function of the cGAS pathway. Nature. 2019;567: 262–266.

39. Dong B, Silverman RH. A bipartite model of 2-5A-dependent RNase L. J Biol Chem. 1997;272: 22236–22242.

40. Silverman RH. Viral encounters with 2',5'-oligoadenylate synthetase and RNase L during the interferon antiviral response. J Virol. 2007;81: 12720–

12729.

41. Kristiansen H, Gad HH, Eskildsen-Larsen S, Despres P, Hartmann R. The
oligoadenylate synthetase family: an ancient protein family with multiple
antiviral activities. J Interferon Cytokine Res. 2011;31: 41–47.

42. Severin GB, Ramliden MS, Hawver LA, Wang K, Pell ME, Kieninger A-K, et
al. Direct activation of a phospholipase by cyclic GMP-AMP in El Tor Vibrio
cholerae. Proc Natl Acad Sci U S A. 2018;115: E6048–E6055.

43. Millman A, Melamed S, Amitai G, Sorek R. Diversity and classification of
cyclic-oligonucleotide-based anti-phage signalling systems. Nat Microbiol.
2020;5: 1608–1615.

44. Tsang WH, Shek KF, Lee TY, Chow KL. An evolutionarily conserved nested
gene pair - Mab21 and Lrba/Nbea in metazoan. Genomics. 2009;94: 177–
187.

45. Wu X, Wu F-H, Wang X, Wang L, Siedow JN, Zhang W, et al. Molecular
evolutionary and structural analysis of the cytosolic DNA sensor cGAS and
STING. Nucleic Acids Res. 2014;42: 8243–8257.

46. Woznica A, Kumar A, Sturge CR, Xing C, King N, Pfeiffer JK. STING
mediates immune responses in the closest living relatives of animals. Elife.
2021;10. doi:10.7554/eLife.70436

47. Ye Q, Lau RK, Mathews IT, Birkholz EA, Watrous JD, Azimi CS, et al.
HORMA Domain Proteins and a Trip13-like ATPase Regulate Bacterial
cGAS-like Enzymes to Mediate Bacteriophage Immunity. Molecular Cell.
2020. pp. 709–722.e7. doi:10.1016/j.molcel.2019.12.009

48. Govande AA, Duncan-Lowey B, Eaglesham JB, Whiteley AT, Kranzusch PJ.
Molecular basis of CD-NTase nucleotide selection in CBASS anti-phage
defense. Cell Rep. 2021;35: 109206.

49. Hogrel G, Guild A, Graham S, Rickman H, Grüschow S, Bertrand Q, et al.
Author Correction: Cyclic nucleotide-induced helical structure activates a TIR
immune effector. Nature. 2023;614: E15.

50. Du Y, Hu Z, Luo Y, Wang HY, Yu X, Wang R-F. Function and regulation of
cGAS-STING signaling in infectious diseases. Front Immunol. 2023;14:
1130423.

51. Ishikawa H, Barber GN. STING is an endoplasmic reticulum adaptor that

facilitates innate immune signalling. Nature. 2008;455: 674–678.

52. Nishimura MT. TIR Domains of Plant Immune Receptors are NAD plus
Consuming Enzymes that Promote Cell Death. MOLECULAR PLANT-
MICROBE INTERACTIONS. AMER PHYTOPATHOLOGICAL SOC 3340
PILOT KNOB ROAD, ST PAUL, MN 55121 USA; 2019. pp. 217–218.

53. Horsefield S, Burdett H, Zhang X, Manik MK, Shi Y, Chen J, et al. NAD+
cleavage activity by animal and plant TIR domains in cell death pathways.
Science. 2019;365: 793–799.

54. Margolis SR, Wilson SC, Vance RE. Evolutionary Origins of cGAS-STING
Signaling. Trends Immunol. 2017;38: 733–743.

55. Zhang G, Fang X, Guo X, Li L, Luo R, Xu F, et al. The oyster genome
reveals stress adaptation and complexity of shell formation. Nature.
2012;490: 49–54.

56. Gizzi AS, Grove TL, Arnold JJ, Jose J, Jangra RK, Garforth SJ, et al. A
naturally occurring antiviral ribonucleotide encoded by the human genome.
Nature. 2018;558: 610–614.

57. Fenwick MK, Li Y, Cresswell P, Modis Y, Ealick SE. Structural studies of
viperin, an antiviral radical SAM enzyme. Proc Natl Acad Sci U S A.
2017;114: 6806–6811.

58. Rivera-Serrano EE, Gizzi AS, Arnold JJ, Grove TL, Almo SC, Cameron CE.
Viperin Reveals Its True Function. Annu Rev Virol. 2020;7: 421–446.

59. Seifert M, Bera SC, van Nies P, Kirchdoerfer RN, Shannon A, Le T-T-N, et
al. Inhibition of SARS-CoV-2 polymerase by nucleotide analogs from a
single-molecule perspective. Elife. 2021;10. doi:10.7554/eLife.70968

60. Lachowicz JC, Gizzi AS, Almo SC, Grove TL. Structural Insight into the
Substrate Scope of Viperin and Viperin-like Enzymes from Three Domains of
Life. Biochemistry. 2021;60: 2116–2129.

61. Eme L, Gentekaki E, Curtis B, Archibald JM, Roger AJ. Lateral Gene
Transfer in the Adaptation of the Anaerobic Parasite Blastocystis to the Gut.
Curr Biol. 2017;27: 807–820.

62. Richards TA, Soanes DM, Jones MDM, Vasieva O, Leonard G, Paszkiewicz
K, et al. Horizontal gene transfer facilitated the evolution of plant parasitic
mechanisms in the oomycetes. Proc Natl Acad Sci U S A. 2011;108: 15258–

15263.

63. Gabaldón T. Patterns and impacts of nonvertical evolution in eukaryotes: a
paradigm shift. Ann N Y Acad Sci. 2020;1476: 78–92.

64. Leger MM, Eme L, Stairs CW, Roger AJ. Demystifying Eukaryote Lateral
Gene Transfer (Response to Martin 2017 DOI: 10.1002/bies.201700115).
Bioessays. 2018;40: e1700242.

65. Bernheim A, Sorek R. The pan-immune system of bacteria: antiviral defence
as a community resource. Nat Rev Microbiol. 2020;18: 113–119.

66. Koonin EV, Makarova KS, Wolf YI. Evolutionary Genomics of Defense
Systems in Archaea and Bacteria. Annu Rev Microbiol. 2017;71: 233–261.

67. van Houte S, Buckling A, Westra ER. Evolutionary Ecology of Prokaryotic
Immune Mechanisms. Microbiol Mol Biol Rev. 2016;80: 745–763.

68. Hochhauser D, Millman A, Sorek R. The defense island repertoire of the
Escherichia coli pan-genome. PLoS Genet. 2023;19: e1010694.

69. LeGault KN, Hays SG, Angermeyer A, McKitterick AC, Johura F-T, Sultana
M, et al. Temporal shifts in antibiotic resistance elements govern phage-
pathogen conflicts. Science. 2021;373. doi:10.1126/science.abg2166

70. Keeling PJ, Palmer JD. Horizontal gene transfer in eukaryotic evolution. Nat
Rev Genet. 2008;9: 605–618.

71. Katoh K, Standley DM. MAFFT multiple sequence alignment software
version 7: improvements in performance and usability. Mol Biol Evol.
2013;30: 772–780.

72. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for
automated alignment trimming in large-scale phylogenetic analyses.
Bioinformatics. 2009;25: 1972–1973.

73. Price MN, Dehal PS, Arkin AP. FastTree 2--approximately maximum-
likelihood trees for large alignments. PLoS One. 2010;5: e9490.

74. Chernomor O, Minh BQ, Forest F, Klaere S, Ingram T, Henzinger M, et al.
Split diversity in constrained conservation prioritization using integer linear
programming. Methods in Ecology and Evolution. 2015. pp. 83–91.
doi:10.1111/2041-210x.12299

75. Edgar RC. MUSCLE v5 enables improved estimates of phylogenetic tree

confidence by ensemble bootstrapping. bioRxiv. 2021. p. 2021.06.20.449169. doi:10.1101/2021.06.20.449169

76. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et al. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. Mol Biol Evol. 2020;37: 1530–1534.

77. Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. RAxML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. Bioinformatics. 2019;35: 4453–4455.

78. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. Nucleic Acids Res. 2021;49: W293–W296.

79. Hulsen T. DeepVenn -- a web application for the creation of area-proportional Venn diagrams using the deep learning framework Tensorflow.js. arXiv [cs.HC]. 2022. Available: http://arxiv.org/abs/2210.04597

80. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate protein structure prediction with AlphaFold. Nature. 2021;596: 583–589.

81. Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al. AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-sequence space with high-accuracy models. Nucleic Acids Res. 2022;50: D439–D444.

82. Shang G, Zhang C, Chen ZJ, Bai X-C, Zhang X. Cryo-EM structures of STING reveal its mechanism of activation by cyclic GMP–AMP. Nature. 2019;567: 389–393.

Figures

Figure 1: HMM searches to find homologs across the eukaryotic Tree of Life

(A) A schematic of the HMM search process. Starting from initial, animal-dominated HMM
profiles for each protein family, we used iterative HMM searches of the EukProt database to
generate pan-eukaryotic HMMs. These were combined with bacterial sequences to enable
discovery of bacteria-like homologs in eukaryotes. Each set of searches was repeated 3-5 times
until few or no additional eukaryotic sequences were recovered. (B) Phylogenetic tree of
eukaryotes, with major supergroups color-coded. The height of the colored rectangles for each
group is proportional to its species representation in EukProt. Horizontal, colored bars mark
each eukaryotic species in which we found homologs of STINGs, CD-NTases, or Viperins.
White space indicates species where we did not recover any homologs. The CD-NTase hits are
divided into the three eukaryotic superfamilies defined in Fig. 2.

752
Figure 2: Independent HGT events gave rise to multiple CD-NTase superfamilies
(A) Maximum likelihood phylogenetic tree generated by IQtree of CD-NTases spanning
eukaryotic and bacterial diversity. The Mab21 superfamily (red, top left) is largely an animal-

specific innovation, with many paralogs including cGAS. In contrast, most other eukaryotic
lineages encode CD-NTases from the OAS superfamily (multicolor, top right). The relatively
small eSMODS superfamily (pink, bottom left) is a recent HGT between clade D bacteria and
eukaryotes. Bacterial CD-NTase sequences shown in gray. Eukaryotic sequences are colored
according to eukaryotic group as in Fig. 1. The tree is arbitrarily rooted on a branch separating
clades A, B, G, and H, which did not typically have associated eukaryotic sequences, from the
rest of the bacterial CD-NTases. (B) Venn diagrams showing the number of species where we
detected at least one homolog of STING, Mab21 superfamily CD-NTases, and/or OAS
superfamily CD-NTases in Metazoa (left) or non-metazoan eukaryotes (right). (C) Within the
CD-NTase phylogenetic tree in A, the OAS superfamily branches within clade C bacterial CD-
NTases (gray branches). (D) Clade D CD-NTases (gray branches) have been horizontally
transferred into eukaryotes multiple times, giving rise to both the Mab21 superfamily and the
eSMODS superfamily. Ultrafast bootstraps determined by IQtree shown at key nodes. See
Supplementary Figure 4 for full CD-NTase phylogenetic tree.

Figure 3: Diverse eukaryotic STING proteins bridge the gap between metazoans and bacteria

(A) Graphical depiction of common domain architectures of STING proteins. (B) Maximum likelihood unrooted phylogenetic tree of STING domains from Metazoa and bacteria, which are separated by one long branch. Black dot (•) indicates proteins that have been previously experimentally characterized. Bacterial sequences are in gray and animal sequences are in red. (C) Maximum likelihood unrooted phylogenetic tree of hits from iterative HMM searches for diverse eukaryotic STING domains. The STING domains from bacteria-like STINGs (blSTINGs) from diverse eukaryotes break up the long branch between bacterial and animal STINGs.

Structures of the indicated STING proteins are shown above, with those predicted by AlphaFold
indicated by an asterisk. Homologs with X-ray crystal structures are from[3,82]. Colored regions
show two domain architectures in bacteria and eukaryotes (STING linked to a TIR domain and
STING linked to a transmembrane domain), each of which have evolved convergently in
bacteria and eukaryotes. Ultrafast bootstraps determined by IQtree shown at key nodes. See
Supplementary Figure 5 for full STING phylogenetic tree.

786
Figure 4: Viperin is a deeply conserved innate immune module
(A) Maximum likelihood phylogenetic tree generated by IQtree of Viperins from eukaryotes,
bacteria, and archaea. All major eukaryotic supergroups have at least two species that encode

a Viperin homolog (colored supergroups). Bacterial Viperin sequences shown in gray and
archaeal sequences in dark gray. There are two clades of Chloroplastida (a group within
Archaeplastida) sequences that branch robustly within the bacteria clade. Ultrafast bootstraps
determined by IQtree shown at key nodes. Tree is arbitrarily rooted between the major
eukaryotic and bacterial clades. See Supplementary Figure 6 for fully annotated Viperin
phylogenetic tree.

796
797

Figure 5: Proposed model of evolutionary history of CD-NTases, STING, and Viperin
Proposed summary models of the evolutionary history of innate immune components. (A) We
define two distinct superfamilies of CD-NTases that likely arose from bacteria-eukaryote HGT:
eSMODS and Mab21. Within the Mab21 superfamily (which contains cGAS), a number of
animal-specific duplications gave rise to numerous paralogs. The OAS superfamily of CD-
NTases are abundant across diverse eukaryotic taxa and were likely present in the LECA. (B)
Drawing on a shared ancient repertoire of protein domains that includes STING, TIR, and
transmembrane (TM) domains, bacteria and eukaryotes have convergently evolved similar
STING proteins through domain shuffling. (C) Viperins are widespread across the eukaryotic
tree and likely were present in the LECA. In addition, two sets of recent HGT events from
bacteria have equipped algal species with new viperins.

809
Supplementary Figure 1: Collectors curves and full search strategy
810

(A) Detailed schematic outlining the iterative HMM search strategy. Blue boxes and blue shaded

region show eukaryotic-only searches to create pan-eukaryotic HMMs and yellow indicates

eukaryotic-bacterial searches to create universal HMMs. For the combined bacterial/eukaryotic
searches (yellow box), bacterial and eukaryotic sequences were each downsampled to 50
sequences (phylogenetic tree downsampled via PDA) to maintain equal contributions from
bacteria and eukaryotic sequences. Separately, bacterial sequences were aligned and used to
make an HMM which was used to search EukProt as a 'bacteria only search' and for STING we
searched with PF15009 for a comparable Eukaryotic PFAM search (not shown in flowchart). We
did this extra search for STING as PF15009 contains part of the eukaryotic STING
transmembrane domain and so our first search with STING was with a STING-domain-only
HMM (See Materials and Methods). Pink (MUSCLE) and orange (MAFFT) boxes show the final
alignments and phylogenetic trees that were constructed. (B) STING, CD-NTase, and Viperin
collector's curves showing the number of cumulative protein sequences that were found after
each iterative search. Results from eukaryotic searches are shown in blue and the combined
searches in yellow. Solid black line indicates the number of hits from the starting Pfam HMM
alone and the dotted gray line shows the number of hits from a bacteria-only HMM. Note that
some searches yielded hits that were members of more distant protein families, which were
later removed from the analysis and are not counted here.

Supplementary Figure 2: Phylogenetic trees of EukProt species by data type

Phylogenetic trees derived from Fig. 1 separating species represented in EukProt v3 by

genomes (A) or transcriptomes (B). Supergroups are color-coded as in Figure 1. Colored bars

833 mark each eukaryotic species in which the HMM search found a homolog sequence of STING,
834 CD-NTase, or Viperin. Black bar chart shows BUSCO completeness score for each
835 genome/transcriptome.
836
837
838

839
Supplementary Figure 3: Phylogenetic trees from different alignments and tree building
methods show robust topologies
Unrooted maximum likelihood phylogenetic trees generated from two separate alignments
(MUSCLE and MAFFT) and with three different tree inference programs (FastTree, IQtree, and
RaxML-ng). Scale bar of 1 shown beneath each tree represents the number of amino acid
substitutions per position in the underlying alignment. Colored branches show eukaryotic
sequences with the same color scheme as Fig. 1, while gray lines are bacterial sequences. For
the majority of relationships discussed here, we recovered the same tree topology at key nodes
regardless of alignment or tree reconstruction algorithm used. The only exception was in the
STING FastTree phylogenies, wherein the TM-STING clade moved to multiple positions in the
phylogeny, depending on alignment algorithm used.

851
852

853
Supplementary Figure 4: CD-NTase phylogenetic tree
Maximum likelihood phylogenetic tree generated by IQtree of hits from iterative HMM searches
for diverse eukaryotic CD-NTases. Scale bar represents the number of amino acid substitutions

857   per position in the underlying MUSCLE alignment. Ultrafast bootstrap values calculated by
858   IQtree at all nodes with support >70 are shown. Branches with support values <70 were
859   collapsed to polytomies.
860

Supplementary Figure 5: STING phylogenetic tree

Maximum likelihood phylogenetic tree of hits from iterative HMM searches for diverse eukaryotic STING domains. Scale bar represents the number of amino acid substitutions per position in the

underlying MUSCLE alignment. Ultrafast bootstrap values calculated by IQtree at all nodes with

support >70 are shown. Branches with support values <70 were collapsed to polytomies.

Supplementary Figure 6: Viperin phylogenetic tree

Maximum likelihood phylogenetic tree generated by IQtree of hits from iterative HMM searches for diverse eukaryotic Viperins. Scale bar represents the number of amino acid substitutions per position in the underlying MUSCLE alignment. Ultrafast bootstrap values calculated by IQtree at all nodes with support >70 are shown. Branches with support values <70 were collapsed to polytomies.

Supplementary Figure 7: TIR domain of *Crassostrea gigas'* TIR-STING is closely related to metazoan TIR domains

Unrooted maximum likelihood tree of diverse TIR domains. Scale bars on the phylogenetic tree represent the number of amino acid substitutions per position in the underlying MUSCLE alignment. Ultrafast bootstrap values calculated by IQtree at key nodes are shown.

Supplementary Data

Supp. Data Fasta 1: CD-NTase
 Fasta file with all CD-NTase amino acid sequences analyzed.
Supp. Data Fasta 2: STING
 Fasta file with all STING amino acid sequences analyzed.

Supp. Data Fasta 3: Viperin
Fasta file with all Viperin amino acid sequences analyzed.
Supp. Data Table 1: Hmmscan excel file
Hmmscan data for each CD-NTase, STING, and Viperin protein sequence.
Supp. Data Tree 1: CD-NTase
Newick file of maximum likelihood phylogenetic tree generated from a MUSCLE
alignment with IQtree. Newick file is used in Fig. 2 and Supp. Fig. 3 and 4. Node support
values calculated from ultrafast bootstraps.
Supp. Data Tree 2: STING
Newick file of maximum likelihood phylogenetic tree generated from a MUSCLE
alignment with IQtree. Newick file is used in Fig. 3 and Supp. Fig. 3 and 5. Node support values
calculated from ultrafast bootstraps.
Supp. Data Tree 3: Viperin
Newick file of maximum likelihood phylogenetic tree generated from a MUSCLE
alignment with IQtree. Newick file is used in Fig. 4 and Supp. Fig. 3 and 6. Node support values
calculated from ultrafast bootstraps.