1   **Shifts in diversification influence rates of song evolution in honeyeaters**

4   Eleanor M. Hay[1,2]*, Steven L. Chown[1], Matthew D. McGee[1]

7   [1] School of Biological Sciences, Monash University, Melbourne, VIC 3800, Australia

8   [2] Department of Biology, University of Miami, Coral Gables, FL 33146, USA

11   *Corresponding author; email: eleanor.m.hay@outlook.com

ABSTRACT

Birdsongs are acoustic signals that can play a role in sexual selection. Despite the established

role of birdsong in selection and reproductive isolation at microevolutionary scales, the

macroevolutionary relationships between song evolution and clade diversification remain largely

unexplored. We test the hypothesis that shifts in diversification influence rates of song evolution

in honeyeaters, a diverse clade of songbirds restricted to Australasia. Using a song dataset for

163 honeyeater species, we identify lineage-specific shifts in diversification, and test the

influence of these shifts on song evolution with state-dependent, relaxed, multivariate Brownian

motion models. We also explore the underlying patterns of song evolution and phylogenetic

signal of song. Shifts to lower rates of diversification are associated with lower rates of song

evolution. This suggests the two factors are correlated, but is contrary to expectations that

elevated rates of diversification will be associated with elevated rates of song evolution. We also

found that song follows a punctuated mode of evolution, implying that birdsong has an important

role in speciation events, although this signal likely erodes over time and is subject to

phylogenetic constraints.

Keywords: sexual signals, speciation, extinction, macroevolution, RevBayes, MuSSCRat, trait

evolution

INTRODUCTION

Determining what factors cause species diversification is a central goal of many evolutionary studies. The influence of morphological traits (Cooney & Thomas, 2021; Rabosky et al., 2013; Simões et al., 2020) and environmental factors such as ecological opportunity (Dumont et al., 2012; Mahler et al., 2010), on rates of speciation and diversification have been a major focus, but other factors including sexual signals, can also play a role (Maan & Seehausen, 2011; Ritchie, 2007; Wagner et al., 2012). Divergence and development of reproductive isolation are key initial phases in speciation events (Coyne & Orr, 2004; Hernández-Hernández et al., 2021; Jarvis, 2021). Sexual signals frequently emerge as the most divergent traits between closely related species and often experience strong selection (Ritchie, 2007; Servedio & Boughman, 2017). The evolution and divergence of sexual signals can outpace ecological divergence (Arnegard et al., 2010). Consequently, it has been hypothesized that rapid changes in sexual signals could drive species diversification (Maan & Seehausen, 2011; Servedio & Boughman, 2017).

Vocalizations may act as sexual signals that influence speciation. Acoustic signals are used by a wide range of organisms and are complex behavioral traits that enable individuals of the same species to identify one another (Laidre & Johnstone, 2013). Acoustic signals have broad-scale functions in species recognition and fine-scale functions in mate choice, thus potentially influencing speciation (Catchpole & Slater, 2003; Wilkins et al., 2013). Divergence of acoustic signals has played a role in the early stages of speciation in a wide range of organisms including cicadas (Marshall et al., 2008), crickets (Gray & Cade, 2000; Tinghitella et al., 2018; Zuk et al., 2006), frogs (Boul et al., 2007), and birds (Grant & Grant, 2006). In the context of reproductive character displacement, however, the strength of selection on signals depends on range overlap, with sexual signals expected to be under greater divergence when species are

sympatric rather than allopatric (Maan & Seehausen, 2011). Therefore, at macroevolutionary

scales (i.e., clade or family-level), the strength of selection on acoustic signals is expected to

vary between species and among clades. Whether the processes of divergence, assortative

mating, and selection of acoustic traits, which occur at the microevolutionary scale, can have

macroevolutionary consequences for clade diversification, remains unclear.

Birdsong is one of the best-known acoustic signals and is an important sexual signal.

Male birds sing to attract females. Consequently, song can act as an important premating

isolation barrier and influence speciation (Price, 2008; Tobias et al., 2020). Indeed, song

evolution has long thought to be a factor that could explain the diversity of passerine birds

(Baptista & Trail, 1992; Fitzpatrick, 1988). The role of birdsong in speciation and reproductive

isolation has been studied at fine scales: empirical evidence supports the theoretical expectations.

Divergence of song has contributed to reproductive isolation in warblers (Irwin et al., 2001) and

is thought to have influenced the early stages of peripatric speciation in antbirds in Amazonia

(Seddon & Tobias, 2007). Additionally, long-term studies on Darwin's finches have

demonstrated that sexual selection on differences in vocalizations, resulting from changes in

beak morphology, have impacted mate choice and contributed to reproductive isolation and

speciation (Grant & Grant, 2006; Podos, 2001; Podos & Nowicki, 2004). Although acoustic

signals are commonly hypothesized to influence diversification (Wilkins et al., 2013), most

studies have focused on population or sister species comparisons. Empirical studies testing the

influence of signal divergence on speciation at broader scales remain scarce (Mason et al., 2017).

Where the relationship between birdsong and speciation has been investigated at

macroevolutionary scales, bursts in rates of speciation coincide with bursts in song evolution,

however, no significant correlations have been uncovered at the family-level (Mason et al.,

2017).

Indeed, the relationship between diversification and birdsong has proven difficult to

disentangle, largely due to song itself being a complex trait, which in birds is influenced by

developmental mode and constrained by factors such as beak morphology, body size, and the

environment (Derryberry et al., 2018; Friedman et al., 2019; Friis et al., 2022). The majority of

macroevolutionary studies on birdsong have therefore focused on the underlying drivers of song

evolution. Like most sexual signals, acoustic signals are expected to diverge between close

relatives, especially if ranges overlap, potentially leading to a lack of phylogenetic signal in the

trait (Pfennig & Pfennig, 2010). However, song frequency is understood to be phylogenetically

conserved (Mikula et al., 2021), largely due to constraints imposed by morphological traits,

especially body size (Derryberry et al., 2018; Pearse et al., 2018). Additionally, the

developmental mode of song is also thought to influence song evolution and could complicate its

relationship with speciation. Birdsong ranges from learned to innate, and learning is understood

to be a form of phenotypic plasticity that is influenced by cultural transmission and can impact

evolutionary outcomes (Pfennig et al., 2010). Learned songs are typically associated with higher

rates of evolution (Mason et al., 2017) and divergence (Lachlan & Servedio, 2004) than innate

songs, although the evidence is mixed (Freeman et al., 2017). Such complexity in the evolution

of acoustic traits therefore provides a challenge for diversification approaches and can result in

disparity when investigations are conducted over varying phylogenetic and spatial scales.

To overcome such limitations, we test the relationship between birdsong evolution and

diversification using a monophyletic clade restricted to a particular region, leveraging the power

that concentrating on a diverse, single clade can bring (Marquet et al., 2004). Thus, we focus on

honeyeaters (Aves: Meliphagidae), a phenotypically diverse clade of birds (192 species; Gill et

al., 2020) that range throughout Australasia and are known for their distinctive vocalizations

(Higgins et al., 2008). Song in honeyeaters is learned and varies greatly among species, with

some species exhibiting simple and monotonal songs, whereas others have complex and variable

songs (Fig. 1). This variation in vocalizations, combined with the high diversity and endemism

of honeyeaters, makes the group ideal for studying the evolution of vocalizations and their

potential role in species diversification.

122   Specifically, we use honeyeaters as a model clade to explore the relationship between

diversification and birdsong evolution. Using a similar framework to past studies that have

investigated the relationship between song evolution and diversification across bird clades

(Mason et al., 2017), we first test for shifts in honeyeater diversification and then assess if shifts

in diversification are correlated with differential rates of song evolution. We predict that if

birdsong plays a role in diversification, then shifts in diversification will influence rates of song

evolution: species with elevated rates of diversification should have elevated rates of song

evolution. To further disentangle the patterns of song evolution, we also compare fit of different

evolutionary models to assess patterns of song evolution. We assess whether there is evidence of

changes in song occurring at speciation events, which would be expected if song has a role in

speciation. Lastly, we test for phylogenetic signal of song. By doing so, we can assess the

theoretical expectation that if bird song is a sexually selected trait and undergoing selection from

reproductive character displacement, then there will be a lack of phylogenetic signal.

METHODS

Song and phylogenetic data

We used a previously collated dataset of male honeyeater song variables (Hay et al., 2024).

Containing data for 163 honeyeater species, the song dataset considers five aspects of song (Fig.

1; Table S1) that have been explored in past macroevolutionary birdsong studies (e.g.,

Derryberry et al., 2018; Mason & Burns, 2015; Mason et al., 2017): peak frequency (frequency

in which the most sound energy was produced), maximum frequency (maximum frequency of

the highest pitched note in song), minimum frequency (minimum frequency of the lowest pitched

note in song), song bandwidth (maximum frequency - minimum frequency), and song pace

(number of notes/song duration). Song variables were $\log_{10}$-transformed prior to statistical

analysis to meet parametric assumptions of normality and homogeneity of variance, and because

logarithmic scales of sound frequency correspond to how birds perceive and modulate sound

(Cardoso, 2013).

To estimate diversification rates and model trait evolution, we used a honeyeater

phylogeny from Hay et al. (2022). This phylogeny contains all 192 species of honeyeater (Gill et

al., 2020), and was constructed using a combination of nuclear genes, mitochondrial genes, and

topological constraints from previous phylogenomic studies.

RevBayes models

A range of phylogenetic comparative methods implemented through the program RevBayes

(Höhna et al., 2016) were used to test the relationship between diversification and birdsong

evolution. RevBayes enables Bayesian inference for phylogenetic comparative methods, and

uses a probabilistic graphical model framework, which is a powerful generic framework for

specifying and analyzing statistical models (Höhna et al., 2014; Höhna et al., 2016).

First, to explore patterns of diversification in honeyeaters, we estimated branch-specific

speciation and extinction rates (Höhna et al. 2019) using RevBayes (version 1.2.4; Höhna et al.,

2016). This model uses birth-death processes, in which diversification rates vary among

branches, to estimate lineage-specific rates of diversification. Because speciation and extinction

models are subject to bias when species are missing (Mynard et al., 2023), we estimated branch-

specific speciation and extinction rates using the full honeyeater phylogeny containing 192

species (Hay et al., 2022). Estimates of the number of diversification rate shifts are sensitive to

priors (Höhna et al. 2019), therefore we repeated this analysis using several priors on the number

of rate shifts (3, 5, 8, and 10 rate shifts). The MCMC sampler was set for 4 runs of 2,500

generations and a burn-in of 10%. Model convergence was determined using Tracer v1.7.1

(Rambaut et al., 2018) by ensuring ESS were above 200. Output from RevBayes models was

visualized in R (R Core Team, 2013) using the package 'RevGadgets' (Tribble et al., 2021). The

number of shifts in diversification rate and the lineages associated with shifts were identified by

extracting the 'num_shifts' variable in R. Priors had little influence on the number of

diversification rate shifts and species associated with shifts in diversification (Fig. S1-4): we

present the results from the model with 5 rate shifts.

To test the influence of shifts in diversification rates on rate of song evolution, we then

used state-dependent, relaxed, multivariate Brownian motion models of trait evolution

(MuSSCRat; May and Moore 2020) in RevBayes (Höhna et al., 2016). These models enabled us

to test the influence of shifts in diversification identified by the previous model (a discrete

character: species associated with shift in diversification vs not associated with shift in

diversification) on the evolution of the five song variables (multivariate continuous characters:

peak frequency, maximum frequency, minimum frequency, song bandwidth, and song pace). The

MuSSCRat model is advantageous because it jointly estimates the evolutionary history of the

discrete and continuous characters, while also considering background rate variation (variation in

the evolution of the continuous character, not due to the discrete character of interest). This

method permits rates to vary along branches and among continuous characters and reduces

biases in rate estimates. For this analysis, phylogeny was trimmed to the 163 honeyeater species

we have song data for using the '*drop.tip*' function from the ape package (Paradis et al. 2004) in

R (R Core Team, 2013). The MCMC was run for 500,000 generations with a 10% burn-in.

Tracer v1.7.1 (Rambaut et al., 2018) was used to visualize output and determine convergence. To

evaluate the sensitivity of posterior parameter estimates, we repeated the MCMC across different

priors on the number of rate shifts of the continuous characters (5, 10, and 15 shifts). The R

package 'RevGadgets' (Tribble et al., 2021) was used to visualize the state-dependent branch

rates with the '*plotTrace*' function, and summary statistics were calculated using

'*summarizeTrace*' (Table S2). To further explore the underlying patterns of song evolution, we

also extracted the background rates and overall Brownian motion rates of song evolution

estimated from these models.

Correlations between shifts in diversification and rates of song evolution were visualized

using the R package 'phytools' (Revell, 2012).

Kappa models and phylogenetic signal of song

To further explore underlying patterns of song evolution and how this might be linked with

diversification, we compared fit of different evolutionary models and tested for phylogenetic

signal of song.

We fitted different models of trait evolution, specifically testing for trait changes at

speciation events, which would be expected if song influences speciation. We used Kappa

models, which transform branch lengths based on trait changes, and compared the fit of these

models against Brownian motion models in which traits are assumed to evolve at a constant rate.

Higher Kappa values typically indicate gradual evolution, or anagenesis, whereas Kappa values

below 0.5 suggest a punctuated mode of evolution, and that character states change at speciation

events. To fit these models, we used the function '*transformPhylo.ML*' from the 'MOTMOT'

package (Thomas and Freckleton, 2012) in R.

Trait values are predicted to be more similar between closely related species, however

sexual signals are often expected to be the most divergent traits between species (Servedio &

Boughman, 2017). Therefore, we tested how conserved song variables are in honeyeaters. The

phylogenetic signal of each song variable was estimated using Pagel's lambda (Pagel, 1999) and

a likelihood ratio test using the '*phylosig'* function from the 'phytools' package (Revell, 2012) in

R. Pagels lambda is a scaling parameter that ranges from 0 to 1. Using this measure, a lambda of

0 indicates no correlation between species, and a lambda of 1 indicates that the correlation of

trait values between species is equal to expectations of Brownian motion (Revell et al., 2008).

RESULTS

Branch-specific speciation and extinction models identified several patterns of honeyeater

diversification. Overall, there were two diversification rate categories found for honeyeaters.

Much of the tree follows a relatively constant rate of diversification; however, the model

identifies several species associated with a rate shift to a lower rate of diversification (Fig. 2).

Species associated with shifts in diversification are typically from the more basal linages within

each clade, many of which are monotypic genera (e.g., *Glycicharea fallax*, *Certhionyx*

variegatus, *Oreornis chrysogenys*, *Purnella albifrons*).

MuSSCRat models revealed that shifts in diversification have a significant influence on

rates of honeyeater song evolution (posterior probability = 1.0). This was consistent across

different priors (Fig. S5-7). The state-dependent branch rates show that shifts to lower rates of

diversification are associated with lower rates of song evolution (Fig. 3a). There is no overlap of

confidence intervals and over a 10-fold difference in song evolution rate estimates between these

two diversification rate categories (mean rate of song evolution$_{\text{diversification shift}}$ = 0.17, 95 % CI =

[0.091, 0.31]; mean rate of song evolution$_{\text{no diversification shift}}$ = 1.83, 95 % CI = [1.69, 1.91]; Fig.

3c).

The overall Brownian motion rates of song evolution extracted from MuSSCRat models

uncover further patterns of song evolution (Fig. 3b). Most of the phylogeny exhibits a relatively

constant rate of birdsong evolution, however increased rates of evolution are notable in some

young sister species. For example, *Philemon kisserensis* and *Philemon citreogularis* are

associated with increased rates of song evolution, as well as *Lichmera limbata* and *Lichmera*

indistincta, and *Manorina melanocephala*, *Manorina melanotis*, and *Manorina flavigula* (Fig.

3b). *Sugomel niger* is also identified as having high overall rates of song evolution.

Comparison of different trait evolution models provides further insight into honeyeater

song evolution. The Kappa models perform better than the Brownian motion models in all cases

(Table S3). Values of Kappa extracted from models are all below 0.5, suggesting that song

follows a punctuated mode of evolution, with changes in trait values being associated with

cladogenesis. Maximum frequency returned the lowest Kappa statistic (0.34), followed by peak

frequency (0.35), song pace (0.38), bandwidth (0.46), and minimum frequency (0.50).

Lastly, we found that the phylogenetic signal varied substantially among the five song

variables. Values of Pagel's lambda ranged from 0.31 to 0.92 (Table S4). Song bandwidth had the

lowest signal ($\lambda = 0.31$), followed by maximum frequency ($\lambda = 0.58$), pace ($\lambda = 0.66$), peak

frequency ($\lambda = 0.75$) and then minimum frequency ($\lambda = 0.92$).

DISCUSSION

We found that shifts in rates of diversification for honeyeaters are correlated with significant

differences in rates of song evolution. Theory suggests that if song has a role in speciation and

diversification of clades then bursts of diversification should correlate with bursts in song

evolution (Wilkins et al., 2013). Past studies following a similar approach to the one adopted

here found relationships between bursts in song evolution and speciation across 581 species of

tanagers and ovenbirds, but this was not significant when each family was considered separately

(Mason et al., 2017). We found that shifts to a lower rate of diversification are significantly

correlated with lower rates of song evolution, providing evidence that song evolution and

diversification are associated within families.

The directionality of this finding is, however, unusual given that expectations are

typically for increased rates of diversification to be associated with increased rates of song

evolution if song is acting as an isolating mechanism (Baptista & Trail, 1992; Price, 2008; Tobias

et al., 2020). Honeyeater species associated with shifts to lower diversification rates are

comparatively older lineages to the rest of the tree and consist of many monotypic genera. Older

lineages have more time to accumulate species, but they also have more time for extinction. In

birds, rates of speciation tend to slow over time as niches become filled (Ricklefs 2006; Price

2008). The lower rates of diversification we observe in these older lineages have likely resulted

from speciation rates decreasing and extinction remaining constant over time. It's important to

note that the directionality of our finding is partly due to the way that the analyses have been

conducted: the diversification model found shifts to a lower rate of speciation, and this was

correlated to lower rates of song evolution. Using this approach there was technically no way to

relate shifts to higher diversification rates (since we did not find these) to higher rates of song

evolution. Nonetheless, we still find two overall rate categories of diversification and song

evolution in honeyeaters, in which lower rates of diversification are correlated with lower rates

of song evolution and higher rates of diversification are correlated with higher rates of song

evolution.

Our results do raise the intriguing question of what might cause rates of song evolution to

slow down in honeyeaters. At the lineage level, complexity and diversity of honeyeater songs is

known to be impacted by population density. For example, the critically endangered Regent

Honeyeater (*Anthochaera phrygia*) has had the complexity of their songs decline for decades

(Crates et al. 2021). Song in honeyeaters is learned and young juveniles must co-occur with other

breeding adult males to learn songs (Crates et al. 2021). Regent Honeyeaters are becoming so

rare that young birds are unable to learn their own songs. This has consequences for mating

success of individuals since males sing to attract female birds and can cause further population

declines and even extinction. One explanation for our findings may be that such processes at the

population level can scale up in the extinction-speciation interaction to have the

macroevolutionary consequences on clade richness we observe in our study.

Exploring underlying patterns of song evolution enabled us to further unveil the

relationship between song evolution and diversification. We find evidence that song follows a

punctuated rather than gradual model of evolution, suggesting that changes in honeyeater song

are important for speciation events. Overall Brownian motion rates of song evolution extracted

from MuSSCRat models (Fig. 3b) corroborate this punctuated mode of song evolution and show

increased rates of song evolution associated with some recent speciation events, however, we do

not see increased rates with older speciation events. One possible explanation is that past signal

of song divergence and evolution at speciation events has eroded over time, which could be why

we observe increased rates of song evolution in young sister species. In general, success in

linking micro with macroevolution has been variable (Schluter 2024): past studies linking

birdsong and diversification have had mixed outcomes. For example, a study of 175 taxon pairs

of New World passerine birds found no association between faster rates of song evolution and

higher speciation rates (Freeman et al., 2022). Schluter (2024) suggests that difficulties in linking

rates of reproductive trait evolution and species diversification could be because present day trait

values might not accurately represent the past. It has been proposed that initial rate of species

accumulation could be strongly influenced by the rate of evolution of reproductive isolation,

however, over time other ecological processes that influence clade diversity intervene (Schluter,

2024). This could provide an explanation for mixed outcomes from past studies, and why the

overall Brownian motion rates of song evolution only show elevated rates in young species.

The overall Brownian motion rates of evolution highlight elevated rates of song evolution

in *Sugomel niger* and three young sister species groups: The Gray Friarbird (*Philemon*

kisserensis) and the Little Friarbird (*P. citreogularis*); The Indonesian Honeyeater (*Lichmera*

limbata) and The Brown Honeyeater (*L. indistincta*); and the Noisy Miner (*Manorina*

melanocephala), Black-eared Miner (*M. melanotis*), and Yellow-throated Miner (*M. flavigula*).

Songs are thought to diverge slowly in allopatry, and only rapidly increase on secondary contact,

contributing to reinforcement (Lachlan & Servedio, 2004; Servedio & Boughman, 2017).

Interestingly, the *Lichmera* and *Philemon* species are not currently sympatric, but do have

adjacent ranges to some extent: The Indonesian Honeyeater ranges throughout Timor and the

Lesser Sunda Islands, whereas the Brown Honeyeater is found across Australia, New Guinea,

and other parts of Indonesia (Higgins et al., 2022), while the little friarbird ranges on Australia

and New Guinea (Higgins et al., 2020a) and the gray friarbird is restricted to small islands

offshore of East Timor (Higgins et al., 2020b). In saying that, honeyeaters have high dispersal

ability, and there has been shifting biomes on mainland Australia and connection and

disconnection of Australia and New Guinea (Andersen et al., 2014; Hay et al., 2022; Marki et al.,

2017). Furthermore, Peñalba et al. (2018) have shown that population structure and geneflow is

very dynamic in the little friarbird species complex and the brown honeyeater species complex

and does not adhere to geographic barriers across northern Australia and New Guinea. Thus,

these species have likely had contact during the speciation process, and the increased rates of

song evolution we observe in these species could have contributed to divergence. The three

Manorina species all have overlapping ranges across Australia: The Yellow-throated Miner

ranges across nearly all of Australia, except for some areas along the East Coast (Higgins et al.,

2020c); The Noisy Miner ranges along a large proportion of the East side of Australia (Higgins

et al., 2020d); and the Black-eared Miner has the smallest range but is found inland around the

intersection between South Australia, Victoria, and New South Wales (Higgins et al., 2020e).

These species arguably have the highest rates of song evolution and could imply that sympatry

has contributed to divergence of song. As an important caveat to this study, many of these effects

are influenced by how species are classified. For example, the Indonesian Honeyeater was

previously identified as a distinct species (Gill et al. 2020) but has now been synonymized with

the Brown Honeyeater (Gill et al. 2023). How one classifies species and subspecies can influence

outcomes of macroevolutionary analysis, although this is likely to have little effect in this

context, given the strength of the relationship we found, and that our models controlled for such

potential background rate variation.

We also find that song exhibits phylogenetic signal, and that most of the interspecific

diversity of song frequency and pace in honeyeaters can be explained by evolutionary history.

Theory suggests that divergence is expected for sexually selected traits between closely related

species (Maan & Seehausen, 2011; Servedio & Boughman, 2017). Thus, the level of

phylogenetic signal uncovered for frequency and pace is contrary to such theoretical

expectations. This is unsurprising, however, given other studies have found similar levels of

phylogenetic signal for song traits in birds: peak frequency was found to have a lambda of 0.87

for 5085 species of passerine birds (Mikula et al., 2021), and a lambda of 0.74 for 1022 passerine

species (Friis et al., 2022). Our results are consistent with these previous studies and confirm that

peak frequency of song is phylogenetically conserved. Phylogenetic conservatism of song is, to

some extent, driven by allometric constraints imposed by body size (Fletcher, 2004). Body size

itself is a conserved trait that directly impacts the length of the vocal tract and the size of the

syrinx (Bertelli & Tubaro, 2002; Suthers & Zollinger, 2008; Friis et al. 2021), and is a driver of

honeyeater song evolution (Hay et al., 2024). These other underlying processes likely contribute

to the difficulty in disentangling the links between song evolution and diversification.

Nonetheless, we find some evidence that song evolution has contributed to diversification

within this family of birds. Understanding the influence of reproductive isolation on

diversification has been recognized as an urgent question in evolutionary biology (Matute &

Cooper, 2021), especially because the evolution of reproductive isolation often defines

speciation (Coyne & Orr, 2004). Here, we contribute to understanding of this question by testing

the hypothesis that shifts in diversification are associated with increased rates of song evolution

in honeyeaters. We found that shifts to lower rates of diversification are correlated with lower

rates of song evolution, and that song follows a punctuated model of evolution rather than

gradual, suggesting that changes in song occur at speciation events and this is followed by

periods of stasis. Ultimately, we find that song evolution and diversification are correlated in

honeyeaters, and changes in song play an important role in speciation events, however this signal

likely erodes over time due to underlying constraints of song evolution and other ecological

processes.

REFERENCES

Andersen, M. J., Naikatini, A., & Moyle, R. G. (2014). A molecular phylogeny of Pacific

honeyeaters (Aves: Meliphagidae) reveals extensive paraphyly and an isolated Polynesian

radiation. *Molecular Phylogenetics and Evolution, 71*, 308-315.

https://doi.org/10.1016/j.ympev.2013.11.014

Arnegard, M. E., McIntyre, P. B., Harmon, L. J., Zelditch, M. L., Crampton, W. G., Davis, J. K.,

Sullivan, J. P., Lavoue, S., & Hopkins, C. D. (2010). Sexual signal evolution outpaces

ecological divergence during electric fish species radiation. *The American Naturalist,*

176(3), 335-356. https://doi.org/10.1086/655221

Baptista, L. F., & Trail, P. W. (1992). The role of song in the evolution of passerine diversity.

Systematic Biology, 41(2), 242-247. https://doi.org/10.2307/2992524

Bertelli, S., & Tubaro, P. L. (2002). Body mass and habitat correlates of song structure in a

primitive group of birds. *Biological Journal of the Linnean Society, 77*, 423-430.

https://doi.org/10.1046/j.1095-8312.2002.00112.x

Boul, K. E., Funk, W. C., Darst, C. R., Cannatella, D. C., & Ryan, M. J. (2007). Sexual selection

drives speciation in an Amazonian frog. *Proceedings of the Royal Society B: Biological*

Sciences, 274(1608), 399-406. https://doi.org/10.1098/rspb.2006.3736

Cardoso, G. C. (2013). Using frequency ratios to study vocal communication. *Animal Behaviour,*

85(6), 1529-1532. https://doi.org/10.1016/j.anbehav.2013.03.044

Catchpole, C. K., & Slater, P. J. B. (2003). *Bird Song: Biological Themes and Variations*.

Cambridge University Press.

Cooney, C. R., & Thomas, G. H. (2021). Heterogeneous relationships between rates of speciation

and body size evolution across vertebrate clades. *Nature Ecology and Evolution, 5*(1),

101-110. https://doi.org/10.1038/s41559-020-01321-y

Coyne, J. A., & Orr, H. A. (2004). *Speciation*. Sinauer Associates.

Crates, R., Langmore, N., Ranjard, L., Stojanovic, D., Rayner, L., Ingwersen, D., & Heinsohn, R.

(2021). Loss of vocal culture and fitness costs in a critically endangered

songbird. *Proceedings of the Royal Society B*, *288*(1947), 20210225.

https://doi.org/10.1098/rspb.2021.0225

Derryberry, E. P., Seddon, N., Derryberry, G. E., Claramunt, S., Seeholzer, G. F., Brumfield, R.

T., & Tobias, J. A. (2018). Ecological drivers of song evolution in birds: disentangling the

effects of habitat and morphology. *Ecology and Evolution, 8*(3), 1890-1905.

https://doi.org/10.1002/ece3.3760

Dumont, E. R., Davalos, L. M., Goldberg, A., Santana, S. E., Rex, K., & Voigt, C. C. (2012).

Morphological innovation, diversification and invasion of a new adaptive zone.

Proceedings of the Royal Society B: Biological Sciences, 279(1734), 1797-1805.

https://doi.org/10.1098/rspb.2011.2005

Fitzpatrick, J. W. (1988). Why so many passerine birds? A response to Raikow. *Systematic

Zoology, 37*(1), 71-76. https://doi.org/10.2307/2413195

Fletcher, N. H. (2004). A simple frequency-scaling rule for animal communication. *The Journal

of the Acoustical Society of America, 115*(5), 2334-2338.

https://doi.org/10.1121/1.1694997

Freeman, B. G., Montgomery, G. A., & Schluter, D. (2017). Evolution and plasticity: divergence

of song discrimination is faster in birds with innate song than in song learners in

423   Neotropical passerine birds. *Evolution, 71*(9), 2230-2242.

424   https://doi.org/10.1111/evo.13311

425 Freeman, B. G., Rolland, J., Montgomery, G. A., & Schluter, D. (2022). Faster evolution of a

426   premating reproductive barrier is not associated with faster speciation rates in New World

427   passerine birds. *Proceedings of the Royal Society B: Biological Sciences, 289*(1966),

428   20211514. https://doi.org/10.1098/rspb.2021.1514

429 Friedman, N. R., Miller, E. T., Ball, J. R., Kasuga, H., Remes, V., & Economo, E. P. (2019).

430   Evolution of a multifunctional trait: shared effects of foraging ecology and

431   thermoregulation on beak morphology, with consequences for song evolution.

432   *Proceedings of the Royal Society B: Biological Sciences, 286*(1917), 20192474.

433   https://doi.org/10.1098/rspb.2019.2474

434 Friis, J. I., Sabino, J., Santos, P., Dabelsteen, T., & Cardoso, G. C. (2021). The allometry of

435   sound frequency bandwidth in songbirds. *The American Naturalist, 197*(5), 607-614.

436   https://doi.org/10.1086/713708

437 Friis, J. I., Sabino, J., Santos, P., Dabelsteen, T., Cardoso, G. C., & Jennions, M. D. (2022).

438   Ecological adaptation and birdsong: how body and bill sizes affect passerine sound

439   frequencies. *Behavioral Ecology, 33*(4), 798-806. https://doi.org/10.1093/beheco/arac042

440 Gill, F., Donsker, D., & Rasmussen, P. (2020). *IOC World Bird List* (v 10.2).

441   https://doi.org/10.14344/IOC.ML.10.2.

442 Gill, F., D. Donsker, and P. Rasmussen. 2023. IOC World Bird List (v 13.1).

443 Grant, P. R., & Grant, B. R. (2006). Evolution of character displacement in Darwin's finches.

444   *Science, 313*, 224-226. https://doi.org/10.1126/science.1128374

Gray, D. A., & Cade, W. H. (2000). Sexual selection and speciation in field crickets. *Proceedings*

of the National Academy of Sciences of the USA, 97, 14449-14454.

https://doi.org/10.1073/pnas.97.26.14449

Harvey, M. G., Singhal, S., & Rabosky, D. L. (2019). Beyond reproductive isolation:

demographic controls on the speciation process. *Annual Review of Ecology, Evolution,*

and Systematics, 50(1), 75-95. https://doi.org/10.1146/annurev-ecolsys-110218-024701

Hay, E. M., McGee, M. D., & Chown, S. L. (2022). Geographic range size and speciation in

honeyeaters. *BMC Ecology and Evolution, 22*, 1-14. https://doi.org/10.1186/s12862-022-

02041-6

Hay, E. M., McGee, M. D., White, C. R., & Chown, S. L. (2024). Body size shapes song in

honeyeaters. *Proceedings of the Royal Society B: Biological Sciences*. https://doi.org/

10.1098/rspb.2024.0339

Hernández-Hernández, T., Miller, E. C., Roman-Palacios, C., & Wiens, J. J. (2021). Speciation

across the Tree of Life. *Biological Reviews, 96*(4), 1205-1242.

https://doi.org/10.1111/brv.12698

Higgins, P. J., Christidis, L., & Ford, H. (2020a). Little Friarbird (*Philemon citreogularis*),

version 1.0. In J. del Hoyo, A. Elliott, J. Sargatal, D. A. Christie, and E. de Juana (Eds.),

Birds of the World. Cornell Lab of Ornithology. https://doi.org/10.2173/bow.litfri1.01

Higgins, P. J., Christidis, L., & Ford, H. (2020b). Gray Friarbird (*Philemon kisserensis*), version

1.0. In J. del Hoyo, A. Elliott, J. Sargatal, D. A. Christie, & E. de Juana (Eds.), *Birds of*

the World. Cornell Lab of Ornithology. https://doi.org/10.2173/bow.gryfri1.01

Higgins, P. J., Christidis, L., & Ford, H. (2020c). Yellow-throated Miner (*Manorina flavigula*),

version 1.0. In J. del Hoyo, A. Elliott, J. Sargatal, D. A. Christie, & E. de Juana (Eds.).

Birds of the World. Cornell Lab of Ornithology. https://doi.org/10.2173/bow.yetmin1.01

Higgins, P. J., Christidis, L., & Ford, H. (2020d). Noisy Miner (*Manorina melanocephala*),

version 1.0. In J. del Hoyo, A. Elliott, J. Sargatal, D. A. Christie, & E. de Juana (Eds.).

Birds of the World. Cornell Lab of Ornithology. https://doi.org/10.2173/bow.noimin1.01

Higgins, P. J., Christidis, L., & Ford, H. (2020e). Black-eared Miner (*Manorina melanotis*),

version 1.0. In J. del Hoyo, A. Elliott, J. Sargatal, D. A. Christie, & E. de Juana (Eds.).

Birds of the World. Cornell Lab of Ornithology. https://doi.org/10.2173/bow.blemin1.01

Higgins, P. J., Christidis, L., & Ford, H. (2022). Brown Honeyeater (*Lichmera indistincta*),

version 1.1. In S. M. Billerman (Ed.), *Birds of the World*. Cornell Lab of Ornithology,

Ithaca, NY, USA. https://doi.org/10.2173/bow.brohon1.01.1

Higgins, P. J., Ford, H. A., & Christidis, L. (2008). Family Meliphagidae. In J. del Hoyo, A.

Elliott, & D. A. Christie (Eds.), *Handbook of the Birds of the World: Penduline-tits to

shrikes* (Vol. 13, pp. 498–691). Lynx Edicions.

Höhna, S., Heath, T. A., Boussau, B., Landis, M. J., Ronquist, F., & Huelsenbeck, J. P. (2014).

Probabilistic graphical model representation in phylogenetics. *Systematic Biology, 63*(5),

753-771. https://doi.org/10.1093/sysbio/syu039

Höhna, S., Landis, M. J., Heath, T. A., Boussau, B., Lartillot, N., Moore, B. R., Huelsenbeck, J.

P., & Ronquist, F. (2016). RevBayes: Bayesian phylogenetic inference using graphical

models and an interactive model-specification language. *Systematic Biology, 65*(4), 726-

736. https://doi.org/10.1093/sysbio/syw021

Höhna, S., Freyman, W. A., Nolen, Z., Huelsenbeck, J. P., May, M. R., and Moore, B. R. (2019).

A Bayesian approach for estimating branch-specific speciation and extinction rates.

bioRxiv. https://doi.org/10.1101/555805

Irwin, D. E., Bensch, S., & Price, T. D. (2001). Speciation in a ring. *Nature, 409*, 333-337.

https://doi.org/10.1038/35053059

Jarvis, E. D. (2021). At the beginning of speciation. *Science, 371*(6536), 1312.

https://doi.org/10.1126/science.abg5454

Lachlan, R. F., & Servedio, M. R. (2004). Song learning accelerates allopatric speciation.

Evolution, 58(9), 2049-2063. https://doi.org/10.1111/j.0014-3820.2004.tb00489.x

Laidre, M. E., & Johnstone, R. A. (2013). Animal signals. *Current Biology, 23*(18), R829-833.

https://doi.org/10.1016/j.cub.2013.07.070

Maan, M. E., & Seehausen, O. (2011). Ecology, sexual selection and speciation. *Ecology Letters,*

14(6), 591-602. https://doi.org/10.1111/j.1461-0248.2011.01606.x

Mahler, D. L., Revell, L. J., Glor, R. E., & Losos, J. B. (2010). Ecological opportunity and the

rate of morphological evolution in the diversification of Greater Antillean anoles.

Evolution, 64(9), 2731-2745. https://doi.org/10.1111/j.1558-5646.2010.01026.x

Marki, P. Z., Jonsson, K. A., Irestedt, M., Nguyen, J. M. T., Rahbek, C., & Fjeldsa, J. (2017).

Supermatrix phylogeny and biogeography of the Australasian Meliphagides radiation

(Aves: Passeriformes). *Molecular Phylogenetics and Evolution, 107*, 516-529.

https://doi.org/10.1016/j.ympev.2016.12.021

Marquet, P. A., Fernández, M., Navarette, A. A., & Valdovinos, C. (2004). Diversity emerging:

towards a deconstruction of biodiversity patterns. In L. M. & H. L. (Eds.), *Frontiers of*

biogeography: New directions in the geography of nature (pp. 191-209). Sinauer

Associates.

Marshall, D. C., Slon, K., Cooley, J. R., Hill, K. B. R., & Simon, C. (2008). Steady Plio-

Pleistocene diversification and a 2-million-year sympatry threshold in a New Zealand

cicada radiation. *Molecular Phylogenetics and Evolution, 48*(3), 1054-1066.

https://doi.org/10.1016/j.ympev.2008.05.007

Mason, N. A., & Burns, K. J. (2015). The effect of habitat and body size on the evolution of

vocal displays in Thraupidae (tanagers), the largest family of songbirds. *Biological*

Journal of the Linnean Society, 114, 538-551. https://doi.org/10.1111/bij.12455

Mason, N. A., Burns, K. J., Tobias, J. A., Claramunt, S., Seddon, N., & Derryberry, E. P. (2017).

Song evolution, speciation, and vocal learning in passerine birds. *Evolution, 71*(3), 786-

796. https://doi.org/10.1111/evo.13159

Matute, D. R., & Cooper, B. S. (2021). Comparative studies on speciation: 30 years since Coyne

and Orr. *Evolution, 75*(4), 764-778. https://doi.org/10.1111/evo.14181

May, M. R., & Moore, B. R. 2020. A Bayesian approach for inferring the impact of a discrete

character on rates of continuous-character evolution in the presence of background-rate

variation. *Systematic Biology*, 69:530–544. https://doi.org/10.1093/sysbio/syz069

Mikula, P., Valcu, M., Brumm, H., Bulla, M., Forstmeier, W., Petruskova, T., Kempenaers, B., &

Albrecht, T. (2021). A global analysis of song frequency in passerines provides no

support for the acoustic adaptation hypothesis but suggests a role for sexual selection.

Ecology Letters, 24(3), 477-486. https://doi.org/10.1111/ele.13662

Mynard, P., Algar, A., Lancaster, L., Bocedi, G., Fahri, F., Gubry-Rangin, C., Lupiyaningdyah,

P., Nangoy, M., Osborne, O., Papadopulos, A., Sudiana, I. M., Juliandi, B., Travis, J., &

Herrera-Alsina, L. (2023). Impact of phylogenetic tree completeness and misspecification

534   of sampling fractions on trait dependent diversification models. *Systematic Biology*.

535   https://doi.org/10.1093/sysbio/syad001

536   Oliveros, C. H., Field, D. J., Ksepka, D. T., Barker, F. K., Aleixo, A., Andersen, M. J., Alstrom,

537   P., Benz, B. W., Braun, E. L., Braun, M. J., Bravo, G. A., Brumfield, R. T., Chesser, R. T.,

538   Claramunt, S., Cracraft, J., Cuervo, A. M., Derryberry, E. P., Glenn, T. C., Harvey, M. G.,

539   Hosner, P. A., Joseph, L., Kimball, R. T., Mack, A. L., Miskelly, C. M., Peterson, A. T.,

540   Robbins, M. B., Sheldon, F. H., Silveira, L. F., Smith, B. T., White, N. D., Moyle, R. G.,

541   & Faircloth, B. C. (2019). Earth history and the passerine superradiation. *Proceedings of*

542   *the National Academy of Sciences of the USA, 116*(16), 7916-7925.

543   https://doi.org/10.1073/pnas.1813206116

544   Pagel, M. (1999). Inferring the historical patterns of biological evolution. *Nature, 401*(6756),

545   877-884.

546   Paradis, E., Claude, J., & Strimmer, K. (2004). APE: analyses of phylogenetics and evolution in

547   R language. *Bioinformatics, 20*(2), 289-290.

548   https://doi.org/10.1093/bioinformatics/btg412

549   Pearse, W. D., Morales-Castilla, I., James, L. S., Farrell, M., Boivin, F., & Davies, T. J. (2018).

550   Global macroevolution and macroecology of passerine song. *Evolution, 72*(4), 944-960.

551   https://doi.org/10.1111/evo.13450

552   Peñalba, J. V., Joseph, L., & Moritz, C. (2019). Current geography masks dynamic history of

553   gene flow during speciation in northern Australian birds. *Molecular Ecology,* 28, 630–

554   643. https://doi.org/10.1111/mec.14978

Pfennig, D. W., & Pfennig, K. S. (2010). Character displacement and the origins of diversity. *The*
American Naturalist, 176, 26-44. https://doi.org/10.1086/657056

Pfennig, D. W., Wund, M. A., Snell-Rood, E. C., Cruickshank, T., Schlichting, C. D., & Moczek,
A. P. (2010). Phenotypic plasticity's impacts on diversification and speciation. *Trends in*
Ecology & Evolution, 25(8), 459-467. https://doi.org/10.1016/j.tree.2010.05.006

Podos, J. (2001). Correlated evolution of morphology and vocal signal structure in Darwin's
fnches. *Nature, 409*, 185-188. https://doi.org/10.1038/35051570

Podos, J., & Nowicki, S. (2004). Beaks, adaptation, and vocal evolution in Darwin's finches.
Bioscience, 54, 501-510. https://doi.org/10.1641/0006-
3568(2004)054[0501:BAAVEI]2.0.CO;2

Price, T. (2008). *Speciation in Birds*. Roberts and Co.

R Core Team. (2013). R: a language and environment for statistical computing. https://www.r-
project.org/

Rabosky, D. L., Santini, F., Eastman, J., Smith, S. A., Sidlauskas, B., Chang, J., & Alfaro, M. E.
(2013). Rates of speciation and morphological evolution are correlated across the largest
vertebrate radiation. *Nature Communications, 4*, 1958.
https://doi.org/10.1038/ncomms2958

Rambaut, A., Drummond, A. J., Xie, D., Baele, G., & Suchard, M. A. (2018). Posterior
summarization in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology, 67*(5),
901-904. https://doi.org/10.1093/sysbio/syy032

Revell, L. J. (2012). phytools: an R package for phylogenetic comparative biology (and other
things). *Methods in Ecology and Evolution, 3*(2), 217-223. https://doi.org/10.1111/j.2041-
210X.2011.00169.x

Revell, L. J., Harmon, L. J., & Collar, D. C. (2008). Phylogenetic signal, evolutionary process,

and rate. *Systematic Biology, 57*(4), 591-601.

https://doi.org/10.1080/10635150802302427

Ricklefs, R. E. (2006). Global variation in the diversification rate of passerine birds. *Ecology*,

87(10), 2468-2478. https://doi.org/10.1890/0012-9658(2006)87[2468:GVITDR]2.0.CO;2

Ritchie, M. G. (2007). Sexual selection and speciation. *Annual Review of Ecology, Evolution,*

and Systematics, 38, 79-102. https://doi.org/10.1146/annurev.ecolsys.38.091206.095733

Schluter, D. 2024. Variable success in linking micro and macroevolution. *Evolutionary Journal*

of the Linnean Society, kzae016. https://doi.org/10.1093/evolinnean/kzae016

Seddon, N., & Tobias, J. A. (2007). Song divergence at the edge of Amazonia: an empirical test

of the peripatric speciation model. *Biological Journal of the Linnean Society, 90*, 173-

188. https://doi.org/10.1111/j.1095-8312.2007.00753.x

Servedio, M. R., & Boughman, J. W. (2017). The role of sexual selection in local adaptation and

speciation. *Annual Review of Ecology, Evolution, and Systematics, 48*, 85-109.

https://doi.org/10.1146/annurev-ecolsys-110316-022905

Simões, T. R., Vernygora, O., Caldwell, M. W., & Pierce, S. E. (2020). Megaevolutionary

dynamics and the timing of evolutionary innovation in reptiles. *Nature Communications,*

11(1), 3322. https://doi.org/10.1038/s41467-020-17190-9

Singhal, S., Huang, H., Grundler, M. R., Marchan-Rivadeneira, M. R., Holmes, I., Title, P. O.,

Donnellan, S. C., & Rabosky, D. L. (2018). Does population structure predict the rate of

speciation? a comparative test across Australia's most diverse vertebrate radiation. *The*

American Naturalist, 192(4), 432-447. https://doi.org/10.1086/699515

Suthers, R. A., & Zollinger, S. A. (2008). From brain to song: the vocal organ and vocal tract. In

P. H. Zeigler & P. Marler (Eds.), *Neuroscience of Birdsong* (pp. 78-98). Cambridge

University Press.

Tinghitella, R. M., Broder, E. D., Gurule-Small, G. A., Hallagan, C. J., & Wilson, J. D. (2018).

Purring crickets: the evolution of a novel sexual signal. *The American Naturalist, 192*(6),

773-782. https://doi.org/10.1086/700116

Thomas, G. H., and R. P. Freckleton. 2012. MOTMOT: models of trait macroevolution on trees.

Methods in Ecology and Evolution, 3:145–151. https://doi.org/10.1111/j.2041-

210X.2011.00132.x

Tobias, J. A., Ottenburghs, J., & Pigot, A. L. (2020). Avian diversity: speciation, macroevolution,

and ecological function. *Annual Review of Ecology, Evolution, and Systematics, 51*(1),

533-560. https://doi.org/10.1146/annurev-ecolsys-110218-025023

Tribble, C. M., Freyman, W. A., Landis, M. J., Lim, J. Y., Barido-Sottani, J., Kopperud, B. T.,

Höhna, S., & May, M. R. (2021). RevGadgets: an R package for visualizing Bayesian

phylogenetic analyses from RevBayes. *Methods in Ecology and Evolution, 13*(2), 314-

323. https://doi.org/10.1111/2041-210x.13750

Wagner, C. E., Harmon, L. J., & Seehausen, O. (2012). Ecological opportunity and sexual

selection together predict adaptive radiation. *Nature, 487*(7407), 366-369.

https://doi.org/10.1038/nature11144

Wilkins, M. R., Seddon, N., & Safran, R. J. (2013). Evolutionary divergence in acoustic signals:

causes and consequences. *Trends in Ecology & Evolution, 28*(3), 156-166.

https://doi.org/10.1016/j.tree.2012.10.002

Zuk, M., Rotenberry, J. T., & Tinghitella, R. M. (2006). Silent night: adaptive disappearance of a

sexual signal in a parasitized population of field crickets. *Biology Letters, 2*(4), 521-524.

https://doi.org/10.1098/rsbl.2006.0539

FIGURE CAPTIONS

Figure 1. Song spectrograms displaying frequency (kHz) over time (s) for five species of

honeyeater: a) New Zealand Bellbird (*Anthornis melanura*; xeno-canto accession number =

XC153005), b) Eastern Spinebill (*Acanthorhynchus tenuirostris;* xeno-canto accession number =

XC490800), c) Blue-faced Honeyeater (*Entomyzon cyanotis*; xeno-canto accession number =

XC439288), d) Noisy Friarbird (*Philemon corniculatus*; xeno-canto accession number =

XC287056), and e) Singing Honeyeater (*Gavicalis virescens*; xeno-canto accession number =

XC334251). The five song variables considered in this study are indicated with annotations in a):

peak frequency represents the frequency of maximum amplitude across the entire song,

maximum frequency is the highest pitched note in song, minimum frequency is the lowest

pitched note in song, bandwidth is the difference between maximum and minimum frequency

across the song, and pace is the number of notes over song duration.

Figure 2. Lineage-specific net diversification rates for honeyeaters. This is modeled across the

full phylogeny of 192 honeyeater species. Branches on the phylogeny are colored according to

diversification rate, with the warmer colors representing high rates of diversification and cooler

colors representing lower rates of diversification. Lineages associated with shifts in

diversification are indicated with the dots at the tips of the phylogeny.

Figure 3. Patterns of honeyeater song evolution extracted from MuSSCRat models. Here,

analysis was completed using the 163 honeyeater species we had song data for. Branches on the

phylogeny on the left (a) indicate state-specific branch rates. The states tested are indicated with

the dots at tip labels, these are the species that were found to have shifts in diversification. Rates

of song evolution for the two states are displayed in the density plot (c). The phylogeny on the

right (b) displays the overall Brownian motion rates of song evolution extracted from models.

691 **FIGURES**

693 **Figure 1.**

704 **Figure 2.**

Figure 3.

720