1 **No evidence for song evolution influencing speciation patterns in honeyeaters**

4 Eleanor M. Hay[1], Steven L. Chown[1], Matthew D. McGee[1]

7 [1] School of Biological Sciences, Monash University, Melbourne, VIC 3800, Australia

10 *Corresponding author; email: eleanor.m.hay@outlook.com

**Abstract**

Birdsongs are well-known acoustic signals and are thought to play a key role in sexual

selection. Despite the established role of birdsong in sexual selection and reproductive

isolation at microevolutionary scales, its macroevolutionary impacts on speciation and

diversification remain largely unexplored. Here, we test the hypothesis that song influences

speciation on macroevolutionary scales, using honeyeaters, a diverse clade of songbirds

restricted to Australasia. Using song data for 163 honeyeater species, we employ

phylogenetic comparative models of trait evolution to detect shifts in optima for song

evolution and use hidden state speciation and extinction models to assess the influence of

these shifts on diversification. Contrary to theoretical expectations, we find no evidence that

song influences speciation or diversification rates, and models suggest other factors play a

role. Overall, song does not significantly contribute to speciation in honeyeaters, likely due to

other ecological factors including range size and dispersal ability.

Keywords: sexual signals, birdsong, speciation, macroevolution, RevBayes, trait evolution

## Introduction

Determining what factors cause species diversification is a central goal of many evolutionary studies. The influence of morphological traits (Rabosky et al. 2013; Simões et al. 2020; Cooney and Thomas 2021) and environmental factors, such as ecological opportunity (Mahler et al. 2010; Dumont et al. 2012), on rates of speciation and diversification have been a major focus, but other factors, including sexual signals, can also play a role (Ritchie 2007; Maan and Seehausen 2011; Wagner et al. 2012). Divergence and development of reproductive isolation are key initial phases in speciation events (Coyne and Orr 2004; Hernández-Hernández et al. 2021; Jarvis 2021). Sexual signals frequently emerge as the most divergent traits between closely related species and often experience strong selection (Ritchie 2007; Servedio and Boughman 2017). Evolution and divergence of sexual signals can outpace ecological divergence (Arnegard et al. 2010) and it has been commonly hypothesised that rapid changes in sexual signals could drive species diversification (Maan and Seehausen 2011; Servedio and Boughman 2017).

Vocalisations are in part sexual signals that have the potential to influence speciation. Acoustic signals are used by a wide range of organisms and are complex, behavioural traits that enable individuals of the same species identify one another (Laidre and Johnstone 2013). Acoustic signals have broad-scale functions in species recognition and fine-scale functions in mate choice, thus potentially influencing speciation (Catchpole and Slater 2003; Wilkins et al. 2013). Divergence of acoustic signals has played a role in the early stages of speciation in a wide range of organisms including cicadas (Marshall et al. 2008), crickets (Gray and Cade 2000; Zuk et al. 2006; Tinghitella et al. 2018), frogs (Boul et al. 2007), and birds (Grant and Grant 2006). However, in the context of reproductive character displacement, the strength of selection on signals depends on range overlap, with sexual signals expected to be under greater divergence when species are sympatric rather than allopatric (Maan and Seehausen

2011). Therefore, at macroevolutionary scales (i.e., clade or family-level), the strength of

selection on acoustic signals is expected to vary between species and among clades. It

remains unclear, however, whether the processes of divergence, assortative mating, and

selection of acoustic traits, which occur at the microevolutionary scale, can have

macroevolutionary consequences.

Birdsong is one of the best-known acoustic signals and is an important sexual signal.

Male birds sing to attract females: consequently, song is hypothesised to act as an important

premating isolation barrier in birds and influence speciation (Price 2008; Tobias et al. 2020).

The role of birdsong in speciation and reproductive isolation has been studied at fine scales,

and empirical evidence supports theoretical expectations. Divergence of song has contributed

to reproductive isolation in warblers (Irwin et al. 2001) and has influenced the early stages of

peripatric speciation in antbirds in Amazonia (Seddon and Tobias 2007). Additionally, long-

term studies on Darwin's finches have demonstrated that sexual selection on differences in

vocalisations, resulting from changes in beak shape morphology, have impacted mate choice

and contributed to reproductive isolation and speciation (Podos 2001; Podos and Nowicki

2004; Grant and Grant 2006). Although acoustic signals have are commonly hypothesised to

influence diversification (Wilkins et al. 2013), most studies focus on population or sister

species comparisons, and empirical studies testing the influence of signal divergence on

speciation at broader scales remain scarce (Mason et al. 2017). Where the relationship

between song and speciation has been investigated at macroevolutionary scales, bursts in

rates of speciation coincide with bursts in song evolution, however, no significant

correlations have been uncovered at the family-level (Mason et al. 2017), and the influence of

song on speciation remains to be explored in other clades and other regions of the world.

The relationship between speciation and birdsong has proven increasingly difficult to

disentangle, largely due to song itself being a complex trait, which in birds is influenced by

developmental mode and constrained by factors such as beak morphology, body size, and the

environment (Derryberry et al. 2018; Friedman et al. 2019; Friis et al. 2022). The majority of

macroevolutionary studies on birdsong have therefore focussed on the underlying drivers of

song evolution. Like most sexual signals, acoustic signals are expected to diverge between

close relatives, especially if ranges overlap, potentially leading to a lack of phylogenetic

signal in the trait (Pfennig and Pfennig 2010). However, song frequency is understood to be

phylogenetically conserved (Mikula et al. 2021), largely due to constraints imposed by

morphological traits, especially body size (Derryberry et al. 2018; Pearse et al. 2018).

Moreover, the developmental mode of song is thought to influence song evolution and could

complicate its relationship with speciation. Birdsong is either learned or innate, and learning

is understood to be a form of phenotypic plasticity which is influenced by cultural

transmission and can impact evolutionary outcomes (Pfennig et al. 2010). Learned songs are

typically associated with higher rates of evolution (Mason et al. 2017) and divergence

(Lachlan and Servedio 2004) than innate songs, although conflicting evidence exists for this

(Freeman et al. 2017). Such complexity in the evolution of acoustic traits provides a

challenge for speciation approaches and can result in disparity when investigations are

conducted over varying phylogenetic and spatial scales.

To overcome such limitations, we test the relationship between song and speciation

using a monophyletic clade restricted to a particular region, recognising the power that

focusing on a diverse single clade can bring (Marquet et al. 2004). Thus, we focus on

honeyeaters (Aves: Meliphagidae), a phenotypically diverse clade of birds (192 species; Gill

et al. 2020) that range throughout Australasia and are known for their distinctive

vocalisations (Higgins et al. 2008). Song in honeyeaters is learned and varies greatly among

species, with some species exhibiting complex and variable songs such as the Tūī (Hill et al.

2015), while others have simpler and more stereotyped songs, such as the noisy miner

(Jurisevic and Sanderson 2016). This variation in vocalisations, combined with the high

diversity and endemism of honeyeaters, makes this group an ideal model for studying the

evolution of vocalisations and their potential role in speciation. Specifically, we use

honeyeaters as a model clade to test the hypothesis that song influences speciation at

macroevolutionary scales and that rates of song evolution are correlated rates of speciation.

Using a dataset of five song variables for 163 species of honeyeater, we implement an

approach where we first model shifts in trait evolution and identify species associated with

shifts in song evolution, and test whether these are associated with increased rates of

speciation and diversification.

**Methods**

*Song and phylogenetic data*

We use a song dataset from a previous honeyeater study (Hay et al. 2023). Containing song

data for 163 honeyeater species, the song dataset considers five relevant aspects of song (Fig.

1; Table S1) which have commonly been explored in past song studies (Mason and Burns

2015; Mason et al. 2017; Derryberry et al. 2018): peak frequency (frequency which the most

sound energy was produced), maximum frequency (maximum frequency of the highest

pitched note in song), minimum frequency (minimum frequency of the lowest pitched note in

song), song bandwidth (difference between maximum and minimum frequency), and song

pace (number of notes/song duration). Song variables were $\log_{10}$-transformed prior to

statistical analysis to meet parametric assumptions of normality and homogeneity of variance,

and because logarithmic scales of sound frequency correspond to how birds perceive and

modulate sound (Cardoso 2013).

To model trait evolution and estimate speciation rate we used a honeyeater phylogeny

from Hay et al. (2022), which contains 192 species of honeyeater. The phylogeny was

trimmed to represent the 163 honeyeater species for which we had song data and using the

'*drop.tip*' function from the 'ape' package (Paradis et al. 2004) in R (R Core Team 2013).

Trait values are predicted to be more similar between closely related species, however sexual

signals are often expected to be the most divergent traits between species (Servedio and

Boughman 2017). To test for how conserved song traits are, we used the '*phylosig*' function

from the 'phytools' package (Revell 2012) in R (R Core Team 2013) and calculated the

phylogenetic signal of each song variable, using lambda and a likelihood ratio test. When

lambda is equal to 1 variation along the phylogenetic tree is equal to that of a Brownian

motion model, whereas values lower than one suggest less variation than expected under

Brownian motion (Revell et al. 2008).

*Analyses*

A range of phylogenetic comparative methods, implemented through the program RevBayes

(Höhna et al. 2016) were used to test the relationship between song and speciation. RevBayes

enables Bayesian inference for phylogenetic comparative methods, and uses a probabilistic

graphical model framework which is a powerful generic framework for specifying and

analysing statistical models (Höhna et al. 2014; Höhna et al. 2016). All RevBayes models

were implemented through the CIPRES Science Gateway v3.3 online server (Miller et al.

2010). First, to explore patterns of song evolution in honeyeaters, we used relaxed Ornstein-

Uhlenbeck (OU) models of trait evolution to identify shifts in trait evolution of the different

song traits in RevBayes. OU models were also implemented in a multivariate framework

using RevBayes and verified using multivariate OU models in the package 'PhylogeneticEM'

(Bastide et al. 2018) in R (R Core Team 2013). The influence of these shifts on speciation

was then tested with hidden state-dependent speciation and extinction models (Beaulieu and

O'Meara 2016) in RevBayes (Höhna et al. 2016).

Relaxed Ornstein-Uhlenbeck models of trait evolution were used to model song

evolution along the honeyeater phylogeny. Relaxed OU models allow the optimal phenotype

to vary over branches of the phylogenetic tree to estimate tip-specific measures of OU

evolution (Uyeda and Harmon 2014) and enabled us to infer branch-specific evolutionary

optima of song evolution in honeyeaters. Here, we first assessed patterns of OU trait

evolution in each of the five song traits (peak frequency, maximum frequency, minimum

frequency, bandwidth, and pace) separately. Relaxed OU models were implemented in

RevBayes using reversible-jump Markov chain Monte Carlo. Examples from the online

tutorial (May and Höhna 2022) were followed, and the MCMC sampler was set for 4 runs of

50,000 generations with a burn-in of 10%. Tracer v1.7.1 (Rambaut et al. 2018) was used to

visualise log files and to ensure convergence by verifying ESS values were above 200 for all

parameters. Because priors can influence performance of OU models, particularly the number

of expected shifts, all models were repeated over a number of shifts (15, 30, and 45). Model

fit was assessed using stepping-stone and path sampler analyses, and the best model was

selected based on marginal likelihood values. In all cases the preferred models had 45 shifts

(Table S2). Branch-specific rates of OU evolution were visualised by plotting output using

the package 'RevGadgets' (Tribble et al. 2021) in R (R Core Team 2013). Branch theta

values were extracted for each species and shifts in OU trait evolution were determined based

on plots and theta values (Table S3).

Trait evolution is inherently a multivariate process (Walsh and Blows 2009; Revell

and Harmon 2022). Thus, we also used multivariate relaxed OU models in RevBayes. The

function '*dnPhyloOrnsteinUhlenbeckREML*' for the relaxed OU model in RevBayes does not

enable the inclusion of a correlation matrix to account for covariation between multivariate

traits. To overcome this limitation, we performed a phylogenetic principal component

analysis on the five song variables to ensure no correlation between traits, and considered

PC1 to PC3, which accounted for 97.4 % of the total variation, in analysis. Multivariate

model parameters, model fit, and visualisation followed the same approach and settings as

univariate models above.

As an additional measure to test for shifts in trait evolution, and to verify the results

from the multivariate relaxed OU models in RevBayes, we also used the package

'PhylogeneticEM' (Bastide et al. 2018) in R (R Core Team 2013). PhylogeneticEM simulates

a scalar OU model of evolution that allows for maximum likelihood estimation of shifts in

OU trait evolution and uses this approach to automatically detect position of shifts along a

phylogeny, while accounting for correlations among traits (Bastide et al. 2018). The function

'phyloEM' was used to test for OU shifts in multivariate trait evolution, using PC1-PC3, the

same dataset which was used for the multivariate OU model in RevBayes.

Finally, to test the influence of shifts in song evolution on speciation rates in

honeyeaters we used hidden state-dependent speciation and extinction models (HiSSE;

Beaulieu and O'Meara 2016) which were also implemented in RevBayes (Höhna et al. 2016).

The HiSSE framework enables testing of character dependent diversification and additionally

models an unobserved hidden character that potentially impacts species diversification rates.

HiSSE relies on testing the influence of binary characters on speciation but can disentangle

rate variation due to the character of interest with a hidden state. HiSSE represents a

substantial improvement on previous state-dependent diversification models (Maddison and

FitzJohn 2015; Rabosky and Goldberg 2015). Species associated with shifts in song evolution

were coded with a '1' and all other species were coded as '0'. This was done to test whether

shifts in song evolution are also associated with increases in diversification rate. Speciation

models can be subject to bias when species are missing (Mynard et al. 2023), therefore we

implemented all HiSSE models across the full honeyeater phylogeny containing all 192

species of honeyeater (Hay et al. 2022). The 29 honeyeater species with no song data were

treated as having unknown tip states and were coded with '?' rather than '0' or '1'. The

MCMC sampler was set for 4 runs of 3,000 generations with a burn-in of 10%. Tracer v1.7.1

(Rambaut et al. 2018) was used to visualise output and determine convergence by verifying

ESS values were above 200 for all parameters.

**Results**

Phylogenetic signal varied substantially among the five song variables. Values of lambda

varied from 0.27 to 0.91 and the significance values of the likelihood ratio tests were all

below $p < 0.001$ (Table S4). Song bandwidth had the lowest signal ($\lambda = 0.27$), followed by

maximum frequency ($\lambda = 0.61$), pace ($\lambda = 0.68$), peak frequency ($\lambda = 0.75$) and then

minimum frequency ($\lambda = 0.91$).

Investigation of the evolution of song variables with univariate relaxed OU models

revealed several patterns. Overall, the three frequency variables (peak frequency, maximum

frequency, and minimum frequency) had very similar estimates for relaxed models of OU

trait evolution. Here, the clade *Myzomela + Sugomel* is associated with a shift in OU trait

evolution for all three frequency variables, and *Philemon* is identified with shifts in peak and

maximum frequency (Fig. 2). Despite this, we uncovered little evidence that shifts in trait

evolution influence speciation rates for peak frequency (mean $\lambda_{no\ shift} = 2.44$, 95 % CI = [1.80,

3.18]; mean $\lambda_{shift} = 3.46$, 95 % CI = [2.31, 4.74]; Fig. 2; Table S5), maximum frequency

(mean $\lambda_{no\ shift} = 2.49$, 95 % CI = [1.83, 3.23]; mean $\lambda_{shift} = 3.17$, 95 % CI = [2.18, 4.28]; Fig.

2; Table S5), and minimum frequency (mean $\lambda_{no\ shift}$ = 2.55, 95 % CI = [1.88, 3.24]; mean

$\lambda_{shift}$ = 3.35, 95 % CI = [2.10, 4.77]; Fig. 2; to S5). Shifts in bandwidth (mean $\lambda_{no\ shift}$ = 2.78,

95 % CI = [2.11, 3.48]; mean $\lambda_{shift}$ = 1.56, 95 % CI = [0.50, 2.93]) and pace (mean $\lambda_{no\ shift}$ =

2.81, 95 % CI = [2.08, 3.57]; mean $\lambda_{shift}$ = 1.45, 95 % CI = [1.01E-6, 5.33]) appear to be

associated with lower rates of speciation (Fig. 2; Table S5), but this is also non-significant,

and could be due to the lower sample sizes for shifts in bandwidth (14 species associated with

shifts) and pace (7 species identified with shifts).

The two methods of multivariate OU trait evolution (RevBayes and PhylogeneticEM)

uncovered similar shifts in optima of song in honeyeaters. Species from the genera

*Myzomela*, *Sugomel*, *Philemon*, *Nesoptilotis* and *Gymnomyza* are all associated with shifts in

OU trait evolution in both approaches (Fig. 3). The PhylogeneticEM method does reveal a

shift associated with the larger clade containing *Melidectes* and relatives, which was not

recovered in the RevBayes relaxed multivariate OU model (Fig. 3).

We found no evidence that shifts in song evolution are associated with increased rates

of speciation or net diversification for both the RevBayes (mean $\lambda_{no\ shift}$ = 2.54, 95 % CI =

[1.81, 3.31]; mean $\lambda_{shift}$ = 3.03, 95 % CI = [2.09, 4.00]; mean diversification$_{no\ shift}$ = 2.52, 95

% CI = [1.78, 3.29]; mean diversification$_{shift}$ = 2.98, 95 % CI = [2.05, 3.98]) and

PhylogeneticEM approaches (mean $\lambda_{no\ shift}$ = 2.24, 95 % CI = [1.53, 3.01]; mean $\lambda_{shift}$ = 3.29,

95 % CI = [2.28, 4.44] ; mean diversification$_{no\ shift}$ = 2.21, 95 % CI = [1.50, 2.99]; mean

diversification$_{shift}$ = 3.25, 95 % CI = [2.28, 4.47]), and the 95 % confidence intervals overlap

substantially (Fig. 3; Table S5). Both methods uncovered similar estimates for speciation and

net diversification (Fig. 3; Table S5).

**Discussion**

In this study, we used honeyeaters to explore the macroevolutionary consequences of song

evolution, a sexually selected behavioural trait, which has commonly been hypothesised to

influence speciation in birds. We find no evidence that shifts in song evolution are associated

with increases in speciation or diversification rates. The significant overlap between the

confidence intervals of states is present in all tests, and hidden state speciation and extinction

models suggest that there are other factors (hidden states) which play a role. Ultimately, song

does not appear to have contributed to speciation on a macroevolutionary scale in

honeyeaters, possibly due to other factors including geographical processes and dispersal to

islands dominating speciation history in honeyeaters (Hay et al. 2022), and any apparent

influence of song is likely mediated by these factors and underlying variation in body size.

Most of the interspecific diversity of song frequency and pace in honeyeaters can be

explained by evolutionary history. Theory suggests that divergence is expected for sexually

selected traits between closely related species (Maan and Seehausen 2011; Servedio and

Boughman 2017). Thus, the level of phylogenetic signal uncovered for frequency and pace is

contrary to such theoretical expectations. This is unsurprising, however, given other studies

have found similar levels of phylogenetic signal for song traits in birds. For example, peak

frequency was found to have a lambda of 0.87 for 5085 species of passerine birds (Mikula et

al. 2021), and a lambda of 0.74 for 1022 passerine species (Friis et al. 2022). Our results are

consistent with these previous studies and confirm that peak frequency of song is

phylogenetically conserved. This is likely driven by allometric constraints imposed by body

size (Fletcher 2004), which itself is a very conserved trait, and directly impacts the length of

the vocal tract and the size of the syrinx (Bertelli and Tubaro 2002; Suthers and Zollinger

2008). Certainly, body size has been demonstrated to have a prominent influence on

honeyeater song and has been found to structure variation in frequency and song pace (Hay et

al. 2023). Interestingly, minimum frequency was found to have the highest phylogenetic

signal rather than peak frequency. It is unclear if this is the case in other clades as past studies

have typically focussed on peak frequency. Nonetheless, body size is understood to place a

limit on the lowest frequency that a bird can effectively produce (Fletcher 2004; Friis et al.

2021), so perhaps minimum frequency is just particularly constrained by body size.

We found no evidence that shifts in song evolution are associated with higher rates of

speciation and net diversification. Macroevolutionary influences of song evolution on

speciation have been explored in 581 species of tanagers and ovenbirds and shifts in song

were found to be correlated with shifts in diversification, but this was not significant when

each family was considered separately (Mason et al. 2017). This is consistent with the

findings of this study, which is restricted to the family-level, and could imply that

macroevolutionary signals of song on speciation and diversification emerge only when very

different clades of birds are examined simultaneously, as is the case for other taxa too

(Rabosky et al. 2013; Rabosky and Huang 2016). Additionally, the song dataset used in this

study was missing data for 29 honeyeater species. As is often the case with missing data,

species inhabiting the most remote areas were data deficient. This skewed our dataset towards

common species and especially those found in Australia and New Zealand. Areas such as

New Guinea, which has high honeyeater species richness (Marki et al. 2019), and islands

throughout the Indo-Pacific, which contain endemic species, were lacking in data. Given

previous work highlighting island settings for speciation in the group (Hay et al. 2022), this

potential effect of our data set should be considered a caveat for the study.

Nonetheless, our results indicate that song is not a major driver of speciation at

macroevolutionary scales in honeyeaters. Recent studies have found that faster rates of song

evolution do not result in higher speciation rates in New World passerine birds (Freeman et

al. 2022), and ecological and demographic factors may provide a better explanation for

speciation (Singhal et al. 2018; Harvey et al. 2019; Freeman et al. 2022). State-dependent

speciation and extinction models used in the present study indicate hidden states are

associated with differential rates of speciation and net diversification, implying that other

factors not considered here could contribute to patterns of speciation. Honeyeaters have a

dynamic evolutionary history: Australia is the centre of honeyeater diversity (Driskell and

Christidis 2004; Norman et al. 2007; Oliveros et al. 2019), however, dispersal to islands

throughout the Indo-Pacific, and vicariant speciation due to connection and disconnection of

New Guinea and the changing biomes of Australia have played key roles in the

diversification of this group (Toon et al. 2010; Andersen et al. 2014; Marki et al. 2017; Hay et

al. 2022). Moreover, past studies have found range size and dispersal ability to significantly

contribute to speciation in honeyeaters when applying state-dependent speciation and

extinction models (Hay et al. 2022). Allopatric speciation is the predominant mode of

speciation in honeyeaters and could explain the lack of influence of song evolution on

speciation because selection on signal divergence is expected to be stronger for sympatry

(Pfennig and Pfennig 2010). Songs are thought to diverge slowly in allopatry, and only

rapidly increase on secondary contact, contributing to reinforcement (Lachlan and Servedio

2004; Servedio and Boughman 2017). Thus, contributions of song to speciation likely

depends on the mode of speciation, so although song could be aiding microevolutionary

processes such as reproductive isolation, other factors including range size and dispersal

ability have contributed more to speciation more at the macroevolutionary scale (Hay et al.

2022).

Across all models of trait evolution, shifts in song evolution are notable in clades of

*Myzomela + Sugomel* and *Philemon*. These clades have previously been identified to have

increased rates of speciation (Hay et al. 2022) and are also the most speciose genera in

honeyeaters (Gill et al. 2020). However, past studies have shown that small range sizes

resulting from dispersal to islands has been a primary driver of speciation in these clades

(Hay et al. 2022), and the (non-significant) slight increase in speciation rate observed with

these shifts in song evolution could be due to these underlying factors. It is also important to

consider the influence of body size on these patterns because honeyeater song is under strong

allometric constraints (Hay et al. 2023), and these clades show great differences in body size.

For example, all species of *Myzomela* are between 6 to 16 g, whereas species of *Philemon*

range from 50 g to 194 g, in which the majority of species are over 100 g (Dunning 2008;

Tobias et al. 2022). Additionally, body size evolution has been correlated with speciation and

species diversity across different vertebrate clades (Ricklefs 2004; Rabosky and Adams 2012;

Rabosky et al. 2013) and has contributed to speciation in honeyeaters (Brennan and Keogh

2018; Marki et al. 2019). Therefore, it is possible that the shifts in song evolution could be

due to shifts in body size evolution, and underlying variation in body size could be

contributing to both the observed patterns of song evolution and speciation in this study.

Understanding the influence of reproductive isolation on diversification has been

recognised as an urgent question in evolutionary biology (Matute and Cooper 2021),

especially because the evolution of reproductive isolation often defines speciation (Coyne

and Orr 2004). Here, we contribute to understanding of this question by testing the

hypothesis that song has influenced speciation on macroevolutionary scales in honeyeaters.

We uncovered several shifts in song evolution but found no evidence that these shifts were

associated with significant increases in diversification or speciation rates. Models also

indicated that other factors (hidden states) contribute to differential rates of speciation and

diversification, implying that speciation history in honeyeaters is better explained by other

factors.

**Literature Cited**

Andersen MJ, Naikatini A, Moyle RG. 2014. A molecular phylogeny of Pacific honeyeaters

(Aves: Meliphagidae) reveals extensive paraphyly and an isolated Polynesian radiation.

Molecular Phylogenetics and Evolution, 71:308-315.

Arnegard ME, McIntyre PB, Harmon LJ, Zelditch ML, Crampton WG, Davis JK, Sullivan

JP, Lavoue S, Hopkins CD. 2010. Sexual signal evolution outpaces ecological

divergence during electric fish species radiation. The American Naturalist, 176:335-

356.

Bastide P, Ané C, Robin S, Mariadassou M. 2018. Inference of adaptive shifts for

multivariate correlated traits. Systematic Biology, 67:662-680.

Beaulieu JM, O'Meara BC. 2016. Detecting hidden diversification shifts in models of trait-

dependent speciation and extinction. Systematic Biology, 65:583-601.

Bertelli S, Tubaro PL. 2002. Body mass and habitat correlates of song structure in a primitive

group of birds. Biological Journal of the Linnean Society, 77:423-430.

Boul KE, Funk WC, Darst CR, Cannatella DC, Ryan MJ. 2007. Sexual selection drives

speciation in an Amazonian frog. Proceedings of the Royal Society B: Biological

Sciences, 274:399-406.

Brennan IG, Keogh JS. 2018. Miocene biome turnover drove conservative body size

evolution across Australian vertebrates. Proceedings of the Royal Society B: Biological

Sciences, 285:20181474.

Cardoso GC. 2013. Using frequency ratios to study vocal communication. Animal Behaviour,

85:1529-1532.

Catchpole CK, Slater PJB. 2003. Bird Song: Biological Themes and Variations. Cambridge

University Press: New York.

Cooney CR, Thomas GH. 2021. Heterogeneous relationships between rates of speciation and

body size evolution across vertebrate clades. Nature Ecology and Evolution, 5:101-110.

Coyne JA, Orr HA. 2004. Speciation. Sinauer Associates: Sunderland, Massachusetts USA.

Derryberry EP, Seddon N, Derryberry GE, Claramunt S, Seeholzer GF, Brumfield RT,

Tobias JA. 2018. Ecological drivers of song evolution in birds: disentangling the effects

of habitat and morphology. Ecology and Evolution, 8:1890-1905.

Driskell AC, Christidis L. 2004. Phylogeny and evolution of the Australo-Papuan

honeyeaters (Passeriformes, Meliphagidae). Molecular Phylogenetics and Evolution,

31:943-960.

Dumont ER, Davalos LM, Goldberg A, Santana SE, Rex K, Voigt CC. 2012. Morphological

innovation, diversification and invasion of a new adaptive zone. Proceedings of the

Royal Society B: Biological Sciences, 279:1797-1805.

Dunning JB. 2008. CRC Handbook of Avian Body Masses. Taylor & Francis Group, LLC:

Fletcher NH. 2004. A simple frequency-scaling rule for animal communication. The Journal

of the Acoustical Society of America, 115: 2334-2338.

Freeman BG, Montgomery GA, Schluter D. 2017. Evolution and plasticity: divergence of

song discrimination is faster in birds with innate song than in song learners in

Neotropical passerine birds. Evolution, 71:2230-2242.

Freeman BG, Rolland J, Montgomery GA, Schluter D. 2022. Faster evolution of a premating

reproductive barrier is not associated with faster speciation rates in New World

passerine birds. Proceedings of the Royal Society B: Biological Sciences,

289:20211514.

Friedman NR, Miller ET, Ball JR, Kasuga H, Remes V, Economo EP. 2019. Evolution of a

multifunctional trait: shared effects of foraging ecology and thermoregulation on beak

morphology, with consequences for song evolution. Proceedings of the Royal Society

B: Biological Sciences, 286:20192474.

Friis JI, Sabino J, Santos P, Dabelsteen T, Cardoso GC. 2021. The allometry of sound

frequency bandwidth in songbirds. The American Naturalist, 197:607-614.

Friis JI, Sabino J, Santos P, Dabelsteen T, Cardoso GC, Jennions MD. 2022. Ecological

adaptation and birdsong: how body and bill sizes affect passerine sound frequencies.

Behavioral Ecology, 33:798-806.

Gill F, Donsker D, Rasmussen P. 2020. IOC World Bird List.

Grant PR, Grant BR. 2006. Evolution of character displacement in Darwin's finches. Science,

313:224-226.

Gray DA, Cade WH. 2000. Sexual selection and speciation in field crickets. Proceedings of

the National Academy of Sciences of the USA, 97: 14449-14454.

Harvey MG, Singhal S, Rabosky DL. 2019. Beyond reproductive isolation: demographic

controls on the speciation process. Annual Review of Ecology, Evolution, and

Systematics, 50:75-95.

Hay EM, McGee MD, Chown SL. 2022. Geographic range size and speciation in

honeyeaters. BMC Ecology and Evolution, 22:1-14.

Hay EM, McGee MD, White CR, Chown SL. 2023. Body size shapes song in honeyeaters.

BioRxiv.

Hernández-Hernández T, Miller EC, Roman-Palacios C, Wiens JJ. 2021. Speciation across

the Tree of Life. Biological Reviews, 96:1205-1242.

Higgins PJ, Ford HA, Christidis L. 2008. Family Meliphagidae. In: Handbook of the Birds of

the World: Penduline-tits to shrikes. Lynx Edicions: Barcelona, Spain.

Hill SD, Amiot C, Ludbrook MR, Ji W. 2015. Seasonal variation in the song structure of tui

(Prosthemadera novaeseelandiae). New Zealand Ecological Society, 39:110-115.

Höhna S, Heath TA, Boussau B, Landis MJ, Ronquist F, Huelsenbeck JP. 2014. Probabilistic

graphical model representation in phylogenetics. Systematic Biology, 63:753-771.

Höhna S, Landis MJ, Heath TA, Boussau B, Lartillot N, Moore BR, Huelsenbeck JP,

Ronquist F. 2016. RevBayes: Bayesian phylogenetic inference using graphical models

and an interactive model-specification language. Systematic Biology, 65:726-736.

Irwin DE, Bensch S, Price TD. 2001. Speciation in a ring. Nature, 409:333-337.

Jarvis ED. 2021. At the beginning of speciation. Science, 371:1312.

Jurisevic MA, Sanderson KJ. 2016. The vocal repertoires of six honeyeater (Meliphagidae)

species from Adelaide, South Australia. Emu - Austral Ornithology, 94:141-148.

Lachlan RF, Servedio MR. 2004. Song learning accelerates allopatric speciation. Evolution,

58:2049-2063.

Laidre ME, Johnstone RA. 2013. Animal signals. Current Biology, 23:R829-833.

Maan ME, Seehausen O. 2011. Ecology, sexual selection and speciation. Ecology Letters,

14:591-602.

Maddison WP, FitzJohn RG. 2015. The unsolved challenge to phylogenetic correlation tests

for categorical characters. Systematic Biology, 64:127-136.

Mahler DL, Revell LJ, Glor RE, Losos JB. 2010. Ecological opportunity and the rate of

morphological evolution in the diversification of Greater Antillean anoles. Evolution,

64:2731-2745.

Marki PZ, Jonsson KA, Irestedt M, Nguyen JMT, Rahbek C, Fjeldsa J. 2017. Supermatrix

phylogeny and biogeography of the Australasian Meliphagides radiation (Aves:

Passeriformes). Molecular Phylogenetics and Evolution, 107:516-529.

Marki PZ, Kennedy JD, Cooney CR, Rahbek C, Fjeldsa J. 2019. Adaptive radiation and the

evolution of nectarivory in a large songbird clade. Evolution, 73:1226-1240.

Marquet PA, Fernández M, Navarette AA, Valdovinos C. 2004. Diversity emerging: towards

a deconstruction of biodiversity patterns. In: Frontiers of biogeography: New directions

in the geography of nature. Sinauer Associates: Sunderland.

Marshall DC, Slon K, Cooley JR, Hill KBR, Simon C. 2008. Steady Plio-Pleistocene

diversification and a 2-million-year sympatry threshold in a New Zealand cicada

radiation. Molecular Phylogenetics and Evolution, 48:1054-1066.

Mason NA, Burns KJ. 2015. The effect of habitat and body size on the evolution of vocal

displays in Thraupidae (tanagers), the largest family of songbirds. Biological Journal of

the Linnean Society, 114:538-551.

Mason NA, Burns KJ, Tobias JA, Claramunt S, Seddon N, Derryberry EP. 2017. Song

evolution, speciation, and vocal learning in passerine birds. Evolution, 71:786-796.

Matute DR, Cooper BS. 2021. Comparative studies on speciation: 30 years since Coyne and

Orr. Evolution, 75:764-778.

May MR, Höhna S. 2022. RevBayes: relaxed Ornstein-Uhlenbeck models. Available from

https://revbayes.github.io/tutorials/cont_traits/relaxed_ou.html.

Mikula P, Valcu M, Brumm H, Bulla M, Forstmeier W, Petruskova T, Kempenaers B,

Albrecht T. 2021. A global analysis of song frequency in passerines provides no

support for the acoustic adaptation hypothesis but suggests a role for sexual selection.

Ecology Letters, 24:477-486.

Miller MA, Pfeiffer W, Schwartz T. 2010. Creating the CIPRES Science Gateway for

inference of large phylogenetic trees. 2010 Gateway Computing Environments

workshop (GCE). New Orleans, LA, Ieee.

Mynard P, Algar A, Lancaster L, Bocedi G, Fahri F, Gubry-Rangin C, Lupiyaningdyah P,

Nangoy M, Osborne O, Papadopulos A, Sudiana IM, Juliandi B, Travis J, Herrera-

Alsina L. 2023. Impact of phylogenetic tree completeness and misspecification of

489  sampling fractions on trait dependent diversification models. Systematic Biology,

490  72:106-119

491 Norman JA, Rheindt FE, Rowe DL, Christidis L. 2007. Speciation dynamics in the Australo-

492  Papuan Meliphaga honeyeaters. Molecular Phylogenetics and Evolution, 42:80-91.

493 Oliveros CH, Field DJ, Ksepka DT, Barker FK, Aleixo A, Andersen MJ, Alstrom P, Benz

494  BW, Braun EL, Braun MJ, Bravo GA, Brumfield RT, Chesser RT, Claramunt S,

495  Cracraft J, Cuervo AM, Derryberry EP, Glenn TC, Harvey MG, Hosner PA, Joseph L,

496  Kimball RT, Mack AL, Miskelly CM, Peterson AT, Robbins MB, Sheldon FH, Silveira

497  LF, Smith BT, White ND, Moyle RG, Faircloth BC. 2019. Earth history and the

498  passerine superradiation. Proceedings of the National Academy of Sciences of the

499  USA, 116:7916-7925.

500 Paradis E, Claude J, Strimmer K. 2004. APE: analyses of phylogenetics and evolution in R

501  language. Bioinformatics, 20:289-290.

502 Pearse WD, Morales-Castilla I, James LS, Farrell M, Boivin F, Davies TJ. 2018. Global

503  macroevolution and macroecology of passerine song. Evolution, 72:944-960.

504 Pfennig DW, Pfennig KS. 2010. Character displacement and the origins of diversity. The

505  American Naturalist, 176:26-44.

506 Pfennig DW, Wund MA, Snell-Rood EC, Cruickshank T, Schlichting CD, Moczek AP. 2010.

507  Phenotypic plasticity's impacts on diversification and speciation. Trends in Ecology &

508  Evolution, 25:459-467.

509 Podos J. 2001. Correlated evolution of morphology and vocal signal structure in Darwin's

510  fnches. Nature, 409:185-188.

511 Podos J, Nowicki S. 2004. Beaks, adaptation, and vocal evolution in Darwin's finches.

512  Bioscience, 54:501-510.

513 Price T. 2008. Speciation in birds. Roberts and Co.: Colorado.

R Core Team. 2013. R: a language and environment for statistical computing.

Rabosky DL, Adams DC. 2012. Rates of morphological evolution are correlated with species

richness in salamanders. Evolution, 66:1807-1818.

Rabosky DL, Goldberg EE. 2015. Model inadequacy and mistaken inferences of trait-

dependent speciation. Systematic Biology, 64:340-355.

Rabosky DL, Huang H. 2016. A robust semi-parametric test for detecting trait-dependent

diversification. Systematic Biology, 65:181-193.

Rabosky DL, Santini F, Eastman J, Smith SA, Sidlauskas B, Chang J, Alfaro ME. 2013.

Rates of speciation and morphological evolution are correlated across the largest

vertebrate radiation. Nature Communications, 4:1958.

Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. 2018. Posterior summarization in

Bayesian phylogenetics using Tracer 1.7. Systematic Biology, 67:901-904.

Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other

things). Methods in Ecology and Evolution, 3:217-223.

Revell LJ, Harmon LJ. 2022. Phylogenetic comparative methods in R. Princeton University

Press: Princeton, NJ.

Revell LJ, Harmon LJ, Collar DC. 2008. Phylogenetic signal, evolutionary process, and rate.

Systematic Biology, 57:591-601.

Ricklefs RE. 2004. Cladogenesis and morphological diversification in passerine birds.

Nature, 430:338-341.

Ritchie MG. 2007. Sexual selection and speciation. Annual Review of Ecology, Evolution,

and Systematics, 38:79-102.

Seddon N, Tobias JA. 2007. Song divergence at the edge of Amazonia: an empirical test of

the peripatric speciation model. Biological Journal of the Linnean Society, 90:173-188.

Servedio MR, Boughman JW. 2017. The role of sexual selection in local adaptation and

speciation. Annual Review of Ecology, Evolution, and Systematics, 48:85-109.

Simões TR, Vernygora O, Caldwell MW, Pierce SE. 2020. Megaevolutionary dynamics and

the timing of evolutionary innovation in reptiles. Nature Communications, 11:3322.

Singhal S, Huang H, Grundler MR, Marchan-Rivadeneira MR, Holmes I, Title PO,

Donnellan SC, Rabosky DL. 2018. Does population structure predict the rate of

speciation? a comparative test across australia's most diverse vertebrate radiation. The

American Naturalist, 192:432-447.

Suthers RA, Zollinger SA. 2008. From brain to song: the vocal organ and vocal tract. In:

Neuroscience of birdsong. Cambridge University Press: Cambridge.

Tinghitella RM, Broder ED, Gurule-Small GA, Hallagan CJ, Wilson JD. 2018. Purring

crickets: the evolution of a novel sexual signal. The American Naturalist, 192:773-782.

Tobias JA, Ottenburghs J, Pigot AL. 2020. Avian diversity: speciation, macroevolution, and

ecological function. Annual Review of Ecology, Evolution, and Systematics, 51:533-

560.

Tobias JA, Sheard C, Pigot AL, Devenish AJM, Yang J, Sayol F, Neate-Clegg MHC,

Alioravainen N, Weeks TL, Barber RA, Walkden PA, MacGregor HEA, Jones SEI,

Vincent C, Phillips AG, Marples NM, Montano-Centellas FA, Leandro-Silva V,

Claramunt S, Darski B, Freeman BG, Bregman TP, Cooney CR, Hughes EC, Capp

EJR, Varley ZK, Friedman NR, Korntheuer H, Corrales-Vargas A, Trisos CH, Weeks

BC, Hanz DM, Topfer T, Bravo GA, Remes V, Nowak L, Carneiro LS, Moncada RA,

Matysiokova B, Baldassarre DT, Martinez-Salinas A, Wolfe JD, Chapman PM, Daly

BG, Sorensen MC, Neu A, Ford MA, Mayhew RJ, Fabio Silveira L, Kelly DJ,

Annorbah NND, Pollock HS, Grabowska-Zhang AM, McEntee JP, Carlos TGJ,

Meneses CG, Munoz MC, Powell LL, Jamie GA, Matthews TJ, Johnson O, Brito GRR,

Zyskowski K, Crates R, Harvey MG, Jurado Zevallos M, Hosner PA, Bradfer-

Lawrence T, Maley JM, Stiles FG, Lima HS, Provost KL, Chibesa M, Mashao M,

Howard JT, Mlamba E, Chua MAH, Li B, Gomez MI, Garcia NC, Packert M, Fuchs J,

Ali JR, Derryberry EP, Carlson ML, Urriza RC, Brzeski KE, Prawiradilaga DM,

Rayner MJ, Miller ET, Bowie RCK, Lafontaine RM, Scofield RP, Lou Y, Somarathna

L, Lepage D, Illif M, Neuschulz EL, Templin M, Dehling DM, Cooper JC, Pauwels

OSG, Analuddin K, Fjeldsa J, Seddon N, Sweet PR, DeClerck FAJ, Naka LN, Brawn

JD, Aleixo A, Bohning-Gaese K, Rahbek C, Fritz SA, Thomas GH, Schleuning M.

2022. AVONET: morphological, ecological and geographical data for all birds.

Ecology Letters, 25:581-597.

Toon A, Hughes JM, Joseph L. 2010. Multilocus analysis of honeyeaters (Aves:

Meliphagidae) highlights spatio-temporal heterogeneity in the influence of

biogeographic barriers in the Australian monsoonal zone. Molecular Ecology, 19:2980-

2994.

Tribble CM, Freyman WA, Landis MJ, Lim JY, Barido-Sottani J, Kopperud BT, Höhna S,

May MR. 2021. RevGadgets: an R package for visualizing Bayesian phylogenetic

analyses from RevBayes. Methods in Ecology and Evolution, 13:314-323.

Wagner CE, Harmon LJ, Seehausen O. 2012. Ecological opportunity and sexual selection

together predict adaptive radiation. Nature, 487:366-369.

Walsh B, Blows MW. 2009. Abundant genetic variation + strong selection = multivariate

genetic constraints: a geometric view of adaptation. Annual Review of Ecology,

Evolution, and Systematics, 40:41-59.

Wilkins MR, Seddon N, Safran RJ. 2013. Evolutionary divergence in acoustic signals: causes

and consequences. Trends in Ecology & Evolution, 28:156-166.

Zuk M, Rotenberry JT, Tinghitella RM. 2006. Silent night: adaptive disappearance of a

sexual signal in a parasitized population of field crickets. Biology Letters, 2:521-524.

**Figures**

**Figure 1.** Song spectrogram for the New Zealand Bellbird (*Anthornis melanura*; xeno-cento

recording accession number: XC153005), displaying frequency (kHz) over time (s). The five

song variables considered in this study are indicated with annotations; peak frequency

represents the frequency of maximum amplitude across the entire song, maximum frequency

is the highest pitched note in song, minimum frequency is the lowest pitched note in song,

bandwidth is the difference between maximum and minimum frequency across the song, and

pace is the number of notes over song duration. NZ Bellbird illustration by Eleanor Hay.

**Figure 2.** Results from relaxed OU and HiSSE models for each of the five song traits; peak

frequency (a), maximum frequency (b), minimum frequency (c), bandwidth (d) and pace (e),

Phylogenetic trees display branch specific theta values from relaxed OU models in

RevBayes. Dots at the tips of the trees indicate which species were identified as associated

with shifts in song evolution and were tested to influence speciation rates. Respective

speciation rates extracted from HiSSE models associated with the coding at the tips are

displayed in the density plots to the right of each phylogeny.

**Figure 3.** Honeyeater phylogenetic tree (a), the branches of the phylogeny display estimated

branch theta values from the multivariate relaxed OU model in RevBayes. Dots on the

branches of the phylogeny indicate shifts revealed by PhylogeneticEM. Dots at the tip of the

tree show the binary classifications for HiSSE models from the RevBayes and

PhylogeneticEM (labelled as PhyloEM) approaches. Density plots below the phylogeny show

the estimated speciation and net diversification rates from HiSSE models for shifts identified

by RevBayes (b) and PhylogeneticEM (c). Birds are illustrations by Eleanor Hay.