OrthoPhyl – A turn-key solution for large scale whole genome bacterial phylogenomics

Earl A Middlebrook[1], Robab Katani[2], Jeanne M Fair[1]
[1] Genomics and Bioanalytics Group, Los Alamos National Lab
[2] Huck Institutes of Life Sciences, Pennsylvania State University

Abstract

Motivation – There are a staggering number of publicly available bacterial genome sequences (at writing, 1.5 million assemblies in NCBI's GenBank alone), and the deposition rate continues to increase. This wealth of data begs for phylogenetic analyses to place these sequences within an evolutionary context. A phylogenetic placement not only aids in taxonomic classification, but informs the evolution of novel phenotypes, targets of selection, and horizontal gene transfer. Building trees from multi-gene codon alignments is a laborious task that requires bioinformatic expertise, rigorous curation of orthologs, and heavy computation. Compounding the problem is the lack of tools that can streamline these processes for building trees from large scale genomic data.

Results - Here we present OrthoPhyl, which takes as input bacterial genome assemblies and reconstructs trees from whole genome codon alignments. The workflow can analyze an arbitrarily large number of input genomes (>1200 tested here) by identifying a diversity spanning subset of assemblies and using these genomes to build gene models to identify orthologs in the full dataset. OrthoPhyl is an easy-to-run, turn-key solution for generating high-resolution bacterial trees from datasets with a wide range of divergences. To illustrate the versatility of OrthoPhyl, we show three use-cases: E. coli/Shigella, Brucella/Ochrobactrum, and the order Rickettsiales. We compare trees generated with our

software to published trees from alternative methods. We show that OrthoPhyl trees are

consistent with other methods while incorporating more data, allowing for greater numbers

of input genomes, and more flexibility of analysis.

Availability and Implementation - Code used in this manuscript is available at

https://github.com/eamiddlebrook/OrthoPhyl.git. To aid in usability, a Singularity container

is available at *https://cloud.sylabs.io/library/earlyevol/default/orthophyl*. Installation and

execution instructions are provided in the associated github readme file. Assemblies used

with this manuscript are available from https://www.ncbi.nlm.nih.gov/assembly/ with

accessions in **Supplementary table 1-3**.

Contact – Earl A. Middlebrook – earlm@lanl.gov

<u>1 Introduction</u>

Underpinning every aspect of an organism's biology is its evolutionary history. Phylogenetic

methods strive to reconstruct these histories using genetic data. With the advent of "Next-"

and "Third-gen" sequencing methodologies, the amount of genetic data in public databases

has increased dramatically. At writing there are currently 1,502,852 bacterial genome

assemblies available through NCBI alone (ncbi.nlm.nih.gov/assembly). The wealth of

available genetic data has transformed the fields of bacterial phylogenetics and taxonomy

(Konstantinidis and Tiedje, 2007; Jain *et al.*, 2018; Varghese *et al.*, 2015). Unfortunately,

most whole genome phylogenetic analysis methods require specialized bioinformatic data

processing and expertise (Smith, 2013). To compound the issue, many available large-scale

trees are under-resolved (NCBI taxonomy) or are generated with methods unsuited for

narrow and broad evolutionary distances (Hördt *et al.*, 2020) reducing their utility at multiple

evolutionary scales. This flood of genetic data necessitates an easy-to-use phylogenetic

analysis pipeline to help reveal the evolutionary context for the myriad of sequenced

bacteria.

1.1 Analysis options

1.1.1 Single and Multi-locus

A common way to generate bacterial trees is to compare single homologous loci (e.g.,

16S ribosomal, recA, gyrA, rpoB, or dnaK genes). This method has advantages which include

the ability to use conserved primer sites to amplify sequences, very low computational

intensity, and the availability of curated databases (SILVA, NCBI GenBank, etc.). Using genes

other than rDNA allows for greater resolution at certain evolutionary distances but still lacks

broad resolution (i.e., Near resolution for swiftly evolving genes, far for conserved) (Yang,

1998). These trees are prone to misleading topologies due to horizontal gene transfers,

incomplete lineage sorting, paralogous gene conversions, and limited phylogenetic

information (Huerta-Cepas *et al.*, 2007). Using multi-locus alignments to build phylogenetic

trees increases the total number of phylogenetically informative sites (more sequence used),

reduces the effects of incongruent topologies, and widens the breadth of resolvable

evolutionary resolution (Gontcharov *et al.*, 2004). However, in multi-locus analyses, the

effort and cost of primer design and amplification optimization scales with the number of

sequences used, which can rapidly become prohibitive.

1.1.3 Whole genome

Classically, whole genome alignments were required to identify Single Nucleotide

Polymorphisms (SNPs) of assemblies (see Shakya *et al.*, 2020; Treangen *et al.*, 2014 for

method details). These methods add considerable phylogenetic data for tree building and

can produce different topologies than multi-locus alignments, illustrating their utility (Baltrus

et al., 2014). A major drawback of whole genome alignments is that they tend to fail for

genomes with <85% identity depending on the aligner used (Shakya *et al.*, 2020). Thus,

phylogenetic information drops quickly as a function of evolutionary distance. The practical

result is a lack of resolution at even moderate evolutionary distances. Additionally, having

the "missing" data enriched for the more divergent sites will possibly lead to an

underestimation of the sequence to reference evolutionary distances for more divergent

sequences (Bertels *et al.*, 2014; Spencer *et al.*, 2007; Shavit Grievink *et al.*, 2013).

1.1.4 K-mer overlap

An emergent method to quickly estimate evolutionary histories from genome

assemblies is to use *k*-mer content. This involves identifying subsequences of length *k* within

each assembly and calculating the shared *k*-mer content (for details see Bussi *et al.*, 2021).

The advantage of this class of methods is that they are extremely fast relative to alignment-

based methods. The comparison of *k*-mer overlap between sequences naturally leads to

pairwise distance matrices, thus, neighbor-joining trees can be generated directly from the

results (Saitou and Nei, 1987). Crucially, *k*-mer distances quickly become non-linearly

associated with percent identity, leading to erroneous estimated branch lengths (Jain *et al.*,

2018).

An interesting advance in *k*-mer based approaches is kSNP (Gardner and Hall, 2013;

Gardner *et al.*, 2015), which identifies SNPs by flanking conserved *k*-mers. This approach

allows the use of tree building methods including parsimony, maximum likelihood, and

Baysian. Like whole genome alignments and *k*-mer overlap methods, kSNP suffers from a

rapid decline in phylogenetically informative data as sequences diverge. Additionally, the

signal/noise ratio degrades quickly because non-homologous *k*-mers are increasingly

common as sequences diverge (Gardner and Hall, 2013).

1.1.5 Whole transcriptome/proteome alignments

To use the wealth of data generated by NGS methods and simultaneously alleviate the

problem of whole genome alignments, whole transcriptome/proteome alignment can be

used. This method breaks the problem of identifying homologous loci into two tractable

parts: gene identification, then identification of homologous sequences. This approach

identifies many phylogenetically informative sites (like whole genome), but alignments are

more tractable (like single loci). However, gene-based alignments are negatively affected by

paralogs, i.e., duplicated genes or contamination. Thus, most methods filter out gene

families with paralogs. However, see ASTRAL-Pro for a method that is "paralog aware"

(Zhang *et al.*, 2020).

A trade-off arises while using predicted transcript/protein sequences to build species

trees. Transcripts have the most phylogenetic information (silent mutation) but are hard to

align at high divergencies (States *et al.*, 1991). Proteins remain alignable at high divergences

but lack the information content of transcripts. This can be addressed by converting protein

alignments to corresponding codon alignments, leveraging nucleotide phylogenetic

information and protein alignment accuracy (Wernersson and Pedersen, 2003; Bininda-

Emonds, 2005).

1.2 Benefits of an automated workflow

The skills required to generate bacterial phylogenetic trees from whole genomes are

extensive. Some of the many steps in building robust species trees from whole genome

assemblies include gathering and annotating assemblies, identifying and filtering orthologs,

sequence alignment, trimming and concatenation, and finally, tree inference (Ashford *et al.*,

2020). Each one of these steps, especially for large data sets, requires expertise in picking

parameters, file management, and data format manipulation and filtering. Most of these

steps also require familiarity with UNIX command-line.

The phylogenetic context of organisms is becoming increasingly important for standard

molecular and evolutionary studies. Even with the available data and a clear need for the

phylogenetic placement of focal study species, technical barriers preclude many researchers

from inferring their evolutionary histories. An easy-to-use phylogenomic pipeline will

encourage wide adoption of phylogenomic methods to complement ongoing environmental

and clinical microbiological research, and perhaps help standardize the estimation of

phylogenetic trees within and between research labs.

To meet this demand, we present OrthoPhyl, a phylogenomic pipeline that takes bacterial

genomes as input, annotates them, identifies orthologs, converts protein to nucleotide

alignments, and builds species trees with both concatenated alignments and gene tree to

species tree reconciliation. The workflow accepts an arbitrarily large number of input

genomes (tested up to 1200). To speed up analysis, a subset of samples representing the

whole dataset's diversity are identified, their proteomes are used to build hidden Markov

models and then identify orthologs in the full dataset. This strategy allows the generation of

trees for 689 Brucella assemblies in ~48hrs using 30 cpus and 58.3GB memory with no

hands-on time required. This pipeline is designed to be an easy to install, scalable, turn-key

solution for generating high-resolution bacterial trees from diverse clades.

2 Methods

2.1 Main Pipeline

OrthoPhyl's workflow can be broken down into four main steps. Genome assemblies are

annotated (**Figure 2.A**), resulting proteins are assigned to orthogroups (**Figure 2.B**),

orthogroup proteins are aligned and converted to codon alignments (**Figure 2.C**), and finally,

concatenated codon alignments are used to infer phylogenetic trees (**Figure 2.D**).

2.1.1 Annotation

OrthoPhyl starts by annotating assemblies with Prodigal (Hyatt *et al.*, 2010). This

provides both protein and transcript sequences that are used in subsequent steps.

Redundant transcripts (> 99.9% nucleotide identity) found in a genome are removed using

bbmap's *dedupe* (BBMap, 2022) with the logic they are likely the result of recent gene

duplications and provide no or little phylogenetic information. This removes unnecessary

paralogs from the dataset to preserve the usefulness of the containing orthogroup for

downstream analysis.

Figure 1 - Visual workflow diagram of OrthoPhyl. Grey boxes indicate processes. Programs used in each step are listed unless a custom script was used. Orange, tan, and purple boxes represent user input, intermediate files, and species tree outputs, respectively. The workflow is divided into four main tasks **A)** annotate assemblies and remove identical transcripts, **B)** if more than "N" assemblies are to being analyzed, identify orthologs in a diversity spanning subset of assemblies, then expand to the full dataset with iterative HMM searches, **C)** align protein sequences and generate codon alignments, finally **D)** estimate species tree topologies with concatenated codon alignment supermatrix along with a gene tree to species tree consensus method.

2.1.2 Orthogroup assignment

To identify orthogroups (OGs) to use for tree generation, *OrthoFinder* is used (Emms and

Kelly, 2015, 2019). When trees are being generated for large numbers (default: > 30) of

genome assemblies, finding orthogroups directly becomes intractable. Thus, OrthoPhyl

identifies a diversity spanning subset of assemblies to identify OGs. Briefly, average

nucleotide identity (ANI) is estimated for all assembly pairs with *FastANI* (Jain *et al.*, 2018). A

custom hierarchical clustering-like algorithm is used to cluster assemblies based on pairwise

ANI values, then single representatives for each of the N groups (default: 30) are chosen as

input for OrthoFinder. *FastANI* fails to give an ANI percent for genome comparisons with ANI

less than ~75%. For these instances, an arbitrary value of 50% is used. The representative

genome picking will be unaffected if there are less than 30 clades at this or greater distances

to all other clades. Even with several clades at greater than 75% ANI, the ultimate list of OGs

is unlikely to be affected.

To expand identified OGs to the full assembly dataset, OG proteins are realigned with

Mafft (Katoh and Standley, 2013), then *hmmer* (Eddy, 2008, 2009, 2011) generates hidden

Markov models and searches against all predicted proteins. OrthoPhyl optimizes the HMM

hit score cutoffs to maximize the number of taxa with proteins in the OG, while removing

paralogous hits. The expanded protein set for each OG is then realigned and HMMs are

again generated and searched against all proteins to capture additional, more divergent

orthologues.

2.1.3 Codon Alignment generation

Once OGs are identified for the full set of assemblies, protein sequences are realigned

with *Mafft* (Katoh and Standley, 2013). These alignments are then used as the basis for

codon alignment using *PAL2NAL* (Suyama *et al.*, 2006). Codon alignments are then trimmed

with *TRIMAL* (Capella-Gutiérrez *et al.*, 2009) to reduce noise associated with alignment

edges. *Alignment_Assessment* (Portik *et al.*, 2016), is used to assess the quality of codon

alignments, including their phylogenetic signals, taxa per alignment, and alignments per taxa

(Figure 2). This is critical for ensuring high-quality data are used for generating trees and can

aid in troubleshooting phylogenies with low branch support.

Our tool then classifies OGs as relaxed or strict single-copy orthologs (SCOs). Strict SCOs

are genes found in every taxon with no paralogs, while relaxed SCOs are found in X% of

assemblies still with no paralogs, with X being a tunable parameter (default:30%). Moving

forward, these two sets of OGs are used in parallel to generate species trees.

Figure 2 – Alignment metric visualization generated by OrthoPhyl. A) Histograms of the number of taxa in each alignment is shown. Dotted red line indicates the >30% taxa cutoff for alignments used to generate the "SCO relaxed" trees. B) Amount of missing data (including gaps and ambiguous bases) per alignment is shown with a historgram. This is exclusive of genomes without detected orthologs. C) The number of phylogenetically informative sites (variation in > 1 taxa) compared to alignment length for each orthogroup are shown. A linear regression is shown with a solid red line with associated slopes. A one-to-one relationship is shown as a black dotted line.

2.1.4 Species Tree Estimation

Two methods are used to generate species trees for the input assemblies, gene-tree to

species-tree estimation with *ASTRAL-III* (Zhang *et al.*, 2018) and gene alignment supermatrix

to maximum likelihood (ML). First, to use *ASTRAL*, ML gene-trees are generated with either

FastTree (Price *et al.*, 2010) or *IQtree* (Nguyen *et al.*, 2015; Kalyaanamoorthy *et al.*, 2017;

Hoang *et al.*, 2018) per user input. Then strict and relaxed SCO trees are used as input to

ASTRAL to generate species trees. Trees are also generated from concatenated codon

alignments and maximum likelihood methods. The user can specify any combination of

IQtree, *FastTree* or *RAxML* (Stamatakis, 2014) to infer tree structure. Finally, these trees are

compared with generalized Robinson-Foulds metrics provided by ETEtoolkit (Huerta-Cepas

et al., 2007).

2.2 Datasets

As a proof of concept, OrthoPhyl was used to build trees for three bacterial clades. We

analyzed well characterized *E. coli*/*Shigella* strains. These included a list of 34 complete

genomes analyzed in (Shakya *et al.*, 2020), obtained from NCBI.

The genus *Brucella* was used to show OrthoPhyl's ability to resolve very closely related

sequences while dealing with very long relative branch lengths in the same tree

(*Ochrobactrum* group). Assemblies were acquired using NCBI Taxon number 234 with the

utility script "*gather_genomes.sh*" (supp. Methods - Gathering and Filtering Assemblies and

Supplementary Figure 1) which is packaged with OrthoPhyl

(github.com/eamiddlebrook/OrthoPhyl/utils/). This resulted in 689 assemblies after filtering

for >98% completion, < 0.1% duplication, and < 1% contamination with CheckM (Parks *et al.*,

2015). It is important to note that *Ochrobactrum* species were recently moved to the genus

Brucella (Hördt *et al.*, 2020). However, for clarity and due to controversy within the field, we

chose to keep the *Ochrobactrum* labeling for these species. An assembly of *Mycoplana*

dimorpha (GCA_003046475.1) was added to the dataset as an outgroup.

Finally, a tree for all NCBI Rickettsiales assemblies was generated to illustrate

OrthoPhyl's utility in dealing with bacterial order level divergences. For Rickettsiales

assemblies, we again used *gather_genomes.sh,* this time with NCBI taxon number 766. To

illustrate a more challenging scenario, filtering of Rickettsiales assemblies was less stringent

with cut-offs at > 95% completion, < 0.2% duplication, and < 1.5% contamination, again

using CheckM (Parks *et al.*, 2015). This resulted in 1202 assemblies. Seven *Pelagibacter*

assemblies were added to this dataset to serve as an outgroup. Accessions and stats for all

genome assemblies used in this manuscript are available in **Supplementary Table 1-3**.

3 Implementation

3.1 Small Tree of Closely Related Assemblies: *E. coli and Shigella*

To illustrate Orthophyl's ability to build trees from moderate numbers of samples on a

local Linux computer, we ran our *E. coli* analysis (34 genomes) on a desktop with a 12-core

Intel processor with 16gb ram running RHEL8 (centOS8). Run statistics are provided in **Table**

2. The full pipeline took less than 4 hours to build 2 maximum likelihood trees with *RAxML*

and 2 ASTRAL trees, one each from the strict and relaxed SCO gene sets.

OrthoPhyl identified 2142 strict and 3426 relaxed SCOs (found in 10 genomes) in the

analyzed *E. coli* and *Shigella* assemblies. Alignment of these sequences constitutes 2.04 and

3.19 MB with 6 and 7.2% phylogenetically informative sites for the SCO and relaxed SCO

sets, respectively (**Table 1**).

The resulting tree build by FastTree2 from the SCO-relaxed dataset shows successful

differentiation of the *E. coli* phylotypes **(Figure 3)**. This tree shows B1 and A as sister

phylotypes, with *S. sonnel, S. boydii* and *S. flexneri* between them. That clade is in turn next

to phylotype E and *S. dysenteriae*, then D1 and D2. Finally, B2 is sister to all of them. This

topology is identical to the reported tree from (Shakya *et al.*, 2020) using whole genome

alignments, except for the placement of the root (*E. fergusonii)*, where they have the root

placed between D2/B2 and the rest while OrthoPhyl's tree indicates it should be placed

between B2 and the other phylotypes.

Table 1 - OrthoPhylo output, runtime, ram usage, storage

Metric	Assembly Data Set		
	E. coli + Shigella	Brucella	Rickettsiales
Number of assemblies	34	689	1208
Number SCOs - strict	2142	897	42
Total Length (bp)	2,040,000	855,924	29,248
% Informative SNPs	6	41.4	81.5
% missing data	0.3	0.5	2.3
ANI % (avg/med/min)*	97.6/99.2/89.9	95.4/99.8/63.7	83.2/57.3/48.1
Number SCOs − relaxed	3426	1762	348
Total Length (bp)	3,190,000	1,627,004	295,820
% Informative SNPs	7.2	43.3	78.9
% missing data	9.3	2.7	10.6
ANI % (avg/med/min)*	97.3/99.0/89.1	95.0/99.8/63.2	82.6/55.7/48.0
CPU time (hr:min)	50:25:00	373:09:00	152:00:00
CPU efficiency	19%	24%	41%
Total Runtime (hr:min)	9:47	48:34:00	12:10
Max Mem Used (GB)	>16	58.34	17.3

*ANI percentage is calculated with ExANIFA (github.com/eamiddlebrook/ExANIFA) on the final
concatenated codon alignments excluding gapped positions. See Supplemental Methods for details.

Figure 3 - Strict single copy ortholog based maximum likelihood phylogeny of 34 Escherichia and Shigella assemblies. The tree was constructed with 2142 genes totaling 2.04MB of sequence. Tips are labeled with the given species and strain from NCBI's BioSample database (details in supplementary table 1). Bars next to tree indicate *E.coli* phylotypes [Shakya 2017]. The tree was rooted at *E. fergusonii* then root node was removed. The red arrow indicates the alternative root placement from Shakya 2017.

3.2 Genus level Species trees of 689 assemblies: *Brucella/Ochrobactrum*

For the 689 Brucella/Ochrobactrum assemblies analyzed (plus *Mycoplana dimorpha*

outgroup), OrthoPhyl was run on a RedHat8 compute node with 30 CPUS and 500 GB of ram.

The full analysis (generating strict and relaxed SCO with *FastTree2* and *ASTRAL* trees) took

just over 2 days (48hrs 31 min). Total ram usage was moderate at 58GB. CPU usage

efficiency was 25%, with a total CPU time of approx. 373 hours **(Table 1)**.

OrthoPhyl identified 785 strict and 1635 relaxed SCOs in the full dataset of assemblies.

The distribution of taxon number represented in each alignment shows that most

orthogroup alignments have all taxa represented, with approximately 1400 OGs having 680-

689 taxa (**Figure 2 A**). The OG alignments have every little missing data (gaps or ambiguous

bases) with the vast majority showing less than 10% (**Figure 2 B**). **Figure 2 C** shows the

number of informative sites vs. each alignment's length. A slope of 0.42, indicates the

alignments are far from the value that would constitute noisy alignments or over-saturation

of mutations.

The maximum likelihood (ML) *Brucella* phylogeny generated by FastTree based on the

1635 relaxed SCOs **(Figure 4)** shows very close agreement with trees generated by (Ashford

et al., 2020) for the *Ochrobactrum* clades (see **Supp. Figure 2** for high resolution tree with all

accessions and species labeled). For instance, it supports two major *Ochrobactrum* clades,

Group A - *O. anthropi, O. lupini, O. tritici. O. pecoris, O. oryzae, O, cicero, O. intermediaum,*

O. pseudointermedium, and *O.daejeonense,* and Group B - *O. grignonenense, O.*

pituitosa(pituitosum), O. quorumnocens, O.rhizosphaerae, O. pseudo grignonenense, O.

thiophenivorans, and *O. gallinifaecis* **(Figure 4 orange bars)**. Like Ashford et al., our tree

Figure 4 – A) Maximum likelihood tree of Brucella/Ochrobactrum assemblies generated by FastTree within OrthoPhyl. If provided by NCBI, species are labeled. Branch colors. To aid in visualization of mono/polyphyly, branches are colored by reported species. Internal branch colors are given based on majority rule of offspring. Branch lengths represent relative number of mutations per site. The tree was rooted with *Mycoplana dimorpha* that was subsequently removed, and the classic Brucella clade was collapsed. **B)** Subtree of the collapsed classic Brucella clade. A high-resolution version of the full tree showing Species, strain, accession, and bootstrap support is available in **Supplementary figure 2**.

supports *O. endophytica* as basal to the other *Ochrobactrum*. However, the clade with *O.*

haematophila, O. soli and *O. teleogrylli* is shown as sister to Group B *Ochrobactrum* instead

of Group A. A major distinction between our tree topology and the one presented in Ashford

et al. is that our tree shows the monophyletic Brucella clade splitting the *Ochrobactrum*

clade between *O. endophytica* and all other *Ochrobactrum*. In contrast, the Brucellae in

Ashford et al. split *Ochrobactrum* between Group A and B, as in (Hördt *et al.*, 2020). This

difference could be a product of having different outgroups (*Mycoplana ramose, M.*

dimorpha, and *Rhizobium etli* for Ashford et al. and just *M. dimorpha* for OrthoPhyl), and our

tree being subject for long branch attraction between *Brucella* and the root. Additionally,

our tree indicates that one *O. thiophenivorans* and *O. rhizosphaerae* sample were likely

mislabeled when uploaded, as both have examples rendered polyphyletic by assemblies of

many other species with high support.

Ochrobactrum lupini and *O. ciceri* are closely associated with *O. anthropi* and *O.*

intermedia (respectively). Like Ashford et al. comparing field isolates to type strains, we had

many *O. anthropi* and *O. intermedia* assemblies which show that *O. lupini* and *O. ciceri* arose

from within their respective clades and are likely misclassified as species, but instead should

be classified as *O. anthropi* and *O. intermedia*, respectively. For *O. lupini* and *O. anthropi*, this

is supported by previous work (Gazolla Volpiano *et al.*, 2019).

The clade comprising the classical *Brucella* strongly agrees with published phylogenies

(Suárez-Esquivel *et al.*, 2020; Ashford *et al.*, 2020). *Brucella vulpis* at the base*, B.*

pinnipedialis arising from within *B. ceti* (rendering *B. ceti* paraphyletic) (Orsini *et al.*, 2022), *B.*

canis arising from within *B. suis* (rendering *B. suis* paraphyletic), and *B. melitensis* and *B.*

abortus being sister groups. The classical *Brucella* have very short branches at the crown (*B.*

microti, B.neotomae, B.ovis, B. suis/canis, B. abortus/melitensis and *B. pinnipedialis/ceti*),

indicating very rapid divergence times for these clades, as seen elsewhere (Ashford *et al.*,

2020; Suárez-Esquivel *et al.*, 2020; Orsini *et al.*, 2022).

3.2.1 kSNP vs. OrthoPhyl

To compare performance of OrthoPhyl with an alternative assembly-to-tree method,

kSNP4 (Gardner *et al.*, 2015) was run with the same hardware as OrthoPhyl. See

supplementary methods for details. The resulting kSNP4 alignment shows 7.15 million SNPs

were identified, considerably longer than the maximum genome length of the dataset. As

expected, much of this SNP alignment consisted of missing data (~78%) indicating *k*-mers not

being shared between assemblies.

FastTree2 was used externally to generate a tree with the *k*-mer based SNPs. To reduce

runtime, memory usage, and gap-based noise, SNPs found in less than 30% of assemblies

were removed, with a minimum of 20% being preserved, this left 1,569,787 SNPs (exactly

20% of original SNPs, with ~13% total missing data). This SNP file was then used to generate

a tree with FastTree2 using the GTRGAMMA model. For a high-resolution tree with species,

strain, and assembly accessions labeled, see **(Supp. Figure 3)**. For deep branches outside of

the classic Brucella, this tree shows largely the same topology as the tree generated by

OrthoPhyl. Some notable exceptions within the Ochrobactrum include: 1) the placement of

O. gallinifaecis and one of the *O. thiophenivorans* being sister to *O. pseudogrignonensis* in

the OrthoPhyl tree while it is sister to the clade of *O. pseudogrignonensis, O. quorumnocens,*

O, pituitosa (among others) in the kSNP based tree **(Figure 4A, orange arrow)**, and 2) *O.*

daejeonensis is placed one node more basal in the kSNP tree **(Figure 4A, red arrow)**. For the

Brucella clade, there are 2 major differences. The first is the placement of the root, with

kSNP showing it coming in-between B. ovis and the rest of Brucella **(Figure 4B, black arrow)**.

The second is (given the OrthoPhyl root) *B. neotomae* being place, along with two B. suis

samples **(Figure 4B, asterisk)** on the short branch leading to the *B. melitensis, B. abortus, B.*

ovis, B. pinnipedialis, B. ceti clade, instead of sister to all *Brucella* internal to *B. microti*

(Figure 4B, purple arrow). Compared to kSNP4, OrthoPhyl had favorable runtime and

resource usage requirements. Major differences include max memory usage (58 vs 204 GB),

max storage (53 vs 480GB) and runtime (48 vs > 216 hours) for OrthoPhyl and kSNP4,

respectively.

3.2.3 Order Level Divergence Tree with 1200 samples: Rickesiales

Rickettsiales was chosen as a challenging taxon to test OrthoPhyl's ability to deal with

wide phylogenetic distances and extensive genome reduction. Additionally, this set of

assemblies was not filtered stringently, leaving a total of 1209 assemblies (7 outgroup) of

varying quality (see Datasets section above). As with the *Brucella* analysis, OrthoPhyl was run

on a 30 cpu compute node with 500 GB of available ram. The program took just 12 hours

and 10 minutes to complete using a total of 17.3 GB memory. The CPU usage efficiency was

41%, with a total CPU time of approx. 152 hours **(Table 1)**.

Even with this challenging genomic dataset, OrthoPhyl identifies > 29,000 homologous

base pairs from strict SCOs and 295,000 bp from relaxed SCOs (found in 366 or more

assemblies). These homologous sequences come from for 42 and 348 genes respectively

(**Table 1**). The relaxed SCO FastTree ML tree reconstructed from the 1202 Rickettsiales

assemblies recovers all valid genera as monophyletic (**Figure 5A**). The sole paraphyletic

genus recovered is *Candidatus Jidaibacter*. Other than *Jidaibacter*, the recovered topology of

genera is consistent with previously published analyses (Salje, 2021; Schön *et al.*, 2022), with

Figure 6 - A) Maximum likelihood tree for Rickettsiales generated by FastTree within OrthoPhyl. Branch colors show reported genera for each sample according to NCBI metadata. To aid in identification, labels are also provided. Samples with no genus level identification are labeled in grey. Internal branch colors are given based on majority rule of offspring. Note the red Wolbachia clade has been compressed vertically (1/5 ratio) thus clade widths are not proportional to number of samples across the tree. Branch lengths are relative probability of a mutation per site. The tree was rooted with a *Pelagibacter* outgroup that was subsequently removed. Candidate genera (Candidatus) are labeled Can. The single, genus level polytomy is labeled in red (*Canidatus-Jidibacter*). **B)** The *Rickettsia* subtree is shown with monophyletic species identifiers collapsed. Following tip labels in parentheses is the number of collapsed tips. The two most basal branches are omitted, as they are only identified to genus. All assemblies with species level classification show monophyly with the exception of *R. conorii (red arrows)* and *R. massiliae/R. rhipicephali (orange arrow).*

two major clades 1) *Wolbachia* sister to *Ehrlichia* and *Anaplasma*, with *Neoricketsia* basal to

them and 2) *Rickettsia* and *Orientia* separated from the rest by the root. A high-resolution

tree is provided in **Supplementary Figure 4,** which provides species, assembly accession and

number of SCOs per sample.

To illustrate OrthoPhyl's ability to resolve narrow evolutionary windows along with

order level divergences in the same tree, the species rich *Rickettsia* genus subtree is shown

(Figure 5B). Only three *Rickettsia* with species level identification (provided by NCBI

metadata) show polytomy. First, samples labeled *R. conorii* are placed in 3 very different

clades across the tree **(Figure 5B, red arrows)**, indicating that they are likely misidentified.

Analysis in Hördt et al, using digital DNA:DNA hybridization, also indicates *R. conorii* and

related species need revision. Additionally, the placement of *R. massiliae* and *R. rhipicephali*

samples render each other polyphyletic (**Figure 5B**, orange arrow). Excluding these samples,

the estimated *Rickettsia* tree is identical to the one presented in [Diop 2017].

<u>**4 Key features**</u>

4.1 Codon alignment based

Most ortho-gene methods use protein sequences, which are easy to align. However, our

software creates codon alignments of orthogroups, giving it two major advantages. Using

codon alignments to infer phylogenetic trees leverages protein alignment accuracy. Because

proteins have a 20 characters while nucleotides have only 4 in their "alphabet", protein

alignments have a much better signal-to-noise ratio (States *et al.*, 1991). Using nucleotide

alignments adds phylogenetic signal through codon degeneracy, allowing greater amounts of

information per gene (Wernersson and Pedersen, 2003; Bininda-Emonds, 2005).

4.2 Many input assemblies

This tool can generate trees with >1000 assemblies. It does this by identifying

orthogroups in a diversity-spanning subset of assemblies, then assigning proteins to these

orthogroups for the full set of predicted proteins by iterative HMM searches. This reduces

the computationally demanding all-vs-all protein searching that would otherwise be

required while retaining the ability to identify the majority of orthogroups spanning the

dataset.

4.3 User friendly

OrthoPhyl manages the formatting and organization of intermediate files, which is a

non-trivial task in a phylogenetic workflow such as this. As with any phylogenetic method,

alignment and tree estimation parameters will likely need to be tuned to obtain robust

results, thus we exposed these variables for altering at runtime. If OrthoPhyl fails due to

input error or lack of memory recourses, it can restart at the last completed step. Being

robust to restarting also allows users to tune parameters without rerunning the entire

pipeline.

<u>5 Discussion</u>

OrthoPhyl successfully reconstructs phylogenetic trees for clades with broad

evolutionary divergences. This is achieved with little user input; only requiring one command

line input pointing the pipeline to the assembly and output directories. OrthoPhyl can

generate trees for >30 bacterial assemblies on a laptop or >1000 assemblies on a

workstation or single compute node with moderate resources (30 cpus and 100GB RAM).

Many useful options are provided by OrthoPhyl, including alignment trimming parameters,

number of assemblies from which to identify initial SCOs, tree building software to use, and

evolutionary models for tree estimation.

With few exceptions, the trees estimated for the *E. coli/Shigella,*

Brucella/Ochrobactrum, and Rickettsiales datasets are consistent with published topologies.

This illustrates OrthoPhyl's ability to resolve broad evolutionary relationships: from highly

similar genomes with ANIs of >99% for *E.coli/Shigella* and *Brucella/Ochrobactrum* to highly

dissimilar assemblies with ANIs of 57% for the Rickettsiales. With OrthoPhyl being able to

generate whole genome-based phylogenies for more than 1000 assemblies, it fills a large

gap in phylogenomic analysis software, providing an easy to use, assembly-to-tree tool that

will help many researchers incorporate evolutionary analysis into their ongoing bacterial

studies.

The *Brucella/Ochrobactrum* and Rickettsiales trees are, to our knowledge, the largest

phylogenies published for these clades, let alone using whole genome methods. While more

work is necessary to validate the topologies presented here, they are a starting point for

assessing metadata-based species labels and expected clade monophyly. Future work will

test the robustness of topologies to different evolutionary models and maximum-likelihood

methods.

5.1 Pipeline limitations

There are many parameters to tune during the various steps within this pipeline. Since we focus on usability there is an inevitable tradeoff with optimization, mainly through automated parameter choice. Therefore, this pipeline is useful for "democratizing" the generation of large bacterial phylogenetic trees but does not produce publication ready trees without user inspection of intermediate file such as alignment quality, single copy ortholog numbers, percent phylogenetically informative sites, and of course bootstrap values. Ultimately, trees should be compared to existing data to ensure major clades are consistent, and if not, total evidence for novel branching should be investigated.

5.2 Future

Future versions of this software package will include several additional evolutionary analyses. Bayesian tree inference methods will be incorporated into the pipeline to allow more flexibility in tree estimation methods. Since codon alignments of single copy orthologs are generated while running this pipeline, we also plan on integrating a software package, e.g., ETE3 (Huerta-Cepas *et al.*, 2016), to test evolutionary models such as positive/negative selection and neutrality. A "forest of life" based horizontal gene transfer analysis will be added using gene tree comparisons. This analysis will look at k-means clustering of gene trees to build multiple consensus trees to identify if blocks of genes do not agree with a single consensus species tree hypothesis (Puigbò *et al.*, 2019).

In the current iteration of OrthoPhyl, very ancient paralogs (originating before the most recent common ancestor of the dataset species) might be clustered into the same orthogroup, leading to the orthogroup's removal during filtering even though the 2 sub-trees of the orthogroup could be consistent with the species tree. Future iterations of this

software will filter the orthogroups in a deep paralog aware manner, using a rapidly

generated neighbor-joining species tree to identify and rescue such orthogroups.

Although this pipeline is geared towards prokaryotes, only two changes are needed to

adapt it to eukaryotic phylogenetics: 1) allowing users to input precomputed annotations

directly, avoiding the complex task of Eukaryotic annotation within the workflow itself and 2)

taking into consideration paralog subtrees which are consistent with a consensus species

tree. The next version of OrthoPhyl will incorporate these changes to expand the use cases

to Eukaryotic phylogenetics.

OrthoPhyl identifies gene family groups for the dataset provided. While this allows the

workflow to adapt to any set of genome assemblies, it requires considerable compute

resources to complete. Alternatively, predefined gene family models could be used as a

starting point, then the un-clustered protein sequences would be fed into the standard

OrthoPhyl orthogene identification pipeline to capture additional gene families with

phylogenetic signal.

5.3 Conclusion

Phylogenetic reconstruction is critical to understanding the evolution of pathogenicity,

novel traits, horizontal gene transfer, and metabolic potentials of bacteria. Unfortunately,

large-scale phylogenomic analyses of diverse bacterial groups require deep understanding of

bioinformatic methods and great care in dealing with the myriad intermediate files and

formats used during the analysis. To our knowledge, OrthoPhyl is the only software to take

>1000 bacterial assemblies with order level divergence, generate ortholog codon

alignments, then reconstruct phylogenetic trees without user input or management of

intermediate files being necessary. Thus, OrthoPhyl will allow many research groups,

including those with modest computing resources, to leverage of the wealth of publicly

available genomic data to enrich their ongoing analyses with phylogenomic inferences across

a broad swath of bacterial diversity.

7 Acknowledgments

The authors would like to thank Migun Shakya, Taehyung Kwon and John Gillece for

their valuable insight and comments and the whole Disease Surveillance and Molecular

Epidemiology of Brucella project team members in Tanzania and Rwanda, especially Prof.

Joram Buza for comments and suggestions.

This research was funded by the Defense Threat Reduction Agency through Triad

National Security, LLC, operator of the Los Alamos National Laboratory under Contract No.

89233218CNA000001 with the U.S. Department of Energy.

8 Conflict of interest
The authors declare no conflict of interest.
460
461
462   **9 Bibliography**
463

Ashford,R.T. *et al.* (2020) Application of Whole Genome Sequencing and Pan-Family Multi-Locus
Sequence Analysis to Characterize Relationships Within the Family Brucellaceae. *Front.*
Microbiol., **11**.
Baltrus,D.A. *et al.* (2014) Incongruence between multi-locus sequence analysis (MLSA) and
whole-genome-based phylogenies: Pseudomonas syringae pathovar pisi as a cautionary
tale. *Mol. Plant Pathol.*, **15**, 461–465.
BBMap (2022) *SourceForge*.
Bertels,F. *et al.* (2014) Automated Reconstruction of Whole-Genome Phylogenies from Short-
Sequence Reads. *Mol. Biol. Evol.*, **31**, 1077–1088.
Bininda-Emonds,O.R. (2005) transAlign: using amino acids to facilitate the multiple alignment of
protein-coding DNA sequences. *BMC Bioinformatics*, **6**, 156.

Bussi,Y. *et al.* (2021) Large-scale k-mer-based analysis of the informational properties of
genomes, comparative genomics and taxonomy. *PLOS ONE*, **16**, e0258693.
Capella-Gutiérrez,S. *et al.* (2009) trimAl: a tool for automated alignment trimming in large-scale
phylogenetic analyses. *Bioinformatics*, **25**, 1972–1973.
Eddy,S.R. (2009) A new generation of homology search tools based on probabilistic inference.
Genome Inform. Int. Conf. Genome Inform., **23**, 205–211.
Eddy,S.R. (2008) A Probabilistic Model of Local Sequence Alignment That Simplifies Statistical
Significance Estimation. *PLOS Comput. Biol.*, **4**, e1000069.
Eddy,S.R. (2011) Accelerated Profile HMM Searches. *PLOS Comput. Biol.*, **7**, e1002195.
Emms,D.M. and Kelly,S. (2019) OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biol.*, **20**, 238.
Emms,D.M. and Kelly,S. (2015) OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. *Genome Biol.*, **16**,
157.
Gardner,S.N. *et al.* (2015) kSNP3.0: SNP detection and phylogenetic analysis of genomes
without genome alignment or reference genome. *Bioinformatics*, **31**, 2877–2878.
Gardner,S.N. and Hall,B.G. (2013) When Whole-Genome Alignments Just Won't Work: kSNP v2
Software for Alignment-Free SNP Discovery and Phylogenetics of Hundreds of Microbial
Genomes. *PLoS ONE*, **8**, e81760.
Gazolla Volpiano,C. *et al.* (2019) Reclassification of Ochrobactrum lupini as a later heterotypic
synonym of Ochrobactrum anthropi based on whole-genome sequence analysis. *Int. J.*
Syst. Evol. Microbiol., **69**, 2312–2314.
Gontcharov,A.A. *et al.* (2004) Are Combined Analyses Better Than Single Gene Phylogenies? A
Case Study Using SSU rDNA and rbcL Sequence Comparisons in the Zygnematophyceae
(Streptophyta). *Mol. Biol. Evol.*, **21**, 612–624.
Hoang,D.T. *et al.* (2018) UFBoot2: Improving the Ultrafast Bootstrap Approximation. *Mol. Biol.*
Evol., **35**, 518–522.
Hördt,A. *et al.* (2020) Analysis of 1,000+ Type-Strain Genomes Substantially Improves
Taxonomic Classification of Alphaproteobacteria. *Front. Microbiol.*, **11**.
Huerta-Cepas,J. *et al.* (2016) ETE 3: Reconstruction, Analysis, and Visualization of Phylogenomic
Data. *Mol. Biol. Evol.*, **33**, 1635–1638.
Huerta-Cepas,J. *et al.* (2007) The human phylome. *Genome Biol.*, **8**, R109.
Hyatt,D. *et al.* (2010) Prodigal: prokaryotic gene recognition and translation initiation site
identification. *BMC Bioinformatics*, **11**, 119.
Jain,C. *et al.* (2018) High throughput ANI analysis of 90K prokaryotic genomes reveals clear
species boundaries. *Nat. Commun.*, **9**, 5114.
Kalyaanamoorthy,S. *et al.* (2017) ModelFinder: fast model selection for accurate phylogenetic
estimates. *Nat. Methods*, **14**, 587–589.
Katoh,K. and Standley,D.M. (2013) MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. *Mol. Biol. Evol.*, **30**, 772–780.
Konstantinidis,K.T. and Tiedje,J.M. (2007) Prokaryotic taxonomy and phylogeny in the genomic
era: advancements and challenges ahead. *Curr. Opin. Microbiol.*, **10**, 504–509.
Nguyen,L.-T. *et al.* (2015) IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating
Maximum-Likelihood Phylogenies. *Mol. Biol. Evol.*, **32**, 268–274.

Orsini,M. *et al.* (2022) Brucella ceti and Brucella pinnipedialis genome characterization unveils
genetic features that highlight their zoonotic potential. *MicrobiologyOpen*, **11**, e1329.

Parks,D.H. *et al.* (2015) CheckM: assessing the quality of microbial genomes recovered from
isolates, single cells, and metagenomes. *Genome Res.*, **25**, 1043–1055.

Portik,D.M. *et al.* (2016) An evaluation of transcriptome-based exon capture for frog
phylogenomics across multiple scales of divergence (Class: Amphibia, Order: Anura).
Mol. Ecol. Resour., **16**, 1069–1083.

Price,M.N. *et al.* (2010) FastTree 2 – Approximately Maximum-Likelihood Trees for Large
Alignments. *PLOS ONE*, **5**, e9490.

Puigbò,P. *et al.* (2019) Genome-Wide Comparative Analysis of Phylogenetic Trees: The
Prokaryotic Forest of Life. *Methods Mol. Biol. Clifton NJ*, **1910**, 241–269.

Saitou,N. and Nei,M. (1987) The neighbor-joining method: a new method for reconstructing
phylogenetic trees. *Mol. Biol. Evol.*, **4**, 406–425.

Salje,J. (2021) Cells within cells: Rickettsiales and the obligate intracellular bacterial lifestyle.
Nat. Rev. Microbiol., **19**, 375–390.

Schön,M.E. *et al.* (2022) The evolutionary origin of host association in the Rickettsiales. *Nat.*
Microbiol., **7**, 1189–1199.

Shakya,M. *et al.* (2020) Standardized phylogenetic and molecular evolutionary analysis applied
to species across the microbial tree of life. *Sci. Rep.*, **10**, 1723.

Shavit Grievink,L. *et al.* (2013) Missing Data and Influential Sites: Choice of Sites for
Phylogenetic Analysis Can Be As Important As Taxon Sampling and Model Choice.
Genome Biol. Evol., **5**, 681–687.

Smith,D.R. (2013) The battle for user-friendly bioinformatics. *Front. Genet.*, **4**, 187.

Spencer,M. *et al.* (2007) Conditioned Genome Reconstruction: How to Avoid Choosing the
Conditioning Genome. *Syst. Biol.*, **56**, 25–43.

Stamatakis,A. (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinforma. Oxf. Engl.*, **30**, 1312–1313.

States,D.J. *et al.* (1991) Improved sensitivity of nucleic acid database searches using application-
specific scoring matrices. *Methods*, **3**, 66–70.

Suárez-Esquivel,M. *et al.* (2020) Brucella Genomics: Macro and Micro Evolution. *Int. J. Mol. Sci.*,
21, 7749.

Suyama,M. *et al.* (2006) PAL2NAL: robust conversion of protein sequence alignments into the
corresponding codon alignments. *Nucleic Acids Res.*, **34**, W609–W612.

Treangen,T.J. *et al.* (2014) The Harvest suite for rapid core-genome alignment and visualization
of thousands of intraspecific microbial genomes.

Varghese,N.J. *et al.* (2015) Microbial species delineation using whole genome sequences.
Nucleic Acids Res., **43**, 6761–6771.

Wernersson,R. and Pedersen,A.G. (2003) RevTrans: multiple alignment of coding DNA from
aligned amino acid sequences. *Nucleic Acids Res.*, **31**, 3537–3539.

Yang,Z. (1998) On the best evolutionary rate for phylogenetic analysis. *Syst. Biol.*, **47**, 125–133.

Zhang,C. *et al.* (2018) ASTRAL-III: polynomial time species tree reconstruction from partially
resolved gene trees. *BMC Bioinformatics*, **19**, 153.

Zhang,C. *et al.* (2020) ASTRAL-Pro: Quartet-Based Species-Tree Inference despite Paralogy. *Mol.*
Biol. Evol., **37**, 3292–3307.

563
564
565