A member of a new dynamin superfamily modulates mitochondrial

membrane branching in *Trypanosoma brucei*

Chloé Alexandra Morel[1], Corinne Asencio[1], David Moreira[2], Corinne Blancard[3],

Bénédicte Salin[3], Etienne Gontier[4], Stéphane Duvezin-Caubet[3], Manuel Rojo[3],

Frédéric Bringaud[1] and Emmanuel Tetaud[1]*

[1]Univ. Bordeaux, CNRS, MFP, UMR 5234, F-33000, Bordeaux, France

[2]Ecologie Systématique Evolution, CNRS, Université, Paris-Sud, Université Paris-

Saclay, AgroParisTech, Gif-sur-Yvette, France.

[3]Univ. Bordeaux, CNRS, IBGC, UMR 5095, F-33000, Bordeaux, France

[4]Univ. Bordeaux, CNRS, BIC, UAR 3420, F-33000, Bordeaux, France

*Correspondence: emmanuel.tetaud@u-bordeaux.fr

SUMMARY

Unlike most other eukaryotes, where mitochondria continuously fuse and divide,

the mitochondrion of trypanosome cells forms a single and continuously

interconnected network that divides only during cytokinesis. However, the

machinery governing mitochondrial remodeling and interconnection of

trypanosome mitochondrion remain largely unknown. We functionally characterize

a novel dynamin-superfamily protein (DSP) from *T. brucei* (*Tb*MfnL) which shares

close similarity with a family of homologs present in various eukaryotic and

prokaryotic phyla, but not in opisthokonts like mammals and budding yeast. The

sequence and domain organization of *Tb*MfnL is distinct and it is phylogenetically

very distant from the yeast and mammalian dynamin-related proteins involved in

mitochondrial fusion/fission dynamincs, such as Opa1 and Mfn. *Tb*MfnL localizes to

the inner mitochondrial membrane facing the matrix, and upon overexpression,

induces a strong increase in the interconnection and branching of mitochondrial

filaments in a GTPase-dependent manner. *Tb*MfnL is a component of a novel

membrane remodeling machinery with an unprecedented matrix-side localization

that is able to modulate the degree of intermitochondrial connections.

Keywords: Mitochondrial dynamics – Dynamin-superfamily protein - Membrane

remodeling - Evolution – *Trypanosomes*

INTRODUCTION

Trypanosoma brucei is a parasite responsible for African sleeping sickness as well as the related cattle disease Nagana, affecting sub-Saharan Africa. *T. brucei* is transmitted to mammals by an insect vector: the tsetse fly (*Glossina* spp.). During the trypanosome life cycle, the parasite exists in at least two replicative forms; the procyclic form (PCF), transmitted by the tsetse fly, and the bloodstream form (BSF) responsible for diseases in vertebrates. Unlike most other eukaryotes that contain numerous mitochondria, trypanosomes have a single mitochondrion, which is highly elongated and reticulated. In contrast to yeast and mammals, with tens to hundreds of mitochondrial DNA (mtDNA) nucleoids distributing throughout the mitochondrial compartment, the mitochondrial genome of trypanosome is restricted to a specific structure at the basis of the flagellum known as kinetoplast (kDNA) [1]. During the trypanosome life cycle and its adaptation to different hosts and environments, the shape of its mitochondrial compartment undergoes spectacular changes that reflect its functional plasticity [2]. Indeed, the mitochondrion of this parasite exists in at least two major forms: (i) the fully active and developed one characteristic for the PCF that harbors the oxidative phosphorylation complexes (OXPHOS) for energy production [3, 4] and (ii) the functionally less active and morphologically reduced form found in the bloodstream, with energy produced exclusively through glycolysis since OXPHOS is repressed [5]. These modifications correlate with the adaptation of the parasite to the proline-rich hemolymph and tissue fluids of the blood-feeding tsetse fly and the glucose-rich blood of a mammalian host [6]. Since studies in mammals and in yeast having unraveled tight links between mitochondrial bioenergetics, morphology and dynamics [7-9], it is tempting to assume that such close relationships not only regulate mitochondrial function and energy metabolism, but also the life cycle and pathogenic potential of trypanosomes. However, in contrast to fungi and metazoa, where mitochondrial morphology and fusion/fission dynamics have been thoroughly characterized [10, 11], little is known about the morphology and dynamics of mitochondria in trypanosomes.

In most eukaryotes, mitochondria are very dynamic, alternating between two major events, fusion and fission. Mitochondrial dynamics allows to maintain mitochondrial morphology, distribution and size [8]. The main proteins involved in these events are large GTPases belonging to the dynamin superfamily protein (DSP), such as Dnm1/Drp1 in yeast and mammalian cells, which are soluble

protein transiently recruited to mitochondrial and peroxisomal membranes to
mediate organelle division [12, 13]. Mammalian mitochondrial fusion requires
fusion of outer mitochondrial membrane (OMM) followed by fusion of inner
mitochondrial membrane (IMM) which is carried out by mitofusins 1 and 2 (Mfn1
and Mfn2) and optic atrophy 1 (Opa1), respectively [14, 15]; while mitofusins are
anchored to the OMM and face the cytosol, Opa1 localizes to the mitochondrial
inter-membrane space (IMS). Alteration of fusion/fission dynamics and of the
equilibrium between them provokes mitochondrial fragmentation or hyperfusion
[12, 16], and is linked to diseases, notably neuropathies [17].

The investigation of mitochondrial fission and fusion in trypanosomes has been
relatively limited until now. Available evidence indicates that mitochondrial fission
is restricted to cell division [18, 19] and may not occur continuously [18, 20] but
can be induced artificially [19, 21]. Evidence for mitochondrial fusion in
trypanosomes came 25 years ago from *in vivo* genetic exchange of mtDNA [22].
Finally, a recent study of mitochondrial dynamics in another trypanosomatid,
Crithidia fasciculata, unveiled the occurrence of mitochondrial fission, fusion, and
sliding throughout the cell cycle, pointing to the existence of mitochondrial fusion
and fission machineries in trypanosomatids [23].

Two decades of research have revealed that numerous mitochondrial fission and
fusion factors that, despite significant differences, appear conserved in fungi and
mammals [24]. In plants, mitochondria also fuse and divide. While plant DSPs
involved in fission are known, factors directly involved in fusion have not been
identified, suggesting the existence of a fusion machinery differing from that of
yeast and mammals [25, 26], [27]. So far, only two dynamin-like proteins (*Tb*Dlp)
have been described in *T. brucei* (Tb927.3.4720 and Tb927.3.4760): their ablation
impairs mitochondrial division and endocytosis and leads to cytokinesis arrest [19,
28, 29]. *Tb*Dlp thus appears to be the functional homolog of Drp1/Dnm1 in
mammals and yeast, but its fission activity on mitochondria seems restricted to
the cytokinesis [19]. Another presumed DSP (*Tb*MfnL, Tb927.7.2410) has been
also reported to be involved in mitochondrial fission [30]. To date, however, no
protein involved in mitochondrial fusion has been identified in trypanosomes.

In this study, we revisited the function of the trypanosome DSP *Tb*MfnL [30].
Phylogenic analysis reveals that this protein defines a new family of DSP found in
both eukaryotes and prokaryotes but absent from organisms containing Mfn or
Opa1 homologs (fungi and mammals) and from plants. We find that excess *Tb*MfnL

leads to increased mitochondrial branching and interconnexion and we show that
it is anchored to the mitochondrial inner membrane, facing the matrix. Taken
together, these findings suggest that *Tb*MfnL is not involved in fission, but
mediates or modulates mitochondrial interconnection and branching.
110

RESULTS

1 - Sequence analysis of *Tb*MfnL identifies it as a member of a novel
dynamin family.

To identify potential fusion players, we started our search by careful re-
examination of trypanosome genomic and proteomic databases for proteins
showing sequence homology to known fusion factors (Opa1/Mfn1-2/Fzo1). We
were only able to identify the previously reported dynamin-related protein, named
*Tb*Mfn-Like or *Tb*MfnL (Tb927.7.2410) [30], that displays a domain organization
that has more similarities with mammalian and yeast fusion factors of the outer
membrane (Mfn2/Fzo1) than of the inner membrane (Opa1) (**Figure 1A and**
Table S1). Analysis of the *Tb*MfnL sequence revealed several motifs and domains,
including an N-terminal region containing a predicted mitochondrial targeting
signal (MTS) with potential cleavage sites (**Figure S1**), followed by a GTPase
domain with conserved G1 to G4 motifs and two consecutive C-terminal
transmembrane domains (TMs) (**Figure 1A and Figures S1/S2**). The presence
of an N-terminal MTS makes it unlikely for *Tb*MfnL to localize to the outer
membrane, as Mfn/Fzo, and the absence of an N-proximal TM domain, which
prevents Opa1 import into the matrix, suggests that *Tb*MfnL is imported into the
matrix and does not represent a functional homolog of the fusion factors Opa1 or
Mfn.

We next investigated whether this DSP could be conserved in other organisms.
Using Blastp with *Tb*MfnL as a query to search various protein databases, we
identified well-conserved sequences in relatives of *Trypanosoma* belonging to the
class Kinetoplastea, including free-living bodonid species (**Figure S1 and Table**
S1 for examples). Surprisingly, we also discovered homologous proteins with high
similarity across the entire sequence in other very diverse Euglenozoa species,
including the classes Euglenida and Diplonemea, and also in various much more
distantly related eukaryotic lineages, particularly within the SAR clade, including
Stramenopiles and Alveolata (**Figure 1B, Figure S2 and Table S1** for example).
However, no proteins with the same general architecture were identified in the

Opisthokonta (the clade that includes fungi and animals) or in plants, the only sequences showing some similarity being restricted, as anticipated, to the well-conserved GTPase domain. Unexpectedly, we also identified several sequences similar to *Tb*MfnL in bacteria, mainly in the PVC group and in Proteobacteria (**Figure 1B, Figure S2 and Table S1** for example). Analysis of several of these proteins showed again a very conserved structural and domain organization along the whole sequence, with the exception of the absence of the potential mitochondrial targeting sequence (MTS) in prokaryotes (**Figure 1B and Figure S2**).

We conducted a maximum likelihood phylogenetic analysis of the identified sequences, incorporating several DSPs from mammals and yeast, as well as the cyanobacterial homolog DBLP as outgroup sequences (**Figure 1C and Figure S3**). Most eukaryotic sequences branched within a well-supported (99% ultrafast bootstrap support –UFBS-) group. Despite the fact that sequences within this clade tended to group according to their taxonomic origin, we observed several unexpected relationships (e.g., dinoflagellates branched within the Euglenozoa, and cryptophytes and haptophytes within the Stramenopiles). Moreover, colponemids (a group of deep-branching alveolates) formed a fully-supported (100% UFBS) group with two stramenopile sequences. Bacterial sequences formed a paraphyletic group at the base of this main eukaryotic clade, with two ameobozoan species (*Acanthamoeba castellanii* and *Dracoamoeba jomungandri*) robustly branching (97% UFBS) within one of the bacterial subgroups. This unexpected relationship strongly suggested that these amoebae acquired this gene by horizontal gene transfer (HGT) from bacterial donors. By contrast, the anomalous relationships among eukaryotic taxa observed within the main eukaryotic clade were more difficult to explain, although multiple gene duplications and losses appeared to have played a role to shape the current distribution of this gene in eukaryotes. The class Diplonemea provided evidence for this hypothesis, with two highly supported (100% UFBS) groups of sequences of the same genera (*Diplonema*, *Lacrimia*, and *Rhynchopus*) branching paraphyletically at the base of a large clade of dinoflagellate (Alveolata) sequences (**Figure S3**). A small group of sequences containing stramenopile, alveolate, and haptophyte species branched sister to all the other prokaryotic and eukaryotic sequences and most likely represented an ancient paralog of this gene, giving additional evidence for the role of duplication in the evolution of this gene. Furthermore, it is very likely that HGT

between different eukaryotic groups also occurred, as in the case of the dinoflagellate sequences cited above, which appear to have acquired this gene from euglenozoan donors, and in the case of the haptophyte and cryptophytes sequences that branched within the large group of stramenopile sequences.

Given the specific architecture of this DSP, its large phylogenetic distance to other previously known protein families, and its wide and distinct taxonomic representation in prokaryotes and eukaryotes –but its absence in animals and plants- strongly indicate that it represents a novel DSP associated with membrane dynamics.

2 - *Tb*MfnL is localized in the mitochondrion

To determine the subcellular localization of *Tb*MfnL (Tb927.7.2410), we added a 10xHA tag at the C-terminal extremity of the endogenous *Tb*MfnL (*Tb*MfnL$_{::10HA}$) in both PCF and BSF trypanosomes (**Figure 2A**). C-terminal tagging was used to avoid disruption of the potential N-terminal MTS. Western blot analysis with anti-HA antibody showed that *Tb*MfnL$_{::10HA}$ is about 2-fold more expressed in PCF than BSF, which is in agreement with the data obtained through SILAC and transcriptomic analysis for this protein [31, 32] (**Figure 2B**). The localization of the *Tb*MfnL protein was studied in PCF by immunofluorescence using threonine dehydrogenase (Tdh, [33]) as a mitochondrial matrix marker (**Figure 2C**). *Tb*MfnL$_{::10HA}$ and Tdh clearly colocalized in the mitochondrial compartment, but *Tb*MfnL depicted a less homogeneous mitochondrial distribution (**Figure 2C, Merge**). Unfortunately, we could not use the anti-Tdh immune serum for immunofluorescence analysis on BSF. Consequently, we performed a labeling of the mitochondria, with an anti-HSP60 (mitochondrial maker). Like observed in PCF, *Tb*MfnL$_{::10HA}$ and HSP60 colocalized in the mitochondrial compartment (**Figure 2C, Merge)**. The presence of a N-terminal targeting sequence and its localization by fluorescence microscopy demonstrated that *Tb*MfnL is a mitochondrial protein. These data are consistent with the mitochondrial localization of the C-terminal GFP-tagged *Tb*MfnL reported by the TrypTag database (cellular localization of trypanosome proteins [34]). Of note, the N-terminal GFP-Tagged *Tb*MfnL exhibited a cytoplasmic localization in TrypTag database [34], likely due to masking of the MTS pre-sequence.

3 – Inactivation and silencing of *Tb*MfnL does not alter mitochondrial shape

Live microscopy of yeasts and cultured mammalian cells has revealed that mitochondrial morphology is continuously remodeled by antagonistic fission and fusion reactions [16, 35]. Under normal conditions, the fusion/fission balance is established in favor of fusion and the mitochondria appear filamentous. Inhibition of fusion leads to mitochondrial fragmentation by ongoing fission [16] and reversely enhancing fusion leads to hyperfused mitochondria with increased branching and interconnection [12].

To investigate the role of *Tb*MfnL, we first inactivated the *Tb*MfnL gene by CRISPR/Cas9 [36]. Both *Tb*MfnL alleles were inactivated by integration of a puromycin resistance marker cassette into both PCF and BSF cloned cells (**Figure 3A**). Cloned parasites were analyzed by PCR to select cell lines carrying puromycin resistance gene insertions in both *Tb*MfnL alleles (*Tb*MfnL$^{-/-}$) (**Figure 3B**). No significant impact on cell growth was observed in these PCF and BSF null-mutants over a period of 7 days compared to parental cells (**Figure S4**) and no significant difference in the structure of the *Tb*MfnL$^{-/-}$ mitochondrion was observed by staining mitochondrion with rhodamine-123 or immunofluorescence analyses with the anti-HSP60 immune serum, in both PCF and BSF mutant cell lines, respectively (**Figure 3C and 3D**). To confirm that the CRISPR/Cas9 strategy leads to *Tb*MfnL gene inactivation, we repeated this process in a strain endotagged with a C-terminal 3xTy1 tag in order to follow the expression of the protein by western blot (**Figure S5A-D**) upon insertion of the puromycin resistance gene as above (**Figure S5E-G**) and deletion of part of the *Tb*MfnL gene (**Figure S5H-J**), as above. In both cases, the protein was not detected in the inactivated clones and no modification of the mitochondrial structure was observed (**Figure S5G/J**).

In order to detect and quantify discrete modifications of the mitochondrial shape that may escape detection upon qualitative visual analysis, we developed an ImageJ macro (**Data S1**) for automatic quantification of mitochondrial junctions from fluorescence images (**Figure 3C/D/E**). This allowed us to establish that the number of mitochondrial junctions is equivalent in *Tb*MfnL$^{-/-}$ and parental PCF cells (8.5±4.9 vs 9.4±4.3) confirming the qualitative analysis of the microscopy images (**Figure 3C**). The mitochondrial and cristae structures were also examined by transmission electron microscopy (TEM) in *Tb*MfnL$^{-/-}$ PCF cells cultured with either proline or glucose as the sole carbon source. Under these conditions, no significant

alterations in mitochondrial or cristae structure were observed, nor were there any noticeable effects on the growth of the mutant cells (**Figure S6**).

These findings regarding the consequences of *Tb*MfnL inactivation in PCF and BSF are not in line with a previous report describing a highly fenestrated mitochondrial network consecutive to a partial but significant inhibition (70%) of *Tb*MfnL expression by RNAi in BSF [30]. To exclude that the divergence arose from the different inactivation and silencing strategies, we conditionally down-regulated expression of *Tb*MfnL using a stem-loop RNAi construct. This approach led to a significant reduction (~70%) of the expression of endogenously tagged *Tb*MfnL$_{::10HA}$ in PCF (as observed in BSF [30]), but this was not paralleled by a modification of mitochondrial shape (**Figure S7A-C**). Since mitochondrial fenestration upon *Tb*MfnL silencing was observed in BSF [30], we decided to repeat the experiment using the same strain (*T. brucei* 427 BSF), the same RNAi system (p2T7-177) and the same sequence used for down-regulation [30]. As previously reported [30], we observed a significant reduction of *Tb*MfnL expression (~80% after 4 days) of the endogenously tagged *Tb*MfnL$_{::3HA}$ in BSF (**Figure S7D-F**) that was not accompanied by mitochondrial shape modifications (**Figure S7G**). In our hands, neither inactivation nor silencing of *Tb*MfnL alters mitochondrial morphology or induces a fenestration phenotype.

4 - *Tb*MfnL overexpression increases mitochondrial branching

We also investigated the possible role of *Tb*MfnL by conditionally overexpressing a C-terminally-tagged (3xTy1) *Tb*MfnL in PCF (^{oe}*Tb*MfnL$_{::3Ty1}$, **Figure 4A**). Expression was confirmed by western blot analysis after 3 to 5 days of tetracycline induction (**Figure 4B**). Rhodamine-123 staining showed that the parental and non-induced ^{oe}*Tb*MfnL$_{::3Ty1}$ cell lines harbor the expected reticulated mitochondrial structure, while the mitochondrion appears significantly more reticulated and branched in the tetracycline-induced ^{oe}*Tb*MfnL$_{::3Ty1}$ cells (**Figure 4C, Figure S8**). After 5 days of induction, the average number of mitochondrial junctions, quantified with the ImageJ macro, appeared more than 2 times higher in induced ^{oe}*Tb*MfnL$_{::3Ty1}$ cells compared to non-induced or parental cells (**Figure 4D**). In order to estimate the level of *Tb*MfnL overexpression achieved in *T. brucei* PCF, we overexpressed *Tb*MfnL$_{::3Ty1}$ in the *Tb*MfnL$_{::3Ty1}$ background, in which the endogenous *Tb*MfnL alleles were 3Ty1-tagged (clone 2A11; **Figure S5A-D**). Western blot analysis with the anti-Ty1 antibodies revealed that expression of *Tb*MfnL$_{::3Ty1}$ is 5.5-times increased

in the resulting tetracycline-induced TbMfnL$_{::3Ty1}$/^{oe}TbMfnL$_{::3Ty1}$ compared to the

parental TbMfnL$_{::3Ty1}$ clone, which only expressed the endogenously tagged

TbMfnL$_{::3Ty1}$ (**Figure S9A**). Overexpression of TbMfnL$_{::3Ty1}$ once again leads to

alterations in mitochondrial morphology (**Figure S9B**). It is also noteworthy that

the expression of TbMfnL without the C-terminal 3Ty1 tag induced the same

phenotype as the tagged TbMfnL, confirming that the phenotype is not due to the

use of a C-terminal tag (**Figure S9A/B**). These data clearly demonstrated that

overexpression of TbMfnL stimulates inter-mitochondrial connections.

To characterize further inter-mitochondrial branching with higher resolution we

used serial block face scanning electron microscopy (SBF-SEM), an approach

allowing to characterize the overall structure of membranes and organelles within

the entire cell volume [18]. With this approach, hundreds of images are collected

to perform a 3D reconstruction of the whole cell, as well as spatial organization of

the individual organelles within the cell. Four to five parental and ^{oe}TbMfnL$_{::3Ty1}$ PCF

cells were reconstructed in 3D, with a particular emphasis on the mitochondrion,

the nucleus, the flagellar pocket and the cell outline. We have not been able to

correctly identify the kinetoplast DNA on the electron microscopy images.

However, since kDNA is in close contact with the flagellar pocket, we have assumed

its position. Other organelles (glycosomes, acidocalcisomes, Golgi, flagellum, etc.)

are not represented either in order to not overload the 3D representation. As

expected, the parental PCF cells showed a reticulated mitochondrion along the

whole cell (**Figure 4E**), a rather central nucleus and the flagellar pocket on the

posterior side. The length of the parental and ^{oe}TbMfnL$_{::3Ty1}$ cells, varied between

20.4±3.3 and 23±4.7 μm, respectively. As observed by fluorescence microscopy,

^{oe}TbMfnL$_{::3Ty1}$ mitochondrion showed more reticulation (**Figure 4E**). Unexpectedly,

in some cells this reticulation was even more prominent at the edges of the flagellar

pocket, where the kDNA is located, as illustrated in **Figure 4E**, an observation that

we had not identified by immunofluorescence microscopy approaches. This hyper-

reticulated structure (leaving a few gaps in the mitochondrion) formed a globular

structure and probably encapsulated the kDNA (**Figure 4E**). This structure called

kDNA pocket has been already described in BSF where kDNA replicates [20], but

is even more prominent in the ^{oe}TbMfnL$_{::3Ty1}$ cells. We determined that the

mitochondrial volume in PCF cells (4.9 ± 1 μm³, **Figure 4F**) aligns with previous

reports (2.5–3 μm³, [2]) and as expected, is nearly twice that of BSF cells (1.1–3

μm³, [18, 20]). As anticipated, the mitochondrial volume in $^{oe}TbMfnL_{::3Ty1}$ cells is

nearly doubled compared to parental cells (8.3 ± 1.9 µm³, **Figure 4F**). Additionally, TEM analysis of the mitochondrial and cristae structures in ^{oe}TbMfnL$_{::3Ty1}$ cells cultured with either proline or glucose as the sole carbon source revealed no significant alterations (**Figure S6**).

We also investigated the effect of TbMfnL expression in the BSF trypanosomes, where the mitochondrial compartment is smaller and composed of a shorter and poorly interconnected mitochondrion. Expression of ^{oe}TbMfnL$_{::3Ty1}$ was confirmed by western blot (**Figure 4G**), but unlike in PCF, the mitochondrial structure (labeled here with mitochondrial HSP60) was not altered, remaining tubular like in the parental cells (**Figure 4H**). Induction over a longer period (up to 21 days) did not show mitochondrial alterations.

5 - TbMfnL overexpression increases mitochondrial branching in a GTPase dependent manner

All dynamin-like proteins possess a GTP-binding domain with highly conserved G1-G4 motifs. They can be inactivated by mutations in the G1 motif that lower the efficiency of GTP-binding and hydrolysis. To determine whether TbMfnL activity requires a functional GTPase domain, the key lysine 141 in the highly conserved G1 motif was converted to alanine (K141A), according to previously characterized GTP-binding domains from other organisms [37-39]. The K141A mutation was introduced into the TbMfnL$_{::3Ty1}$ sequence, which was overexpressed in the PCF ^{oe}TbMfnL-K141A$_{::3Ty1}$ (**Figure 5A**) and TbMfnL$_{::3Ty1}$ PCF backgrounds (TbMfnL$_{::3Ty1}$/^{oe}TbMfnL-K141A$_{::3Ty1}$, **Figure S9**). Expression of TbMfnL-K141A$_{::3Ty1}$ was confirmed by western blot analyses after 3 to 5 days of tetracycline induction in the ^{oe}TbMfnL-K141A$_{::3Ty1}$ cell line (**Figure 5B**) and its overexpression was estimated as 3-fold in the TbMfnL$_{::3Ty1}$/^{oe}TbMfnL-K141A$_{::3Ty1}$ mutant compared to the parental TbMfnL$_{::3Ty1}$ clone (**Figure S9A**). TbMfnL-K141A$_{::3Ty1}$ localized to the mitochondria (**Figure 5C**), as expected, but did not alter mitochondrial shape (**Figure 5C/D, Figure S8**) nor the number of mitochondrial junctions (**Figure 4D**). To exclude the possibility that concentration-dependent effects (due to the twofold lower expression of TbMfnL-K141A$_{::3Ty1}$ compared to the parental TbMfnL$_{::3Ty1}$) might prevent K141A from inducing membrane branching, rather than its inactivated GTPase activity, we equalized the expression levels of wild-type TbMfnL$_{::3Ty1}$ and mutant TbMfnL-K141A$_{::3Ty1}$. The tetracycline concentration used for TbMfnL$_{::3Ty1}$ expression was reduced from 1 µg/ml to 0.05 µg/ml. Under

these conditions, both variants displayed comparable expression levels at 3 and 5
days (**Figure S10A**); however, hyper-reticulation was observed exclusively in cells
overexpressing TbMfnL$_{::3Ty1}$ (**Figure S10B**). This clearly demonstrates that TbMfnL
is a dynamin-related protein that relies on a functional GTPase domain to modulate
mitochondrial branching in trypanosomes.

The mitochondrial localization of ^{oe}TbMfnL$_{::3Ty1}$ and ^{oe}TbMfnL-K141A$_{::3Ty1}$ was also
confirmed by electron microscopy and immunogold-labeling using an anti-Ty1
antibody. After induction and immunogold-labeling, both proteins are clearly
localized in the mitochondrion (**Figure S11A/B-c/d**) as compared to controls with
non-induced cells (**Figure S11A/B-a/b**), and confirmed by gold-particles
quantification (**Figure S11C**).

6 - Submitochondrial localization of TbMfnL
Immunofluorescence analyses showed that ^{oe}TbMfnL$_{::3Ty1}$ co-localized with Tdh
(**Figure 6A**), as previously described for the endogenously-tagged TbMfnL$_{::10HA}$
(**Figure 2C**). To increase the resolution of TbMfnL localization we performed
Ultrastructure Expansion Microscopy (U-ExM), which increases the size of a sample
while preserving its ultrastructure [40-42]. Using this approach, we showed that
both ^{oe}TbMfnL$_{::3Ty1}$ (**Figure 6B**) and the endogenous TbMfnL$_{::10HA}$ (**Figure 6C**)
displayed mitochondrial localization with punctate, non-homogeneous distribution
patterns, which differed from the localization of the matrix marker (Tdh). These
findings suggest an association of the TbMfnL protein with mitochondrial
membranes.

7 - Mitochondrial targeting and membrane anchoring of TbMfnL
To investigate the mechanisms ensuring mitochondrial targeting of TbMfnL, we
first addressed the two domains that may determine TbMfnL localization: (i) the
N-terminal MTS which may facilitate targeting to the inner membrane,
intermembrane space and matrix localization, and (ii) the C-proximal/terminal TMs
which could mediate membrane anchoring. To investigate their capacity to target
proteins to mitochondria, we fused the MTS domain, or the two transmembrane
domains of TbMfnL, to GFP. Fusion of the MTS at the N-terminus of GFP ($_{MTS::}$GFP)
fully targeted GFP to the mitochondria (**Figure 7A**). Interestingly, western blot
analyses showed that $_{MTS::}$GFP and cGFP (cytosolic GFP) have the same apparent
size (**Figure 7B**), suggesting that the predicted MTS (**Figure S1**) was indeed

cleaved by matrix processing peptidase upon import of $_{MTS::}$GFP across the inner membrane. In the same line, ^{oe}TbMfnL$_{::3Ty1}$ and ^{oe}TbMfnL-K141A$_{::3Ty1}$ showed the same apparent size as ^{oe}TbMfnL-ΔMTS$_{::3Ty1}$ by western blot analyses, suggesting that the MTS is also cleaved in its natural context (**Figure 7C**). In contrast, western blot analyses showed that addition of the two C-proximal TMs at the C-terminal extremity of GFP (GFP$_{::TM}$) induces a size shift (**Figure 7B**). Immunofluorescence analyses revealed that GFP$_{::TM}$ was not targeted to the mitochondrion or cytosol, but instead to others distinct structures (**Figure 7A**). However, it is important to note that this localization could simply represent unspecific membrane-targeting by hydrophobic TMs. This inability of TbMfnL TMs to mediate mitochondrial targeting indicates that they differ from the TMs of other mitofusins, which are responsible for mitochondrial outer membrane localization [43].

To confirm the role and relevance of these domains on TbMfnL localization and function, we overexpressed 3Ty1-tagged TbMfnL-mutants without MTS (^{oe}TbMfnL-ΔMTS$_{::3Ty1}$) or without TMs (^{oe}TbMfnL-ΔTM$_{::3Ty1}$) in PCF trypanosomes (**Figure 7D and 7G**). Tetracycline-induced overexpression of both recombinant TbMfnL was confirmed by western blot, 3 to 5 days after induction (**Figure 7E/7H, Figure S9**). In contrast to ^{oe}TbMfnL$_{::3Ty1}$, ^{oe}TbMfnL-ΔMTS$_{::3Ty1}$ did not colocalize with Tdh revealing that the N-terminal MTS is indeed required for proper mitochondrial targeting of TbMfnL (**Figure 7F**). The MTS is therefore essential for the correct mitochondrial localization of TbMfnL. In addition, overexpression of ^{oe}TbMfnL-ΔMTS$_{::3Ty1}$ did not alter the mitochondrial structure, indicating that its proper, MTS-mediated, mitochondrial localization is essential for TbMfnL-function (**Figure 7F/Anti-Tdh, Figure S9**). Overexpression of a TbMfnL truncated version missing the C-proximal TMs (^{oe}TbMfnL-ΔTM$_{::3Ty1}$, **Figure 7G-H, Figure S9**) did not affect its mitochondrial localization (**Figure 7I**), which confirms that the N-terminal MTS sequence was sufficient for mitochondrial targeting of TbMfnL. However, in contrast to full length TbMfnL (**Figures 4C and 6A/B**), TbMfnL lacking TMs did not induce a modification of the mitochondrial shape (**Figure 7I, Figure S9**), demonstrating that the transmembrane domains are required for proper function. Altogether, our data showed that TbMfnL is targeted to the mitochondrion via an N-terminal MTS and that C-proximal TMs are required for the cellular function of TbMfnL.

Finally, we investigated the precise localization of $TbMfnL_{::3Ty1}$ and its variants ($TbMfnL-\Delta TM_{::3Ty1}$ and $TbMfnL-\Delta MTS_{::3Ty1}$) in the clone 2A11, where the endogenous $TbMfnL$ alleles were tagged with 3Ty1 (**Figure S5A-D**). This was achieved using a protease protection assay, a method previously used to determine the submitochondrial localization of various mitochondrial proteins in trypanosomes [44, 45]. Mitochondria are enriched by differential centrifugation, converted into mitoplasts by digitonin-mediated permeabilization of the outer membrane and finally incubated with proteinase K in the presence or absence of Triton X-100 to permeabilize the inner membrane.

Western blot analysis revealed that the cytosolic protein Enolase was almost entirely recovered in the supernatant, confirming that the generation of mitoplasts had effectively released cytosolic components (**Figure 8A, Figure S12**). In contrast, the $TbMfnL_{::3Ty1}$ protein and its variants exhibited fractionation patterns consistent with those of mitochondrial proteins, such as Atom40, Prohibitin, and Tdh, which serve as markers for the outer mitochondrial membrane (OMM), inner mitochondrial membrane (IMM, facing the intermembrane space), and mitochondrial matrix, respectively (**Figure 8A/C, Figure S12**). Proteinase K treatment of mitoplasts led to the near-complete digestion of outer membrane Atom40, intermembrane space Prohibitin (bound to the inner membrane), and $TbMfnL-\Delta MTS_{::3Ty1}$, which is no longer targeted to the mitochondrion and thus unprotected. In contrast, the size and levels of $TbMfnL_{::3Ty1}$, $TbMfnL-\Delta TM_{::3Ty1}$ and Tdh remained unaffected, indicating protease protection by the inner membrane. The intact state of Tdh further confirmed the integrity of the mitochondrial inner membrane and the reliability of the assay (**Figure 8A/C, Figure S12**). Digestion of the OMM-embedded β-barrel protein Atom40 and IMS Prohibitin, along with the preservation of $TbMfnL_{::3Ty1}$ and $TbMfnL-\Delta TM_{::3Ty1}$, suggests that $TbMfnL$ faces the matrix and is protected by the inner membrane. The addition of Triton X-100, which disrupts membrane integrity, exposed matrix proteins to proteinase K, resulting in the degradation of $TbMfnL_{::3Ty1}$, $TbMfnL-\Delta TM_{::3Ty1}$, and Tdh, thereby confirming their localization and orientation on the matrix side (**Figure 8A/C, Figure S12**). It should be noted that western blot revealed a size difference between the full-length $TbMfnL_{::3Ty1}$ and $TbMfnL-\Delta TM_{::3Ty1}$. Specifically, we observed two bands: the upper band corresponding to the endogenously tagged protein and the lower band representing to the overexpressed $TbMfnL-\Delta TM_{::3Ty1}$ protein.

To further investigate the localization of *Tb*MfnL, we separated the membrane and
matrix fractions by performing repeated freeze-thaw lysis on mitoplast
preparations. The membrane fraction was further divided into integral and
peripheral fractions via carbonate extraction. *Tb*MfnL$_{::3Ty1}$ and *Tb*MfnL-ΔTM$_{::3Ty1}$
were analyzed alongside control proteins, Tdh (matrix-localized) and Prohibitin
(inner membrane, IMS oriented). As expected, *Tb*MfnL$_{::3Ty1}$ was exclusively
retained in the membrane fraction, whereas *Tb*MfnL-ΔTM$_{::3Ty1}$, lacking its
transmembrane domains, was released from membrane (**Figure 8B/C, Figure**
S12). These data are consistent with a recent analysis of the mitochondrial
proteome, which proposed that the *Tb*MfnL protein is located in the inner
mitochondrial membrane [46].

Interestingly, structural predictions of *Tb*MfnL using Alphafold, supported by a very
high pLDDT score (**Figure S13**), suggest that the two previously identified
transmembrane domains form a loop. This loop may act as an anchor within the
inner membrane, thereby orientating the protein towards the matrix (**Figure 8C**),
consistent with the findings from proteinase K assay and carbonate extraction.

DISCUSSION

Membrane fusion and fission are often controlled by proteins of the dynamin superfamily (DSP) [47, 48]. These proteins consistently have a GTPase domain and contain domains required for membrane binding and oligomerization. In trypanosomes, only DSPs involved in fission have been characterized, and no fusion proteins similar to those of yeasts and mammals (Mfn and Opa1) have been identified so far. This may suggest that trypanosome mitochondria fuse via a machinery that differs from that characterized in yeasts and mammals. We have identified a novel DSP, named Mfn-Like (MfnL), which first representative characterized from the protozoan parasite *T. brucei* (*Tb*MfnL). We dit not establish a direct role in fusion, but demonstrated that it is able to increase mitochondrial interconnectivity in a GTPase-dependent manner, as previously reported for fusion factors Mfn/Opa1 [43, 49]. Interestingly, *Tb*MfnL domain organization and sub-mitochondrial localization differ significantly from that of Mfn and Opa1, suggesting that *Tb*MfnL and possibly other MfnL members mediate and/or enhance membrane branching and interconnection through a different molecular mechanism.

As observed for trypanosomes, many organisms including plants, lack genes encoding protein similar to known fusion factors. Interestingly, the search for homologous sequences to *Tb*MfnL allowed us to identify close DSP homologs in several eukaryotic phyla lacking Mfn and Opa1. These new DSPs are widely distributed in eukaryotes, which suggests that they represent an ancient protein family. Most of these phyla belong to the Diphoda, which together with the Opimoda are the two major supergroups of eukaryotes based on the most accepted position of the root of the tree of eukaryotes [50]. The only opimodan sequences found were the two amoeba cited above, most likely representing HGTs from bacteria, and the malawimonad *Gefionella okellyi* (**Figure 1C and Figure S3**). This malawimonad sequence may also reflect an HGT acquisition, in this case from a diphodan ancestor. The taxonomic distribution of this protein and its phylogeny support that it was ancestral at least in Diphoda. However, its presence also in bacteria, with any clear evidence of an HGT-mediated eukaryotic origin of the bacterial sequences, opens the possibility for an even more ancient origin before the diversification of Diphoda and Opimoda, which would imply a massive loss in opimodan taxa except in a few ones such as *Gefionella*. Future analysis of other deep-branching opimodan taxa will help to test this hypothesis. Homologs of *Tb*MfnL were also identified in several bacterial phyla, suggesting the existence in

these organisms of a novel machinery involved in membrane shaping and/or
remodeling that is different from the one found in yeasts and mammals. While
studies of yeast and mammalian DSPs provide information about their functions
and structures, little is known about bacterial DSPs, especially their cellular role.
Indeed, some bacterial DSPs have been associated with a variety of processes
involving membranes *in vivo*, such as a surveillance mechanism for membrane
punctures caused by antibiotics and bacteriophages [51, 52], membrane vesicle
formation [53] or cytokinesis by promoting membrane curvature at the septum
[54, 55]. In addition, studies of the *Bacillus subtilis* DSP DynA have shown that
this protein can promote membrane fusion *in vitro* [56]. The mechanism by which
the fusion operates is not yet known, but bacterial DSPs are probably recruited to
the sites where membrane fusion is needed [55, 57]. Interestingly, structural
analysis of *Campylobacter jejuni* DLP1/DLP2 proteins allowed to propose a
mechanism explaining how these proteins attach and bind distant and opposing
membranes [58]. Unfortunately, the cellular function of *C. jejuni* DLP1/DLP2 is
currently unknown. DSPs identified in this study have not yet been characterized
and no information on their localization and function is available. However, among
the identified organisms, we found bacteria of the phylum Planctomycetes, which
are unusual organisms. Indeed, these bacteria contain a compartmentalized
cytosol, separated by an intracytoplasmic membrane [59], in some cases
surrounding the DNA [60] or forming an anammoxosome, a membrane-bound
organelle responsible for the production of energy [61, 62]. It is then conceivable
that the identified protein in Planctomycetes (*Pb*DBF, **Figure 1 and S2**) is involved
in membrane structuring.
The role of *Tb*MfnL in mitochondrial shaping was observed by its overexpression in
PCF trypanosomes, which induced a very strong increase in number of connections
in the single branched mitochondrion. Moreover, the inactivation of its GTPase
domain via mutation of the G1 motif also confirms its belonging to the large family
of dynamins. Interestingly, a link between hyperconnection of a branching network
and increased fusion processes has been reported in a highly different biological
system, filamentous fungi, which grow as interconnected branching networks [63].
This raises the intriguing possibility that an excess of *Tb*MfnL in PCF trypanosomes
enhances the interconnectivity of mitochondrial networks by inducing
mitochondrial fusion.

The absence of fragmentation observed upon partial or total inactivation of *Tb*MfnL expression is in agreement with the absence or a very low level of constitutive mitochondrial fission. In contrast, in mammals and yeast, inhibition of fusion results in mitochondrial fragmentation by constitutive fission [16, 64]. Thus, our results are in agreement with previous studies suggesting that fission only occurs during cell division [19, 28]. The lack of observable mitochondrial alterations may also be attributed to the involvement of other proteins in mitochondrial structuring, as suggested by studies on the mitochondrial outer membrane proteome in *T. brucei* [65, 66] and the presence of mitochondrial complexes such as the recently characterized MICOS [44]. Additionally, subtle changes in mitochondrial structure may exist but could have gone undetected using our current approaches. It is worth noting that *Tb*MfnL seems to participate in a unique membrane-structuring mechanism, and its inactivation may not produce effects similar to those observed for Mfn or Opa1.

Another intriguing point was the discrepancy with the results reported by Vanwalleghem et al. [30]. Indeed, they describe fenestration of the mitochondrion in BSF cells with down-regulated *Tb*MfnL, whereas in our studies, involving inactivation by CRISPR/Cas9 or down-regulation by RNAi in BSF cells, we did not observe any alterations in mitochondrial structure. The only modification (hyper-reticulation) of the mitochondrial structure was observed exclusively during the overexpression of *Tb*MfnL and only in PCF forms. This discrepancy may arise from differences in cell lines, culturing conditions, sample preparation methods, or mitochondrial staining protocols. We used rhodamine staining on live cells and immunofluorescence labeling to visualize the mitochondrial network, whereas Vanwalleghem et al. [30] employed FIB-SEM tomography. It is also possible that *Tb*MfnL knockdown affects mitochondrial import, since disruptions in mitochondrial import have been reported to influence mitochondrial structure [65].

Using several microscopy approaches, we have demonstrated that *Tb*MfnL is localized within the mitochondrial membranes, with two domains involved in its accurate mitochondrial addressing, a canonical N-terminal MTS and two adjacent C-terminal transmembrane domains. Expression of MTS-less *Tb*MfnL or MTS-GFP recombinant proteins in PCF trypanosomes confirmed the role of this 41 aa-long MTS for mitochondrial import. Proteins with an N-terminal MTS are typically imported into the mitochondrial matrix, with a few exceptions, such as Opa1 and Apoptosis-inducing factor (AIF) face the intermembrane space [67]. Indeed, Opa1

and AIF localize mainly to the IMS due to their TM domains located in close proximity to the MTS (**Figure 1A**), interrupting transport across the inner membrane. The C-proximal localization of the transmembrane domains of *Tb*MfnL is not compatible with such import interruption, but more presumably suggested that its C-terminus is also exposed to the matrix, as predicted by Alphafold (**Figure 8C**). This model is consistent with the proteinase K resistance of *Tb*MfnL in mitoplasts. This orientation is also probably similar to that of the prokaryotic MfnLs identified in this study, which could also be anchored in the plasma membrane and orientated towards the cytosol. Interestingly, recent publications describe mitochondrial remodeling originating from the matrix side of the mitochondria. Sheikh *et al.* [68] demonstrated that expressing a potential DSP from a giant virus in *T. brucei* PCF significantly affects mitochondrial morphology within the matrix, with a notable association with the inner membrane. Additionally, Kumar *et al.* [69, 70] showed that a dynamin superfamily-like pseudoenzyme, a distant relative of the dynamin superfamily, stabilizes cristae architecture through interactions on the matrix side of the mitochondrial inner membrane. This contrasts with the mode of action of human or yeast mitofusins, which are involved in the remodeling of mitochondrial outer and inner membranes, but never via the matrix side.

In conclusion, the data presented here uncover a new family of DSPs and suggest the existence of a novel membrane-structuring mechanism found in both eukaryotes and prokaryotes, but distinct from those involved in mitochondrial membrane fusion in mammals and fungi.

ACKNOWLEDGMENTS AND FUNDING SOURCES

We thank Hassan Hashimi (University of South Bohemia) for providing us the anti-Prohibitin antibody, André Schneider (University of Bern) for providing us the anti-Atom40 antibody, Keith Gull (University of Manchester) for providing us the anti-Ty1 antibody and Derrick Robinson, Mélanie Bonhivers, Elina Casas and Nicolas Landrein (University of Bordeaux) for the pPOTv7 expression vector and extremely valuable help in the expansion microscopy experiment. Cell sorter analyses were performed at the TBMCore facility (FACSility) on BD FACSAria™ III Sorter and we thank Atika Zouine for data acquisition and interpretation. We also thanks Bordeaux Imaging Center (BIC), which is a member of the FranceBioImaging national infrastructure (ANR10-INBS-04), for helping us to design the ImageJ macro and microscopy acquisitions. The Bringaud team is supported by the Centre National de la Recherche Scientifique (CNRS, https://www.cnrs.fr/), the Université de Bordeaux (https://www.u-bordeaux.fr/) and the Agence Nationale de la Recherche (ANR, https://anr.fr/) through the ParaFrap "Laboratoire d'Excellence" (LabEx, https://www.enseignementsup-recherche.gouv.fr/cid51355/laboratoires-d-excellence.html) (ANR-11-LABX-0024). This work was supported by the "Fondation pour la Recherche Médicale" (FRM, https://www.frm.org/) ("Equipe FRM", grant n°EQU201903007845) and the ANR grant ADIPOTRYP (ANR19-CE15-0004-01) to FB and the ERC Advanced grant 787904 to D.M. Portions of this manuscript were based on Dr. Morel's thesis defended at the University of Bordeaux.

AUTHOR CONTRIBUTIONS

C.A.M. conceptualized the study, performed experiments, contributed to analysis and writing. C.A. performed experiments and contributed to analysis. D.M. performed all the phylogenic analysis. C.B., B.S. and E.G. performed the EM and SBF-SEM experiments and contributed to their analysis. S.D-C., M.R., F.B. and E.T. conceptualized the study, contributed to analysis and wrote the manuscript, with also an input from all the authors.

COMPETING INTEREST STATEMENT

The authors declare no competing interest.

 FIGURE LEGENDS

Figure 1 - Sequences homologous to *Tb*MfnL and domain organization of
the proteins. (A) Structure comparison between mitochondrial dynamin-
superfamily proteins (DSPs): *T. brucei Tb*MfnL, human mitofusin 1 or 2
(*Hs*Mfn1/2), yeast Fzo1 (*Sc*Fzo1) and human Opa1 (*Hs*Opa1). MTS, predicted
mitochondrial targeting sequence; CC, coiled coil region; green box, GTPase
domain with the four GTP-binding motifs in yellow; TM, transmembrane span; BSE,
bundle signaling element; Stalk domain; Paddle domain. Two TM domains were
initially proposed for Mfn1/2, but recently, Mattie et al. (2018) [71] reported the
presence of only a single TM domain, with the C-terminus extending into the IMS.
This contrasts with Fzo1 and *Tb*MfnL, which are proposed to have a U-shaped
structure. Therefore, we have indicated this as TM? in the figure. Structure
predictions were performed using PSIPRED [72]. More details and sequence
alignment are shown in **Figure S2**. **(B)** Domain organization of several identified
proteins from various organisms. MTS, predicted mitochondrial targeting
sequence; CC, coiled-coil region; green box, GTPase domain with the four GTP-
binding motifs in yellow; TM, transmembrane span; Structure predictions were
performed using PSIPRED. More details and sequence alignment are shown in
Figure S2. *Tb*MfnL, *Trypanosoma brucei* (ID: Tb927.7.2410); *El*MfnL, *Euglena*
longa (ID: EP00668_Euglena_longa_P029018); *Dp*MfnL, *Diplonema papillatum*
(ID: KAJ9468802.1); *Pg*MfnL, *Polarella glacialis* (ID: CAE8737761.1); *Ah*MfnL,
Achlya hypogyna (ID: OQR970001); *Ct*MfnL, *Chaetoceros tenuissimus* (ID:
GFH50037.1); *Hf*MfnL, *Hondaea fermentalgiana* (ID: GBG23909.1); *Cht*MfnL,
Chrysochromulina tobinii (ID: KOO35358.1); *Pb*MfnL, *Planctomycetaceae*
bacterium (ID: MBV8881356.1); *Ps*MfnL, *Pedosphaera* sp. (ID: MCH2381658.1);
*Rc*MfnL, *Rubrimonas cliftonensis* (ID: WP_093253540.1); *Gp*MfnL,
Gammaproteobacteria bacterium (ID: RTZ59088.1); *Mb*MfnL, *Myxococcales*
bacterium (ID: MCA9558847.1); *Ld*MfnL, *Lentisphaerae* bacterium (ID:
NLB69313.1); *Kb*MfnL, *Kiritimatiellae* bacterium (ID: MBO7223041.1). The
accession number for *Euglena longa* was obtained from Eukprot V3 web site. **(C)**
Maximum likelihood phylogenetic tree of eukaryotic and prokaryotic DSPs. The
tree was reconstructed using the LG+C60+G4 model of sequence evolution.
Ultrafast bootstrap support values are shown on each branch. Taxa are colored

according to their taxonomic affiliation. Groups of sequences belonging to the same phylum are collapsed (see **Figure S3** for the complete tree).

Figure 2 - Subcellular localization of *Tb*MfnL. (A) Schematic representation of *Tb*MfnL endogenously tagged at its C-terminus extremity with 10xHA. Only one of the two alleles is tagged. **(B)** Western blotting of whole-cell extracts ($5x10^6$ cells) of *T. brucei* PCF or BSF wild-type cells and PCF or BSF cells expressing *Tb*MfnL$_{::10HA}$ revealed with an anti-HA antibody. It should be noted that Enolase used as a loading control is about 3 to 4 times more expressed in BSF than in PCF according to Hannaert *et al*. 2003 [73]. **(C)** Subcellular localization of *Tb*MfnL$_{::10HA}$ in PCF and BSF. Colocalization of *Tb*MfnL$_{::10HA}$ (anti-HA antibody) with matrix mitochondrial threonine dehydrogenase for PCF (Anti-Tdh) and mitochondrial HSP60 for BSF (Anti-HSP60) analyzed by standard immunofluorescence. n, nucleus; k, kinetoplast. The scale bar represents 5 μm.

Figure 3 - Inactivation of the *Tb*MfnL gene. (A) Schematic representation of *Tb*MfnL inactivation by insertion of the puromycine resistant marker (PAC). **(B)** PCR confirmation of *Tb*MfnL gene inactivation on both alleles in BSF and PCF. **(C)** The mitochondrial shape of living parental (WT) and *Tb*MfnL$^{-/-}$ PCF cells (right panel) were designed by an ImageJ macro (Mitochondrial Junctions) from rhodamine 123 staining (left panel). **(D)** Antibody directed against HSP60 was used to label and visualized mitochondrial shape in BSF cells. Top panel: BSF parental cell (WT); bottom panel: BSF *Tb*MfnL$^{-/-}$ cell. n, nucleus; k, kinetoplast. **(E)** Corresponding mitochondrial junctions' quantification. Small red circles in the skeleton image indicate the junctions where three tubules converge. The scale bar represents 5 μm. Statistic: t test - Confidence interval 95% - P-value style: 0.1234 (ns); 0.0332 (*); 0.0021 (**); 0.0002 (***); <0.0001 (****).

Figure 4 - Overexpression of *Tb*MfnL and GTPase mutant. (A) Schematic representation of the overexpressing *Tb*MfnL protein tagged with 3Ty1 peptide at the C-terminus of the protein (^{oe}*Tb*MfnL$_{::3Ty1}$). **(B)** Western blotting of whole-cell extracts of *T. brucei* PCF cells overexpressing *Tb*MfnL$_{::3Ty1}$. Non-induced (-Tet) and induced (+Tet) cells after 3 to 5 days (d3 and d5) were revealed with an anti-Ty1 antibody. Antibody directed against Enolase protein was used as loading control. **(C)** Mitochondrial structure analysis using rhodamine 123 staining on living cells;

mitochondrial structure of PCF parental (WT) and PCF ^{oe}TbMfnL$_{::3Ty1}$ cells before (-Tet) and 5 days after (+Tet) tetracycline induction. **(D)** Quantification of mitochondrial junctions' number using Rhodamine 123 microscopy images. Mitochondrial junctions number: WT, 9.4 ± 4.3; -Tet ^{oe}TbMfnL$_{::3Ty1}$ 8.7 ± 3.6; +Tet ^{oe}TbMfnL$_{::3Ty1}$ 20.4 ± 8.9; -Tet ^{oe}TbMfnL-K141A$_{::3Ty1}$ 8.6 ± 4.8 and +Tet ^{oe}TbMfnL-K141A$_{::3Ty1}$ 10 ± 5.2. **(E)** Surface rendering of segmented PCF using SBF-SEM. Surface rendering of segmented PCF WT and ^{oe}TbMfnL cells, with ^{e}TbMfnL induced by tetracycline for 5 days, showing the cell body (transparent), nucleus (blue), mitochondrion (green), and flagellar pocket (red). 5 WT and 4 ^{oe}TbMfnL cells were analyzed. A magnified view of the upper images, enclosed in a box, was shown. Small white arrows mark the mitochondria-free areas within the mitochondrial network. **(F)** Determination of mitochondrial volume (µm^3) in both WT and ^{oe}TbMfnL cells; WT 4.9±1 and ^{oe}TbMfnL 8.3±1.9. Statistic: t test - Confidence interval 95% - P-value style: 0.1234 (ns); 0.0332 (*); 0.0021 (**); 0.0002 (***); <0.0001 (****). **(G)** Western blotting of whole-cell extracts of *T. brucei* BSF overexpressing TbMfnL$_{::3Ty1}$ cells. Non-induced (-Tet) and induced (+Tet) cells after 3 to 7 days (d3, d5 and d7) were revealed with an anti-Ty1 antibody. Antibody directed against Enolase protein was used as loading control. **(H)** Subcellular localization of ^{oe}TbMfnL$_{::3Ty1}$ in BSF by immunofluorescence. Antibody directed against HSP60 was used to label and visualized mitochondrial shape in parental (WT) and TbMfnL$_{::3Ty1}$ overexpressing cells. n, nucleus; k, kinetoplast. The scale bar represents 5 µm.

Figure 5 - Overexpression of GTPase mutant(A) Schematic representation of the overexpressing TbMfnL-K141A$_{::3Ty1}$ mutant protein tagged with 3Ty1 peptide at the C-terminus of the protein (^{oe}TbMfnL-K141A$_{::3Ty1}$). **(B)** Western blotting of whole-cell extracts of *T. brucei* PCF overexpressing TbMfnL-K141A$_{::3Ty1}$ cells; (-Tet) non-induced and (+Tet) induced cells for 3 or 5 days (d3 and d5). Antibody directed against glycosomal Fumarate reductase protein (Frdg) was used as loading control. **(C)** Submitochondrial localization of ^{oe}TbMfnL-K141A$_{::3Ty1}$ in PCF by immunofluorescence; Colocalization of ^{oe}TbMfnL-K141A$_{::3Ty1}$ (anti-Ty1) with matrix mitochondrial Threonine dehydrogenase (Anti-Tdh) was analyzed by standard immunofluorescence, after 5 days induction (d5+Tet). n, nucleus; k, kinetoplast. **(D)** Mitochondrial structure analysis using rhodamine 123 staining on living cells; mitochondrial structure of PCF parental (WT) and PCF ^{oe}TbMfnL-

K141A$_{::3Ty1}$ cells before (-Tet) and 5 days after (+Tet) tetracycline induction. The

scale bar represents 5 µm.

Figure 6 – Submitochondrial localization of *Tb*MfnL. Subcellular localization

of ^{oe}*Tb*MfnL$_{::3Ty1}$ in PCF by immunofluorescence and Ultra Expansion Microscopy

(UExM); Colocalization of ^{oe}*Tb*MfnL$_{::3Ty1}$ (anti-Ty1) with matrix mitochondrial

Threonine dehydrogenase (Anti-Tdh) was analyzed by standard

immunofluorescence **(A)** and by UExM with an expansion factor of ~4.4 **(B)**, after

5 days induction (d5+Tet). **(C)** Subcellular localization of ^{oe}*Tb*MfnL$_{::10HA}$ in PCF

analyzed by UExM. n, nucleus; k, kinetoplast. The scale bar represents 5 µm.

Figure 7 - Role of *Tb*MfnL domains. (A) Localization of the GFP according to

the different tags fused. Colocalization of GFP with the mitochondrial matrix

marker threonine dehydrogenase (Anti-Tdh) was assessed by standard

immunofluorescence after 5 days of tetracycline induction for MTS$_{::GFP}$ and GFP$_{::TM}$.

The "c" in cGFP stands for constitutive expression, resulting in the cytosolic

localization of GFP. MTS, mitochondrial targeting signal; TM, transmembrane

domains. **(B)** Western blot analysis of GFP expression. cGFP, constitutive and

cytosolic GFP. Non-induced (-Tet) and induced (+Tet) cells after 5 days. **(C)**

*Tb*MfnL size analysis by western blot using anti-Ty1 antibody. **(D)** Schematic

representation of N-terminus truncated *Tb*MfnL protein (first 41 amino acids)

tagged with 3Ty1 peptide at the C-terminus of the protein (^{oe}*Tb*MfnL-ΔMTS$_{::3Ty1}$).

(E) Western blotting of whole-cell extracts of *T. brucei* PCF overexpressing

*Tb*MfnL-ΔMTS$_{::3Ty1}$ cells; (-Tet) non-induced and (+Tet) induced cells for 3 or 5

days. **(F)** Subcellular localization of ^{oe}*Tb*MfnL-ΔMTS$_{::3Ty1}$ in PCF after 5 days

induction. Colocalization of ^{oe}*Tb*MfnL-ΔMTS$_{::3Ty1}$ (anti-Ty1) with matrix

mitochondrial Threonine dehydrogenase (Anti-Tdh) analyzed by standard

immunofluorescence. **(G)** Schematic representation of truncated transmembrane

domains of *Tb*MfnL protein (45 amino acids) tagged with 3Ty1 peptide at the C-

terminus of the protein (^{oe}*Tb*MfnL-ΔTM$_{::3Ty1}$). **(H)** Western blotting of whole-cell

extracts of *T. brucei* PCF overexpressing *Tb*MfnL-ΔTM$_{::3Ty1}$ cells; (-Tet) non-induced

and (+Tet) induced cells for 3 days. **(I)** Subcellular localization of ^{oe}*Tb*MfnL-

ΔTM$_{::3Ty1}$ in PCF after 3 days induction, analyzed by standard immunofluorescence.

n, nucleus; k, kinetoplast. The scale bar represents 5 µm.

Figure 8 – Membrane topology of *Tb*MfnL. (A) The Proteinase K protection
assay involved incubating mitoplasts with Proteinase K and Triton X100, as
denoted by the "+" symbol. TF represents the total fraction containing both
cytoplasmic and membrane components; S denotes the supernatant, comprising
the cytosolic contents; and Pellet represents the organelle fraction. **(B)** Carbonate
extraction of mitochondria isolated from ^{oe}*Tb*MfnL$_{::3Ty1}$ and ^{oe}*Tb*MfnL-ΔTM$_{::3Ty1}$ cells.
Mem, membrane fraction; Matrix, matrix fraction; Int. Mem, integral membrane
fraction; Per. Mem, peripheral membrane fraction. **(C)** Predicted structure of
*Tb*MfnL, from AlphaFold protein structure database. *Tb*MfnL was embedded in
mitochondrial inner membrane (IMM). The color coding corresponds to the
predicted secondary structure depicted on the top of the figure. The first 41 N-
terminal amino acids corresponding to the MTS are not shown as they are cleaved.
The N- and C-termini are marked by small black spheres. The AlphaFold model
score of *Tb*MfnL is between confident to very high for most of the structured
domains (https://alpha- fold.ebi.ac.uk/entry/Q57XN3, **Figure S13**).

MATERIALS AND METHODS

Trypanosomes and cell cultures

The PCF of *T. brucei* EATRO1125.T7T (TetR-HYG-T7RNAPOL-NEO, where TetR stands for tetracycline resistance, HYG for hygromycin, T7RNAPOL for RNA polymerase T7, and NEO for neomycin) was cultured at 27°C with 5% CO_2 in SDM79 medium containing 10% (vol/vol) heat-inactivated fetal calf serum, 5 µg/ml hemin, 25 µg/ml hygromycin and 10 µg/mL neomycin. PCF cells were also cultured in SDM80 medium containing either glucose (10 mM) as the sole carbon source or proline (5 mM) supplemented with N-acetyl-glucosamine (50 mM) to inhibit the glucose transporter. The bloodstream form of *T. brucei* 427 90-13 (TetR-HYG-T7RNAPOLNEO) was cultured at 37°C with 5% CO_2 in Iscove's modified Dulbecco's medium (IMDM) supplemented with 10% (vol/vol) heat-inactivated fetal calf serum, 0.2 mM β-mercaptoethanol, 36 mM $NaHCO_3$, 1 mM hypoxanthine, 0.16 mM thymidine, 1 mM sodium pyruvate, 0.05 mM bathocuproine, 1.5 mM L-cysteine [74], 5 µg/ml hygromycin and 2.5 µg/mL neomycin. Overexpression cell lines were induced with tetracycline (10 µg/mL for BSF and 1 µg/mL for PCF). Growth was monitored by daily cell counting with the cytometer Guava® Muse® and Guava® easyCyte™.

Endogenous tagging and inactivation of *Tb*MfnL by CRISPR-Cas9

Endogenous tagging and inactivation of *Tb*MfnL were achieved by CRISPR Cas9 technology [36]. Briefly, inactivation of the *Tb*MfnL was achieved by inserting the resistant marker puromycin (Pac) or a small sequence encoding for a *Bam*HI restriction site plus 6 successive stop codons, flanked by 50 bp homologous to the 5' and 3' *Tb*MfnL sequences from the Cas9 cutting site. Similar approaches have been performed for endogenous tagging with the insertion of a sequence encoding 3xTy1 or 3xHA. The EATRO1125.T7T PCF and the 427 90.13 BSF ($1x10^6$ cells) were respectively transfected, using Amaxa nucleofectorII, with 1 µg of purified cassette (puromycin resistance marker, Stop*Bam*H1Stop, 3xTy1 or 3xHA), 30 µg Cas9 protein from IDT and pre-annealing TracrRNA (0.4 µmol) and gRNA (0.4 µmol). Cells were transfected using program X-001 and selected or not with puromycin (SDM79, 1 µg/mL, or IMDM 0.1 µg)/mL). Cells were cloned by using a cell sorter (TBM Core facility), and selection of double inactivated *Tb*MfnL gene (*Tb*MfnL$^{-/-}$) or endogenously tagged clones was done by DNA extraction, with NucleoSpin Blood (Macherey-Nagel) and PCR amplification, see supplemental

Table S2. Guide RNAs were designed using EuPaGDT, from http://tritrypdb.org.
Primers and guide RNAs used were synthetized by Integrated DNA Technologies
(IDT) and are listed in supplemental **Table S2**.

Immunofluorescence

Cells were washed twice with PBS, then fixed with 2% paraformaldehyde (PFA) for
10 min at room temperature and 0.1mM glycine was added 10 min to stop the
reaction. The cells were spread on slides and permeabilized with 0.05% triton X-
100. After incubation in PBS containing 4% bovine serum albumin (BSA) 20 min,
cells were incubated for 1 hour with primary antibodies diluted in PBS-BSA 4%,
washed 4 times with PBS and incubated for 45 min with secondary antibodies
diluted in PBS-BSA 4% followed by three washes. Then, kinetoplasts and nuclei
were labelled with DAPI (10 µg/mL) for 5 min. Slides were washed 3 times with
PBS and mounted with SlowFade Gold (Molecular probes). Images were acquired
with MetaMorph software on Zeiss Imager Z1 or an Axioplan 2 microscope and
processed with ImageJ.

Mitochondria staining in live cells

Rhodamine-123 (30 µg/mL) was added on cell culture (5×10^6 - 1×10^7 cells per mL)
for 15 min at room temperature, then cells were washed twice with PBS and spread
on slides. As an alternative to labeling mitochondria in living cells with Rhodamine-
123, Mitotracker (MitoTracker Green FM) was used. Cells were washed once with
PBS, incubated for 10 minutes with 10 µM MitoTracker Green FM, washed again
with PBS, and then spread onto slides. Images were acquired with MetaMorph
software on an Axioplan 2 microscope and processed with ImageJ.

Ultrastructure Expansion Microscopy (UExM)

The protocol of UExM was realized exactly as described by Casas et *al.* 2022
(dx.doi.org/10.17504/protocols.io.bvwqn7dw). An expansion factor was
determined using the ratio between the size of the coverslip (12 mm) and the size
of the gels after the first expansion. Images were acquired on a confocal Leica
SP5-MP (Bordeaux Imaging Center) using a 63X oil objective and processed with
ImageJ.

Quantification of Mitochondrial junctions - ImageJ macro

To quantify the number of mitochondrial junctions, an ImageJ macro was
developed with the help of the Bordeaux Imaging Center (BIC). The script is
presented in supplemental (**Data S1**).

Endogenous tagging and overexpression of *Tb*MfnL

For endogenous gene tagging, primers were designed as described in [75] and PCR was performed using pPOTv7 vector as template. The pPOT used here was pPOTv7 for C-terminus 10xHA tagging (blasticidin resistance). For overexpression, *MfnL* gene (Tb427.7.2410) was inserted in both pLew100 (phleomycine resistance) [76] and pHD1336 (blasticidin resistance) [77] expression vectors. Fragments were amplified and cloned into *Hin*dIII and *Xba*I restrictions sites of the pLew100 and into *Hin*dIII and *Bam*HI restrictions sites of the pHD1336 expression vectors. In addition, a 3xTy1 tag was added at the C-terminus of the protein in the pLew100-*Tb*MfnL, using the *Xba*I and *Bam*HI restrictions sites. The truncated version of *Tb*MfnL without the first 41 amino acids (*Tb*MfnL-DMTS) and without the two trans-membranes domain (*Tb*MfnL-DTM) were amplified and cloned in the pLew100 with a 3xTy1 tag as previously described. Catalytic lysine (K141) was replaced by an alanine, using PCR approach [78] and PCR product cloned in the pLew100 with a 3xTy1 tag (*Tb*MfnL-K141A). EATRO1125.T7T PCF was transfected using Amaxa Nucleofector®II, program X-001 and selected in SDM79 containing 25 µg/ml hygromycin 10 µg/mL neomycin and 5 µg/mL phleomycin. 427 90-13 BSF was transfected using the same conditions and selected in IMDM containing 5 µg/ml hygromycin, 2.5 µg/mL neomycin and 5 µg/mL blasticidin. Primers used for the constructions are presented in supplemental **Table S2**.

Down-regulation of *Tb*MfnL gene expression

Down-regulation of *Tb*MfnL expression by RNAi in PCF was achieved by expression of stemloop "sense/antisense" RNA molecules targeting a 400-bp fragment of the *Tb*MfnL gene introduced into the pLew100 tetracycline-inducible expression vector. A PCR-amplified 450-bp fragment, containing the antisense *Tb*MfnL sequence was inserted between *Xho*I and *Bam*HI restriction sites of the pLew100 plasmid. Then, the separate 400-bp PCR-amplified fragment containing the sense *Tb*MfnL sequence was inserted upstream of the antisense sequence, using *Hin*dIII and *Xho*I restriction sites. The resulting plasmid, pLew100-*Tb*MfnL-SAS, contains a sense and antisense version of the *Tb*MfnL fragment separated by a 50-bp fragment. The EATRO1125 PCF was transfected with the pLew100-*Tb*MfnL-SAS and cells were selected in SDM79 medium containing 25 µg/ml hygromycin, 10 µg/mL neomycin and 5 µg/mL phleomycin. Expression of the RNAi was induced by tetracycline (1 µg/mL). Down-regulation of *Tb*MfnL expression by RNAi in BSF was achieved exactly as described [30] with a 280-bp fragment derived from the *Tb*MfnL open reading frame inserted into the p2T7-177 plasmid [79]. Linearized

891 plasmid was transfected in *T. brucei* 427 90-13 (TetR-HYG-T7RNAPOL-NEO) BSF

892 cells. Expression of the RNAi was induced by tetracycline (10 μg/mL). Primers used

893 for the constructions are presented in supplemental **Table S2**.

Serial Block-Face Scanning Electron Microscopy (SBF-SEM)

895 The entire procedure is detailed and has been carried out as described in the

896 article: Advancing yeast cell analysis: A cryomethod for serial block-face scanning

897 electron microscopy imaging in mitochondrial morphology studies [80].

Immunoelectron microscopy

Harvested cells were placed on the surface of formvar-coated copper grids (400

mesh). Each loop was quickly submersed in liquid propane (−180°C) and

transferred into a precooled solution of 0.1% uranyl acetate in dry acetone for 3

days at -82°C. Samples were rinsed with acetone at -20°C, and embedded

progressively at -20°C in LR Gold resin (EMS, USA). Resin polymerization was

carried out at -20°C for 7 days under UV illumination. Ultrathin LR Gold sections

were collected on nickel grids coated with formvar. Sections were first incubated

for 15 min with 500mM NH_4Cl in Tris-buffered saline (TBS) pH7.8, blocked 2x10

min with 2% BSA in TBS pH 7.8. The grids were incubated 1 hour at room

temperature with anti Ty1 antibody diluted to 1:200 in TBS containing 2% BSA

rinsed with TBS containing 2% BSA and with TBS containing 1% BSA. The samples

were then incubated for 1 hour at room temperature with anti-mouse IgG diluted

to 1:20 conjugated to 10 nm gold particles (BioCell). The sections were rinsed with

TBS containing 1% BSA, and fix with 1% glutaraldehyde in TBS. After rinsing with

TBS, grids were contrasted through a 5 min incubation with 2% uranyl acetate in

water, followed by 1 min incubation with 1% lead citrate. Observations were

performed on a HITACHI H7650 transmission electron microscope operated at 80

KV with an Orius 1000-11 MPx camera (Gatan, Abingdon, UK).

Transmission electron microscopy (TEM)

918 The entire procedure is thoroughly detailed and has been carried out precisely as

919 described in the article: Plunge Freezing: A Tool for the Ultrastructural and

920 Immunolocalization Studies of Suspension Cells in Transmission Electron

921 Microscopy [81].

Western blot analyses

923 Total protein extracts ($5x10^6$ cells) were separated by SDS-PAGE (10%) and

924 immunoblotted on TransBlot Turbo Midi-size PVDF Membranes (Bio-Rad).

925 Immunodetection was performed using the primary antibodies, diluted in PBS-

Tween Milk (0.05% Tween20, 5% skimmed milk powder), summarized in supplemental **Table S3**. Revelation was performed using a second antibody coupled to the horseradish peroxidase and the Clarity western enhanced-chemiluminescence (ECL) substrate as describes by the manufacturer (Bio-Rad). Images were acquired and analyzed with the ImageQuant Las 4000 luminescent image analyzer.

Mitochondria-enriched preparation, proteinase K protection assays and carbonate extraction

This procedure has been well established and allows the determination of the submitochondrial localization of trypanosome proteins [44, 45, 82]. 10^8 cells were lysed in SoTE buffer (20 mM Tris-HCl, pH 7.5, 0.6 mL sorbitol, and 2 mM ethylenediaminetetraacetic acid), supplemented with 0.02% (w/v) digitonin (D141-100MG, Sigma-Aldrich) for 5 min at 4°C. A portion of the resulting suspension was collected as the total fraction (TF), while the remaining suspension underwent centrifugation at 6,000 x g for 3 minutes at 4°C to isolate a mitochondria-enriched pellet. This pellet was then resuspended in SoTE supplemented with 0.2% (w/v) digitonin and incubated on ice for 15 minutes. The resuspended mixture was divided into three tubes and subjected to centrifugation at 6,000 × g for 3 minutes at 4°C, resulting in the separation of mitoplasts and soluble protein in the supernatant (S). Each mitoplast pellet was resuspended in SoTE and subjected to the proteinase K protection assay. Proteinase K (50 µg) was added to two tubes, one supplemented with 1% Triton-X100 (v/v), while the third tube remained untreated as a control. All tubes were then incubated on ice for 30 minutes, followed by addition of 5 mM phenylmethylsulfonyl fluoride to stop the reaction. Proteins from all fractions were precipitated with trichloroacetic acid before undergoing resolution by SDS-PAGE.

For the extraction of integral membrane proteins using the carbonate method, the procedure begins with a mitochondria-enriched pellet. The pellet is resuspended in 500 µL of 10 mM $MgCl_2$ and subjected to a freeze-thaw cycle repeated ten times, involving freezing in liquid nitrogen and thawing at room temperature. Subsequently, the suspension is divided into two fractions: 50 µL and 450 µL. The 450 µL fraction is centrifuged at 10,000 × g for 5 minutes at 4 °C, with the supernatant designated as the matrix fraction. The 50 µL fraction is centrifuged under the same conditions, and the resulting pellet is resuspended in 50 µL of 10 mM $MgCl_2$, representing the membrane fraction. The remaining pellet from the 450

μL sample is further processed by resuspending it in 400 μL of 0.1 M Na$_2$CO$_3$ (pH 11). This mixture is incubated on ice for 30 minutes, sonicated four times, and centrifuged at 20,000 × g for 20 minutes at 4 °C. The supernatant is collected as the peripheral membrane protein fraction. The pellet is washed by resuspension in 400 μL of 0.1 M Na$_2$CO$_3$ and subjected to another centrifugation under the same conditions. This final pellet constitutes the integral membrane protein fraction. All protein fractions are precipitated using trichloroacetic acid, resuspended in 50 to 100 μL of 2% SDS, and analyzed by SDS–PAGE.

Sequence database searches and phylogenetic analysis

Two databases were used to identify sequences homologous to the *T. brucei* *Tb*MfnL sequence (Tb927.7.2410). The first search was conducted using Blastp [83] (protein-protein BLAST, https://blast.ncbi.nlm.nih.gov/Blast.cgi) on the NCBI nr protein database, with *Tb*MfnL as the query and kinetoplastids excluded from the search. The parameters included a maximum of 5000 target sequences and the Blosum62 matrix. For the EuKProt v3 database [84] search (https://evocellbio.com/eukprot/), the Blastp server was used with the entire database (993 species) and *Tb*MfnL as the query, without applying advanced parameters. The selection of sequences was then performed manually, retaining only those sequences that exhibited homology across the entire *Tb*MfnL protein. Sequences corresponding solely to the highly conserved GTPase domain were excluded. A total of 315 sequences were selected for phylogenetic analysis. An additional 23 sequences were included from well-characterized dynamins involved in fission (Dnm1), fusion (Opa1, Mfn1/2, Fzo1), BDLP (the cyanobacterial Mfn homolog), and dynamins associated with clathrin-mediated endocytosis (Dyn1/2) to be used as an outgroup.

The sequences were aligned with MAFFT L-INS-i v7.450 [85] and trimmed with BMGE v1.12 (-m BLOSUM30 -b 3 -g 0.2 -h 0.5) [86]. A maximum likelihood phylogenetic tree was then reconstructed with IQ-TREE v.2.0.3 [87] using its model finder (-m TEST) option, including complex mixture models, to select the best fit substitution model, as well as 1000 ultrafast bootstrap replicates to estimate branch statistical support.

Structural prediction software and sequence analysis

Structure predictions were performed using PSIPRED ("http://bioinf.cs.ucl.ac.uk/psipred/") [72] and prediction of mitochondrial

targeting sequences by various algorithms (MitoFates, PSORT II and TargetP-2.0) [88].

Statistical analysis

Experiments were performed at least in triplicates. Statistical analyses were performed using Prism (GraphPad) software. The results are presented as mean ± S.D. Where indicated the results were subjected to two-sided Student's t-test to determine statistical differences against the indicated group (Confidence interval 95% - P-value style: 0.1234 (ns); 0.0332 (*); 0.0021 (**); 0.0002 (***); <0.0001 (****)).

SUPPLEMENTAL INFORMATION

Figure S1: Alignment of various kinetoplastid MfnL proteins.

Figure S2: Alignment between *Tb*MfnL and various eukaryotic and prokaryotic MfnLs.

Figure S3: Complete maximum likelihood phylogenetic tree of eukaryotic and prokaryotic DSPs.

Figure S4: Growth curves of parental, over-expressing, and mutant cells.

Figure S5: Inactivation of *Tb*MfnL expression using the CRISPR/Cas9 system.

Figure S6: TEM analysis of cells cultured on glucose or proline.

Figure S7: Down-regulation of *Tb*MfnL expression by RNAi.

Figure S8: Results of the ImageJ plugin applied to rhodamine-stained cells.

Figure S9: Overexpression levels of different *Tb*MfnL variants.

Figure S10: Variation in *Tb*MfnL expression levels upon tetracycline induction.

Figure S11: Mitochondrial localization of *Tb*MfnL as shown by immunoelectron microscopy.

Figure S12: Full gel results shown for the PK and carbonate assays.

Figure S13: pLDDT score for *Tb*MfnL protein derived from AlphaFold predictions.

Data S1: ImageJ code used in the study.

Table S1: BLASTp analysis results for several identified organisms.

Table S2: List of all primers and guide RNAs utilized in this study.

Table S3: List of all antibodies used in this study.

1022

REFERENCES

1. Bohringer, S., and Hecker, H. (1975). Quantitative ultrastructural investigations of the life cycle of Trypanosoma brucei: a morphometric analysis. J Protozool *22*, 463-467.

2. Bily, T., Sheikh, S., Mallet, A., Bastin, P., Perez-Morga, D., Lukes, J., and Hashimi, H. (2021). Ultrastructural Changes of the Mitochondrion During the Life Cycle of Trypanosoma brucei. The Journal of eukaryotic microbiology, e12846.

3. Priest, J.W., and Hajduk, S.L. (1994). Developmental regulation of Trypanosoma brucei cytochrome c reductase during bloodstream to procyclic differentiation. Mol Biochem Parasitol *65*, 291-304.

4. van Hellemond, J.J., Opperdoes, F.R., and Tielens, A.G. (2005). The extraordinary mitochondrion and unusual citric acid cycle in Trypanosoma brucei. Biochem Soc Trans *33*, 967-971.

5. Michels, P.A., Bringaud, F., Herman, M., and Hannaert, V. (2006). Metabolic functions of glycosomes in trypanosomatids. Biochim Biophys Acta *1763*, 1463-1477.

6. Bringaud, F., Riviere, L., and Coustou, V. (2006). Energy metabolism of trypanosomatids: adaptation to available carbon sources. Mol Biochem Parasitol *149*, 1-9.

7. Duvezin-Caubet, S., Jagasia, R., Wagener, J., Hofmann, S., Trifunovic, A., Hansson, A., Chomyn, A., Bauer, M.F., Attardi, G., Larsson, N.G., et al. (2006). Proteolytic processing of OPA1 links mitochondrial dysfunction to alterations in mitochondrial morphology. J Biol Chem *281*, 37972-37979.

8. Sauvanet, C., Duvezin-Caubet, S., di Rago, J.P., and Rojo, M. (2010). Energetic requirements and bioenergetic modulation of mitochondrial morphology and dynamics. Semin Cell Dev Biol *21*, 558-565.

9. Silva Ramos, E., Larsson, N.G., and Mourier, A. (2016). Bioenergetic roles of mitochondrial fusion. Biochim Biophys Acta *1857*, 1277-1283.

10. Chan, D.C. (2012). Fusion and fission: interlinked processes critical for mitochondrial health. Annual review of genetics *46*, 265-287.

11. Westermann, B. (2010). Mitochondrial fusion and fission in cell life and death. Nat Rev Mol Cell Biol *11*, 872-884.

12. Sesaki, H., Adachi, Y., Kageyama, Y., Itoh, K., and Iijima, M. (2014). In vivo
functions of Drp1: lessons learned from yeast genetics and mouse
knockouts. Biochim Biophys Acta *1842*, 1179-1185.

13. Otera, H., Ishihara, N., and Mihara, K. (2013). New insights into the function
and regulation of mitochondrial fission. Biochim Biophys Acta *1833*, 1256-
1268.

14. Malka, F., Guillery, O., Cifuentes-Diaz, C., Guillou, E., Belenguer, P.,
Lombes, A., and Rojo, M. (2005). Separate fusion of outer and inner
mitochondrial membranes. EMBO Rep *6*, 853-859.

15. Song, Z., Ghochani, M., McCaffery, J.M., Frey, T.G., and Chan, D.C. (2009).
Mitofusins and OPA1 mediate sequential steps in mitochondrial membrane
fusion. Mol Biol Cell *20*, 3525-3532.

16. Legros, F., Lombes, A., Frachon, P., and Rojo, M. (2002). Mitochondrial
fusion in human cells is efficient, requires the inner membrane potential,
and is mediated by mitofusins. Mol Biol Cell *13*, 4343-4354.

17. Bertholet, A.M., Delerue, T., Millet, A.M., Moulis, M.F., David, C., Daloyau,
M., Arnaune-Pelloquin, L., Davezac, N., Mils, V., Miquel, M.C., et al. (2016).
Mitochondrial fusion/fission dynamics in neurodegeneration and neuronal
plasticity. Neurobiol Dis *90*, 3-19.

18. Hughes, L., Borrett, S., Towers, K., Starborg, T., and Vaughan, S. (2017).
Patterns of organelle ontogeny through a cell cycle revealed by whole-cell
reconstructions using 3D electron microscopy. J Cell Sci *130*, 637-647.

19. Chanez, A.L., Hehl, A.B., Engstler, M., and Schneider, A. (2006). Ablation
of the single dynamin of T. brucei blocks mitochondrial fission and
endocytosis and leads to a precise cytokinesis arrest. J Cell Sci *119*, 2968-
2974.

20. Jakob, M., Hoffmann, A., Amodeo, S., Peitsch, C., Zuber, B., and
Ochsenreiter, T. (2016). Mitochondrial growth during the cell cycle of
Trypanosoma brucei bloodstream forms. Sci Rep *6*, 36565.

21. Esseiva, A.C., Chanez, A.L., Bochud-Allemann, N., Martinou, J.C., Hemphill,
A., and Schneider, A. (2004). Temporal dissection of Bax-induced events
leading to fission of the single mitochondrion in Trypanosoma brucei. EMBO
Rep *5*, 268-273.

22.   Gibson, W., Crow, M., and Kearns, J. (1997). Kinetoplast DNA minicircles
1090   are inherited from both parents in genetic crosses of Trypanosoma brucei.
1091   Parasitol Res *83*, 483-488.

1092   23.   DiMaio, J., Ruthel, G., Cannon, J.J., Malfara, M.F., and Povelones, M.L.
1093   (2018). The single mitochondrion of the kinetoplastid parasite Crithidia
1094   fasciculata is a dynamic network. PLoS ONE *13*, e0202711.

1095   24.   van der Bliek, A.M., Shen, Q., and Kawajiri, S. (2013). Mechanisms of
1096   mitochondrial fission and fusion. Cold Spring Harb Perspect Biol *5*.

1097   25.   Logan, D.C. (2006). Plant mitochondrial dynamics. Biochim Biophys Acta
1098   *1763*, 430-441.

1099   26.   Arimura, S.I. (2018). Fission and Fusion of Plant Mitochondria, and Genome
1100   Maintenance. Plant Physiol *176*, 152-161.

1101   27.   Gao, H., Sage, T.L., and Osteryoung, K.W. (2006). FZL, an FZO-like protein
1102   in plants, is a determinant of thylakoid and chloroplast morphology. Proc
1103   Natl Acad Sci U S A *103*, 6759-6764.

1104   28.   Morgan, G.W., Goulding, D., and Field, M.C. (2004). The single dynamin-
1105   like protein of Trypanosoma brucei regulates mitochondrial division and is
1106   not required for endocytosis. J Biol Chem *279*, 10692-10701.

1107   29.   Benz, C., Stribrna, E., Hashimi, H., and Lukes, J. (2017). Dynamin-like
1108   proteins in Trypanosoma brucei: A division of labour between two paralogs?
1109   PLoS ONE *12*, e0177200.

1110   30.   Vanwalleghem, G., Fontaine, F., Lecordier, L., Tebabi, P., Klewe, K., Nolan,
1111   D.P., Yamaryo-Botte, Y., Botte, C., Kremer, A., Burkard, G.S., et al. (2015).
1112   Coupling of lysosomal and mitochondrial membrane permeabilization in
1113   trypanolysis by APOL1. Nature communications *6*, 8078.

1114   31.   Urbaniak, M.D., Guther, M.L., and Ferguson, M.A. (2012). Comparative
1115   SILAC proteomic analysis of Trypanosoma brucei bloodstream and procyclic
1116   lifecycle stages. PLoS ONE *7*, e36619.

1117   32.   Siegel, T.N., Hekstra, D.R., Wang, X., Dewell, S., and Cross, G.A. (2010).
1118   Genome-wide analysis of mRNA abundance in two life-cycle stages of
1119   Trypanosoma brucei and identification of splicing and polyadenylation sites.
1120   Nucleic Acids Res *38*, 4946-4957.

1121   33.   Millerioux, Y., Ebikeme, C., Biran, M., Morand, P., Bouyssou, G., Vincent,
1122   I.M., Mazet, M., Riviere, L., Franconi, J.M., Burchmore, R.J., et al. (2013).
1123   The threonine degradation pathway of the Trypanosoma brucei procyclic

form: the main carbon source for lipid biosynthesis is under metabolic
control. Mol Microbiol *90*, 114-129.

34.   Billington, K., Halliday, C., Madden, R., Dyer, P., Barker, A.R., Moreira-Leite,
F.F., Carrington, M., Vaughan, S., Hertz-Fowler, C., Dean, S., et al. (2023).
Genome-wide subcellular protein map for the flagellate parasite
Trypanosoma brucei. Nat Microbiol *8*, 533-547.

35.   Nunnari, J., Marshall, W.F., Straight, A., Murray, A., Sedat, J.W., and
Walter, P. (1997). Mitochondrial transmission during mating in
Saccharomyces cerevisiae is determined by mitochondrial fusion and fission
and the intramitochondrial segregation of mitochondrial DNA. Mol Biol Cell
8, 1233-1242.

36.   Asencio, C., Herve, P., Morand, P., Oliveres, Q., Morel, C.A., Prouzet-
Mauleon, V., Biran, M., Monic, S., Bonhivers, M., Robinson, D.R., et al.
(2024). Streptococcus pyogenes Cas9 ribonucleoprotein delivery for
efficient, rapid and marker-free gene editing in Trypanosoma and
Leishmania. Mol Microbiol.

37.   Hales, K.G., and Fuller, M.T. (1997). Developmentally regulated
mitochondrial fusion mediated by a conserved, novel, predicted GTPase. Cell
90, 121-129.

38.   Hermann, G.J., Thatcher, J.W., Mills, J.P., Hales, K.G., Fuller, M.T., Nunnari,
J., and Shaw, J.M. (1998). Mitochondrial fusion in yeast requires the
transmembrane GTPase Fzo1p. J Cell Biol *143*, 359-373.

39.   Santel, A., Frank, S., Gaume, B., Herrler, M., Youle, R.J., and Fuller, M.T.
(2003). Mitofusin-1 protein is a generally expressed mediator of
mitochondrial fusion in mammalian cells. J Cell Sci *116*, 2763-2774.

40.   Gambarotto, D., Hamel, V., and Guichard, P. (2021). Ultrastructure
expansion microscopy (U-ExM). Methods in cell biology *161*, 57-81.

41.   Amodeo, S., Kalichava, A., Fradera-Sola, A., Bertiaux-Lequoy, E., Guichard,
P., Butter, F., and Ochsenreiter, T. (2021). Characterization of the novel
mitochondrial genome segregation factor TAP110 in Trypanosoma brucei. J
Cell Sci *134*.

42.   Gorilak, P., Pruzincova, M., Vachova, H., Olsinova, M., Schmidt
Cernohorska, M., and Varga, V. (2021). Expansion microscopy facilitates
quantitative super-resolution studies of cytoskeletal structures in
kinetoplastid parasites. Open Biol *11*, 210131.

43.   Rojo, M., Legros, F., Chateau, D., and Lombes, A. (2002). Membrane
1160   topology and mitochondrial targeting of mitofusins, ubiquitous mammalian
1161   homologs of the transmembrane GTPase Fzo. J Cell Sci *115*, 1663-1674.

1162   44.   Kaurov, I., Vancova, M., Schimanski, B., Cadena, L.R., Heller, J., Bily, T.,
1163   Potesil, D., Eichenberger, C., Bruce, H., Oeljeklaus, S., et al. (2018). The
1164   Diverged Trypanosome MICOS Complex as a Hub for Mitochondrial Cristae
1165   Shaping and Protein Import. Curr Biol *28*, 3393-3407 e3395.

1166   45.   Wenger, C., Oeljeklaus, S., Warscheid, B., Schneider, A., and Harsman, A.
1167   (2017). A trypanosomal orthologue of an intermembrane space chaperone
1168   has a non-canonical function in biogenesis of the single mitochondrial inner
1169   membrane protein translocase. PLoS Pathog *13*, e1006550.

1170   46.   Pyrih, J., Hammond, M., Alves, A., Dean, S., Sunter, J.D., Wheeler, R.J.,
1171   Gull, K., and Lukes, J. (2023). Comprehensive sub-mitochondrial protein
1172   map of the parasitic protist Trypanosoma brucei defines critical features of
1173   organellar biology. Cell reports *42*, 113083.

1174   47.   Daumke, O., and Praefcke, G.J. (2016). Invited review: Mechanisms of GTP
1175   hydrolysis and conformational transitions in the dynamin superfamily.
1176   Biopolymers *105*, 580-593.

1177   48.   Ramachandran, R., and Schmid, S.L. (2018). The dynamin superfamily.
1178   Curr Biol *28*, R411-R416.

1179   49.   Shepard, K.A., and Yaffe, M.P. (1999). The yeast dynamin-like protein,
1180   Mgm1p, functions on the mitochondrial outer membrane to mediate
1181   mitochondrial inheritance. J Cell Biol *144*, 711-720.

1182   50.   Derelle, R., Torruella, G., Klimes, V., Brinkmann, H., Kim, E., Vlcek, C.,
1183   Lang, B.F., and Elias, M. (2015). Bacterial proteins pinpoint a single
1184   eukaryotic root. Proc Natl Acad Sci U S A *112*, E693-699.

1185   51.   Sawant, P., Eissenberger, K., Karier, L., Mascher, T., and Bramkamp, M.
1186   (2016). A dynamin-like protein involved in bacterial cell membrane
1187   surveillance under environmental stress. Environ Microbiol *18*, 2705-2720.

1188   52.   de Sousa Borges, A., and Scheffers, D.J. (2016). Bacterial dynamin as a
1189   membrane puncture repair kit. Environ Microbiol *18*, 2298-2301.

1190   53.   Michie, K.A., Boysen, A., Low, H.H., Moller-Jensen, J., and Lowe, J. (2014).
1191   LeoA, B and C from enterotoxigenic Escherichia coli (ETEC) are bacterial
1192   dynamins. PLoS ONE *9*, e107211.

54.   Schlimpert, S., Wasserstrom, S., Chandra, G., Bibb, M.J., Findlay, K.C.,
Flardh, K., and Buttner, M.J. (2017). Two dynamin-like proteins stabilize
FtsZ rings during Streptomyces sporulation. Proc Natl Acad Sci U S A *114*,
E6176-E6183.

55.   Bramkamp, M. (2012). Structure and function of bacterial dynamin-like
proteins. Biol Chem *393*, 1203-1214.

56.   Burmann, F., Ebert, N., van Baarle, S., and Bramkamp, M. (2011). A
bacterial dynamin-like protein mediating nucleotide-independent membrane
fusion. Mol Microbiol *79*, 1294-1304.

57.   Jimah, J.R., and Hinshaw, J.E. (2019). Structural Insights into the
Mechanism of Dynamin Superfamily Proteins. Trends Cell Biol *29*, 257-273.

58.   Liu, J., Noel, J.K., and Low, H.H. (2018). Structural basis for membrane
tethering by a bacterial dynamin-like pair. Nature communications *9*, 3345.

59.   Lindsay, M.R., Webb, R.I., and Fuerst, J.A. (1997). Pirellulosomes: a new
type of membrane-bounded cell compartment in planctomycete bacteria of
the genus Pirellula. Microbiology (Reading) *143*, 739-748.

60.   Fuerst, J.A., and Webb, R.I. (1991). Membrane-bounded nucleoid in the
eubacterium Gemmata obscuriglobus. Proc Natl Acad Sci U S A *88*, 8184-
8188.

61.   Jogler, C. (2014). The bacterial 'mitochondrium'. Mol Microbiol *94*, 751-755.

62.   Neumann, S., Wessels, H.J., Rijpstra, W.I., Sinninghe Damste, J.S., Kartal,
B., Jetten, M.S., and van Niftrik, L. (2014). Isolation and characterization of
a prokaryotic cell organelle from the anammox bacterium Kuenenia
stuttgartiensis. Mol Microbiol *94*, 794-802.

63.   Herzog, S., Schumann, M.R., and Fleissner, A. (2015). Cell fusion in
Neurospora crassa. Current opinion in microbiology *28*, 53-59.

64.   Sesaki, H., and Jensen, R.E. (1999). Division versus fusion: Dnm1p and
Fzo1p antagonistically regulate mitochondrial shape. J Cell Biol *147*, 699-
706.

65.   Niemann, M., Wiese, S., Mani, J., Chanfon, A., Jackson, C., Meisinger, C.,
Warscheid, B., and Schneider, A. (2013). Mitochondrial outer membrane
proteome of Trypanosoma brucei reveals novel factors required to maintain
mitochondrial morphology. Mol Cell Proteomics *12*, 515-528.

66. Povelones, M.L., Tiengwe, C., Gluenz, E., Gull, K., Englund, P.T., and Jensen, R.E. (2013). Mitochondrial shape and function in trypanosomes requires the outer membrane protein, TbLOK1. Mol Microbiol *87*, 713-729.

67. Otera, H., Ohsakaya, S., Nagaura, Z., Ishihara, N., and Mihara, K. (2005). Export of mitochondrial AIF in response to proapoptotic stimuli depends on processing at the intermembrane space. EMBO J *24*, 1375-1386.

68. Sheikh, S., Panek, T., Gahura, O., Tyc, J., Zahonova, K., Lukes, J., Elias, M., and Hashimi, H. (2023). A Novel Group of Dynamin-Related Proteins Shared by Eukaryotes and Giant Viruses Is Able to Remodel Mitochondria From Within the Matrix. Mol Biol Evol *40*.

69. Kumar, A., Gok, M.O., Nguyen, K.N., Connor, O.M., Reese, M.L., Wideman, J.G., Munoz-Gomez, S.A., and Friedman, J.R. (2024). A dynamin superfamily-like pseudoenzyme coordinates with MICOS to promote cristae architecture. Curr Biol *34*, 2606-2622 e2609.

70. Mears, J.A. (2024). Mitochondrial biology: Unique membrane remodeling from the matrix. Curr Biol *34*, R581-R583.

71. Mattie, S., Riemer, J., Wideman, J.G., and McBride, H.M. (2018). A new mitofusin topology places the redox-regulated C terminus in the mitochondrial intermembrane space. J Cell Biol *217*, 507-515.

72. Jones, D.T. (1999). Protein secondary structure prediction based on position-specific scoring matrices. J Mol Biol *292*, 195-202.

73. Hannaert, V., Albert, M.A., Rigden, D.J., da Silva Giotto, M.T., Thiemann, O., Garratt, R.C., Van Roy, J., Opperdoes, F.R., and Michels, P.A. (2003). Kinetic characterization, structure modelling studies and crystallization of Trypanosoma brucei enolase. Eur J Biochem *270*, 3205-3213.

74. Hirumi, H., and Hirumi, K. (1989). Continuous cultivation of Trypanosoma brucei blood stream forms in a medium containing a low concentration of serum protein without feeder cell layers. J Parasitol *75*, 985-989.

75. Dean, S., Sunter, J., Wheeler, R.J., Hodkinson, I., Gluenz, E., and Gull, K. (2015). A toolkit enabling efficient, scalable and reproducible gene tagging in trypanosomatids. Open Biol *5*, 140197.

76. Wirtz, E., Leal, S., Ochatt, C., and Cross, G.A. (1999). A tightly regulated inducible expression system for conditional gene knock-outs and dominant-negative genetics in Trypanosoma brucei. Mol Biochem Parasitol *99*, 89-101.

77.　Biebinger, S., Wirtz, L.E., Lorenz, P., and Clayton, C. (1997). Vectors for
inducible expression of toxic gene products in bloodstream and procyclic
Trypanosoma brucei. Mol Biochem Parasitol *85*, 99-112.

78.　Ling, M.M., and Robinson, B.H. (1997). Approaches to DNA mutagenesis:
an overview. Anal Biochem *254*, 157-178.

79.　Wickstead, B., Ersfeld, K., and Gull, K. (2002). Targeting of a tetracycline-
inducible expression system to the transcriptionally silent minichromosomes
of Trypanosoma brucei. Mol Biochem Parasitol *125*, 211-216.

80.　Blancard, C., Decoeur, F., Duvezin-Caubet, S., Giraud, M.F., and Salin, B.
(2024). Advancing yeast cell analysis: A cryomethod for serial block-face
scanning electron microscopy imaging in mitochondrial morphology studies.
Biology of the cell / under the auspices of the European Cell Biology
Organization, e202400038.

81.　Blancard, C., and Salin, B. (2017). Plunge Freezing: A Tool for the
Ultrastructural and Immunolocalization Studies of Suspension Cells in
Transmission Electron Microscopy. J Vis Exp, e54874.

82.　Bochud-Allemann, N., and Schneider, A. (2002). Mitochondrial substrate
level phosphorylation is essential for growth of procyclic Trypanosoma
brucei. J Biol Chem *277*, 32849-32854.

83.　Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. (1990).
Basic local alignment search tool. J Mol Biol *215*, 403-410.

84.　Richter, D.J., Berney, C., Strassert, J.F.H., Poh, Y.-P., Herman, E.K., Muñoz-
Gómez, S.A., Wideman, J.G., Burki, F., and de Vargas, C. (2022). EukProt:
A database of genome-scale predicted proteins across the diversity of
eukaryotes. bioRxiv, 2020.2006.2030.180687.

85.　Katoh, K., and Standley, D.M. (2013). MAFFT multiple sequence alignment
software version 7: improvements in performance and usability. Mol Biol
Evol *30*, 772-780.

86.　Criscuolo, A., and Gribaldo, S. (2010). BMGE (Block Mapping and Gathering
with Entropy): a new software for selection of phylogenetic informative
regions from multiple sequence alignments. BMC Evol Biol *10*, 210.

87.　Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D.,
von Haeseler, A., and Lanfear, R. (2020). IQ-TREE 2: New Models and
Efficient Methods for Phylogenetic Inference in the Genomic Era. Mol Biol
Evol *37*, 1530-1534.

88.   Claros, M.G., and Vincens, P. (1996). Computational method to predict
mitochondrially imported proteins and their targeting sequences. Eur J
Biochem *241*, 779-786.

1299

Figure

Figure 1

Figure 2

Figure 3

Figure 4

A

MTS — CC — ▮ — GTPase — CC — TM TM — CC — CC — 3Ty1

B

^{oe}TbMfnL-K141A$_{::3Ty1}$

d3		d5	
-Tet	+Tet	-Tet	+Tet

70 —   Anti-Ty1

130 —   Anti-Frdg

C

^{oe}TbMfnL-K141A$_{::3Ty1}$ (IF)

Anti-Tdh   Anti-Ty1

d5+Tet

DAPI   Merge

n   k

D

Rhodamine 123

WT

5 μm

^{oe}TbMfnL-K141A$_{::3Ty1}$

-Tet

5 μm

d5+Tet

5 μm

Figure 5

Figure 6

Figure 7

Figure 8

SUPPLEMENTAL INFORMATION

Data S1 - Code ImageJ: Mitochondrial junctions (MJ)

```
run("Duplicate...", "title=ORI duplicate channels=2");
run("Duplicate...", "title=Estfond");
run("Gaussian Blur...", "sigma=10");
imageCalculator("Divide create 32-bit", "ORI","Estfond");
selectWindow("Result of ORI");
rename("OriCorr");
selectWindow("Estfond");
setAutoThreshold("Li dark");
run("Create Selection");
selectWindow("OriCorr");
run("Restore Selection");
run("Enlarge...", "enlarge=10 pixel");
setAutoThreshold("Li dark");
run("Analyze Particles...", "size=15-500 show=Masks");
rename("Mask");
run("Options...", "iterations=1 count=4 pad do=Open");
run("Duplicate...", "title=Skeleton");
setOption("BlackBackground", false);
run("Skeletonize");
run("Duplicate...", "title=Ends");
run("Options...", "iterations=1 count=7 pad do=Erode");
imageCalculator("Subtract create", "Skeleton","Ends");
selectWindow("Result of Skeleton");
rename("Endpoints");
selectWindow("Skeleton");
run("Duplicate...", "title=Jonctions");
run("Options...", "iterations=1 count=6 pad do=Erode");
run("Analyze Particles...", "size=0-5000 summarize");
imageCalculator("Add create", "Endpoints","Jonctions");
selectWindow("Result of Endpoints");
rename("Combo");
imageCalculator("Subtract create", "Skeleton","Combo");
selectWindow("Result of Skeleton");
rename("Segments");
selectWindow("Combo");
close();
selectWindow("ORI");
setOption("ScaleConversions", true);
run("8-bit");
run("Merge Channels...", "c1=Jonctions c2=Endpoints c3=Segments c4=ORI create ignore"); close("\\Others");
```

TABLE S1 TO S3

Table S1 – Results from BLASTp and EukProt v3 (query: *Tb*MfnL) of several identified organisms

Name	Organism	Domain Clade	Sequence ID	Amino acid	E-Value	Cover (%)	Identity (%)	Presence of Opa1 /Mgm1	Presence of Mfn /Fzo1	Presence of MidX**	Mitochondrial targeting signal prediction TargetP-2.0 PSORT II MitoFates
*Tb*MfnL	*Trypanosoma brucei*	Eukaryota Discoba; Euglenozoa Kinetoplastea	Tb927.7.2410 (XP_845864.1)	600	0.0	100	100	NO	NO	NO	0.6068 0.3040 0.2000
*El*MfnL	*Euglena longa*	Eukaryota Discoba; Euglenozoa Euglenida	EP00668_Euglena_ longa_P029018	530	8.9e-115 (7e-111)	83	42.8	NO	NO	NO	ND*
*Dp*MfnL	*Diplonema papillatum*	Eukaryota Discoba; Euglenozoa Diplonemea	KAJ9468802.1	479	1e-87 (2e-84)	76	36.3	NO	NO	NO	0.7375 0.3910 0.9300
*Pg*MfnL	*Polarella glacialis*	Eukaryota Sar; Alveolata Dinophyceae	CAE8737761.1	601	2e-79 (3e-80)	73	39.1	NO	NO	NO	0.0063 0.1110 0.0030
*Hf*MfnL	*Hondaea fermentalgiana*	Eukaryota Sar; Stramenopiles Bigyra	GBG23909.1	536	4e-77 (1e-78)	82	35.6	NO	NO	NO	0.5606 0.2170 0.9540
*Ah*MfnL	*Achlya hypogyna*	Eukaryota Sar; Stramenopiles Oomycota	OQR97000.1	498	2e-81 (2e-82)	82	36.6	NO	NO	NO	0.9793 0.7390 0.9990
*Ct*MfnL	*Chaetoceros tenuissimus*	Eukaryota Sar; Stramenopiles Ochrophyta	GFH50037.1	552	1e-77 (6e-78)	74	35.6	NO	NO	NO	0.6911 0.3040 0.8800
*Cht*MfnL	*Chrysochromulina tobinii*	Eukaryota Haptista; Haptophyta Pavlovales	KOO35358.1	514	2e-65 (4e-72)	51	42.5	NO	NO	NO	0.5910 0.2610 0.8740
*Pb*MfnL	*Planctomycetota* bacterium	Prokaryota Planctomycetota	HLY10620.1	448	2e-81 (3e-81)	74	36.8	NO	NO	NO	0.0000 0.1110 0.0000
*Ps*MfnL	*Pedosphaera* sp.	Prokaryota Verucomicrobia	MCH2381658.1	479	2e-75 (1e-76)	79	35.3	NO	NO	NO	0.0099 0.2170 0.6920
*Rc*MfnL	*Rubrimonas cliftonensis*	Prokaryota α-Proteobacteria	WP_093253540.1	450	6e-74 (6e-80)	74	35.6	NO	NO	NO	0.0002 0.1110 0.0100
*Gp*MfnL	*Gammaproteobacteria* bacterium	Prokaryota γ-Proteobacteria	RTZ59088.1	457	1e-71 (6e-80)	75	34.7	NO	NO	NO	0.0000 0.2220 0.0070
*Mb*MfnL	*Myxococcales* bacterium	Prokaryota δ-Proteobacteria	MCA9558847.1	452	5e-70 (4e-71)	77	33.3	NO	NO	NO	0.0001 0.3330 0.0010
*Ld*MfnL	*Lentisphaerae* bacterium	Prokaryota Lentisphaerae	NLB69313.1	465	2e-54 (5e-55)	64	33.1	NO	NO	NO	0.0168 0.0000 0.2470
*Kb*MfnL	*Kiritimatiellae* bacterium	Prokaryota Kiritimatiellae	MBO7223041.1	459	3e-52 (2e-55)	42	43.0	NO	NO	NO	0.0003 0.2610 0.0730
*Hs*Mfn	*Homo sapiens mitofusin-2*	Eukaryota Homo sapiens	NP_001121132.1	757	<1e-2	70	15.4	YES (Opa1)	YES (Fzo1)	NO	0.0787 0.2170 0.0880
*Sc*Fzo1	*Saccharomyces cerevisiae*	Eukaryota Fungi	NP_009738.1	855	<1e-2	61	16.8	YES (Mgm1)	YES (Mfn)	NO	0.0000 0.0430 0.0000
*Nc*BDLP	*Nostoc punctiforme PCC 73102*	Bacteria Cyanobacteriota	WP_2412711.1	693	<1e-2	69	18.2	NO	YES (Mfn)	NO	0.0001 0.1740 0.0240
*Tc*MfnL	*Trypanosoma cruzi*	Eukaryota Discoba; Euglenozoa Kinetoplastea	RNC57734.1	611	0.0	83	54.2	NO	NO	NO	0.6242 0.3910 0.7960
*Le*MfnL	*Leishmania enriettii*	Eukaryota Discoba; Euglenozoa Kinetoplastea	XP_067693110.1	602	2e-130	97	39.5	NO	NO	NO	0.4552 0.1110 0.8820
*Phs*MfnL	*Phytomonas sp. EM1*	Eukaryota Discoba; Euglenozoa Kinetoplastea	CCW65548.1	590	5e-89	78	32.2	NO	NO	NO	0.8692 0.3910 0.9800

ND*, not determined as the N-terminus sequence is unavailable. Sequence IDs beginning with "EP" originate from EukProt v3 database. Parentheses in the E-Value column indicate the E-value obtained from the reciprocal BLASTp analysis, using the putative MfnL ortholog as the query against the *T. brucei* proteome. E-score value below 1e-2 could be considered as "not significant". **We were unable to identify the MidX sequence in the genomes of species that contain the *Tb*MfnL sequence.

Table S2 – Primers used in this study

Name	Sequence	Used for
ET198	TTCGTGTTGCCAACTACAAGGTGAAAAATTTTCTGCGAGCTGGTGAACGGCAACCGCTGGGTAATGAGGTGAGGGTCAAAGGTTCTGGTAGTGGTTCC	Endotag
ET199	CATGCTCGGCAACCTAAGCATGATAGTCCGCGGTAAGAGCAAAAGAACCAGCGCCCCACGTAAATTTGCTTCCAACATTTCCAATTTGAGAGACCTGTGC	
ET160	CATAACGTTACCCACCACATCATCACTGAAGCGACTGTTGCGCAGAAGTCATGACCGAGTACAAGCCCACGGTG	CRISPR/Cas9
ET161	TGTAGTAGTCGCAAAACACATGCGGTTAGTCGAACAGTTGGCACCCAACTTCAGGCACCGGGCTTGCGGGTCATG	
ET211	ATTGACACTCCTATTCATTTGAATGATCGGACATCAGTTGAGGGTCAGCTATGACCGAGTACAAGCCCACGGTG	
ET242	CACACCGTGAAGCAAACCAACGTGTCACGGCGAACAGATCATACCCCCTGTCAGGCACCGGGCTTGCGGGTCATG	
ET92	GCGGCGAGGGTGCGTACGGCCCGG	
ET93	TCACCGTCACCGCCGACGTCGAGG	
ET109	CTGTGGTTCCCAAGGACAATCACC	
MFN_102	ACTGTTGCGCAGAAGTCAGT(NGG)	
MFN_721	TCAGTTGAGGGTCAGCTCAG(NGG)	
MFN_1772	AACCGCTGAGTGATGAGGCG(NGG)	
ET189	GATCAAGCTTCCACATCATCACTGAAGCGACTGT	RNAi
ET190	GATCCTCGAGGTTGGAGATACGCCCGATCGCTGC	
ET191	GATCAAGCTTCTCGAGCCTCGGTTATGTCGTCTTCACCAC	
ET192	GATCGGATCCCCACATCATCACTGAAGCGACTGT	
ET362	GATCGGATCCTTATTTTGATGTTTGACCCC	
ET363	GATCCTCGAGCGGTTGAGGAACTCTTATTG	
ET94	GATCCATATGGGATCCAAGCTTATGAGGCAGTTAGGAAGTGCAG	*Tb*MfnL
ET95	GATCGGTACCGCGGCCGCTCTAGATTATTTGACCCTCACCTCATTACC	
ET96	GATCGGTACCTCTAGATTTGACCCTCACCTCATTACCCAG	
ET149	GAACCACAGCGCGGGAGCGTCTACGATGATTAATC	K141A
ET150	GATTAATCATCGTAGACGCTCCCGCGCTGTGGTTC	
ET197	GATCGCGGCCGCAAGCTTCATATGGGTGCCAACTGCTCGACTAACC	ΔMTS
ET292	CCGGAAGATGAAGGTTAAGTCATTGGTTGACTTTG	ΔTM
ET293	CAAAGTCAACCAATGACTTAACCTTCATCTTCCGG	

Table S3 – Antibodies used in this study

Name	Species	Dilution	Source/Ref
Anti-Ty1	Mouse	WB 1:800	iMET lab
		IF 1:500	
		UExM 1:250	
Anti-HA	Mouse	WB 1:1000	Biolegend
		IF 1:1000	
		UExM 1:500	
Anti-HA	Rat	IF 1:1000	Roche
Anti-Tdh	Rabbit	WB 1:500	iMET lab
		IF 1:200	
		UExM 1:100	
Anti-Enolase	Rabbit	WB 1:100000	Gift from P. Michels, Edinburgh, UK
Atom40	Rabbit	WB 1:10000	Gift from A. Schneider, Bern, Switzerland
Anti-Frdg	Rabbit	WB 1:50	iMET lab
Anti-Hsp60 H7	Mouse	IF 1:500	iMET lab
Anti-PPDK	Rabbit	WB 1:1000	iMET lab
Anti-GFP	Rabbit	WB 1:1000	Abcam
Anti-Prohibitin	Rabbit	WB 1:1000	Gift from H. Hashimi, Czech Republic
Anti-Mouse HRP	Goat	WB 1:5000	Bio-Rad
Anti-Rabbit HRP	Goat	WB 1:10000	Bio-Rad
Anti-Mouse Alexa Fluor 594	Donkey	IF 1:400	ThermoFisher
		UExM 1:250	
Anti-mouse Alexa Fluor 488	Donkey	IF 1:400	ThermoFisher
		UExM 1:250	
Anti-Rabbit Alexa Fluor 594	Donkey	IF 1:400	ThermoFisher
		UExM 1:250	
Anti-Rabbit Alexa Fluor 488	Donkey	IF 1:400	ThermoFisher
		UExM 1:250	
Anti-Rat Alexa Fluor 594	Donkey	IF 1:400	ThermoFisher

Supplemental figures

Figure S1 - Sequence alignement between *Tb*MfnL and various kinetoplastids MfnL. MTS, predicted mitochondrial targeting sequence are indicated by grey boxes; GTPase domain is indicated by a green box, and the four GTP-binding motifs are indicated by yellow boxes and numbered; The coil coil domain are indicated by light brown boxes. TM, transmembrane span are indicated by purple boxes. Identities and similarities are represented by white characters on a black background and black characters on a grey background respectively. *Tb*MfnL, *Trypanosoma brucei* DRP (ID: Tb927.7.2410); *Phy*MfnL, *Phytomonas* sp. EM1(ID: CCW65548); *Pc*MfnL, *Paratrypanosoma confusum* (ID: PCON_0070060); *Bs*MfnL, *Bodo saltans* (ID: BSAL_56690); *Ba*MfnL, *Blechomonas ayalai* (ID: Baya_002_1420); *Cf*MfnL, *Crithidia fasciculata* (ID: CFAX1_240011500); *Ls*MfnL, *Leptomonas seymouri* (ID: Lsey_0356_0020); *Li*MfnL, *Leishmania infantum* (ID: LINF_220010300); *Em*MfnL, *Endotrypanum monterogeii* (ID: EMOLV88_220009500); *Ph*MfnL, *Porcisia hertigi* (ID: JKF63_04376); *Tc*MfnL, *Trypanosoma cruzi* (ID: TcCLB.508321.60); *Tev*MfnL, *Trypanosoma evansi* (ID: TevSTIB805.7.2480); *Te*MfnL, *Trypanosoma equiperdum* (ID: TEOVI_000623300); *Tv*MfnL, *Trypanosoma vivax* (ID: TvY486_0702250); *Tc*MfnL, *Trypanosoma congolense* (ID: TcIL3000_7_1710); *Tr*MfnL, *Trypanosoma rangeli* (ID: TRSC58_06192); *Tg*MfnL, *Trypanosoma grayi* (ID: DQ04_08691020); *Tm*MfnL, *Trypanosoma melophagium* (ID: LSM04_004715); *Tt*MfnL, *Trypanosoma theileri* (ID: TM35_000191450).

Figure S2 - Sequence and structure alignement between *Tb*MfnL and homologous proteins identified in various eukaryotes and prokaryotes with BLASTp. MTS, predicted mitochondrial targeting sequence are highlighted in grey; GTPase domain is highlighted in green, and the four GTP-binding motifs are highlighted in yellow and numbered; The coil coil domain are highlighted in light brown. TM, transmembrane span are highlighted in purple. Identities and similarities are represented by white characters on a black background and black characters on a grey background respectively. Structure predictions were performed using PSIPRED and mitochondrial targeting sequences were predicted by various algorithms (TargetP2.0, PSORTII, MitoFates, MitoprotII and iPSORT). **Eukayota:** *Tb*MfnL, *Trypanosoma brucei* DRP (ID: Tb927.7.2410); *El*MfnL, *Euglena longa* (ID: EP00668_Euglena_longa_P029018); DpMfnL, *Diplonema papillatum* (ID: KAJ9468802.1)*;* *Pg*MfnL, *Polarella glacialis* (ID: CAE8737761.1); *Ah*MfnL, *Achlya hypogyna* (ID: OQR970001); *Ct*MfnL, *Chaetoceros tenuissimus* (ID: GFH50037.1); *Hf*MfnL, *Hondaea fermentalgiana* (ID: GBG23909.1); *Cht*MfnL, *Chrysochromulina tobinii* (ID: KOO35358.1). **Prokaryota:** *Pb*MfnL, *Planctomycetaceae* bacterium (ID: MBV8881356.1); *Ps*MfnL, *Pedosphaera* sp. (ID: MCH2381658.1); *Rc*MfnL, *Rubrimonas cliftonensis* (ID: WP_093253540.1); *Gp*MfnL, *Gammaproteobacteria* bacterium (ID: RTZ59088.1); *Mb*MfnL, *Myxococcota* bacterium (ID: MBU0550280.1); *Ld*MfnL, *Lentisphaerae* bacterium (ID: NLB69313.1); *Kb*MfnL, *Kiritimatiellae* bacterium (ID: MBO7223041.1). The accession number for *Euglena longa* was obtained from Eukprot V3 database. The alignment was performed using Geneious software with the BLOSUM62 matrix.

Figure S3 - Maximum likelihood phylogenetic tree of eukaryotic and eukaryotic DSPs. The tree was reconstructed using the LG+C60+G4 model of sequence evolution. Ultrafast bootstrap support values (1000 replicates) are shown on each branch. Taxa are colored according to their taxonomic affiliation.

Figure S4 – Growth curves of parental, over-expressing and mutant cells. Cells were maintained in the exponential growth phase and cumulative cell numbers reflect normalization for dilution during cultivation. (.ni) non-induced and (.i) induced cells. PCF, procyclic forms; BSF, bloodstream forms. n=3 to 6.

Figure S5 - *Tb*MfnL endotagged with a 3xTy1 sequence and inactivation of *Tb*MfnL expression by CRISPR/-Cas9 in PCF. (A) Design of the endotagged strain with addition of the 3Ty1 tags at the C-terminus of the protein using the CRISPR/Cas9 system. **(B)** Selection of homozygous clones by PCR (2A11, 1E4 and 1G6) using the PCR approach shown in panel A. **(C)** Western-blot analysis of whole cell extracts (5×10^6 cells) from *T. brucei* PCF clones expressing *Tb*MfnL$_{::3Ty1}$ revealed with anti-Ty1 antibody and anti-Enolase as a control, and expression levels between homozygous and heterozygous cells. **(D)** Analysis of mitochondrial structure by rhodamine 123 staining in live cells (clone 2A11); **(E)** CRISPR/Cas9-mediated inactivation of *Tb*MfnL using a single guide RNA and insertion of a puromycin resistance marker (PAC) in the *Tb*MfnL$_{::3Ty1}$ (2A11 clone). **(F)** Western blot of whole cell extracts (5×10^6 cells) of *Tb*MfnL$_{::3Ty1}$ (clone 2A11 background) and *Tb*MfnL$_{::3Ty1::PAC}$ (A12 and D9) revealed by anti-Ty1 antibody. **(G)** Rhodamine staining of *Tb*MfnL$_{::3Ty1::PAC}$ (clone A12). **(H)** CRISPR/Cas9-mediated inactivation of *Tb*MfnL using two different guide RNAs that deletes approximately 600 bp of the coding sequence in the *Tb*MfnL$_{::3Ty1}$ (2A11 background). **(I)** Western blot of whole cell extracts (5×10^6 cells) of *Tb*MfnL$_{::3Ty1}$ (2A11 background) and Δ102/720-*Tb*MfnL$_{::3Ty1}$ clones (1C4, 1E6 and 1E7) revealed by anti-Ty1 and anti-Enolase antibodies. **(J)** Rhodamine staining of Δ102/720-*Tb*MfnL$_{::3Ty1}$ (clone 1C4).

Figure S6 - Mitochondrial ultrastructure analyzed by transmission electron microscopy (TEM). Cells cultured with either glucose or proline, including those overexpressing *Tb*MfnL::3Ty1 and inactivated TbMfnL⁻/⁻, were examined by TEM. Overexpression of *Tb*MfnL::3Ty1 was induced for 5 days with tetracycline (Tet) prior to TEM analysis. Panels A–F provide an overview, while panels G–L focus on the mitochondrion and cristaes. n, nucleus; m, mitochondrion; k, kinetoplast; fp, flagellar pocket; white arrows point to the cristae.

Figure S7 - Inhibition of *Tb*MfnL expression by RNAi in PCF (A/B/C) and BSF (D/E/F/G). (A) Shematic representation of the sequence used for down-regulation of *Tb*MfnL expression in PCF. **(B)** *Tb*MfnL::10HA down-regulation by RNA interference in PCF. Western blot analysis of total extract (5x10^6 PCF cells) of non-induced (-Tet) and induced (+Tet) cells using anti-HA antibody after 3 days of tetracycline induction. Antibody against Enolase was used as a loading control. **(C)** Rhodamine 123 labelled mitochondrial network of cells incubated with tetracycline for 3 days. **(D)** Shematic representation of the sequence used for down-regulation of *Tb*MfnL expression in BSF. **(E)** *Tb*MfnL::3HA down-regulation by tetracycline-induced RNA interference in BSF. Western blot analysis of the total extract (5x10^6 BSF cells) of non-induced (-Tet) and induced (+Tet) cells using anti-HA antibody after 4 days of tetracycline induction. Antibody against Enolase was used as a loading control. **(F)** Relative expression of down-regulation of *Tb*MfnL expression by RNAi in BSF. **(G)** Mitochondrial network labelled with anti-HSP60 from cells incubated or not with tetracycline for 3 days. The scale bar represents 5 μm.

Figure S8 - Mitochondrial shape of parental WT, ^{oe}*Tb*MfnL_{::3Ty1} and ^{oe}*Tb*-MfnL-K141A_{::3Ty1} PCF cells. The top panel depicts live cells stained with rhodamine 123. Bottom panel was obtained using the ImageJ macro (Mitochondrial Junctions). Tet refers to tetracycline, d5, tetracycline induced cells for 5 days. The scale bar corresponds to 5 μm. ni , non-induced cells; d5 .i, induced cells for 5 days. The scale bar corresponds to 5 μm. Red circles in the skeleton image represent the junctions of three mitochondrial tubules.

Figure S9 - Overexpression of *Tb*MfnL in endotagged cells. (A) Western-blot of whole-cell extracts (5x10⁶ cells) of *T. brucei* parental WT PCF and expressing various *Tb*MfnL in *Tb*MfnL::3Ty1 background (clone 2A11) revealed with an anti-Ty1 antibody. Quantification of the expression were performed using ImageJ with PPDK as a loading control. **(B)** Mitochondrial structure analysis using rhodamine 123 staining on living cells; mitochondrial structure of parental WT PCF and expressing various *Tb*MfnL in *Tb*MfnL::3Ty1 background without tetracycline (-Tet) and 5 days after addition of tetracycline (+Tet). The scale bar represents 5 µm.

**Figure S10 - Variation in the expression level of *Tb*MfnL:
:3Ty1 induced by tetracycline..** (A) Western blot analysis of
whole-cell extracts (5 x 10⁶ cells) from *T. brucei* PCF
expressing *Tb*MfnL::3Ty1 and *Tb*MfnL-141A::3Ty1 in the
*Tb*MfnL::3Ty1 background (clone 2A11) probed with an anti-
Ty1 antibody. Expression levels were quantified using
ImageJ, with PPDK as a loading control, at day 3 and day 5
following tetracycline induction. (B) Corresponding
mitochondrial structures were visualized using Mitotracker
staining on live cells. The scale bar represents 5 µm.

Figure S11 - Immuno localization of ^{oe}TbMfnL$_{::3Ty1}$ by electron microscopy. Immuno localization of ^{oe}TbMfnL$_{::3Ty1}$ **(A)** and ^{oe}TbDBFK141A$_{::3Ty1}$ **(B)** expressed in PCF by electron microscopy. Non-induced cells (-Tet, a and b) and 5 days induced cells (+Tet, c and d) were labelled with an anti-Ty1 antibody and an anti-mouse IgG-gold particle conjugate; black dots represent gold particles highlighted by white arrows; b/d are magnification of a and c, respectively. **(C)** Quantification of mitochondrial gold particles. For non-induced (-Tet) and induced (+Tet) ^{oe}TbMfnL$_{::3Ty1}$ cells, 221 and 3076 gold particles were counted, respectively, and for non-induced (-Tet) and induced (+Tet) ^{oe}TbMfnL-K141A$_{::3Ty1}$ cells, 78 and 1123 gold particles were counted, respectively. A mean of the non-induced (-Tet) cells from the two strains is represented on the graph. Statistic: t test - Confidence interval 95% - P-value style: 0.1234 (ns); 0.0332 (*); 0.0021 (**); 0.0002 (***); <0.0001 (****).

Figure S12 - Membrane topology of *Tb*Mfnl (entire western blot). (A) The Proteinase K protection assay involved incubating mitoplasts with Proteinase K and Triton X100, as denoted by the"+" symbol. TF represents the total fraction containing both cytoplasmic and membrane components; S denotes the supernatant, comprising the cytosolic contents; and Pellet represents the organelle fraction. **(B)** Carbonate extraction of mitochondria isolated from ᵒᵉ*Tb*MfnL ₍₎₃Ty1 and ᵒᵉ*Tb*MfnL-ΔTM₍₎₃Ty1 cells. Mem., membrane fraction; Matrix, matrix fraction; Int. Mem, integral membrane fraction; Per. Mem, peripheral membrane fraction. The star indicates the corresponding protein. The expected sizes of the proteins are as follows: *Tb*MfnL - 62 kDa, Enolase - 46 kDa, Tdh - 37 kDa, Prohibitin - 31 kDa, and Atom40 - 38 kDa.

Figure S13 - AlphaFold structure of *Tb*MfnL with pLDDT scores for each amino acid. The pLDDT scores are color-coded as blue (>90), light blue (>70), yellow (>50), and orange (<50). The MTS sequence is highlighted by a black box, the GTPase domain by a green box, and the transmembrane domains by purple boxes. The average pLDDT score for the entire protein was 72.5.