Lanthanide-dependent isolation of phyllosphere methylotrophs selects for a

phylogenetically conserved but metabolically diverse community

Alekhya M. Govindaraju[a], Colleen A. Friel[b], Nathan M. Good[a], Sidney L. Banks[a], Kenan S.

Wayne[a], N. Cecilia Martinez-Gomez[,#]

a. Department of Plant and Microbial Biology, University of California, Berkeley, Berkeley, CA,

USA

b. Department of Microbiology and Molecular Genetics, Michigan State University, East

Lansing, MI, USA

#Address correspondence to: N. Cecilia Martinez-Gomez, cecimartinez@berkeley.edu

Running Title

Lanthanide-dependent isolation of phyllosphere methylotrophs

Abstract

The influence of lanthanide biochemistry during methylotrophy demands a reassessment of how

the composition and metabolic potential of methylotrophic phyllosphere communities are

affected by the presence of these metals. To investigate this, methylotrophs were isolated from

soybean leaves by selecting for bacteria capable of methanol oxidation with lanthanide cofactors.

Of the 344 pink-pigmented facultative methylotroph isolates, none were obligately lanthanide-

dependent. Phylogenetic analyses revealed that all strains were nearly identical to each other and

to model strains from the *extorquens* clade of *Methylobacterium*, with *rpoB* providing higher

resolution than 16s *r*RNA for strain-specific identification. Despite the low species diversity, the

metabolic capabilities of the community diverged greatly. Strains encoding identical PQQ-

dependent alcohol dehydrogenases displayed significantly different growth from each other on

alcohols in the presence and absence of lanthanides. Several strains also lacked well-

characterized lanthanide-associated genes thought to be important for phyllosphere colonization.

Additionally, 3% of our isolates were capable of growth on sugars and 23% were capable of

growth on aromatic acids, substantially expanding the range of multicarbon substrates utilized by

members of the *extorquens* clade in the phyllosphere. Whole genome sequences of eleven novel

strains are reported. Our findings suggest that the expansion of metabolic capabilities, as well as

differential usage of lanthanides and their influence on metabolism among closely related strains,

point to evolution of niche partitioning strategies to promote colonization of the phyllosphere.

Importance

Lanthanides are essential metals for life. The identification of lanthanide-associated processes

has been well-studied in methylotrophic bacteria, which are plant symbionts that utilize reduced

one-carbon compounds for growth. Yet, the importance of lanthanides in plant-microbe

interactions and the effects of lanthanides on microbial physiology and colonization in plants

remains poorly understood. Here, we characterize the first methylotrophic bacterial community

isolated from the phyllosphere in a lanthanide-dependent manner. We have identified strains

encoding identical lanthanide-dependent enzymes yet exhibiting differences in lanthanide-

associated growth, and have identified strains lacking lanthanide-associated genes thought to be

important for phyllosphere colonization. In addition, we have identified many strains capable of

metabolisms that were thought to be rare within this clade. Overall, our isolates serve as a

microcosm by which to interrogate how lanthanides influence methylotrophic physiology in

plant environments and highlights how phylogenetically similar strains can diverge greatly in

metabolic potential.

Introduction

The phyllosphere, the aerial portion of plants dominated by leaves (1), is colonized by a

diverse community of microorganisms (2–4). Despite the variety of potential substrates provided

by plant leaves (carbohydrates, organic acids, amino acids, volatile compounds), microorganisms

in the phyllosphere must overcome waxy leaf cuticles, UV light, temperature fluctuations, low

water availability and other biotic and abiotic impediments to colonization (2,3). Cultivation-

independent studies of the phyllosphere have only emerged within the last decade (3). A

proteogenomic study characterized the phyllosphere community of *Glycine max* (soybean) and

showed high levels of proteins characteristic of the *Alphaproteobacteria* class and,

specifically, the *Methylobacterium* genus (5). Methylotrophs of the *Methylobacterium* genus are

so-named due to their capacity to utilize reduced one-carbon compounds such as methane,

methanol, methylamine, formaldehyde, or formate as their sole source of carbon and energy (6) .

In the phyllosphere, methylotrophs are well-suited to take advantage of methanol released daily

by pectin methylesterases during routine plant cell wall breakdown and repair (100 Tg y^{-1}) (4,7).

Methylotrophs may even stimulate plants into releasing additional methanol by secreting the

plant hormone cytokinin, which triggers plant cell division (2).

In Gram-negative methylotrophic bacteria, pyrroloquinoline quinone (PQQ) methanol

dehydrogenases convert methanol to formaldehyde (8). Although the first-identified methanol

dehydrogenase from Gram-negative methylotrophs was the calcium (Ca^{2+}) -dependent MxaFI

(6,8), recent studies have highlighted the importance of an alternative methanol dehydrogenase,

XoxF, which coordinates a lanthanide atom in complex with PQQ (8–10). Prior to the discovery

of XoxF, the calcium (Ca^{2+})-dependent methanol dehydrogenase MxaFI was thought to be the

only enzyme responsible for methanol oxidation in this system. XoxF is widespread in the

environment (11,12) and phylogenetically ancestral to MxaFI (8). Metaproteomics analysis of

the soybean phyllosphere showed comparable levels of XoxF to MxaFI (5,13). In addition to

XoxF, ExaF is a lanthanide-dependent alcohol dehydrogenase with sub-nanomolar K_M towards

ethanol and auxiliary activity with formaldehyde, a toxic intermediate of methylotrophic

metabolism (14). Induction of XoxF can repress MxaFI expression at lanthanide concentrations

higher than 100 nM (15). Although the importance of lanthanide-dependent alcohol

dehydrogenases in alcohol and aldehyde metabolism is clear, the extent to which lanthanides

influence other metabolisms and the diverse set of proteins that regulate lanthanide-dependent

processes are open areas of research.

Lanthanides have expanded our understanding of methylotrophic metabolism and

ecology even beyond the phyllosphere. The addition of lanthanides to the isolation process of

microbial communities has led to the discovery of new cultivable methylotrophs in coal slag (16)

and marine environments (17), but this phenomenon has yet to be tested on phyllosphere

communities. Lanthanides have also unlocked previously unknown metabolic capabilities of

well-characterized plant symbionts. For example, a *Bradyrhizobium* strain isolated from legumes

displayed little to no growth on methanol, despite maintaining genes for methanol metabolism,

until the addition of lanthanides to the growth medium (18).

Lanthanide utilization during alcohol oxidation in methylotrophs has been well-

characterized, yet methylotrophs can also take advantage of other substrates in the phyllosphere

whose metabolism may or may not be influenced by lanthanides. For example, methylotrophs

have several mechanisms by which they can grow on the C_1 compound, methylamine.

Methylamine dehydrogenases (*mauFBEDACJGLMN*) catalyze the oxidation of methylamine to

formaldehyde, which can be further oxidized to formate for assimilation or dissimilation (6).

Alternatively, methylamine can be converted via the *N*-methylglutamate pathway (*mgdDCBA*,

mgsABC, *gmaS*) to methylene-tetrahydrofolate for assimilation or dissimilation, although the

extent of formaldehyde as an obligate intermediate in this pathway remains unknown (19). The

model organism *M. extorquens* AM1 encodes both pathways yet primarily uses methylamine

dehydrogenase during growth on methylamine, whereas the model organism *M. extorquens* PA1

only encodes the N-methylglutamate pathways (19,20). Organisms solely utilizing the *N*-

methylglutamate pathway have markedly slower growth on methylamine than those that can

catalyze the direct oxidation to formaldehyde (19,20).

Methylobacterium species can be further classified as pink-pigmented facultative

methylotrophs (PPFMs) (4) that can utilize multi-carbon substrates available in the phyllosphere,

including acetate (C2), pyruvate (C3), and succinate (C4). Recently, facultative methylotrophy

has expanded to include species capable of growth on sugars within the *extorquens* clade (21)

and species capable of growth on methoxylated aromatic acids within the *nodulans* and

aquaticums clades of *Methylobacterium* (22). Methylotrophic growth on the latter is predicted to

occur via an initial demethoxylation of the methoxy group that is released as formaldehyde to

generate protocatechuic acid; formaldehyde can be assimilated or dissimilated via

methylotrophic pathways (22). The capability to utilize different carbon sources, such as

methanol, ethanol, and multicarbon substrates· may explain why methylotrophs are so well-suited

to life in the phyllosphere (2,5,23) where metabolite availability changes dynamically (2).

Despite extensive characterizations of the microbial community in the phyllosphere as

well as recent advances in lanthanide-dependent metabolisms, the effect of lanthanides on

methylotrophic community composition in the phyllosphere remains unknown. To investigate

this, methylotrophs were isolated from the phyllosphere of soybean plants and selected for

growth on methanol in the presence and absence of lanthanides, and then grown on different

carbon substrates. Growth parameters were measured with and without the addition of lanthanum

(La^{3+}). These strains were found to be phylogenetically similar but phenotypically distinct, with

growth rates differentially affected by substrate identity and the presence or absence of

lanthanum.

Results

The soybean phyllosphere methylotrophic community is not obligately lanthanide-dependent

Methylotrophic bacterial strains were isolated from the soybean phyllosphere by selection

on minimal salts media with methanol as the sole carbon source. Previous studies have shown

that the presence of lanthanides allows for the isolation of novel methylotrophic strains (16,17);

thus, we also included lanthanum in our media. Of 344 total isolates, 158 strains were isolated

with the addition of La^{3+} and 186 strains without the addition of La^{3+} to the selection medium.

After isolating single colonies, all 344 strains were retested for growth on methanol in the

presence and absence of La^{3+} to determine if lanthanides were necessary for growth. All strains

were pink, showed growth on methanol in the presence and absence of La^{3+} in the growth

medium, and showed growth on succinate and were therefore all classified as PPFMs. Each

strain was assigned a unique SLI (**S**oybean **L**eaf **I**solate) number, which is used to reference

specific strains throughout this study (**Table 1**).

Methylotrophic communities from the soybean phyllosphere are phylogenetically similar

Chromosomal DNA was extracted from the isolates and *rpoB* regions were PCR

amplified and Sanger sequenced. Using NCBI BLAST, *rpoB* sequences from each isolate were

compared against the Joint Genome Institute's (JGI) Integrated Microbial Genomes (IMG)

database to putatively assign species-level taxonomic classifications to each isolate. All isolates

matched to *Methylobacterium* species within group B (24) , with the closest species being

Methylobacterium extorquens AM1, *Methylobacterium extorquens* PA1, *Methylobacterium*

extorquens TK001, and *Methylobacterium zatmanii* 135. Untrimmed *rpoB* sequences from each

isolate, as well as from reference *Methylobacterium* and related species, were aligned using

MUSCLE, and Maximum-Likelihood trees with 100 bootstrap replications were constructed

using MEGA (**Figure 1A**). The *rpoB*-based trees show a high degree of relatedness amongst all

isolates, integrating SLI strains among other reference methylotroph strains within the

extorquens clade.

We sought to augment our understanding of the SLI community composition at the

taxonomic level by employing pairwise average nucleotide identity (ANI) comparisons of all

orthologous genes shared among all assembled SLI genomes (see **Table 1**) and representative

methylotrophic genomes using the IMG database. Results from the complete pairwise ANI

comparisons can be found in the **Table S1**, but have been condensed to include only SLI strains

and representative species from the *extorquens, aquaticum,* and *nodulans* clades in **Figure 1B**.

Based on ANI comparisons, all SLI strains were most similar to *Methylobacterium* species from

the *extorquens* clade (93-99% ANI) when compared to the *nodulans* (79% ANI) and *aquaticum*

clades (78% ANI), indicating that our community likely belongs to the *extorquens* clade. Based

on the 95% standard species cutoff (25), all SLI strains except for SLI 516, 575, and 576 are

genetically identical to the model organisms *Methylobacterium extorquens* AM1 and PA1.

Additionally, SLI 231, 233, 274, 285, 499, and 505 appear to be nearly 100% genetically

identical to each other; yet, as will be discussed in detail later, they exhibit distinct metabolic

differences (see **Table S2** for full comparison of genomes).

Whole-genome sequencing of representative SLI strains reveals new* extorquens *strains

The genomes of SLI strains 223, 231, 233, 274, 285, 384, 499, 505, 516, 575, 576 were

sequenced through the Joint Genome Institute and annotated using the IMG Annotation Pipeline

v.5.0.22. Summaries of the genome information for each strain compared to *M. extorquens* AM1

and PA1 are represented in **Table 1**. Notably, all strains but SLI 223 encoded at least one

plasmid in addition to the chromosome, with most strains (SLI 231, 233, 274, 285, 499, 505,

516, 575, 576) encoding at least three additional plasmids. The size, % GC, and number of genes

per chromosome and per plasmid were very similar for SLI 231, 233, 274, 285, 499, and 505 and

for SLI 516, 575, 576; this underscores the high degree of ANI between the strains (**Figure 1B**).

SLI 516, 575, and 576 – which are the least similar by ANI to *M. extorquens* AM1 and PA1

(**Figure 1B**) – also have the smallest chromosome sizes despite having the broadest metabolic

capabilities (**Figure 2A**). None of the strains encoded the megaplasmid (1261460 bp) found in

M. extorquens AM1, and the plasmid sizes of the SLI strains were very different from the

plasmid sizes of *M. extorquens* AM1.

Predictions about metabolic potential can be made from genomic analysis. All genomes

encode tetrahydromethanopterin and tetrahydrofolate-dependent pathways for C_1 oxidation and

marker genes for the serine cycle for C_1 assimilation, indicating that all strains are Type II

methylotrophs (6). All genomes encode four formate dehydrogenases, save for SLI 516, 575, and

576 which lack formate dehydrogenase 1. All genomes also encode the *N*-methylglutamate

pathway for methylamine utilization (*mgdDCBA*, *mgsABC*, *gmaS*) (22,26). Strains SLI 231, 233,

274, 285, 499, 505, 516, 575, and 576 encode a previously identified gene island that confers the

ability to grow on methoxylated aromatic acids (22) .

The SLI genomes can be grouped into two categories based on the presence or absence of

lanthanide-associated genes: group 1 consists of SLI 223, 231, 233, 274, 285, 384, 499, 505 and

group 2 consists of SLI 516, 575, and 576. Genomes from both groups encode calcium-

dependent methanol dehydrogenases and cognate cytochromes (*mxaFIG*) and lanthanide-

dependent methanol dehydrogenases and cognate cytochromes (*xoxFG*) indicating that all strains

are capable of lanthanide-independent and lanthanide-dependent methanol metabolism. Yet,

genes for the lanthanide-dependent alcohol dehydrogenase and cognate cytochrome (*exaFG*) are

lacking in group 2 genomes. In addition, group 2 genomes lack the gene for the lanthanide-

binding protein, lanmodulin, despite encoding the rest of the genes involved in the lanthanide

utilization and transport cluster (27). Thus, the influence of lanthanides on the metabolisms of

group 2 strains is of particular interest. The differences between groups 1 and 2 is also reflected

in the average nucleotide identities in **Figure 1B**. A complete summary of all metabolic and

lanthanide-related genes found in the SLI strains is included in **Table S2**.

Methylotroph isolates possess broad metabolic capabilities independent of lanthanide

availability

Although phyllosphere isolates were phylogenetically similar to domesticated research

strains such as *M. extorquens* AM1 and PA1, we hypothesized that environmental methylotrophs

might possess expanded substrate repertoires and broader metabolic capabilities. To test our

hypothesis, all isolates were screened for growth on C_1 substrates (methanol, methylamine,

dimethylsulfide), C_2 substrates (ethanol, oxalate), sugars (fructose, glucose), aromatic acids

(vanillic acid), and complex insoluble substrates (lignin, cellulose, tyrosine) **(Table 2)**.

Lanthanides were also added to growth media for each substrate, but this addition did not unlock

novel substrate utilization capabilities for any SLI tested. All strains exhibited growth on

methanol, methylamine, and ethanol. No strains grew on dimethylsulfide, oxalate, glucose,

lignin, cellulose, or tyrosine. Notably, 10 out of 344 strains exhibited moderate to substantial

growth on fructose and 78 out of 344 strains grew on vanillic acid.

The subset of SLI strains with completely sequenced genomes (SLI 223, 231, 233, 274,

285, 384, 499, 505, 516, 575, 576) that were shown to have broad metabolic capabilities from the

screen described above were spotted onto solid media containing five different carbon sources

without La^{3+}, with growth of *M. extorquens* PA1 serving as a comparison **(Figure 2A),** to

visualize relative differences in metabolic capabilities. From this subset, all grew on methanol,

confirming the initial growth phenotype, and all strains also grew on ethanol and methylamine.

SLI 231, 233, 499, 505, and 575 grew on vanillic acid. Isolates that grew on vanillic acid also

grew more robustly on methylamine, as evidenced by larger and more opaque colony

phenotypes. SLI 516, 575, and 576 were the only strains of this subset capable of growth on

fructose. Interestingly, these three strains also exhibited biofilm formation during growth in

liquid cultures (**Figure 2B, C**) and had heterogeneous colony morphology compared to all other

strains (**Figure 2A**). This initial screen revealed that aromatic acid catabolism (22) and fructose

utilization (21) are more widespread within the *extorquens* clade than previously assumed.

Addition of lanthanides influences the catabolism of alcohols

Based on our initial screen for substrate utilization on solid media, six representative

isolates (SLI 231, 233, 384, 499, 505, 575) were chosen as a test cohort for downstream growth

analyses because they displayed expanded metabolic capabilities compared to model organisms

M. extorquens AM1 or PA1, which cannot grow on sugars or aromatic acids. To further

investigate the impact of La^{3+} on this community, the growth rates and total biomass yields of

the test cohort strains were compared during growth on methanol and ethanol as lanthanides have

been shown to impact the oxidation of both alcohols in methylotrophic bacteria .

Growth rates (**Figure 3A, 3C**) and yields (**Figure 3B, 3D**) on 20 mM methanol and 34

mM ethanol, both with and without La^{3+} in the growth medium, were measured. Growth rates on

methanol were significantly higher in the presence of La^{3+} for 5 out of 6 strains (SLI 231, SLI

233, SLI 384, SLI 499, SLI 505; **Figure 3A**). Final yields were significantly lower in the

presence of La^{3+} for SLI 384 but significantly higher in the presence of La^{3+} for SLI 499 and SLI

505; thus, lanthanides can have variable effects on growth yields despite similar trends in growth

rates. Even amongst nearly identical strains encoding identical lanthanide-independent and -

dependent methanol dehydrogenases, there was significant variability in the final yields achieved

in the presence of La^{3+} between SLI 575 and SLI 231, SLI 233, SLI 499, and SLI 505 and in the

absence of La^{3+} between SLI 233 and SLI 575 (p-values for significance of all strains from one-

way ANOVA in **Table S3**).

Interestingly, growth rates on ethanol were higher in the presence of La^{3+} for SLI 231,

SLI 233, SLI 384, SLI 499, and SLI 505 but *not* SLI 575, the only strain in the test cohort

lacking ExaFG. Thus, we hypothesize that XoxFG must be acting as the primary ethanol

dehydrogenase during growth on ethanol with lanthanides in this strain (14). Although SLI 231,

SLI 233, SLI 384, SLI 499, and SLI 505 possess high degrees of genomic similarity (**Figure 1B**)

and all strains encode identical lanthanide-dependent enzymes, SLI 384 exhibits significantly

higher growth rates than the other strains in the presence of La^{3+} (p-values for significance of all

strains from one-way ANOVA in **Table S3**). Unlike the phenotypes with methanol (**Figure 3A-**

B), the presence of La^{3+} in the growth medium does not appear to have significant effects on the

final yields obtained, save for SLI 505 which exhibited significantly lower yields with La^{3+}.

For most strains, the presence of La^{3+} increases growth rate in methanol but the effect on

yields during growth on methanol and ethanol appear to be strain-dependent. Addition of

lanthanides did not result in a universal phenotypic change during growth on alcohols despite

most strains possessing nearly identical lanthanide-dependent enzymes. This raises interesting

hypotheses about the differential regulation of lanthanide-dependent enzymes and metabolisms

in methylotrophs and the role of lanthanide-dependent alcohol dehydrogenases during growth on

ethanol specifically.

SLI strains lacking lanthanide-associated genes reveal insights into lanthanide physiology

Genomic analysis identified several genomes (SLI 516, 575, 576) that lack lanmodulin

(**Table S2**) and do not have significant differences during growth on methanol or ethanol in the

presence of lanthanides (**Figure 3**). Lanmodulin is one of the most abundant proteins in the

phyllosphere (23) and can bind to lanthanides with very high affinity and selectivity (28) yet no

physiological role for lanmodulin has been identified; thus, the absence of lanmodulin in a subset

of SLI genomes from the phyllosphere was intriguing. To investigate the role of lanmodulin in

our SLI strains, the gene *lanM* (MexAM1_META1p1786) was cloned into an IPTG-inducible

expression vector and transformed into electrocompetent SLI 575. To replicate lanthanide

conditions in the phyllosphere – where lanthanides can exist in soluble or poorly soluble forms at

varying concentrations – lanmodulin expression was phenotyped during growth on alcohols in

the presence of excess (2 µM) and limiting (50 nM) concentrations of $LaCl_3$ and La_2O_3 (**Figure**

4A, B).

SLI 575 also lacks ExaF, so ethanol oxidation in excess lanthanide conditions must be

carried out by XoxF but could be facilitated by either MxaFI or XoxF in limiting lanthanide

concentrations (14,15). Therefore, we hypothesized that growth phenotypes of *lanM* expression

in these lanthanide conditions would be identical during growth on both methanol and ethanol.

During growth on methanol, expression of *lanM* had no effect in the presence of excess soluble

La^{3+} but resulted in an 8% significant decrease in growth rate with limiting soluble La^{3+} and 18-

24% significant increase in growth rate with excess and limiting poorly soluble La^{3+} (**Figure**

4A). In contrast, during growth on ethanol, significant differences are only observed during

growth on limiting soluble La^{3+} (8% increase) or excess poorly soluble La^{3+} (14% increase)

(**Figure 4B**) . Our preliminary results indicate a putative supplementary role for lanmodulin

during growth on poorly soluble lanthanide sources, but differences during growth on methanol

and ethanol require further investigation.

Robust aromatic acid utilization by SLI strains reveals novel lanthanide-related phenotypes

Genes that confer the ability to grow on methoxylated aromatic acids (ferulic acid,

vanillic acid, protocatechuic acid, *p*-hydroxybenzoic acid) have been recently identified in

methylotrophs primarily from the *aquaticum* and *nodulans* clade (22) . These genes exist as a

horizontally-transferred genetic island that is notably absent from the *extorquens* clade, save for

their presence in *Methylobacterium sp.* AMS5 (22) . Our screen isolated 78 strains capable of

growth on the methoxylated aromatic acid, vanillic acid. Of the SLI strains with assembled

genomes, those capable of growth on vanillic acid encode gene islands identical to each other

(**Table S2**). Aromatic acid gene islands in SLI strains include all of the genes found in the

aromatic acid gene island in *Methylobacterium sp.* AMS5 although specific regulatory genes are

in a different order. To demonstrate the robustness of SLI growth on aromatic acids, SLI strains

from the test cohort were grown on a low (5 mM) and high (12 mM) concentration of vanillic

acid (22,29,30) in the presence and absence of LaCl$_3$ and their growth was compared to that of

Methylobacterium sp. AMS5 (**Figure 5**). SLI 384 was used as a control, as it does not encode

genes for vanillic acid catabolism and therefore does not grow on vanillic acid (**Figure 2**).

Two striking phenotypes emerge when comparing growth on different concentrations of

vanillic acid. First, all strains exhibit slower growth rates on high concentrations of vanillic acid

compared to low concentrations of vanillic acid (**Figure 5A, C**). Despite encoding nearly

identical aromatic acid gene islands, *Methylobacterium sp.* AMS5 exhibits poor growth on high

concentrations of vanillic acid compared to the SLI strains (**Figure 5C)** and there are significant

differences in growth between the SLI strains as well (p-values for significance of all strains

from one-way ANOVA in **Table S3**). Second, addition of lanthanides influences aromatic acid

utilization based on substrate concentration despite all SLI strains encoding identical aromatic

acid gene islands. SLI 505 has higher growth rates with lanthanides on 5 mM vanillic acid but

lower growth rates with lanthanides on 12 mM vanillic acid (**Figure 5A, C**). SLI 231 has lower

growth rates with lanthanides only during growth on 12 mM vanillic acid, a trend also seen in

SLI 499 although the differences are not significant (**Figure 5C**). Conversely, SLI 233 has

marginally higher growth rates during growth on 12 mM vanillic acid (**Figure 5C**). Lanthanides

variably affect final yields during growth on low concentrations of vanillic acid (**Figure 5B**), but

have little effect on yield during growth on high concentrations of vanillic acid (**Figure 5D**).

SLI strains encode identical methylamine oxidation genes but display different growth

phenotypes

All SLI genomes only encode genes for the *N*-methylglutamate pathway similar to what

is found in *M. extorquens* PA1 for methylamine utilization (19). To determine if the growth of

SLI strains on methylamine is influenced by lanthanides, SLI strains were grown in 15 mM

methylamine in the presence and absence of $LaCl_3$. Growth rates (**Figure 6A**) and final yields

(**Figure 6B**) are reported. Addition of $LaCl_3$ did not significantly impact the growth rate or yield

for any strain tested. Interestingly, SLI 575 had significantly lower yields than SLI 231, 233,

499, and 509, despite having a similar growth rate to all four strains; this trend was not

lanthanide-dependent. Also of note, SLI 384 had nearly triple the lag time and half the growth

rate and final yield as other SLI strains despite encoding identical *N*-methylglutamate pathway

genes. This suggests additional details about methylamine catabolism in SLI strains beyond the

N-methylglutamate pathway that result in the diminished growth phenotypes observed.

SLI strains expand fructose utilization in the extorquens clade

Most methylotrophs, including *M. extorquens* AM1 and PA1, encode all of the genes

necessary for sugar oxidation but lack sugar-specific transporters and assimilatory pathways

required for their utilization as a primary substrate (26). Notably, complete sugar catabolism has

only been demonstrated in endophytic *extorquens* clade members (21). Thus, it was striking to

isolate ten strains capable of robust growth on fructose as the sole carbon source (**Table 2**). Of

the strains with assembled genomes, SLI 516, SLI 575, and SLI 576 were capable of growth on

fructose. Interestingly, these three SLI strains also displayed the lowest sequence similarity to

other SLI strains and to *M. extorquens* AM1 and PA1 based on ANI (**Figure 1B**), hinting that the

differences in genomes might be due in part to unknown genes specific to sugar transport and

catabolism. To quantify growth on fructose, SLI 575 was grown on 25 mM fructose in the

presence and absence of LaCl$_3$ (**Figure 7**). Growth rates and yields in both conditions were

nearly identical. Growth rates on fructose were comparable to rates for methylamine and

substantially lower than rates for methanol or vanillic acid. To our knowledge, SLI 516, 575, and

576 are the first identified epiphytic *extorquens* strains capable of growth on a sugar substrate.

Discussion

Methylotrophs that inhabit the phyllosphere influence both the plants that they colonize

and global biogeochemical processes (1,4). Yet, studies investigating the physiology,

biodiversity, and ecology of methylotrophic communities of the phyllosphere have only emerged

within the last decade (3,31) and do not take into account the role of lanthanides in influencing

their community composition. Furthermore, despite the abundance of methylotrophs in the

phyllosphere, the bulk of phylogenomic studies on *Methylobacterium* species use genomes

isolated from non-phyllosphere environments (24). Here, we describe the first community-level

characterization of the role of lanthanides during the metabolisms of natural methylotrophic

populations in soybean plants, revealing novel insights into methylotrophic biodiversity,

metabolic capabilities, and lanthanide-dependent metabolisms.

Marker gene sequencing for assigning taxonomic groups has proven difficult in

Methylobacterium, as16s *r*RNA has poor phylogenetic resolution in *Methylobacterium* (24). We

characterized our soybean methylotroph community composition using *rpoB* (**Figure 1A**), a

highly polymorphic single-copy gene with sub-species resolution with *Methylobacterium* (24).

Our phylogenetic analysis revealed two insights: 1) all of our isolates are closely related to each

other (**Figure 1A**) and fall within the *extorquens* clade of *Methylobacterium* (**Figure 1A, B**), and

2) even amongst nearly identical strains (**Figure 1B**), there is broad diversity in terms of the

possession of lanthanide-related genes and metabolic capabilities, and that these differences are

reflected in how strains group based on *rpoB* sequences (**Figure 8**). Overall, our results

substantiate *rpoB* as a superior marker gene for phylogenetic analysis of *Methylobacterium*

species, and, importantly, highlight how phenotypic characterizations can reveal novel strains

that appear identical by *rpoB* sequence alone.

Crops, such as soybeans, can sequester lanthanides from the soil through repeated cycles

of harvest limiting lanthanide availability in the soil and affecting lanthanide availability for

methylotrophs in the phyllosphere. Thus, it is not surprising that our microbial isolation from

soybean plants did not reveal any novel lanthanide-dependent phenotypes despite previous

studies identifying organisms that have increased heterogeneity in the presence of lanthanides

(32) or that are strictly dependent on lanthanides for growth (17,18). The maintenance of both

calcium-dependent and lanthanide-dependent methanol dehydrogenase systems in our SLI

strains could suggest that 1) lanthanides are not always present or bioavailable in the

phyllosphere and therefore MxaFI remains essential for methylotrophic growth; 2) lanthanides

are present but extensively competed for, so calcium-dependent methanol oxidation systems

offer an alternative for methylotrophic growth; and/or 3) lanthanide-dependent enzymes have

additional non-metabolic or non-lanthanide-related functions that necessitate their maintenance.

These hypotheses could be examined further by repeating our isolations in non-domesticated

phyllosphere environments where lanthanides are more prevalent or bioavailable and where the

isolation of obligately lanthanide-dependent methylotrophs might be more likely.

Notably, of the test cohort reported here, SLI 575 does not encode ExaF or lanmodulin

and is the only strain that does not exhibit significant lanthanide-dependent growth phenotypes

on alcohols (**Figure 3**). As lanmodulin is reported to be one of the most highly expressed

peptides in the phyllosphere (23), has been proposed to function as a lanthanide biosensor, and

has been shown to bind lanthanides (28) it is possible that SLI 575 is lacking the sensor required

to coordinate lanthanide availability with metabolic processes and the role of lanmodulin in the

phyllosphere specifically remains unknown. A previous study in *Methylobacterium aquaticum*

22A, which natively encodes *lanM*, found that overexpression of *lanM* results in faster growth on

methanol with 20 nM $LaCl_3$ and slower growth with 100 nM $LaCl_3$ (33). In our SLI strain

lacking lanmodulin, we see a similar trend during growth on methanol where *lanM* expression *in*

trans decreases growth rates at higher lanthanide concentrations but increases growth rates at

lower lanthanide concentrations. However, our results using poorly soluble lanthanide

concentrations and ethanol as a substrate are novel, and additional experiments involving ICP-

MS are required to determine the effect of lanmodulin mutants on intracellular lanthanide

concentrations. Here, SLI 575 and related strains (SLI 516, 576) emerge as an interesting model

by which to interrogate the role of ExaF or lanmodulin in methylotrophs both in laboratory and

plant environments.

Previous studies have reported that the model organism *M. extorquens* PA1 has an

average growth rate of 0.042 h^{-1} and an average final yield of OD_{600} 0.712 on 15 mM

methylamine (19,20). While our SLI strains exhibit similar growth rates regardless of LaCl$_3$

addition (**Figure 6A**), the final yields are nearly half of what is achieved by PA1 (**Figure 6B**);

this suggests metabolic bottlenecks towards efficient methylamine assimilation that are not

immediately obvious from a preliminary genomic analysis of C1 assimilation genes. One

possibility is that methylamine that is incorporated via the N-methylglutamate pathway

preferentially serves as a nitrogen source rather than a carbon source (19,20) but future studies

are required to validate this hypothesis in our SLI strains.

Although the ability to utilize methanol as a substrate has provided a competitive

advantage for methylotrophs in the phyllosphere, sugars and aromatic acid are abundant enough

to support the growth of diverse microorganisms (2,22,34). It is therefore advantageous for

microbes to be able to utilize both multicarbon sources, such as fructose and vanillic acid, and

single carbon sources like methanol. Detailed investigations into methylotrophic aromatic acid

catabolism in the presence and absence of lanthanides is necessary to understand how this

metabolism functions in carbon cycling of lignin by-products in natural environments, as well as

identify additional roles for lanthanides in the metabolism of non-alcohol substrates. The

decrease in growth rates in the presence of lanthanides only during growth on high

concentrations of aromatic acid is surprising, considering that all known lanthanide-dependent

enzymes are in the periplasm yet all enzymes used for aromatic acid utilization are in the

cytoplasm; the role of lanthanides during this metabolism is an ongoing area of research. The

substantial yield but decreased growth rate achieved by SLI strains on high concentrations vs low

concentrations of vanillic acid demonstrates that the ability to grow on methoxylated aromatic

acids is widely distributed among members of the *extorquens* clade, more robust than what has

been reported in other model strains, and could reveal novel insights about lanthanide

metabolism.

Genomic analyses for fructose transporters or assimilatory pathways in SLI 516, 575, or

576 (SLI strains with assembled genomes capable of growth on fructose) identified genes

encoding for phosphofructokinase that were absent in all other SLI strains. Phosphofructokinase

is a key regulatory enzyme of glycolysis, responsible for the phosphorylation of fructose to

fructose-6-phosphate. The end-product of this reaction is also an intermediate of the assimilatory

RuMP cycle; however, none of our SLI strains employ the RuMP cycle for assimilation.

Taken together, the methylotrophic community in the phyllosphere of soybeans is

metabolically diverse and heavily influenced by lanthanide availability when consuming

methanol. We hypothesize that the capacity to expand substrate repertoires might also be

important drivers of efficient colonization. Whether the selection for low species diversity is at

the level of the plant or arises through natural selection within methylotrophic communities, this

finding points to the evolution of niche partitioning strategies among a single species in order to

maximize resources in shared habitats (31,35,36). Results from this study pave the way for future

work to investigate 1) expanded metabolic capabilities in *extorquens* clade methylotrophs, 2)

tradeoffs associated with lanthanide-dependent and -independent alcohol metabolism, and 3)

niche partitioning among genetically identical strains during plant colonization.

Materials & Methods

Methylotroph isolation

Environmental methylotrophic bacterial strains were isolated from soybean plants

(*Glycine max*) growing on the Michigan State University Agronomy Farm (East Lansing,

Michigan, U.S.A. 42.6908, -84.4866). Four leaves were harvested from four different plants on

September 7, 2018, and each leaf was placed into a sterile 50 mL conical tube. 50 ml of sterile 50

mM phosphate buffer (pH 7.3) was added to each tube, the tubes were vortexed for 5 minutes,

and 100 μL of buffer was spread onto solidified (1.5% wt/vol agar) MP minimal salts media

(37). The isolation medium contained 125 mM methanol, RPMI 1640 Vitamins Solution (Sigma

Aldrich, St. Louis, MO, USA) diluted to 1X, and 50 μg/mL cycloheximide. Two sets of isolation

media were prepared; one with and one without 2 μM $LaCl_3$. Isolation plates were grown at 30

°C for several days until bacterial colonies were visible. Pink colonies were selected from the

plates and struck out for isolated colonies onto solidified media of the same composition to

confirm growth. Single colonies of isolated strains were then inoculated into 650 μL of MP

medium with 125 mM methanol and 1X RPMI 1640 Vitamins Solution in a 48 well microplate.

All isolated strains were inoculated into medium with the same concentration of La^{3+} as they

were isolated with, and then incubated shaking for 48 hours at 200 rpm on an Innova 2300

platform shaker (Eppendorf, Hamburg, Germany) at 30°C. Strains were frozen for future use by

adding 25 μL of sterile dimethyl sulfoxide to 500 μL of late-exponential phase culture in a sterile

screw-cap vial, flash-freezing in liquid nitrogen, and storing at -80°C.

Strains cultivation

Cultures used for growing and maintaining strains, extracting DNA, and subculturing for

growth phenotypic analyses were prepared as follows: Freezer stocks of each strain were struck

out onto MP agar supplemented with 15 mM succinate and 1X RPMI 1640 Vitamins Solution

and incubated at 30°C until single colonies emerged (2-4 days). Single colonies of each strain

were used to inoculate 3 mL cultures of MP with 15 mM succinate and 1X RPMI 1640 Vitamins

Solution and grown overnight in round-bottom glass culture tubes (ThermoFisher Scientific,

Waltham, MA, USA) at 30°C, shaking at 200 rpm on an Innova S44i shaker (Eppendorf,

Hamburg, Germany) to an OD_{600} of approximately 1.5. See sections below for downstream

analyses performed on these cultures.

Chromosomal DNA extraction for whole-genome and marker gene sequencing

483   Genomic DNA was extracted from 3 mL overnight cultures of each strain (see ***Strains***

Cultivation) using the DNeasy PowerSoil Pro Kit from Qiagen (Hilden, Germany) per

manufacturer's protocol. Genomic DNA was quantified using a Take3 Microvolume Plate

Spectrophotometer (BioTek, Winooski, VT, USA). Whole-genome sequencing via PacBio

Sequel II was performed by the Department of Energy's Joint Genome Institute (JGI, Walnut

Creek, CA, USA). Genomes can be accessed through JGI Integrated Microbial Genomes &

Microbiomes (IMG) (see **Table 1** for IMG Genome IDs). Marker gene sequencing was

performed by PCR amplifying regions of *16s* rRNA or *rpoB* using universal primers[18] (16s_fwd:

5'GAGTTTGATCCTGGCTCA3', 16s_rev: 5'TACCTTGTTACGACTT3'; rpoB_fwd:

5'AAGGACATCAAGGAGCAGGA3', rpoB_rev: 5'ACSCGGTAKATGTCGAACAG3'),

confirming products via gel electrophoresis on a 1% agarose gel, purifying PCR products using

the GeneJET PCR Purification Kit (ThermoFisher Scientific, Waltham, MA, USA) , and Sanger

sequencing the PCR products. Untrimmed sequences were aligned using MAFFT and MUSCLE,

and a Maximum-Likelihood tree using a Tamura-Nei model and 100 bootstrap replications was

constructed using MEGA. Average nucleotide identity scores were calculated through JGI IMG

interface for pairwise genome comparisons using whole-genome sequences from the IMG

database as inputs.

Growth phenotypic analyses

Isolate strains were cultivated as described in ***Strains Cultivation*** and were screened for

their ability to grow on diverse substrates in the presence and absence of La^{3+}. To screen for the

ability to grow on diverse substrates, overnight cultures of each strain were washed twice in MP

media with no carbon source at 2000 x g for 10 minutes, resuspended in MP media to an OD600

of 0.1, and spotted (10 uL) onto MP agar plates supplemented with 1X 1640 RPMI Vitamins

Solution, and either a soluble substrate (20 mM methanol, 34 mM ethanol, 15 mM methylamine,

5 mM potassium oxalate, 25 mM glucose, 25 mM fructose) or an insoluble substrate (5 mM

vanillic acid); the insoluble substrates were added as non-sterilized powders directly to the

melted agar prior to pouring the plate. Agar plates were incubated at 30C for 2-4 days or until

visible colonies formed.

To screen for La^{3+} effects during growth on methanol, overnight cultures of each strain

were washed twice in MP media with no carbon source at 2000 x g for 10 minutes and

inoculated at an OD600 of 0.1 into a transparent 96-well plate (Corning, Corning, NY, USA)

containing 200 uL of MP media supplemented with 125 mM methanol, 1X RPMI 1640 Vitamins

Solution, and +/- 2 μM $LaCl_3$ for endpoint growth assays. Strains with demonstrated growth on

solid media of different substrates were additionally phenotyped in liquid media in the presence

and absence of La^{3+}. Overnight cultures of strains of interest were washed twice in MP media

with no carbon source at 2000 x g for 10 minutes and inoculated at an OD600 of 0.1 into a

transparent 48-well plate (Corning, Corning, NY, USA) containing 650 uL of MP media

supplemented with 1X 1640 RPMI Vitamins Solution, +/- 2 μM $LaCl_3$, and appropriate carbon

source (20 mM methanol, 15 mM methylamine, 34 mM ethanol, 12 mM vanillic acid, 25 mM

fructose). All endpoint assays and growth curves were carried out at 30°C and 548 rpm using a

Synergy HTX plate reader (BioTek, Winooski, VT, USA) with OD600 readings measured every

30 minutes. Data was analyzed using Microsoft Excel. 2-4 replicates were run for each strain in

each condition, and at least 10 exponential-phase data points were used for linear regression

analysis to calculate the growth rates of each strain. Paired t-tests were performed to identify

statistically significant ($p < 0.05$) differences between growth in the presence or absence of

$LaCl_3$ for each strain. One-way ANOVA with post-hoc Tukey HSD was performed to identify

statistically significant ($p < 0.05$) differences among all strains in the presence or absence of

$LaCl_3$..

Design and construction of lanM expression vector

The expression vector encoding the IPTG-inducible promoter $P_{L/04/A1}$ and mCherry (38)

was modified to include the ribosome binding site (agggagagaccccga) for *fae*

(MexAM1_META1p1766) and a KpnI cut site between the XbaI and HindIII cut sites natively

on the plasmid. Briefly, complementary single-stranded oligos were synthesized encoding an

XbaI cut site, the entire sequence of the ribosome binding site, KpnI cut site sequence, and

HindIII cut site sequence. Oligos were annealed to each other to generate a double-stranded

DNA fragment, which underwent a restriction enzyme digest with XbaI (New England Biolabs,

Ipswich, MT, USA) and HindIII (New England Biolabs, Ipswich, MT, USA) at 37°C for one

hour. The expression vector was also digested with XbaI and HindIII under the same conditions

to remove the gene for mCherry. The digested DNA fragment and vector were ligated using T4

DNA ligase (New England Biolabs, Ipswich, MA, USA) at room temperature for 10 minutes and

electroporated into 10β *E. coli* competent cells (New England Biolabs, Ipswich, MA, USA).

Modified plasmids encoding a ribosome-binding site were verified by colony PCR and full

plasmid sequencing (Primordium Labs, South San Francisco, CA, USA). The gene for

lanmodulin (*lanM,* MexAM1_META1p1786) was amplified by PCR with primers including

regions overlapping the expression vector between the ribosome-binding site and HindIII site.

The expression vector was linearized by PCR with primers including regions overlapping with

20 bp of the start and end of *lanM* and treated with DpnI at 37°C for one hour. NEBuilder HiFi

DNA Assembly (New England Biolabs, Ipswich, MA, USA) was used per manufacturer's

protocols to assemble the expression plasmid encoding *lanM*. The assembly was electroporated

into 10β *E. coli* competent cells (New England Biolabs, Ipswich, MA, USA) . The *lanM*

expression vector was verified by colony PCR and full plasmid sequencing (Primordium Labs,

South San Francisco, CA, USA) and electroporated into SLI 575 at 2500 V with an outgrowth in

Nutrient Broth for approximately 12 hours, and SLI 575 containing the *lanM* expression vector

was selected for on minimal media plates supplemented with succinate and 50 µg/mL

kanamycin. All growth experiments with SLI 575/ *lanM* were carried out in media supplemented

with 50 µg/mL kanamycin and, if induced, 1 mM IPTG.

Acknowledgements

We thank Allison Hunt and Zachary Jensen—previous members of the Martinez-Gomez Lab—

for their invaluable efforts during the early stages of this project. We thank members of Tobias

Erb's lab at the Max Planck Institute for Terrestrial Microbiology for generously providing an

IPTG-inducible expression vector. We would also like to acknowledge Jean-Baptiste Leducq,

Christopher Marx, Alexander Alleman, and all members of the Marx Lab at the University of

Idaho for their thoughtful comments and suggestions during experimental planning and

manuscript drafting. The information, data, or work presented herein was funded by the United

States Department of Energy, Office of Science, Office of Biological and Environmental

Research, under Award Number DE-SC0022318 subaward SH5849-772894; by the National

Science Foundation under Grant 2142154; by the National Science Foundation under Grant

2127732. A.G. was supported in part by the National Institutes of Health Genetic Dissection of

Cells and Organisms Training Grant 1T32GM132022-01.

Competing Interests

The authors declare no competing financial interests.

Data Availability

The data generated and/or analyzed during the current study as well as stocks of novel reported

strains are available from the corresponding author upon reasonable request. Genomes from

novel strains reported in this study can be found through JGI IMG using the IMG Genome IDs

indicated in **Table 1**.

References

1. Vacher C, Hampe A, Porté AJ, Sauer U, Compant S, Morris CE. The Phyllosphere: Microbial

Jungle at the Plant–Climate Interface. Annu Rev Ecol Evol Syst. 2016 Nov 1;47(1):1–24.

2. Lindow SE, Brandl MT. Microbiology of the phyllosphere. Appl Environ Microbiol. 2003

Apr;69(4):1875–83.

3. Vorholt JA. Microbial life in the phyllosphere. Nat Rev Microbiol. 2012 Dec 1;10(12):828–40.

4. Yurimoto H, Shiraishi K, Sakai Y. Physiology of Methylotrophs Living in the Phyllosphere.

Microorganisms. 2021;9(4).

5. Delmotte N, Knief C, Chaffron S, Innerebner G, Roschitzki B, Schlapbach R, et al. Community

proteogenomics reveals insights into the physiology of phyllosphere bacteria. Proc Natl Acad Sci.

2009;106(38):16428–33.

6. Chistoserdova L, Lidstrom ME. Aerobic Methylotrophic Prokaryotes. In: Rosenberg E, DeLong

EF, Lory S, Stackebrandt E, Thompson F, editors. The Prokaryotes: Prokaryotic Physiology and

Biochemistry [Internet]. Berlin, Heidelberg: Springer Berlin Heidelberg; 2013. p. 267–85.

Available from: https://doi.org/10.1007/978-3-642-30141-4_68

7. Galbally IE, Kirstine W. The Production of Methanol by Flowering Plants and the Global Cycle

of Methanol. J Atmospheric Chem. 2002 Nov 1;43(3):195–229.

8. Keltjens JT, Pol A, Reimann J, Op den Camp HJM. PQQ-dependent methanol dehydrogenases:

rare-earth elements make a difference. Appl Microbiol Biotechnol. 2014 Jul 1;98(14):6163–83.

9. Good NM, Moore RS, Suriano CJ, Martinez-Gomez NC. Contrasting in vitro and in vivo methanol

oxidation activities of lanthanide-dependent alcohol dehydrogenases XoxF1 and ExaF from

Methylobacterium extorquens AM1. Sci Rep. 2019 Mar 12;9(1):4248.

10. Nakagawa T, Mitsui R, Tani A, Sasa K, Tashiro S, Iwama T, et al. A Catalytic Role of XoxF1 as

La3+-Dependent Methanol Dehydrogenase in *Methylobacterium extorquens* Strain AM1. PLOS

ONE. 2012 Nov 27;7(11):e50480.

11. Huang J, Yu Z, Groom J, Cheng JF, Tarver A, Yoshikuni Y, et al. Rare earth element alcohol

dehydrogenases widely occur among globally distributed, numerically abundant and

environmentally important microbes. ISME J. 2019 Aug 1;13(8):2005–17.

12. Taubert M, Grob C, Howat AM, Burns OJ, Dixon JL, Chen Y, et al. XoxF encoding an alternative

methanol dehydrogenase is widespread in coastal marine environments. Environ Microbiol. 2015

Oct;17(10):3937–48.

13. Knief C, Delmotte N, Chaffron S, Stark M, Innerebner G, Wassmann R, et al. Metaproteogenomic

analysis of microbial communities in the phyllosphere and rhizosphere of rice. ISME J. 2012 Jul

1;6(7):1378–90.

14. Good NM, Vu HN, Suriano CJ, Subuyuj GA, Skovran E, Martinez-Gomez NC. Pyrroloquinoline

Quinone Ethanol Dehydrogenase in *Methylobacterium extorquens* AM1 Extends Lanthanide-

Dependent Metabolism to Multicarbon Substrates. J Bacteriol. 2016 Nov 15;198(22):3109–18.

15. Vu Huong N., Subuyuj Gabriel A., Vijayakumar Srividhya, Good Nathan M., Martinez-Gomez

N. Cecilia, Skovran Elizabeth. Lanthanide-Dependent Regulation of Methanol Oxidation Systems

in *Methylobacterium extorquens* AM1 and Their Contribution to Methanol Growth. J Bacteriol.

2016 Mar 31;198(8):1250–9.

16. Wegner Carl-Eric, Gorniak Linda, Riedel Stefan, Westermann Martin, Küsel Kirsten.

Lanthanide-Dependent Methylotrophs of the Family Beijerinckiaceae: Physiological and Genomic

Insights. Appl Environ Microbiol. 2019 Dec 13;86(1):e01830-19.

17. Howat AM, Vollmers J, Taubert M, Grob C, Dixon JL, Todd JD, et al. Comparative Genomics

and Mutational Analysis Reveals a Novel XoxF-Utilizing Methylotroph in the Roseobacter Group

Isolated From the Marine Environment. Front Microbiol [Internet]. 2018;9. Available from:

https://www.frontiersin.org/articles/10.3389/fmicb.2018.00766

18. Wang L, Suganuma S, Hibino A, Mitsui R, Tani A, Matsumoto T, et al. Lanthanide-dependent

methanol dehydrogenase from the legume symbiotic nitrogen-fixing bacterium *Bradyrhizobium*

diazoefficiens strain USDA110. Enzyme Microb Technol. 2019 Nov;130:109371.

19. Nayak DD, Marx CJ. Methylamine utilization via the N-methylglutamate pathway in

Methylobacterium extorquens PA1 involves a novel flow of carbon through C1 assimilation

and dissimilation pathways. J Bacteriol. 2014 Dec;196(23):4130–9.

20. Nayak DD, Marx CJ. Experimental Horizontal Gene Transfer of Methylamine Dehydrogenase

Mimics Prevalent Exchange in Nature and Overcomes the Methylamine Growth

Constraints Posed by the Sub-Optimal N-Methylglutamate Pathway. Microorganisms. 2015 Mar

10;3(1):60–79.

21. Van Aken B, Peres CM, Doty SL, Yoon JM, Schnoor JL. *Methylobacterium populi* sp. nov., a

novel aerobic, pink-pigmented, facultatively methylotrophic, methane-utilizing bacterium isolated

from poplar trees (Populus deltoides×nigra DN34). Vol. 54, International Journal of Systematic

and Evolutionary Microbiology,. Microbiology Society,; 2004. p. 1191–6.

22. Lee JA, Stolyar S, Marx CJ. Aerobic Methoxydotrophy: Growth on Methoxylated Aromatic

Compounds by Methylobacteriaceae. Front Microbiol [Internet]. 2022;13. Available from:

https://www.frontiersin.org/article/10.3389/fmicb.2022.849573

23. Ochsner AM, Hemmerle L, Vonderach T, Nüssli R, Bortfeld-Miller M, Hattendorf B, et al. Use

of rare-earth elements in the phyllosphere colonizer *Methylobacterium extorquens* PA1. Mol

Microbiol. 2019 May 1;111(5):1152–66.

24. Leducq JB, Sneddon D, Santos M, Condrain-Morel D, Bourret G, Martinez-Gomez NC, et al.

Comprehensive Phylogenomics of *Methylobacterium* Reveals Four Evolutionary Distinct Groups

and Underappreciated Phyllosphere Diversity. Genome Biol Evol. 2022 Aug 3;14(8).

25. Figueras MJ, Beaz-Hidalgo R, Hossain MJ, Liles MR. Taxonomic affiliation of new genomes

should be verified using average nucleotide identity and multilocus phylogenetic analysis. Genome

Announc. 2014 Dec 4;2(6).

26. Chistoserdova L, Kalyuzhnaya MG, Lidstrom ME. The expanding world of methylotrophic

metabolism. Annu Rev Microbiol. 2009;63:477–99.

27. Roszczenko-Jasińska P, Vu HN, Subuyuj GA, Crisostomo RV, Cai J, Lien NF, et al. Gene

products and processes contributing to lanthanide homeostasis and methanol metabolism in

Methylorubrum extorquens AM1. Sci Rep. 2020 Jul 29;10(1):12663.

28. Cotruvo JA Jr, Featherston ER, Mattocks JA, Ho JV, Laremore TN. Lanmodulin: A Highly

Selective Lanthanide-Binding Protein from a Lanthanide-Utilizing Bacterium. J Am Chem Soc.

2018 Nov 7;140(44):15056–61.

29. Varman AM, He L, Follenfant R, Wu W, Wemmer S, Wrobel SA, et al. Decoding how a soil

bacterium extracts building blocks and metabolic energy from ligninolysis provides road map for

lignin valorization. Proc Natl Acad Sci. 2016 Oct 4;113(40):E5802.

30. Kamimura N, Takahashi K, Mori K, Araki T, Fujita M, Higuchi Y, et al. Bacterial catabolism of

lignin-derived aromatics: New findings in a recent decade: Update on bacterial lignin catabolism.

Environ Microbiol Rep. 2017 Dec 1;9(6):679–705.

31. Schäfer M, Vogel CM, Bortfeld-Miller M, Mittelviefhaus M, Vorholt JA. Mapping phyllosphere

microbiota interactions in planta to establish genotype–phenotype relationships. Nat Microbiol.

2022 Jun 1;7(6):856–67.

32. Skovran E, Martinez-Gomez NC. Just add lanthanides. Science. 2015 May 22;348(6237):862–3.

33. Fujitani Y, Shibata T, Tani A. A Periplasmic Lanthanide Mediator, Lanmodulin, in

Methylobacterium aquaticum Strain 22A. Front Microbiol [Internet]. 2022;13. Available from:

https://www.frontiersin.org/articles/10.3389/fmicb.2022.921636

34. Welte CU, de Graaf R, Dalcin Martins P, Jansen RS, Jetten MSM, Kurth JM. A novel

methoxydotrophic metabolism discovered in the hyperthermophilic archaeon *Archaeoglobus*

fulgidus. Environ Microbiol. 2021 Jul 1;23(7):4017–33.

35. Schlechter RO, Kear EJ, Bernach M, Remus DM, Remus-Emsermann MNP. Metabolic resource

overlap impacts competition among phyllosphere bacteria. ISME J. 2023 Sep 1;17(9):1445–54.

36. Schäfer M, Pacheco AR, Künzler R, Bortfeld-Miller M, Field CM, Vayena E, et al. Metabolic

interaction models recapitulate leaf microbiota ecology. Science. 381(6653):eadf5121.

37. Delaney NF, Kaczmarek ME, Ward LM, Swanson PK, Lee MC, Marx CJ. Development of an

Optimized Medium, Strain and High-Throughput Culturing Methods for *Methylobacterium*

extorquens. PLOS ONE. 2013 Apr 30;8(4):e62957.

38. Carrillo M, Wagner M, Petit F, Dransfeld A, Becker A, Erb TJ. Design and Control of

Extrachromosomal Elements in *Methylorubrum extorquens* AM1. ACS Synth Biol. 2019 Nov

15;8(11):2451–6.

Table 1. Summary of SLI strain genomes. Whole-genome sequencing performed by Joint

Genome Institute and annotations obtained using IMG Annotation Pipeline v.5.0.22

Strain Information			Chromosome			Plasmid		
Strain	**Proposed Name**	**IMG Genome ID**	**Size (bp)**	**% GC**	**No. of genes**	**Size (bp)**	**% GC**	**No. of genes**
M. extorquens AM1	-	644736386	5511322	68.71	5029	1261460 44195 37858 24943	67.65 67.93 65.25 66.94	1168 33 34 30
M. extorquens PA1	-	641228497	5471154	68.18	4939	-	-	-
SLI 223	*M. extorquens* SLI 223	2917523761	5504362	68.23	5222	-	-	-
SLI 231	*M. extorquens* SLI 231	2918956041	5373034	68.29	5165	166358 27292 29644	67.47 61.51 58.78	176 25 27
SLI 233	*M. extorquens* SLI 233	2932223180	5373474	68.29	5287	166399 27296 29645	67.47 61.51 58.79	181 28 27
SLI 274	*M. extorquens* SLI 274	2947147188	5374052	68.29	5202	166386 27292 29646	67.47 61.51 58.79	178 26 27
SLI 285	*M. extorquens* SLI 285	2932228704	5375017	68.30	5295	166399 27292 29655	67.47 61.51 58.78	182 25 28
SLI 384	*M. extorquens* SLI 384	2918961435	5504365	68.23	5233	28462	67.45	48
SLI 499	*M. extorquens* SLI 499	2947152626	5373409	68.29	5218	166383 27292 29645	67.47 61.51 58.79	178 25 27
SLI 505	*M. extorquens* SLI 505	2918966707	5372985	68.29	5170	166379 27292 29645	67.47 61.51 58.79	176 26 27
SLI 516	*M. extorquens* SLI 516	2918972107	5009907	68.66	4762	199519 28884 20194	67.26 66.85 60.93	215 39 30
SLI 575	*M. extorquens* SLI 575	2918977154	5009893	68.66	4764	199519 28890 20194	67.26 66.84 60.93	214 39 30
SLI 576	*M. extorquens* SLI 576	2918982202	5009856	68.66	4762	199518 28890 20194	67.26 66.84 60.93	215 39 31

Figure 1. Phylogenetic and taxonomic characterization of a methylotrophic community isolated

from the soybean phyllosphere. (**A**) Phylogenetic tree based on *rpoB* sequences. Untrimmed

rpoB sequences from each isolate as well as from reference *Methylobacterium* and related

species were obtained from IMG, aligned using MUSCLE, and the Maximum-Likelihood trees

were constructed with 100 bootstrap replications using MEGA. Groupings indicate clade identity

for each strain (**B**) Average nucleotide identity of all SLI strains compared to representative

species from the *extorquens, nodulans,* and *aquaticum* clades. All pairwise comparison ANIs

found in **Table S1**

Table 2. Distribution of substrate-utilization capabilities of SLI community. All 344 isolates

were grown in liquid minimal salts media containing either C_1 (20 mM methanol, 15 mM

methylamine, 1 mM dimethylsulfide), C_2 (34 mM ethanol, 5 mM potassium oxalate), sugar (25

mM fructose, 25 mM glucose), or complex insoluble substrates (vanillic acid, cellulose, lignin,

tyrosine)

	Substrate	No. of SLI strains
C_1 Substrates	Methanol	344
	Methylamine	344
	Dimethylsulfide	0
C_2 Substrates	Ethanol	344
	Potassium Oxalate	0
Sugars	Glucose	0
	Fructose	10
Insoluble C_{2+} Substrates	Vanillic Acid	78
	Lignin	0
	Cellulose	0
	Tyrosine	0

Figure 2. (A) Spotting of a subset of SLI strains on various substrates to demonstrate differential

metabolic capabilities of SLI test cohort. All strains were spotted on minimal agar supplemented

with either 20 mM methanol, 34 mM methanol, 15 mM methylamine, 5 mM vanillic acid, 25

mM fructose, or no carbon. *Methylobacterium extorquens* PA1Δ*cel* as a comparison of known

substrate utilization capabilities in a model *extorquens* clade strain. Biofilm formation of SLI 575

in liquid culture (**B**) and in microplates (**C**)

Figure 3. Growth phenotypes of SLI test cohort on alcohols in the presence and absence of La^{3+}.
(**A**) Growth rates on 20 mM methanol +/- 2 μM LaCl$_3$ (**B**) Final yields on 20 mM methanol +/- 2 μM LaCl$_3$ (**C**) Growth rates on 34 mM ethanol +/-2 μM LaCl$_3$ (**D**) Final yields on 34 mM ethanol +/- 2 μM LaCl$_3$. Black bars (bottom) represent cultures grown in the absence of LaCl$_3$; hatched bars (top) represent cultures grown in the presence of LaCl$_3$. N=3. Error bars represent standard deviation. Significant differences for each strain +/- LaCl$_3$ determined by Student's paired t-test (*, $p<0.05$; **, $p<0.01$). Significant differences between all strains +/- LaCl$_3$ determined using one-way ANOVA with post-hoc Tukey HSD (**Table S3**)

Figure 4. Growth phenotypes of lanmodulin expression in SLI 575 (**A**) Growth rates of IPTG-

induced *lanM* expression in SLI 575 on 20 mM methanol with 2 uM or 50 nM LaCl$_3$ or La$_2$O$_3$

(**B**) Growth rates of IPTG-induced *lanM* expression in SLI 575 on 34 mM ethanol with 2 uM or

50 nM LaCl$_3$ or La$_2$O$_3$. Striped bars (right) indicate *lanM* expression induced with 1 mM IPTG.

N=2-4. Error bars represent standard deviation. Significant differences for each strain +/- 1 mM

IPTG determined by Student's paired t-test (*, $p<0.05$; **, $p<0.01$; ***, $p<0.001$)

Figure 5. Growth phenotypes of SLI test cohort and *Methylobacterium sp.* AMS5 on vanillic

acid in the presence and absence of La^{3+}. (**A**) Growth rates on 5 mM vanillic acid +/- 2 μM LaCl$_3$

(**B**) Final yields on 5 mM vanillic acid +/- 2 μM LaCl$_3$ (**C**) Growth rates on 12 mM vanillic acid

+/- 2 μM LaCl$_3$ (**D**) Final yields on 12 mM vanillic acid +/- 2 μM LaCl$_3$. n.g., no growth; m.g.,

minimal growth (<OD$_{600}$ 0.2). Black bars (bottom) represent cultures grown in the absence of

LaCl$_3$; hatched bars (top) represent cultures grown in the presence of LaCl$_3$. N=3. Error bars

represent standard deviation. Significant differences for each strain +/- LaCl$_3$ determined by

Student's paired t-test (*, p<0.05; **, p<0.01). Significant differences between all strains +/-

LaCl$_3$ determined using one-way ANOVA with post-hoc Tukey HSD (**Table S3**)

Figure 6. Growth phenotypes of SLI test cohort on methylamine in the presence and absence of

La^{3+}. (**A**) Growth rates on 15 mM methylamine +/- 2 μM $LaCl_3$ (**B**) Final yields on 15 mM

methylamine +/- 2 μM $LaCl_3$. Black bars (bottom) represent cultures grown in the absence of

$LaCl_3$; hatched bars (top) represent cultures grown in the presence of $LaCl_3$. N=3. Error bars

represent standard deviation. Significant differences for each strain +/- $LaCl_3$ determined by

Student's paired t-test. Significant differences between all strains +/- $LaCl_3$3 determined using

one-way ANOVA with post-hoc Tukey HSD (**Table S3**)

835 **Figure 7.** Growth curves of SLI 575 on 25 mM fructose +/- 2 μM LaCl₃. N=4. Error bars

836 represent standard deviation

Figure 8. Metabolic capabilities (growth on vanillic acid, growth on fructose, slow growth on

methylamine) and lanthanide-related gene products (MxaFI, XoxF, ExaF, lanmodulin) present in

SLI strains and model *extorquens* clade species referenced in this study (*M. extorquens* AM1, *M.*

extorquens PA1, *M. sp.* AMS5). Phylogenetic tree modified from **Figure 1A**

Table S1. Average nucleotide identity of all 11 SLI strains compared to evolutionarily-related

methylotroph and non-methylotrophic species, including all species represented in phylogenetic

tree from **Figure 1.** A, *Afipia felis* 76713; B, *Bradyrhizobium diazoefficiens* USDA 110; C,

Methylobacterium aquaticum DSM 16371; D, *Methylobacterium aquaticum* MA-22A;

E, *Methylobacterium brachiatum* 111MFTsu3.1M4; F, *Methylobacterium chloromethanicum*

CM4; G, *Methylobacterium extorquens* AM1; H, *Methylobacterium extorquens* DM4;

I, *Methylobacterium extorquens* PA1; J, *Methylobacterium indicum* NS230; K,

Methylobacterium indicum SE2,11; L, *Methylobacterium komagatae* DSM 19563;

M, *Methylobacterium nodulans* ORS 2060; N, *Methylobacterium organophilum* DSM 760;

O, *Methylobacterium oryzae* CBMB20; P, *Methylobacterium phyllosphaerae* CBMB27;

Q, *Methylobacterium platani* JCM 14648; R, *Methylobacterium populi* BJ001;

S, *Methylobacterium pseudosasicola* BL36; T, *Methylobacterium radiotolerans* JCM 2831;

U, *Methylobacterium* sp. 275MFSha3.1; V, *Methylobacterium* sp. 4-46; W, *Methylobacterium*

sp. AMS5; X, *Methylobacterium* sp. UNCCL125; Y, *Methylobacterium* sp. WSM2598; Z,

Methylobacterium tarhaniae DSM 25844; AA, *Methylobacterium* variabile DSM 16961; AB,

Methylobacterium zatmanii PSBB041; AC, *Pseudomonas putida* KT2440; AD,

Rhodopseudomonas palustris DSM 126; AE, SLI 223; AF, SLI 231; AG, SLI 233; AH, SLI 274;

AI, SLI 285; AJ, SLI 384; AK, SLI 499; AL, SLI 505; AM, SLI 516; AN, SLI 575; AO, SLI 576

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S	T	U	V	W	X	Y	Z	AA	AB	AC	AD	AE	AF	AG	AH	AI	AJ	AK	AL	AM	AN	AO
A	100	78	75	75	75	75	75	76	76	75	75	75	75	75	75	78	76	75	75	75	75	76	75	75	76	76	75	72	78	76	76	75	75	75	76	76	76	75	75	75	75
B	78	100	76	76	76	76	76	76	76	76	75	76	76	75	75	78	76	76	76	76	76	76	75	76	76	76	76	72	80	76	76	76	76	76	76	76	76	76	76	76	76
C	75	76	100	89	79	80	79	80	79	89	89	79	82	80	79	79	93	80	79	79	79	82	79	79	82	91	89	79	73	76	79	79	79	79	79	79	79	79	79	79	79
D	75	76	89	100	79	79	79	80	79	79	93	79	82	80	79	79	93	80	79	80	80	83	79	80	83	90	89	79	73	76	79	79	79	79	79	79	79	79	79	79	79
E	75	76	79	79	100	80	80	80	80	79	80	81	79	86	86	86	84	80	92	86	86	79	80	86	79	79	80	80	72	76	80	80	80	80	80	80	80	80	80	80	80
F	75	76	80	79	80	100	97	97	97	79	79	80	79	80	80	80	84	90	80	80	80	79	95	80	79	79	80	97	72	76	97	97	97	97	97	97	97	97	93	93	93
G	75	76	79	79	80	97	100	97	97	79	79	80	79	80	80	80	84	90	80	80	80	79	95	80	79	79	80	97	72	76	100	97	97	97	97	100	97	97	93	93	93
H	75	76	80	79	80	97	97	100	97	79	79	80	79	80	80	80	84	90	80	80	80	79	95	80	79	79	80	97	72	76	97	97	97	97	97	97	97	97	93	93	93
I	76	76	79	79	80	97	97	97	100	79	79	79	79	80	80	80	84	90	80	80	80	79	95	80	79	79	80	97	72	76	97	97	97	97	97	97	97	97	93	93	93
J	76	76	89	93	79	79	79	79	79	100	99	79	83	80	79	79	93	80	79	80	79	83	79	79	83	90	89	79	73	76	79	79	79	79	79	79	79	79	79	79	79
K	75	76	89	79	80	79	79	79	79	99	100	79	83	80	79	79	93	80	79	80	80	83	79	79	83	90	89	79	73	76	79	79	79	79	79	79	79	79	79	79	79

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
L	75	75	79	79	81	80	80	80	79	79	79	100	78	83	81	81	84	80	81	82	82	78	79	82	78	79	79	79	72	75	79	79	79	79	79	79	79	79	79	79
M	75	76	82	82	79	79	79	79	79	83	83	78	100	79	79	79	87	79	79	79	79	86	79	79	86	82	83	79	72	76	79	79	79	79	79	79	79	79	79	79
N	75	76	80	80	86	80	80	80	80	80	80	83	79	100	91	91	84	81	86	99	95	80	80	91	80	80	80	73	76	80	80	80	80	80	80	80	80	80	80	80
O	75	75	79	79	86	80	80	80	80	79	79	81	79	91	100	99	84	80	86	91	91	79	80	99	79	79	80	80	72	76	80	80	80	80	80	80	80	80	80	80
P	75	76	79	79	86	80	80	80	80	79	79	81	79	91	99	100	84	80	86	91	91	79	80	99	79	79	80	80	73	76	80	80	80	80	80	80	80	80	80	80
Q	78	78	93	93	84	84	84	84	84	93	93	83	87	84	84	84	100	84	84	85	84	87	84	84	87	93	92	84	75	78	84	84	84	84	84	84	84	84	84	84
R	76	76	80	80	80	90	90	90	90	80	80	80	79	81	80	80	84	100	80	81	80	79	90	80	79	80	80	90	72	76	90	90	90	90	90	90	90	90	90	90
S	75	76	79	79	92	80	80	80	80	79	79	81	79	86	86	86	84	80	100	86	86	79	80	86	79	79	79	80	73	76	80	80	80	80	80	80	80	80	80	80
T	75	76	79	80	86	80	80	80	80	80	80	82	79	99	91	91	85	81	86	100	95	80	80	91	80	80	80	80	73	76	80	80	80	80	80	80	80	80	80	80
U	75	76	79	80	86	80	80	80	80	80	80	82	79	95	91	91	84	80	86	95	100	80	80	91	80	80	80	80	73	76	80	80	80	80	80	80	80	80	80	80
V	75	76	82	83	79	79	79	79	79	83	83	78	86	80	79	79	87	79	79	80	80	100	79	79	79	83	83	79	72	76	79	79	79	79	79	79	79	79	79	79
W	75	76	79	79	80	95	95	95	95	79	79	79	79	80	80	80	84	90	80	80	80	79	100	80	79	79	79	95	72	76	95	95	95	95	95	95	95	93	93	93

X	75	76	79	80	86	80	80	80	80	79	79	82	79	91	99	99	84	80	86	91	91	79	80	100	79	79	80	80	73	76	80	80	80	80	80	80	80	80	80	80
Y	75	76	82	83	79	79	79	79	79	83	83	78	86	80	79	79	87	79	79	80	80	99	79	79	100	83	83	79	72	76	79	79	79	79	79	79	79	79	79	79
Z	76	76	91	90	79	79	79	79	79	90	90	79	82	80	79	79	93	80	79	80	80	83	79	79	83	100	90	79	73	76	79	79	79	79	79	79	79	79	79	79
AA	76	76	89	89	80	80	80	80	79	89	89	79	83	80	80	80	92	80	79	80	80	83	79	80	83	90	100	80	73	76	79	79	79	79	79	79	79	79	79	79
AB	75	76	79	79	80	97	97	97	97	79	79	79	79	80	80	80	84	80	80	80	79	95	80	79	79	79	80	100	72	76	97	97	97	97	97	97	97	93	93	93
AC	72	72	73	73	72	72	72	72	73	73	72	72	73	72	73	75	72	73	73	73	72	72	73	72	73	73	72	72	100	72	72	72	72	72	72	72	72	72	72	72
AD	78	80	76	76	76	76	76	76	76	76	76	75	76	76	76	76	78	76	76	76	76	76	76	76	76	76	76	72	76	100	76	76	76	76	76	76	76	76	76	76
AE	76	76	79	79	80	97	100	97	97	79	79	79	79	80	80	80	84	80	80	80	79	95	80	79	79	79	97	72	76	100	97	97	97	97	100	97	97	93	93	93
AF	76	76	79	79	80	97	97	97	97	79	79	79	79	80	80	80	84	80	80	80	79	95	80	79	79	79	97	72	76	97	100	100	100	100	97	100	100	94	94	94
AG	76	76	79	79	80	97	97	97	97	79	79	79	79	80	80	80	84	80	80	80	79	95	80	79	79	79	97	72	76	97	100	100	100	100	97	100	100	94	94	94
AH	76	76	79	79	80	97	97	97	97	79	79	79	79	80	80	80	84	80	80	80	79	95	80	79	79	79	97	72	76	97	100	100	100	100	97	100	100	94	94	94
AI	76	76	79	79	80	97	97	97	97	79	79	79	79	80	80	80	84	80	80	80	79	95	80	79	79	79	97	72	76	97	100	100	100	100	97	100	100	94	94	94

AJ	76	76	79	79	80	97	100	97	97	79	79	79	79	80	80	80	84	90	80	80	80	79	95	80	79	79	79	97	72	76	100	97	97	97	97	100	97	97	93	93	93
AK	76	76	79	79	80	97	97	97	97	79	79	79	79	80	80	80	84	90	80	80	80	79	95	80	79	79	79	97	72	76	97	100	100	100	100	97	100	100	94	94	94
AL	76	76	79	79	80	97	97	97	97	79	79	79	79	80	80	80	84	90	80	80	80	79	95	80	79	79	79	97	72	76	97	100	100	100	100	97	100	100	94	94	94
AM	75	76	79	79	80	93	93	93	93	79	79	79	79	80	80	80	84	90	80	80	80	79	93	80	79	79	79	93	72	76	93	94	94	94	94	93	94	94	100	100	100
AN	75	76	79	79	80	93	93	93	93	79	79	79	79	80	80	80	84	90	80	80	80	79	93	80	79	79	79	93	72	76	93	94	94	94	94	93	94	94	100	100	100
AO	75	76	79	79	80	93	93	93	93	79	79	79	79	80	80	80	84	90	80	80	80	79	93	80	79	79	79	93	72	76	93	94	94	94	94	93	94	94	100	100	100

878 **Table S2**. Summary of metabolic and lanthanide-related genes present in SLI strains, using *M. extorquens* AM1 and PA1 as reference

879 strains

		Strain												
		M. extorquens AM1	*M. extorquens* PA1	SLI 223	SLI 231	SLI 233	SLI 274	SLI 285	SLI 384	SLI 499	SLI 505	SLI 516	SLI 575	SLI 576
PQQ-dep. alcohol dehydrogenases	*mxaF*	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mxaI	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mxaG	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	xoxF1	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	xoxG	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	xoxF2	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	exaF	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	-	-	-
	exaG	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	-	-	-
Formaldehyde oxidation pathway	*fae1*	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	fae2	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	fae3	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mtdA	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mtdB	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mch	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	fhcA	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	fhcB	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	fhcC	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	fhcD	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Formate dehydrogenases	*fdh1A*	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	-	-	-
	fdh1B	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	-	-	-
	fdh2A	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	fdh2B	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	fdh2C	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	fdh2D	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	fdh3A	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	fdh3B	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	fdh4A	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	fdh4B	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Formate assimilation & serine cycle	*mtdA*	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	fch	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	ftfL	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	sga	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	glyA	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	hpr	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
RuMP cycle	*hps*	-	-	-	-	-	-	-	-	-	-	-	-	-
	phi	-	-	-	-	-	-	-	-	-	-	-	-	-
	pfk	-	-	-	-	-	-	-	-	-	-	✓	✓	✓
	fba	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

		Strain												
		M. extorquens AM1	*M. extorquens* PA1	SLI 223	SLI 231	SLI 233	SLI 274	SLI 285	SLI 384	SLI 499	SLI 505	SLI 516	SLI 575	SLI 576
Methylamine dehydrogenase	*mauF*	✓	-	-	-	-	-	-	-	-	-	-	-	-
	mauB	✓	-	-	-	-	-	-	-	-	-	-	-	-
	mauE	✓	-	-	-	-	-	-	-	-	-	-	-	-
	mauD	✓	-	-	-	-	-	-	-	-	-	-	-	-
	mauA	✓	-	-	-	-	-	-	-	-	-	-	-	-
	mauC	✓	-	-	-	-	-	-	-	-	-	-	-	-
	mauJ	✓	-	-	-	-	-	-	-	-	-	-	-	-
	mauG	✓	-	-	-	-	-	-	-	-	-	-	-	-
	mauL	✓	-	-	-	-	-	-	-	-	-	-	-	-
	mauM	✓	-	-	-	-	-	-	-	-	-	-	-	-
	mauN	✓	-	-	-	-	-	-	-	-	-	-	-	-
N-methylglutamate pathway	*mgdD*	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mgdC	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mgdB	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mgdA	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mgsA	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mgsB	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mgsC	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	gmaS	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Aromatic acid catabolism gene island	*ech*	-	-	-	✓	✓	✓	✓	-	✓	✓	✓	✓	✓
	vdh	-	-	-	✓	✓	✓	✓	-	✓	✓	✓	✓	✓
	vanA	-	-	-	✓	✓	✓	✓	-	✓	✓	✓	✓	✓
	vanB	-	-	-	✓	✓	✓	✓	-	✓	✓	✓	✓	✓
	pobA	-	-	-	✓	✓	✓	✓	-	✓	✓	✓	✓	✓
	pcaG	-	-	-	✓	✓	✓	✓	-	✓	✓	✓	✓	✓
	pcaH	-	-	-	✓	✓	✓	✓	-	✓	✓	✓	✓	✓
	vanR	-	-	-	✓	✓	✓	✓	-	✓	✓	✓	✓	✓
	pcaK	-	-	-	✓	✓	✓	✓	-	✓	✓	✓	✓	✓
	pcaB	-	-	-	✓	✓	✓	✓	-	✓	✓	✓	✓	✓
	pcaC	-	-	-	✓	✓	✓	✓	-	✓	✓	✓	✓	✓
	pcaD	-	-	-	✓	✓	✓	✓	-	✓	✓	✓	✓	✓
	pcaIJ	-	-	-	✓	✓	✓	✓	-	✓	✓	✓	✓	✓
	pcaF	-	-	-	✓	✓	✓	✓	-	✓	✓	✓	✓	✓

		M. extorquens AM1	M. extorquens PA1	SLI 223	SLI 231	SLI 233	SLI 274	SLI 285	SLI 384	SLI 499	SLI 505	SLI 516	SLI 575	SLI 576
							Strain							
Methylolanthanin Cluster	*mluA*	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mluR	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mluI	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mllA	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mllBC	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mllDE	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mllF	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mllG	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mllH	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mllJ	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Lanthanide Utilization and Transport	*lutA*	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	lutB	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	lutC	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	lutD	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	lutE	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	lutF	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	lutG	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	lutH	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	lanM	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	-	-	-
	lutI	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Ln Switch	*mxcQ*	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mxcE	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mxbD	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
	mxbM	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

886 **Table S3.** Significant differences in growth rates and final yields between all strains on 20 mM

887 methanol (**Figure 3**) **,** 34 mM ethanol (**Figure 3**)**,** 5 and 12 mM vanillic acid (**Figure 5**), 15 mM

888 methylamine **(Figure 6**) +/- LaCl$_3$ determined using one-way ANOVA with post-hoc Tukey

889 HSD. Significant differences for each strain +/- LaCl$_3$ or +/- IPTG for lanmodulin expression

890 phenotypes (**Figure 4**) determined by Student's paired t-test

20 mM Methanol ± LaCl$_3$			
Growth Rate		**Final Yield**	
Comparison	**p-value**	**Comparison**	**p-value**
SLI 231 ± LaCl$_3$	0.005752	SLI 231 ± LaCl$_3$	0.1264
SLI 233 ± LaCl$_3$	0.002731	SLI 233 ± LaCl$_3$	0.4189
SLI 384 ± LaCl$_3$	0.6598	SLI 384 ± LaCl$_3$	0.6513
SLI 499 ± LaCl$_3$	0.00311	SLI 499 ± LaCl$_3$	0.2585
SLI 505 ± LaCl$_3$	0.004484	SLI 505 ± LaCl$_3$	0.2158
SLI 575 ± LaCl$_3$	0.00002213	SLI 575 ± LaCl$_3$	0.2463
SLI 231 x SLI 233 - LaCl$_3$	1	SLI 231 x SLI 233 - LaCl$_3$	0.9232
SLI 231 x SLI 384 - LaCl$_3$	0.133	SLI 231 x SLI 384 - LaCl$_3$	0.9546
SLI 231 x SLI 499 - LaCl$_3$	1	SLI 231 x SLI 499 - LaCl$_3$	0.9966
SLI 231 x SLI 505 - LaCl$_3$	0.9998	SLI 231 x SLI 505 - LaCl$_3$	0.6545
SLI 231 x SLI 575 - LaCl$_3$	0.7947	SLI 231 x SLI 575 - LaCl$_3$	0.05092
SLI 233 x SLI 384 - LaCl$_3$	0.1335	SLI 233 x SLI 384 - LaCl$_3$	1
SLI 233 x SLI 499 - LaCl$_3$	1	SLI 233 x SLI 499 - LaCl$_3$	0.9949
SLI 233 x SLI 505 - LaCl$_3$	0.9998	SLI 233 x SLI 505 - LaCl$_3$	0.9861
SLI 233 x SLI 575 - LaCl$_3$	0.7929	SLI 233 x SLI 575 - LaCl$_3$	0.01978
SLI 384 x SLI 499 - LaCl$_3$	0.1097	SLI 384 x SLI 499 - LaCl$_3$	0.9987
SLI 384 x SLI 505 - LaCl$_3$	0.1007	SLI 384 x SLI 505 - LaCl$_3$	0.9702
SLI 384 x SLI 575 - LaCl$_3$	0.03403	SLI 384 x SLI 575 - LaCl$_3$	0.02226
SLI 499 x SLI 505 - LaCl$_3$	1	SLI 499 x SLI 505 - LaCl$_3$	0.8666
SLI 499 x SLI 575 - LaCl$_3$	0.8691	SLI 499 x SLI 575 - LaCl$_3$	0.03202
SLI 505 x SLI 575 - LaCl$_3$	0.8975	SLI 505 x SLI 575 - LaCl$_3$	0.01123
SLI 231 x SLI 233 + LaCl$_3$	0.5873	SLI 231 x SLI 233 + LaCl$_3$	0.9674
SLI 231 x SLI 384 + LaCl$_3$	0.8102	SLI 231 x SLI 384 + LaCl$_3$	0.9996
SLI 231 x SLI 499 + LaCl$_3$	0.9683	SLI 231 x SLI 499 + LaCl$_3$	0.9961
SLI 231 x SLI 505 + LaCl$_3$	0.6435	SLI 231 x SLI 505 + LaCl$_3$	0.07812
SLI 231 x SLI 575 + LaCl$_3$	0.9997	SLI 231 x SLI 575 + LaCl$_3$	0.00001957
SLI 233 x SLI 384 + LaCl$_3$	0.9966	SLI 233 x SLI 384 + LaCl$_3$	0.8901
SLI 233 x SLI 499 + LaCl$_3$	0.28	SLI 233 x SLI 499 + LaCl$_3$	0.82
SLI 233 x SLI 505 + LaCl$_3$	1	SLI 233 x SLI 505 + LaCl$_3$	0.1797
SLI 233 x SLI 575 + LaCl$_3$	0.7234	SLI 233 x SLI 575 + LaCl$_3$	0.00001466
SLI 384 x SLI 499 + LaCl$_3$	0.4465	SLI 384 x SLI 499 + LaCl$_3$	1

Comparison	p-value	Comparison	p-value
SLI 384 x SLI 505 + LaCl$_3$	0.9991	SLI 384 x SLI 505 + LaCl$_3$	0.05721
SLI 384 x SLI 575 + LaCl$_3$	0.9146	SLI 384 x SLI 575 + LaCl$_3$	0.00002191
SLI 499 x SLI 505 + LaCl$_3$	0.3154	SLI 499 x SLI 505 + LaCl$_3$	0.04763
SLI 499 x SLI 575 + LaCl$_3$	0.8993	SLI 499 x SLI 575 + LaCl$_3$	0.00002345
SLI 505 x SLI 575 + LaCl$_3$	0.7776	SLI 505 x SLI 575 + LaCl$_3$	0.00000566

34 mM Ethanol ± LaCl$_3$			
Growth Rate		**Final Yield**	
Comparison	**p-value**	**Comparison**	**p-value**
SLI 231 ± LaCl$_3$	0.09398	SLI 231 ± LaCl$_3$	0.9407
SLI 233 ± LaCl$_3$	0.07046	SLI 233 ± LaCl$_3$	0.6408
SLI 384 ± LaCl$_3$	0.08337	SLI 384 ± LaCl$_3$	0.8935
SLI 499 ± LaCl$_3$	0.3597	SLI 499 ± LaCl$_3$	0.2088
SLI 505 ± LaCl$_3$	0.3525	SLI 505 ± LaCl$_3$	0.008165
SLI 575 ± LaCl$_3$	0.5879	SLI 575 ± LaCl$_3$	0.694
SLI 231 x SLI 233 - LaCl$_3$	0.8923	SLI 231 x SLI 233 - LaCl$_3$	1
SLI 231 x SLI 384 - LaCl$_3$	0.1229	SLI 231 x SLI 384 - LaCl$_3$	1
SLI 231 x SLI 499 - LaCl$_3$	0.9989	SLI 231 x SLI 499 - LaCl$_3$	0.9171
SLI 231 x SLI 505 - LaCl$_3$	0.9319	SLI 231 x SLI 505 - LaCl$_3$	1
SLI 231 x SLI 575 - LaCl$_3$	0.2853	SLI 231 x SLI 575 - LaCl$_3$	0.06206
SLI 233 x SLI 384 - LaCl$_3$	0.5211	SLI 233 x SLI 384 - LaCl$_3$	0.9999
SLI 233 x SLI 499 - LaCl$_3$	0.9805	SLI 233 x SLI 499 - LaCl$_3$	0.9522
SLI 233 x SLI 505 - LaCl$_3$	0.4079	SLI 233 x SLI 505 - LaCl$_3$	0.9998
SLI 233 x SLI 575 - LaCl$_3$	0.8289	SLI 233 x SLI 575 - LaCl$_3$	0.05244
SLI 384 x SLI 499 - LaCl$_3$	0.2151	SLI 384 x SLI 499 - LaCl$_3$	0.9858
SLI 384 x SLI 505 - LaCl$_3$	0.02644	SLI 384 x SLI 505 - LaCl$_3$	1
SLI 384 x SLI 575 - LaCl$_3$	0.9918	SLI 384 x SLI 575 - LaCl$_3$	0.07454
SLI 499 x SLI 505 - LaCl$_3$	0.7844	SLI 499 x SLI 505 - LaCl$_3$	0.9889
SLI 499 x SLI 575 - LaCl$_3$	0.4549	SLI 499 x SLI 575 - LaCl$_3$	0.2034
SLI 505 x SLI 575 - LaCl$_3$	0.06793	SLI 505 x SLI 575 - LaCl$_3$	0.07879
SLI 231 x SLI 233 + LaCl$_3$	0.9987	SLI 231 x SLI 233 + LaCl$_3$	0.9963
SLI 231 x SLI 384 + LaCl$_3$	0.004637	SLI 231 x SLI 384 + LaCl$_3$	0.9995
SLI 231 x SLI 499 + LaCl$_3$	0.93	SLI 231 x SLI 499 + LaCl$_3$	0.2218
SLI 231 x SLI 505 + LaCl$_3$	0.2374	SLI 231 x SLI 505 + LaCl$_3$	0.1215
SLI 231 x SLI 575 + LaCl$_3$	0.6082	SLI 231 x SLI 575 + LaCl$_3$	0.111
SLI 233 x SLI 384 + LaCl$_3$	0.008673	SLI 233 x SLI 384 + LaCl$_3$	0.9655
SLI 233 x SLI 499 + LaCl$_3$	0.7734	SLI 233 x SLI 499 + LaCl$_3$	0.1068
SLI 233 x SLI 505 + LaCl$_3$	0.1337	SLI 233 x SLI 505 + LaCl$_3$	0.05601
SLI 233 x SLI 575 + LaCl$_3$	0.4013	SLI 233 x SLI 575 + LaCl$_3$	0.05096
SLI 384 x SLI 499 + LaCl$_3$	0.001055	SLI 384 x SLI 499 + LaCl$_3$	0.3409
SLI 384 x SLI 505 + LaCl$_3$	0.000131	SLI 384 x SLI 505 + LaCl$_3$	0.1958
SLI 384 x SLI 575 + LaCl$_3$	0.000385	SLI 384 x SLI 575 + LaCl$_3$	0.1798
SLI 499 x SLI 505 + LaCl$_3$	0.7021	SLI 499 x SLI 505 + LaCl$_3$	0.9985
SLI 499 x SLI 575 + LaCl$_3$	0.9815	SLI 499 x SLI 575 + LaCl$_3$	0.9971
SLI 505 x SLI 575 + LaCl$_3$	0.9678	SLI 505 x SLI 575 + LaCl$_3$	1

15 mM Methylamine ± LaCl$_3$			
Growth Rate		**Final Yield**	
Comparison	**p-value**	**Comparison**	**p-value**
SLI 231 ± LaCl$_3$	0.2622	SLI 231 ± LaCl$_3$	0.8076
SLI 233 ± LaCl$_3$	0.3036	SLI 233 ± LaCl$_3$	0.3688
SLI 384 ± LaCl$_3$	0.1904	SLI 384 ± LaCl$_3$	0.2502
SLI 499 ± LaCl$_3$	0.5452	SLI 499 ± LaCl$_3$	0.6538
SLI 505 ± LaCl$_3$	0.8891	SLI 505 ± LaCl$_3$	0.5779
SLI 575 ± LaCl$_3$	0.6923	SLI 575 ± LaCl$_3$	0.687
SLI 231 x SLI 233 - LaCl$_3$	0.9982	SLI 231 x SLI 233 - LaCl$_3$	1
SLI 231 x SLI 384 - LaCl$_3$	0.000869	SLI 231 x SLI 384 - LaCl$_3$	0.00843
SLI 231 x SLI 499 - LaCl$_3$	0.8468	SLI 231 x SLI 499 - LaCl$_3$	0.9972
SLI 231 x SLI 505 - LaCl$_3$	0.7692	SLI 231 x SLI 505 - LaCl$_3$	0.9984
SLI 231 x SLI 575 - LaCl$_3$	0.9989	SLI 231 x SLI 575 - LaCl$_3$	0.02364
SLI 233 x SLI 384 - LaCl$_3$	0.001113	SLI 233 x SLI 384 - LaCl$_3$	0.008092
SLI 233 x SLI 499 - LaCl$_3$	0.9659	SLI 233 x SLI 499 - LaCl$_3$	0.9984
SLI 233 x SLI 505 - LaCl$_3$	0.9256	SLI 233 x SLI 505 - LaCl$_3$	0.9991
SLI 233 x SLI 575 - LaCl$_3$	0.9681	SLI 233 x SLI 575 - LaCl$_3$	0.02556
SLI 384 x SLI 499 - LaCl$_3$	0.001847	SLI 384 x SLI 499 - LaCl$_3$	0.005875
SLI 384 x SLI 505 - LaCl$_3$	0.002082	SLI 384 x SLI 505 - LaCl$_3$	0.00611
SLI 384 x SLI 575 - LaCl$_3$	0.0007	SLI 384 x SLI 575 - LaCl$_3$	0.858
SLI 499 x SLI 505 - LaCl$_3$	1	SLI 499 x SLI 505 - LaCl$_3$	1
SLI 499 x SLI 575 - LaCl$_3$	0.6775	SLI 499 x SLI 575 - LaCl$_3$	0.01568
SLI 505 x SLI 575 - LaCl$_3$	0.5903	SLI 505 x SLI 575 - LaCl$_3$	0.0164
SLI 231 x SLI 233 + LaCl$_3$	1	SLI 231 x SLI 233 + LaCl$_3$	0.9982
SLI 231 x SLI 384 + LaCl$_3$	0.00001482	SLI 231 x SLI 384 + LaCl$_3$	0.0002457
SLI 231 x SLI 499 + LaCl$_3$	0.5225	SLI 231 x SLI 499 + LaCl$_3$	1
SLI 231 x SLI 505 + LaCl$_3$	0.845	SLI 231 x SLI 505 + LaCl$_3$	0.869
SLI 231 x SLI 575 + LaCl$_3$	0.9733	SLI 231 x SLI 575 + LaCl$_3$	0.002645
SLI 233 x SLI 384 + LaCl$_3$	0.00001392	SLI 233 x SLI 384 + LaCl$_3$	0.0003005
SLI 233 x SLI 499 + LaCl$_3$	0.4466	SLI 233 x SLI 499 + LaCl$_3$	0.9939
SLI 233 x SLI 505 + LaCl$_3$	0.7702	SLI 233 x SLI 505 + LaCl$_3$	0.9749
SLI 233 x SLI 575 + LaCl$_3$	0.9913	SLI 233 x SLI 575 + LaCl$_3$	0.003543
SLI 384 x SLI 499 + LaCl$_3$	0.0000301	SLI 384 x SLI 499 + LaCl$_3$	0.0002317
SLI 384 x SLI 505 + LaCl$_3$	0.00002318	SLI 384 x SLI 505 + LaCl$_3$	0.0004394
SLI 384 x SLI 575 + LaCl$_3$	0.00001136	SLI 384 x SLI 575 + LaCl$_3$	0.0548
SLI 499 x SLI 505 + LaCl$_3$	0.9809	SLI 499 x SLI 505 + LaCl$_3$	0.8198
SLI 499 x SLI 575 + LaCl$_3$	0.2506	SLI 499 x SLI 575 + LaCl$_3$	0.002429
SLI 505 x SLI 575 + LaCl$_3$	0.4965	SLI 505 x SLI 575 + LaCl$_3$	0.006196
12 mM Vanillic Acid ± LaCl$_3$			
Growth Rate		**Final Yield**	
Comparison	**p-value**	**Comparison**	**p-value**
SLI 231 ± LaCl$_3$	0.02098	SLI 231 ± LaCl$_3$	0.1654

Comparison	p-value	Comparison	p-value
SLI 233 ± LaCl$_3$	0.06785	SLI 233 ± LaCl$_3$	0.35
SLI 499 ± LaCl$_3$	0.1018	SLI 499 ± LaCl$_3$	0.2796
SLI 505 ± LaCl$_3$	0.01559	SLI 505 ± LaCl$_3$	0.01442
SLI 575 ± LaCl$_3$	0.8571	SLI 575 ± LaCl$_3$	0.6117
SLI 231 x SLI 233 - LaCl$_3$	0.05422	SLI 231 x SLI 233 - LaCl$_3$	0.9758
SLI 231 x SLI 499 - LaCl$_3$	0.000385	SLI 231 x SLI 499 - LaCl$_3$	1
SLI 231 x SLI 505 - LaCl$_3$	0.1099	SLI 231 x SLI 505 - LaCl$_3$	1
SLI 231 x SLI 575 - LaCl$_3$	0.002493	SLI 231 x SLI 575 - LaCl$_3$	0.03123
SLI 233 x SLI 499 - LaCl$_3$	0.004892	SLI 233 x SLI 499 - LaCl$_3$	0.9737
SLI 233 x SLI 505 - LaCl$_3$	0.8575	SLI 233 x SLI 505 - LaCl$_3$	0.9763
SLI 233 x SLI 575 - LaCl$_3$	0.0807	SLI 233 x SLI 575 - LaCl$_3$	0.01756
SLI 499 x SLI 505 - LaCl$_3$	0.001474	SLI 499 x SLI 505 - LaCl$_3$	1
SLI 499 x SLI 575 - LaCl$_3$	0.1419	SLI 499 x SLI 575 - LaCl$_3$	0.03167
SLI 505 x SLI 575 - LaCl$_3$	0.02013	SLI 505 x SLI 575 - LaCl$_3$	0.01953
SLI 231 x SLI 233 + LaCl$_3$	0.003649	SLI 231 x SLI 233 + LaCl$_3$	0.5874
SLI 231 x SLI 499 + LaCl$_3$	0.05221	SLI 231 x SLI 499 + LaCl$_3$	0.9987
SLI 231 x SLI 505 + LaCl$_3$	1	SLI 231 x SLI 505 + LaCl$_3$	0.7883
SLI 231 x SLI 575 + LaCl$_3$	0.006383	0.SLI 231 x SLI 575 + LaCl$_3$	0.0005726
SLI 233 x SLI 499 + LaCl$_3$	0.1471	SLI 233 x SLI 499 + LaCl$_3$	0.4589
SLI 233 x SLI 505 + LaCl$_3$	0.002128	SLI 233 x SLI 505 + LaCl$_3$	0.98
SLI 233 x SLI 575 + LaCl$_3$	0.9563	SLI 233 x SLI 575 + LaCl$_3$	0.0002413
SLI 499 x SLI 505 + LaCl$_3$	0.03278	SLI 499 x SLI 505 + LaCl$_3$	0.6414
SLI 499 x SLI 575 + LaCl$_3$	0.3161	SLI 499 x SLI 575 + LaCl$_3$	0.0006731
SLI 505 x SLI 575 + LaCl$_3$	0.003754	SLI 505 x SLI 575 + LaCl$_3$	0.0001875

5 mM Vanillic Acid ± LaCl$_3$

Growth Rate		Final Yield	
Comparison	**p-value**	**Comparison**	**p-value**
SLI 231 ± LaCl$_3$	0.1459	SLI 231 ± LaCl$_3$	0.0005
SLI 233 ± LaCl$_3$	0.7953	SLI 233 ± LaCl$_3$	0.002389
SLI 499 ± LaCl$_3$	0.1194	SLI 499 ± LaCl$_3$	0.04408
SLI 505 ± LaCl$_3$	0.01286	SLI 505 ± LaCl$_3$	0.02268
SLI 575 ± LaCl$_3$	0.6059	SLI 575 ± LaCl$_3$	0.7737
sp. AMS5 ± LaCl$_3$	0.08744	sp. AMS5 ± LaCl$_3$	0.4056
SLI 231 x SLI 233 - LaCl$_3$	0.788	SLI 231 x SLI 233 - LaCl$_3$	1
SLI 231 x SLI 499 - LaCl$_3$	0.9988	SLI 231 x SLI 499 - LaCl$_3$	0.9978
SLI 231 x SLI 505 - LaCl$_3$	0.6593	SLI 231 x SLI 505 - LaCl$_3$	0.9547
SLI 231 x SLI 575 - LaCl$_3$	1	SLI 231 x SLI 575 - LaCl$_3$	0.000497
SLI 231 x sp. AMS5 - LaCl$_3$	0.001096	SLI 231 x sp. AMS5 - LaCl$_3$	0.02093
SLI 233 x SLI 499- LaCl$_3$	0.5811	SLI 233 x SLI 499- LaCl$_3$	0.9905
SLI 233 x SLI 505- LaCl$_3$	0.1241	SLI 233 x SLI 505- LaCl$_3$	0.9811
SLI 233 x SLI 575- LaCl$_3$	0.727	SLI 233 x SLI 575- LaCl$_3$	0.000395
SLI 233 x sp. AMS5 - LaCl$_3$	0.000169	SLI 233 x sp. AMS5 - LaCl$_3$	0.01604
SLI 499 x SLI 505- LaCl$_3$	0.8523	SLI 499 x SLI 505- LaCl$_3$	0.8003

Comparison	p-value	Comparison	p-value
SLI 499 x SLI 575 - LaCl$_3$	0.9998	SLI 499 x SLI 575 - LaCl$_3$	0.000938
SLI 499 x sp. AMS5- LaCl$_3$	0.001957	SLI 499 x sp. AMS5- LaCl$_3$	0.0428
SLI 505 x SLI 575 - LaCl$_3$	0.7242	SLI 505 x SLI 575 - LaCl$_3$	0.000151
SLI 505 x sp. AMS5 - LaCl$_3$	0.0128	SLI 505 x sp. AMS5 - LaCl$_3$	0.005178
SLI 575 x sp. AMS5- LaCl$_3$	0.001309	SLI 575 x sp. AMS5- LaCl$_3$	0.2517
SLI 231 x SLI 233 + LaCl$_3$	0.8618	SLI 231 x SLI 233 + LaCl$_3$	
SLI 231 x SLI 499 + LaCl$_3$	0.9976	SLI 231 x SLI 499 + LaCl$_3$	0.9983
SLI 231 x SLI 505 + LaCl$_3$	0.04592	SLI 231 x SLI 505 + LaCl$_3$	0.8716
SLI 231 x SLI 575 + LaCl$_3$	0.9691	SLI 231 x SLI 575 + LaCl$_3$	0.0002235
SLI 231 x sp. AMS5 + LaCl$_3$	3.07E-08	SLI 231 x sp. AMS5 + LaCl$_3$	4.54E-07
SLI 233 x SLI 499 + LaCl$_3$	0.6388	SLI 233 x SLI 499 + LaCl$_3$	0.000003121
SLI 233 x SLI 505 + LaCl$_3$	0.2714	SLI 233 x SLI 505 + LaCl$_3$	0.6685
SLI 233 x SLI 575+ LaCl$_3$	0.9989	SLI 233 x SLI 575+ LaCl$_3$	1.28E-04
SLI 233 x sp. AMS5 + LaCl$_3$	1.42E-08	SLI 233 x sp. AMS5 + LaCl$_3$	3.16E-07
SLI 499 x SLI 505 + LaCl$_3$	0.02222	SLI 499 x SLI 505 + LaCl$_3$	2.06E-06
SLI 499 x SLI 575 + LaCl$_3$	0.8327	SLI 499 x SLI 575 + LaCl$_3$	0.001131
SLI 499 x sp. AMS5 + LaCl$_3$	4.21E-08	SLI 499 x sp. AMS5 + LaCl$_3$	0.000001286
SLI 505 x SLI 575 + LaCl$_3$	0.1584	SLI 505 x SLI 575 + LaCl$_3$	0.00001043
SLI 505 x sp. AMS5 + LaCl$_3$	3.60E-09	SLI 505 x sp. AMS5 + LaCl$_3$	0.001483
SLI 575 x sp. AMS5 + LaCl$_3$	1.81E-08	SLI 575 x sp. AMS5 + LaCl$_3$	0.03846

SLI 575/*lanM* ± IPTG

Growth Rate in 20 mM MeOH		Growth Rate in 34 mM EtOH	
Comparison	p-value	Comparison	p-value
2 µM LaCl$_3$ ± IPTG	0.9924	2 µM LaCl$_3$ ± IPTG	0.2604
50 nM LaCl$_3$ ± IPTG	0.02077	50 nM LaCl$_3$ ± IPTG	0.00736
1 µM La$_2$O$_3$ ± IPTG	0.00772	1 µM La$_2$O$_3$ ± IPTG	0.000704
25 nM La$_2$O$_3$ ± IPTG	0.01731	25 nM La$_2$O$_3$ ± IPTG	0.09511