Transferrin Receptor is Required for CD4+ T Cell Iron-Dependent Response During Acute *Toxoplasma gondii* Infection[1]

Stephen L. Denton*, Tathagato Roy*, Martin Akandawen*, Hunter K. Keplinger*, Sai K. Ng*, and Jason P. Gigley*†

*Department of Molecular Biology, University of Wyoming, Laramie, WY, U.S.A.
†Corresponding Author: 307-766-6151, jgigley@uwyo.edu

Abstract

Elemental iron is an essential nutrient involved in many biological processes including infection and immunity. How iron impacts *Toxoplasma gondii* (*T. gondii*) *in vivo* and development of immunity during infection is unclear. We found that although iron is required for parasite proliferation *in vitro,* paradoxically, iron restriction *in vivo* increased parasite burdens during acute and persistent infection stages and decreased survival of mice. Iron restriction lowered IL-12 and IFNγ in spleen and serum, but ratios of myeloid cells and the number and function of Natural Killer cells were unchanged. Iron restriction significantly impaired the development of CD4+ and CD8+ T cell responses to *T. gondii* during replicating type II and attenuated vaccine strain *cps1-1* infection. Low iron conditions reduced the percent and absolute numbers of antigen experienced CD11a+CD49d+, functional IFNγ+, and CD62L-KLRG1+ effector T cells. Iron restriction also decreased vaccine efficacy of *cps1-1* strain against secondary lethal challenge. Antigen experienced CD4+ and CD8+ T cells both upregulated their iron transporter Transferrin receptor 1 (CD71) during infection regardless of iron restriction. Mice whose CD4+ T cells were deficient in CD71 had reduced CD4+ T cell antigen experience and polyfunctionality, yet CD8+ T cell responses remained intact and their long term survival

[1] Funding Sources: NIH 2P20GM103432, R21AI1592, and R21AI1612.

was not affected compared to wild type litter mate controls. This study highlights that iron

acquisition by T cells is required for activation and vaccine induced long-term protection

against *T. gondii*. Understanding how iron affects multiple immune compartments will be

essential to define iron regulation of immunity to *T. gondii*.

List of Abbreviations

Transferrin Receptor 1, CD71; Deferiprone, DFP; Toxoplasma Lysate Antigen, TLA;

(Days, Weeks) Post Infection, (D,W)PI; T Cell Receptor, TCR; Mean Fluorescent

Intensity, MFI; Radical Oxygen Species, ROS; Conditional Knockout of CD71 in CD4+ T

Cells; CD4cCD71KO

Introduction

Toxoplasma gondii (T. gondii) is an obligate intracellular protozoan parasite infecting about 30% of people world-wide which persists for life and poses a significant health risk (1-3). Understanding the factors that contribute to pathogenesis and development of immunity to this pathogen are critical to develop better vaccines and therapies to treat this infection. T. gondii has evolved to rely on scavenging nutrients from the host (4-6). Many of these nutrients are also required for proper development of immune responses to control this infection (7-9). The interplay between parasite nutritional scavenging and immune response development dependent on the same nutrients is poorly understood.

Essential metal iron is a nutrient that is utilized both by T. gondii and the development of immunity to the parasite. T. gondii is highly sensitive to exogenous control of iron levels (10, 11). Treatment of infected cells with IFNγ impairing parasite replication is rescued by iron salt and iron-bearing transferrin (12). A natural function of IFNγ treatment of cells is downregulation of iron uptake receptor Transferrin Receptor (CD71) (13); CD71 in turn is upregulated by T. gondii secretions (14). Modulating iron levels in mice infections has been shown to affect intestinal inflammatory damage as well as parasite load (15). However, the long term effects of iron modulation on development of protective immunity to T. gondii is not known.

Immunity to T. gondii is triggered by recognition of the parasite by TLR4 and 11 and the production of the proinflammatory cytokine IL-12 (16-20). IL-12 activates group 1 Innate Lymphoid Cells including Natural Killer (NK) Cells and ILC1 for the initial production of IFNγ required for early control of T. gondii (21-23). After presentation of T.

gondii antigens to T cells by CD8a+ Dendritic cells, both CD4+ and CD8+ T cells mount

a robust response producing high levels of IFNγ (24). CD8+ T cells, *via* their

IFNγ production, emerge to become the dominant cell required for long-term protection

against the infection. CD4+ T cells are required continuously to help maintain CD8+ T

cell dependent protection (25, 26). Despite induction of a robust CD8+ T cell response,

immunity to *T. gondii* is not sterilizing and the parasite escapes to develop a persistent

infection in skeletal muscle and the central nervous system (27). Part of the inability of

CD8+ T cells to clear the persistent infection is the development of immune exhaustion

(27-29). The immune exhaustion of CD8+ T cells maybe partially dependent upon CD4+

T cells responding to the infection, highlighting that CD4+ T cells can be essential but

also detrimental to long term CD8+ T cell dependent protection (26, 30). The

immunological reasons why the parasite is not completely cleared is still not well defined.

Available iron could be an important factor required in this process.

Iron-dependent roles in immunity include neutrophil extracellular trap formation

(31), macrophage polarization (32, 33), dendritic cell maturation (34), and NK cell target

recognition (35, 36). T Cell proliferation and IL-2R upregulation during activation is also

dependent on intracellular iron levels (37) and CD71-dependent iron uptake (38). T-helper

type 1 (Th1) but not Th2 cytokine production is also dependent on iron status (39, 40),

indicating that relative iron levels could be a sensing mechanism for T-helper type

differentiation. Much of what is known about iron and immunity is in the absence of

infection. How iron regulates infection pathogenesis and development of immunity during

infection including *T. gondii* infection *in vivo* is unclear.

We investigated how limiting iron during infection in a murine model impacted

infection pathogenesis and immunity to *T. gondii*. Similar to previously published studies,

we observed that limiting iron *in vitro* slowed the proliferation of the parasite. However,

unlike other studies we observed that there was increased morbidity and mortality of the

animals in iron limiting conditions, and the parasite disseminated better to the CNS to

develop a much higher persistent cyst burden than normal iron conditions. To examine

the role of iron in *T. gondii* resistance, we observed that splenic and serum production of

IL-12p40 and IFNγ by 5 days after infection were inhibited by iron restriction, but the

proportions of spleen leukocytes and NK production of IFNγ were unaffected. However,

the expansion of antigen experienced CD4+ and CD8+ T cells responding to infection

was dependent on iron availability. Consequently, the Th1 polyfunction (TNFα and IFNγ

production) of the CD4+ T cells was inhibited by iron restriction in *T. gondii* infection.

Inhibition of T cell responses by iron restriction was independent from antigen load and

impacted development of long-term vaccine induced protection. CD4+ T cell specific

expression of CD71 was intrinsically required for antigen experience and polyfunction of

CD4+ and not CD8+ T cells, yet conditional CD71 knockout mice were not more

susceptible to *T. gondii* infection. These results demonstrate the importance of iron in

global development of protective immunity to *T. gondii* but also that iron-sensing by CD71

regulates CD4+ T cell activation is a critical component of long-term immunity to *T. gondii*

infection.

Materials and Methods

Ethics Statement

All animal husbandry and experiments were performed by the Gigley Laboratory

at the University of Wyoming in strict accordance to the guidelines approved by the

University of Wyoming Institutional Animal Care and Use Committee.

Mouse strains, parasite propagation, and infections

C57BL/6J (#000664), RAG1-/- (B6.129S7-Rag1$^{tm1Mom/J}$, #002216), and CD4-

CreER^{T2} (B6(129X1)-Tg(Cd4-cre/ERT2)11Gnri/J, #022356) mice were purchased from

Jackson Laboratories. CD71$^{Flx/Flx}$ mice were a kind gift from Mark Andrews, Harvard

University. CD4-CreER^{T2} mice were crossed with CD71$^{Flx/Flx}$ for two generations and the

resulting progeny were interbred to produce CD4-CreER^{T2} +/-::CD71$^{Flx/Flx}$. The genomes

of these mice contained 75% C57BL/6 heritage from the CD4-CreER^{T2} parentage and

25% 129 heritage from the CD71$^{Flx/Flx}$ parentage, thus, all experiments using these mice

used internal littermate controls. Mouse genotypes were assessed by PCR from genomic

DNA isolated from ear clips *via* digestion in DirectPCR Digestion Buffer + Proteinase K

(Viagen Biotech) using the following primer pairs: CD4-CreER^{T2} Forward (0.5µM): 5'-

GCG GTC TGG CAG TAA AAA CTA TC -3', CD4-CreER^{T2} Reverse (0.5µM): 5'- GTG

AAA CAG CAT TGC TGT CAC TT -3', and separately TfR^{Flx} Forward (0.3µM): (5'-TTC

AGT TCC CAG TGA CCA CA -3'), TfR^{Flx} Reverse (0.3µM): 5'- TCC TTT CTG TGC CCA

GTT CT -3'. The CD4-CreER^{T2} transgene was detected by a 100nt PCR product. The

310nt lox-containing CD71 allele was differentiated from the 165nt wild-type CD71 allele

by gel electrophoresis. Female mice aged 8-12 weeks old were used for all experiments.

125   Type II *T. gondii* ME49 strain was maintained by serial passage in CBA mice

126 (Jackson Laboratories #000656). Cyst numbers were counted in 10µL of brain

127 homogenate from previously infected animals placed on a glass microscope slide. Mice

128 were inoculated intragastrically (i.g.) with 7-10 cysts in 200µL Phosphate-Buffered Saline

129 (PBS) via oral gavage.

130   Type I *T. gondii* RH strain was maintained by routine passage in MRC5 fibroblasts

131 in 1 % Dulbecco's Modified Eagles Medium (1% DMEM) (1% Fetal Bovine Serum, 1X

132 Penicillin/Streptomycin, 1X Amphotericin B). Upon lysis of MRC5 monolayers,

133 tachyzoites were purified by 3µm filtration and enumerated on a hemocytometer. 1×10^3

134 parasites were injected intraperitoneally (i.p.) in 200µL PBS. Attenuated Type I *T. gondii*

135 strain *cps1-1* (CPS) was maintained by routine passage in MRC5 fibroblasts in 1% DMEM

136 supplemented with 300mM uracil. Following filtration and enumeration similar to RH

137 cultures, CPS was injected i.p. with a 27G needle at parasite concentrations as indicated

138 in 1 X PBS.

139 *Drug Formulation and Treatment*

140   Deferiprone (3-Hydroxy-1,2-dimethyl-4(1H)-pyridone "DFP", Sigma-Aldrich

141 #379409, CAS #30652-11-0) was dissolved at 10mg/mL in distilled water at 37˚C and

142 sterilized by filtration through a 0.2µm membrane. Each preparation was tested for Iron-

143 Binding Units using a colorimetric Iron Assay Kit (Abcam Cat #ab83366) according to the

144 manufacturer's instructions. Briefly, a standard curve was created using a known amount

145 of iron was prepared in Assay Buffer, and variable volumes of DFP preparations were

146 added to replicates of the top standard. Iron reducer was added to every sample to

147 measure total free iron in the solution. After addition of the iron probe, the absorbance of

the samples was measured at 593nm and the amount of free iron remaining in the sample

was extrapolated from the standard curve. The volume of DFP from each preparation

required to reduce the iron in the samples by 50% was deemed an Iron-Binding Unit (U).

DFP was administered at 75U per mouse intraperitoneally with a 27G needle every other

day starting the day preceding infection, unless otherwise indicated.

Tamoxifen (Sigma-Aldrich #T5648, CAS #10540-29-1) was dissolved at 20mg/mL

in corn oil (Sigma-Aldrich #C8267, CAS #8001-30-7) overnight by agitation at 37°C in the

dark. Tamoxifen was administered at 100mg/kg to mice intraperitoneally with a 27G

needle daily for the five days preceding infection.

ELISA

Blood samples were collected by retroorbital bleeding of isoflurane anesthetized

mice, followed by a coagulation period for 30 minutes at room temperature. Serum was

collected after centrifugation. Splenic samples were suspended at $5*10^6$ cells per well in

FACS Media and incubated for 48 hours at 37°C and 5% CO_2. Cells were lysed by 3X

freeze/thaw cycles at -80°C, and supernatant isolated for ELISA. Both serum and spleen

samples were diluted at a volume of 1:20 for IL-12p40 and 1:15 for IFNγ ELISAs, which

were performed according to manufacturer instructions using kits for IL-12p40 (Biolegend

431604) and IFNγ (Biolegend 430804).

Parasite Burdens by Genome Equivalence

Genomic DNA from specified organs was purified from the samples using the DNeasy

Blood & Tissue Kit (Qiagen 69506) according to manufacturer instructions. A standard

curve of known quantity of RH gDNA was assessed alongside 100ng-400ng of sample

DNA as template to query for the *T. gondii* unique gene B1 using SSO Advanced SYBR

Master Mix (Bio-Rad 1725017) and 0.5µM primers (F: 5'- CGT CCG TCG TAA TAT CAG

-3', R: 5'- GAC TTC ATG GGA CGA TAT G -3'). The thermocycler conditions were 1:

95°C for 10 minutes, 2: 95°C for 5 seconds, 3: 60°C for 5 seconds, 4: 72°C for 5 seconds

+ plate read, 5: Cycle 2-4, 44 repetitions, 6: Melt Curve 65°C to 95°C with 0.5°C, 5 second

increments + plate read. Parasite concentrations were extrapolated from linear regression

of the standard curve.

Plaque Assay

MRC5 fibroblasts were seeded into 6-well tissue culture plates and treated with

DFP or ferric citrate as indicated immediately prior to infection. 100 RH parasites were

inoculated per well and incubated for 7 days at 37°C and 5% CO_2. Monolayers were fixed

with 100% EtOH and stained with Plaque Assay Stain (125mL of 10% Crystal Violet in

EtOH w/v + 500mL 1% Ammonium Oxalate w/v). Plaques were visualized by projection

using a Bellco Glass, Inc. Slide Magnifier and manually counted. Plaque size was

estimated as a measurement of A = π*m((LA+TA)/4)^2, where A = Area (mm2), m =

magnification factor, LA = length of longest axis of plaque (mm), and TA = length of

transverse axis perpendicular to the longest axis (mm). Representative images were

taken with an Apple® iPhone Xr and background digitally removed with Adobe®

Photoshop.

Single Cell Suspension

For peritoneal exudate cells (PEC), the peritoneum of mice was exposed intact,

and injected with 10mL Stain/Wash Buffer with EDTA (SWBE, 5% w/v Bovine Calf Serum

and 2mM EDTA in PBS) *via* a 21G needle. Cells were dislodged by external agitation of

the cavity during gradual injection of the SWBE and redrawn in the syringe. Spleens were

extracted and crushed through a 70μm cell strainer with SWBE washing, followed by

osmotic lysis of red blood cells (RBC) by suspension in 1mL 1X RBC Lysis Buffer (Tonbo

Biosciences #TNB-4300-L100) for 3 minutes. RBC lysis was quenched by addition of

20mL SWBE, and samples were passed through a 70μm cell strainer. Single cell

suspension concentrations were estimated with a hemocytometer.

Flow Cytometry- Minimal Information About T Cell / Natural Killer Cell Assays (MIATA-

MIANKA)

Single cell suspensions were aliquoted to a clear FACS V-bottom plate at one

million cells per well. Cell viability was estimated using LIVE/DEAD Fixable Aqua dye

(ThermoFisher L34957), followed by extracellular antibody staining in SWBE, fixation and

permeabilization by BD Cytofix/Cytoperm (BD Biosciences 554714), and intracellular /

secondary staining in BD Perm/Wash buffer, all according to manufacturer instructions.

Samples were processed in Guava EasyCyte 12HT Flow Cytometer and analyzed in

FlowJo v.9.7.7 software.

For NK Cell assays, wells were coated with 0.5μg per well of NK1.1 (PK136,

BioXCell BE0036) overnight at 4°C the day prior to staining. After plating in coated wells,

cells were stimulated at 37°C and 5% CO_2 four 4 hours with 0.66X Protein Transport

Inhibitor Cocktail (PTIC, eBioscience 00-4980-03) in FACS Media (Isocove's DMEM with

L-Glutamine and 25mM HEPES supplemented with 1X Penicillin/ Streptomycin,

Amphotericin B, Non-Essential Amino Acids, 0.1mM Sodium Pyruvate, 0.2mM Glutamine

XL, 10% Fetal Bovine Serum, and 0.1mM 2-mercaptoethanol). Extracellular stain

included CD3-FITC (17A2, Biolegend 100204), CD335/NKp46-APC (29A1.4, Biolegend

137608), and CD49b-Biotin (DX5, Biolegend 108904). Intracellular / secondary stain

included IFNγ-PE (XMG1.2, Biolegend 505806) and Streptavidin-PE/Cy7 (Biolegend

405206). cNK cells were boolean gated on live cells by Live/Dead Aqua, lymphocytes by

scatter profile, and as CD3-, CD49b+, and NKp46+.

For T Cell phenotype assays, cells were plated and proceeded with direct *ex vivo*

staining. Extracellular stain included CD4-BV421 (GK1.5, Biolegend 100438), CD8-

PE/Cy5 (53-6.7, Biolegend 100710), CD11a-PE/Cy7 (I21/7, Biolegend 153108), CD62L-

APC/Cy7 (MEL-14, Biolegend 104428), CD49d-PE (9C10/MFR4.B, Biolegend 103706),

KLRG1-PE (2F1/KLRG1, Biolegend 138408), KLRG1-APC (2F1/KLRG1, Biolegend

138412), CD71/TfR1-APC (RI7217, Biolegend 113820), and intracellular stain Ki67-PE

(16A8, Biolegend 652404). Cells were boolean gated on live cell by Live/Dead Aqua,

lymphocytes by scatter profiling, and either CD4+/CD8- or CD4-/CD8+.

For T cell function assays, cells were plated in FACS media supplemented with

4µg Toxoplasma Lysate Antigen (TLA) per test and stimulated for 8 hours at 37˚C and

5% CO_2. For the last 4 hours of stimulation, PTIC was added to each well at a final

concentration of 1X. Extracellular staining included CD4-BV421 (GK1.5, Biolegend

100438) and CD8-PE/Cy5 (53-6.7, Biolegend 100710), and intracellular staining included

IFNγ-PE (XMG1.2, Biolegend 505806), TNFα-APC (MP6-XT22, Invitrogen 17-7321-82),

and GZB-FITC (QA16A02, Biolegend 372296). TLA was produced by sonication of

purified RH strain *T. gondii* parasites in PBS, and the concentration was determined by

Bradford Assay. Cells were boolean gated on live cell by Live/Dead Aqua, lymphocytes

by scatter profiling, and either CD4+/CD8- or CD4-/CD8+.

For fluorescence activated cell sorting, 20 million cells per tube were assessed for

cell viability by Live/Dead Aqua and stained without fixation with CD3-PE (17A2,

Biolegend 100206) and CD4-FITC (GK1.5, Biolegend 100406). Samples were processed

in a BD FACSMelody Cell Sorter before immediate RNA isolation. Sorting strategy is

defined in Figure S4.

For splenic profiling screens, extracellular stain proceeded immediately *ex vivo*

and the following antibodies were used: CD3-BV421 (17A2, Biolegend 100228), CD19-

BV605 (6D5, Biolegend 115540), NK1.1-FITC (PK136, Biolegend 108706), CD11b-APC

(M1/70 Biolegend 101212), Ly6C-PE (HK1.4, Biolegend 128008), Ly6G-PE/Cy7 (1A8,

Biolegend 127618), CD11c-FITC (N418, Biolegend 117306), and I-E/I-A-PE

(M5/114.15.2, Biolegend 107608). Gating strategy is defined in Figure 3A.

RNA Isolation and qRT-PCR

Samples collected in FACS experiments were pelleted and resuspended in TRI Reagent

(Sigma T9424) for RNA Isolation according to manufacturer instructions. RNA was further

purified by RNA Cleanup Protocol of RNeasy Kit (Qiagen 74104). RNA samples were

quantified by UV absorbance and cDNA synthesis was performed on normalized mass

(50ng P1 samples, 20ng P2 samples) of RNA according to manufacturer instructions with

Superscript IV Reverse Transcriptase (Invitrogen 18091050). Equal volumes of cDNA

samples were used in qRT-PCR reactions using Sso Advanced Universal SYBR Master

Mix (Bio-Rad 1725271) and validated primers (41, 42): Actin Forward (0.5µM) 5'- ACC

CAC ACT GTG CCC ATC TA -3', Actin Reverse (0.5µM) 5'- CAC GCT CGG TCA GGA

TCT TC -3', CD71 Forward (0.5µM) 5'- TCA AGC CAG ATC AGC AAT TCT C -3', and

CD71 Reverse (0.5µM) 5'- AGC CAG TTT CAT CTC CAC ATG - 3'. The PCR protocol

was 1: 95˚C for 3:00, 2: 95˚C for 0:15, 60˚C for 0:15, 72˚C for 1:20 + Plate Read, 5:

Repeat 2-5 44X, 6: Melt Curve 65.0˚C to 95.0˚C, 0.5˚C increment for 0:05 + Plate Read.

mRNA ratios were estimated using the ΔΔCt method.

Statistical Analysis

Statistics and graph generation were completed using Graphpad Prism v.9 and

Microsoft Excel v16.54 software. Experiments are pooled from at least two independent

technical repeats containing 3-5 replicates per experiment. Statistics were performed

between two groups by Student's t-test or between multiple groups by One-Way ANOVA,

or by Kaplan-Meier Survival analysis. Significance is indicated by $*p<0.05$, $**p<0.01$,

$***p<0.005$, and $****p<0.001$. Figures were assembled in Adobe Illustrator 2021,

references in Endnote X8.

Results

Iron is an essential nutrient for Toxoplasma gondii

To study the roles of iron in *Toxoplasma gondii* pathogenesis and immunity, it was

necessary to induce a rapid depletion of iron at the onset of an infection. To this end, we

selected use of the small molecule cell permeable iron chelator deferiprone (DFP) (43).

Based on lot-to-lot differences of iron chelation ability, we developed experimental

reproducibility by normalization of iron-binding efficiency in each formulation. In this

assay, the redox ability of free iron reacts with Ferense S, a chromogen to produce a

colorimetric readout linearly correlated with free iron concentration. We challenged a

known amount of iron, 10nM, with a range of DFP amounts by batch. The required mass

of each formulation to decrease the free iron to 5nM, or 50% binding, was deemed an

Iron-Binding Unit (U) (Figure 1A).

Dose titrating DFP into an infection setting showed a dose-dependent inhibition of

parasite replication by plaque assay (Figure S1A) both in plaque number (Figure S1B)

and plaque size (Figure S1C). The minimal dose tested that was sufficient to completely

prevent plaque formation was 1.37U DFP, therefore, we tested this amount of DFP in a

titration of a range of ferric citrate supplementation. A minimal dose of 20µg ferric citrate

was sufficient to rescue the inhibition of parasite growth (Figure S1D). Lower doses of

ferric citrate alone did not increase numbers of plaques observed (Figure S1E). However,

at higher doses, which overcame DFP inhibitory effects, ferric citrate alone displayed

toxicity to the host monolayer integrity (Figure S1A). This agrees with previous studies

demonstrating the role of vacuolar Iron transporter (VIT) being important for regulating

iron homeostasis in *T. gondii* (10). In summary experiments, DFP prevented *T. gondii*

plaque formation in an iron-dependent manner (Figure 1B-C).

Iron restriction "therapy" increases morbidity and mortality of T. gondii infection

Given the parasite-inhibitory properties of DFP *in vitro* we tested how iron chelation

with DFP impacted infection pathogenesis *in vivo*. DFP is currently used in human

patients as a safe iron overload therapy (43). During Type II ME49 strain infection in

mice, treatment with DFP resulted in increased weight loss and mortality compared to

non-treated controls (Figure 1D-E). The difference between iron chelation inhibition of

proliferation *in vitro* and increased morbidity and mortality *in vivo* was surprising.

Therefore we measured acute parasite burden in tissues of infected mice acute (14 DPI)

and chronic (5 weeks PI, 5WPI) infection (20). DFP unexpectedly increased

dissemination by 4-fold as assessed by parasite burden in the lung at 14DPI (Figure 1F),

and by encystation in the brain at 5WPI (Figure 1G). These results indicate DFP restriction

of iron impacts not only parasite proliferation, but dissemination and possibly

development of immunity to *T. gondii*.

IL-12 and IFNγ production is reliant on available iron in acute infection

IL-12 dependent IFNγ production is the key pathway for resistance to *T. gondii*, but

also may contribute to inflammatory tissue damage (18, 22, 44). Therefore, changes in

expression of these cytokines could correlate with the increased pathogenesis due to

DFP-induced iron restriction. To determine how DFP treatment impacted development of

immune control to *T. gondii* we compared IL-12 and IFNγ levels in naïve mice, ME49

infected mice and mice infected with ME49 and treated with DFP. As expected, *T. gondii*

infection significantly increased IL-12 and IFNγ in spleen cells (IL-12: 17-fold, IFNγ: 32-

fold) and serum (IL-12: 4-fold, IFNγ: 9-fold) compared to naïve animals (Figure 2). Iron

restriction by DFP treatment significantly lowered the levels of these cytokines in both

spleen cells and serum. These observations suggest that iron is a key factor in the

activation of the immune response to *T. gondii*.

Iron restriction does not impact splenocyte composition in acute infection

Because we observed significantly lower IL-12 and IFNγ during the early acute

infection we further explored innate immune cell recruitment at this stage of infection.

Neutrophils, Monocytes, Macrophages and Dendritic cells are all important producers of

IL-12 (45, 46). Group 1 ILCs including natural killer cells and ILC1 are important early

sources of IFNγ to initiate control of *T. gondii* (21-23). Analysis of spleen immune cell

composition during acute ME49 infection with and without iron restriction by DFP revealed

that restricting iron availability with DFP treatment (Figure 3A-B) did not have a significant

impact on the frequencies of (CD19+ CD3-) B cells (Figure S2B), (CD19- CD3+ NK1.1-)

T Cells (Figure S2C), (CD19- CD3- NK1.1+) NK Cells (Figure S2D), (Lineage- CD11b+

Ly6Chi Ly6G-) inflammatory monocytes (Figure S2E), (Lineage- CD11b+ Ly6Cint Ly6G+)

neutrophils (Figure S2F), and (Lineage- CD11c+ MHCII+) Dendritic cells (Figure S2G).

Notably, the total live lymphocyte numbers in spleen were significantly reduced at 5DPI

only in iron restricted and infected animals, from $5.5\text{-}6*10^7$ cells to $4*10^7$ (Figure S2A).

These results suggested that the overall composition of splenocytes could not explain the

differences in morbidity, mortality and dissemination.

To further test if the innate cell composition is affected by iron restriction, we

infected Rag1[tm1Mom] (RAG1-/-), mice which are deficient in adaptive B or T cells (47), with

cysts of ME49. RAG1-/- mice started to succumb to the infection starting at 2-3 WPI,

however, iron restriction by DFP treatment during infection of these animals did not

significantly change the mortality of the infection (Figure 3C). Taken together, these

results suggest that the level of available iron impacts some innate immune cell function,

but not cellular composition during acute *T. gondii* infection.

Iron does not regulate NK cell number or IFNγ production.

NK cells are critical for early T cell independent control of *T. gondii* infection (18,

21, 22, 48, 49). Even though there was no difference in mortality due to iron restriction in

RAG1-/- animals who have an intact NK cell compartment, there was a significantly

impaired systemic IFNγ production in WT animals. Therefore, we assayed NK cell (CD3-

CD49b+ NKp46+) function in the presence or absence of iron restriction with DFP at 5

DPI in the spleen. The frequency and number of the NK cells dropped significantly during

acute infection compared to naïve animals. However, the decrease in NK cells was

similar regardless of DFP treatment (Figure 4A). NK cells were activated to produce IFNγ

regardless of iron restriction by DFP suggesting that NK cells do not rely on available iron

levels for their response (Figure 4B).

Iron is required to generate antigen experienced CD4+ and CD8+ T cells

Our results above suggest innate immune cell activity during iron restriction with

DFP is relatively intact leading us to hypothesize that iron regulates and is essential for

development of adaptive immune responses to *T. gondii*. We tested this hypothesis by

examining the CD4+ and CD8+ T cell responses at the beginning of the adaptive

response at 10 DPI with ME49 cysts by flow cytometry (50, 51). As expected, the

frequency and number of total splenic helper CD4+ and cytotoxic CD8+ T cells was

reduced after *T. gondii* infection compared to naïve animals (Figure 5A). In the iron-

restricted infection group, this reduction in CD4+ and CD8+ T cells was comparable

(Figure 5A). To better determine whether and how iron regulates T cell responses to *T.*

gondii infection, antigen experienced T cells were assayed by flow cytometry. T cell

upregulation of CD11a and tandem expression of CD49d is strongly correlated with

markers of activation, thus, CD11a and CD49d (CD11a(Hi) CD49d+) serve as efficient

surrogate markers of antigen experience (52). The frequency of the CD11a(Hi) CD49d+

population of antigen experienced CD4+ T cells increases from 5.7% in naïve animals to

47.3% in *T. gondii* infected animals without iron restriction (Figure 5B). Similarly, the

frequency of CD11a(Hi) CD49d+ CD8+ antigen experienced T cells increased from 5.6%

to 43.8% after *T. gondii* infection without iron restriction (Figure 5C). Absolute numbers

of antigen experienced CD4+ and CD8+ T cells also increased significantly after infection

in non-DFP treated animals compared to naïve controls (Figure 5B-C). However, the

frequency of antigen experienced cells only reached 27.4% in CD4+ T cells and 22.1%

in CD8+ T cells (Figure 5B-C). Moreover, the absolute number of antigen experienced

CD4+ and CD8+ T cells under iron restriction was not significantly higher than in the naïve

group. Iron restriction by treating animals with DFP during infection thus significantly

inhibited increases in both the frequency and absolute number of antigen experienced

CD11a(Hi) CD49d+ T cells compared to naïve and infected non-DFP treated animals.

Iron is required for functional CD4+ T cell polyfunctional response

Due to the inhibitory effect of iron restriction on increases in T cell antigen

experience, it was likely that the functions of these cells were also inhibited. Therefore

we measured function of CD4+ (IFNγ and TNFα) and CD8+ T cells (IFNγ) *ex vivo* from

mice infected with ME49 and treated or not with DFP to restrict iron (53-57). The

frequency of polyfunctional IFNγ+ / TNFα+ CD4+ T cells increased from 0.017% to 13.8%

after infection (Figure 5D). Additionally, the frequency of IFNγ-producing CD8+ T cells

increased from 0.44% in naïve to 8.7% after infection (Figure 5E). The inhibitory effect of

iron restriction on antigen experience in CD4+ T cells translated to inhibited function, as

the frequency of polyfunctional CD4+ T cells was 6.5%, significantly lower than untreated

infected animals, but still significantly higher than naïve animals (Figure 5D). In contrast,

iron restricted animals still mounted a functional CD8+ T cell response, as the frequency

of IFNγ+ CD8+ T cells (6.1%) was not significantly different than in untreated infection

(Figure 5E).

Antigen experienced CD4+ and CD8+ T cells upregulate CD71 expression

CD71 mediates iron uptake in most mammalian cell types, including immune cells

(58). Additionally, CD71 is both upregulated and required for T cell activation *in vitro* (59-

61). To link the inhibitory effect of iron restriction on T cell responses to T cell intrinsic iron

uptake, we measured CD71 expression levels on CD4+ and CD8+ T cells during *T. gondii*

infection using flow cytometry (Figure 5F-I). We observed a significant 2.7-fold increase

in CD71 MFI on CD4+ T cells and a 2.4-fold increase on CD8+ T cells during *T. gondii*

infection. In iron restricted animals, CD71 MFI of total CD4+ or CD8+ T cells was not

significantly different compared to untreated infection groups (Figure 5F-G). We next

measured CD71 MFI in antigen experienced CD11dHi CD49a+ T cells. Similar to total

CD4+ and CD8+ T cells, CD71 increased in expression after *T. gondii* infection on

CD11dHi CD49a+ antigen experienced CD4+ and CD8+ T cells compared to naïve

conditions (Figure 5H-I). The MFI of CD71 was significantly higher on antigen

experienced CD4+ and CD8+ T cells on a per cell basis (Figure 5H-I) than total CD4+

and CD8+ T cells (Figure 5F-G) from both non-DFP treated and iron restricted DFP

treated infected mice.   Iron restriction by DFP treatment also significantly increased CD71

MFI on antigen experienced CD4+ T cells compared to cells from non-treated infected

animals (Figure 5H). Interestingly, iron restriction by DFP treatment did not impact the

CD71 MFI of antigen experienced CD8+ T cells (Figure 5I). These results show that CD71

is upregulated on T cells during *T. gondii* infection, and that iron restriction by DFP

treatment further upregulates CD71 only in antigen experienced CD4+ T cells.

Iron restriction reduces frequency of effector CD8+ T cell subsets

Iron restriction significantly decreased the number of antigen experienced CD8+ T

cells during *T. gondii* infection. To provide control of the parasite, CD8+ T cells

differentiate from naïve (CD62L+KLRG1-, FI) into memory precursor (CD62L-KLRG1-,

FII), short lived effector cell (CD62L-KLRG1Hi, FIII), and CD62L+ effector cells

(CD62L+KLRG1+, FIV) (Figure 5J) (62). Therefore we measured CD8+ T cell

differentiation in the presence or absence of iron restriction by DFP treatment during *T.*

gondii infection defined by functional output (Figure 5J). The frequency of the naïve FI

population decreased after infection from 93.55% of naïve animals to 62.65% in normal

iron condition (non-DFP treated) animals. In the iron restricted (DFP-treated) group, there

was a significantly greater proportion of CD8+ T cells remaining naïve at 75.73%

compared to non-treated infected mice. After infection in non-DFP treated mice, CD8+ T

cells differentiated into FII memory precursor subset (CD62L- KLRG1-, 13.38%), the FIII

effector subset (CD62L- KLRG1+, 13.06%), and the FIV CD62L+ effector subset

(CD62L+ KLRG1+, 10.89%). Iron restriction by DFP treatment inhibited differentiation of

FIII (6.01%) and FIV (5.61%) effector subsets, both of which were significantly lower than

in untreated infection and higher than naïve mice. However, the proportion CD8+ T cells

belonging to the FII memory precursor subset (12.65%) was not significantly different

between the DFP-treated or non-treated infection groups.

Reduced T cell activation during iron restriction is independent of antigen load and

occurs during vaccination.

One of the reasons we may observe decreased antigen experienced CD4+ and

CD8+ T cells and CD4+ T cell polyfunctional responses is the lack of sufficient *T. gondii*

antigen due to the possibility that iron restriction slows *T. gondii* growth *in vivo*. The

quantity and quality of antigen load is linked to differences in T cell activation and

protection in *T. gondii* infection (25, 63, 64). Therefore to control for differences in antigen

levels that might cause defects in T cell responses during DFP induced iron restriction,

we immunized mice i.p. using 1×10^6 tachyzoites of the non-replicating vaccine strain of

T. gondii, CPS (24, 65) with or without DFP induced iron restriction. We analyzed the

CD4+ and CD8+ T cell responses at the site of infection (PEC, Figure 6) and the systemic

response on day 7 p.i. (spleen, Supplemental Figure 3).

In the PEC, the frequency and absolute number of total CD4+ and CD8+ T cells

increased after vaccination (Figure 6A). Iron restriction using DFP significantly reduced

those increases and were not significantly greater than naïve mice (Figure 6A). The

frequency and number of antigen experienced (CD11aHi CD49d+) CD4+ and CD8+ T

cells were then measured in the PEC of vaccinated mice with and without DFP iron

restriction. In naïve animals, there was a high baseline frequency of CD11a(Hi) / CD49d+

antigen experienced CD4+ and CD8+ cells. The frequency and absolute number of

antigen experienced CD4+ and CD8+ T cells significantly increased in response to

vaccination with CPS compared to naïve controls (CD4+ frequency: 2.33-fold, number:

7.31-fold and CD8+ frequency: 2.13-fold, number: 7.43-fold). In contrast, DFP induced

iron restriction resulted in a near complete inhibition of increased antigen experienced

CD4+ T cell populations with only 1.13-fold (frequency) and 1.38-fold (number) non-

significant increases after vaccination (Figure 6B). Similarly, DFP induced iron restriction

inhibited CD8+ T cell populations with 1.67-fold (frequency) and 2.15-fold (number) non-

significant changes compared to unvaccinated groups (Figure 6C). Consequently, we

measured CD4+ T cell polyfunction (IFNγ+/TNFα+) and CD8+ T cell single function

(IFNγ+) after CPS vaccination. Both the frequency and absolute number of single and

polyfunctional CD4+ and CD8+ T cells were significantly increased after vaccination in

the PEC. DFP induced iron restriction significantly reduced CD4+ and CD8+ T cell

functions in both frequency and absolute number, and for CD8+ T cells, frequency and

absolute number of functional cells were not significantly different from naïve animals

(Figure 6D-E).

We next measured the ratio of naïve (CD62L+KLRG1-, FI), memory precursor

(CD62L-KLRG1-, FII), short lived effector (CD62L-KLRG1+, FIII), and CD62L+effector

(CD62L+KLRG1+, FIV) CD8+ T cells in PEC of naïve, CPS immunized/non-treated and

CPS immunized/DFP-treated iron restricted mice (Figure 6F). We observed an increase

in naïve CD8+ T cells (FI) after CPS vaccination compared to non-immunized animals

and iron restriction further increased the frequency of these cells compared to naïve and

CPS alone. Memory precursor CD8+ T cells (FII) decreased significantly after CPS

vaccination with or without DFP treatment and DFP did not affect this decrease. The

frequency of effector CD8+ T cells (FIII) increased significantly after CPS vaccination

compared to naïve animals and DFP induced iron restriction significantly decreased this

cell population to lower levels than naïve and CPS infected mice. There was only a

moderate effect on CD8+ CD62L+ effector T cells with CPS immunization and CPS with

DFP treatment compared to naïve mice.

In the spleen, the response to CPS is more subtle due to the localized immune

response induced by this parasite strain in the PEC (62, 66). Measuring the CD4+ and

CD8+ T cell responses in the periphery revealed that even though the frequency and

absolute number of total T cells were unchanged (Figure S3A), antigen experience and

function increased significantly after vaccination (Figure S3B-E). Similar to PEC these

increases were inhibited in DFP induced iron restriction conditions and were not greater

than naïve mouse cells. These results demonstrate that iron-restriction in CPS

vaccination when antigen availability is normalized still significantly decreases T cell

responses similar to ME49 infection, revealing the role of iron as a critical factor involved

for the proper activation of T cell immunity to *T. gondii* independent of antigen level.

Iron-restricted inhibition of adaptive immunity also reduces vaccine induced long-

term protective immunity against lethal T. gondii rechallenge.

Iron-restriction inhibits CD4+ and CD8+ T cells responses during primary infection

of both ME49 and CPS strains. How iron impacts development of long term memory and

control of *T. gondii after* vaccination with CPS has not been tested. CPS vaccination

induces robust T cell memory that is highly protective against lethal challenge (25). Mice

were immunized with a dose titration of CPS (5 X 10^4, 1 X 10^5, and 5 X 10^5) tachyzoites

and treated or not with DFP for the first three weeks after vaccination. Mice were then

allowed to recover without DFP treatment to re-establish normal iron homeostasis for two

weeks. All mice were then challenged with a lethal dose (1×10^3) of the Type I CPS

parental strain RH i.p and monitored for survival. As shown in Figure 6E, 5×10^4

tachyzoites were not protective and 1×10^5 were partially protective against lethal

challenge. 5×10^5 CPS tachyzoites induced near-complete protective immunity to lethal

challenge. DFP induced iron restriction decreased CPS vaccine dose efficacy in both 1

$\times 10^4$ and 5×10^5 CPS animals. This result demonstrates that iron is critical during

immunization and iron restriction prevents development of immune memory to *T. gondii*.

CD4+ T cell intrinsic CD71 is required for development of CD4+ T cell responses

during T. gondii infection

Our results demonstrate that iron is critical for CD4+ and CD8+ T cell responses

during *T. gondii* infection and this could be why DFP induced iron restriction results in

higher morbidity, mortality and cyst burdens in the CNS. However, iron restriction

appeared to impact CD4+ T cells more than CD8+ T cells. In addition, CD4+ T cells also

upregulated CD71 more than CD8+ T cells after infection, and was further increased with

DFP induced iron restriction. Therefore, we hypothesized that CD4+ T cell intrinsic CD71

was required for their activation and function. To test this hypothesis, we generated CD4-

CreER^{T2} :: CD71$^{Flx/Flx}$ mice for tamoxifen induced conditional knockout of CD71 in CD4+

T cells (Figure S4).

CD4+ conditional CD71 knockout mice (CD4cCD71KO) were infected with ME49

cysts and the CD4+ and CD8+ T cell frequency and absolute numbers were measured

by flow cytometry. The frequency and number of total CD4+ T cells in CD4cCD71KO

compared to WT CreER^{T2} negative littermate controls was significantly reduced (Figure

7A, C). Surprisingly, the total CD8+ T cell population, not non-T cell populations,

increased in frequency and absolute number in the CD4cCD71KO mice (Figure 7A, B,

D). As further validation of the model, CD71 expression was significantly reduced on the

CD4+ T cells in the CD4cCD71KO compared to WT littermate controls (Figure 7E). There

was no change in TfR1 expression in CD8+ T cells (Figure 7F). To determine why the

CD4+ T cell population was reduced we measured their proliferation using the

proliferation marker Ki67 (67). Ki67 expression was reduced in CD4+ T cells from the

CD4cCD71KO group by 52.7% compared to WT littermate controls (Figure 7G).

To examine how iron uptake via CD71 in CD4+ T cells impacted the immune

response to *T. gondii*, we measured antigen experience (CD11aHiCD49d+) and

functionality of CD4+ and CD8+ T cell subsets in CD4cCD71KO mice and WT littermate

controls after ME49 infection. The frequency of antigen experienced CD4+ T cells in the

CD4cCD71KO + tamoxifen group was 38.90%, significantly lower than the non-tamoxifen

treated CD4cCD71KO (57.20%) and WT littermate controls with or without tamoxifen

(53.56% and 58.01%, respectively) (Figure 8A). CD4+ T cells are required to maintain

CD8+ T cell responses and memory development (26). To our surprise, the frequency

and absolute number of antigen experienced CD11aHi CD49d+ CD8+ T Cells were not

significantly different comparing CD4cCD71KO (58.33% in untreated and 64.84% in

tamoxifien-treated) WT littermate controls (59.51% in untreated and 55.23% in tamoxifen-

treated) (Figure 8B). The frequency and absolute number of polyfunctional CD4+ T cells

was highly dependent on CD71 (Figure 8C). The polyfunctional (IFNγ+/TNFα+) CD4+ T

cell proportions were reduced to 1.13% in tamoxifen-treated CD4cCD71KO from 11.46%

and 6.78% in the littermate controls and 10.76% non-tamoxifen treated CD4cCD71KO

(Figure 8C). There was no significant difference in the functional CD8+ T cell response

across all groups (Figure 8D) suggesting that iron largely regulates the ability of the CD4+

T cells to expand and function during infection independent of CD4+ T cell help to

developing CD8+ T cell responses. To further test this possibility we infected all mouse

groups with ME49 and measured survival. No significant difference in survival was

observed comparing CD4cCD71KO to WT litter mate controls with or without tamoxifen

treatment (Figure 8E), supporting the importance of iron uptake for CD4 function, but not

for helping CD8+ T cell responses develop.

Discussion

Competition between host and pathogen for nutritional iron shapes the

pathogenesis and outcome of infections. Here, we investigated this complex interaction

during *T. gondii* infection. Our results demonstrate that iron is not only essential for

Toxoplasma gondii replication, but also for optimal host immune responses because

restricting iron *in vivo* results in higher morbidity and mortality in mice. During iron

restriction with DFP, splenic and systemic IL-12 and IFNγ levels were significantly

reduced in acute infection. However, in the absence of adaptive immune cells in RAG1-

/- mice, iron restriction does not further enhance mortality due to infection. The immune

cells impacted the most by iron restriction were CD4+ and CD8+ T cells, as antigen

experience of both and CD4+ functionality were dependent on iron. Importantly, iron

restriction inhibition of CD4+ and CD8+ T cells was independent of antigen load and was

essential for development of vaccine induced long-term protection against lethal *T. gondii*

challenge. Both CD4+ and CD8+ T cells upregulate CD71 protein expression in response

to *T. gondii*, and especially so by antigen experienced CD4+ T cells when iron is limited.

Lastly, we find that CD71 expression is required for expansion of antigen experienced

and polyfunctional CD4+ T cell in a cell intrinsic manner. Taken together, these results

suggest that iron homeostasis is a required nutrient for efficient CD4+ and CD8+ T cell

responses to *T.* gondii and regulated by CD71 in CD4+ T cells. The implications of this

study suggest that iron nutrition is an extenuating factor not only for resisting primary

infections, but with long lasting effects even after iron balance is restored for vaccine

efficacy.

The absolute requirement for iron in *T. gondii* biology remains an attractive target

for therapeutic intervention, however, restricting iron could also have a negative effect on

the development of immunity to this infection. The levels of IL-12 and IFNγ at 5DPI were

reduced by iron restriction in both serum and spleen, but not completely absent. IL-12 is

produced by the myeloid cell populations including neutrophils, monocytes, macrophages

and dendritic cells during *T. gondii* infection (45, 46). Iron overload can affect the

polarization and differentiation of monocytes into macrophages via mitochondrial

oxidative stress (32, 33). Treatment of iron deficiency with intravenous iron can affect the

differentiation of monocytes into macrophages and DCs (34). Although iron homeostasis

could be important in these roles, we did not observe any significant difference in the

ratios of these cells in the spleen during acute infection, suggesting that DFP induced iron

restriction may not significantly impair myeloid cell cellular composition during *T. gondii*

infection. IL-12-dependent IFNγ produced by NK cells is critical for early control of *T.*

gondii and reaches its peak by day 5 post infection (22). Our results reveal NK cell

responses in DFP induced iron restriction conditions remained intact at that time point

compared to normal iron conditions. These results further support that iron may not have

an important role in innate immunity to *T. gondii*. Taken together, this could indicate that

parasite detection by the innate immune system was impaired and translates to lower

cytokine production *in vivo,* but the potential capacity of NK cell IFNγ production is

independent of iron homeostasis. Further exploration of the impact of iron in innate

immunity to *T. gondii* is warranted to answer these questions.

Available iron and CD4+ T cell intrinsic CD71 are critical factors for the

development of adaptive immunity to *T. gondii* infection. Blockade of CD71 transporting

iron on human CD4+ T cells prevents their proliferation after anti-CD3/CD28 activation

(38).   Iron status can also affect the differentiation of T helper cell type, potentially

impairing development of proper T helper populations important for controlling disease

(37, 39, 40). Our results demonstrate that iron levels do impact the development of T cell

responses during *T. gondii* infection. Expansion of antigen experienced CD11a(Hi)

CD49d+ CD4+ and CD8+ T cells and CD4+ T cell functionality were inhibited in iron

restriction conditions during *T. gondii* infection and after vaccination. CD4+ T cell intrinsic

CD71 was essential for antigen experienced cell expansion and optimal polyfunctional

responses. There are several possibilities that may explain the role of iron and CD71 in

T cell expansion and activation during *T. gondii* infection.

Iron levels in a T cell can be sensed and could provide post transcriptional

regulation of critical genes for T cell activation, proliferation, function, and differentiation.

Iron is a known cofactor for Iron-Responsive Element (IRE) Binding Protein 1 (IRP1),

which is a regulator of translation for many proteins, such as CD71, by differential binding

of mRNA IRE's (68). T cell transcription factors such as Tbx21, Eomesoderm, PRDM1,

Akt/mTor, NFκB, TCF1, and STATs, all critical for T cell activation, could contain IREs in

their 5' and 3' UTRs that IRP1 could then bind to regulate their expression depending on

cellular iron levels. There is evidence that at least Tbx21 and STAT1 may be mitigated by

iron levels (69, 70).

T cell activation requires a metabolic shift to support their activation, proliferation,

differentiation, and function to control diseases. Naïve, memory, and effector T cells

exhibit distinct metabolic profiles (71-75) and undergoing a metabolic shift once they

receive proper activation signals for antigen recognition, costimulation, and cytokines (76,

77). Iron and uptake via CD71 may supply sufficient iron levels for energetic demands of

T cell activation. In addition to regulating mRNA stability and translation, IRP1 also works

to regulate the citric acid cycle (TCA cycle) as aconitase when iron sulfur clusters are

bound (78). Our study may have revealed how iron uptake is a critical fourth signal to

turn on metabolic activity for CD4+ and CD8+ T cell expansion and function and how this

is also critical for vaccine efficacy against this infection. Interestingly, our results also

demonstrate that iron and CD71 mediated iron uptake is more important for CD4+ T cells

to become activated than CD8+ T cells.

The activation, proliferation, differentiation and function of T cells can be regulated

by critical second messengers: reactive oxygen species (ROS) produced inside of T cells.

Iron and CD71 dependent iron uptake could impact ROS levels thereby affecting their

intracellular signaling for activation. Iron reacts with hydrogen peroxide to produce

hydroxy radicals (OH) by the Fenton reaction (79). The level of ROS and hydroxy radicals

can be sensed by glutathione which then in turn regulates T cell proliferation (80). ROS

can also be detected by the Nrf2-KEAP1-Cul3 which releases Nrf2 from the complex to

become an active suppressor of T cell activation (80). Maintenance of moderate levels of

ROS is also essential for proper TCR signaling that then can enhance T cell activation,

proliferation, differentiation, and function (81-84). Understanding how iron and CD71

uptake impacts ROS signaling in T cells during *T. gondii* infection will be important future

research.

CD71 may function independently of iron uptake in T cell activation. CD71 has

been observed to co-localize with the T Cell Receptor (TCR) complex and to be a

functional signaling component of the immune synapse (85). The mechanism(s) of how

CD71 interacting with the TCR complex influence the level of T cell activation are

unknown. CD71 is phosphorylated by Protein Kinase C (PKC) at position serine 24 in the

cytoplasmic domain of CD71 away from the transferrin binding site (86, 87). Endocytosis

of CD71 is required for iron uptake, but the serine 24 residue is dispensable for it,

suggesting a potential still-undiscovered function of CD71 phosphorylation (88-90). Since

PKC is central to TCR engagement downstream of CD3 and CD28 signaling (91),

phosphorylation of CD71 by PKC would spatiotemporally position CD71 to contribute to

signaling as part of the TCR complex. CD71 expression is ultimately controlled by IRP1-

dependent iron sensing, therefore the secondary potential signaling function of CD71 in

the TCR could act as another checkpoint for adequate iron levels to support T cell

activation.

CD4+ and CD8+ T cells' reliance on iron was different depending on strain of *T.*

gondii infection. Our results indicate that iron may be more essential for regulating CD4+

T cell effector responses than CD8+ T cell effector responses during Type II *T. gondii*

strain acute infection, but that effector responses of both CD4+ and CD8+ T cells are

equally reliant on available iron after vaccination with the attenuated vaccine strain CPS.

Antigen processing *T. gondii* infection can occur through cross presentation,

phagocytosis, or autophagy (92-94). In typical type II ME49 infections, proliferation of the

parasite is coupled with antigen processing, but in DFP induced iron limiting conditions,

the parasite proliferation and dissemination are altered. This may differentially affect

modes of antigen processing and presentation to CD4+ T cells compared to CD8+ T cells.

For example, since iron levels influence autophagic potential of cells (95) and autophagy

is a major route for MHCII antigen processing (96), it is plausible that modulation of

autophagy by iron restriction preferentially inhibits antigen processing important for

antigen experience for both CD4+ and CD8+ T cells but inhibits function of only CD4+ T

cells in ME49 studies. In contrast, in CPS vaccination studies, the parasite cannot

proliferate, thus, the potential differing effects of iron restriction on antigen processing and

presentation to CD4+ or CD8+ T cells would be equal, providing some explanation as to

why we observed inhibition of both CD4+ and CD8+ T cells in terms of antigen experience

and function. Additionally, in both CPS and ME49 strain infections, the proportion of CD8+

FIII and FIV effector subpopulations, not the FII memory precursor subpopulation, were

inhibited by iron restriction, suggesting that the type and strength of the CD8+ T cell

response relies on iron as much as CD4+ T cell polyfunctional response.

In addition, CD4+ T cells may be more reliant on iron uptake by CD71 than CD8+

T cells. Our data may support this possibility because we observed that only antigen

experienced CD4+ T cells upregulated their CD71 to a higher level in iron limiting

conditions with DFP. CD4+ T cells are required to provide help for the activation,

maintenance of activation, and memory cell differentiation of CD8+ T cells during *T. gondii*

infection (24, 26, 66). The conditional knockout of CD71 in CD4+ T cells reduced overall

frequencies and number of CD4+ T cells, CD4+ T cell antigen experience, and CD4+

functionality. These results demonstrate that CD71 is critical for development of CD4+ T

cell responses to *T. gondii*. Based on these results we were expecting that there would

be a decrease in CD8+ T cell responses due to the lack of CD4+ T cell help in the absence

of CD71. However, this was not the case; within the CD8+ T cell compartment, there was

no change of antigen experience or functionality in the CD4cCD71KO mouse treated with

tamoxifen compared to all other groups. Intriguingly, the reduced frequency of total CD4+

T cells in the spleen in the CD4cCD71KO mice treated with tamoxifen was compensated

by an increase in total CD8+ T cell frequencies and numbers. It is possible that the

compensatory increase of overall CD8+ numbers could additionally compensate for

reduced protection provided by CD4+ T cells in the conditional knockout, resulting in

similar disease outcomes. A recent study has demonstrated that CD71 antibody blockade

that inhibits iron uptake by human CD4+ T cells still results in generation of accessory

helper T cells *in vitro* (38). An alternative explanation for our results then is that by

knocking down expression of CD71 in CD4+ T cells, the generation of these accessory

CD4+ T cell helper cells is retained. While overall functionality and expansion of antigen

experienced CD4+ T cells is reduced, these cells might provide CD8+ T cells with the

help they need to continue to function and develop memory. This is further supported by

our survival data that shows no difference in long-term survival between CD4cCD71KO

tamoxifen treated and WT litter mate controls. Further studies of the role of CD71 in the

T cell responses to *T. gondii*, whether iron-dependent or -independent, will be helpful to

uncover additional roles for iron as a fourth signal regulating T cell immunity during this

infection.

In summary, the results of this study confirm the nutritional requirement of iron for

both *Toxoplasma gondii* and its corresponding immune response and defined the adverse

outcome of mice that do not meet this nutritional need. The mechanism driving the

adverse outcome is explained by defective CD4+ and CD8+ T cell responses. Optimal

CD4+ T cells responses were mediated by iron uptake receptor Transferrin Receptor 1.

Many questions remain and will be addressed in future studies.

Acknowledgements

The principal investigator and corresponding author of this project is J.P.G. Project

conception was conducted by S.L.D. and J.P.G. Experiments were designed by S.L.D.

and performed by S.L.D., T.R., M.A., H.K.K., and S.K.N. Data analysis was conducted by

S.L.D., T.R., and M.A. Intellectual contributions were made by S.L.D. and J.P.G.

Technical expertise was provided by J.P.G. Manuscript preparation was performed by

S.L.D. and J.P.G., with contributions by all authors on revisions. Funding securement and

project management was performed by J.P.G.

References

1.   Pappas, G., N. Roussos, and M. E. Falagas. 2009. Toxoplasmosis snapshots:

global status of Toxoplasma gondii seroprevalence and implications for pregnancy

and congenital toxoplasmosis. *International journal for parasitology* 39: 1385-

1394.

2.   Luft, B. J., and J. S. Remington. 1988. AIDS commentary. Toxoplasmic

encephalitis. *The Journal of infectious diseases* 157: 1-6.

3.   Tenter, A. M., A. R. Heckeroth, and L. M. Weiss. 2000. Toxoplasma gondii: from

animals to humans. *International journal for parasitology* 30: 1217-1258.

4.   Fox, B. A., J. P. Gigley, and D. J. Bzik. 2004. Toxoplasma gondii lacks the enzymes

required for de novo arginine biosynthesis and arginine starvation triggers cyst

formation. *International journal for parasitology* 34: 323-331.

5.   Krug, E. C., J. J. Marr, and R. L. Berens. 1989. Purine metabolism in Toxoplasma

gondii. *J Biol Chem* 264: 10601-10607.

6.   Romano, J. D., S. Sonda, E. Bergbower, M. E. Smith, and I. Coppens. 2013.

Toxoplasma gondii salvages sphingolipids from the host Golgi through the

rerouting of selected Rab vesicles to the parasitophorous vacuole. *Mol Biol Cell*

24: 1974-1995.

7.   Ma, E. H., G. Bantug, T. Griss, S. Condotta, R. M. Johnson, B. Samborska, N.

Mainolfi, V. Suri, H. Guak, M. L. Balmer, M. J. Verway, T. C. Raissi, H. Tsui, G.

Boukhaled, S. Henriques da Costa, C. Frezza, C. M. Krawczyk, A. Friedman, M.

Manfredi, M. J. Richer, C. Hess, and R. G. Jones. 2017. Serine Is an Essential

Metabolite for Effector T Cell Expansion. *Cell metabolism* 25: 345-357.

8.   Tanaka, Y., Y. Nagai, M. Okumura, M. I. Greene, and T. Kambayashi. 2020.

PRMT5 Is Required for T Cell Survival and Proliferation by Maintaining Cytokine

Signaling. *Front Immunol* 11: 621.

9.   Nagafuku, M., K. Okuyama, Y. Onimaru, A. Suzuki, Y. Odagiri, T. Yamashita, K.

Iwasaki, M. Fujiwara, M. Takayanagi, I. Ohno, and J. Inokuchi. 2012. CD4 and

CD8 T cells require different membrane gangliosides for activation. *Proceedings*

of the National Academy of Sciences of the United States of America 109: E336-

342.

10.   Aghabi, D., M. Sloan, Z. Dou, A. J. Guerra, and C. R. Harding. 2021. The vacuolar

iron transporter mediates iron detoxification in Toxoplasma gondii. *bioRxiv*.

11.   Almeida, M. P. O., E. A. V. Ferro, M. P. P. Briceño, M. C. Oliveira, B. F. Barbosa,

and N. M. Silva. 2019. Susceptibility of human villous (BeWo) and extravillous

(HTR-8/SVneo) trophoblast cells to Toxoplasma gondii infection is modulated by

intracellular iron availability. *Parasitol Res* 118: 1559-1572.

12.   Dimier, I. H., and D. T. Bout. 1998. Interferon-gamma-activated primary
enterocytes inhibit Toxoplasma gondii replication: a role for intracellular iron.
Immunology 94: 488-495.

13.   Byrd, T. F., and M. A. Horwitz. 1993. Regulation of transferrin receptor expression
and ferritin content in human mononuclear phagocytes. Coordinate upregulation
by iron transferrin and downregulation by interferon gamma. *The Journal of clinical*
investigation 91: 969-976.

14.   Gail, M., U. Gross, and W. Bohne. 2004. Transferrin receptor induction in
Toxoplasma gondii-infected HFF is associated with increased iron-responsive
protein 1 activity and is mediated by secreted factors. *Parasitol Res* 94: 233-239.

15.   Oliveira, M. C., L. B. Coutinho, M. P. O. Almeida, M. P. Briceno, E. C. B. Araujo,
and N. M. Silva. 2020. The Availability of Iron Is Involved in the Murine
Experimental Toxoplasma gondii Infection Outcome. *Microorganisms* 8.

16.   Plattner, F., F. Yarovinsky, S. Romero, D. Didry, M. F. Carlier, A. Sher, and D.
Soldati-Favre. 2008. Toxoplasma profilin is essential for host cell invasion and
TLR11-dependent induction of an interleukin-12 response. *Cell host & microbe* 3:
77-87.

17.   Yarovinsky, F., S. Hieny, and A. Sher. 2008. Recognition of Toxoplasma gondii by
TLR11 prevents parasite-induced immunopathology. *Journal of immunology*
(Baltimore, Md. : 1950) 181: 8478-8484.

18.   Gazzinelli, R. T., M. Wysocka, S. Hayashi, E. Y. Denkers, S. Hieny, P. Caspar, G.
Trinchieri, and A. Sher. 1994. Parasite-induced IL-12 stimulates early IFN-gamma

synthesis and resistance during acute infection with Toxoplasma gondii. *Journal*

of immunology (Baltimore, Md. : 1950) 153: 2533-2543.

19.   Chen, M., F. Aosai, K. Norose, H. S. Mun, O. Takeuchi, S. Akira, and A. Yano.

2002. Involvement of MyD88 in host defense and the down-regulation of anti-heat

shock protein 70 autoantibody formation by MyD88 in Toxoplasma gondii-infected

mice. *The Journal of parasitology* 88: 1017-1019.

20.   Hitziger, N., I. Dellacasa, B. Albiger, and A. Barragan. 2005. Dissemination of

Toxoplasma gondii to immunoprivileged organs and role of Toll/interleukin-1

receptor signalling for host resistance assessed by in vivo bioluminescence

imaging. *Cell Microbiol* 7: 837-848.

21.   Hunter, C. A., C. S. Subauste, V. H. Van Cleave, and J. S. Remington. 1994.

Production of gamma interferon by natural killer cells from Toxoplasma gondii-

infected SCID mice: regulation by interleukin-10, interleukin-12, and tumor

necrosis factor alpha. *Infection and immunity* 62: 2818-2824.

22.   Gazzinelli, R. T., S. Hieny, T. A. Wynn, S. Wolf, and A. Sher. 1993. Interleukin 12

is required for the T-lymphocyte-independent induction of interferon gamma by an

intracellular parasite and induces resistance in T-cell-deficient hosts. *Proceedings*

of the National Academy of Sciences of the United States of America 90: 6115-

6119.

23.   López-Yglesias, A. H., E. Burger, E. Camanzo, A. T. Martin, A. M. Araujo, S. F.

Kwok, and F. Yarovinsky. 2021. T-bet-dependent ILC1- and NK cell-derived IFN-

γ mediates cDC1-dependent host resistance against Toxoplasma gondii. *PLoS*

pathogens 17: e1008299.

24.   Gigley, J. P., B. A. Fox, and D. J. Bzik. 2009. Cell-mediated immunity to
806   Toxoplasma gondii develops primarily by local Th1 host immune responses in the
807   absence of parasite replication. *Journal of immunology (Baltimore, Md. : 1950)*
808   182: 1069-1078.

809   25.   Gigley, J. P., B. A. Fox, and D. J. Bzik. 2009. Long-term immunity to lethal acute
810   or chronic type II Toxoplasma gondii infection is effectively induced in genetically
811   susceptible C57BL/6 mice by immunization with an attenuated type I vaccine
812   strain. *Infection and immunity* 77: 5380-5388.

813   26.   Casciotti, L., K. H. Ely, M. E. Williams, and I. A. Khan. 2002. CD8(+)-T-cell
814   immunity against Toxoplasma gondii can be induced but not maintained in mice
815   lacking conventional CD4(+) T cells. *Infection and immunity* 70: 434-443.

816   27.   Bhadra, R., J. P. Gigley, L. M. Weiss, and I. A. Khan. 2011. Control of Toxoplasma
817   reactivation by rescue of dysfunctional CD8+ T-cell response via PD-1-PDL-1
818   blockade. *Proceedings of the National Academy of Sciences of the United States
819   of America* 108: 9196-9201.

820   28.   Ivanova, D. L., R. Krempels, S. L. Denton, K. D. Fettel, G. M. Saltz, D. Rach, R.
821   Fatima, T. Mundhenke, J. Materi, I. R. Dunay, and J. P. Gigley. 2020. NK Cells
822   Negatively Regulate CD8 T Cells to Promote Immune Exhaustion and Chronic
823   Toxoplasma gondii Infection. *Frontiers in cellular and infection microbiology* 10:
824   313.

825   29.   Xiao, J., Y. Li, R. H. Yolken, and R. P. Viscidi. 2018. PD-1 immune checkpoint
826   blockade promotes brain leukocyte infiltration and diminishes cyst burden in a
827   mouse model of Toxoplasma infection. *Journal of neuroimmunology* 319: 55-62.

828   30.   Hwang, S., D. A. Cobb, R. Bhadra, B. Youngblood, and I. A. Khan. 2016. Blimp-1-
829   mediated CD4 T cell exhaustion causes CD8 T cell dysfunction during chronic
830   toxoplasmosis. *The Journal of experimental medicine* 213: 1799-1818.

831   31.   Kono, M., S. Matsuhiroya, A. Obuchi, T. Takahashi, S. Imoto, S. Kawano, and K.
832   Saigo. 2020. Deferasirox, an iron-chelating agent, alleviates acute lung
833   inflammation by inhibiting neutrophil activation and extracellular trap formation. *J
834   Int Med Res* 48: 300060520951015.

835   32.   Agoro, R., M. Taleb, V. F. J. Quesniaux, and C. Mura. 2018. Cell iron status
836   influences macrophage polarization. *PloS one* 13: e0196921.

837   33.   Cui, Y., S. Gutierrez, S. Ariai, L. Öberg, K. Thörn, U. Gehrmann, S. M. Cloonan, T.
838   Naessens, and H. Olsson. 2022. Non-heme iron overload impairs monocyte to
839   macrophage differentiation via mitochondrial oxidative stress. *Frontiers in
840   Immunology* 13.

841   34.   Fell, L. H., S. Seiler-Mussler, A. B. Sellier, B. Rotter, P. Winter, M. Sester, D. Fliser,
842   G. H. Heine, and A. M. Zawada. 2016. Impact of individual intravenous iron
843   preparations on the differentiation of monocytes towards macrophages and
844   dendritic cells. *Nephrol Dial Transplant* 31: 1835-1845.

845   35.   Flynn, M. G., L. Mackinnon, V. Gedge, M. Fahlman, and T. Brickman. 2003.
846   Influence of iron status and iron supplements on natural killer cell activity in trained
847   women runners. *Int J Sports Med* 24: 217-222.

848   36.   Baines, M. G., F. L. Lafleur, and B. E. Holbein. 1983. Involvement of transferrin
849   and transferrin receptors in human natural killer effector:target interaction.
850   *Immunol Lett* 7: 51-55.

37.   Yarosz, E. L., C. Ye, A. Kumar, C. Black, E. K. Choi, Y. A. Seo, and C. H. Chang.

2020. Cutting Edge: Activation-Induced Iron Flux Controls CD4 T Cell Proliferation

by Promoting Proper IL-2R Signaling and Mitochondrial Function. *Journal of*

immunology (Baltimore, Md. : 1950) 204: 1708-1713.

38.   Berg, V., M. Modak, J. Brell, A. Puck, S. Kunig, S. Jutz, P. Steinberger, G. J.

Zlabinger, and J. Stockl. 2020. Iron Deprivation in Human T Cells Induces

Nonproliferating Accessory Helper Cells. *Immunohorizons* 4: 165-177.

39.   Pfeifhofer-Obermair, C., P. Tymoszuk, M. Nairz, A. Schroll, G. Klais, E. Demetz,

S. Engl, N. Brigo, and G. Weiss. 2021. Regulation of Th1 T Cell Differentiation by

Iron via Upregulation of T Cell Immunoglobulin and Mucin Containing Protein-3

(TIM-3). *Front Immunol* 12: 637809.

40.   Leung, S., A. Holbrook, B. King, H. T. Lu, V. Evans, N. Miyamoto, C. Mallari, S.

Harvey, D. Davey, E. Ho, W. W. Li, J. Parkinson, R. Horuk, S. Jaroch, M. Berger,

W. Skuballa, C. West, R. Pulk, G. Phillips, J. Bryant, B. Subramanyam, C.

Schaefer, H. Salamon, E. Lyons, D. Schilling, H. Seidel, J. Kraetzschmar, M.

Snider, and D. Perez. 2005. Differential inhibition of inducible T cell cytokine

secretion by potent iron chelators. *J Biomol Screen* 10: 157-167.

41.   Rodriguez, R., C. L. Jung, V. Gabayan, J. C. Deng, T. Ganz, E. Nemeth, and Y.

Bulut. 2014. Hepcidin induction by pathogens and pathogen-derived molecules is

strongly dependent on interleukin-6. *Infection and immunity* 82: 745-752.

42.   Chen, A. C., A. Donovan, R. Ned-Sykes, and N. C. Andrews. 2015. Noncanonical

role of transferrin receptor 1 is essential for intestinal homeostasis. *Proceedings of*

the *National Academy of Sciences of the United States of America* 112: 11714-11719.

43. Hatcher, H. C., R. N. Singh, F. M. Torti, and S. V. Torti. 2009. Synthetic and natural iron chelators: therapeutic potential and clinical use. *Future Med Chem* 1: 1643-1670.

44. Adams, L. B., J. B. Hibbs, Jr., R. R. Taintor, and J. L. Krahenbuhl. 1990. Microbiostatic effect of murine-activated macrophages for Toxoplasma gondii. Role for synthesis of inorganic nitrogen oxides from L-arginine. *Journal of immunology (Baltimore, Md. : 1950)* 144: 2725-2729.

45. Liu, C.-H., Y.-t. Fan, A. Dias, L. Esper, R. A. Corn, A. Bafica, F. S. Machado, and J. Aliberti. 2006. Cutting Edge: Dendritic Cells Are Essential for In Vivo IL-12 Production and Development of Resistance against Toxoplasma gondii Infection in Mice. *The Journal of Immunology* 177: 31-35.

46. Mercer, H. L., L. M. Snyder, C. M. Doherty, B. A. Fox, D. J. Bzik, and E. Y. Denkers. 2020. Toxoplasma gondii dense granule protein GRA24 drives MyD88-independent p38 MAPK activation, IL-12 production and induction of protective immunity. *PLoS pathogens* 16: e1008572.

47. Mombaerts, P., J. Iacomini, R. S. Johnson, K. Herrup, S. Tonegawa, and V. E. Papaioannou. 1992. RAG-1-deficient mice have no mature B and T lymphocytes. *Cell* 68: 869-877.

48. Hunter, C. A., R. Chizzonite, and J. S. Remington. 1995. Il-1-Beta Is Required for Il-12 to Induce Production of Ifn-Gamma by Nk-Cells - a Role for Il-1-Beta in the T-

Cell-Independent Mechanism of Resistance against Intracellular Pathogens.

Journal of Immunology 155: 4347-4354.

49.   Goldszmid, R. S., P. Caspar, A. Rivollier, S. White, A. Dzutsev, S. Hieny, B. Kelsall,

G. Trinchieri, and A. Sher. 2012. NK cell-derived interferon-gamma orchestrates

cellular dynamics and the differentiation of monocytes into dendritic cells at the site

of infection. *Immunity* 36: 1047-1059.

50.   Lee, Y. H., J. Y. Channon, T. Matsuura, J. D. Schwartzman, D. W. Shin, and L. H.

Kasper. 1999. Functional and quantitative analysis of splenic T cell immune

responses following oral toxoplasma gondii infection in mice. *Experimental*

parasitology 91: 212-221.

51.   Carrof, B., M. Levacher-Clergeot, F. Chau, J. J. Pocidalo, and F. Derouin. 1994.

Toxoplasma gondii: kinetics of lymphocyte subsets in blood and spleen of perorally

infected mice. *Experimental parasitology* 78: 410-417.

52.   McDermott, D. S., and S. M. Varga. 2011. Quantifying antigen-specific CD4 T cells

during a viral infection: CD4 T cell responses are larger than we think. *Journal of*

immunology (Baltimore, Md. : 1950) 187: 5568-5576.

53.   Haque, S., J. Franck, H. Dumon, L. H. Kasper, and A. Haque. 1999. Protection

against lethal toxoplasmosis in mice by an avirulent strain of Toxoplasma gondii:

stimulation of IFN-gamma and TNF-alpha response. *Experimental parasitology* 93:

231-240.

54.   Gazzinelli, R. T., I. Eltoum, T. A. Wynn, and A. Sher. 1993. Acute cerebral

toxoplasmosis is induced by in vivo neutralization of TNF-alpha and correlates with

the down-regulated expression of inducible nitric oxide synthase and other

918 markers of macrophage activation. *Journal of immunology (Baltimore, Md. : 1950)*

919 151: 3672-3681.

920 55. Shirahata, T., T. Yamashita, C. Ohta, H. Goto, and A. Nakane. 1994. CD8+ T

921 lymphocytes are the major cell population involved in the early gamma interferon

922 response and resistance to acute primary Toxoplasma gondii infection in mice.

923 *Microbiol Immunol* 38: 789-796.

924 56. Jongert, E., A. Lemiere, J. Van Ginderachter, S. De Craeye, K. Huygen, and S.

925 D'Souza. 2010. Functional characterization of in vivo effector CD4(+) and CD8(+)

926 T cell responses in acute Toxoplasmosis: an interplay of IFN-gamma and cytolytic

927 T cells. *Vaccine* 28: 2556-2564.

928 57. Denkers, E. Y., and R. T. Gazzinelli. 1998. Regulation and function of T-cell-

929 mediated immunity during Toxoplasma gondii infection. *Clinical microbiology*

930 *reviews* 11: 569-588.

931 58. Giannetti, A. M., P. J. Halbrooks, A. B. Mason, T. M. Vogt, C. A. Enns, and P. J.

932 Bjorkman. 2005. The molecular mechanism for receptor-stimulated iron release

933 from the plasma iron transport protein transferrin. *Structure* 13: 1613-1623.

934 59. Bayer, A. L., P. Baliga, and J. E. Woodward. 1998. Transferrin receptor in T cell

935 activation and transplantation. *Journal of leukocyte biology* 64: 19-24.

936 60. Brekelmans, P., P. van Soest, P. J. Leenen, and W. van Ewijk. 1994. Inhibition of

937 proliferation and differentiation during early T cell development by anti-transferrin

938 receptor antibody. *European journal of immunology* 24: 2896-2902.

61.   Brekelmans, P., P. van Soest, J. Voerman, P. P. Platenburg, P. J. Leenen, and W.

van Ewijk. 1994. Transferrin receptor expression as a marker of immature cycling

thymocytes in the mouse. *Cellular immunology* 159: 331-339.

62.   Wilson, D. C., S. Matthews, and G. S. Yap. 2008. IL-12 signaling drives CD8+ T

cell IFN-gamma production and differentiation of KLRG1+ effector subpopulations

during Toxoplasma gondii Infection. *Journal of immunology (Baltimore, Md. : 1950)*

180: 5935-5945.

63.   Sanecka, A., N. Yoshida, E. M. Kolawole, H. Patel, B. D. Evavold, and E. M.

Frickel. 2018. T Cell Receptor-Major Histocompatibility Complex Interaction

Strength Defines Trafficking and CD103(+) Memory Status of CD8 T Cells in the

Brain. *Front Immunol* 9: 1290.

64.   Moretto, M. M., J. Chen, M. Meador, J. Phan, and I. A. Khan. 2023. A Lower Dose

of Infection Generates a Better Long-Term Immune Response against

Toxoplasma gondii. *Immunohorizons* 7: 177-190.

65.   Fox, B. A., and D. J. Bzik. 2002. De novo pyrimidine biosynthesis is required for

virulence of Toxoplasma gondii. *Nature* 415: 926-929.

66.   Jordan, K. A., E. H. Wilson, E. D. Tait, B. A. Fox, D. S. Roos, D. J. Bzik, F.

Dzierszinski, and C. A. Hunter. 2009. Kinetics and phenotype of vaccine-induced

CD8+ T-cell responses to Toxoplasma gondii. *Infection and immunity* 77: 3894-

3901.

67.   Gerdes, J., H. Lemke, H. Baisch, H. H. Wacker, U. Schwab, and H. Stein. 1984.

Cell cycle analysis of a cell proliferation-associated human nuclear antigen defined

by the monoclonal antibody Ki-67. *Journal of immunology (Baltimore, Md. : 1950)*

133: 1710-1715.

68.   Hentze, M. W., and L. C. Kuhn. 1996. Molecular control of vertebrate iron

metabolism: mRNA-based regulatory circuits operated by iron, nitric oxide, and

oxidative stress. *Proceedings of the National Academy of Sciences of the United*

States of America 93: 8175-8182.

69.   Van Den Ham, K. M., M. T. Shio, A. Rainone, S. Fournier, C. M. Krawczyk, and M.

Olivier. 2015. Iron prevents the development of experimental cerebral malaria by

attenuating CXCR3-mediated T cell chemotaxis. *PloS one* 10: e0118451.

70.   Regis, G., M. Bosticardo, L. Conti, S. De Angelis, D. Boselli, B. Tomaino, P.

Bernabei, M. Giovarelli, and F. Novelli. 2005. Iron regulates T-lymphocyte

sensitivity to the IFN-gamma/STAT1 signaling pathway in vitro and in vivo. *Blood*

105: 3214-3221.

71.   Rathmell, J. C., M. G. Vander Heiden, M. H. Harris, K. A. Frauwirth, and C. B.

Thompson. 2000. In the absence of extrinsic signals, nutrient utilization by

lymphocytes is insufficient to maintain either cell size or viability. *Mol Cell* 6: 683-

692.

72.   Frauwirth, K. A., J. L. Riley, M. H. Harris, R. V. Parry, J. C. Rathmell, D. R. Plas,

R. L. Elstrom, C. H. June, and C. B. Thompson. 2002. The CD28 signaling pathway

regulates glucose metabolism. *Immunity* 16: 769-777.

73.   Verbist, K. C., C. S. Guy, S. Milasta, S. Liedmann, M. M. Kaminski, R. Wang, and

D. R. Green. 2016. Metabolic maintenance of cell asymmetry following division in

activated T lymphocytes. *Nature* 532: 389-393.

74.   Menk, A. V., N. E. Scharping, R. S. Moreci, X. Zeng, C. Guy, S. Salvatore, H. Bae,

985   J. Xie, H. A. Young, S. G. Wendell, and G. M. Delgoffe. 2018. Early TCR Signaling

986   Induces Rapid Aerobic Glycolysis Enabling Distinct Acute T Cell Effector

987   Functions. *Cell Rep* 22: 1509-1521.

988   75.   Loschinski, R., M. Bottcher, A. Stoll, H. Bruns, A. Mackensen, and D.

989   Mougiakakos. 2018. IL-21 modulates memory and exhaustion phenotype of T-

990   cells in a fatty acid oxidation-dependent manner. *Oncotarget* 9: 13125-13138.

991   76.   Wang, R., Christopher P. Dillon, Lewis Z. Shi, S. Milasta, R. Carter, D. Finkelstein,

992   Laura L. McCormick, P. Fitzgerald, H. Chi, J. Munger, and Douglas R. Green.

993   2011. The Transcription Factor Myc Controls Metabolic Reprogramming upon T

994   Lymphocyte Activation. *Immunity* 35: 871-882.

995   77.   Frauwirth, K. A., J. L. Riley, M. H. Harris, R. V. Parry, J. C. Rathmell, D. R. Plas,

996   R. L. Elstrom, C. H. June, and C. B. Thompson. 2002. The CD28 Signaling

997   Pathway Regulates Glucose Metabolism. *Immunity* 16: 769-777.

998   78.   Gray, N. K., S. Quick, B. Goossen, A. Constable, H. Hirling, L. C. Kühn, and M. W.

999   Hentze. 1993. Recombinant iron-regulatory factor functions as an iron-responsive-

1000   element-binding protein, a translational repressor and an aconitase. A functional

1001   assay for translational repression and direct demonstration of the iron switch. *Eur*

1002   *J Biochem* 218: 657-667.

1003   79.   Silva-Gomes, S., S. Vale-Costa, R. Appelberg, and M. S. Gomes. 2013. Iron in

1004   intracellular infection: to provide or to deprive? *Frontiers in cellular and infection*

1005   *microbiology* 3: 96.

80.   Yarosz, E. L., and C. H. Chang. 2018. The Role of Reactive Oxygen Species in

Regulating T Cell-mediated Immunity and Disease. *Immune Netw* 18: e14.

81.   Sena, L. A., S. Li, A. Jairaman, M. Prakriya, T. Ezponda, D. A. Hildeman, C.-R.

Wang, P. T. Schumacker, J. D. Licht, and H. Perlman. 2013. Mitochondria are

required for antigen-specific T cell activation through reactive oxygen species

signaling. *Immunity* 38: 225-236.

82.   Lepez, A., T. Pirnay, S. Denanglaire, D. Perez-Morga, M. Vermeersch, O. Leo, and

F. Andris. 2020. Long-term T cell fitness and proliferation is driven by AMPK-

dependent regulation of reactive oxygen species. *Sci Rep* 10: 21673.

83.   Kaminski, M. M., S. W. Sauer, C. D. Klemke, D. Süss, J. G. Okun, P. H. Krammer,

and K. Gülow. 2010. Mitochondrial reactive oxygen species control T cell activation

by regulating IL-2 and IL-4 expression: mechanism of ciprofloxacin-mediated

immunosuppression. *Journal of immunology (Baltimore, Md. : 1950)* 184: 4827-

4841.

84.   Arbore, G., E. E. West, R. Spolski, A. A. B. Robertson, A. Klos, C. Rheinheimer,

P. Dutow, T. M. Woodruff, Z. X. Yu, L. A. O'Neill, R. C. Coll, A. Sher, W. J. Leonard,

J. Köhl, P. Monk, M. A. Cooper, M. Arno, B. Afzali, H. J. Lachmann, A. P. Cope,

K. D. Mayer-Barber, and C. Kemper. 2016. T helper 1 immunity requires

complement-driven NLRP3 inflammasome activity in CD4[+] T cells. *Science* 352:

aad1210.

85.   Salmeron, A., A. Borroto, M. Fresno, M. J. Crumpton, S. C. Ley, and B. Alarcon.

1995. Transferrin receptor induces tyrosine phosphorylation in T cells and is

physically associated with the TCR zeta-chain. *Journal of immunology (Baltimore, Md. : 1950)* 154: 1675-1683.

86. Davis, R. J., G. L. Johnson, D. J. Kelleher, J. K. Anderson, J. E. Mole, and M. P. Czech. 1986. Identification of serine 24 as the unique site on the transferrin receptor phosphorylated by protein kinase C. *J Biol Chem* 261: 9034-9041.

87. Cheng, Y., O. Zak, P. Aisen, S. C. Harrison, and T. Walz. 2004. Structure of the human transferrin receptor-transferrin complex. *Cell* 116: 565-576.

88. Rothenberger, S., B. J. Iacopetta, and L. C. Kuhn. 1987. Endocytosis of the transferrin receptor requires the cytoplasmic domain but not its phosphorylation site. *Cell* 49: 423-431.

89. Zerial, M., M. Suomalainen, M. Zanetti-Schneider, C. Schneider, and H. Garoff. 1987. Phosphorylation of the human transferrin receptor by protein kinase C is not required for endocytosis and recycling in mouse 3T3 cells. *The EMBO journal* 6: 2661-2667.

90. Davis, R. J., and H. Meisner. 1987. Regulation of transferrin receptor cycling by protein kinase C is independent of receptor phosphorylation at serine 24 in Swiss 3T3 fibroblasts. *J Biol Chem* 262: 16041-16047.

91. Isakov, N., and A. Altman. 2002. Protein kinase C(theta) in T cell activation. *Annu Rev Immunol* 20: 761-794.

92. Goldszmid, R. S., I. Coppens, A. Lev, P. Caspar, I. Mellman, and A. Sher. 2009. Host ER-parasitophorous vacuole interaction provides a route of entry for antigen cross-presentation in Toxoplasma gondii-infected dendritic cells. *The Journal of experimental medicine* 206: 399-410.

93.   Lee, Y., M. Sasai, J. S. Ma, N. Sakaguchi, J. Ohshima, H. Bando, T. Saitoh, S.

Akira, and M. Yamamoto. 2015. p62 Plays a Specific Role in Interferon-γ-Induced

Presentation of a Toxoplasma Vacuolar Antigen. *Cell Rep* 13: 223-233.

94.   Yano, A., S. Ohno, K. Norose, T. Baba, K. Yamashita, F. Aosai, and K. Segawa.

1992. Antigen presentation by Toxoplasma-infected cells: antigen entry through

cell membrane fusion. *Int Arch Allergy Immunol* 98: 13-17.

95.   Wang, Y., M. Wang, Y. Liu, H. Tao, S. Banerjee, S. Srinivasan, E. Nemeth, M. J.

Czaja, and P. He. 2022. Integrated regulation of stress responses, autophagy and

survival by altered intracellular iron stores. *Redox Biol* 55: 102407.

96.   Romao, S., N. Gasser, A. C. Becker, B. Guhl, M. Bajagic, D. Vanoaica, U. Ziegler,

J. Roesler, J. Dengjel, and J. Reichenbach. 2013. Autophagy proteins stabilize

pathogen-containing phagosomes for prolonged MHC II antigen processing.

Journal of Cell Biology 203: 757-766.

1064

1065

1066

Figure 1. Iron restriction reduces *T. gondii* replication *in vitro* yet increases morbidity and mortality *in vivo*.

A) Separate lots of DFP formulations were tested for Iron-Binding Units as described. Line graph shows titration curves of DFP used to determine 50% quenching of iron. **B)** Representative plaque assays are presented of RH strain *T. gondii* infection of fibroblasts that were non-treated (NT) or treated with 1.37U DFP, 30µg ferric citrate (Fe), or 1.37U DFP and 30µg ferric citrate (DFP + Fe). **C)** Bar graphs present pooled data from two independent replicates of N=6 each for total plaque number (left) and size of plaques (right, minimum 30 plaques measured per condition if possible). **D-H)** C57BL/6 mice infected with 8 cysts ME49 strain *T. gondii* i.g. and treated i.p. with 75U DFP every other day starting the day before infection were monitored or sacrificed at the timepoints indicated. **D)** Survival of infected mice after treatment with 75U DFP. Survival graph presents data pooled from three independent experiments of N=3-5 per group. **E)** Line graph presents percentage of weight loss over time of mice after treatment with 75U DFP. Data pooled from two independent repeats of N=4-5 per group. **F)** Parasite burden at 14DPI or time of death in infected organs after treatment with 75U DFP. Bar graphs

present data from lung and brain pooled from two independent experiments of N=4 each.

G) Brain cysts enumerated from infected mice treated with 75U DFP i.p. every other day

starting on day -1 and surviving until day 35. Bar graph presents data pooled from four

independent experiments but due to death of mice prior to day 35 the total sample size

was 20 for NT and 11 for DFP treated groups. Bar graphs depict mean $\pm$ SD. Significance

is denoted with * where $p \leq 0.05$, ** where $p \leq 0.01$, *** where $p \leq 0.005$, **** where $p \leq$

0.001, and NS where $p > 0.05$.

1090

Figure 2. Iron restriction lowers IL-12 and IFNγ at 5DPI of *T. gondii* infection.

Spleen cells and serum from C57BL/6 mice infected with 8 cysts ME49 I.G. and treated

with 75U DFP were harvested at 5DPI. Bar graphs present results of ELISA

measurements of serum levels and splenocyte production of IL-12p40 and IFNγ pooled

from two independent experiments of N=4 each. All graphs depict mean ± SD.

Significance is denoted with * where p ≤ 0.05, ** where p ≤ 0.01, *** where p ≤ 0.005,

**** where p ≤ 0.001, and NS where p > 0.05.

1098

Figure 3. Iron restriction does not impact spleen cell proportions during early acute

***T. gondii* infection.**

C57BL/6 mice were infected orally with 8 cysts ME49 and treated with 75U DFP i.p. on

days -1, 1, and 3. Splenocytes were isolated on day 5 for flow cytometric analysis. **A)**

Gating strategy presents Boolean gates (denoted G and in blue boxes) and population

gates (denoted P and in red boxes). Lineage negative populations were gated as

Live/Dead-, CD19-, CD3-, and NK1.1-. Population gates included P1 (Total Live Cells),

P2 (Live, CD19+), P3 (Live, CD19+, CD3+, NK1.1-), P4 (Live, CD19-, CD3-, NK1.1+), P5

(Lineage-, CD11b+, Ly6CHi, Ly6G-, confirmed by side scatter profile), P6 (Lineage-,

CD11b+ Ly6Cint, Ly6G+, confirmed by side scatter profile), P7 (Lineage-, CD11c+, I-A/I-

E+). **B)** Pie charts depict the frequency means of indicated cell populations derived from

raw data (Supplemental Figure 2) pooled from three independent experiments of N=4 per

1111   group. **C)** Survival of RAG1-/- mice infected orally with 8 cysts ME49 and treated with

1112   75U DFP i.p. every other day starting the day before infection. Survival data is pooled

1113   from two independent experiments where NS indicates $p > 0.05$ by Kaplan-Meier

1114   analysis.

1115

Figure 4. NK cell number and function are not affected by iron restriction during *T. gondii* infection.

C57BL/6 mice were orally infected with 8 cysts ME49 and treated with 75U DFP i.p. on days -1, 1, 3. Splenocytes were stimulated with α-NK1.1 according to the MIANKA. **A)** Bar graphs and contour plots present frequency and absolute numbers of splenic cNK cells (Live, CD3-, CD49b+, NKp46+). **B)** Bar graphs and contour plots present frequency and absolute numbers of IFNγ-producing cNK cells (Live, CD3-, CD49b+, NKp46+, IFNγ+). Data pooled from two independent experiments of N=4 per group. All graphs depict mean ± SD. Significance is denoted with * where p ≤ 0.05, ** where p ≤ 0.01, *** where p ≤ 0.005, **** where p ≤ 0.001, and NS where p > 0.05.

1126

Figure 5. Iron is required for CD4+ and CD8+ T cell activation, expansion, function,

and effector cell differentiation during Type II strain ME49 infection.

C57BL/6 mice were orally infected with 8 cysts ME49 strain *T. gondii* and treated with

75U DFP i.p. every other day starting the day before infection. Splenocytes were prepared

for stimulation with TLA or for direct *ex vivo* flow cytometry.. **A)** Direct *ex vivo* staining of

CD4 and CD8 is represented by contour plots and bar graphs presenting frequency and

absolute numbers of CD4+ T cells (Live/CD4+/CD8-) or CD8+ T Cells (Live/CD4-/CD8+).

B) Direct *ex vivo* staining of CD11a and CD49d is represented by contour plots and bar

graphs presenting frequency and absolute numbers of antigen experienced CD4+ T cells (Live/CD4+/CD8-/CD11aHi/CD49d+). **C)** Direct *ex vivo* staining of CD11a and CD49d is represented by contour plots and bar graphs presenting frequency and absolute numbers of antigen experienced CD8+ T cells (Live/CD4-/CD8+/CD11aHi/CD49d+). **D)** TLA-stimulated staining of IFNγ and TNFα is represented by contour plots and bar graphs of frequency and absolute numbers of polyfunctional CD4+ T Cells (Live/CD4+/CD8-/IFNγ+/TNFα+). **E)** TLA-stimulated staining of IFNγ is represented by contour plots and bar graphs of frequency and absolute numbers of functional CD8+ T Cells (Live/CD4-/CD8+/IFNγ+). **F-I)** Direct *ex vivo* staining of CD71 is represented by histograms and bar graphs presenting the geometric mean of CD71 intensity of F) Total CD4+ T cells (Live/CD4+/CD8-), G) Total CD8+ T cells (Live/CD4-/CD8+), H) Antigen experienced CD4+ T cells (Live/CD4+/CD8-/CD11aHi/CD49d+), and I) Antigen experienced CD8+ T cells (Live/CD4-/CD8+/CD11aHi/CD49d+). **J)** Direct *ex vivo* staining of KLRG1 and CD62L on CD8+ T cells is represented by contour plots and bar graphs presenting the frequency of FI: Naïve (CD62L+/KLRG1-), FII: Memory Precursor (CD62L-/KLRG1-), FIII: Short lived Effector (CD62L-/KLRG1+), and FIV: CD62L+ Effector (CD62L+/KLRG1+) subpopulations within the Live/CD4-/CD8+ population. All data pooled from 2-5 independent experiments of N=3-5 per group. All graphs depict mean $\pm$ SD. Significance is denoted with * where $p \leq 0.05$, ** where $p \leq 0.01$, *** where $p \leq 0.005$, **** where $p \leq 0.001$, and NS where $p > 0.05$.

1155

Figure 6. Inhibition of CD4+ and CD8+ T cell activation, expansion, function, and effector cell differentiation by iron restriction during vaccination is independent of antigen load.

A-F) C57BL/6 mice were infected with 1 X 10^6 tachyzoites CPS i.p. and treated with 75U DFP as described. At 7DPI, peritoneal Exudate Cells (PEC) were prepared for simulation or direct *ex vivo* staining. **A)** Direct *ex vivo* staining of CD4 and CD8 is represented by contour plots and bar graphs presenting frequency and absolute numbers of CD4+ T cells (Live/CD4+/CD8-) or CD8+ T Cells (Live/CD4-/CD8+). **B)** Direct *ex vivo* staining of CD11a

and CD49d is represented by contour plots and bar graphs presenting frequency and

absolute numbers of antigen experienced CD4+ T cells (Live/CD4+/CD8-

/CD11aHi/CD49d+). **C)** Direct ex vivo staining of CD11a and CD49d is represented by

contour plots and bar graphs presenting frequency and absolute numbers of antigen

experienced CD8+ T cells (Live/CD4-/CD8+/CD11aHi/CD49d). **D)** TLA-stimulated

staining of IFNγ and TNFα is represented by contour plots and bar graphs presenting

frequency and absolute numbers of polyfunctional CD4+ T Cells (Live/CD4+/CD8-

/IFNγ+/TNFα+). **E)** TLA-stimulated staining of IFNγ is represented by contour plots and

graphs presenting frequency and absolute numbers of functional CD8+ T Cells

(Live/CD4-/CD8+/IFNγ+). **F)** Direct *ex vivo* staining of KLRG1 and CD62L on CD8+ T cells

is represented by contour plots and graphs presenting the frequency of FI: Naïve

(CD62L+/KLRG1-), FII: Memory Precursor (CD62L-/KLRG1-), FIII: Short lived Effector

(CD62L-/KLRG1+), and FIV: CD62L+ Effector (CD62L+/KLRG1+) subpopulations within

the Live/CD4-/CD8+ population. **G)** C57BL/6 mice were injected with 1 X 10^4 (left), 5 X

10^4 (middle), or 1 X 10^5 (right) tachyzoites of CPS strain *T. gondii* while being treated or

not with 75U DFP for the first three weeks of vaccination. DFP treatment was then

stopped for the last 2 weeks of vaccination allowing iron levels to recover. Graphs present

survival of mice lethally challenged with 1 X 10^3 RH strain *T. gondii* two weeks after

cessation of DFP treatment. All flow cytometry data are pooled from three independent

experiments of N=3-4 per group. All bar graphs depict mean $\pm$ SD. Vaccination studies

were conducted twice. Significance is denoted with * where $p \leq 0.05$, ** where $p \leq 0.01$,

*** where $p \leq 0.005$, **** where $p \leq 0.001$, and NS where $p > 0.05$.

1186

Figure 7. CD71 is required for activation and expansion of CD4+ T cells in a cell

intrinsic manner during *T. gondii* infection.

CD4cCD71KO mice and littermate controls were genotyped for possession (Cre+) or not

(Cre-) of CreER[T2] recombinase under the CD4 promotor, and all possessed flanking lox

sites of the endogenous CD71 gene. Cre+ and Cre- littermates were alternatively treated

(Tam+) or not (Tam-) with 100mg/kg tamoxifen i.p. daily for five days prior to oral infection

with 8 cysts ME49. Splenocytes were harvested 10DPI for direct *ex vivo* staining

according to the MIATA. **A-D)** Direct *ex vivo* staining of CD4 and CD8 is represented by

contour plots and bar graphs presenting frequency and absolute numbers of B) non-T

Cells (Live/CD4-/CD8-), C) CD4+ T cells (Live/CD4+/CD8-), and D) CD8+ T cells

(Live/CD4-/CD8+). **E-F)** Direct *ex vivo* staining of CD71 is represented by histograms and

bar graphs presenting geometric mean of CD71 intensity in E) CD4+ T Cells

(Live/CD4+/CD8-) and F) CD8+ T cells (Live/CD4-/CD8+). **G-H)** Direct *ex vivo* staining of

Ki67 is represented by histograms and bar graphs presenting geometric mean of Ki67

intensity in G) CD4+ T Cells (Live/CD4+/CD8-) and H) CD8+ T cells (Live/CD4-/CD8+).

All data pooled from two independent experiments of N=3-4 per group. Bar graphs depict

mean $\pm$ SD. Significance is denoted with * where $p \leq 0.05$, ** where $p \leq 0.01$, *** where

$p \leq 0.005$, **** where $p \leq 0.001$, and NS where $p > 0.05$.

1205

Figure 8. CD71 is required to increase antigen experience and polyfunctional

responses of CD4+ T cells in *T. gondii* infection

CD4cCD71KO mice and littermate controls were genotyped for possession (Cre+) or not

(Cre-) of CreER[T2] recombinase under the CD4 promotor, and all possessed flanking lox

sites of the endogenous CD71 gene. Cre+ and Cre- littermates were alternatively treated

(Tam+) or not (Tam-) with 100mg/kg tamoxifen i.p. daily for five days prior to oral infection

with 8 cysts ME49. Splenocytes were harvested 10DPI for stimulation or direct *ex vivo*

staining according to the MIATA. **A)** Direct *ex vivo* staining of CD11a and CD49d is

represented by contour plots and bar graphs presenting frequency and absolute numbers

of antigen experienced CD4+ T cells (Live/CD4+/CD8-/CD11aHi/CD49d+). **B)** Direct *ex*

vivo staining of CD11a and CD49d is represented by contour plots and bar graphs

presenting frequency and absolute numbers of antigen experienced CD8+ T cells

(Live/CD4-/CD8+/CD11aHi/CD49d). **C)** TLA-stimulated staining of IFNγ and TNFα is

represented by contour plots and bar graphs presenting frequency and absolute numbers

of polyfunctional CD4+ T Cells (Live/CD4+/CD8-/IFNγ+/TNFα+). **D)** TLA-stimulated

staining of IFNγ is represented by contour plots and bar graphs presenting frequency and

absolute numbers of polyfunctional CD8+ T Cells (Live/CD4-/CD8+/IFNγ+/GZB+). **E)**

Survival of ME49 infected CD4cCD71KO mice and littermate controls treated with

tamoxifen. Survival data pooled from two independent experiments of N=4-5 per group

and analyzed by Kaplan-Meier statistics where NS indicates $p > 0.05$. All bar graphs

pooled from two independent experiments of N=3-4 per group and depict mean $\pm$ SD.

Bar graph significance is indicated with * where $p \leq 0.05$, ** where $p \leq 0.01$, *** where p

≤ 0.005, **** where $p \leq 0.001$, and NS where $p > 0.05$.

1229

Supplemental Figure 1. Titration of deferiprone and ferric citrate in plaque assays.

The growth rate of RH strain *T. gondii* treated with indicated amounts of DFP or ferric

citrate was determined by plaque assay. **A)** Shown are representative plaques formed by

RH at indicated levels of DFP *(top),* ferric citrate *(middle)*, and ferric citrate + 1.37U DFP

(bottom). **B)** Bar graph presents the number of RH plaques formed when treated with

indicated doses of DFP. **C)** Bar graph presents the size of RH plaques formed when

treated with indicated doses of DFP. **D)** Bar graph shows the number of plaques formed

after supplementation of 1.37U DFP with indicated amounts of ferric citrate. **E)** Bar graph

presents the number of plaques formed when supplemented with indicated amounts of

ferric citrate alone. Data representative of two independent experiments. Bar graphs show

1240   mean $\pm$ SD. Significance is denoted with * where $p \leq 0.05$, ** where $p \leq 0.01$, *** where

1241 $p \leq 0.005$, **** where $p \leq 0.001$, and NS where $p > 0.05$.

Supplemental Figure 2. Iron restriction does not impact splenic proportions at 5DPI

of *T. gondii* infection.

C57BL/6 mice were infected orally with 8 cysts ME49 and treated with 75U DFP i.p. on days -1, 1, and 3. Splenocytes were isolated on day 5 for flow cytometric analysis. Bar graphs present frequency and absolute numbers of the following populations gated according to Figure 3A: **A)** P1 (Live Cells), **B)** P2 (Live, CD19+), **C)** P3 (Live, CD19+, CD3+, NK1.1-), **D)** P4 (Live, CD19-, CD3-, NK1.1+), **E)** P5 (Lineage-, CD11b+, Ly6CHi, Ly6G-, confirmed by side scatter profile), **F)** P6 (Lineage-, CD11b+ Ly6Cint, Ly6G+, confirmed by side scatter profile), **G)** P7 (Lineage-, CD11c+, I-A/I-E+), and **H)** P8 (Lineage- CD11b- CD11c-). Data pooled from three independent experiments. Bar graphs depict mean $\pm$ SD. Significance is denoted with * where $p \leq 0.05$, ** where $p \leq 0.01$, *** where $p \leq 0.005$, **** where $p \leq 0.001$, and NS where $p > 0.05$.

1255

Supplemental Figure 3. Inhibition of splenic T cell antigen experience and

functionality by iron restriction during vaccination is independent of antigen load.

C57BL/6 mice infected with 1×10^6 tachyzoites CPS i.p. and treated with 75U DFP as

described in the Materials and Methods. Animals were sacrificed and splenocytes were

prepared for stimulation and for direct *ex vivo* staining. **A)** Direct *ex vivo* staining of CD4

and CD8 is represented by contour plots and bar graphs presenting frequency and

absolute numbers of CD4+ T cells (Live/CD4+/CD8-) or CD8+ T Cells (Live/CD4-/CD8+).

B) Direct *ex vivo* staining of CD11a and CD49d is represented by contour plots and bar

graphs presenting frequency and absolute numbers of antigen experienced CD4+ T cells

(Live/CD4+/CD8-/CD11aHi/CD49d+). **C)** Direct *ex vivo* staining of CD11a and CD49d is

represented by contour plots and bar graphs presenting frequency and absolute numbers

of antigen experienced CD8+ T cells (Live/CD4-/CD8+/CD11aHi/CD49d+). **D)** TLA-

stimulated staining of IFNγ and TNFα is represented by contour plots and bar graphs of

frequency and absolute numbers of polyfunctional CD4+ T Cells (Live/CD4+/CD8-

/IFNγ+/TNFα+). **E)** TLA-stimulated staining of IFNγ is represented by contour plots and

bar graphs of frequency and absolute numbers of functional CD8+ T Cells (Live/CD4-

/CD8+/IFNγ+). All data pooled from three independent experiments of N=4 per group. Bar

graphs depict mean $\pm$ SD. Significance is denoted with * where p $\leq$ 0.05, ** where p $\leq$

0.01, *** where p $\leq$ 0.005, **** where p $\leq$ 0.001, and NS where p $>$ 0.05.

Supplemental Figure 4. Conditional knockout of iron uptake receptor CD71 in

CD4+ T cells

CD4cCD71KO mice and littermate controls were genotyped for possession (Cre+) or

not (Cre-) of CreER^{T2} recombinase under the CD4 promotor, and all possessed flanking

lox sites of the endogenous CD71 gene. Cre+ and Cre- littermates were alternatively

treated (Tam+) or not (Tam-) with 100mg/kg tamoxifen i.p. daily for five days prior to

oral infection with 8 cysts ME49. Splenocytes were harvested 10DPI for FACS and RT-

PCR according to the MIATA. **A)** FACS gating strategy used for sorting of non CD4+ T

cells (P1: Live, CD3+/-, CD4-) and CD4+ T cells (P2: Live, CD3+, CD4+). **B)**

Pseudocolored dotplots represent FACS sorting populations for each group. **C)** Bar

graphs present relative mRNA expression of CD71 compared to actin and normalized to

the Cre- Tam- littermate control. The left graph shows CD71 mRNA expression of P1

sorted cells and the right graph shows CD71 mRNA expression of P2 sorted cells. All

data was pooled from two independent experiments of N=1-4 per group and graphs are

mean $\pm$ SD. No comparisons were statistically significant.