Title:

Memory-specific encoding activities of the ventral tegmental area dopamine and
GABA neurons

Authors: Vasileios Glykos[1,2,*], Shigeyoshi Fujisawa[1,*]

Affiliations: [1]Laboratory for Systems Neurophysiology, RIKEN Center for Brain

Science, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan

[2]Present address: Synapse Biology Unit, Okinawa Institute of Science and Technology,
1919-1 Tancha, Onna-son, Kunigami-gun, Okinawa, Japan 904-0495

*Corresponding author: shigeyoshi.fujisawa@riken.jp, vasileios.glykos@oist.jp

Abstract

Although the midbrain dopamine (DA) system plays a crucial role in higher cognitive

functions, including updating and maintaining short-term memory, the encoding

properties of the somatic spiking activity of ventral tegmental area (VTA) DA neurons

for short-term memory computations have not yet been identified. Here, we probed and

analyzed the activity of optogenetically identified DA and GABA neurons while mice

engaged in short-term memory-dependent behavior in a T-maze task. Single-neuron

analysis revealed that significant subpopulations of DA and GABA neurons responded

differently between left and right trials in the memory delay. With a series of control

behavioral tasks and regression analysis tools, we show that firing rate differences are

linked to short-term memory-dependent decisions and cannot be explained by reward-

related processes, motivated behavior, or motor-related activities. This evidence

provides novel insights into the mnemonic encoding activities of midbrain DA and

GABA neurons.

KEYWORDS: Short-term memory, ventral tegmental area, dopamine, GABA

Introduction

Dopamine (DA) neurons originating in the ventral tegmental area (VTA) project to diverse forebrain regions, forming distinct but interacting neuromodulatory systems that are thought to play pivotal roles in the regulation of reward-related learning, motivation, and cognition (Sawaguchi and Goldman-Rakic, 1991; Schultz et al., 1993; Goldman-Rakic, 1995; Schultz et al., 1997; Tzschentke, 2001; Schultz, 2002; Pierce and Kumaresan, 2006; Berridge, 2007; Vijayraghavan et al., 2007; Lammel et al., 2008; Robbins and Arnsten, 2009; Hauber, 2010; Cohen et al., 2012; Salamone and Correa, 2012; Howe et al., 2013; Matsumoto and Takada, 2013; Hamid et al., 2016; Mohebi et al., 2019). A wealth of electrophysiological recordings from midbrain DA neurons, complemented by *in vivo* microdialysis data indicate that midbrain DA activity promotes behaviors associated with motivation (Wise, 2004; Berridge, 2007; Salamone and Correa, 2012; Howe et al., 2013; Matsumoto and Takada, 2013; Hamid et al., 2016; Mohebi et al., 2019) and supports reward-based learning by encoding reward prediction error (RPE) signals (Schultz et al., 1993; Schultz et al., 1997; Cohen et al., 2012).

Also, DA is of central importance to higher cognitive functions, such as updating and maintaining short-term memory (Sawaguchi and Goldman-Rakic, 1991; Miller and Cohen, 2001; Ott and Nieder, 2019). Pioneering behavioral studies which pharmacologically manipulated the activity of DA receptors in the PFC revealed the significant role of DA signals on short-term memory. In fact, an inverted-U-shape effect was discovered, where too little or too much DA receptor stimulation impairs PFC-engaging short-term memory (Sawaguchi and Goldman-Rakic, 1991; Vijayraghavan et al., 2007; Robbins and Arnsten, 2009). Moreover, at the origin of the DA system, electrophysiological recordings at the VTA showed that DA neurons are not active in the delay period of memory tasks (Ljungberg et al., 1991; Schultz, 2002; Phillips et al., 2004; Matsumoto and Takada, 2013; Choi et al., 2020).

Motivated by the response of DA neurons to reward-related stimuli and memory delays, several lines of computational modeling studies sought to answer when and how DA signals support short-term memory "update" and "maintenance". They proposed the "gating theory", which provided a unified computational framework for reward

prediction and short-term memory (Cohen et al., 2002; Dreher and Burnod, 2002;
Montague et al., 2004; Ott and Nieder, 2019). According to the model, reward-
predicting cues, elicit phasic DA release which opens the gate for the afferent signals to
be stored in memory (update). But, in the delay period, low, tonic DA levels close the
gate for interfering signals to enter the PFC and overwrite the short-term memory
component (maintenance). Although the "gating theory" fits adequately the behavior-
unique responses of DA neurons to the coding schemes of short-term memory, it relies
mainly upon empirical evidence of the population activity of DA neurons and the
longstanding consensus that short-term memory depends on the unbroken chain of
persistent neuronal activity (Durstewitz et al., 2000; Curtis and D'Esposito, 2003).

However, recent advances in the study of the brain's functional organization suggest
that persistent neuronal activity might not be the only candidate mechanism for the
active maintenance of goal representation over short delays, leading to the proposal of
new coding schemes for short-term memory (Stokes, 2015; Miller et al., 2018). One of
these candidate mechanisms regards the memory-dependent dynamic changes in
functional connectivity. Neural oscillations are abundant in the mammalian brain and
are thought to offer the networking framework for the temporal organization of neuronal
activity and information processing in short-term memory (Uhlhaas and Singer, 2006;
Buschman et al., 2012; Miller et al., 2018). Calculating the phase coherence of neural
oscillations between distributed brain regions provides an estimation of the functional
connectivity between them (Fries, 2005). Among other basal ganglia regions, the VTA
engages dynamically in the large-scale network of brain systems that support memory-
related information processing. Simultaneous electrophysiological recordings were
performed in the PFC and the VTA while rodents executed memory-guided behavioral
choices in a T-maze task (Fujisawa and Buzsáki, 2011). Neural oscillations (4Hz) were
prominent in both regions throughout the task, but their power and coherence were
adaptively increased in memory delay. In the same task, another short-term memory-
related coding scheme was reported, this time at the single neuronal level. It was shown
that while rodents navigate the maze, performing memory-guided decisions, PFC and
parietal neurons differentiate their firing activities between opposite behavioral choices
(Fujisawa et al., 2008; Harvey et al., 2012). Neuronal preference for choices was

prominent in the delay period of the memory task but diminished in control non-
memory behavioral tasks. To summarize, this novel empirical evidence from rodent
studies on the Tmaze behavioral apparatus complements the coding framework of short-
term memory with more dynamic and adaptive information-processing mechanisms
other than persistent activity. Notably, by averaging neuronal population activities, none
of these computational schemes would have been revealed.

In studying the role of DA neurons in short-term memory, we should take into
consideration that the DA neuronal circuit is by no means self-contained and therefore it
should not be investigated in isolation. Neurons utilizing GABA as a neurotransmitter
constitute approximately 30% of the VTA neuronal population. The memory-related
encoding properties of these inhibitory neurons have been largely overlooked, despite
evidence of a strong inhibitory influence on neighboring DA neurons (Nair-Roberts et
al., 2008; Omelchenko and Sesack, 2009; Tan et al., 2012; van Zessen et al., 2012) and
well-established interconnections with the PFC circuit (Carr and Sesack, 2000a, b).

In light of the above, we wished to investigate with fine temporal and spatial resolution
the firing activity of optogenetically identified DA and GABA neurons while mice
performed a T-maze reward-seeking task with memory load. We took into consideration
that (i) earlier studies analyzed either the population activity of putative DA neurons or
drew inferences of the population activity from voltammetry measurements (Ljungberg
et al., 1991; Schultz, 2002; Phillips et al., 2004; Matsumoto and Takada, 2013; Choi et
al., 2020), and (ii) field potentials (like the 4Hz oscillations recorded in the VTA) stem
mainly from phase-aligned excitatory or inhibitory post-synaptic potentials, whereas
spiking activity is sparse (Traub et al., 2004; Buzsaki, 2006). (iii) A recent report
revealed the causal relationship of DA activity with short-term memory, by inhibiting
DA neurons with optogenetic tools. However, they did not report the encoding
properties of single VTA DA neurons (Choi et al., 2020). Therefore, we looked for hints
of memory-related encoding activities in single DA and GABA neurons by
characterizing their firing preference for opposite behavioral choices.

Results

Optogenetic identification of DA and GABA neurons in the VTA.

In the present study, we sought to investigate the encoding properties of DA and GABA neurons of the VTA while mice engage in memory-dependent reward-seeking behavior. To identify neurons, we expressed the light-gated cation channel, channelrhodopsin-2 (ChR2), in DA and GABA neurons by injecting an adeno-associated virus containing FLEX-ChR2 into DAT-Cre and VGAT-Cre transgenic mice, respectively (Bäckman et al., 2006; Tsai et al., 2009; Vong et al., 2011) (Figure 1A). Optogenetic identification and parallel electrophysiological recordings were performed using a custom-made diode-probe system (diode-fiber assemblies attached to high-density silicon probes(Stark et al., 2012), (Figures 1A and S1). For each neuron, we assessed the response to light pulse trains delivered before and after behavioral sessions (Figures 1 and fig. S1 and S2). We identified 104 neurons recorded from five DAT-Cre mice (hereafter referred to as DA neurons) and 74 neurons recorded from four VGAT-Cre mice (GABA neurons) with significant excitatory responses to light pulses (Figure 1B). Light-induced spikes from these neurons were almost identical to spontaneous spikes (waveform correlation coefficient > 0.9, Figure S1E and 1F). In addition, the electrophysiological profiles of the identified neuronal populations resembled those of previous studies (i.e., DA neurons fired action potentials with both wider waveforms and slower spontaneous firing rates than GABA neurons; Figure S1G), confirming the selective expression of ChR2 in DA and GABA neurons (Cohen et al., 2012; Tan et al., 2012).

Behavioural Performance in a memory-dependent decision-making task.

Mice were trained to perform sensory-guided and memory-dependent decisions in the "Memory Task" (Figures 2A, S3A and S3B). This task required animals to associate a visual cue presented at the beginning of the trial with a rewarded side arm of a figure-eight T-maze. A short memory delay was introduced between cue presentation and

action selection. Following a correct response, they received water (5 μl) from a

waterspout located at the end of each arm. Depending on the individual features of

cognitive demand, the maze apparatus was divided into separate sections (i.e., "start,"

"cue," "delay," "side arms," and "reward"). To ensure that the mice made choices

guided by the visual cues and had minimal influence from other behavioral parameters

on decisions, we eliminated imbalances between the left and right trials in key task

parameters (e.g., reward amount, visual environment, effort, and motor skill

requirements).

At the time of neurophysiological data collection, all mice performed memory task trials

with high accuracy. Averaging across sessions, the total correct rate was $86.8 \pm 7.9\%$

(mean ± standard deviation [SD]; left: $88.1 \pm 10.0\%$, right: $87.2 \pm 11.6\%$; paired t-test

evaluating left vs right performance rate: $t(59) = 0.46$, $P = 0.65$, 60 sessions in nine

mice, Figure 2B). In addition, performance was independent of individual preference for

the left-or-right arm visits in any of the recorded sessions (test of independence, $\chi^2(1) <$

3.84, $P > 0.05$, Ho: correct rate is independent of arm choice, Figure S3C).

We also assessed the contribution of memory-related processing to task performance.

To achieve this, we delivered blocks of trials with different memory loads in separate

training sessions. Across all sessions, the correct performance rate dropped with higher

memory load demands (mean ± SD; low load: $83.1 \pm 8.3\%$, high load: $73.9 \pm 9.5\%$;

paired t-test on correct performance rate: $t(12) = 4.33$, $P < 0.001$, 13 sessions in seven

mice, Figure 2C). This result is consistent with earlier reports (Floresco and Phillips,

2001; Floresco and Magyar, 2006) and highlights the important role of memory in

supporting decisions in the present task.

The population activity of DA neurons is not elevated during the memory task

trials.

DA neurons are not known to be active in the delay period of short-term memory tasks

(Ljungberg et al., 1991; Schultz et al., 1993; Phillips et al., 2004; Matsumoto and

Takada, 2013; Choi et al., 2020), even though DA is a key neurotransmitter in the

regulation of prefrontal cortical mnemonic functions (Goldman-Rakic et al., 1989;

Smiley et al., 1992; Smiley and Goldman-Rakic, 1993; Goldman-Rakic, 1997;

Tzschentke, 2001). This well-established notion has been mainly inferred from the

average analysis of putative neuronal population activities and striatal voltammetry

recordings (Ljungberg et al., 1991; Schultz, 2002; Phillips et al., 2004; Matsumoto and

Takada, 2013; Choi et al., 2020). Corroborating these earlier reports, the average

discharge rate of identified DA neurons in the present study remained essentially

constant (Figure 2D). Simple linear regression analysis, with the neuronal firing rate as

the response variable and the animal's position on the maze as the single predictor

variable, showed that from the beginning until the end of the trial (a 1.5-meter distance),

the population activity of DA neurons deviated slightly by 0.17 ± 0.62 Hz (mean $\pm$

standard error of the mean [SEM], did not differ from a distribution with a mean equal

to zero; one-sample t-test on the position coefficient, $t_{(103)} = 0.275$, $P = 0.78$). Notably,

in the memory-delay period, the discharge rate of DA neurons declined by $- 0.72 \pm 2.3$

Hz (mean $\pm$ SEM, one-sample t-test on the position coefficient, $t_{(103)} = -0.31$, $P = 0.75$).

On the other hand, the GABA neurons elevated their discharge rate by 4.29 ± 1.10 Hz in

the delay period (mean $\pm$ SEM, one-sample t-test on the position coefficient, $t_{(73)} = 4.09$,

$P < 0.001$), confirming evidence from an earlier report (Cohen et al., 2012).

DA and GABA neurons in the VTA show trajectory-specific encoding preferences

in short-term memory-dependent behaviors

Making interpretations of the encoding properties of single neurons from population

rate averages is highly challenging in tasks with many behavioral choices, especially for

functionally heterogeneous populations such as the DA neurons. To overcome this

limitation, we analyzed the firing activity of single neurons, by taking into consideration

two important behavioral parameters. First, in every trial, the animals visited either the

left or right rewarded side arms. Therefore, we grouped and averaged trial spike trains

of single neurons by the corresponding lap trajectories (left or right; see also Methods

and Figure S4A). Also, in the present task, significant behavioral events (including

visual cue presentation, memory delay, and reward delivery) were inherently bound to

fixed positions in the maze (Figure 2A, S3A and S3B). Thus, we arranged spiking

events according to their position, to get an estimate of the behavioral correlates of

neuronal activity. To this end, individual trial trajectories were linearized and

represented as a one-dimensional vector consisting of 100 linearly spaced points (trial

start: point 0; trial reward: point 100).

Examples of discharge patterns arranged by position and trajectory are shown in Figures

3A, S5A and S5B. These representative neurons differentiate their responses between

left and right trials at certain maze positions in a robust manner. To assess the

trajectory-specific effects on neuronal firing activity, we used the permutation method

(Figure S4). First, we calculated the original difference between the average firing rates

in the left and right trials. We then randomly reassigned the trajectory labels (left or

right) on the trial spike trains and produced the permuted average firing rate differences.

If neurons were modulated by trajectory, the original and permuted firing rate

differences were significantly different. Since spiking events are arranged by position,

the permutation method can also detect positions with significant differences.

Figure 3B summarizes the results from the permutation analysis applied to the

populations of 104 DA and 74 GABA neurons. In the left and middle heatmaps, we

organized the normalized mean firing rates for the preferred and non-preferred lap

trajectories (i.e., trajectories with higher and lower firing rates respectively). The right

heatmap contains the maze positions with significant firing rate differences. In both

neuronal populations, there was abundant trajectory-specific activity, concentrated

mostly in the delay and side-arm sections. Almost 20% of DA neurons differentiated

their response between left or right trajectories in those maze sections (21% in the delay

section and 22% in the side-arm section, 104 neurons, permutation test, $P < 0.05$, Figure

3B and Table S1). In GABA neurons, the percentage was even higher, with almost

50% of these cells eliciting trajectory-specific activities (47% in the delay section and

47% in the side-arm section, 74 neurons, permutation test, $P < 0.05$, Figure 3B and

Table S1).

There have been reports of DA neurons discriminating between visual cues in a T-maze

task (Engelhard et al., 2019) or choice selections in delayed-match-to-sample tasks

(Matsumoto and Takada, 2013; Choi et al., 2020). However, in the present study, VTA

neurons did not exhibit different levels of activity in the cue section. Furthermore,

neuronal preference for trajectories was not restricted to the turning point, which could

indicate neuronal engagement in motor preparation for choice execution. Instead, it was

spread in a wider area, covering a distance from the memory delay onset until the end of

the side arms.

A plausible explanation for the trajectory-specific responses in the side arms is that

neurons were under the control of the sensory, motor, or goal-directed behavioral

processes triggered by the opposite trajectories (Howe et al., 2013; Hamid et al., 2016;

Mohebi et al., 2019). However, in the memory-delay section, trajectories were identical

for the left and right trials, which could be suggestive of the engagement of these

neurons in short-term memory processing. Neuronal preferences to arm visits in

memory delay are not uncommon in T-maze tasks. They have been reported in

prefrontal and post-parietal cortical neurons and have been attributed to short-term

memory-dependent decisions (Fujisawa et al., 2008; Harvey et al., 2012). So, is the

trajectory-specific activity in our task reminiscent of internally generated, memory

representations, or can be attributed to the well-known DA-linked neuronal

computations (Schultz, 2002; Cohen et al., 2012; Berke, 2018; Engelhard et al., 2019)?

To test this hypothesis, we proceeded to a series of statistical analyses and control

behavioral tasks.

Multiple regression analysis confirms the trajectory-specific effect on DA and

GABA neurons.

We discovered that significant proportions of VTA neurons fired preferentially for left

or right trajectories at specific locations on the maze when we arranged discharge

patterns by arm visit and position. This result does not attest that trajectory and position

alone contribute to the neuronal firing rate. Midbrain DA neurons are known to respond

to a wide variety of behavioral parameters (i.e., choice accuracy, reward history,

running speed, and distance to rewards (Engelhard et al., 2019)) which could also exert

a significant effect on neuronal firing activity. However, their effect could be dampened

due to the specific firing range arrangement.

Since these behavioral variables are difficult to control with behavioral tasks, we

assessed their contribution to neuronal responses using multiple regression analysis

(Figure S6 and Supplementary Text 1). We found that all the examined variables (lap

trajectory, trial number, speed, trial accuracy, and reward history) contributed to the

firing activities of neuronal subpopulations; however, only the lap trajectory predictor

could explain better the trajectory-specific activities observed in the ~20% of DA and

~50% of GABA neurons that were identified with the permutation analysis.

276   Memory-dependent but not motivated behavior is related to trajectory-specific

277   activity in VTA neurons

Next, we investigated the contribution of short-term memory in decision-making on the

trajectory-specific activity of VTA neurons. Memory-dependent decision-making

depends on three major computational components. These are (i) sensory input gating,

(ii) maintaining and manipulating memory contents and (iii) generating and executing

appropriate motor plans (Cohen et al., 2002; Dreher and Burnod, 2002; Montague et al.,

2004; Ott and Nieder, 2019).

We eliminated all three components in a variation of the memory task. Specifically, we

trained mice in the no-cue-no-choice task, in which they were not presented with a

visual cue and, therefore, could not make predictions about the location of the reward

(Figures 4A, S3A and S3B). Furthermore, the choice selection was prevented by the

presence of blocked side arms when they arrived at the T-intersection. After a short

delay (approximately 1 s), access to one of the side arms (chosen pseudo-randomly) was

permitted, which always led to a reward.

In recording sessions, mice received mixed protocols composed of randomly interleaved

memory task and no-cue-no-choice task trials. We evaluated the trajectory-specific

activities on each task separately using the permutation method. In the delay section of

the no-cue-no-choice task, we observed a significant reduction in the number of

positions with a significant firing rate difference (mean ± SD; DA: memory task 5.9 ±

3.8 points, no-cue-no-choice task 1.5 ± 3.4 points, paired t-test, $P = 0.002$, four animals;

GABA: memory task 10.7 ± 3.9 points, no-cue-no-choice task 1.5 ± 2.0 points, paired t-

test, $P < 0.001$, three animals, Figures 4B to 4E). The attenuating effect on trajectory-

specific activity was also reflected by a marked reduction in the number of trajectory-

specific neurons (Figure 4E, numbers in parentheses). However, the firing rate

difference between the left and right-side arms was strong in both tasks (DA: memory

task 8.5 ± 7.8 points, no-cue-no-choice task 5.9 ± 7.4 points, paired t-test, $P = 0.139$,

four animals; GABA: memory task 12.6 ± 8.1 points, no-cue-no-choice task 9.9 ± 8.8

points, paired t-test, $P = 0.234$, three animals, Figures 4B to 4E).

However, the significant reduction in trajectory-specific encoding preference in the no-

cue-no-choice task could not be entirely attributed to the absence of the memory

component. This is because important running speed, motor responses, and motivational

discrepancies exist between memory and no-cue-no-choice tasks. With regard to

motivation, the important role of DA in adaptive decision-making is widely recognized

(Hamid et al., 2016; Berke, 2018; Mohebi et al., 2019). We did not observe animal

choice bias in memory task performance (Figures 2B and S3C), but we cannot rule out

the possibility that individual neurons were modulated differently by effortful actions to

reach the left- and right-sided rewards (Figure S7). Unlike the memory task, in the no-

cue-no-choice task, the mice could not direct behavior towards the left- or right-side

arms due to the absence of a visual cue. As a result, they were unable to allocate

incentive motivational drives to the left-or-right trials (Howe et al., 2013; Hamid et al.,

2016; Berke, 2018; Mohebi et al., 2019). Also, the initial access denial to the side arms

in the control task eliminates any potential differences in the motor preparation coding

schemes (according to the "gating theory") for the opposite arm visits in the memory

task (Engelhard et al., 2019; Ott and Nieder, 2019). Finally, regarding speed, a

representative example of running speed differences between the two tasks within a

single session is shown in Figure 4F.

To dissociate the short-term memory component of neuronal activity from the

modulatory effects of running speed, incentive motivation, and motor-related signaling,

we trained mice in a second control task. The cue-no-choice task preserved the same

running speed parameters (Figure S8), motor skill requirements, and physical effort

demands (i.e., visual cues, maze shape, arm length, and reward amount were the same)

as the memory task, but it prevented animals from making decisions. Accordingly, the

animals were presented with the same visual cue as in the memory task, which indicated

the side arm that was rewarded and enabled them to allocate incentive motivational

drive to the left-or-right trials; however, they were always forced to visit the rewarded

arm by blocked access to the unrewarded arm (Figures 5A, S3A and S3B). In the same

recording session, the mice performed a separate block of memory task trials. Similar to

the first control task, in the delay section of the cue-no-choice task we observed a

significant reduction in the spatial extent of the firing rate difference (DA: memory task

5.4 ± 3.6 points, cue-no-choice task: 0.4 ± 1.1, paired t-test, $P = 0.011$, four animals;

GABA: memory task 10.7 ± 6.3 points, cue-no-choice task 3.5 ± 4.2 points, paired t-

test, $P < 0.001$, 1 animal, Figures 5B to 5E). In the side arms, however, the trajectory-

specific effect remained strong and was not significantly different from the effect

observed in the memory task (DA: memory task 8.1 ± 10.3 points, cue-no-choice task

5.0 ± 4.6 points, paired t-test, $P = 0.146$, four animals; GABA: memory task 8.7 ± 5.9

points, cue-no-choice 5.2 ± 6.2 points, paired t-test, $P = 0.086$, 1 animal, Figures 5B to

5E).

Together, these results suggest that trajectory-specific responses in the delay period of

the memory task could reflect short-term memory representations linked to decision-

making behavior and cannot be explained by running speed, motor, and motivation-

related signaling differences.

Neuronal activities in delay and reward are unrelated

It is widely recognized that DA and GABA neurons of the VTA respond to reward and

reward-predicting cues (Schultz et al., 1993; Schultz et al., 1997; Cohen et al., 2012;

Matsumoto and Takada, 2013; Engelhard et al., 2019; Choi et al., 2020). Therefore, we

examined whether the selective neuronal responses to left-or-right trials in the memory

delay and side-arm sections could be attributed to encoding biases related to rewards.

We defined the first second of reward consumption as the reward section.

DA neurons are known to be excited by rewards (Schultz et al., 1993; Schultz et al.,

1997; Cohen et al., 2012; Matsumoto and Takada, 2013; Engelhard et al., 2019; Choi et

al., 2020). In agreement with this notion, we discovered that 27 DA neurons (28% of

104 neurons, Figure 6B column 4) responded to reward with significant excitation,

which is a high proportion considering the low uncertainty of the task due to the high-

performance rate and overtraining.

We subsequently examined whether VTA neurons could discriminate between left- and

right-arm rewards. DA neurons are known to discriminate between rewards with

different magnitudes and predictabilities(Tobler et al., 2005; Morris et al., 2006). The

animals were offered equivalent options in terms of reward magnitude, uncertainty, and

effort. Thus, we predicted the presence of a small number of reward-discriminating

neurons. However, we found that 23% of DA neurons and 46% of GABA neurons

differed significantly in their responses to left-or-right rewards (Figures 6A, 6B column

3, and Table S1; paired t-test comparing mean firing rates, $P < 0.05$). Therefore, we

assessed the relationship between encoding preferences in the reward section and those

in the remainder of the maze by correlating the average firing rate difference in the

reward section with the average rate difference in the preceding maze sections. In both

neuronal populations, we found a significant positive relationship between the reward

and side-arm sections (Figure 6C; Pearson's correlation; DA: $R = 0.31$, $P < 0.001$;

GABA: $R = 0.67$, $P < 0.001$) but not between the reward and delay sections (Figure 6C;

Pearson's correlation; DA: $R = -0.05$, $P > 0.05$; GABA: $R = -0.03$, $P > 0.05$). These

results demonstrate that trajectory-specific firing activities in memory delay do not

reflect reward preference during consumption.

Discussion

In the present study, we performed extracellular recordings from optogenetically identified DA and GABA neurons in the VTA while mice performed reward-seeking tasks on a T-maze apparatus. Mice were trained to choose between two spatially separate goals under the instruction of visual cues presented at the beginning of the trials. A short memory delay was introduced between cue presentation and choice selection. We discovered that subpopulations of DA and GABA neurons showed differential responses between the left and right trials, starting from the onset of the memory delay period in the main arm, where the trajectories were indistinguishable. Trajectory-specific preference was not correlated with reward history, running speed, the incentive motivational drive of physical effort, or reward-related encoding differences, and diminished significantly when the memory-dependent decision component was eliminated in control behavioral tasks. This evidence indicates that populations of DA and GABA neurons in the VTA encode internally generated signals that support short-term memory in decision-making.

Activities of midbrain DA neurons in short-term memory

The well-established notion that DA somatic spiking activity is low in short-term memory stemmed mainly from firing rate averages of putative neuronal populations (Ljungberg et al., 1991; Schultz, 2002; Phillips et al., 2004; Matsumoto and Takada, 2013; Choi et al., 2020). Consistent with those reports we did not observe a profound variation in the population activity of DA neurons during the memory task. However, a wealth of recent studies has shown that DA neurons are functionally and genetically segregated (Lammel et al., 2008; Lammel et al., 2011; Engelhard et al., 2019). Moreover, in many real-life situations, animals have to choose between many options for behavioral responses. In such behavioral conditions, averaging firing rates cannot reveal the fine computational processes of single neurons. To address this limitation, we analyzed the firing activities of identified single neurons, focusing on different discharge patterns between behavioral choices. Here, we demonstrated that memory-

specific activities by midbrain DA neurons can be represented as trajectory-specific

responses in the delay period of the memory task.

The "gating theory" unifies the signaling activities of DA neurons in reward prediction

and short-term memory (Cohen et al., 2002; Dreher and Burnod, 2002; Montague et al.,

2004; Ott and Nieder, 2019). Our evidence does not question the validity of this

computational model, since we do not provide evidence of how the selective preference

for one response over the other translates into the release site. However, it is worth

noting that trajectory-specific activities we observed here were rare in the visual-cue

section when the updating of sensory information takes place. Moreover, firing rate

differences between left and right-arm responses declined in a control behavioral task

(cue-no-choice) without memory load, but with the same motor skill requirements as the

memory task. Thus, motor preparation coding schemes for different responses (Cohen et

al., 2002; Ott and Nieder, 2019) cannot be accounted for the trajectory-specific

activities in the memory delay. Instead, our evidence indicates that trajectory-specific

activities by DA (also GABA) neurons are functionally linked to the maintaining and

manipulating of memory contents.

Memory-related, trajectory-specific neuronal activities have also been reported for PFC

and post-parietal neurons while rodents perform reward-seeking responses on the T-

maze (Fujisawa et al., 2008; Harvey et al., 2012). Furthermore, in the same task, both

the PFC and VTA circuits are functionally engaged with coherent 4-Hz oscillations in

the memory delay (Fujisawa and Buzsáki, 2011). Our results complement this line of

work, demonstrating that also DA and GABA midbrain neurons elicit memory-related,

trajectory-specific activities.

The present study also corroborates important findings from a recent report, which

demonstrated that optogenetic perturbations in DA neuron excitability exert a strong

effect on short-term memory performance, highlighting the causal role of DA neuronal

firing activity in memory-dependent behavior (Choi et al., 2020). Also, in agreement

with a previous report (Engelhard et al., 2019) by the same laboratory, we show that

subpopulations of VTA neurons are modulated by running speed, cumulative

performance rate, current choice accuracy, and reward history. Disparities between this

and our study in the proportions of modulated neurons could be attributed to the
different recording techniques applied as well as the maze regions of interest; for
example, Engelhard et al. analyzed neuronal firing activities in the visual-cue period
(Engelhard et al., 2019), whereas we focused on memory delay.

Overall, our results are in agreement with the notion that DA neurons encode a variety
of behavioral parameters in complex environments. In addition, we confirmed that in
memory-dependent behaviors, DA neuronal populations did not elicit sustained
increases in their discharge rate. However, in the present task, DA neurons individually
encoded internal representations by differentiating their responses to lap trajectories in
memory delay.

Role of motivated behavior in trajectory-specific encoding properties of VTA
neurons

Midbrain DA activity is known to be involved in motivated behavior while rodents
navigate mazes to receive rewards in trials lasting a few seconds (Hamid et al., 2016;
Berke, 2018; Mohebi et al., 2019). When mice approach rewards, striatal DA
concentrations increase, scaling flexibly with reward size and proximity, which is
proposed to reflect a neural correlate of a sustained motivational drive (Howe et al.,
2013). To evaluate the role of motivated behavior in the trajectory-specific preference
of midbrain neurons, we compared firing activities between a memory task and a
control task without memory-dependent decisions (cue-no-choice task). Although in the
cue-no-choice task, the behavioral parameters that determined the incentive
motivational drives were the same as in the memory task (visual cues, maze shape, and
reward amount), neuronal responses did not differ between the left and right trials. This
result strongly indicates that incentive motivational drives (at least for physical effort)
do not contribute to trajectory-specific activities of midbrain neurons during the delay
period of the memory task.

Memory-specific activities of the VTA neurons are not attributed to reward prediction error signaling

We also assessed the role of reward-related processing in the trajectory-specific activity of the midbrain neurons. In behavioral tasks in which animals estimate the spatial proximity of distant rewards, it has been suggested that DA neurons calculate RPE signals from state-value functions (Hamid et al., 2016; Berke, 2018; Engelhard et al., 2019; Mohebi et al., 2019; Kim et al., 2020). In the present study, the animals received ongoing visual input, facilitating the continuous estimation of reward proximity. Thus, DA neurons can potentially estimate RPE signals from successive state values assigned to each position on the maze track (Figure S7). Therefore, the difference in firing activity between the left and right trials could be the result of differences in the state-value functions assigned to these trajectories (Hamid et al., 2016; Berke, 2018). However, significant evidence contradicts this hypothesis.

First, the behavioral parameters that determine the state-value functions for the left and right trajectories were set to be identical in the cue-no-choice task and memory task, by preserving the same maze configurations and delivering equal amounts of reward. In addition, behavior in both tasks was cue-driven; therefore, animals could make predictions about the reward location and orchestrate behavior accordingly. However, we observed a prominent reduction in the firing rate difference between the left and right trials in the cue-no-choice task (Figure 5). Second, a significant subset of DA neurons (approximately 20%) responded differently to the left and right rewards in the memory task, although the same amount of reward was delivered. This unexpected finding raised the hypothesis that the encoding preference for reward could be reflected in the values of the preceding states in the maze and, therefore, could account for the trajectory-specific effect in memory delay. However, the differences in the firing activity elicited by the consumption of left-or-right rewards were unrelated to the firing rate difference in the delay section (Figure 6C). In conclusion, these findings indicate that the encoding preference for lap trajectories exhibited by midbrain DA and GABA neurons cannot be simply explained by discrepancies in RPE signaling.

GABA neurons of the VTA and short-term memory

With the advent of highly selective identification and perturbation techniques, new evidence has emerged regarding the encoding properties and functional roles of local VTA inhibitory networks in reward processing and motivation. There are reports demonstrating that GABA neurons of the VTA suppress reward consummatory behavior (van Zessen et al., 2012), facilitate aversive behavior (Matsumoto and Hikosaka, 2007; Tan et al., 2012), and elicit sustained activities in the delay period between conditioned and unconditioned stimuli (Cohen et al., 2012). During these behaviors, the responses of DA and GABA neurons are often inverse, such that when GABA neurons are excited, neighboring DA neurons decrease their discharge rate. In particular, aversive stimuli excite GABA neurons, which then suppress the neighboring DA neurons (Tan et al., 2012). In addition, during reward consumption, GABA neurons are inactive (Cohen et al., 2012; van Zessen et al., 2012); however, when excited, they inhibit DA neurons and disturb consummatory behavior (van Zessen et al., 2012). Finally, in classical conditioning tasks, DA neurons respond to rewards and reward-predicting stimuli, whereas GABA neurons remain silent during such events (Cohen et al., 2012). However, we demonstrated here that midbrain DA and GABA neurons elicit remarkably similar encoding properties. Both neuronal populations respond to short-term memory-specific activities manifested by encoding preferences for lap trajectories. Notably though, GABA neurons are more strongly engaged in this dynamic encoding activity since almost twice as many inhibitory neurons responded differently to the left and right trials.

This result presents an activity paradox. Given the abundant and potent synaptic inhibition of DA neurons by neighboring GABA neurons (Omelchenko and Sesack, 2009), it was unexpected that both populations were highly active and similarly engaged in tasks. However, anatomical evidence provides a plausible explanation. Local inhibitory neurons form a dense network of local synaptic innervations that target the dendritic sites of DA and other GABA neurons (Traub et al., 2004; Buzsaki, 2006; Omelchenko and Sesack, 2009). Although potent and well-suited for coordinated network activity, this synaptic inhibition is not as strong as somatic inhibition (Jhou et al., 2009; Omelchenko and Sesack, 2009), and it has been suggested that it is not

sufficient to suppress DA neurons when they receive a strong excitatory drive from

extrinsic sources (van Zessen et al., 2012).

Conclusion

In summary, we optogenetically probed DA and GABA neurons in the VTA while mice

performed a decision-making task with memory load. We discovered that both of these

neuronal populations elicited memory-dependent preferences for left-or-right

trajectories that could not be explained by motor activity, motivated behavior, or

reward-related processes. This evidence indicates that VTA neurons encode mental

representations to support short-term memory-dependent decisions and provides

insights into novel sophisticated coding strategies employed by the midbrain DA and

GABA neurons in reward-related behavior.

553 **Methods & Materials**
554

555 **Key resources table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti-Tyrosine Hydroxylase	EMD Millipore	
Anti-GFP	Aves Labs	
Alexa Fluor 546 IgG	Thermo Fisher Scientific	
Alexa Fluor 488 IgG	Jackson Immunoresearch Laboratories, Inc.	
DAPI Fluoromount-G	SouthernBiotech	
Bacterial and virus strains		
AAV5-EF1a-DIO-hChR2(H134R)-EYFP-WPRE-pA	UNC Vector Core	N/A
Experimental models: Organisms/strains		
Mouse: C57BL/6J-Slc6a3$^{tm1.1(cre)Bkmn/J}$	The Jackson Laboratory	Jax #006660; RRID: IMSR_JAX:006660
Mouse: C57BL/6J- Slc32a1$^{tm2(cre)Lowl}$	The Jackson Laboratory	Jax #0016962; RRID: IMSR_JAX:0016962
Software and algorithms		
MATLAB	MathWorks	https://www.mathworks.com/
KlustaKwik2	(Kadir et al., 2014)	https://github.com/kwikteam/klustakwik2
RPvdsEX	Tucker-Davis Technologies	https://www.tdt.com/component/rpvdsex/
Other		
Silicon probe: 64-site, 6-shank	NeuroNexus	Buzsaki64spL

Optical fiber AFS50/125Y	Thorlabs, Inc.	https://www.thorlabs.co.jp/thorproduct.cfm?partnumber=AFS50/125Y
Blue laser diode	OSRAM Opto Semiconductors	PL450B
256-channel Multiplexed Biosignal Amplifier	Amplipex, Ltd.	KJE-1001
Figure-eight T-maze apparatus	O'Hara & Co., Ltd.	http://ohara-time.co.jp/

CONTACT FOR REAGENT AND RESOURCE SHARING

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and

will be fulfilled by the Lead Contact, Dr. Shigeyoshi Fujisawa

(shigeyoshi.fujisawa@riken.jp).

Animals

All experiments were approved by the RIKEN Institutional Animal Care and Use

Committee. We used five adult male DAT-ires-Cre (Jackson's Lab; stock #6660;

(Bäckman et al., 2006) and four Vgat-ires-Cre (Jackson's Lab; stock #16962; (Vong et

al., 2011) mice backcrossed to C57BL/6J. Animals were housed in separate cages on a

12-h dark/light cycle and each performed the behavioral tasks at the same time of the

day, between 11:00 and 17:00. In the cage, they were provided *ad libitum* food access

but were restrained from water availability.

Behavioral tasks

All behavioral tasks took place on a T-maze apparatus. More information about the

maze configuration is provided in Figure S3.

Memory task. Behavioral sessions commenced with the animal being placed at the

"starting position" (Figures 2A, S3A and S3B). Then, access to the main corridor was

provided and the animal had to run through the "start" section (0-50 cm) before it

arrived at the maze segment surrounded by two PC monitors ("visual-cue" section, 50-

80 cm). In this section, it was presented with a distinctive visual object (vertical black

and gray bars) in one of the two monitors (the other monitor remained dark) indicates

which side arm to visit to obtain the reward (i.e., left cue → left reward, right cue →

right reward). In the third region of the central arm ("delay" section, 80-120 cm) both

monitors turned dark. While running in the delay section, the animal had to maintain the

reward-related information and based on that perform the action selection at the T-

intersection. The intersection at the end of the main arm designated the end of the delay

section and the beginning of the "side-arms" section (120-150 cm) where the animal

runs towards the reward position in anticipation of the reward. Reward (5 μl water) was

delivered on correct trials at the end of the side-arms section from a waterspout. The

first activation of the light-beam sensor at the waterspout triggered the water-delivery

pump (Burkert, Ingelfingen, Germany), followed by reward consumption ("reward"

section). After consuming the reward, the animal could return of its own will to the

starting position, to commence a new trial.

Daily behavioral sessions consisted of 80-100 trials. Only animals with at least three

successive sessions with an 80% performance ratio or more in the training phase were

been proceeded to surgical operations.

No-Cue-No-Choice task. Trials of this control task were delivered in recording

sessions, interleaved with memory task trials. When the animal entered the visual-cue

section it was not instructed by the visual cue (Figures 4A, S3A and S3B). Also, access

to both side arms was initially denied by closed sliding doors. Approximately 1 s after

the animal arrived at the turning point, one of the sliding doors opened (pseudo-

randomly) providing access to the reward. On every trial, the presentation or absence of

the visual cue could instruct the animal about the task rules (i.e., memory task or no-

cue-no-choice task).

Cue-No-Choice-task. The settings of this control task were the same as the memory
task settings, except for the blockade of the unrewarded side arm (Figures 5A, S3A and
S3B). Thus, the animals were always forced to perform correct choices. Because in both
tasks, the same cue was presented, the animals could potentially be confused about the
trial's task rules. Therefore, memory task and cue-no-choice task trials were delivered in
separate sets within the same recording session. Accordingly, when the animals
completed the set of cue-no-choice task trials (approximately 50 trials), they were
automatically delivered with another set of memory task trials (approximately 50 trials).

Recording sessions lasted approximately 20-30 minutes.

Intracranial surgeries and electrophysiological recording

The surgical process consisted of two separate operations. First, mice (DAT-ires-Cre or
Vgat-ires-Cre) were surgically injected with 200-500 nl of adeno-associated virus
AAV5-EF1a-DIO-hChR2(H134R)-EYFP-WPRE-pA (University of North Carolina
vector core facility; (Tsai et al., 2009) into the VTA stereotaxically (from inferior
cerebral vein AP: ~6.65 mm, from midline ML: - 0.55 mm on the left hemisphere, from
surface 4-4.5 mm, Figure 1A). Ten to fifteen days later, mice were implanted with a
silicon probe in the same AP and ML coordinates (vertical insertion was intended, 0
degrees; Figures 1A, S1A to S1D). We used Buzsaki64spL (NeuroNexus, Ann Arbor,
MI, USA) silicon probes which are composed of 6 shanks (10 mm long, 15 μm thick,
200 μm shank separation) and each shank has 10 recording sites (160 μm^2 each site 0.6-
1.0 MΩ impedance). The silicon probe was attached to a custom-made
micromanipulator and moved gradually to the desired depth position. On every probe
shank, optic fibers were firmly attached to secure an accurate and firm insertion of the
recording channels in the deep midbrain area (Figure S1B). For experiments where light
delivery was required, two of the optic fibers (shanks 2 and 5) were coupled with blue
(450 nm) laser diodes (PL450B, OSRAM Opto Semiconductors). Light dispersion
could potentially cover the axial and transverse span of all 64 channels (Stark et al.,
2012).

During recording sessions, the wide-band neurophysiological signals were acquired

continuously at 20 kHz on 256-channel Amplipex systems (KJE-1001, Amplipex Ltd,

Hungary; (Berényi et al., 2014)). Following surgery, the probe was inserted 45 μm

deeper in intoe brain on a daily basis, until it reached the VTA. Thereafter, the probe

was moved deeper by 20 μm / day. The average recording coordinates for the DAT-Cre

animals are 3.32 ± 0.32 mm (mean $\pm$ standard deviation) rostrocaudal and 0.82 ± 0.17

mm mediolateral, and for the VGAT-Cre animals, 3.52 ± 0.29 mm rostrocaudal and

0.90 ± 0.17 mm mediolateral (Figure S1A).

We cannot exclude the possibility that some neurons were recorded in successive

sessions because clustering analysis was performed on individual sessions.

Data analysis

Unless otherwise stated, data analysis was performed with custom-made programs

designed in MATLAB with Signal-processing and Statistics toolboxes.

Light-stimulation protocols for optogenetic identification. Light stimulation protocols

were delivered before and after the behavioral tasks. They were composed of 1, 2, 3,

and 4 mW blocks of 450 nm light pulses. Each block consisted of 150 square pulses (12

ms pulse duration; 0-1 ms and 11-12 ms contained artifacts) delivered at 1, 2, 3 to 10

Hz. Electrophysiological data recorded during light stimulation and behavioral

protocols within a single session were merged and clustered together.

Statistical analysis for detection of light-responsive units. Neurons with light-induced

responses exceeding the average spontaneous activity were classed as light-responsive.

To identify light-responsive neurons we applied the statistical analysis described in

detail in Figure S2.

Estimation of firing activity during behaviour. To estimate the neuronal firing activity
while animals performed the behavioural task, we took into consideration the primary
goal of this study; that is to look for trajectory-specific encoding properties, as well as
the inherent limitation of the task; that is the experimenter could not control the
temporal precision of the behavioural events. To overcome this limitation, we arranged
firing activity by the animal's position on the maze. To do so, first, we linearized the
trial trajectories and assigned them with a lap trajectory label (left or right). Then, the
linearized products were divided into 100 position points and normalized so that
position 0 corresponded to the starting point of the trial and position 1 to the waterspout.
Second, we constructed post-distance histograms, analogous to the peri-stimulus-time-
histograms (PSTHs), although the time of spiking events was replaced by the position
they occurred (for the purpose of simplicity, also by habit, we will call the post-distance
histograms as PSTHs). To construct accurate PSTHs we considered the exact position
the spikes were discharged and the time the animal occupied this certain position. Let
$n^{(k)}(x)$ be the number of spikes of a single neuron and $t^{(k)}(x)$ be the occupation time
in the x_{th} position point of the k_{th} trial (Figure S4A). Then, $\hat{\lambda}(x) = \frac{1}{K} * \sum_{k=1}^{K}[n^{(k)}(x) /$
$t^{(k)}(x)]$ where K is the number of trials, represents the average firing rate probability
(spikes / sec) at position point x. To examine the trajectory-specific encoding properties
of VTA neurons we produced average firing rate histograms for correct left and right
trials, separately. Then, both histograms were smoothed with a Gaussian Kernel
function (σ= 0.5, length of 20 position points).

Firing rate heatmap construction. To construct the normalized firing rate heatmaps
shown in Figures 3B, 4C, 4D, 5C, 5D, 6B, S5C to S5E and S6B we took the following
steps. First, for every neuron we produced the average firing rate for left and right
correct trials. Second, we normalized both rates by dividing them with the maximum
firing rate of the strongest trajectory response (e.g., for the example shown in Figure
S4A we divided both average firing rates by the maximum rate of the response to the
left trials). Then, the normalized rate of the stronger trajectory response was assigned to
the "preferred" heatmap and the rate of the weaker trajectory response to the "non-

preferred" heatmap (e.g., for the example shown in Figure S4A, the left normalized rate
was assigned to the "preferred" heatmap and the right rate to the "non-preferred"
heatmap). Both rates occupied the same row. The row ordering was determined by the
position of maximum rate.

Identifying trajectory-specific neurons with the permutation method. To identify
neurons with trajectory-specific encoding properties we applied the permutation test
reported elsewhere (Fujisawa et al., 2008) and described in detail in Figure S4.

Regression analysis. We designed a generalized linear regression model (GLM) with
the neuronal firing rate (FR) modelled as a gaussian function of the lap trajectory (T),
speed (S), trial number (TN), performance (R), current trial accuracy (A_0) and previous
trial accuracy (A_{-1}) behavioural variables. With the permutation analysis we observed
that the trajectory-specific effect on the firing activity was dependent on position. Thus,
we examined the joint effect of trajectory with position (P) on spiking activity. All
dependent and independent variables were arranged by position. The values of the
trajectory (1 for left and 2 for right), trial number, performance (cumulative correct
rate), current trial and previous trial accuracy (1 for correct trial, 0 for error trial)
variables remained constant throughout the whole trial. The firing rate, position and
speed variables changed their values on every position.
The GLM was:

$$FR = \beta_0 + \sum_{k=1}^{6} \beta_{T*P} T \cdot P^k + \sum_{k=1}^{6} \beta_S \cdot S^k + \beta_{TN} \cdot TN + \beta_R \cdot R + \beta_{A_0} \cdot A_0 + \beta_{A_{-1}} \cdot A_{-1} + \varepsilon$$

where the β values are the regressor coefficients for the different predictors (including
the intercept β_0) and ε is the Gaussian noise term. The 6th degree order polynomials of
position and speed were chosen for model optimization with the Bayes information
criterion.

First, we generated model predictions of the average firing rates for left ($\hat{L}_0$) and right
($\hat{R}_0$) trials, and from those we calculated the predicted firing rate difference ($\hat{D}_0$). Then,

we shuffled the trajectory labels assigned to the tested variable (the assigned labels to

the rest of the independent variables remained intact) and assessed the effect on the

firing rate difference. For every predictor we produced 500 shuffled rate differences, $\widehat{D}_j$.

If the absolute mean value of $\widehat{D}_0$ exceeded the top 5% of the $\widehat{D}_j$ values (including

Bonferroni correction), then the hypothesis was rejected, and the predictor was

significantly contributing to the firing rate difference. We examined every maze region

individually, but, here we report only for the delay region.

Reward-related excitation or inhibition. The reward section was defined as the first

second of reward consumption. To assess neuronal response to reward consumption and

categorize it as excitatory, inhibitory or non-responsive we performed the following

analysis. First, we produced the smoothed mean firing rate response in the time domain

(as we did in the maze sections in the space domain) for left, $\hat{\lambda}_{Left}(t)$, and

right, $\hat{\lambda}_{Right}(t)$ trials. For the preferred arm of each neuron, we compared the mean

firing rate in the reward section to the mean rate in the 100 ms epoch preceding reward

delivery (paired t-test on mean firing rates; *P < 0.05*; Figure 6B column 4).

Encoding preferences in the reward section. The difference in the intensity of neuronal

firing activity between left and right rewards was assessed by comparing the mean

firing rate of neuronal activity elicited in the reward section of left and right trials

(paired t-test on mean firing rates; *P < 0.05*; Figure 6B column 3).

Relationship of encoding preferences in the reward section to those in the remainder

of the maze. To assess whether the trajectory-specific firing activity in the maze was

linked to discrepancies in the response to left and right rewards, we followed the next

steps of analysis. First, for every neuron and every maze section, we calculated the

mean value of the relative firing rate difference between left and right trials ($\widehat{D}_{start}(x)$,

$\widehat{D}_{cue}(x)$, $\widehat{D}_{delay}(x)$, $\widehat{D}_{arms}(x)$, $\widehat{D}_{reward}(t)$). Then, for each neuronal group, we

calculated the linear relationship (Pearson's correlation) between the reward section

mean values, to those in the remainder of the maze (Pearson's correlation; Figure 6C).

Immunohistochemistry

After completion of the recording sessions, which lasted about a month, mice were

anesthetized with isoflurane and perfused transcardially with 10 ml PBS and 10 ml

paraformaldehyde (4%), before they were decapitated. Brains were then removed, post-

fixed and coronal slices (100 μm) were prepared. The primary antibodies used were

rabbit anti-tyrosine hydroxylase (TH) and chicken anti-GFP. The secondary antibodies

used were AlexaFluor 549 anti-rabbit and 488 IgG anti-chicken, respectively. Sections

were further stained with DAPI to visualize nuclei. Image acquisition was performed

with a fluorescence microscope NanoZoomer (Hamamatsu, Japan) system.

References

Goldman-Rakic, P.S., Leranth, C., Williams, S.M., Mons, N., and Geffard, M. (1989). Dopamine synaptic complex with pyramidal neurons in primate cerebral cortex. Proceedings of the National Academy of Sciences *86*, 9015-9019.

Ljungberg, T., Apicella, P., and Schultz, W. (1991). Responses of monkey midbrain dopamine neurons during delayed alternation performance. Brain research *567*, 337-341.

Sawaguchi, T., and Goldman-Rakic, P.S. (1991). D1 dopamine receptors in prefrontal cortex: involvement in working memory. Science *251*, 947-950.

Smiley, J.F., Williams, S.M., Szigeti, K., and Goldman‑Rakic, P.S. (1992). Light and electron microscopic characterization of dopamine‑immunoreactive axons in human cerebral cortex. Journal of Comparative Neurology *321*, 325-335.

Schultz, W., Apicella, P., and Ljungberg, T. (1993). Responses of monkey dopamine neurons to reward and conditioned stimuli during successive steps of learning a delayed response task. Journal of neuroscience *13*, 900-913.

Smiley, J.F., and Goldman-Rakic, P.S. (1993). Heterogeneous targets of dopamine synapses in monkey prefrontal cortex demonstrated by serial section electron microscopy: a laminar analysis using the silver-enhanced diaminobenzidine sulfide (SEDS) immunolabeling technique. Cerebral cortex *3*, 223-238.

Goldman-Rakic, P.S. (1995). Cellular basis of working memory. Neuron *14*, 477-485.

Goldman-Rakic, P.S. (1997). The cortical dopamine system: role in memory and cognition. Advances in pharmacology *42*, 707-711.

Schultz, W., Dayan, P., and Montague, P.R. (1997). A neural substrate of prediction and reward. Science *275*, 1593-1599.

Carr, D.B., and Sesack, S.R. (2000a). Projections from the rat prefrontal cortex to the
ventral tegmental area: target specificity in the synaptic associations with
mesoaccumbens and mesocortical neurons. Journal of neuroscience *20*, 3864-3873.

Carr, D.B., and Sesack, S.R. (2000b). GABA‐containing neurons in the rat ventral
tegmental area project to the prefrontal cortex. Synapse *38*, 114-123.

Durstewitz, D., Seamans, J.K., and Sejnowski, T.J. (2000). Neurocomputational models
of working memory. Nature neuroscience *3*, 1184-1191.

Floresco, S.B., and Phillips, A.G. (2001). Delay-dependent modulation of memory
retrieval by infusion of a dopamine D_1 agonist into the rat medial prefrontal cortex.
Behavioral neuroscience *115*, 934.

Miller, E.K., and Cohen, J.D. (2001). An integrative theory of prefrontal cortex
function. Annual review of neuroscience *24*, 167-202.

Tzschentke, T.M. (2001). Pharmacology and behavioral pharmacology of the
mesocortical dopamine system. Progress in neurobiology *63*, 241-320.

Cohen, J.D., Braver, T.S., and Brown, J.W. (2002). Computational perspectives on
dopamine function in prefrontal cortex. Current opinion in neurobiology *12*, 223-229.

Dreher, J.-C., and Burnod, Y. (2002). An integrative theory of the phasic and tonic
modes of dopamine modulation in the prefrontal cortex. Neural networks *15*, 583-602.

Schultz, W. (2002). Getting formal with dopamine and reward. Neuron *36*, 241-263.

Curtis, C.E., and D'Esposito, M. (2003). Persistent activity in the prefrontal cortex
during working memory. Trends in cognitive sciences *7*, 415-423.

Montague, P.R., Hyman, S.E., and Cohen, J.D. (2004). Computational roles for
dopamine in behavioural control. Nature *431*, 760-767.

Phillips, A.G., Ahn, S., and Floresco, S.B. (2004). Magnitude of dopamine release in

medial prefrontal cortex predicts accuracy of memory on a delayed response task.

Journal of Neuroscience *24*, 547-553.

Traub, R.D., Bibbig, A., LeBeau, F.E., Buhl, E.H., and Whittington, M.A. (2004).

Cellular mechanisms of neuronal population oscillations in the hippocampus in vitro.

Annu Rev Neurosci *27*, 247-278.

Wise, R.A. (2004). Dopamine, learning and motivation. Nature reviews neuroscience *5*,

483-494.

Fries, P. (2005). A mechanism for cognitive dynamics: neuronal communication

through neuronal coherence. Trends in cognitive sciences *9*, 474-480.

Tobler, P.N., Fiorillo, C.D., and Schultz, W. (2005). Adaptive coding of reward value

by dopamine neurons. Science *307*, 1642-1645.

Bäckman, C.M., Malik, N., Zhang, Y., Shan, L., Grinberg, A., Hoffer, B.J., Westphal,

H., and Tomac, A.C. (2006). Characterization of a mouse strain expressing Cre

recombinase from the 3′ untranslated region of the dopamine transporter locus.

genesis *44*, 383-390.

Buzsaki, G. (2006). Rhythms of the Brain (Oxford University Press).

Floresco, S.B., and Magyar, O. (2006). Mesocortical dopamine modulation of executive

functions: beyond working memory. Psychopharmacology *188*, 567-585.

Morris, G., Nevet, A., Arkadir, D., Vaadia, E., and Bergman, H. (2006). Midbrain

dopamine neurons encode decisions for future action. Nature neuroscience *9*, 1057-

1063.

Pierce, R.C., and Kumaresan, V. (2006). The mesolimbic dopamine system: the final

common pathway for the reinforcing effect of drugs of abuse? Neuroscience &

biobehavioral reviews *30*, 215-238.

Uhlhaas, P.J., and Singer, W. (2006). Neural synchrony in brain disorders: relevance for
cognitive dysfunctions and pathophysiology. neuron *52*, 155-168.

Berridge, K.C. (2007). The debate over dopamine's role in reward: the case for
incentive salience. Psychopharmacology *191*, 391-431.

Matsumoto, M., and Hikosaka, O. (2007). Lateral habenula as a source of negative
reward signals in dopamine neurons. Nature *447*, 1111-1115.

Vijayraghavan, S., Wang, M., Birnbaum, S.G., Williams, G.V., and Arnsten, A.F.
(2007). Inverted-U dopamine D1 receptor actions on prefrontal neurons engaged in
working memory. Nature neuroscience *10*, 376-384.

Fujisawa, S., Amarasingham, A., Harrison, M.T., and Buzsáki, G. (2008). Behavior-
dependent short-term assembly dynamics in the medial prefrontal cortex. Nature
neuroscience *11*, 823.

Lammel, S., Hetzel, A., Häckel, O., Jones, I., Liss, B., and Roeper, J. (2008). Unique
properties of mesoprefrontal neurons within a dual mesocorticolimbic dopamine
system. Neuron *57*, 760-773.

Nair-Roberts, R.G., Chatelain-Badie, S., Benson, E., White-Cooper, H., Bolam, J., and
Ungless, M. (2008). Stereological estimates of dopaminergic, GABAergic and
glutamatergic neurons in the ventral tegmental area, substantia nigra and retrorubral
field in the rat. Neuroscience *152*, 1024-1031.

Jhou, T.C., Geisler, S., Marinelli, M., Degarmo, B.A., and Zahm, D.S. (2009). The
mesopontine rostromedial tegmental nucleus: a structure targeted by the lateral habenula
that projects to the ventral tegmental area of Tsai and substantia nigra compacta. Journal
of Comparative Neurology *513*, 566-596.

Omelchenko, N., and Sesack, S.R. (2009). Ultrastructural analysis of local collaterals of
rat ventral tegmental area neurons: GABA phenotype and synapses onto dopamine and
GABA cells. Synapse *63*, 895-906.

Robbins, T.W., and Arnsten, A.F. (2009). The neuropsychopharmacology of fronto-
executive function: monoaminergic modulation. Annual review of neuroscience *32*,
267-287.

Tsai, H.-C., Zhang, F., Adamantidis, A., Stuber, G.D., Bonci, A., De Lecea, L., and
Deisseroth, K. (2009). Phasic firing in dopaminergic neurons is sufficient for behavioral
conditioning. Science *324*, 1080-1084.

Hauber, W. (2010). Dopamine release in the prefrontal cortex and striatum: temporal
and behavioural aspects. Pharmacopsychiatry *43*, S32-S41.

Fujisawa, S., and Buzsáki, G. (2011). A 4 Hz oscillation adaptively synchronizes
prefrontal, VTA, and hippocampal activities. Neuron *72*, 153-165.

Lammel, S., Ion, D.I., Roeper, J., and Malenka, R.C. (2011). Projection-specific
modulation of dopamine neuron synapses by aversive and rewarding stimuli. Neuron
70, 855-862.

Vong, L., Ye, C., Yang, Z., Choi, B., Chua Jr, S., and Lowell, B.B. (2011). Leptin
action on GABAergic neurons prevents obesity and reduces inhibitory tone to POMC
neurons. Neuron *71*, 142-154.

Buschman, T.J., Denovellis, E.L., Diogo, C., Bullock, D., and Miller, E.K. (2012).
Synchronous oscillatory neural ensembles for rules in the prefrontal cortex. Neuron *76*,
838-846.

Cohen, J.Y., Haesler, S., Vong, L., Lowell, B.B., and Uchida, N. (2012). Neuron-type-
specific signals for reward and punishment in the ventral tegmental area. nature *482*, 85-
88.

Harvey, C.D., Coen, P., and Tank, D.W. (2012). Choice-specific sequences in parietal
cortex during a virtual-navigation decision task. Nature *484*, 62-68.

Salamone, J.D., and Correa, M. (2012). The mysterious motivational functions of
mesolimbic dopamine. Neuron *76*, 470-485.

Stark, E., Koos, T., and Buzsáki, G. (2012). Diode probes for spatiotemporal optical
control of multiple neurons in freely moving animals. Journal of neurophysiology *108*,
349-363.

Tan, K.R., Yvon, C., Turiault, M., Mirzabekov, J.J., Doehner, J., Labouèbe, G.,
Deisseroth, K., Tye, K.M., and Lüscher, C. (2012). GABA neurons of the VTA drive
conditioned place aversion. Neuron *73*, 1173-1183.

van Zessen, R., Phillips, J.L., Budygin, E.A., and Stuber, G.D. (2012). Activation of
VTA GABA neurons disrupts reward consumption. Neuron *73*, 1184-1194.

Howe, M.W., Tierney, P.L., Sandberg, S.G., Phillips, P.E., and Graybiel, A.M. (2013).
Prolonged dopamine signalling in striatum signals proximity and value of distant
rewards. nature *500*, 575-579.

Matsumoto, M., and Takada, M. (2013). Distinct representations of cognitive and
motivational signals in midbrain dopamine neurons. Neuron *79*, 1011-1024.

Niv, Y. (2013). Dopamine ramps up. Nature *500*, 533-535.

Berényi, A., Somogyvári, Z., Nagy, A.J., Roux, L., Long, J.D., Fujisawa, S., Stark, E.,
Leonardo, A., Harris, T.D., and Buzsáki, G. (2014). Large-scale, high-density (up to
512 channels) recording of local circuits in behaving animals. Journal of
neurophysiology *111*, 1132-1149.

Stokes, M.G. (2015). 'Activity-silent'working memory in prefrontal cortex: a dynamic
coding framework. Trends in cognitive sciences *19*, 394-405.

Hamid, A.A., Pettibone, J.R., Mabrouk, O.S., Hetrick, V.L., Schmidt, R., Vander
Weele, C.M., Kennedy, R.T., Aragona, B.J., and Berke, J.D. (2016). Mesolimbic
dopamine signals the value of work. Nature neuroscience *19*, 117-126.

Berke, J.D. (2018). What does dopamine mean? Nature neuroscience *21*, 787-793.

Miller, E.K., Lundqvist, M., and Bastos, A.M. (2018). Working Memory 2.0. Neuron
100, 463-475.

Engelhard, B., Finkelstein, J., Cox, J., Fleming, W., Jang, H.J., Ornelas, S., Koay, S.A.,

Thiberge, S.Y., Daw, N.D., and Tank, D.W. (2019). Specialized coding of sensory,

motor and cognitive variables in VTA dopamine neurons. Nature *570*, 509-513.

Mohebi, A., Pettibone, J.R., Hamid, A.A., Wong, J.-M.T., Vinson, L.T., Patriarchi, T.,

Tian, L., Kennedy, R.T., and Berke, J.D. (2019). Dissociable dopamine dynamics for

learning and motivation. Nature *570*, 65-70.

Ott, T., and Nieder, A. (2019). Dopamine and cognitive control in prefrontal cortex.

Trends in cognitive sciences *23*, 213-234.

Choi, J.Y., Jang, H.J., Ornelas, S., Fleming, W.T., Fürth, D., Au, J., Bandi, A., Engel,

E.A., and Witten, I.B. (2020). A comparison of dopaminergic and cholinergic

populations reveals unique contributions of VTA dopamine neurons to short-term

memory. Cell reports *33*, 108492.

Kim, H.R., Malik, A.N., Mikhael, J.G., Bech, P., Tsutsui-Kimura, I., Sun, F., Zhang, Y.,

Li, Y., Watabe-Uchida, M., and Gershman, S.J. (2020). A unified framework for

dopamine signals across timescales. Cell *183*, 1600-1616. e1625.

Acknowledgments

This study was supported by the Ministry of Education, Culture, Sports, Science, and Technology (Grants-in-Aid for Scientific Research 18H02711 and 18H05525), the Mitsubishi Foundation, the Naito Foundation, and the Japan Agency for Medical Research and Development (AMED).

In memory of Miles Adrian Whittington. A true mentor and friend.

Author contributions

V.G. and S.F. designed the experiments. V.G. performed the experiments and analyzed the data. V.G. and S.F. wrote the paper.

Competing Interest Statements

The authors declare that they have no competing interests.

Fig 1

Figure 1. Identifying midbrain dopaminergic and GABAergic neurons

(A) Left: We confined ChR2 expression to DA and GABA neurons by injecting locally
into the VTA the adeno-associated virus FLEX-ChR2 into transgenic mice expressing
the Cre recombinase under the control of the promoter of the DA transporter
(DAT::Cre) or the vesicular GABA transporter (VGAT::Cre). Approximately 10 days
after the virus injection, the silicon probe was inserted into the brain in the same AP and
ML coordinates. On a daily basis, the probe was inserted deeper into the brain by a few
microns. Therefore, recording sessions were performed on different DV coordinates.

Right: High-pass filtered voltage trace recorded during a light-stimulation session.
Thick blue lines indicate light pulses (450 nm, 12 ms). Two light-induced spikes are
shown below.

(B) Light response patterns of representative DA (red), GABA (blue), and unidentified
(gray) neurons. (Top) Raster plots of spikes discharged during light stimulation (colored
dots) and in the inter-stimulus baseline period (baseline, black dots). (Bottom) PSTHs
extracted from the light-induced spikes. The black dashed line indicates the upper
confidence limit of the baseline activity. If it is exceeded by the light-induced PSTH,
then the unit is identified as light-responsive (See Figure S2 for an explanation of this
term). Right inset shows, superimposed, the mean waveforms of spontaneous (black)
and light-induced (colored) spikes recorded by a single probe shank.

1000

1001

Fig 2

1002

Figure 2. Memory task, performance and population activity.

(A) Schematic representation of the T-maze apparatus configuration in the memory task. Depending on the individual features of cognitive demand, the maze was divided into sections. Every trial commenced when the animal left the starting point, running along the central arm. In the visual cue section, a visual cue signaled the rewarded location. Between the visual cue section and the turning point at the end of the main arm, a brief memory delay was introduced. After reward consumption, the animals returned of their free will to the starting point to commence a new trial.

(B) Correct performance rates in sessions with electrophysiological recordings. Gray lines illustrate performance rates for left and right trials in every session. Colored lines illustrate performance averages across sessions (mean ± standard deviation) for DAT-Cre (red) and VGAT-Cre (blue) animals.

(C) In some training sessions animals received two blocks of trials with different memory loads. Gray lines illustrate correct performance rates for each block in every session and black lines show the average performance rates (mean ± standard deviation) across sessions, for all animals.

1019   **(D)** Mean firing rates (thick lines) ± 1 standard error of the mean (shaded areas) of the

1020   population activities of DA (red) and GABA (blue) neurons. The averaged population

1021   firing activity of GABA neurons increased in the cue and delay sections. However, the

1022   averaged population activity of DA neurons did not deviate from the beginning until the

1023   end of the trials.

1024

1025

1026

1027

1028

1029

1030

1031

Fig 3

Figure 3. Trajectory-specific activities by DA and GABA neurons in the memory task

(A) Firing patterns of representative DA and GABA neurons. In each example: (Top) Raster plots of spiking events, for every correct trial, and their corresponding firing rate heatmaps as a function of position during right (purple) and left (green) trials. (Bottom) Average firing rates for correct left and right trials. Note that the trial and average firing rates (spikes/sec) are plotted as a function of position but normalized by the amount of time the mouse occupied each position on every trial. Thick lines above the average firing rates represent segments with significantly different firing rates between right and left correct trials (See also Figure S4; permutation test; $P < 0.05$). It is evident in these examples that midbrain neurons differentiate their discharge rates between left and right trajectories in certain positions.

(B) Heatmaps of neuronal population responses organized by preferred lap trajectory (i.e., the trajectory with the stronger response; first column) and non-preferred lap

trajectory (i.e., the trajectory with the weaker response; second column) for DA neurons

(Top; $n = 104$ units, 35 sessions in five mice) and GABA neurons (Bottom; $n = 74$ units,

25 sessions in four mice). Each row contains preferred and non-preferred trajectory

responses of the same neuron. In every row, both responses are normalized by the

maximum rate of the preferred trajectory. The third column shows maze segments with

significantly different discharge rates between preferred and non-preferred trajectories.

1053

Fig 4

Figure 4. VTA neuronal responses in a T-maze task without visual cues and

memory-dependent decisions (no-cue-no-choice task)

(A) Schematic representation of the T-maze apparatus illustrating the sequence of

events in the no-cue-no-choice task.

(B) Firing patterns of representative DA (left) and GABA (right) neurons during the

memory and no-cue-no-choice tasks. Both examples illustrate that the trajectory-

specific firing rate difference in the delay section of the memory task declines

prominently in the control task when animals do not receive visual cues which indicates

the reward location, or make memory-dependent decisions.

(C and D) The firing patterns of DA neurons (C; $n = 96$ units, 30 sessions in four mice)

and GABA neurons (D; $n = 32$ units, 12 sessions in three mice) in the memory task and

the no-cue-no-choice task recorded in the same sessions. (Left and Middle columns)

Normalized average neuronal responses for preferred (left) and non-preferred (middle)

trajectories. The right column represents the maze segments with significantly different

discharge rates. The row order of the neurons is the same for the memory task and the

control task heatmaps. The data shown here for the memory task are a subset of those

shown in Figure 3B.

(E) Average number of position points (mean ± standard deviation) with a significant

rate difference, arranged by maze section and behavioral task for DA (left) and GABA

(right) neurons (** $P < 0.01$, *** $P < 0.001$, paired t-test comparing numbers of

significant position points between tasks. Also, the numbers in parentheses describe the

number of neurons with a significant rate difference).

(F) Representative example showing the prominent difference in running speeds (cm/s)

between the memory (black) and no-cue-no-choice (brown) task trials in a single

session.

1080

Fig 5

1081

Figure 5. VTA neuronal responses in a T-maze task with visual cues but no memory-dependent decisions (cue-no-choice task)

(A) Schematic representation of the T-maze apparatus illustrating the sequence of events in the cue-no-choice task.

(B) Firing patterns of representative DA (left) and GABA (right) neurons during the memory and cue-no-choice tasks. Both examples illustrate that the trajectory-specific firing rate difference in the delay section of the memory task becomes notably weaker in the cue-no-choice task when animals do not make memory-dependent decisions, although running speed activities and incentive motivational drives of physical effort are the same between tasks.

(C and D) The firing patterns of DA neurons (C; $n = 31$ units, 28 sessions in four mice) and GABA neurons (D; $n = 28$ units, 11 sessions in one mouse) in the memory task and the cue-no-choice task recorded during the same sessions. (Left and Middle columns) Normalized average neuronal firing rates associated with the preferred (left) and non-preferred (middle) trajectories. The right column represents the maze segments with significantly different discharge rates. The row order of the neurons is the same for the memory task and the control task heatmaps. The data shown here for the memory task are a subset of those shown in Figure 3B.

(E) Average number of position points (mean ± standard deviation) with significant rate differences, arranged by maze section and behavioral task for (left) DA and (right) GABA neurons (** $P < 0.01$, *** $P < 0.001$, paired t-test comparing numbers of significant position points between tasks. Also, the numbers in parentheses describe the number of neurons with a significant rate difference).

Fig 6

Figure 6. Neuronal responses during reward consumption are not related to the trajectory-specific activities in the memory delay

(A) Firing patterns of representative DA and GABA neurons in the memory task for the period from trial start until the first lick of the waterspout, which triggered the water pump (top, space domain) and during reward consumption (bottom, time domain).

In each example: (Top) Raster plots of the spike trains and their corresponding firing rate heatmaps arranged by trajectory and position (maze) or time (reward) in right (purple) and left (green) trials. (Bottom) Average firing rates for correct left and right trials. The thick lines above the firing rates represent segments with significantly different firing rates between the correct right and left trials.

(B) Firing patterns of DA (top) and GABA (bottom) neurons in the time domain during reward consumption (from the first lick until 1 s later) for preferred (first column) and

non-preferred (second column) rewards (DA: $n = 104$ units, 35 sessions in five mice;

GABA: $n = 74$ units, 25 sessions in four mice). Each row represents the normalized

average firing rates (preferred and non-preferred) of a single neuron on a color scale.

Neurons were ordered according to the time point of the maximum rate in the preferred

arm. The third column shows neurons with significant discrepancies between the left

and right reward-related responses (paired t-test for mean firing rates, $P < 0.05$). The

fourth column shows post-delivery reward segments (100 ms each) with significant

excitation or inhibition compared with the 100-ms pre-reward segment (paired t-test

comparing firing rates, $P < 0.05$).

(C) Correlations between the mean firing rate difference in the reward section and the

difference in every other maze section for DA (left) and GABA (right) neurons

(Pearson's R values with P-values, *** $P < 0.001$). Only the trajectory-specific firing

rate difference in the side arms correlated with the reward-specific rate difference.

1132

1133

1134

1135

1136

1137

1138

1139

1140

1141

Supplementary Materials for

Memory-specific encoding activities of the ventral tegmental area dopamine and

GABA neurons

1146
Vasileios Glykos[1], Shigeyoshi Fujisawa[1]*

Correspondence to: shigeyoshi.fujisawa@riken.jp

This PDF file includes:

- Supplementary Text 1

- Figs 1–8

- Table 1

Supplementary Texts:

Supplementary Text 1. Evaluating the contribution of behavioral variables in

trajectory-specific activity with multiple regression analysis

We designed a generalized linear regression model in which the dependent variable was

the position-arranged firing rate, and the independent variables (predictors) were the

animal's running speed, trial number, performance rate (cumulative correct rate),

current trial accuracy (reward or not), previous trial accuracy (reward history), and lap

trajectory (left or right). Since the permutation analysis of the original spike trains

revealed that the trajectory-specific effect on neuronal firing activity was highly

dependent on the animal's position (occurring mainly in the delay and side-arm

sections), we included the joint effect of lap trajectory and position in the training model

instead of testing the effect of trajectory alone. A major advantage of regression analysis

is that one can dissociate the inherently bound effects of the independent variables. For

example, we can examine the influence of the lap trajectory variable on neuronal firing
activity without including the effect of the running speed variable (which could
potentially differ between left and right trials).

For each neuron, we produced model predictions for the correct left and right trial
average firing rates (Figure. S6A, dashed lines) from which we calculated the predicted
firing rate difference. We subsequently examined the contribution of each independent
variable to the trajectory-specific firing activity by shuffling the trajectory labels
assigned only to this particular variable. If the tested variable exerts a significant effect
on the firing rate, then shuffling the trajectory labels would produce a significant
reduction in the predicted firing rate difference. We examined the same pool of neurons
that we reported on the memory task, as shown in Figure 3B (n=104 DA and n=74
GABA-identified neurons). In the delay region, the joint effect of lap trajectory and
position (trajectory × position predictor) contributed significantly to the predicted
average rate difference of 23 DA neurons (Figure S6B and S6C top), 17 of which
overlapped with the 22 trajectory-specific neurons identified with the permutation
analysis (Figure S6C bottom). Of the GABA neurons, 34 were modulated by trajectory
and position (Figure S6B and S6C top), 30 of which were also trajectory-specific
(Figure S6C bottom). These results confirm that the trajectory factor was responsible for
the firing rate difference shown in Figure 3B. The speed variable significantly
modulated 24 DA and 22 GABA neurons (Figure S6B and S6C top), but only four DA
and four GABA neurons were co-modulated by trajectory (Figure S6C bottom). The
performance and accuracy variables modulated smaller numbers of neurons (Figure S6B
and S6C top), and the reward outcome of the previous trial did not co-modulate any of
the trajectory-specific DA and GABA neurons (Figure S6C bottom). The trial variable
modulated 30 DA neurons and 32 GABA neurons, co-modulating with the trajectory
variable of 8 DA and 15 GABA neurons. However, the distribution of the trial predictor
coefficient did not differ from a distribution with a mean equal to zero (one-sample t-
test, $P > 0.05$, Figure S6D), indicating that the effect of successive trials on firing rate
did not reflect cognitive processing, but was caused by mechanical reasons; due to the
animal's movements, the distance of the recording channel from the targeted neurons
changed continuously, which affected the signal-to-noise ratio and eventually spike

detection. In agreement with Engelhard et al., a notable proportion of DA neurons

(36%) and GABA neurons (74%) were co-modulated by more than one behavioral

variable Figure S6E (Engelhard et al., 2019).

Overall, the regression analysis confirmed the results of the permutation analysis

regarding the significant effect of trajectory on midbrain neuronal activity during

memory-dependent decisions. In addition, it demonstrated that the remaining

independent variables included in our model cannot fully explain trajectory-specific

firing activities in the delay period of the memory task.

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219 **Supplementary Figures:**

1220

Fig S1

1221

Figure S1. Validation of optogenetic identification results. Related to Figure 1.

(A) Serial reconstructions of recording sites along the rostro-caudal axis of coronal midbrain slices for DAT-Cre (red) and VGAT-Cre (blue) animals. *fr*: fasciculus retroflexus, IF: interfascicular nucleus, IP: interpeduncular nucleus, *ml*: medial lemniscus, PBP: parabrachial pigmented nucleus, PN: paranigral nucleus, RN: red nucleus, *rs*: rubrospinal tract, SNc: substantia nigra pars compacta, SNr: substantia nigra pars reticulata.

(B) Photograph of the silicon probe with attached optical fibers. Optical fibers (core diameter 50 μm) were glued to the shanks to ensure a firm, accurate, and durable insertion into the deep midbrain. Optical fibers on shanks 2 and 5 were coupled to blue laser diodes (450 nm), maintaining a certain distance from the tip of the shank to ensure that blue light could irradiate the whole span of the recording electrode array.

(C) Schematic drawing of the mouse brain in the sagittal plane, illustrating the insertion of the silicon probe into the VTA.

(D) Immunofluorescence microscopy image of a coronal brain slice of a DAT-Cre mouse injected with the optogenetic virus in the left VTA. DAT: dopamine transporter (red); YFP: yellow fluorescent protein (green); SN: substantia nigra. NOTE: the white dashed line highlights the trace by the probe shank on the tissue.

(E) Mean waveforms of light-induced (colored) and spontaneous (black) spikes discharged by identified DA (red) and GABA (blue) neurons in a single light stimulation session.

(F) Pearson's correlation coefficient between the waveforms of spontaneous and light-induced spikes for the identified DA (red) and GABA (blue) neurons. The light-induced spike waveforms were almost identical to the spontaneous waveforms for all the identified neurons. This was reflected by the high mean correlation coefficient (mean $\pm$ standard error of the mean; DA: $r = 0.98 \pm 0.09$, *P < 0.05*; GABA: $r = 0.98 \pm 0.03$, *P < 0.05*).

(G) Trough-to-peak spike duration vs. average spontaneous firing rate raster plot for the

identified DA (red) and GABA (blue) neurons. The side plots show the distributions of

the X- and Y-axis values in the main plot (pmf: probability mass function). In

agreement with earlier reports (2) DA neurons fired spikes with lower rates (DA: 4.50 ±

5.00 spikes/sec, GABA: 17.34 ± 15.72 spikes/sec; unpaired t-test comparing the

spontaneous rates: $t_{(176)} = 7.80$, $P < 0.001$) and wider waveforms (DA: 0.46 ± 0.08 ms,

GABA: 0.34 ± 0.10 ms; unpaired t-test comparing trough-to-peak values: $t_{(176)} = 5.96$, P

< 0.001) than the GABA neurons.

(H) Latency of the first spike discharged during light stimulation for the identified DA

(red) and GABA (blue) neurons.

(I) Number of light-induced spikes per light pulse for DA (red) and GABA (blue)

neurons.

1261

A

Light-stimulation protocol and Neuronal response

B

Produce jittered windows from spontaneous activity

C

Produce peri-stimulus-time histograms

Light Response
$\lambda_{light}(t)$, PSTH

Jittered window surrogates
$\lambda_{i,spont}(t)$, PSTH

Jitter #1
min: 0.01 / max: 0.30

D

Compute 95% confidence limits

E

Null hypothesis test

Fig S2

Figure S2. Statistical method for optogenetic identification. Related to Figure 1.

To identify the light-responsive units, we compared the light-induced average responses to average spontaneous firing activity. To this end, we used the following jittering method.

(A) Schematic representation of a spike train during a light stimulation session. Vertical black bars represent spiking events. The blue shaded areas indicate the light-stimulation windows. The firing activity between light stimulations is spontaneous.

(B) To identify light-responsive units we raised the hypothesis that the relative firing rate during light stimulation, $\hat{\lambda}_{light}(t)$, did not differ from the spontaneous firing rate, $\hat{\lambda}_{spont}(t)$. To do so, we employed a jittering method in which a randomly jittered window (12 ms) fitted in the spontaneous activity period preceding the light-pulse (that is, the jittered window onset never preceded the offset of the preceding light-pulse and the offset never succeeded the onset of the succeeding light pulse). We produced as many jittered windows as the light-pulses and we repeated this process 500 times in total.

(C) For every neuron we produced peri-stimulus-time-histograms (PSTHs, 12 ms, 1 ms bins) of the relative firing rate, by superimposing the spike trains of every light stimulation pulse. If $n^{(k)}(t)$ is the number of spikes of a single neuron in post-light onset time t of the k_{th} stimulation pulse, then, the average firing rate, $\hat{\lambda}(t) = \frac{1}{K} * \frac{1}{T} *$ $\sum_{k=1}^{K} n^{(k)}(t - t_0)$, where K is the number of stimulation pulses, T is the bin size (1 ms) and t_0 the light-pulse onset time, represents the relative firing rate at time t . In the end we produced one $\hat{\lambda}_{light}(t)$ and 500 $\hat{\lambda}_{spont}(t)$ firing rates.

(D) From every, $\hat{\lambda}_{i,spont}(t)$ histogram we extracted the maximum and minimum values and stored them in two separate vectors. Vectors were sorted, and from them, the 95% confidence limits were calculated. These were the 2.5% largest value from the maximum PSTH vector and the 2.5% smallest value from the minimum PSTH vector

(that is 13rd from the 500 rows). Units with $\hat{\lambda}_{light}(t)$ PSTH exceeding at any time point

(between 1 ms and 11ms) the high confidence limit, were identified as light-responsive.

(E) If the $\hat{\lambda}_{light}(t)$ PSTH fell below the low confidence limit at any time point, then the

unit was characterized as light-inhibited due to synaptic transmission (we did not find

any light-inhibited units). Otherwise, those units whose PSTH did not violate any of the

confidence limits were characterized as unidentified.

1295

Fig S3

Figure S3. T-maze configuration, behavior and running speed assessment in the memory task. Related to Figure 2.

(A) Photograph of the T-maze apparatus with maze sections labeled. The mice were trained on different variations of an associative T-maze task with visual instruction cues. The behavioural apparatus was a Figure-eight T-maze (O'Hara & CO., LTD, Tokyo, Japan), 50 cm tall, with a main arm (120 cm) and side arms (30 cm each). Infrared light-beam sensors placed at key position points on the maze defined the beginning and end of the successive maze sections. The apparatus was equipped with sliding doors, which prevented the animals from developing unwanted behavior (e.g. moving backwards). Sensor activations were on-line monitored, and significant behavioural events (e.g. visual-cue presentation, reward delivery, door opening/closing) were dictated by an in-house behavioural software written in MATLAB (Mathworks, MA, USA), through a multi-signal processor interface system (RX6; Tucker-Davis Technologies, FL, USA). The moment-by-moment position of the animal on the maze apparatus was monitored continuously by a video tracking system, composed of a red light-emitting diode mounted on the mouse head stage and a video camera (39 frames/sec) hung by the room ceiling. Video data were stored for offline analysis.

(B) Detailed schematic representation of the T-maze apparatus. Infrared light-beam sensors (s1–s6) were mounted securely at key positions on the maze to define the beginning and end of the maze sections (start: s1–s2; visual cue: s2–s3; delay: s3–s4; side-arms: s4–s5/s6; reward: s5/s6). The first activation of s5 or s6 sensors (i.e., 1st lick) triggered the water-delivery pump. Five sliding doors restrained the animals from unwanted moves. Every trial began with the animal activating s1. At this point, the doors d1, d3, and d5 remained closed, isolating the animal in the starting location for approximately 2 s. This served two purposes: (1) to drain the waterspouts and (2) to prevent the animal from making decisions on the next trial guided by the previous trial outcome. After door d1 opened, the animal could run freely along the main arm. Activating s2 and s3 resulted in the visual cue onset and offset, respectively. Activating s5 triggered the left water-delivery pump, and door d3 opened while d1 closed. Activating s6 triggered the right water-delivery pump, and door d5 opened while d1

closed. In the memory task, doors d2 and d4 were always open. In the no-cue-no-choice

task, doors d2 and d4 were initially closed. However, when the mouse activated s4, one

door opened pseudo-randomly.

(C) To assess the influence of behavioural left-right biasing in decision making we

applied the chi square (χ^2) test of independence. The null hypothesis was that correct

performance was independent of behavioural choices. Black line illustrates the chi-

squared distribution for 1 degree of freedom, along with the χ^2 values from the

independence test for every session (DAT-Cre: red dots and VGAT-Cre: blue dots). The

test was passed in every session (the level of significance was set to 5%, corresponding

to $\chi^2 = 3.84$ for 1 degree of freedom).

(D) Representative example of trial speeds (cm/sec) in a single recording session of the

memory task.

(E) Shown are running speeds (mean ± standard error of the mean) categorized by maze

sections and averaged across sessions for left (green) and right (magenta) trials for

individual VGAT-Cre and DAT-Cre mice. Although differences were small, they

occasionally reached significance. Importantly, we did not detect speed differences

between left and right trials in the delay section (unpaired t-test between left and right

speed per region, * $P < 0.05$).

1345

1346

Figure S4. Statistical method for identifying trajectory-specific neurons
(permutation method). Related to Figure 3.

We applied a permutation method reported elsewhere ([Fujisawa et al., 2008](#)) to identify

neurons with trajectory-specific encoding properties. The motivation behind this

analysis is that if the lap trajectory contributes to the firing rate difference observed at

certain positions, then shuffling the trajectory labels assigned to the spike trains of

individual trials would cause a marked reduction in the rate difference. This process is

described below.

(A) From the spike trains of the correct left and right trials (top; raster plot), we

estimated the trial relative firing rates ($\hat{\lambda}_{Left}(x)$ and $\hat{\lambda}_{Right}(x)$; middle). Importantly,

we considered the variability in occupation time between positions and trials. Hence, to

calculate the firing rates, we divided the number of spikes occurring in every position

and trial (trial PSTHs) by the occupation time in the same position and for the same trial

(trial occupation time). We subsequently produced the original average rate difference

($D_0(x)$, bottom) between the left- and right-correct trials.

(B) We shuffled the trajectory labels assigned to the trial spike trains along with the

respective occupation times and re-calculated the difference $D_i(x)$ of the permuted

labels.

(C) We produced 500 surrogates with permuted differences $D_i(x)$. In the end, we

created a 500-by-100 matrix (shuffles-by-position) of $D_i(x)$ surrogates.

(D, E) From these surrogates we extracted the 95% confidence interval (lower and

upper confidence limits) for the null hypothesis test. To do so, first, we calculated the

confidence interval of each position point x. The difference values $D_i(x)$ in every

position (column) were sorted and every row in the new matrix was treated as a

potential confidence limit with different P-value. We term it as "pointwise acceptance

band". If the original data $D_0(x)$ breaks the pointwise band, it corresponds to rejecting

the null hypothesis at position x. However, if this procedure is repeated for every

position x (i.e., 100 position points) in the maze, it raises the issue of multiple

comparisons.

(F, G) To address this, we constructed the "global band", which can control errors of

any false rejection across multiple indices. We produced another 500 permutations and

calculated the percentage of the surrogate data $D_j(x)$ that broke the pointwise band

candidates at any position points. If the percentage was more than 5%, we replaced the

pointwise band with lower P-value. We repeated this process until a pointwise band

candidate was exceeded by less than 5% of the new surrogates. When this happened, the

pointwise band was used as the "global band" (i.e., 95% confidence interval) for the

hypothesis test.

(H) The null hypothesis was rejected if the original difference $D_0(x)$, exceeded at any

position x the global band. Then, the spatial extent of significance was defined by the

number of position points exceeding the 95% confidence interval.

1387

1390 **Figure S5. Firing activities of midbrain DA and GABA neurons in the memory**

1391 **task. Related to Figure 3.**

All plots included in this figure were obtained from data recorded in the memory task.

(A and B) Representative discharge activity of DA (a) and GABA (b) neurons. In each example: (Top) Raster plots of the spikes, arranged by trial, and their corresponding firing rate heatmaps as a function of position in right (purple) and left (green) trials. (Bottom) Average firing rates for correct left and right trials. Note that the firing rate (spikes/s) is plotted as a function of position but has been normalized by the amount of time the mouse occupied each position in every trial. The thick lines above the firing rates represent segments with significantly different firing rates between the correct right and left trials.

(C and D) Heatmaps of the standardized average firing rates for the preferred (first column) and non-preferred (second column) trajectories of DA (c) and GABA (d) neurons. These plots show the number of standard deviations by which the average firing rate varied above or below the mean rate of correct trials as a function of the position on the maze. The row arrangement was the same as that in Figure 3B. The third column of the significant firing-rate difference is the same as that in the third column in Figure 3B.

(E) Heatmaps of the normalized average firing rate for preferred (first column) and non-preferred (second column) trajectories and the significant difference between them (third column) for the unidentified VTA neurons.

1412

1413 **Figure S6. Evaluating DA and GABA neuronal responses to specific behavioral**

1414 **variables in the memory task through regression analysis. Related to Figure 3.**

(A) Average firing rates of representative DA and GABA neurons extracted from real firing rate data (solid lines) and encoding model predictions (dashed lines) for left (green) and right (magenta) trials.

(B) (Left and Middle columns) Normalized firing rate heatmaps for preferred and non-preferred trajectories for DA and GABA neurons recorded in the memory task (the same as the ones in Figure 3B). (Right column) Significant contributions in the firing rate difference (yellow lines) by individual behavioral variables in the memory delay extracted with the permutation analysis of the encoding model predictions.

(C) (Top) Number of neurons significantly modulated by one of the independent variables. (Bottom) Number of neurons co-modulated by trajectory (trajectory-specific) and one of the individual variables. (Note: In B and C, the trajectory-specific neurons correspond to the significant neurons extracted using the permutation analysis of the original firing rates; third column in Figure 3B).

(D) The distribution of the trial coefficient for DA and GABA neurons along with the mean ± standard deviation values.

(E) Histogram of the number of independent behavioral variables co-modulating neurons in the delay region.

Figure S7. The dopamine signaling model of incentive motivational drive. Related to Figures 2, 3 and 4.

Striatal DA levels ramp up when mice navigate a maze or corridor in search for reward (Howe et al., 2013; Hamid et al., 2016; Kim et al., 2020). Initially, this evidence challenged theoretical neuroscientists, since the time-course of this ramping activity (lasting a few seconds) did not conform either with the prediction-error (RPE, phasic activity) or reward rate (tonic activity) theories (Niv, 2013). However, further research on this topic, provided empirical evidence to support a theory that DA conveys a signal for incentive motivational drive in the form of state-action value (value of work; (Hamid et al., 2016), or its derivative RPE; (Kim et al., 2020)).

Accordingly, the motivational value of future rewards is exponentially discounted with time or distance (the theory was developed on an adaptive decision-making framework; Hence the value of reward is defined by reward probability; figure 4 in (Hamid et al., 2016)). Different reward values correspond to different functions of discounted state-action values. If a cue predicts a reward with higher value, then the state-action value jumps to the discounted value function of that reward.

In our study, key parameters that could potentially influence incentive motivational drives were equal between left and right trials (reward amount, lap trajectory distance, etc.) and moreover, behavioral performance did not indicate choice biasing (Figures S3C). However, we cannot rule out the possibility that left and right rewards were valued differently, producing distinct value functions. If this is true, then the trajectory-

specific activities elicited by subgroups of DA and GABA neurons could reflect

differences in the state-action values for left and right trials, and not memory-dependent

decisions.

(A) The plot illustrates the hypothesis of different discounted value functions (i.e., the

state-action value is a function of position and trajectory). In the memory task and the

cue-no-choice task the assignment of the state-action value function takes place when

the visual cue signals which reward is available.

(B) However, in the no-cue-no-choice task, the assignment occurs only when the mice

are given access to one of the side arms.

1467

1468

1469

1470

1471

1472

1473

1474

Figure S8. Running speed differences between the memory task and the cue-no-choice task. Related to Figures 3 and 5

1475

1476

(A) Representative example of running speeds (cm/s) for memory (black) and cue-no-choice (orange) trials in a single session. Speed differences were evident at the T-intersection (between delay and arms).

1477

1478

1479

(B) Average running speed values (mean ± standard error of the mean) grouped by maze section and behavioral task for each of the DAT-Cre and VGAT-Cre animals performing the memory and cue-no-choice tasks. Importantly, no differences were detected in the delay and side-arm sections between memory (black) and cue-no-choice (brown) task trials (* $P < 0.05$, unpaired t-test on speed).

1480

1481

1482

1483

1484

1485

1486

1487

1488

1489

1490 **Supplementary Table:**

1491

1492 **Table S1. Number of neurons with trajectory-specific firing activities in the**

1493 **memory task grouped by maze section**

1494 Data are presented for all recorded neurons and individually for optogenetically

1495 identified DA and GABA neurons; DA, dopamine; GABA, gamma-aminobutyric acid.

Memory task: Significant difference in firing rate between left and right trials					
	start	cue	delay	side arms	reward
All (n = 1,191)	n = 49 (4%)	n = 78 (7%)	n = 377 (32%)	n = 505 (42%)	n = 461 (39%)
DA (n = 104)	n = 4 (4%)	n = 3 (3%)	n = 22 (21%)	n = 23 (22%)	n = 24 (23%)
GABA (n = 74)	n = 4 (5%)	n = 11 (15%)	n = 35 (47%)	n = 35 (47%)	n = 34 (46%)

1496

1497

1498 **SI References**

1499 *All references have been already cited in the main manuscript*

1500

1501

1502