A cellular and spatial atlas of *TP53*-associated tissue remodeling in lung adenocarcinoma

Authors

William Zhao[1,2], Benjamin Kepecs[1,2], Navin R. Mahadevan[6,8,10], Asa Segerstolpe[4], Jason L.

Weirather[8], Naomi R. Besson[9], Bruno Giotti[1,2], Brian Y. Soong[1,2], Chendi Li[12,13], Sebastien

Vigneau[7,14], Michal Slyper[4,15], Isaac Wakiro[7,16], Judit Jane-Valbuena[4,15], Orr Ashenberg[4], Asaf

Rotem[7,17], Raphael Bueno[11], Orit Rozenblatt-Rosen[4,15], Kathleen Pfaff[9], Scott Rodig[10], Aaron N.

Hata[12,13], Aviv Regev[4,5,15], Bruce E. Johnson[7], & Alexander M. Tsankov[1,2,3,4,18*]

Affiliations

[1]Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai

(ISMMS), New York, NY, USA

[2]The Precision Immunology Institute, ISMMS, New York, NY, USA

[3]The Tisch Cancer Institute, ISMMS, New York, NY, USA

[4]Klarman Cell Observatory, Broad Institute of Massachusetts Institute of Technology (MIT) and

Harvard, Cambridge, MA, USA

[5]Department of Biology, MIT, Cambridge, MA, USA

[6]Department of Medical Oncology, Dana-Farber Cancer Institute (DFCI), Boston, MA, USA

[7]Center for Cancer Genomics, Department of Medical Oncology, DFCI, Boston, MA, USA

[8]Department of Data Science, DFCI, Boston, MA, USA

[9]Center for Immuno-Oncology, DFCI, Boston, MA, USA

[10]Department of Pathology, Brigham and Women's Hospital (BWH), Boston, MA, USA

[11]Department of Thoracic Surgery, BWH, Boston, MA, USA

[12]Massachusetts General Hospital Cancer Center, Charlestown, MA, USA

[13]Department of Medicine, Harvard Medical School, Boston, MA, USA

[14]Present address: Ginkgo Bioworks, 27 Drydock Avenue, Boston, MA, USA

[15]Present address: Genentech, 1 DNA Way, South San Francisco, CA, USA

[16]Present address: Louis V. Gerstner Jr. Graduate School of Biomedical Sciences at Memorial

Sloan Kettering Cancer Center, New York, NY, USA

[17]Present address: Astrazeneca, 35 Gatehouse Dr., Waltham, MA, USA

[18]Lead contact

*Correspondence: alexander.tsankov@mssm.edu (A.M.T.)

<u>ABSTRACT</u>

TP53 is the most frequently mutated gene across many cancers and is associated with shorter

survival in lung adenocarcinoma (LUAD). To define how *TP53* mutations affect the LUAD tumor

microenvironment (TME), we constructed a multi-omic cellular and spatial tumor atlas of 23

treatment-naïve human lung tumors. We found that *TP53*-mutant (*TP53*^{mut}) malignant cells lose

alveolar identity and upregulate highly proliferative and entropic gene expression programs

consistently across resectable LUAD patient tumors, genetically engineered mouse models, and

cell lines harboring a wide spectrum of *TP53* mutations. We further identified a multicellular tumor

niche composed of *SPP1*$^{+}$ macrophages and collagen-expressing fibroblasts that coincides with

hypoxic, pro-metastatic expression programs in *TP53*^{mut} tumors. Spatially correlated angiostatic

and immune checkpoint interactions, including *CD274-PDCD1* and *PVR-TIGIT*, are also enriched

in *TP53*^{mut} LUAD tumors, which may influence response to checkpoint blockade therapy. Our

methodology can be further applied to investigate mutation-specific TME changes in other

cancers.

INTRODUCTION

Non-small-cell lung cancer (NSCLC) is a heterogeneous disease that has routinely been classified into different histological subtypes[1]. More recently, genomic alterations have been used to characterize molecular subtypes of NSCLC that can be used to select treatments for patients with specific genomic changes[2-4]. Tyrosine kinase inhibitors (TKIs) and other targeted therapies against oncogenic mutations and chromosomal rearrangements (*e.g.*, *EGFR*, *KRAS-G12C, ALK*) have led to a recent improvement in survival in NSCLC patients[5]. Treatments with immune checkpoint inhibitors (ICIs) (*e.g.*, anti-PD-1/PD-L1) with or without chemotherapy have proven efficacious in a subset of patients[6] and are currently part of the first-line treatment in some patients with stage IV NSCLC[7]. However, due to high rates of cancer recurrence, more effective approaches toward patient stratification and precision therapy are urgently needed.

Tumor protein p53 (*TP53*) is the most commonly mutated gene in NSCLC (~50%) as well as across many other cancers[8]. *TP53* mutations are more common in smokers[9], and are associated with tumor progression and metastasis, leading to shorter survival in patients with NSCLC[10,11,12]. Despite being the most extensively studied tumor suppressor, *TP53* has been challenging to target therapeutically due to the diversity of mutations reported and its involvement in a wide range of regulatory pathways[13-15]. *TP53* is known to have an impact on the tumor microenvironment (TME) during tumor development in mouse models[16-18]. Mutant *TP53* is also associated with increased expression of PD-L1 and improved response to anti-PD1 therapy in human lung adenocarcinoma (LUAD)[19-24], motivating us to investigate TME differences in LUAD patients with *TP53*-mutant (*TP53*^{mut}) *vs. TP53*-wild type (*TP53*^{WT}) tumors.

Single cell and spatial profiling of multiple treatment-naïve, resectable NSCLC patient tumors

could advance our fundamental understanding of *TP53*-mutant cancer cells and their relationship

to the TME and provide insights that may yield new therapeutic opportunities but, prior studies

have not yet addressed this question. Single-cell RNA-sequencing (scRNA-seq) has previously

been used to characterize malignant cells and the TME in resectable NSCLCs, focusing on gene

expression differences between the primary tumor and adjacent normal tissue[25-28], during tumor

progression[29,30], or across metastatic sites[31,32]. Studies have also leveraged TME characteristics to

stratify NSCLC patients into clinically-relevant subgroups that could predict response to

immunotherapy[22,33]. One study focused on characterizing the TME and intra-tumoral

heterogeneity in *EGFR*[mut] LUAD[34], but these previous scRNA-seq studies did not determine the

mutational status of *TP53* in tumors and did not investigate its impact on all cellular compartments

in the TME.

Here, we generated the most comprehensive, multi-omic atlas of NSCLC to date, combining

matched whole-exome sequencing, scRNA-seq, spatial transcriptomics, and multiplex

immunofluorescence from *TP53*[mut] and *TP53*[WT] human tumors, to identify *TP53* mutation-

associated changes in malignant cells and the TME. Our integrated atlas allowed us to identify

spatially validated, cell-cell interactions that may contribute to shaping the tumor architecture of

NSCLC. In doing so, we uncovered a hypoxic expression niche in *TP53*[mut] LUAD and identified

possible mechanisms by which malignant tumor cells may limit the growth of blood vessels in the

TME. We further found a T cell compartment with highly exhausted features in *TP53*[mut] *vs.*

TP53[WT] LUAD, which was accompanied by an enrichment in multiple immune checkpoint

interactions between malignant, T, and myeloid cells. Finally, using a systematic approach to

detect cellular landmarks in spatial transcriptomics data, we observed a potential pro-metastatic

tissue niche consisting of *SPP1*-expressing tumor associated macrophages (TAMs) and a collagen-

expressing population of cancer-associated fibroblasts (CAFs) linked to hypoxia in the *TP53*[mut]

LUAD TME. Overall, our approach provides a framework for leveraging multi-omic technologies

to understand the impact of specific genomic alterations[35,36] on the cellular and spatial architecture

that can enable discovery of new biomarkers and more optimal methods for patient stratification

in lung cancer and other tumor types.

RESULTS

A multi-omic atlas of NSCLC

To characterize the molecular, cellular, and spatial heterogeneity in NSCLC, we performed whole-

exome sequencing (WES), scRNA-seq, spatial transcriptomics (ST; 10X Visium), and multiplex

immunofluorescence (mIF) on primary tumor resections from 23 NSCLC patients (**Figure 1A**), of

which 18 were diagnosed as LUAD (**Table S1**). Twenty-one of the 23 patients had a history of

cigarette smoking (Median 40, Range 12-100 pack years) and all patients were confirmed to be

treatment-naïve prior to tumor resection. We used WES analysis to determine the total tumor

mutational burden, copy-number alterations (CNAs), and mutational status of genes commonly

altered in NSCLC, including *TP53*, *EGFR*, *KRAS*, *STK11, KEAP1*, *RBM10*, and *PTPRD* in each

tumor. There was high concordance between predicted CNA profiles from tumor-matched WES

and those inferred from scRNA-seq data (**Figure 1B** and **Figure S1A**), many of which overlapped

with commonly deleted (*e.g.*, *CDKN2A*, *PTPRD*, *B2M*) and amplified (*e.g.*, *NKX2-1*, *TERT*,

MCL1, *EGFR*) genes in LUAD[8]. After filtering, we obtained a total of 167,193 high-quality cell

profiles and used CNA inference to distinguish between malignant and non-malignant epithelial

cells and then integrated[37], clustered, and annotated them into 11 broad cell classes based on

coherent expression of known marker genes (**Figure 1C**,**D**; **Figure S1B**,**C**; **Table S2**).

Distinguishing tumors by their *TP53* mutation status, the different *TP53* mutations had comparable

effects in downregulating the average expression of *TP53* target genes in malignant cells relative

to *TP53*^{WT} LUAD tumors (**Figure 1E**). The remaining five tumors consisted of squamous,

mucinous, and colloid histology, and were not used for comparisons between *TP53*^{mut} and *TP53*^{WT}

tumors because *TP53* mutations have been found to have histology-dependent effects on patient

outcome[38].

***TP53*^{mut} LUAD is associated with a distinct TME composition**

Cell subsets in the TME of *TP53*^{mut} *vs*. *TP53*^{WT} LUAD tumors had distinguishable cell-intrinsic

expression profiles (**Figure 1D**, **right**) and cell compositions (**Figure 1F** and **Figure S1D**). In

particular, there was a pronounced decrease in the proportion of endothelial cells and pericytes in

TP53^{mut} *vs*. *TP53*^{WT} LUAD tumors (**Figure 1F**, **left**), which we confirmed by scoring the

corresponding cell type signatures in 510 bulk RNA-seq profiles of primary LUAD tumors from

The Cancer Genome Atlas (TCGA) (**Figure 1F**, **right**). Endothelial cell abnormalities have been

linked to tumor cell progression and metastasis[39], and pericyte depletion has been linked to

hypoxia-associated epithelial to mesenchymal transition (EMT) in breast cancer[40].

These compositional differences between in *TP53*^{mut} and *TP53*^{WT} LUAD were also supported by

spatial transcriptomic profiling of 14 tissue sections from 6 tumors in our cohort (2 with *TP53*^{mut}

and 4 with *TP53*^{WT} LUAD). Spatial data from 2 additional tumors (6 slides) were excluded from

downstream analysis due to either having an non-LUAD histology or an insufficient number of

malignant cells (**Figure 1G,H** and **Table S1**). We used Tangram[41], a deep learning framework, to

map tumor-matched scRNA-seq profiles to the corresponding ST data and generate a probabilistic

measure of the cellular composition for each spatial measurement (**Figure 1G**). Spatial cell

composition inference was highly concordant with independent, manual annotations of the

corresponding hematoxylin and eosin (H&E) stains by a pulmonary pathologist (N.R.M.; **Figure**

1G and **Figure S1E**-**G**). Consistent with our scRNA-seq, there was also a decrease in the

proportion of endothelial cells and pericytes across ST spots in $TP53^{mut}$ *vs*. $TP53^{WT}$ tumor samples

(**Figure 1H**).

Dichotomous activity of malignant cell programs in LUAD associates with survival

While malignant cells from one tumor can occupy a gradient of expression states rather than

discrete cell subsets[42,43], different cancer expression programs and their usage across cells and

tumors have not been comprehensively defined in LUAD at single-cell resolution. To identify

shared cell states across tumors despite predominant, patient-specific differences in malignant

cells[42,43], we used canonical correlation analysis (CCA)[44] to integrate and cluster malignant cells

into 18 subsets shared across tumors (**Figure 2A** and **Figure S2A**) and annotated cell subsets using

gene set enrichment analysis (**Figure S2B**), highlighting representative markers (**Figure 2B**).

These subsets were enriched for genes associated with hallmark cancer processes (*e.g.*, cell cycle,

hypoxia[45-47], glycolysis[48], partial EMT (pEMT)[49], interferon gamma (IFNG) and TNFA signaling)

and with lung epithelial cell identity (*e.g.*, alveolar type 2 (AT2-like), ciliated, secretory). Other

malignant cell subsets showed marked expression of antigen presentation (MHCII), stress

response[50], metallothionein, and senescence-related genes[51], which have not been previously

identified *de novo* using scRNA-seq in NSCLC primary tumors. For downstream analyses, we

defined LUAD malignant programs by the set of 10 most differentially expressed genes for each

malignant subset and scored each program's gene set across all malignant cells.

The programs clustered into two main groups after correlating their scores across all malignant

cells (**Figure 2C**) and tumor samples (**Figure S2C**): one consisting of cell cycle, hypoxia, pEMT,

and glycolysis programs, and another of alveolar-like, antigen-presentation, and stress response

programs. Malignant program scores in these two groups were strongly anticorrelated, suggesting

that they were regulated by two distinct and opposing pathways (**Figure 2C** and **Figure S2C**).

This is consistent with a previous LUAD study where cells from patient-derived xenografts

consisted of two distinct intra-tumoral subgroups, distinguished by a gene module consisting

primarily of cell cycle genes[52]. High expression of several malignant programs (*e.g.*, pEMT,

hypoxia, cell-cycle, metallothionein, glycolysis) was associated with shorter survival across LUAD

patient tumors in TCGA (**Figure 2D** and **Figure S2D**). In contrast, high expression of the antigen-

presentation, MHCII program was linked to prolonged survival. Previously, various signatures

derived from bulk RNA profiling in TCGA have been shown to be predictive of outcome in

NSCLC[53,54] but our *de novo* approach allowed for the discovery of more granular programs based

on malignant cell expression only, unconfounded by other cells in the tumor microenvironment.

To identify cell-intrinsic differences between malignant cells from *TP53*[mut] and *TP53*[WT] LUAD,

we compared the mean activity score of different malignant programs between the two patient

groups and found higher expression in *TP53*[mut] *vs*. *TP53*[WT] LUAD for several programs associated

with shorter survival (*e.g.*, cell cycle, glycolysis, pEMT), and a decreased expression of the AT2-

like program and *TP53* targets (**Figure 2E**, **left**), consistent with the observation that *TP53*[mut]

tumors have shorter survival in LUAD TCGA (**Figure S2E**). There was not a statistically

significant difference in either the number of single-nucleotide alterations (SNAs) or CNAs

between *TP53*[mut] and *TP53*[WT] tumors in our cohort (**Figure S2F**). The malignant program

differences were corroborated in bulk RNA-seq data from 263 *TP53*[mut] and 247 *TP53*[WT] LUAD

primary tumors in TCGA (**Figure 2E**, **right**). These findings were also replicated when analyzing

a compendium of independent cohorts (**Table S3**), assembled from published and unpublished

scRNA-seq data from 22 additional treatment-naïve LUAD primary tumor samples with annotated

mutational status (7 *TP53*[mut] and 15 *TP53*[WT])[26,32,55] (**Figure S2G**). Notably, spatial co-expression

of cell cycle with glycolysis and of hypoxia with pEMT programs was higher in *TP53*[mut] *vs*.

TP53[WT] LUAD tumors (**Figure 2F**).

The decreased expression of the AT2-like program in *TP53*[mut] LUAD was consistent with a loss

of AT2 cell identity. Scoring each malignant cell profile across tumors for normal lung epithelial

cell signatures[56], *TP53*[WT] LUAD malignant cells scored highly for AT2 markers (**Figure 2G**),

whereas cells from tumors with squamous, neuroendocrine, or mucinous/colloid subtypes

expressed the expected corresponding basal, neuroendocrine, or goblet signatures[57]. Interestingly,

AT2 scores for *TP53*[mut] LUAD malignant cells were significantly lower than *TP53*[WT] LUAD but

were comparable to AT2 scores in the tumors from non-adenocarcinoma subtypes, suggesting a

loss of alveolar cell identity, which associates with tumor progression in LUAD[58].

Consistent cancer-intrinsic changes observed across different *TP53* variant and co-mutation

classes

To understand the potential joint effect of *EGFR* and *KRAS* co-mutations on the observed *TP53*[mut]

LUAD cancer-intrinsic changes, we partitioned the TCGA cohort (n=506) into 6 subsets with

distinct mutational statuses for these three genes (**Figure S3A**, **left**). *TP53*[mut] tumors with or

without co-mutations in *EGFR* or *KRAS* showed significant decreases in *TP53* target and AT2-

like scores, and a significant increase in CC.G2/M score compared to *TP53*[WT]*KRAS*[WT]*EGFR*[WT]

tumors, whereas *TP53*[WT] tumors with either *EGFR* or *KRAS* mutations did not have such

significant differences in these signatures compared to tumors that were *TP53*[WT]*KRAS*[WT]*EGFR*[WT].

Interestingly, tumors with *TP53* mutations alone or in combination with *KRAS* but not with *EGFR*

had   significant   increases   in   Glycolysis.hypoxia   and   pEMT   scores   compared   to

TP53[WT]*KRAS*[WT]*EGFR*[WT] tumors, suggesting that *EGFR* mutations counteracted the *TP53*-

induced effects on these malignant expression programs (**Figure S3A**), consistent with a recent

report[59]. Multiple regression analysis also showed consistent cancer-intrinsic changes associated

with *TP53* mutational status even when regressing out the effects of *EGFR*, *KRAS* (*G12C, G12D,*

G12V, as well as all *KRAS* mutations together)*, STK11*, *KEAP1*, *RBM10*, and *PTPRD* mutations

(**Figure 2H** and **Figure S3B**). Notably, there were no significant interaction effects between

mutational status of *TP53* and *KRAS* or *TP53* and *EGFR* after ANOVA testing of model outputs.

To investigate the potential impact of different *TP53* mutational variants on changes to the

expression of malignant gene programs, we partitioned the TCGA tumor samples based on

dominant negative (DNE), loss of function (LOF), and Impactful *TP53* mutational classes as

defined by previous functional studies overexpressing different *TP53*[mut] variants in a *TP53*[WT]

A549 LUAD cell line[60,61]. Consistent with our tumor scRNA-seq analysis, decreases in *TP53*

target, and AT2-like scores, and increases in CC.G2/M, Glycolysis.hypoxia, and pEMT scores

were associated with TCGA LUAD tumors with different sub-categories of *TP53* DNE and LOF

mutations (**Figure S3C**) or of *TP53* Impactful mutation sub-categories (**Figure S3D**). The

DNE,LOF and Impactful II mutations gave rise to the most pronounced differences, where

Glycolysis.hypoxia and pEMT programs are significantly increased only in DNE,LOF and

Impactful II variant categories in TCGA (**Figure S3C** and **Figure S3D**). Moreover, many of these

malignant cell intrinsic changes were also observed *in vitro* in A549 LUAD cells (*TP53*[WT])

overexpressing each of a broad spectrum of *TP53* mutational variants and profiled by Perturb-

seq[61] (**Figure S3E**). These trends held across all mutational categories, including all four *TP53*

variants also found in our patients' tumors and one variant also found in the scRNA-seq validation

compendium (**Figure S3E**). As in TCGA, the malignant expression program changes were most

pronounced for Impactful II mutations (**Figure S3E**) but were absent from A549 LUAD cells

overexpressing different *KRAS* mutations (**Figure S3F**). Because we see consistent changes in

malignant expression programs across a wide spectrum of *TP53* mutations, we reason that these

phenotypic changes are associated with a loss of *TP53*[WT] activity, rather than specific gain of

function that arises from any specific mutation. In support of our observations, heterogeneous

deletion events resulting in loss of *TP53* activity has recently been shown to result in homogenous

and deterministic patterns genome evolution[62].

Finally, we observed consistent changes in expression associated with *TP53* mutations at the

protein level. Analyzing bulk proteomic data from the Clinical Proteomic Tumor Analysis

Consortium (CPTAC) showed consistent decreases in *TP53* target and AT2-like signature scores,

and an increase in CC.G2M signature scores at the protein level (**Figure S3G**), where effects were

again most pronounced for DNE,LOF and Impactful II mutation categories (**Figure S3H**).

Conserved changes in signaling entropy and malignant programs in *TP53*[mut] human tumors

and mouse models

We have previously reported loss of alveolar cell identity and reversion to a progenitor-like, highly-plastic cell state (HPCS) during LUAD progression in mice with somatic activation of oncogenic **K**RAS-G12D, which was greater in tumors from mice where the **p**53 was deleted (KP model) than those with wild-type p53 (K model)[63]. Consistent with our findings in human tumors, there was a significant decrease in AT2-like program score, and significant increases in CC.S, Hypoxia, Glycolysis.hypoxia, pEMT program scores when comparing KP versus K mice malignant cells (**Figure S4A**). Moreover, the expression of key hypoxia transcription factor *Hif1a* was higher in KP *vs.* K malignant cells (**Figure S4A**, **right**). Interestingly, we did not observe significant increases in the Hypoxia, Glycolysis.hypoxia, or pEMT programs between *TP53*[mut] and *TP53*[WT] A549 LUAD cells *in vitro* (**Figure S3E**), suggesting that the tumor microenvironment is important for shaping these malignant programs *in vivo*.

The consistent loss in alveolar type II cell identity in *TP53*[mut] LUAD human tumors, cell lines, and mouse models prompted us to explore the consequences of this loss on signaling entropy[64], a representative measure of cellular plasticity. We found an increase in signaling entropy in *TP53*[mut] *vs. TP53*[WT] LUAD malignant cells at the single-cell level in our cohort, which we confirmed in bulk RNA-seq and bulk proteomic profiles across a larger cohort size (**Figure 2I**), as well as in the KP *vs.* K mouse models[63] (**Figure 2J**). This increase in entropy was most significant in DNE,LOF and Impactful II *TP53*[mut] tumors in TCGA (**Figure S4B**) and was significant in *TP53*-

only mutants and *TP53-KRAS* co-mutants but not in *TP53-EGFR* co-mutants in TCGA (**Figure**

S4C), consistent with the trends observed in the malignant program activity changes.

Cell cycle and pEMT programs displayed the highest entropy amongst human malignant subsets

(**Figure S4D**) and the greatest similarity to the HPCS cell state described in the KP mouse model

(**Figure S4E**). Moreover, in A549 cells overexpressing mutant *TP53*, there was significantly

increased entropy in cells harboring Impactful I and Impactful II *TP53* mutations, including

variants found in our scRNA-seq and validation cohorts (**Figure 2K**). Conversely, there was no

significant change in entropy in the same cells overexpressing different classes of *KRAS* mutants

(**Figure S4F**), suggesting that the observed increase in entropy is directly linked to *TP53*

mutations. Taken together*,* malignant cells from *TP53*^WT LUAD tumors were enriched for antigen

presentation and AT2-like cell programs, while those from *TP53*^mut LUAD tumors were highly

entropic, plastic, and associated with programs predictive of poor outcome, including cell cycle,

glycolysis, hypoxia, and pEMT.

***TP53*^mut LUAD is characterized by spatial interactions between malignant and stromal cells**

known to inhibit vascularization and promote EMT

We next focused on the decreased proportion of endothelial cells in *TP53*^mut LUAD (**Figure 1F**).

This association is also observed in *TP53* mutant tumors in TCGA LUAD (**Figure 1F**), with or

without co-mutations in *KRAS* and *EGFR* (**Figure S5A**, **left**), and is most significant in DNE,LOF

TP53 mutants (**Figure S5A**, **right**).

To investigate the potential mechanisms for decreased vascularization, we annotated 9 endothelial

cell subsets based on the expression of established markers for endothelial cell subsets in the

normal lung (**Figure 3A**, **Figure S5B-D**)[65]. These included: aerocytes, a specialized capillary cell

responsible for gas exchange in the alveolus[66] (expressing *HPGD*, *EDNRB*, and *SOSTDC1*);

arterial cells (*GJA5*, *ENPP2*); capillary cells (*CA4*, *FCN3*); vascular endothelial cells (VEC),

including *COL4A1*[+] VECs (VEC.COL4A1), cycling (VEC.Cycling) and interferon-stimulated

(VEC.IFN) subsets; lymphatic cells; pulmonary venous cells; and systemic venous cells. The

proportion of aerocytes — and to a lesser degree arterial cells — out of all endothelial cells was

decreased in *TP53*[mut] *vs*. *TP53*[WT] LUAD (**Figure 3B**), which was further supported by bulk RNA-

seq data deconvolved using scRNA-seq-derived markers (**Figure S5E**). This suggests a

corresponding decrease in gas exchange in *TP53*[mut] LUAD and a resulting decrease in oxygen

diffusion to the area directly surrounding the alveoli at the tumor site, providing a selective

advantage for tumor cell survival[67].

Analyzing ligand-receptor pairs, several interactions between malignant and endothelial cells

known to inhibit endothelial cell growth and function were enriched in *TP53*[mut] LUAD, including

SEMA3A-NRP1 and *EPHB2-EFNB1*[68-70] (**Figure 3C** and **Figure S5F)**. Spatial transcriptomics

from matched tumor samples support these observations, showing a significantly higher spatial

correlation in the expression of *SEMA3A-NRP1* and *EPHB2-EFNB1* ligand-receptor pairs in

TP53[mut] *vs*. *TP53*[WT] LUAD sections (**Figure 3D** and **Figure S5G**). Taken together, we observe

spatially correlated expression of known inhibitory interactions enriched between malignant and

endothelial cells in *TP53*[mut] LUAD, which may explain the accompanying decrease in these

subsets and vascularization.

The heterogeneity of mesenchymal cells and their potential role in faster tumor progression and shorter patient survival has been documented in NSCLC[26,71]. Mesenchymal cells in our tumors were partitioned into 13 subsets (**Figure 3E** and **Figure S6A-C**). Two subsets most closely resembled normal lung myofibroblasts[56] (**Figure 3F**): CAF.COLs expressing high levels of collagens (*COL1A2*, *COL3A1*, *COL1A1*), which share markers with recently described *LRRC15*[+] myofibroblasts[72], and myofibroblasts, expressing both fibroblast and smooth muscle cell markers. We also annotated two pericyte-like subsets, one expressing canonical pericyte markers (*HIGD1B*, *COX4I2*), and one expressing *COL4A2* and EMT-related genes (**Figure S6A,B**); airway and vascular smooth muscle subsets, and alveolar (CAF.ADH1B) and adventitial (CAF.Adventitial) fibroblasts, which correspond to recently described human lung alveolar and adventitial fibroblasts[72]; and fibroblast subsets expressing higher levels of lipid metabolism (CAF.APOE), ribosomal (CAF.Ribo), complement and chemotaxis (CAF.Complement), and interferon-stimulated (CAF.ISGs) genes. Of note, we did not observe CAF subsets that resembled normal lung fibromyocytes or lipofibroblasts (**Figure 3F**). The different mesenchymal subsets were associated with different active regulons (**Figure 3G**), as inferred by pySCENIC[73], with high regulon specificity (**Figure S6D**). As expected, cycling CAFs had an active *E2F1* regulon, a cell cycle regulator; CAF.ISGs activated the *IRF7* and *STAT1* regulons, controlling interferon responsive genes; and CAF.COLs activated a *TWIST1* regulon, previously linked to EMT in cancer[74].

Based on the inferred composition of CAF subsets in spatial transcriptomics spots across sections, pericytes were significantly spatially correlated with endothelial cells (**Figure S6E**, Pearson

correlation *P*-value < 10^{-15}); CAF.COLs with vascular smooth muscle and CAF.ADH1B,

CAF.Complement with CAF.Adventitial, pericytes with CAF.ISGs, and airway smooth muscle

cells with CAF.APOE and CAF.Ribo. Spatially, most mesenchymal subsets were negatively

correlated with malignant cells, indicating occlusion of fibroblasts from the tumor core (**Figure**

S6E). Comparing the relative proportions of mesenchymal cell subsets (out of all mesenchymal

cells) between *TP53*^mut *vs*. *TP53*^WT LUAD, pericytes and CAF.ADH1B were decreased in *TP53*^mut

LUAD (as observed for broad cell classes) (**Figure 3H**), also supported by deconvolved bulk

RNA-seq (**Figure 1F** and **Figure S6F**). Similar to endothelial scores, pericyte scores were

significantly decreased in *TP53* mutant tumors in TCGA LUAD, with or without co-mutations in

KRAS and *EGFR*, and this decrease was most significant in the DNE,LOF class of *TP53* mutations

in TCGA (**Figure S6G**).

TGFB2-TGFBR2 had significantly higher spatial correlation in *TP53*^mut *vs*. *TP53*^WT LUAD

(**Figure 3I**), and *TGFB2-TGFBR2* and *TGFBR2-TGFB3* interactions between mesenchymal and

malignant cells were enriched in *TP53*^mut *vs*. *TP53*^WT LUAD (**Figure S6H**), although the

enrichment was not significant. *VEGFA* expression in myofibroblasts is higher in *TP53*^mut *vs*.

TP53^WT LUAD (**Figure S6I**), consistent with previous reports that myofibroblasts can regulate

angiogenesis in response to hypoxia[75]. Overall, changes in both stromal cell composition and

spatial organization in *TP53*^mut LUAD are accompanied by an enrichment of putative interactions

known to inhibit vasculature and induce epithelial-mesenchymal transition.

Increased *SPP1* expression and immunomodulatory interactions in *TP53*^mut LUAD myeloid

cells

Myeloid cells play an important role in tumor progression and may be a compelling therapeutic

target in cancer[76]. While NSCLC myeloid cell populations were generally characterized[22,27], the

impact of *TP53* mutations on myeloid cells, and their interactions with malignant, stromal, and

immune compartments in the TME has not been studied in detail in humans.

Myeloid cells partitioned into 14 subsets, largely consistent with previous NSCLC atlases[22]

(**Figure 4A,B** and **Figure S7A**), but with significant compositional differences between *TP53*^{mut}

and *TP53*^{WT} LUAD. These include a significant decrease in *TP53*^{mut} LUAD in the proportion of

TAM.FABP4 cells, which resemble tissue-resident alveolar macrophages (TAMs), and an increase

in *SPP1*-expressing TAMs (TAM.SPP1) (**Figure 4C**). In agreement, *SPP1* expression was

significantly increased both in TCGA bulk RNA-seq data and more prominently in myeloid

pseudobulk comparisons between *TP53*^{mut} and *TP53*^{WT} LUAD tumors (**Figure 4D**, **left**).

Expression of *CXCL9/10/11* was also increased in *TP53*^{mut} bulk TCGA LUAD tumors (**Figure

S7B**). The proportion of *CXCL9/10/11* expressing TAMs (TAM.CXCLs) was also higher in

TP53^{mut} LUAD tumors both in the scRNA-seq data and in deconvolution analysis of TCGA bulk

RNA-seq data (**Figure S7C**). Notably, increases in *SPP1* and *CXCL9/10/11* were most significant

in *TP53* mutant tumors without co-mutations in *EGFR* or *KRAS*, and in DNE,LOF and Impactful

II classes of *TP53* mutant tumors in TCGA (**Figure S7D,E**).

Myeloid cell subsets in *TP53*^{mut} and *TP53*^{WT} LUAD were also associated with distinct cell-

intrinsic gene programs, such that genes differentially expressed between myeloid subsets from

TP53^{mut} *vs*. *TP53*^{WT} tumors were enriched for EMT, hypoxia, and interferon response pathway,

especially in TAM.SPP1 cells (**Figure 4E** and **Figure S7F**). *SPP1* expression has been previously

associated with EMT, and with early lymph node metastasis in LUAD[77], as well as with shorter

survival in TCGA LUAD independent of *TP53* mutational status (**Figure S7G**). Notably, the genes

most correlated with *SPP1* expression across single myeloid cells were enriched for EMT, hypoxia

and glycolysis functions (**Figure 4D**, **right**) and included VEGF receptors *NRP1* and *FLT1*

(**Figure S7H**). In support of this observation, a recent compendium of 399 published scRNA-seq

studies has shown that *SPP1*[+] macrophages are consistently found in fibrotic lung diseases,

COVID-19 lung, and pancreatic ductal adenocarcinoma (PDAC), suggesting a role of these

fibrosis-associated *SPP1*[+] macrophages in lung injury and ECM remodeling[78]. In summary, we

find increased expression of *SPP1* in *TP53*[mut] LUAD myeloid cells that is tightly linked with

induction of genes involved in EMT and hypoxia programs.

To identify ligand-receptor interactions that may contribute to a distinct myeloid compartment in

TP53[mut] LUAD, we conducted differential ligand-receptor interaction analysis. Putative

interactions enriched in *TP53*[mut] LUAD include *VEGFA-FLT1* and *NRP1-SEMA3A* (**Figure 4F**),

involved in recruitment of myeloid cells to tumor sites[79], and *TGFB2-TGFBR1* interactions

between malignant and myeloid cells, which are known to promote monocyte differentiation into

TAMs[80], and may help explain the relative increase in the monocyte-macrophage population in

TP53[mut] LUAD. Additionally, *NRP1-VEGFA*, *NRP1-VEGFB*, *FLT1-VEGFB*, and *FLT1-PGF*

interactions between myeloid and mesenchymal cells, induced in *TP53*[mut] LUAD, suggest that

mesenchymal cells may contribute to monocyte recruitment to the TME (**Figure 4F**)[79]. In addition,

CXCL11-CXCR3 interactions between myeloid and B/T cells were enriched in *TP53*[mut] LUAD

(**Figure 4F**) with *CXCR3* specifically expressed by NK, B, and T cells (**Figure 4G**) and *CXCL11*

induced in multiple myeloid subsets in *TP53*[mut] LUAD (**Figure 4H**), most highly in TAM.CXCLs

(**Figure S7I**), which constitute a higher proportion of myeloid cells in *TP53*[mut] LUAD. Thus,

myeloid cells may enhance recruitment of *CXCR3*-expressing T cells in *TP53*[mut] LUAD. Finally,

there was a significant increase in putative *SPP1*-mediated interactions between myeloid and

endothelial cells or pericytes in *TP53*[mut] LUAD (**Figure 4F**), also supported by increased spatial

correlation of *SPP1* and *CD44* expression in *TP53*[mut] tumors (**Figure 4I**). Thus, *TP53*[mut] LUAD

has increased expression of putative interactions involved in monocyte and lymphocyte

recruitment, TAM differentiation, hypoxia, EMT, and other downstream effector functions *via*

SPP1, CXCL11, and other TAM-associated immunomodulators.

Enrichment of T cell exhaustion programs and putative immune checkpoint interactions in

***TP53*[mut] LUAD**

Lymphoid cells in the scRNA-seq data partitioned into 13 subsets of T and NK cells (**Figure 5A,B**

and **Figure S8A**), follicular and marginal zone B cells, IgA and IgG plasma cells, plasmacytoid

DCs, and mast cells (**Figure S8B**). While there were no significant differences in B cell

composition, *TP53*[mut] LUAD had a decreased proportion of *CD4*[+] tissue-resident memory T cells

(CD4.TRM) and an increase in pre-exhausted-like[81], *GZMK*-expressing *CD8*[+] (CD8.GZMK) and

exhausted-like (T.Exhausted) T cell populations out of all T and NK cells (**Figure 5C**), which was

supported by our scRNA-seq validation compendium (**Figure S8C**) and by deconvolved bulk

RNA-seq data from TCGA primary tumors (**Figure S8D**). Higher abundance of exhausted T cells

may be mediated by increased chemotaxis *via CXCL11-CXCR3* interactions (**Figure 4F-H**), where

CXCR3 was upregulated in *TP53*[mut] exhausted T cells (**Figure S8E**). We also observed a higher

proportion of TFH cells relative to all T and NK cells in *TP53*[mut] *vs. TP53*[WT] tumors in both our

discovery and validation scRNA-seq cohorts (**Figure S8C**), which was statistically significant

when combining all samples. Expression of *CXCL13*, a top predictor for response to checkpoint

inhibition therapy[82], was most highly expressed in exhausted-like and TFH subsets and

significantly increased in *TP53*[mut] tumors (**Figure S8E**). Furthermore, expression of immune

checkpoint molecules *PDCD1*, *CTLA4*, *HAVCR2*, and *TIGIT* were significantly higher across

multiple T cell subsets in *TP53*[mut] *vs*. *TP53*[WT] tumors, while expression of *KLRB1*, a marker of

favorable prognosis in cancer[83], was downregulated (**Figure S8E**). Taken together, increased

expression of *CXCL13* and immune checkpoint molecules (*PDCD1*, *CTLA4*, *HAVCR2*, and

TIGIT) and proportions of TFH, *GZMK*-expressing *CD8*[+], and exhausted-like T cells all suggest a

heightened immunogenic potential in the *TP53*[mut] LUAD TME that may contribute to a more

favorable ICI immunotherapy response[82,84-86].

Ligand-receptor analysis of putative immune checkpoint interactions[87] between T, myeloid, and

malignant cells showed an enrichment in *PDCD1-CD274* and *TIGIT-PVR* putative interactions

between T and malignant cells in *TP53*[mut] *vs*. *TP53*[WT] LUAD (**Figure 5D**) and in *CTLA4-CD86*

and *HAVCR2-LGALS9* putative interactions between T and myeloid cells. Indeed, we found higher

expression of *CD274*, *CD86*, *PVR*, *PDCD1*, *CTLA4*, and *TIGIT* expression in *TP53*[mut] bulk RNA-

seq LUAD tumor samples from TCGA (**Figure S8F**), higher expression of *PVR* and *CD274* in

TP53[mut] bulk proteomic LUAD tumor samples from CPTAC (**Figure S8G**), and higher expression

of *PDCD1*, *CTLA4*, and *TIGIT* in multiple T cell subsets in our scRNA-seq data (**Figure S8E**,

right), where *TIGIT* was the most frequently expressed across T cells compared to *CTLA4* and

PDCD1 (**Figure 5E**). *PVR* expression was also higher in malignant cells from *TP53*[mut] *vs*. *TP53*[WT]

tumor scRNA-seq data (**Figure 5F**, **left**) and highly correlated with the expression of the cell cycle

program in malignant cells (**Figure 5F**, **right**), suggesting that *PVR* may be activated during cancer

cell proliferation. Furthermore, there was significantly higher spatial colocalization of *TIGIT* and

PVR expression in *TP53*[mut] *vs*. *TP53*[WT] LUAD (**Figure 5G**). In agreement, there was also a higher

proportion of *PVR*[+] malignant cells in *TP53*[mut] *vs*. *TP53*[WT] tumors (**Figure 5H**, **top**) and even more

prominently in cycling malignant cells (**Figure 5H**, **bottom**).

To investigate the *TP53*[mut] enrichment in interactions between PD-L1 (*CD274*) and PD1 (*PDCD1*)

at a single-cell resolution, we applied multiplex immunofluorescence (mIF) to patient-matched

TP53[mut] and *TP53*[WT] formalin-fixed paraffin-embedded (FFPE) slides from our cohort (**Figure 5I**

and **Figure S8H**). There was a significant increase in log count density (cells/mm^2) of PD-

L1[+]cytokeratin[+] cells, as well as PD1[+]CD8[+] cells in *TP53*[mut] *vs*. *TP53*[WT] tumor samples (**Figure**

5J), and, consistent with the scRNA-seq cell-cell interaction inference (**Figure 5D**), an increase in

colocalization between PD-L1[+]cytokeratin[+] cells and PD1[+]CD8[+] cells in *TP53*[mut] *vs*. *TP53*[WT]

tumor samples, across multiple spatial analyses measuring direct contact, adjacency (as defined by

cells within the 6 nearest neighbors), and proximity (<50 μm) (**Figure 5K**). Taken together,

increased expression of *PVR*, *TIGIT*, *CD274*, and *PDCD1* was concurrently observed with

increased colocalization between the ligands and corresponding receptors in *TP53*[mut] LUAD.

Spatially-defined fibroblast-macrophage niche enriched in hypoxia and EMT programs in

***TP53*[mut] LUAD**

To comprehensively map inter-compartment cellular colocalization using the spatial data, we

deconvolved the proportion of cell subsets across all spatial transcriptomics spots and calculated

the average correlation of cell type proportions across all LUAD sections and their difference

between *TP53*[mut] and *TP53*[WT] LUAD tumors (**Figure 6A**, **Figure S9A**). Interestingly, there was a

significantly lower colocalization (*i.e.,* more negative correlation) of endothelial cells with

mesenchymal cells and with myeloid cells in *TP53*[mut] *vs*. *TP53*[WT] LUAD (**Figure 6B**), consistent

with the increase in hypoxia related gene expression in mesenchymal and myeloid subsets in

TP53[mut] tumors. In addition, spatial colocalization of B, Plasma, and malignant cells was increased

in *TP53*[mut] LUAD (**Figure S9B**). B cell infiltration has been associated with increased PD1/PD-

L1 expression, TMB, and prolonged survival in NSCLC[88] and likely contributes to the heightened

immunogenicity of *TP53*[mut] LUAD tumors.

We further identified multicellular communities by an NMF-based method to derive 15 factors

with varying participation of different cell subsets (**Figure 6C**). Factor 7 (NMF7) contained

multiple myeloid and mesenchymal subsets with known significance in tumor progression,

including TAM.SPP1, CAF.COLs, and myofibroblasts, which were highly colocalized with each

other, as well as with hallmark hypoxia and EMT program expression (**Figure 6D**). Spots high in

NMF7 were low in malignant cell proportions and high in myeloid and mesenchymal cell

proportions (**Figure S9C**). NMF7 was also more positively correlated with hallmark hypoxia and

EMT program expression in the tumor periphery (spots with ≤50% malignant cells) and more

negatively correlated with endothelial cells across all spots in *TP53*[mut] *vs*. *TP53*[WT] LUAD (**Figure**

6E). Because CAFs and macrophages are both suggested to induce EMT in tumors[89,90], NMF7

may represent a hypoxic fibroblast-macrophage niche that resides on the periphery of the tumor

core and may contribute to the early EMT phenotype in malignant cells in *TP53*[mut] LUAD.

DISCUSSION

Targeted therapies for NSCLC are currently directed at oncogenic driver mutations and

chromosomal rearrangements (*e.g.*, *EGFR*, *KRAS-G12C, ALK*), yet there are currently no effective

therapies targeting *TP53* mutations, despite being common in tumors and extensively studied.

Restoration of *Trp53* expression has been shown to cause tumor regression in mice[91], leading to

efforts to treat *TP53*^mut tumors by restoring wild-type p53 activity[14]. We hypothesized that cellular

and spatial profiling of the TME in *TP53*^mut LUAD could provide a more detailed understanding

of how *TP53* mutations could contribute to shorter survival and suggest new therapeutic

opportunities. To this end, we characterized the LUAD TME of *TP53*^mut *vs*. *TP53*^WT tumors by

combining scRNA-seq of 167,193 cells across 23 NSCLC patients, 23 matched tumor and normal

WES, 20 spatial transcriptomics and 4 mIF tissue sections. Together, these represent the most

comprehensive multi-omic map of NSCLC to date and provide a valuable resource of cell

annotations, markers, and integrated data sets for the scientific community.

Our integrative analysis revealed a substantial remodeling of the TME associated with *TP53*

mutations (**Figure 6F**), including decreased presence of pericytes and endothelial aerocytes. The

mechanisms of TME remodeling are likely complex but our study uncovers multiple ones that may

play a role. These include putative ligand-receptor interactions between malignant and endothelial

cells (*SEMA3A-NRP1*, *EPHB2-EFNB1*) that were enriched in *TP53*^mut tumors and have known

roles in inhibiting vascularization, and which were significantly more colocalized in the *TP53*^mut

spatial data. *TP53*^mut-specific depletion of endothelial cells was accompanied by increased hypoxia

and pEMT program expression in malignant cells, which was altered across multiple functional

categories of *TP53* mutations in bulk transcriptomic and proteomic data, most prominently in

DNE,LOF and Impactful II *TP53* mutations. This effect was independent of presence or absence

of *KRAS* mutations but was partially reduced in tumors with *TP53-EGFR* co-mutations. Hypoxia

has been shown to induce EMT through *HIF-1α* as well as other mediators[92,93] and to positively

select for p53-deficient cells that have lost apoptotic potential in mice[94]. Thus, a hypoxic TME

may provide a survival advantage for highly plastic *TP53*[mut] LUAD malignant cells, which have

lost alveolar identity and experience increased proliferative capacity and metastatic potential. This

model is supported by our analysis of data from genetically engineered K and KP mouse models,

and from A549 LUAD cell line (*TP53*[WT]) overexpressing different *TP53* mutational variants.

Furthermore, *de novo* discovery of spatial multicellular communities revealed a highly hypoxic,

EMT-promoting spatial niche enriched for *SPP1*[+] TAMs, myofibroblasts, and collagen producing

CAFs. *SPP1* expression was highly upregulated in *TP53*[mut] monocytes and TAMs and tightly

linked with regulation of hypoxia and EMT. Increased TGF-β signaling likely also helps shape an

EMT-supporting niche in *TP53*[mut] tumor samples, where *TGFB2-TGFBR2* interactions involving

malignant cells were significantly enriched in spatial data. Previous studies linking mutant p53 to

metastatic potential have focused on malignant cell-intrinsic processes[11,12,95], as the effects of

mutant p53 on tumor-stromal and tumor-immune crosstalk have been described only recently in

model systems[96]. Our study is the first to profile the cellular and spatial context of *TP53* mutated

LUAD tumors to provide insight into how *TP53* mutations could remodel the TME to promote

tumor survival and metastasis in patients, leading to poorer outcomes.

Interestingly, one aspect of TME remodeling in *TP53*[mut] LUAD is a higher tumor infiltration by B

cells and *CD8*[+] T cells. Increased lymphocyte recruitment may be caused in part by enriched

CXCL11-CXCR3 ligand-receptor interactions between myeloid and B/T cells in *TP53*[mut] tumors,

most notably exhausted T cells. We observed some overlap of the myeloid and T cell phenotypes

in $TP53^{mut}$ LUAD, the anti-tumor immunity hub we reported previously in colorectal cancer[97], and

the stem immunity hub in NSCLC[98], but additionally report here B/Plasma cell tumor infiltration,

T cell exhaustion (supported by an independent scRNA-seq validation compendium), and the

enrichment of $SPP1$-expressing TAMs. Our findings of a potentially more immunogenic TME in

$TP53^{mut}$ LUAD is consistent with previous studies reporting higher levels of anti-tumor immune

signatures[99] in $TP53^{mut}$ vs. $TP53^{WT}$ LUAD using bulk RNA-seq data from TCGA and longer

survival in advanced $TP53^{mut}$ vs. $TP53^{WT}$ NSCLC patients receiving immune checkpoint

therapy[100]. Previous studies have reported that p53 mutations increase expression of PD-L1 via

decreased inhibition by miR-34[101] and that LUAD tumors with $KRAS$-$TP53$ co-mutations have

higher expression of PD-L1 and improved response to anti-PD1 treatment compared to

$KRAS^{mut}TP53^{WT}$ tumors[19,21,23,102]. The link between mutation status and immunotherapy response

has also been shown in the context of other mutations, where $KRAS^{mut}STK11^{mut}KEAP1^{mut}$ LUAD

tumors demonstrate reduced response compared to $KRAS^{mut}STK11^{WT}KEAP1^{WT}$ tumors[103]. In

agreement, our computational screen for immune checkpoint receptor-ligand pairs showed

enrichment in $PDCD1$-$CD274$, $CTLA4$-$CD86$, and $TIGIT$-PVR interactions in $TP53^{mut}$ vs. $TP53^{WT}$

LUAD tumors; $CTLA4$-$CD86$ and $TIGIT$-PVR interactions have not yet been studied in the context

of $TP53$ mutations to the best of our knowledge. Increased expression of $TIGIT$ in T cells and PVR

in malignant cell both contribute to the spatially-resolved enrichment of $TIGIT$-PVR co-

expression, which we observed in all 8 $TP53^{mut}$ LUAD tumors in our cohort, as well as in bulk

RNA-seq data from TCGA and bulk proteomic data from CPTAC. Recent trials demonstrate

benefit to combining anti-PD1 ICI with chemotherapy as a neo-adjuvant in resectable LUAD[104].

Moreover, trials targeting TIGIT alone or in combination with PD1/PD-L1 in lung cancer have yet

to demonstrate substantial survival benefit, highlighting the importance of identifying subsets of

patient tumors that could benefit from therapeutic intervention (*e.g.* *TP53*[mut], PVR-expressing

tumors)[105]. Future studies will be needed to assess the predictive potential of *TP53* mutational

status for response to these and other therapies.

More generally, the methodology introduced in this study can be used to study the effect of *TP53*

or other mutations on the TME in lung cancer and other tumor types and to infer multicellular

pathways and communities to target for tumor suppressors in other contexts, where more effective

therapies are needed.

METHODS

Human subjects

Fresh solid primary tumor tissue was collected using an IRB-approved protocol (IRB protocol 98-063) at DFCI/BWH and MGH. Patients were all confirmed to be treatment-naïve and consented to the study prior to collection. Immediately after resection, the tissue was reviewed by the clinical pathology team and high-quality portions (determined based on tumor content, necrosis, calcification, fat, and hemorrhage) were allocated for WES, scRNA-seq, spatial transcriptomics, and mIF. The protocol was designed to reduce the time between surgical resection, anatomic pathology review, placement in media, and processing at the Broad Institute. Blood was drawn from the same patients and cryo-preserved at -80 °C for subsequent processing. Patient demographic information, including age, sex, and smoking history, are included in **Table S1**.

Whole-exome sequencing sample processing

DNA was extracted from fresh frozen lung cancer tissue embedded into OCT (TissueTek, Sakura) and from PBMCs isolated from preserved patient-matched blood (AllPrep DNA/RNA extraction kit, Qiagen). Library construction was performed as previously described[106]. In brief, DNA input for shearing was diluted to a final concentration of 20–250 ng in 50 µl of solution. Adapter ligation was performed using palindromic forked adapters (Integrated DNA Technologies), containing unique dual-indexed molecular barcode sequences to improve downstream pooling. End repair/A-tailing, adapter ligation, and library enrichment PCR were carried out using a 96-reaction kit format (Kapa HyperPrep). During solid-phase reversible

immobilization (SPRI), a final elution volume of 30 µl was produced to maximize library

concentration, and vortexing was performed to maximize the effluent. Constructed libraries were

first pooled and hybridization and capture were performed (Illumina's Nextera Exome Kit) using

the recommended protocol from the manufacturer, using a skirted PCR plate to facilitate

automation (Agilent Bravo liquid handling system). Library pools then underwent qPCR

quantification and libraries were adjusted to a concentration of 2nM. DNA libraries were cluster-

amplified using exclusion amplification chemistry in patterned flow cells (Illumina) according to

the manufacturer's recommended protocol. Sequencing of flow cells was performed using

sequencing-by-synthesis chemistry and analyzed using RTA v.2.7.3 or later. Library pools were

sequenced on 76 cycle runs using two 8-cycle index reads across the appropriate number of lanes

to attain coverage for all libraries.

Tissue processing, CD45 sorting, and scRNA-sequencing

Tumor tissue resections were transferred in RPMI on ice from the operating room and washed in

cold PBS in the laboratory and transferred to 5ml Eppendorf tubes containing 3ml dissociation

mixture (NSCLC PDEC protocol[107]). Samples were minced in the Eppendorf tube using scissors

into pieces smaller than ~0.4 mm and incubated at 37 °C, rotating at ~14 rpm. for 10 min. Each

sample was pipetted 20 times with a 1ml pipette tip at room temperature and placed back into

incubation for 10 min. The sample was pipetted again 20 times using a 1 ml pipette tip,

transferred to a 1.7 ml Eppendorf tube and centrifuged at 300–580g for 4–7 min at 4 °C. The

pellet was resuspended in 200–500 µl of ACK (ammonium-chloride-potassium) RBC lysis buffer

(Thermo Fisher Scientific, A1049201) and incubated for 1 min on ice. Twice the volume of cold

PBS was added to stop the reaction, and cells were pelleted by a short centrifugation for 8 s at

4 °C, using the short spin setting and ramping up to 11,000g. If RBCs remained, the RBC lysis

step was performed up to two additional times. To assess cell count and viability, 5 µl of Trypan

blue (Thermo Fisher Scientific, cat. no. T10282) was mixed with 5 µl of the sample and loaded

onto an INCYTO C-Chip Disposable Hemocytometer, Neubauer Improved (VWR, cat. no.

82030-468). Depletion of CD45$^+$ cells in NSCLC samples was performed using CD45

MicroBeads (Miltenyi Biotec, cat. no. 130-045-801) according to the manufacturer's protocol.

The single-cell suspension was centrifuged at 500g for 4 min at 4 °C and the pellet was

resuspended in 80 µl of MACS buffer (PBS supplemented with 0.5% BSA, and 2 mM EDTA)

per 10^6 cells. MACS CD45 microbeads were added to the cell suspension (20 µl per 10 million

cells) and the cells were incubated on ice for 15 min. During incubation, a MS column was

attached to a MidiMACS separator and rinsed with 3 ml MACS buffer. After incubation, the

cells-bead conjugate was washed with 900 µl MACS buffer per 10 million cells. The cells were

then centrifuged at 500g for 4 min at 4 °C and the pellet was resuspended in 500 µl MACS

buffer. The cell suspension was transferred to the column and the effluent was collected

(CD45$^-$ fraction). The column was washed three times with 3 ml MACS buffer. The

CD45$^-$ fraction was centrifuged at 500g for 4 min at 4 °C. The pellet was resuspended in 50 µl of

0.4% BSA (Ambion, cat. no. AM2616) in PBS. Cells were re-counted and adjusted to a final

range of 200–2,000 cells per µl.

NSCLC tissue dissociation protocol

The digestion mix contained 2692 µl HBSS (Thermo Fisher Scientific, cat. no. 14170112),

187.5 µl of 20 mg ml^{-1} pronase (Sigma Aldrich, cat. no. 10165921001) diluted to a final

concentration of 1,250 µg ml^{-1}, 27.6 µl of 1 mg ml^{-1} elastase (Thermo Fisher Scientific, cat. no.

NC9301601) diluted to a final concentration of 9.2 µg ml^{-1}, 30 µl of 10 mg ml^{-1} DNase I (Sigma

Aldrich, cat. no. 11284932001) diluted to a final concentration of 100 µg ml^{-1}, 30 µl of

10 mg ml^{-1} Dispase (Sigma Aldrich, cat. no. 4942078001) diluted to a final concentration of

100 µg ml^{-1}, 30 µl of 150 mg ml^{-1} collagenase A (Sigma Aldrich, cat. no. 10103578001) diluted

to a final concentration of 1,500 µg ml^{-1} and 3 µl of 100 mg ml^{-1} collagenase IV (Thermo Fisher

Scientific, cat. no. NC9836075) diluted to a final concentration of 100 µg ml^{-1}.

scRNA-seq library preparation and sequencing

A total of 8,000 cells were loaded onto each channel of the Chromium Controller (10X

Genomics) to generate single-cell gel bead-in-emulsions (GEMs). Libraries were constructed

using the Chromium Single Cell 3′ Library & Gel Bead Kit v.2 or v.3 (PN-120237, 10X

Genomics): barcoded reverse transcription of RNA, cDNA amplification, fragmentation and

adapter and sample index attachment were all performed according to the manufacturer's

recommended protocol. Barcoded libraries from four 10x channels were pooled together and

sequenced on either one lane of an Illumina HiSeq X or one flow cell of a NextSeq, with paired

end reads accordingly: read 1, 26 nt; read 2, 55 nt; index 1, 8 nt; index 2, 0 nt.

Tissue processing for Spatial Transcriptomics

Fresh frozen lung cancer tissue samples were embedded into OCT (TissueTek, Sakura) and

shipped to the Broad Institute for cryosectioning and generation of hematoxylin and eosin-

stained (H&E) tissue sections. H&E-stained tissues were subject to pathology review to assess

tissue quality, structural preservation, cellular viability, tumor content, inflammation, fibrosis,

and necrosis. Samples were excluded based on small tissue size, low cellularity, or extensive

fibrosis. Regions of interest (ROIs) were then selected based on the criteria mentioned and

marked on H&E images for subsequent tissue sectioning and mounting on Visium slides. Visium

sectioning and processing were performed on the eight samples that passed QC. Tissues were

cryosectioned at 10 um thickness at -22°C and placed in the capture areas of Visium Tissue

Optimization slides (3000394, 10X Genomics) and Visium Spatial Gene Expression slides

(2000233, 10X Genomics). The tissue sections were adhered by warming the backside of the

slides and placed at -80°C for 1-3 days.

Visium spatial gene expression library generation

The tissue optimization sample slide and spatial gene expression slide were processed according

to the manufacturer's protocols

(CG000238_VisiumSpatialTissueOptimizationUserGuide_Rev_A.pdf and

CG000239_VisiumSpatialGeneExpression_UserGuide_Rev_C.pdf). In short, following tissue

methanol fixation and H&E staining, brightfield morphology images were obtained with a Zeiss

Axio microscope using the Metafer slide-scanning platform (Metasystems) at 10x resolution.

The images were joined together with the VSlide software (Metasystems) and exported as tiff

files. Optimal tissue permeabilization times were tested with eight different time points (0, 3, 6,

12, 18, 24, 30, 36 minutes) on the tissue optimization sample slide for one of the samples. 12

minutes of permeabilization was set as the optimal time point and used for the spatial gene

expression slide for all samples. The released RNAs from the tissue were reversely transcribed

into complimentary DNA by priming to the spatial barcoded primers on the glass in the presence

of template switching oligo. After second strand synthesis, a denaturation step released the

cDNAs, which were then amplified with 14-15 cycles of PCR amplification. Finally, indexed

sequencing-ready, spatial gene expression libraries were constructed according to the

manufacturer's protocol. The libraries were pooled together to generate >50,000 reads per spatial

spot and sequenced on Illumina NovaSeq sequencers with S1 and SP kits and the settings: read

1, 28 cycles; read 2, 90 cycles; index 1, 10 cycles; index 2, 10 cycles.

Multiplex immunofluorescence

Staining was performed on a BOND RX fully automated stainer (Leica Biosystems). FFPE tissue

sections of 5-μm thick were baked for 3 hours at 60°C before being loaded into the BOND RX.

Slides were deparaffinized (BOND DeWax Solution, Leica Biosystems, Cat. AR9590) and

rehydrated with multiple rounds of graded ethanol to deionized water. Antigen retrieval was

performed in BOND Epitope Retrieval Solution 1 (pH 6) or 2 (pH 9) (ER1, ER2, Leica

Biosystems, Cat. AR9961, AR9640) at 95°C. Deparaffinization, rehydration and antigen

retrieval are carried out by BOND RX for ~ 14 hours. Slides were then serially stained with

primary antibodies with an incubation time of 30 minutes per antibody. Anti-mouse and anti-

rabbit Opal Polymer Horseradish Peroxidase (Opal Polymer HRP Ms + Rb, Akoya Biosciences,

Cat. ARH1001EA) were then applied as a secondary label for 10 minutes. Signal for antibody

complexes was labeled and visualized by their corresponding Opal Fluorophore Reagents

(Akoya) by incubating the slides for 10 minutes. Opal Fluorophore 780 was paired with a TSA-

DIG amplification to ensure analyzable signal. Slides were incubated in Spectral DAPI solution

(Akoya) for 10 minutes, air dried and mounted with Prolong Diamond Anti-fade mounting

medium (Life Technologies, Cat. P36965) and stored in a light-proof box at 4℃ prior to

imaging.

Imaging was performed using the PhenoImager multispectral imaging platform (Akoya

Biosciences, Marlborough, MA). Each slide was scanned at 20x resolution as a whole-slide

image. These images were then accessed through Phenochart viewing software (Akoya

Biosciences) where 4-8 20x regions of interest (ROIs) were selected. After ROI selection was

approved by a pathologist (S.J.R.), the images were spectrally unmixed and analyzed using

Inform 2.6 (Akoya Biociences). Each analyzable ROI was segmented and quantified for

expression of each marker utilizing the Inform analysis tools.

WES data processing

The Picard pipeline (http://picard.sourceforge.net/) was used to align the tumor and normal

whole exome sequences to the hg19 reference human genome build and generate BAM files.

Somatic short variant discovery was performed using the Mutect 2 algorithm

(https://software.broadinstitute.org/gatk/documentation/tooldocs/current/org_broadinstitute_gatk

_tools_walkers_cancer_m2_MuTect2) from the GAT4K pipeline run on the Terra cloud-based

platform. Somatic mutations were annotated using Funcotator

(https://gatk.broadinstitute.org/hc/en-us/articles/360035889931-Funcotator-Information-and-

Tutorial). Copy number ratios for each exon was calculated by comparing mean exon coverage

with expected coverage based on a panel of normal samples and were then segmented for

downstream analysis. Maf and vcf files were generated as outputs, which were then subsequently

processed with maftools[108] and cnvkit[109] packages.

Large-scale CNA inference and correlation

The inferCNV package[110] (https://github.com/broadinstitute/inferCNV) was applied to infer

large-scale CNVs from scRNA-seq data. Normal lung epithelial cells from a separate dataset

were used as a reference. Raw UMI counts, a gene ordering file, and an annotation file

specifying normal and tumor cells and patient sample of origin were used as input. The function

infercnv::run() was performed with a cutoff of 0.1, cluster_by_groups = T, and HMM = T. CNV

values were predicted using a 6-state Hidden Markov Model (HMM) and were normalized to the

3rd quantile of all values per sample. Cells with > 0.7 normalized CNV values were assigned as

malignant cells. CNVs called from WES data were used as a positive control to assess the

accuracy of the inferCNV analysis using scRNA-seq data. Spearman correlation was run

between inferred CNVs from scRNA and from WES data across genes and plotted as a heatmap.

scRNA-seq preprocessing

CellRanger v3.1 was used to align 10X Chromium reads to the GRCh38 human genome

reference and generate barcode, gene, and count matrices. Preprocessed matrices were then

loaded into Seurat v4[111] in R v4.1.0 using the Read10X() function. Cells with less than 1000

unique molecular identifiers (UMIs), less than 400 detected genes (1000 genes for malignant

cells), or greater than 25% mitochondrial genes were excluded from further analyses.

scRNA-seq integration, clustering, annotation, and doublet removal

The UMI counts matrices were batch-corrected by patient sample, using the Seurat v4

SCTransform() (using 2000 variable features and regressing out percentage of mitochondrial

genes), FindIntegrationAnchors(), and IntegrateData(), using the largest sample as the reference

dataset. PCA was then performed in the variable gene space and 15 principal components were

used for Louvain clustering and UMAP dimensionality reduction. Cell type markers for broad

cell classes were identified using the FindAllMarkers() function, using a maximum of 1,000 cells

per cluster. Cells were annotated based on expression of canonical markers that were established

by previous literature. Individual cell classes were then subset and subjected to another series of

integration, clustering, and annotation based on expression of subset-specific markers. Doublets

were removed based on expression of markers specific to more than one broad cell classes.

scRNA-seq signature scoring

To identify broad cell classes signatures, we used the FindAllMarkers() function in Seurat v4 and

generated gene sets using the most differentially expressed genes for each class, based on $\log_2$

fold change and Bonferroni-corrected p-values compared to the other clusters. We identified

malignant program signatures by performing anchor-based integration across patient samples

malignant cells followed by Louvain clustering at resolution 1.2 and using the FindAllMarkers()

function to derive the top 10 most differentially expressed genes from each cluster, based on $\log_2$

fold change and Bonferroni-corrected p-values. Signatures for normal lung epithelial cells were

derived from published healthy lung scRNA-seq data using the same method. The lists of broad

cell classes and cell subset marker genes, as well as final gene set signatures are provided in

Table S2. To define hallmark EMT and hypoxia scores, we used gene sets termed

"HALLMARK_EPITHELIAL_MESENCHYMAL_TRANSITION" and

"HALLMARK_HYPOXIA" downloaded from MSigDB[112]. Healthy lung epithelial cell

signatures were derived using published scRNA-seq data from the human lung cell atlas[113], as

well as *a priori* knowledge: AT2 (*SFTPC, SFTPA2, SFTPA1, SFTPD, SFTPB*), AT1 (*AGER,*

EMP2, CAV1, HOPX), basal (*KRT5, KRT17, KRT15, S100A2*), neuroendocrine (*CALCA,*

CHGA, *GRP*, *SCG5*, *ASCL1*), goblet (*MUC5AC*, *MUC5B*, *S100P*), secretory (*BPIFB1*, *PIGR*,

SCGB3A1, *SCGB1A1*), ciliated (*TMEM190*, *CAPS*, *FOXJ1*, *TPPP3*, *FAM183A*), and ionocyte

(*ASCL3*, *ATP6V0B*, *FOXI1*, *CLCNKB*, *CFTR*). Signature scores were assigned per cell in

scRNA-seq data using the AddModuleScore() function in Seurat v4, which compares the average

expression of the genes in each gene set to the average expression of a control gene set of 100

randomly selected genes within the same expression bin.

scRNA-seq differential expression analysis

The data was first split into different cell subsets, and pseudobulk profiles were generated by

setting the identity of the objects to patient sample ID and running the AverageExpression()

function in Seurat v4. A Mann-Whitney Wilcoxon test was used to test for Bonferroni-corrected

significance in gene differential expression between groups. The average $\log_2$ fold change was

plotted using the ComplexHeatmap package in R, and asterisks were added for genes that were

significantly differentially expressed.

Gene Set Enrichment Analysis

Using the top 100 DEGs between conditions per cell subset or the top 100 specific markers per

subset, we performed gene-set enrichment analysis using the R package clusterProfiler[114] (v4.0),

focusing on GO terms, KEGG, Reactome, and Hallmark Cancer gene sets. Gene ratios and

adjusted *p*-values were visualized as dot plots, after applying BH correction and setting *p*-value

and *q*-value cutoffs of 0.05.

Similarity to normal lung and K/KP mouse cells

To identify transcriptomic similarities between cell subsets in NSCLC and the normal lung, the

Seurat v4 function TransferData() was used. The broad cell compartment object in the NSCLC

scRNA-seq dataset was assigned to the query parameter and the corresponding compartment in a

published normal lung scRNA-seq dataset[56] was assigned to the reference parameter, and the

remaining parameters were set to defaults. Each cell in the NSCLC object was assigned a

predicted identity based on the normal lung subset it was most similar to transcriptionally. A

confusion matrix was created, displaying the number of cells that overlapped between predicted

identities from the normal lung and annotations assigned *de novo*. A similar approach was used

to classify K/KP mice HPCS cells to their closest annotation in human tumor cells.

Signaling Entropy calculation

Entropy was inferred for each epithelial cell using the R package SCENT[64], using the default

protein-protein interaction (PPI) network ("net17Jan2016.m"). Raw UMI counts were used as

the scRNA-seq input. Entropy was calculated separately for malignant and non-malignant

epithelial cells. Quality control was performed using the scater package in R, and low-quality

cells based on library size, number of expressed genes, and mitochondrial gene proportion were

removed. Differential potency estimation was performed using the DoIntegPPI() function, which

generated entropy values per cell. The same method was used to calculate entropy values for

bulk RNA-seq data from TCGA and bulk proteomic data from CPTAC. Entropy values were

compared using the ggviolin() function using the ggpubr package in R.

Cell-cell interaction analysis

CellPhoneDB[115] (v2.1.7) was used to predict significant ligand-receptor interactions using

normalized raw counts and annotation of broad cell classes as input. The method was run with

default parameters separately for each sample ID to account for batch effects between samples.

Interactions with a p-value below 0.05 were classified as significant. The significant means

output was converted into a binary matrix based on presence or absence of a significant

interaction. Normalized enrichment difference (ED) of a specific interaction was calculated as

follows:

$$ED = n_A/N_A - n_B/N_B$$

where nA and nB represent the number of tumor samples with the interaction in groups A and B,

respectively, while NA and NB represent the total number of tumor samples in these groups.

Significance of enrichment for an interaction was calculated using a Fisher's exact test. Negative

$\log_{10}$ p-values and normalized enrichment differences for each ligand-receptor interaction and

cell pair combination were plotted using the ggplot2 package in R.

scRNA-seq regulon analysis

A scRNA-seq transcription factor (TF) gene regulatory network was constructed using

pySCENIC[73] (v0.11.2). Raw UMI counts were used as input and regulons were predicted using

gene module co-expression using the GRNBoost2 package. The number of regulons was pruned

using the feather ranking databases (hg19-500bp-upstream-7species.mc9nr, hg19-tss-centered-

10kb-10species.mc9nr, hg19-tss-centered-5kb-10species.mc9nr). Regulon activity, driving TFs,

and weights for individual target genes were predicted with the cisTarget() function in

pySCENIC. Regulon activity enrichment scores were predicted for each cellular subset using

AUCell and regulon specificity scores were used to identify regulatory networks specific to each

cell subset. Z-score-transformed regulon x subset matrices were visualized using the

ComplexHeatmap package in R.

Additional scRNA-seq datasets

Processed scRNA-seq data from an additional 5 studies (both published and unpublished) with

available *TP53* mutational status were assembled into a validation compendium, consisting of 15

TP53[WT] and 7 *TP53*[mut] tumor samples (ArrayExpress accessions E-MTAB-6149 and E-MTAB-

6653[26]; https://humantumoratlas.org/ (unpublished); GEO accession GSE123904[32]; BIG

accession HRA000154[55]; https://doi.org/10.24433/CO.0121060.v1[33]). *In-vivo* scRNA-seq data

for *KRAS*[mut]*TP53*[WT] and *KRAS*[mut]*TP53*[mut] mice (Smart-Seq2) was obtained from GEO accession

GSE152607[63]. *In-vitro* scRNA-seq data from lung cancer A549 cell lines with different induced

TP53 variants was obtained from GEO accession GSE161824[61].

Signature scoring of bulk expression profiles

Bulk RNA-seq and somatic mutation data from TCGA LUAD and somatic mutations were

downloaded from cBioPortal[116] using the cgdsr package in R. Curated gene sets with highly

specific markers for individual cell types, subsets, or malignant programs were generated from

the scRNA-seq data as described above. The signature scores for these gene sets were defined as

the mean $\log_{10}$ expression of all the genes in a given gene set and were used to estimate the

relative frequency of cell types, cell subsets in bulk RNA-seq data from TCGA, or activity of

malignant cell programs. The same approach was used for calculating expression signatures in

bulk proteomic data from CPTAC[117], which was downloaded from the CPTAC Data

Portal https://cptac-data-portal.georgetown.edu/cptacPublic/. We confirm the reproducibility of

all our bulk deconvolution results using BayesPrism, a deconvolution method based on a

Bayesian framework that incorporates single-cell expression profiles[118].

Survival analysis

To predict the roles of individual genes and gene signatures on patient survival, TCGA LUAD

tumor samples were classified into high (top 25th percentile), medium (25th to 75th percentile),

and low (bottom 25th percentile); or high (top 50th percentile) and low (bottom 50th percentile)

expression groups. Kaplan-Meier (KM) curves were generated using the survival package in R,

and *p*-values were calculated using a log-rank test or bivariate Cox proportional-hazards

regression analysis to evaluate the significance of different model parameters (*e.g.*, expression

level of genes or malignant signatures, *TP53* mutation status) in relation to overall survival

(defined as time to death from any cause). The reported *p*-values from the Cox proportional-

hazards model reflect the significance of one specific covariate while adjusting for *TP53*

mutation status.

Effects of co-mutations and *TP53* variant classes

TCGA tumor samples were stratified into six mutually exclusive categories with sufficiently large

sample sizes for comparative analysis based on combinations of mutations in *KRAS*, *EGFR*, and

TP53 (*KRAS*^{mut}, *EGFR*^{mut}, *TP53*^{mut}, *KRAS*^{mut}*TP53*^{mut}, *EGFR*^{mut}*TP53*^{mut}, and *KRAS*^{WT}*EGFR*^{WT}

TP53^{WT}). TCGA and CPTAC tumor samples were furthermore grouped into different mutually

exclusive missense variant classes based on either dominant-negative/loss of function (DNE,LOF)

status of mutational variants, or impact categories as described by prior studies[60,61]. Mann-

Whitney-Wilcoxon test *p*-values were computed for each group relative to the relevant wildtype

or control group.

Multiple linear regression and ANOVA analysis of interaction effects

Expression level of individual genes or gene signatures was predicted based on the mutation

status of four different genes (*EGFR*, *KRAS*, *STK11*, and *TP53*), and tumor mutational burden

(TMB) using the following model:

$$Y = \beta 0 + \beta 1 X1 + \beta 2 X2 + \beta 3 X3 + \beta 4 X4 + \beta 5 X5 + \epsilon,$$

where *Y* represents the expression level, *β*0 the y-intercept, *β*1, *β*2, *β*3, *β*4, and *β*5 the regression

coefficients associated with each variable, *X*1, *X*2, *X*3, and *X*4 the mutation status of different

genes (*EGFR*, *KRAS*, *STK11*, and *TP53*), *X*5 the TMB, and ϵ the error term. For each mutation

variable *Xi*, a *p*-value was computed to test the null hypothesis that the regression coefficient *βi*

is equal to zero (indicating no effect). A significant *p*-value (≤ 0.05) suggests that the mutation

status of the respective gene is a significant predictor of *Y* (*i.e.,* there exists a significant

relationship between the mutation status and the expression level). We further conducted the

same analysis using a model containing an expanded set of covariates (mutation status of *EGFR*,

KRAS-G12C, *KRAS-G12D*, *KRAS-G12V*, *STK11*, *KEAP1*, *RBM10*, *PTPRD*, and *TP53* as well as

TMB). An analysis of variance (ANOVA) was conducted on the output of the regression

model's output, evaluating the interactions between *KRAS* and *TP53*, and between *EGFR* and

TP53. A significant interaction implies that the combined effect of two individual interactions is

not merely the sum of their individual effects. The output from the ANOVA provides *p*-values

for each interaction, where a significant *p*-value suggests that the effect of one mutation is altered

by the presence or absence of the other mutation (*i.e.,* not additive).

Pathologist annotation of hematoxylin and eosin (H&E) stains from spatial transcriptomics samples

A board-certified pathologist, blinded to all sample identities and spatial transcriptomics measurements, performed manual annotation of 13 H&E stains using Loupe browser. Annotations on the H&E were classified into multiple categories (*e.g.*, cancer, lymphocytes, myeloid, fibroblast, lepidic adenocarcinoma, TLS, vascular endothelium) that were most representative of cells within barcoded spots. Spots in which there were cell types whose identities could not be readily visually resolved were not annotated. These categories were then mapped to three general cell classes (malignant, stromal, and immune) to minimize variability in pathology annotations across samples and allow for comparison with the computationally inferred cellular compositions.

Processing of spatial transcriptomics data, integration with paired scRNA-seq data, and colocalization analysis

Tangram version 0.4.0, scanpy 1.8.1, and anndata 0.7.6 was used to map scRNA-seq data to spatial locations on 10X Visium spatial transcriptomics samples as previously described[41]. UMI counts for scRNA-seq and spatial transcriptomics data were converted into anndata objects and pre-processed using standard scanpy functions[119]. The top 100 genes that best characterized each broad cell classes were selected by sc.tl.rank_genes_groups(). This subset of genes was used afterwards as training genes for Tangram alignment. To more accurately map single-cells onto spatial transcriptomics spots, an estimate of cell density per spot was attained through watershed-based nuclei segmentation of the hematoxylin and eosin-stained serial tissue section. Images

were first imported by squidpy 1.1.0[120], and smoothened using squidpy.im.process with a sigma

= 4, then segmented by squidpy.im.segment with method = "watershed", using parameters

channel = 0 to select the red color channel and thres = 120. A threshold of 120 was selected to

best separate nuclei from background, and sq.im.calculate_features was used to count the

number of segmented nuclei per spot after segmentation. Finally, mapping was performed by

tg.map_cells_to_space using mode = "constrained" and density_prior = # of nuclei per spot /

total # of nuclei to constrain alignment based on segmented nuclei density, returning

probabilities of spatial location on a per-cell basis, which were employed in later analysis. To

project individual gene expression onto spatial transcriptomics spots, tg.project_genes was used.

Cell type and subset proportion were predicted per spot by summing up cell x spot probabilities

and normalizing the total probability scores per spot to 1. Spatial colocalization was inferred

using Pearson correlation between predicted cell type and subset compositions, and additional

continuous attributes from the metadata across spots.

NMF analysis for spatial transcriptomics data

NMF analysis was adapted from cell2location[121] using cell subset proportions per spatial

transcriptomics spot as inputs, enabling analysis of spatially colocalized cellular programs. NMF

was trained five times for a range of k = 3 to 30 factors, and k = 14 was selected based on

stability of training across the five restarts and elucidating discrete spatially-delimited

compartments across spatial transcriptomics samples. Using scanpy, NMF weights were plotted

onto spatial transcriptomics data to investigate spatial-dependent patterning of NMF factors.

NMF weights per subset were visualized using the heatmap function in cell2location.

Multiplex immunofluorescence analysis

InForm-analyzed ROIs were processed using the Pythologist software[122]. Cell colocalization

involved three spatial categories: 1) direct contact, 2) adjacency within 6 nearest neighbors, and

3) proximity within 50um of PD-L1$^+$cytokeratin$^+$ cells. Proportion of PD1$^+$CD8$^+$ cells out of all

PD-L1$^+$cytokeratin$^+$ neighboring cells were computed across these spatially defined categories

when at least 20 neighboring cells were present.

REFERENCES

984   1.   Travis WD, Brambilla E, Nicholson AG, et al. The 2015 World Health Organization

985   Classification of Lung Tumors: Impact of Genetic, Clinical and Radiologic Advances

986   Since the 2004 Classification. J Thorac Oncol 2015;10(9):1243-1260. DOI:

987   10.1097/JTO.0000000000000630.

988   2.   Cancer Genome Atlas Research N. Comprehensive genomic characterization of

989   squamous cell lung cancers. Nature 2012;489(7417):519-25. DOI: 10.1038/nature11404.

990   3.   Cancer Genome Atlas Research N. Comprehensive molecular profiling of lung

991   adenocarcinoma. Nature 2014;511(7511):543-50. DOI: 10.1038/nature13385.

992   4.   Hanna NH, Robinson AG, Temin S, et al. Therapy for Stage IV Non-Small-Cell Lung

993   Cancer With Driver Alterations: ASCO and OH (CCO) Joint Guideline Update. J Clin

994   Oncol 2021;39(9):1040-1091. DOI: 10.1200/JCO.20.03570.

995   5.   Howlader N, Forjaz G, Mooradian MJ, et al. The Effect of Advances in Lung-Cancer

996   Treatment on Population Mortality. N Engl J Med 2020;383(7):640-649. DOI:

997   10.1056/NEJMoa1916623.

998   6.   Herbst RS, Morgensztern D, Boshoff C. The biology and management of non-small cell

999   lung cancer. Nature 2018;553(7689):446-454. DOI: 10.1038/nature25183.

7.   National Cancer Institute. Non-Small Cell Lung Cancer Treatment.

(https://www.cancer.gov/types/lung/patient/non-small-cell-lung-treatment-pdq).

8.   Campbell JD, Alexandrov A, Kim J, et al. Distinct patterns of somatic genome alterations

in lung adenocarcinomas and squamous cell carcinomas. Nat Genet 2016;48(6):607-16.

DOI: 10.1038/ng.3564.

9.   Halvorsen AR, Silwal-Pandit L, Meza-Zepeda LA, et al. TP53 Mutation Spectrum in

Smokers and Never Smoking Lung Cancer Patients. Front Genet 2016;7:85. DOI:

10.3389/fgene.2016.00085.

10.   Kaubryte J, Lai AG. Pan-cancer prognostic genetic mutations and clinicopathological

factors associated with survival outcomes: a systematic review. NPJ Precis Oncol

2022;6(1):27. DOI: 10.1038/s41698-022-00269-5.

11.   Al Bakir M, Huebner A, Martinez-Ruiz C, et al. The evolution of non-small cell lung

cancer metastases in TRACERx. Nature 2023;616(7957):534-542. DOI: 10.1038/s41586-

023-05729-x.

12.   Van Egeren D, Kohli K, Warner JL, et al. Genomic analysis of early-stage lung cancer

reveals a role for TP53 mutations in distant metastasis. Sci Rep 2022;12(1):19055. DOI:

10.1038/s41598-022-21448-1.

13.   Duffy MJ, Synnott NC, O'Grady S, Crown J. Targeting p53 for the treatment of cancer.

Semin Cancer Biol 2020. DOI: 10.1016/j.semcancer.2020.07.005.

14.   Bykov VJN, Eriksson SE, Bianchi J, Wiman KG. Targeting mutant p53 for efficient

cancer therapy. Nat Rev Cancer 2018;18(2):89-102. DOI: 10.1038/nrc.2017.109.

15.   Zhu G, Pan C, Bei JX, et al. Mutant p53 in Cancer Progression and Targeted Therapies.

Front Oncol 2020;10:595187. DOI: 10.3389/fonc.2020.595187.

16.   Uehara I, Tanaka N. Role of p53 in the Regulation of the Inflammatory Tumor

Microenvironment and Tumor Suppression. Cancers 2018;10(7) (In English). DOI:

ARTN 219

10.3390/cancers10070219.

17.   Shi D, Jiang P. A Different Facet of p53 Function: Regulation of Immunity and

Inflammation During Tumor Development. Front Cell Dev Biol 2021;9:762651. DOI:

10.3389/fcell.2021.762651.

18.   Hodis E, Torlai Triglia E, Kwon JYH, et al. Stepwise-edited, human melanoma models

reveal mutations' effect on tumor and microenvironment. Science

2022;376(6592):eabi8175. DOI: 10.1126/science.abi8175.

19.   Skoulidis F, Goldberg ME, Greenawalt DM, et al. STK11/LKB1 Mutations and PD-1

Inhibitor Resistance in KRAS-Mutant Lung Adenocarcinoma. Cancer Discov

2018;8(7):822-835. DOI: 10.1158/2159-8290.CD-18-0099.

20.   Biton J, Mansuet-Lupo A, Pecuchet N, et al. TP53, STK11, and EGFR Mutations Predict

Tumor Immune Profile and the Response to Anti-PD-1 in Lung Adenocarcinoma. Clin

Cancer Res 2018;24(22):5710-5723. DOI: 10.1158/1078-0432.CCR-18-0163.

21.   Dong ZY, Zhong WZ, Zhang XC, et al. Potential Predictive Value of TP53 and KRAS

Mutation Status for Response to PD-1 Blockade Immunotherapy in Lung

Adenocarcinoma. Clin Cancer Res 2017;23(12):3012-3024. DOI: 10.1158/1078-

0432.CCR-16-2554.

22.   Leader AM, Grout JA, Maier BB, et al. Single-cell analysis of human non-small cell lung

cancer lesions refines tumor classification and patient stratification. Cancer Cell

2021;39(12):1594-1609 e12. DOI: 10.1016/j.ccell.2021.10.009.

23.   Davis AP, Cooper WA, Boyer M, Lee JH, Pavlakis N, Kao SC. Efficacy of

immunotherapy in KRAS-mutant non-small-cell lung cancer with comutations.

Immunotherapy-Uk 2021;13(11):941-952. DOI: 10.2217/imt-2021-0090.

24. Scalera S, Mazzotta M, Corleone G, et al. KEAP1 and TP53 Frame Genomic,

Evolutionary, and Immunologic Subtypes of Lung Adenocarcinoma With Different

Sensitivity to Immunotherapy. J Thorac Oncol 2021;16(12):2065-2077. DOI:

10.1016/j.jtho.2021.08.010.

25. Lavin Y, Kobayashi S, Leader A, et al. Innate Immune Landscape in Early Lung

Adenocarcinoma by Paired Single-Cell Analyses. Cell 2017;169(4):750-765 e17. DOI:

10.1016/j.cell.2017.04.014.

26. Lambrechts D, Wauters E, Boeckx B, et al. Phenotype molding of stromal cells in the

lung tumor microenvironment. Nat Med 2018;24(8):1277-1289. DOI: 10.1038/s41591-

018-0096-5.

27. Zilionis R, Engblom C, Pfirschke C, et al. Single-Cell Transcriptomics of Human and

Mouse Lung Cancers Reveals Conserved Myeloid Populations across Individuals and

Species. Immunity 2019;50(5):1317-1334 e10. DOI: 10.1016/j.immuni.2019.03.009.

28. Zhang L, Zhang Y, Wang C, et al. Integrated single-cell RNA sequencing analysis

reveals distinct cellular and transcriptional modules associated with survival in lung

cancer. Signal Transduct Target Ther 2022;7(1):9. DOI: 10.1038/s41392-021-00824-9.

29. Wang Z, Li Z, Zhou K, et al. Deciphering cell lineage specification of human lung

adenocarcinoma with single-cell RNA sequencing. Nat Commun 2021;12(1):6500. DOI:

10.1038/s41467-021-26770-2.

30. Wang Y, Liu B, Min Q, et al. Spatial transcriptomics delineates molecular features and

cellular plasticity in lung adenocarcinoma progression. Cell Discov 2023;9(1):96. DOI:

10.1038/s41421-023-00591-7.

31.   Kim N, Kim HK, Lee K, et al. Single-cell RNA sequencing demonstrates the molecular

and cellular reprogramming of metastatic lung adenocarcinoma. Nat Commun

2020;11(1):2285. DOI: 10.1038/s41467-020-16164-1.

32.   Laughney AM, Hu J, Campbell NR, et al. Regenerative lineages and immune-mediated

pruning in lung cancer metastasis. Nat Med 2020;26(2):259-269. DOI: 10.1038/s41591-

019-0750-6.

33.   Bischoff P, Trinks A, Obermayer B, et al. Single-cell RNA sequencing reveals distinct

tumor microenvironmental patterns in lung adenocarcinoma. Oncogene

2021;40(50):6748-6758. DOI: 10.1038/s41388-021-02054-3.

34.   He D, Wang D, Lu P, et al. Single-cell RNA sequencing reveals heterogeneous tumor and

immune cell populations in early-stage lung adenocarcinomas harboring EGFR

mutations. Oncogene 2021;40(2):355-368. DOI: 10.1038/s41388-020-01528-0.

35.   Alessi JV, Elkrief A, Ricciuti B, et al. Clinicopathologic and Genomic Factors Impacting

Efficacy of First-Line Chemoimmunotherapy in Advanced NSCLC. J Thorac Oncol

2023;18(6):731-743. DOI: 10.1016/j.jtho.2023.01.091.

36.   Camidge DR, Kim HR, Ahn MJ, et al. Brigatinib Versus Crizotinib in ALK Inhibitor-

Naive Advanced ALK-Positive NSCLC: Final Results of Phase 3 ALTA-1L Trial. J

Thorac Oncol 2021;16(12):2091-2108. DOI: 10.1016/j.jtho.2021.07.035.

37.   Stuart T, Butler A, Hoffman P, et al. Comprehensive Integration of Single-Cell Data. Cell

2019;177(7):1888-1902 e21. DOI: 10.1016/j.cell.2019.05.031.

38.   Wang P, Wang F, He H, et al. TP53 and CDKN2A mutations in patients with early-stage

lung squamous cell carcinoma: an analysis of the correlations and prognostic outcomes.

Ann Transl Med 2021;9(16):1330. DOI: 10.21037/atm-21-3709.

39.   Dudley AC. Tumor endothelial cells. Cold Spring Harb Perspect Med

2012;2(3):a006536. DOI: 10.1101/cshperspect.a006536.

40.   Cooke VG, LeBleu VS, Keskin D, et al. Pericyte depletion results in hypoxia-associated

epithelial-to-mesenchymal transition and metastasis mediated by met signaling pathway.

Cancer Cell 2012;21(1):66-81. DOI: 10.1016/j.ccr.2011.11.024.

41.   Biancalani T, Scalia G, Buffoni L, et al. Deep learning and alignment of spatially

resolved single-cell transcriptomes with Tangram. Nat Methods 2021;18(11):1352-1362.

DOI: 10.1038/s41592-021-01264-7.

42.   Tirosh I, Izar B, Prakadan SM, et al. Dissecting the multicellular ecosystem of metastatic

melanoma by single-cell RNA-seq. Science 2016;352(6282):189-96. DOI:

10.1126/science.aad0501.

43.   Patel AP, Tirosh I, Trombetta JJ, et al. Single-cell RNA-seq highlights intratumoral

heterogeneity in primary glioblastoma. Science 2014;344(6190):1396-401. DOI:

10.1126/science.1254257.

44.   Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell

transcriptomic data across different conditions, technologies, and species. Nature

biotechnology 2018;36(5):411.

45.   Cangul H. Hypoxia upregulates the expression of the NDRG1 gene leading to its

overexpression in various human cancers. BMC Genet 2004;5:27. DOI: 10.1186/1471-

2156-5-27.

46.   Zhang JZ, Behrooz A, Ismail-Beigi F. Regulation of glucose transport by hypoxia. Am J

Kidney Dis 1999;34(1):189-202. DOI: 10.1016/s0272-6386(99)70131-9.

47.   Cui XG, Han ZT, He SH, et al. HIF1/2alpha mediates hypoxia-induced LDHA

expression in human pancreatic cancer cells. Oncotarget 2017;8(15):24840-24852. DOI:

10.18632/oncotarget.15266.

48.   Zhang Z, Deng X, Liu Y, Liu Y, Sun L, Chen F. PKM2, function and expression and

regulation. Cell Biosci 2019;9:52. DOI: 10.1186/s13578-019-0317-8.

49.   Puram SV, Tirosh I, Parikh AS, et al. Single-Cell Transcriptomic Analysis of Primary

and Metastatic Tumor Ecosystems in Head and Neck Cancer. Cell 2017;171(7):1611-+.

(In English). DOI: 10.1016/j.cell.2017.10.044.

50.   Baron M, Tagore M, Hunter MV, et al. The Stress-Like Cancer Cell State Is a Consistent

Component of Tumorigenesis. Cell Syst 2020;11(5):536-546 e7. DOI:

10.1016/j.cels.2020.08.018.

51.   Kinker GS, Greenwald AC, Tal R, et al. Pan-cancer single-cell RNA-seq identifies

recurring programs of cellular heterogeneity. Nat Genet 2020;52(11):1208-1218. DOI:

10.1038/s41588-020-00726-6.

52.   Min JW, Kim WJ, Han JA, et al. Identification of Distinct Tumor Subpopulations in

Lung Adenocarcinoma via Single-Cell RNA-seq. PLoS One 2015;10(8):e0135817. DOI:

10.1371/journal.pone.0135817.

53.   Tang H, Wang S, Xiao G, et al. Comprehensive evaluation of published gene expression

prognostic signatures for biomarker-based lung cancer clinical studies. Ann Oncol

2017;28(4):733-740. DOI: 10.1093/annonc/mdw683.

54.   Nagy A, Munkacsy G, Gyorffy B. Pancancer survival analysis of cancer hallmark genes.

Sci Rep 2021;11(1):6047. DOI: 10.1038/s41598-021-84787-5.

55.   Xing X, Yang F, Huang Q, et al. Decoding the multicellular ecosystem of lung

1139   adenocarcinoma manifested as pulmonary subsolid nodules by single-cell RNA

1140   sequencing. Sci Adv 2021;7(5). DOI: 10.1126/sciadv.abd9738.

1141   56.   Travaglini KJ, Nabhan AN, Penland L, et al. A molecular cell atlas of the human lung

1142   from single-cell RNA sequencing. Nature 2020;587(7835):619-625. DOI:

1143   10.1038/s41586-020-2922-4.

1144   57.   Ferone G, Lee MC, Sage J, Berns A. Cells of origin of lung cancers: lessons from mouse

1145   studies. Genes Dev 2020;34(15-16):1017-1032. DOI: 10.1101/gad.338228.120.

1146   58.   LaFave LM, Kartha VK, Ma S, et al. Epigenomic State Transitions Characterize Tumor

1147   Progression in Mouse Lung Adenocarcinoma. Cancer Cell 2020;38(2):212-228 e13. DOI:

1148   10.1016/j.ccell.2020.06.006.

1149   59.   Yang L, He YT, Dong S, et al. Single-cell transcriptome analysis revealed a suppressive

1150   tumor immune microenvironment in EGFR mutant lung adenocarcinoma. J Immunother

1151   Cancer 2022;10(2). DOI: 10.1136/jitc-2021-003534.

1152   60.   Giacomelli AO, Yang X, Lintner RE, et al. Mutational processes shape the landscape of

1153   TP53 mutations in human cancer. Nat Genet 2018;50(10):1381-1387. DOI:

1154   10.1038/s41588-018-0204-y.

1155   61.   Ursu O, Neal JT, Shea E, et al. Massively parallel phenotyping of coding variants in

1156   cancer with Perturb-seq. Nat Biotechnol 2022;40(6):896-905. DOI: 10.1038/s41587-021-

1157   01160-7.

1158   62.   Baslan T, Morris JPt, Zhao Z, et al. Ordered and deterministic cancer genome evolution

1159   after p53 loss. Nature 2022;608(7924):795-802. DOI: 10.1038/s41586-022-05082-5.

63.   Marjanovic ND, Hofree M, Chan JE, et al. Emergence of a High-Plasticity Cell State

during Lung Cancer Evolution. Cancer Cell 2020;38(2):229-+. (In English). DOI:

10.1016/j.ccell.2020.06.012.

64.   Teschendorff AE, Enver T. Single-cell entropy for accurate estimation of differentiation

potency from a cell's transcriptome. Nat Commun 2017;8:15599. DOI:

10.1038/ncomms15599.

65.   Schupp JC, Adams TS, Cosme C, Jr., et al. Integrated Single-Cell Atlas of Endothelial

Cells of the Human Lung. Circulation 2021;144(4):286-302. DOI:

10.1161/CIRCULATIONAHA.120.052318.

66.   Gillich A, Zhang F, Farmer CG, et al. Capillary cell-type specialization in the alveolus.

Nature 2020;586(7831):785-789. DOI: 10.1038/s41586-020-2822-7.

67.   Muz B, de la Puente P, Azab F, Azab AK. The role of hypoxia in cancer progression,

angiogenesis, metastasis, and resistance to therapy. Hypoxia (Auckl) 2015;3:83-92. DOI:

10.2147/HP.S93413.

68.   Acevedo LM, Barillas S, Weis SM, Gothert JR, Cheresh DA. Semaphorin 3A suppresses

VEGF-mediated angiogenesis yet acts as a vascular permeability factor. Blood

2008;111(5):2674-80. DOI: 10.1182/blood-2007-08-110205.

69.   Guttmann-Raviv N, Shraga-Heled N, Varshavsky A, Guimaraes-Sternberg C, Kessler O,

Neufeld G. Semaphorin-3A and semaphorin-3F work together to repel endothelial cells

and to inhibit their survival by induction of apoptosis. J Biol Chem 2007;282(36):26294-

305. DOI: 10.1074/jbc.M609711200.

70.   Pasquale EB. Eph receptors and ephrins in cancer: bidirectional signalling and beyond.

Nat Rev Cancer 2010;10(3):165-80. DOI: 10.1038/nrc2806.

71.   Irvine AF, Waise S, Green EW, Stuart B, Thomas GJ. Characterising cancer-associated

1184   fibroblast heterogeneity in non-small cell lung cancer: a systematic review and meta-

1185   analysis. Sci Rep 2021;11(1):3727. DOI: 10.1038/s41598-021-81796-2.

1186   72.   Buechler MB, Pradhan RN, Krishnamurty AT, et al. Cross-tissue organization of the

1187   fibroblast lineage. Nature 2021;593(7860):575-579. DOI: 10.1038/s41586-021-03549-5.

1188   73.   Van de Sande B, Flerin C, Davie K, et al. A scalable SCENIC workflow for single-cell

1189   gene regulatory network analysis. Nat Protoc 2020;15(7):2247-2276. DOI:

1190   10.1038/s41596-020-0336-2.

1191   74.   Lee KW, Yeo SY, Sung CO, Kim SH. Twist1 is a key regulator of cancer-associated

1192   fibroblasts. Cancer Res 2015;75(1):73-85. DOI: 10.1158/0008-5472.CAN-14-0350.

1193   75.   De Wever O, Demetter P, Mareel M, Bracke M. Stromal myofibroblasts are drivers of

1194   invasive cancer growth. Int J Cancer 2008;123(10):2229-38. DOI: 10.1002/ijc.23925.

1195   76.   Engblom C, Pfirschke C, Pittet MJ. The role of myeloid cells in cancer therapies. Nat

1196   Rev Cancer 2016;16(7):447-62. DOI: 10.1038/nrc.2016.54.

1197   77.   Dong B, Wu C, Huang L, Qi Y. Macrophage-Related SPP1 as a Potential Biomarker for

1198   Early Lymph Node Metastasis in Lung Adenocarcinoma. Front Cell Dev Biol

1199   2021;9:739358. DOI: 10.3389/fcell.2021.739358.

1200   78.   Heimberg G, Kuo T, DePianto D, et al. Scalable querying of human cell atlases via a

1201   foundational model reveals commonalities across fibrosis-associated macrophages.

1202   bioRxiv 2023:2023.07.18.549537. DOI: 10.1101/2023.07.18.549537.

1203   79.   Casazza A, Laoui D, Wenes M, et al. Impeding macrophage entry into hypoxic tumor

1204   areas by Sema3A/Nrp1 signaling blockade inhibits angiogenesis and restores antitumor

1205   immunity. Cancer Cell 2013;24(6):695-709. DOI: 10.1016/j.ccr.2013.11.007.

80.   Zhang F, Wang H, Wang X, et al. TGF-beta induces M2-like macrophage polarization

via SNAIL-mediated suppression of a pro-inflammatory phenotype. Oncotarget

2016;7(32):52294-52306. DOI: 10.18632/oncotarget.10561.

81.   van der Leun AM, Thommen DS, Schumacher TN. CD8(+) T cell states in human

cancer: insights from single-cell analysis. Nat Rev Cancer 2020;20(4):218-232. DOI:

10.1038/s41568-019-0235-4.

82.   Litchfield K, Reading JL, Puttick C, et al. Meta-analysis of tumor- and T cell-intrinsic

mechanisms of sensitization to checkpoint inhibition. Cell 2021;184(3):596-614 e14.

DOI: 10.1016/j.cell.2021.01.002.

83.   Zhou X, Du J, Liu C, et al. A Pan-Cancer Analysis of CD161, a Potential New Immune

Checkpoint. Front Immunol 2021;12:688215. DOI: 10.3389/fimmu.2021.688215.

84.   Hollern DP, Xu N, Thennavan A, et al. B Cells and T Follicular Helper Cells Mediate

Response to Checkpoint Inhibitors in High Mutation Burden Mouse Models of Breast

Cancer. Cell 2019;179(5):1191-1206 e21. DOI: 10.1016/j.cell.2019.10.028.

85.   Gide TN, Quek C, Menzies AM, et al. Distinct Immune Cell Populations Define

Response to Anti-PD-1 Monotherapy and Anti-PD-1/Anti-CTLA-4 Combined Therapy.

Cancer Cell 2019;35(2):238-255 e6. DOI: 10.1016/j.ccell.2019.01.003.

86.   Thommen DS, Schumacher TN. T Cell Dysfunction in Cancer. Cancer Cell

2018;33(4):547-562. DOI: 10.1016/j.ccell.2018.03.012.

87.   Upadhaya S, Hubbard-Lucey VM, Yu JX. Immuno-oncology drug development forges

on despite COVID-19. Nature reviews Drug Discovery 2020.

88. Chen J, Tan Y, Sun F, et al. Single-cell transcriptome and antigen-immunoglobin analysis

reveals the diversity of B cells in non-small cell lung cancer. Genome Biol

2020;21(1):152. DOI: 10.1186/s13059-020-02064-6.

89. Yu Y, Xiao CH, Tan LD, Wang QS, Li XQ, Feng YM. Cancer-associated fibroblasts

induce epithelial-mesenchymal transition of breast cancer cells through paracrine TGF-

beta signalling. Br J Cancer 2014;110(3):724-32. DOI: 10.1038/bjc.2013.768.

90. Bonde AK, Tischler V, Kumar S, Soltermann A, Schwendener RA. Intratumoral

macrophages contribute to epithelial-mesenchymal transition in solid tumors. Bmc

Cancer 2012;12:35. DOI: 10.1186/1471-2407-12-35.

91. Ventura A, Kirsch DG, McLaughlin ME, et al. Restoration of p53 function leads to

tumour regression in vivo. Nature 2007;445(7128):661-5. DOI: 10.1038/nature05541.

92. Tam SY, Wu VWC, Law HKW. Hypoxia-Induced Epithelial-Mesenchymal Transition in

Cancers: HIF-1alpha and Beyond. Front Oncol 2020;10:486. DOI:

10.3389/fonc.2020.00486.

93. Joseph JP, Harishankar MK, Pillai AA, Devi A. Hypoxia induced EMT: A review on the

mechanism of tumor progression and metastasis in OSCC. Oral Oncol 2018;80:23-32.

DOI: 10.1016/j.oraloncology.2018.03.004.

94. Graeber TG, Osmanian C, Jacks T, et al. Hypoxia-mediated selection of cells with

diminished apoptotic potential in solid tumours. Nature 1996;379(6560):88-91. DOI:

10.1038/379088a0.

95. Tang Q, Su Z, Gu W, Rustgi AK. Mutant p53 on the Path to Metastasis. Trends Cancer

2020;6(1):62-73. DOI: 10.1016/j.trecan.2019.11.004.

96.   Capaci V, Mantovani F, Del Sal G. Amplifying Tumor-Stroma Communication: An

1250   Emerging Oncogenic Function of Mutant p53. Front Oncol 2020;10:614230. DOI:

1251   10.3389/fonc.2020.614230.

1252   97.   Pelka K, Hofree M, Chen JH, et al. Spatially organized multicellular immune hubs in

1253   human colorectal cancer. Cell 2021;184(18):4734-4752 e20. DOI:

1254   10.1016/j.cell.2021.08.003.

1255   98.   Chen JH, Nieman LT, Spurrell M, et al. Spatial analysis of human lung cancer reveals

1256   organized immune hubs enriched for stem-like CD8 T cells and associated with

1257   immunotherapy response. bioRxiv 2023. DOI: 10.1101/2023.04.04.535379.

1258   99.   Li L, Li M, Wang, X. Cancer type-dependent correlations between TP53 mutations and

1259   antitumor immunity. DNA Repair 2020;88. DOI:

1260   https://doi.org/10.1016/j.dnarep.2020.102785.

1261   100.   Assoun S, Theou-Anton N, Nguenang M, et al. Association of TP53 mutations with

1262   response and longer survival under immune checkpoint inhibitors in advanced non-small-

1263   cell lung cancer. Lung Cancer 2019;132:65-71. DOI: 10.1016/j.lungcan.2019.04.005.

1264   101.   Cortez MA, Ivan C, Valdecanas D, et al. PDL1 Regulation by p53 via miR-34. J Natl

1265   Cancer Inst 2016;108(1). DOI: 10.1093/jnci/djv303.

1266   102.   Skoulidis F, Byers LA, Diao L, et al. Co-occurring genomic alterations define major

1267   subsets of KRAS-mutant lung adenocarcinoma with distinct biology, immune profiles,

1268   and therapeutic vulnerabilities. Cancer Discov 2015;5(8):860-77. DOI: 10.1158/2159-

1269   8290.CD-14-1236.

1270   103.   Ricciuti B, Arbour KC, Lin JJ, et al. Diminished Efficacy of Programmed Death-

1271   (Ligand)1 Inhibition in STK11- and KEAP1-Mutant Lung Adenocarcinoma Is Affected

by KRAS Mutation Status. J Thorac Oncol 2022;17(3):399-410. DOI:

10.1016/j.jtho.2021.10.013.

104.   Forde PM, Spicer J, Lu S, et al. Neoadjuvant Nivolumab plus Chemotherapy in

Resectable Lung Cancer. N Engl J Med 2022. DOI: 10.1056/NEJMoa2202170.

105.   Rousseau A, Parisi C, Barlesi F. Anti-TIGIT therapies for solid tumors: a systematic

review. ESMO Open 2023;8(2):101184. DOI: 10.1016/j.esmoop.2023.101184.

106.   Jerby-Arnon L, Neftel C, Shore ME, et al. Opposing immune and genetic mechanisms

shape oncogenic programs in synovial sarcoma. Nat Med 2021;27(2):289-300. DOI:

10.1038/s41591-020-01212-6.

107.   Slyper M, Porter CBM, Ashenberg O, et al. A single-cell and single-nucleus RNA-Seq

toolbox for fresh and frozen human tumors. Nat Med 2020;26(5):792-802. DOI:

10.1038/s41591-020-0844-1.

108.   Mayakonda A, Lin DC, Assenov Y, Plass C, Koeffler HP. Maftools: efficient and

comprehensive analysis of somatic variants in cancer. Genome Res 2018;28(11):1747-

1756. DOI: 10.1101/gr.239244.118.

109.   Talevich E, Shain AH, Botton T, Bastian BC. CNVkit: Genome-Wide Copy Number

Detection and Visualization from Targeted DNA Sequencing. PLoS Comput Biol

2016;12(4):e1004873. DOI: 10.1371/journal.pcbi.1004873.

110.   inferCNV of the Trinity CTAT Project. (https://github.com/broadinstitute/inferCNV).

111.   Hao Y, Hao S, Andersen-Nissen E, et al. Integrated analysis of multimodal single-cell

data. Cell 2021;184(13):3573-3587 e29. DOI: 10.1016/j.cell.2021.04.048.

112. Liberzon A, Birger C, Thorvaldsdottir H, Ghandi M, Mesirov JP, Tamayo P. The
Molecular Signatures Database (MSigDB) hallmark gene set collection. Cell Syst
2015;1(6):417-425. DOI: 10.1016/j.cels.2015.12.004.

113. Sikkema L, Ramirez-Suastegui C, Strobl DC, et al. An integrated cell atlas of the lung in
health and disease. Nat Med 2023;29(6):1563-1577. DOI: 10.1038/s41591-023-02327-2.

114. Wu T, Hu E, Xu S, et al. clusterProfiler 4.0: A universal enrichment tool for interpreting
omics data. Innovation (N Y) 2021;2(3):100141. DOI: 10.1016/j.xinn.2021.100141.

115. Efremova M, Vento-Tormo M, Teichmann SA, Vento-Tormo R. CellPhoneDB: inferring
cell-cell communication from combined expression of multi-subunit ligand-receptor
complexes. Nat Protoc 2020;15(4):1484-1506. DOI: 10.1038/s41596-020-0292-x.

116. Gao J, Aksoy BA, Dogrusoz U, et al. Integrative analysis of complex cancer genomics
and clinical profiles using the cBioPortal. Sci Signal 2013;6(269):pl1. DOI:
10.1126/scisignal.2004088.

117. Gillette MA, Satpathy S, Cao S, et al. Proteogenomic Characterization Reveals
Therapeutic Vulnerabilities in Lung Adenocarcinoma. Cell 2020;182(1):200-225 e35.
DOI: 10.1016/j.cell.2020.06.013.

118. Chu T, Danko CG. Bayesian cell-type deconvolution and gene expression inference
reveals tumor-microenvironment interactions. 2020. DOI: 10.1101/2020.01.07.897900.

119. Wolf FA, Angerer P, Theis FJ. SCANPY: large-scale single-cell gene expression data
analysis. Genome Biol 2018;19(1):15. DOI: 10.1186/s13059-017-1382-0.

120. Palla G, Spitzer H, Klein M, et al. Squidpy: a scalable framework for spatial omics
analysis. Nat Methods 2022;19(2):171-178. DOI: 10.1038/s41592-021-01358-2.

121.   Kleshchevnikov V, Shmatko A, Dann E, et al. Cell2location maps fine-grained cell types

in spatial transcriptomics. Nat Biotechnol 2022. DOI: 10.1038/s41587-021-01139-4.

122.   Griffin GK, Weirather JL, Roemer MGM, et al. Spatial signatures identify immune

escape via PD-1 as a defining feature of T-cell/histiocyte-rich large B-cell lymphoma.

Blood 2021;137(10):1353-1364. DOI: 10.1182/blood.2020006464.

1320

1321

ACKNOWLEDGMENTS

We thank Leslie Gaffney for extensive help in editing and graphical design of the figures in the manuscript, the Broad Genomics Platform, Broad Flow Cytometry Facility, Pathology and Surgery Departments at MGH and BWH; members of the Bueno, Rodig, Hata, Regev, Johnson, and Tsankov labs. The project is part of the Human Tumor Atlas Pilot Project (HTAPP) consortium and National Institutes of Health (NIH) HTAN (Human Tumor Atlas Network) consortium paper package. Work was supported by the Klarman Cell Observatory (to O.R.-R. and A. Regev) and also funded in part by federal funds from the National Cancer Institute (NCI), NIH Task Order HHSN261100039 under Contract HHSN261201500003, and U2CCA233195. A.M.T. was supported by Department of the Army Lung Cancer Research Program Career Development Award W81XWH2210079, American Cancer Society Research Scholar Grant RSG-23-1039063-01-MM, and ISMMS seed funding. W.Z. and B.Y.S. were supported by NIH grant 2T32GM007280-41 as part of the ISMMS Medical Scientist Training Program. The graphical summary was created with Biorender.com.

Figure 1

Figure 1. Spatially resolved, multi-omic atlas of NSCLC.

A. Schematic of study design, including cohort selection (left), genomic assays profiled, sample stratification, downstream analyses, and validation platforms (right).

B. Mean $\log_2$ copy number alteration (CNA) ratio inferred using scRNA-seq malignant cells (left) and bulk WES (right) from tumor-matched samples. Common focal amplifications (bottom, green) and deletions (top, orange) in LUAD are highlighted.

C. Expression of representative markers across each annotated broad cell class.

D. Uniform Manifold Approximation and Projection (UMAP) visualization of 167,193 cells, integrated by tumor sample to minimize intertumoral heterogeneity and colored by annotated cell classes (left) and *TP53* mutational status (right).

E. Average *TP53* target gene expression score in malignant cells isolated from *TP53*[WT] (blue) and *TP53*[mut] (red) LUAD primary tumor samples.

F. Left: Proportion of endothelial cells and pericytes from scRNA-seq of *TP53*[WT] (blue) and *TP53*[mut] (red) LUAD tumors. Right: Mean $\log_{10}$ expression of highly specific endothelial and pericyte markers derived from scRNA-seq in bulk RNA-seq data from TCGA. *P*-values were calculated using a two-tailed multiple regression t-test.

G. Left: Integration of ST data with tumor-matched scRNA-seq data using Tangram allows for deconvolution of the proportion of different cell subsets. Middle: Spatial distribution of cell classes using Tangram (top), and a representative H&E stain annotated by a pathologist (bottom). Right: violin plots comparing proportion of computationally inferred cell classes (y-axis) within manually annotated categories (x-axis). Mann-Whitney-Wilcoxon test *p*-values were computed; Kruskal-Wallis FDR, $p \leq 0.05$.

H. Proportion of endothelial cells (top) and pericytes (bottom) in ST spots from $TP53^{WT}$ (blue) and $TP53^{mut}$ (red) LUAD sections. Mann-Whitney-Wilcoxon test p-values were computed.

Figure 2. Enrichment of cell cycle, hypoxia and pEMT programs accompanied by loss of AT2 identity in *TP53*[mut] LUAD malignant cells.

A. UMAP of 33,377 malignant cells integrated across tumors and colored by annotated malignant expression program subsets.

B. Dot plot of two selected markers representative of each malignant program subset.

C. Pearson correlation of malignant program scores across all malignant cells.

D. Lollipop plot showing -$\log_{10}$ *p*-values from Cox proportional-hazards regression analysis linking malignant programs to TCGA patient disease outcome, corrected for *TP53* mutational status. Sign of the y-axis corresponds to improved (positive value) or worse (negative value) outcome.

E. Comparison of mean malignant program expression scores between *TP53*[mut] and *TP53*[WT] tumors in scRNA-seq (left) and in bulk RNA-seq data from TCGA (right). *P*-values were calculated using a Mann-Whitney-Wilcoxon test (scRNA-seq, left) or a two-tailed multiple regression t-test (TCGA, right).

F. Left: Representative example of the spatial distribution of four malignant programs' expression in a *TP53*[mut] patient. Right: Bar plots comparing colocalization (spatial correlation) of pairs of malignant programs between *TP53*[mut] and *TP53*[WT] tumor sections. Mann-Whitney-Wilcoxon test *p*-values were computed.

G. Module scores for normal lung epithelial cell signatures averaged across malignant cells in each tumor, ordered by histology (first) and *TP53* mutational status (second). $\log_{10}$ SNP count for each tumor is shown on the bottom.

H. Dot plot showing association of different mutations (columns) with mean $\log_{10}$ expression of malignant programs, endothelial, and pericyte scores (rows) in TCGA LUAD. Color and size of

dots correspond to -log$_{10}$ p-value assessed by Mann-Whitney-Wilcoxon (uncorrected *TP53*, left) or a two-tailed multiple regression t-test (*EGFR*, *KRAS-G12C*, *KRAS-G12D*, *KRAS-G12V*, *STK11*, *KEAP1*, *RBM10*, *PTPRD*, *TP53*). Black outline indicates p-value ≤ 0.05, and gray outline indicates p-value of ≤ 0.1.

I. Distribution of entropy scores across non-malignant cells (grey), malignant *TP53*^{WT} cells (blue), and malignant *TP53*^{mut} cells (red) in the scRNA-seq data (left), across adjacent normal (grey), *TP53*^{WT} (blue), and *TP53*^{mut} (red) bulk transcriptomic LUAD tumor samples from TCGA (middle), and across *TP53*^{WT} (blue), and *TP53*^{mut} bulk proteomic tumor samples from CPTAC (right). *P*-values were calculated using a Mann-Whitney-Wilcoxon test.

J. Distribution of entropy scores across normal cells (grey), malignant *Kras*^{mut}/ *Trp53*^{WT} cells (K; blue), and malignant *Kras*^{mut}/*Trp53*^{mut} cells (KP; red) in K and KP mouse tumors.

K. Comparison of entropy scores between *TP53*^{WT} and different *TP53*^{mut} impact categories in A549 cells. Rightmost four violin plots show the entropy distributions of mutations also found in our cohort.

Figure 3. Spatially-resolved interactions linked to vascular depletion and stromal remodeling in *TP53*^{mut} LUAD.

A. UMAP of 15,320 endothelial cells, integrated across tumors and colored by annotated subset.

B. Comparison of aerocyte (left) and arterial (right) cell subset proportions in $TP53^{mut}$ and $TP53^{WT}$. Mann-Whitney-Wilcoxon test was used for computing p-values.

C. Dot plot of differential ligand-receptor (rows) interactions between endothelial cells and other cell classes (columns). Red indicates enrichment in $TP53^{mut}$ and blue enrichment in $TP53^{WT}$ tumor samples. Black outline indicates p-value ≤ 0.05, as assessed by Fisher's exact test.

D. Left: Spatial expression of *SEMA3A* ligand and *NRP1* receptor in a representative $TP53^{mut}$ ST sample. Right: Box plot comparing spatial correlation of *SEMA3A* and *NRP1* expression between $TP53^{WT}$ and $TP53^{mut}$ ST tumor samples. Mann-Whitney-Wilcoxon test p-value was computed.

E. UMAP of 31,868 mesenchymal cells integrated across tumors and colored by annotated subset.

F. Confusion matrix of correspondence between our *de novo* mesenchymal subset annotations and predicted identities using a classifier trained on annotated scRNA-seq data from normal lung mesenchymal cells.

G. Z-scores of the most specific SCENIC regulon activity for each mesenchymal subset.

H. Comparison of mesenchymal subset cell proportions in $TP53^{mut}$ and $TP53^{WT}$. Mann-Whitney-Wilcoxon test p-values were computed.

I. Left: Spatial expression of *TGFB2* ligand and *TGFBR2* receptor in a representative $TP53^{mut}$ ST sample. Right: Box plot comparing spatial correlation of *TGFB2* and *TGFBR2* expression between $TP53^{WT}$ and $TP53^{mut}$ ST tumor samples. Mann-Whitney-Wilcoxon test p-values were computed.

Figure 4. *SPP1* and other immunomodulatory factors are enriched in *TP53*^{mut} LUAD myeloid compartment.

A. UMAP of 18,086 myeloid cells integrated across tumors and colored by annotated subset.

B. Feature plot showing the expression of 15 representative myeloid subset markers.

C. Comparison of myeloid subset cell proportions in *TP53*^{mut} and *TP53*^{WT} scRNA-seq tumors. Mann-Whitney-Wilcoxon test *p*-values were computed.

D. Left: Comparison of *SPP1* expression in TCGA bulk RNA-seq (left) and scRNA-seq (middle) data from *TP53*^{mut} *vs.* *TP53*^{WT} primary tumor samples. Right: Horizontal bar plot of negative $\log_{10}$ *q*-values from gene set enrichment analysis of the 100 most positively correlated genes with *SPP1* in myeloid cells. *P*-values were calculated using a two-tailed multiple regression t-test (TCGA, left) or a Mann-Whitney-Wilcoxon test (scRNA-seq, middle).

E. Differential enrichment of hallmark hypoxia and EMT module score for each myeloid subset, comparing pseudobulk subset profiles for *TP53*^{mut} *vs.* *TP53*^{WT} tumors. Mann-Whitney-Wilcoxon test *p*-values were calculated.

F. Dot plot of differential ligand-receptor (rows) interactions between myeloid cells and other cell classes (columns). Red indicates enrichment in *TP53*^{mut} and blue enrichment in *TP53*^{WT} tumor samples. Black outline indicates *p*-value ≤ 0.05, as assessed by Fisher's exact test.

G. Dot plot of the expression of *CXCR3* and *CXCL11* across broad cell class annotations.

H. Differential expression analysis of ligand and receptor genes from the interactions displayed in **Figure 5F,** comparing pseudobulk myeloid subset profiles for *TP53*^{mut} *vs.* *TP53*^{WT} tumors. Mann-Whitney-Wilcoxon test *p*-values were calculated. Red indicates enrichment in *TP53*^{mut} and blue enrichment in *TP53*^{WT} tumor samples.

I. Left: Spatial expression of *SPP1* ligand and *CD44* receptor in a representative *TP53*^{mut} ST sample. Right: Box plot comparing spatial correlation of *SPP1* and *CD44* expression between *TP53*^{WT} and *TP53*^{mut} ST tumor samples. Mann-Whitney-Wilcoxon test *p*-values were computed.

Figure 5. Immune checkpoint interactions and exhausted-like lymphoid cells are enriched in *TP53*mut LUAD.

A. UMAP of 40,293 T and NK cells integrated across tumors and colored by annotated subset.

B. Feature plot showing the expression of 9 representative T and NK cell subset markers.

C. Comparison of T.Exhausted, CD8.GZMK, and CD4.TRM T cell subset proportions relative to all T and NK cells in *TP53*mut and *TP53*WT scRNA-seq tumors. Mann-Whitney-Wilcoxon test *p*-values were computed.

D. Log$_2$ mean gene expression and predicted significance of ICI-targetable ligand-receptor interactions between T cells and malignant cells (top), and T cells and myeloid (bottom) for each tumor. Horizontal bar plots (right) show negative log$_{10}$ *p*-value (Fisher's exact test) of the enrichment in interactions between *TP53*mut and *TP53*WT tumors.

E. Stacked bar plots depicting proportion of T and NK cells expressing different combinations of immune checkpoint molecules *TIGIT*, *CTLA4*, and *PDCD1* in *TP53*WT (left) and *TP53*mut (right) tumors.

F. Left: Boxplot of mean *PVR* expression in *TP53*WT (blue) and *TP53*mut (red) tumor malignant cells. Right: Correlation of cell cycle score *vs*. mean *PVR* expression in malignant cells for each patient, where blue represents *TP53*WT and red represents *TP53*mut tumors.

G. Left: Spatial expression of *PVR* and *TIGIT* in a representative *TP53*mut ST sample. Right: Box plot comparing spatial correlation of *PVR* and *TIGIT* expression between *TP53*WT and *TP53*mut tumor ST samples (right) in the tumor periphery (spots containing ≤50% malignant cells). Mann-Whitney-Wilcoxon test *p*-values were computed.

H. Stacked bar plots depicting proportion of malignant cells expressing different combination of *PVR* and *CD274* amongst all malignant cells (top) and amongst all cycling malignant cells (bottom) in $TP53^{WT}$ (left) and $TP53^{mut}$ (right) tumors.

I. Representative mIF images of $TP53^{WT}$ (left two; BWH01) and $TP53^{mut}$ (right two; BWH06) tumors with and without cytokeratin (pink). Yellow arrows correspond to $PD1^+CD8^+$ cells. Scale bar, 20 μm.

J. Box plots showing the $\log_{10}$ count density (cells/mm^2) of $PD1^+CD8^+$, total $CD8^+$, PD-$L1^+$cytokeratin$^+$, and total cytokeratin$^+$ cells across regions of interest (ROIs) in $TP53^{WT}$ (BWH01, BWH04, blue) and $TP53^{mut}$ (BWH06, BWH11, red) tumor samples used for mIF. Mann-Whitney-Wilcoxon test *p*-values were computed.

K. Grouped scatter plot showing the proportion of $PD1^+CD8^+$ cells out of all cells in proximity to PD-$L1^+$cytokeratin$^+$. Each circle represents an ROI from $TP53^{WT}$ (BWH01, BWH04, blue) *vs.* $TP53^{mut}$ (BWH06, BWH11, red) mIF tumor stains, where the size of the circle represents the proportion of PD-$L1^+$cytokeratin$^+$-neighboring cells, out of all cells within the ROI. Permutation Test *p*-values were computed.

Figure 6. Spatial multicellular CAF-TAM niche enriched in hypoxia and EMT activity in $TP53^{mut}$ LUAD.

A. Overall spatial correlation of broad cell classes across ST spots (left) and differential spatial correlation between $TP53^{mut}$ and $TP53^{WT}$ tumor samples (right). Red indicates either high colocalization (left) or colocalization enrichment in $TP53^{mut}$ (right), while blue represents either low colocalization (left) or colocalization enrichment in $TP53^{WT}$ (right) tumor samples between pairs of cell subsets.

B. Box plots comparing spatial correlation of endothelial cells with mesenchymal (left), myeloid (middle), and malignant (right) cells across spots in $TP53^{WT}$ *vs*. $TP53^{mut}$ ST tumor samples.

C. Heatmap of the 15 factors depicting multicellular programs, as calculated using NMF.

D. Spatial distribution of the enrichment of different myeloid and mesenchymal cell subsets comprising NMF7 (left) and of expression programs spatially correlated with NMF7 (right) in a representative $TP53^{mut}$ tumor ST sample (patient BWH14).

E. Box plots comparing spatial correlation of NMF7 with hallmark hypoxia score (left), hallmark EMT score (middle) and endothelial cell proportion (right) in $TP53^{WT}$ *vs*. $TP53^{mut}$ ST samples in the tumor periphery (spots with ≤50% malignant cells).

F. Graphical summary of the differences between the $TP53^{WT}$ (left) and $TP53^{mut}$ (right) LUAD TME presented in this study.

Figure S1. Integration and analysis of WES, scRNA-seq, and spatial transcriptomics data across tumors.

A. Spearman correlation shows correspondence between the $\log_2$ CNA ratio predicted by scRNA-seq and WES data from the same patient. Correlation was computed across genes for each patient. Low correspondence in two tumor samples (MGH1183 and BWH09) likely resulted from small number of malignant cells (MGH1183) and intratumoral heterogeneity (BWH09).

B. UMAP of 167,193 cells, colored by patient identity shows integration between tumor samples.

C. Stacked bar plot displaying the composition of broad cell classes for each scRNA-seq patient sample.

D. Top: Proportion of broad cell classes (not shown in **Figure 1F**) out of all cells found in the scRNA-seq data from $TP53^{WT}$ (blue) and $TP53^{mut}$ (red) LUAD tumors. Bottom: Average $\log_{10}$ expression of highly specific broad cell class markers derived from the scRNA-seq data for $TP53^{WT}$ (blue) and $TP53^{mut}$ (red) LUAD bulk RNA-seq tumor samples from TCGA. P-values were calculated using a Mann-Whitney-Wilcoxon test (scRNA-seq, top) or a two-tailed multiple regression t-test (TCGA, bottom).

E. Stacked bar plot depicting mean cell composition across ST tumor samples, predicted using Tangram mapping.

F. Spatial feature plots of broad cell classes not shown in **Figure 1G**, predicted by Tangram mapping.

G. Mean proportion of Tangram-predicted cell classes (y-axis) across different manual spot annotations independently performed by a pathologist (NRM) without knowledge of sample and spot identities.

Figure S2. Characterization of the malignant compartment in NSCLC.

A. UMAP of 33,377 malignant cells, integrated across tumors and colored by patient.

B. Dot plot of gene set enrichment outputs (Hallmark) of 100 most differentially expressed markers for each malignant subcluster.

C. Pearson correlation of average malignant program scores across tumors, analogous to **Figure 2C** but performed across patients instead of across cells.

D. Kaplan Meier survival curves showing association of malignant program expression with survival in bulk TCGA LUAD. Tumors (n = 510) were stratified by program expression into high, medium, and low expression categories. Cox proportional-hazards model was used to calculate p-value corrected for *TP53* mutational status.

E. Kaplan Meier survival curves showing association of *TP53* mutation status with survival in bulk TCGA LUAD. *P*-values were calculated using a log-rank test.

F. Comparison of $\log_{10}$ SNP and CNA abundance between *TP53*^{mut} *vs. TP53*^{WT} tumors shows no significant difference in our cohort.

G. Comparison of malignant program mean expression scores between *TP53*^{mut} and *TP53*^{WT} tumors in scRNA-seq validation cohort. *P*-values were calculated using a Mann-Whitney-Wilcoxon test.

Figure S3. Consistent changes in malignant program expression and cellular entropy in *TP53*[mut] LUAD across co-mutations, variant classes, cell lines, mouse models, bulk transcriptomic, and proteomic data.

A. Left: Stacked bar plot of proportion of tumor samples with different co-mutations in TCGA. Right: Comparison of malignant program mean expression between *TP53*[WT] and different single- or co-mutations in *EGFR*, *KRAS*, and *TP53* in bulk RNA-seq data from TCGA tumor samples.

B. Dot plot showing association of common LUAD mutations (columns) with mean $\log_{10}$ expression of malignant programs, endothelial, and pericyte scores (rows) in TCGA LUAD. Color and size of dots correspond to -log10 *p*-value assessed by Mann-Whitney-Wilcoxon (uncorrected *TP53*, left) or a two-tailed multiple regression t-test (*EGFR*, *KRAS, STK11, TP53*). Black outline indicates *p*-value ≤ 0.05, and gray outline indicates *p*-value of ≤ 0.1.

C. Left: Stacked bar plot of proportion of tumor samples with different categories of dominant-negative/loss of function *TP53*[mut] effect in TCGA. Right: Comparison of malignant program mean expression between *TP53*[WT] and different functional *TP53*[mut] categories in TCGA tumor samples. *P*-values were calculated using a Mann-Whitney-Wilcoxon test.

D. Left: Stacked bar plot of proportion of tumor samples with different impact categories in TCGA. Right: Comparison of malignant program mean $\log_{10}$ expression between *TP53*[WT] and different functional *TP53*[mut] impact categories in TCGA tumor samples. *P*-values were calculated using a Mann-Whitney-Wilcoxon test.

E. Comparison of malignant program mean expression between *TP53*[WT] and different *TP53*[mut] impact categories[59] in A549 cells. Rightmost four violin plots in each panel show the effect of mutations also found in tumor samples from our cohort.

F. Comparison of malignant program mean expression scores between $KRAS^{WT}$ and different $KRAS^{mut}$ impact categories in A549 cells.

G. Comparison of malignant program mean expression between $TP53^{mut}$ and $TP53^{WT}$ tumors using bulk proteomic data from CPTAC. *P*-values were calculated using a Mann-Whitney-Wilcoxon test.

H. Comparison of malignant program mean $\log_2$ expression between $TP53^{WT}$ and different dominant-negative/loss of function (left) and impact categories[59] (right) of $TP53^{mut}$ tumor samples in bulk proteomic data from CPTAC. *P*-values were calculated using a Mann-Whitney-Wilcoxon test.

Figure S4. Effect of different co-mutations and *TP53* mutation variants on malignant program activity and signaling entropy.

A. Comparison of malignant program scores and *Hif1a* expression between K (blue) *vs.* KP (red) LUAD mouse model scRNA-seq data malignant cells. *P*-values were calculated using a Mann-Whitney-Wilcoxon test.

B. Distribution of entropy scores across adjacent normal (grey), *TP53*^WT (blue), and different *TP53*^mut categories (reds) in bulk RNA-seq LUAD tumor samples from TCGA. *P*-values were calculated using a Mann-Whitney-Wilcoxon test.

C. Distribution of entropy scores across *TP53*^WT (blue), and different *TP53*^mut single and co-mutations (reds) in bulk RNA-seq LUAD tumor samples from TCGA. *P*-values were calculated using a Mann-Whitney-Wilcoxon test.

D. Heatmap of mean entropy score for malignant subsets in scRNA-seq data.

E. Stacked bar plot of proportion of HPCS cells from K and KP mice, classified into different malignant subsets from this study.

F. Comparison of entropy scores between *KRAS*^WT and different *KRAS*^mut impact categories in A549 cells.

Figure S5. Characterizing endothelial subsets and their differential roles in $TP53^{WT}$ and $TP53^{mut}$ LUAD.

A. Left: Comparison of mean $\log_{10}$ endothelial marker expression between $TP53^{WT}$ and different single- or co-mutations of $EGFR$, $KRAS$, and $TP53$ in bulk RNA-seq data from TCGA tumor samples. Right: Comparison of mean $\log_{10}$ endothelial marker expression between $TP53^{WT}$ and different dominant-negative/loss of function $TP53^{mut}$ variant categories in bulk RNA-seq LUAD tumor samples from TCGA. Mann-Whitney-Wilcoxon test was used for computing p-values.

B. Dot plot showing the expression of three marker genes for each annotated endothelial subset.

C. Feature plots of the expression of nine selected endothelial subset markers.

D. Stacked bar plot displaying the cell composition of endothelial subsets for each scRNA-seq patient sample.

E. Average $\log_{10}$ expression of aerocytes (left) and arterial (right) markers derived from the scRNA-seq data for $TP53^{WT}$ (blue) and $TP53^{mut}$ (red) LUAD bulk RNA-seq tumor samples from TCGA. P-values were calculated using a two-tailed multiple regression t-test.

F. Top: Dot plot of genes involved in ligand-receptor interactions displayed in **Figure 3C,** Bottom: Pseudobulk differential expression analysis performed on the same genes comparing $TP53^{mut}$ vs. $TP53^{WT}$ tumor samples. Mann-Whitney-Wilcoxon test p-values were calculated. Red indicates enrichment in $TP53^{mut}$ and blue enrichment in $TP53^{WT}$ tumor samples.

G. Left: Spatial expression of $EFHB2$ and $EFNB1$ in a representative $TP53^{mut}$ ST sample. Right: Box plot comparing spatial correlation of $EFHB2$ and $EFNB1$ between $TP53^{mut}$ and $TP53^{WT}$ ST tumor samples. Mann-Whitney-Wilcoxon test p-values were computed.

Figure S6. Mesenchymal cell compartment differences between *TP53*^mut and *TP53*^WT LUAD.

A. Dot plot showing the expression of three marker genes for each annotated mesenchymal subset.

B. Feature plot showing the expression of 12 selected mesenchymal subset markers.

C. Stacked bar plot displaying the cell composition of mesenchymal subsets for each patient sample.

D. Regulon specificity scores (RSS) plotted for three different mesenchymal subsets of interest.

E. Spatial correlation (colocalization) of mesenchymal subsets with other broad cell classes across all ST tumor samples. Red indicates high colocalization and blue low colocalization between pairs of cell subsets.

F. Average $\log_{10}$ expression of CAF.ADH1B markers derived from the scRNA-seq data for *TP53*^WT (blue) and *TP53*^mut (red) LUAD bulk RNA-seq tumor samples from TCGA. *P*-values were calculated using a two-tailed multiple regression t-test.

G. Left: Comparison of mean $\log_{10}$ pericyte marker expression between *TP53*^WT and different single- or co-mutations of *EGFR*, *KRAS*, and *TP53* in bulk RNA-seq data from TCGA tumor samples. Right: Comparison of mean $\log_{10}$ pericyte marker expression between *TP53*^WT and different dominant-negative/loss of function *TP53*^mut variant categories in bulk RNA-seq LUAD tumor samples from TCGA. Mann-Whitney-Wilcoxon test was used for computing *p*-values.

H. Dot plot of differential ligand-receptor (rows) interactions between mesenchymal cells and other cell classes (columns). Red indicates enrichment in *TP53*^mut and blue enrichment in *TP53*^WT tumor samples. Black outline indicates *p*-value ≤ 0.05, as assessed by Fisher's exact test.

I. Top: Dot plot of genes involved in ligand-receptor interactions displayed in **Figure S3H.** Bottom: Pseudobulk differential expression analysis performed on the same genes comparing

TP53$^{\text{mut}}$ *vs*. *TP53*$^{\text{WT}}$ tumor samples. Mann-Whitney-Wilcoxon test *p*-values were calculated. Red

indicates enrichment in *TP53*$^{\text{mut}}$ and blue enrichment in *TP53*$^{\text{WT}}$ tumor samples.

Figure S7. Characterization of myeloid subsets and expression of ligand-receptor genes.

A. Stacked bar plot displaying the cell composition of myeloid subsets for each patient sample.

B. Comparison of *CXCL9/10/11* expression in *TP53*^mut *vs*. *TP53*^WT LUAD tumor samples using bulk RNA-seq data from TCGA.

C. Left: Proportion of TAM.CXCLs subset relative to all myeloid cells in *TP53*^WT (blue) *TP53*^mut (red) LUAD scRNA-seq tumor samples. Right: Average $\log_{10}$ expression of TAM.CXCLs markers in LUAD bulk RNA-seq tumor samples from TCGA. *P*-values were calculated using a Mann-Whitney-Wilcoxon test (scRNA-seq, left) or a two-tailed multiple regression t-test (TCGA, right).

D. Comparison of *SPP1*, *CXCL9/10/11* gene expression among *TP53*^WT and different single- or co-mutations in *EGFR*, *KRAS*, and *TP53*.

E. Comparison of gene expression among dominant-negative/loss of function (left), and impact categories (right) of *TP53*^mut LUAD tumor samples in TCGA. *P*-values were calculated using a Mann-Whitney-Wilcoxon test.

F. Dot plot of gene set enrichments using the Hallmark database for the 40 most differentially expressed genes in each myeloid subsets, comparing genes up in *TP53*^WT (left) and up in *TP53*^mut (right).

G. Kaplan-Meier survival curve showing association between *SPP1* expression and overall survival in bulk RNA-seq LUAD data from TCGA. Tumors (n = 510) were stratified into high or low expression of *SPP*1. Cox proportional-hazards model was used to calculate a *p*-value corrected for *TP53* mutational status.

H. Horizontal bar plot showing the most correlated genes with *SPP1* expression in myeloid cells, ranked by Pearson's correlation coefficient.

I. Dot plot showing expression of genes involved in ligand-receptor interactions shown in **Figure 4F.**

Figure S8. Lymphoid immune checkpoint gene expression and spatial distribution.

A. Dot plot showing the expression of two representative marker genes for each annotated T and NK subset.

B. UMAP of 18,520 B, plasma, mast, and plasmacytoid DC cells, integrated across tumors and colored by annotated subset.

C. Comparison of T.Exhausted (left; validation cohort only) and TFH cell proportions (right; discovery, validation, and combined cohorts) relative to all T and NK cells between $TP53^{mut}$ and $TP53^{WT}$ tumors profiled by scRNA-seq. *P*-values were calculated using a Mann-Whitney-Wilcoxon test.

D. Average $\log_{10}$ expression of specific T.Exhausted, and CD8.GZMK markers derived from the scRNA-seq data in $TP53^{WT}$ (blue) and $TP53^{mut}$ (red) LUAD bulk RNA-seq tumor samples from TCGA. *P*-values were calculated using a two-tailed multiple regression t-test.

E. Left: Dot plot of genes involved in ligand-receptor analysis displayed in **Figure 5D**. Right: Pseudobulk differential expression analysis performed on the same genes comparing $TP53^{mut}$ *vs*. $TP53^{WT}$ tumor samples. Mann-Whitney-Wilcoxon test *p*-values were calculated. Red indicates enrichment in $TP53^{mut}$ and blue enrichment in $TP53^{WT}$ tumor samples.

F. $\log_{10}$ expression of *CD274*, *CD86*, *PVR*, *PDCD1, CTLA4,* and *TIGIT* in $TP53^{WT}$ (blue) and $TP53^{mut}$ (red) LUAD bulk RNA-seq tumor samples from TCGA. *P*-values were calculated using a two-tailed multiple regression t-test.

G. Boxplots of PVR and CD274 (PD-L1) protein expression in CPTAC tumor samples split by *TP53* mutational status. Mann-Whitney-Wilcoxon test *p*-values were computed.

H. Representative mIF images of additional *TP53*[WT] (BWH04) and *TP53*[mut] (BWH11) tumor stains. Yellow arrows correspond to PD1$^+$CD8$^+$ cells and white arrows correspond to PD-L1$^+$cytokeratin$^+$ cells. Scale bar, 50 μm.

Figure S9. Spatial organization and cellular landmarks in *TP53*^mut and *TP53*^WT LUAD.

A. Mean spatial correlation of broad cell classes across ST spots averaged across *TP53*^WT (top) and *TP53*^mut (bottom) samples.

B. Mean (top) and differential (bottom) spatial correlation heatmap of lymphoid (B, plasma, T, NK) subsets with other broad cell classes between *TP53*^mut and *TP53*^WT. Red indicates either high overall colocalization (top) or colocalization enrichment in *TP53*^mut (bottom) while blue represents either low overall colocalization (top) or colocalization enrichment in *TP53*^WT (bottom) tumor samples between pairs of cell subsets.

C. Line plot showing proportion of malignant, myeloid, and mesenchymal cells ordered from NMF7 low spots to NMF7 high spots (x-axis).

Table S1. Clinical data for patient cohort

Table S2. High level cell class markers and subset markers

Table S3. Clinical data for validation cohort