Global Diversity in Mammalian Life Histories: Environmental Realms and Evolutionary Adaptations

E. Beccari[1*], P. Capdevila Lanzaco[2,3], R. Salguero-Gómez[4,5], & C. Pérez Carmona[1]

[1] Department of Botany, Institute of Ecology and Earth Sciences, University of Tartu, J. Liivi 2, 50409 Tartu, Estonia

[2] Departament de Biologia Evolutiva, Ecologia i Ciències Ambientals, Facultat de Biologia, Universitat de Barcelona (UB), Barcelona 08028, Spain

[3] Institut de Recerca de la Biodiversitat (IRBio), Universitat de Barcelona (UB), Barcelona 08028, Spain

[4] Department of Biology, University of Oxford, 11a Mansfield Rd, Oxford, OX1, 3SZ, UK

[5] Evolutionary Demography laboratory, Max Plank Institute for Demographic Research, Konrad Zuße Straße1, Rostock 18057, Germany

*Author for correspondence: Eleonora Beccari (eleonora.beccari@ut.ee)

Statement of authorship: P.C.L., R.S-G., and C.P.C. conceived the core ideas behind this paper, all authors provided fundamental inputs for its final development. E.B collected all data and performed the literature search needed to sort species in environmental realms. E.B. analysed the data with fundamental inputs from C.P.C., P.C.L, and R.G-S. All authors contributed to interpretation of the results. E.B. and C.P.C. led the writing of the manuscript which was edited by all authors.

Keywords: Brain mass; ecology; functional diversity; life-history; mode of life

Abstract

Mammalian life history strategies can be characterized by a few axes of variation, which conform a space where species are positioned according to which life history strategies are favoured in the environment they exploit. Yet, we still lack global descriptions of the diversity of realized mammalian life history and of how this diversity is shaped by the environment. We used six life history traits to build a global life history space and explored how major environmental realms (land, air, water) influence mammalian life history strategies. We demonstrate that realms are tightly linked to distinct life history strategies. Predominantly, aquatic and aerial species adhere to slower life history strategies, while terrestrial species tend to exhibit faster life histories. Highly encephalized terrestrial species are a notable exception to these patterns. In addition, species transitioning between the terrestrial and aquatic realms, such as seals, show intermediate life history strategies. Further, different mode-of-life may play a significant role in allowing to expand the set of strategies exploitable in the terrestrial realm. Our results provide compelling evidence of the link between environmental realms and the diversity of life history strategies among mammals.

Introduction:

The diversity of mammalian adaptations, ranging from the fast and semelparous life histories of antechinus (Braithwaite & Crowther 2008) to the slow and long lives of bowhead whales (Perrin *et al.* 2009), have allowed mammals to occupy all land (Burgin *et al.* 2018; Grossnickle *et al.* 2019), exploit air (Babich Morrow *et al.* 2021; Barclay 1994; Maina 2000), and even return to the sea (Davis 2019). Despite this wide ecological radiation (Grossnickle *et al.* 2019), evolutionary history and energetic constraints limit the set of life history strategies (*i.e.,* combinations of life history traits) that are viable in nature (Grime & Pierce 2012; Southwood 1988; Stearns 1992). These limits restrict mammalian investments in survival, growth, and reproduction to two major dimensions of life history trait variation

(Carmona *et al.* 2021b; Gaillard *et al.* 1989; Healy *et al.* 2019; Oli 2004). The first dimension

generally encompasses the time scale of life, referred to as the fast-slow continuum (Bielby *et*

al. 2007; Capdevila *et al.* 2020; Gaillard *et al.* 1989; Healy *et al.* 2019; Stearns 1992). In

contrast, the second dimension depends more on the subset of species and traits considered,

but generally reflects the timing and intensity of a species' reproductive investment (Bielby *et*

al. 2007; Capdevila *et al.* 2020; Gaillard *et al.* 1989; Healy *et al.* 2019). While these two life

history dimensions are consistently observed across studies with various species and trait

subsets (Bielby *et al.* 2007; Capdevila *et al.* 2020; Gaillard *et al.* 1989), we still lack a

comprehensive understanding of how mammals are organized within this broad life history

space. This gap hampers exploring and quantifying realized mammalian life history diversity,

preventing a thorough comprehension of similarities and differences among mammalian

species (Carmona *et al.* 2021b; Pinsky *et al.* 2022).

Environmental conditions profoundly influence species' life history strategies (Grime &

Pierce 2012; Southwood 1988). Variable environments, for instance, affect the pace of life of

species (Tuljapurkar *et al.* 2009) selecting for delayed life cycles and longer lives compared

to constant and stable environments (Grime & Pierce 2012; Tuljapurkar *et al.* 2009). Because

life history strategies are selected by similar environmental pressures (Grime & Pierce 2012;

Southwood 1988), there is a highly redundant occupation of the life history space regardless

of taxonomic affiliation (Carmona *et al.* 2021b; Cooke *et al.* 2019; Cox *et al.* 2021).

However, mammal's widespread distribution across major environmental realms (*i.e.*, land,

water, air), begs the question of whether mammalian life history strategies are equally

represented across these three realms.

Environmental realms are characterized by distinct physical dimensions, structural properties,

and environmental requirements (Pinsky *et al.* 2022; Steele 1985; Webb 2012).

Consequently, species inhabiting each specific environmental realm are under unique eco-

physiological constraints that are not experienced by species from other realms (Pinsky *et al.*

2019; Webb 2012). For instance, the energetically expensive nature of the aquatic and aerial

realms (Barclay 1994; Gearty *et al.* 2018; Goldbogen 2018; Guigueno *et al.* 2019; Maina

2000) has led to longer lives in aerial species (Babich Morrow *et al.* 2021; Gaillard *et al.*

1989) or long age at female maturity in the aquatic realm (Davis 2019; Perrin *et al.* 2009).

Accordingly, inhabiting different environmental realms may select distinct sets of life history

strategies (Capdevila *et al.* 2020; Pinsky *et al.* 2022). Yet, some similarities do exist across

realms. For example, the capacities for climbing trees or burrowing on the ground offer

partial escape routes from the environmental constrains unique to terrestrial living, selecting

for generally longer lifespans compared to other terrestrial species (Bels & Russell 2023;

Healy *et al.* 2014, 2019; Mincer & Russo 2020; Withers *et al.* 2016). Similarly, larger brains

generally allows species to avoid hazardous situations, selecting for slower lives compared to

other terrestrial representatives (Benson-Amram *et al.* 2016; Bertrand *et al.* 2022; Dembitzer

et al. 2022; González-Lagos *et al.* 2010; Seyfarth & Cheney 2002). It becomes clear how a

comprehensive understanding of mammalian adaptations across various environmental

realms is crucial for unravelling the complexities of life history strategies. Without this

understanding, we will remain ill-equipped to reveal overarching ecological patterns or to

understand the driving forces and consequences of ongoing global changes (Pinsky *et al.*

2022).

Here, we use six life history traits representing investments in survival, growth, and

reproduction for 3,438 mammalian species (53.7% of currently recognized living mammals,

Burgin *et al.* 2018) extracted from the Amniote dataset (Myhrvold *et al.* 2015) to examine the

influence of major environmental realms on mammalian life history strategies. In agreement

with previous research (Bielby *et al.* 2007; Gaillard *et al.* 1989), we first show that two axes

of variation relating to the fast-slow continuum and the reproductive strategies continuum are sufficient to explain a large proportion of mammalian life history strategies (Fig. 1a). Using this plane as a springboard, we proceed to test three hypotheses: (H1) species' position in the life history space will associate with the environmental realm they occupy, as a consequence the different eco-physiological pressures acting among major realms (Grime & Pierce 2012; Pinsky *et al.* 2022; Tuljapurkar *et al.* 2009; Webb 2012). Specifically, we expect aerial species to exhibit slower life history strategies compared to terrestrial species, and aquatic species to occupy areas of the life history space skewed toward higher iteroparity (Capdevila *et al.* 2020). (H2) Because species transitioning between aquatic and terrestrial environments (*i.e.,* semi-aquatic species) experience a blend of selective pressures, they will occupy intermediate positions in the life history space between these two realms (Hood 2020; Perrin *et al.* 2009). Additionally, (H3) within the terrestrial realm, species with arboreal, semi-arboreal and fossorial mode-of-life will display slower life history strategies than species confined on land.

Methods

Data collection and trait imputation

To characterise the mammals' life history strategies, we used the Amniote database which comprises information for ca. 18,300 species of mammals, reptiles, and birds (Myhrvold *et al.* 2015). We extracted mammalian species (N = 4,953), and selected the life history traits with the most complete information (*i.e.,* traits with records for at least 1,000 species): litter size (ls, number of offspring per reproductive event), litter per year (ly, number of litters per year), age at female maturity (fmat, time needed for female individuals to reach maturity in days), weaning length (wea, time needed to wean the offspring in days), gestation length

(gest, time passing between conception and birth in days), longevity (long, individual lifespan

in years) and adult body mass (bm, in grams). For finer definition and information of traits'

measurements please refer to Myhrvold et al. (2015).

Because no trait was fully informed for all species, we imputed missing life history trait

values. Recent studies have shown that imputation of individual traits allows to accurately

characterize the position of species in trait spaces that are the result of combining different

correlated traits (see Carmona, Tamme, et al. 2021; Carmona, Bueno, et al. 2021 for

examples). Additionally, recent simulations indicate that examining patterns in space

occupation built considering imputed species, produces results much closer to real patterns in

space occupation compared to considering only species with complete trait information

(Stewart *et al.* 2023). Because accounting for evolutionary relationships improves the

accuracy of the imputation (Penone *et al.* 2014), we downloaded mammalian phylogenies

from Vertlife database (Upham *et al.* 2019). We selected the complete fossil-based set of

10,000 phylogenies and computed a single maximum clade credibility tree (MCC tree) using

the *maxCladeCred* function of *'phangorn'* R package (Schliep 2011). The 135 species absent

in the phylogeny but with life history information were added at the genus' root using

add.species.to.genus function of *'phytools'* R package (Revell 2012). Species that were

missing in the life history traits database were pruned using *drop.tip* function of *'phyltools'* R

package.

For the imputation procedure, we selected mammalian species with information for body size

and at least one of the other considered life history traits (*i.e.,* ls, ly, fmat, wea, gest, long),

resulting in a subset of 3,438 species. Within this subset 1,293 species had complete trait

measurements (c.a. 37.6% of the total). Then, following Carmona, Tamme, et al. 2021, we

included phylogenetic information in the imputation procedure using the first 10 phylogenetic

eigenvectors derived from the MCC tree. For the imputation, we used the *missForest* function

from the *'missForest'* R package(Stekhoven 2022). After the imputation, we estimated the

reliability of the imputation procedure (Supplementary methods 1). The imputation procedure

efficiently retrieved the real position of a species, with an average error of 2.8% of the range

for the first component and 1.7% for the second component. Imputations errors were

consistent across orders with different level of data completeness and coverage.

The final procedure resulted in a dataset of 3,438 mammalian species (53.7% of currently

recognized living mammals, Burgin *et al.* 2018) from 29 orders in which 37.6% of traits

information (1,293 species) derived from empirical measurement and 62.4% (2,145 species)

were imputed.

Ascribing species to environmental realms and mode-of-life

In order to have information on species adscription to the terrestrial, aerial, and aquatic

realms, we downloaded information on the environment exploited by each mammal from the

Worms (WoRMS Editorial Board 2020) and IUCN (IUCN 2017) databases using the R

packages *'worms'*(Holstein 2018) and *'redlist'* (Chamberlain 2020), respectively. We

classified mammalian species as "terrestrial" (*i.e.,* land realm) in case of unambiguous

classification from both databases, whereas we limited "aerial" category (air) to powered

flight species (*i.e.,* Chiroptera order). Although there is no doubt about the classification of

"fully" aquatic species (*i.e.*, species that spent their entire life in water, *e.g.,* dolphins, whales,

dugongs, Perrin *et al.* 2009), there is not a strict classification of semi-aquatic species (Hood

2020). Due to this, the Worms and IUCN databases classification did not match for many

species. For instance, the rhesus monkey, *Macaca mulatta*, is classified as semi-aquatic

according to the Worms database, while IUCN classifies it as terrestrial. To solve this

problem, we performed a finer classification of aquatic species by summarizing species'

dependency on water. First, we selected all species classified as freshwater and marine in at

least one of the IUCN and Worms databases. Then, we created an Aquatic Dependency Index

(AD Index) to classify species as "fully" aquatic, "fully" terrestrial, or semi-aquatic. We built

the AD Index answering a series of binary questions capable of summarizing the amount of

time spent in water during fundamental phases of species life histories (*e.g.,* birth, mating;

Supplementary methods 3). The AD Index was used to refine aquatic classification by

relabelling species scoring zero in the AD Index (*i.e.,* no important parts of life are strictly

constrained by water presence) as terrestrial species. Then, we classified as aquatic (hereafter

aquatic) all species scoring 5 out of a maximum of 5 on the AD Index (*i.e.,* species that spent

their whole life in water, like dolphins), while we classified as semi-aquatic all species

scoring between 1 and 4 (Supplementary methods 3). Out of the 3,438 mammalian species

considered in our analyses, 2,634 species were classified as terrestrial (76.6% of total set of

species), 661 species were classified as aerial (19.2% of the total), 85 species were classified

as aquatic (2.5% of the total), and 58 species were classified as semi-aquatic (*i.e.,* transitional

species, 1.7% of the total).

We further classified terrestrial species according to their arboreal, semiarboreal, and

fossorial mode-of-life by using the same classification applied in Santini, *et al.* 2022. We

manually added missing information throughout a literature review(Bels & Russell 2023;

IUCN 2017; University of Michigan 2022). Out of the 2,634 species classified as terrestrial,

595 species were classified as arboreal (22.6% of terrestrial species), 175 species were

classified as semi-arboreal (6.6% of terrestrial species), 117 species were classified as

fossorial (4.4% of terrestrial species), and 1,747 species were classified as 'fully' terrestrial

(*i.e.,* confined on land, 66.3% of terrestrial species).

Species taxonomies across all previously mentioned data tables (*i.e.,* life history traits,

phylogeny, Worms, IUCN, and mode-of-life databases) were standardized using *gnr_resolve*

function of the *'taxize'* R package (Chamberlain & Szöcs 2013) resolving for the Global

Biodiversity Information Facility (GBIF) Backbone Taxonomy.

Construction of the global life history traits space

Body mass is a strong allometric constraint that affects the values of many other life history

traits (Bielby *et al.* 2007; Jeschke & Kokko 2009; Sibly & Brown 2007). Mammals' body

mass can span from a few grams of shrews to tons for whales, and the aquatic realm strongly

selects for bigger sizes compared to the terrestrial realm (Gearty *et al.* 2018). Therefore, to

address life history strategies without confounding effects from different orders of magnitude,

we accounted for the effect of body mass before computing the mammalian life history space.

For this, we performed a series of ordinary linear regressions between single log-transformed

life history traits and log-transformed species' adult body mass. Then, we extracted body

mass–corrected residuals for each species from each regression (Jeschke & Kokko 2009;

Revell 2009). Body mass–corrected residuals were further centred, scaled, and used as size-

corrected life history traits (Supplementary material Fig. 1).

Mapping species' position in multidimensional spaces based on trait information allows the

identification of major axes of trait variation (Bueno *et al.* 2023; Díaz *et al.* 2016), as well as

summarizing species' ecological strategies. We constructed the mammalian life history space

by performing a principal component analysis (PCA) on the complete set of mammalian

species. We retained the first two axes which together account for 75.1% of variance in

mammalian life history traits.

Then we estimated the reliability of our methodological choices by comparing this life

history space (*i.e.,* the scores of species in the selected number of principal components) with

three other spaces: one built considering only complete set of records (*i.e.,* non-imputed data;

N = 1,293; Supplementary material Fig. 2); one built considering the imputed dataset without

body mass correction (N = 3,438); and one built considering the same body mass–corrected

residuals used to build the life history space (N = 3,438) but accounting for the phylogenetic

history of species (Supplementary material Fig. 3; Supplementary material Table 1). In all

abovementioned cases, the comparisons revealed strong correspondence between spaces,

demonstrating that our methodological choices do not influence the inferred relationships

between life history traits (Supplementary Methods 2).

Exploring patterns within the global life history space

We derived life history structure (*i.e.,* the patterns of organization of species in the life history

space) using the trait probability density (TPD) framework (Carmona *et al.* 2016), which

computes species' probabilistic distribution in the life history space using multivariate kernel

density estimation. The kernel density estimation for each species was estimated as a

multivariate normal distribution cantered at the species' coordinates in the PCA space

(Carmona *et al.* 2016, 2019), and with a standard deviation computed using the unconstrained

bandwidth selectors from the *'Hpi'* function in the *'ks'* package (Duong 2007, 2022). To

estimate the life history structure of the whole set of mammals, all species' kernels were

weighted and aggregated to form a single continuous TPD function across the whole

mammalian life history space. The value of the TPD function at any given point of the life

history space reflects the density of species in that particular area of the life history space

(Carmona *et al.* 2016, 2019). We graphically represented the so-built mammalian life history

structure by highlighting the contours containing 50% and 99% of the total probabilistic

distribution of species (Fig. 1a). We used the *'TPD'* R package (Carmona 2019; Carmona *et*

al. 2019).

We used mammalian global structure to test whether mammals distribute in the life history

space following a random assembly of life history traits (*i.e.,* life history strategies are

randomly selected) or whether certain life history strategies are more prevalent than others

(Carmona *et al.* 2021a; Díaz *et al.* 2016; Southwood 1988). First, we computed functional

richness (the amount of functional space occupied (Carmona *et al.* 2016; Mason *et al.* 2005)

at increasing probability thresholds (*i.e.,* from 0.1% to 99.9% of the TPD function, using

0.1% increments). In this way, we estimated a "profile" of the probabilistic distribution of

species, reflecting what amount of the life history space is occupied at different probability

thresholds. We compared the observed functional richness profile with a null model (999

repetitions) considering a bivariate normal distribution of species with the same mean and

variance as the observed data (see Carmona, Bueno et al., 2021). This analysis was performed

by sampling 3,438 simulated species (*i.e.,* complete set of mammals) from a bivariate normal

distribution with the same mean and covariance of observed data calculated using the

'*rmvnorm*' function of the R package '*tmvtnorm*'(Wilhelm & Manjunath 2010). In the same

way as done for the observed data, for each repetition of the null model we computed life

history structure and measured functional richness at increasing probability thresholds. We

then compared the observed and null functional richness profiles; functional richness values

lower than expected under the null model would suggest that the global set of mammals

occupy a smaller portion of the life history space than expected under a normal distribution

and vice-versa.

Then, using the same null model, we tested whether the portions of life history space that are

mainly exploited by mammalian species are closer or farther than expected from the centre of

the distribution. For this, we estimated functional divergence (*i.e.*, degree to which species'

density is skewed towards the extremes of the life history space) considering 99% of the total

species' probability (Carmona *et al.* 2016; Mason *et al.* 2005) for both observed and

simulated data. High functional divergence would reveal that species are segregated towards

the edges of the life history space, whereas low divergence would indicate that species are

clustered in the centre of the life history space (Carmona *et al.* 2021a).

We estimated the standardized effect size (SES) of both the functional profiles (at each

probability threshold) and the functional divergence measures as: SES = (observed value –

mean (simulated values))/standard deviation (simulated values). SES values express the

number of SD units by which the observed value deviates from the mean of the simulated

ones. A SES value lower than 0 indicates that the observed metric is smaller than the average

of the simulated values and vice versa. Finally, we tested the statistical significance of

observed and simulated differences by estimating two-sided p-values computed by

confronting the SES values with a cumulative normal distribution with mean $= 0$ and

standard deviation $= 1$ (Carmona *et al.* 2021a, Supplementary material Fig.4).

Lastly, to have a clearer picture of how different species are positioned in the life history

space, we computed TPD functions for each mammalian order with at least 6 species (total of

18 orders). We graphically represented the mammalian life history structure of each order by

highlighting the contours containing 50% and 99% of the total probabilistic distribution of

species order (Supplementary material Fig. 5).

Exploring environmental realms' patterns in the life history space

We produced realm-specific TPD functions by aggregating the species that are characteristic

of each realm. We followed the same procedure explained above for the global set of

mammalian species. In the same way as done with the complete mammalian life history

structure, we represented graphically the TPD for each realm and highlighted contours

containing 50% and 99% of the total probability (Fig. 1b-d).

We compared the structure of each of the environmental realms (*i.e.,* air, land, water and

transitional) considering multiple aspects. First, we compared realms in terms of the amount

of life history space that they occupy (*i.e.,* their functional richness). Because functional

richness is known to be positively related to the number of species present in a group, we

built a null model (999 repetitions) randomizing species assignment (Carmona *et al.* 2021a).

For each realm, we randomly extracted the same number of observed species from the

complete mammalian database and computed TPD functions of the random community. We

computed functional richness at the 99% of total probability for both observed and

randomized structures and calculated SES as explained above. Under this null model,

negative SES values would indicate that the species present in the realm are more clustered in

the life history space (*i.e.,* occupy a smaller portion of the life history space, and therefore

species are redundant in their life history strategy) than expected given the same number of

randomly selected species and vice versa. A highly clustered realm would indicate high

redundancy of life history strategies (Carmona *et al.* 2021a), suggesting that the prevalent

conditions in that realm exert a strong selection on life history strategies. This procedure

allows to compare functional richness among realms with very different number of species

(Supplementary material Table 2a).

Then, we statistically tested whether dissimilarities in species positioning were a product of a

realm-driven selection of life history strategies. We estimated how much of the total variation

in the position of the species on life history space was explained by difference in realms. For

this, we first calculated the dissimilarities (using Euclidean distances) between all pairs of

species in the life history space. Then, we performed a PERMANOVA analysis considering

the three major environmental realms (*i.e.,* air, land, and water) as the explanatory variable (R

package *'vegan'* (Oksanen *et al.* 2020)). If the realm explains a large portion of life history

variation, it means that life history variability is mainly related to differences between realms

rather than differences among species within the realm (Carmona *et al.* 2021a).

We estimated pairwise dissimilarities in life history structures between pairs of realms (*i.e.,*

air, land, water, and transitional, Fig. 2). Dissimilarities were calculated as 1 - overlap of the

corresponding TPD functions producing measures varying between 1, indicating a complete

overlap between the considered TPD functions, and 0, when the two functions are completely

disjoint. TPD-based dissimilarities consider both differences in the boundaries and

differences in the density of the occupation of the life history space (Carmona *et al.* 2021b;

Germain *et al.* 2023; Toussaint *et al.* 2021). These dissimilarities can be further decomposed

into two complementary components: one quantifying up to what point the life history

differences between realms are related to the occupation of exclusive regions of the space

(*i.e.,* turnover) and another reflecting how differently the two realms occupy the shared parts

of the life history space (*i.e.,* nestedness Carmona *et al.* 2019).

We tested whether the observed dissimilarities between environmental realms were

significantly higher than expected by chance by means of null models (Micó *et al.* 2020;

Traba *et al.* 2015). In this null model, for each pair of realms, we considered only the set of

species belonging to both realms and we randomized the realm classification. We maintained

the same number of species observed in each considered realm and avoided species

assignment to multiple realms. For each repetition of the null model (n = 999), we estimated

the TPD functions of each realm and estimated their dissimilarity, as well as its nestedness

and turnover components. Finally, we computed SES values for all mentioned metrics as

described above (Supplementary material Table 2b).

Lastly, we explored whether dissimilarities between environmental realms were maintained

also in closely related species. To do this, we computed TPD functions for orders having

species inhabiting in different realms. Because all aerial species belong to the Chiroptera

order, we decided to focus on orders having both terrestrial and aquatic species. We extracted

orders that had at least six species in both the aquatic (in this case identified by both aquatic

and semi-aquatic species) and terrestrial realms. For each of these orders, we calculated a life

history structure for terrestrial species and one for aquatic species. We represented

graphically structures highlighting 50 and 99% of the total probability and we computed

pairwise dissimilarities between the terrestrial and aquatic life history structure of each order

(Supplementary material Fig. 6).

Extracting differences in terrestrial mode-of-life and species life histories

For the terrestrial realm, we further produced specific TPD functions for each of the

considered mode-of-life (*i.e.*, arboreal, semiarboreal, fossorial, and confined on land)

following the same procedure as above. We represented graphically the TPD for each mode-

of-life and highlighted contours containing 50% and 99% of the total probability

(Supplementary material Fig. 7). As above, we computed observed functional richness for

each mode-of-life, and we performed a null model to extract SES values of functional

richness (Supplementary material Table 3a). Then, we estimated pairwise dissimilarities in

life history structures across all different mode-of-life and environmental realms (*i.e.*, aerial,

aquatic, transitional) following the same procedure as above (Supplementary material Fig. 8).

We tested whether the observed dissimilarities were significantly higher than expected by

chances by performing the same null model as done above for environmental realms

(Supplementary material Table 3b).

We tested whether, within taxonomically related species, exploiting the environment with

different mode-of-life allows for distinct set of life history strategy than species confined on

land. For that, we selected all mammalian orders showing arboreal, semiarboreal, or fossorial

species and species confined on land. Within each order, we extracted the centroid of the

functional structure of the species confined on land as well as the centroid for the functional

structure of the mode-of-life exploited in the order (Fig. 4). Then, we produced three datasets

(one for arboreality, one for semi-arboreality, and one for fossoriality), containing the

centroids the behaviours together with the ones of species confined on land for each of the

considered order. We used these datasets to statistically test for difference between species

confined on land and species showing different mode-of-life through a series of linear mixed

effect models using the R package *'lmerTest '* (Kuznetsova *et al.* 2017). For each dataset we

performed two models, one for each principal component, using the mean score of the order

along the PCs as response variables, mode-of-life as fixed effect, and mammalian orders as

random intercept (Supplementary material Table 4).

Similarly to arboreal, semiarboreal, and fossorial mode-of-life, species with high

encephalisation are capable to exploit the environment in a plastic way, avoiding hazards and

stressful situations (González-Lagos *et al.* 2010; Zhu *et al.* 2023). Consequently, we further

explored whether high encephalization (Barton & Capellini 2011; González-Lagos *et al.*

2010; Pérez☐Barbería *et al.* 2007; Reader & Laland 2002) allows to exploit portion of the

life history space otherwise not successful for the terrestrial realm (Supplementary methods

4). The final procedure resulted in two generalized additive models (GAMs) mapping species

observed brain mass (Barton & Capellini 2011; Benson-Amram *et al.* 2016; brain mass

information were extracted from Combine database Soria *et al.* 2021, for a subset of 1,054

species with non-imputed brain mass observations) in response to the species position in the

life history space (Supplementary material Fig. 9).

All statistical analyses were performed using R version 4.0.3 (R Core Team 2020).

Results & Discussion

Global life history strategies are constrained and profoundly divergent

Mammalian life history traits vary mostly along a two-dimensional space, which suggests a

strong trait covariation limiting viable life history strategies. The first two principal

components (PC) of the PCA account for 75.1% of life history variation, conforming a life

history space characterised by two primary trade-offs: litter size *vs.* gestation length, and

number of litters per year *vs.* age at female maturity and weaning length (Fig. 1a). The

configuration of this space is line with previous studies on smaller sets of species (267

species in Bielby et al., 2007; 80 species in Gaillard et al., 1989). The first dimension of this

space (PC1) captures the fast-slow continuum (Stearns 1992), distinguishing fast species with

short lives, gestation and weaning, early maturity, and frequent litters, from slow species with

fewer and infrequent litters, longer lives, and later maturity (Fig. 1a). The second dimension

(PC2) reflects lifetime reproductive effort, illustrating the trade-off between litter size and

frequency, as well as offspring quality (*i.e.,* altricial *vs.* precocial), and reproductive

frequency (*i.e.,* semelparity *vs.* iteroparity). Species' positioning and life history trait

correlations in the life history space remain consistent after accounting for evolutionary

history (Procrustes test r = 0.95; p < 0.001 for species; angles correlation r = 0.85, p < 0.001

for traits; see Supplementary methods 2), suggesting that this two-dimensional life history

strategy space applies to all mammals, independent of shared ancestry (Supplementary

material Fig. 3).

Mammalian species exhibit a remarkably uneven, concentrated distribution within the life

history space. Comparing global mammal occupation patterns along this space to null models

representing bivariate normal distributions with identical mean and covariance (Carmona *et*

al. 2021a), the occupied space is smaller than expected by chance (Supplementary material

Fig. 4a). For example, at the 99% quantile of species probability distribution, observed

functional richness (*i.e.*, amount of space occupied by species) is 32% smaller than the

simulated counterpart ($SES_{Fric99\%}$ = -11.79, p < 0.001, n = 999 iterations). Additionally,

species' distribution is much more divergent than the simulated (SES_{Fdiv} = 18.79, p < 0.001; n

= 999 iterations; Extended data Fig.4b), primarily due to species clustering in two hotspots at

opposite ends of the fast-slow continuum. Our findings reveal a dichotomy in successful

investments to survival, development, and reproduction of global mammals: species investing

on quantity of reproductive investment contrast with those prioritizing quality (Grime &

Pierce 2012; Southwood 1988; Stearns 1992). Species in the "fast hotspot" (Fig. 1), such as

mice and rabbits, optimize fast turnover and high reproductive output, enabling them to

propagate their genes in unpredictable and unfavourable environments (Grime & Pierce

2012; Southwood 1988). Conversely, "slow hotspot" species (Fig. 1a), including primates,

cetaceans and bats, typically make costly parental investments to equip their few offspring

with advantageous adaptations for long-term persistence and environmental exploitation

(Barton & Capellini 2011; Grime & Pierce 2012; Southwood 1988). This bimodal pattern

mirrors that of plant traits (Carmona *et al.* 2021a; Salguero-Gómez *et al.* 2016), suggesting

that, across the tree of life, species tend to cluster around a few contrasting syndromes that

lead to successful strategies (Junker *et al.* 2022).

Divergent life history strategies correspond to different environmental realms

Major environmental realms select for distinct and highly redundant life history strategies

(Fig. 1b-d; Fig. 2). This finding is confirmed by a PERMANOVA analysis, which attributes

35% of variation in life history traits to differences among realms. The marked differences in

life history strategies among realms are further demonstrated by the dissimilarities between

pairs of realms ($D_{land-water}$ = 69%, $D_{land-air}$ = 86%, $D_{air-water}$ = 86%; p < 0.001 in all cases; Fig. 2;

Supplementary material Table 2), as well as by the high levels of species redundancy within

individual realms (*i.e.,* species within an environmental realm tend to have similar trait

values; $R_{land} = 7.99$, $R_{air} = 9.25$, $R_{water} = 4.38$; $p < 0.001$ in all cases; Supplementary material

Table 2). Terrestrial species disproportionally occupy the "fast hotspot" (Fig. 1c), including

representatives of marsupials as well as the Carnivora, Lagomorpha, Rodentia, and

Eulipotyphla orders (Supplementary material Fig. 5). Primates and Perissodactyla are the

only exceptions to this pattern since they cluster in the slowest portion of the life history

space. On the contrary, aerial (Fig. 1a) and aquatic species (Fig. 1d) are confined to the slow

end of the spectrum. Despite the similar pace of life of aquatic and aerial species, their

reproductive strategies differ substantially. Aerial species exhibit long gestation times, while

aquatic species display late female maturity and extended weaning (Fig. 1a). The

predominance of slow strategies in aquatic species contrasts with previous observations

mainly based on baleen whales (Gaillard *et al.* 1989), which are among the longest-lived

aquatic species (Fig. 1d). Our study, however, includes dolphins, displaying a more complete

picture. Dolphins and baleen whales exhibit distinct migratory, dietary, and social habits

(Perrin *et al.* 2009), influencing resource perception and gathering within realms and

affecting life history variation (Famoso *et al.* 2018; Soriano-Redondo *et al.* 2020; Zhu *et al.*

2023).

Despite marked differences and redundancy among realms, characteristic strategies of a given

environmental realm can be observed in others. This shared set of life history strategies is

particularly evident in the case of the aquatic realm, fully contained within the life history

space occupied by terrestrial species (Nestedness$_{land-water}$ = 1, Fig. 2b). The aerial realm is also

highly overlapping with the terrestrial one (Nestedness$_{land-air}$ = 0.89; Fig. 2a); primates, a

small subset of terrestrial species, are the main reason for such an overlap. For example, apes

(including humans), and dolphins have essentially the same life history traits, whereas

terrestrial Lemuriformes and Tarsius species are located in the part of the life history space

occupied by bats (Figure 1; Supplementary material Fig. 5).

Species transitioning between realms have intermediate life history strategies

The life history strategies of semi-aquatic species provide further evidence for the profound influence of environment realms on life history variation. Semi-aquatic species exhibit intermediate life history strategies, slower than terrestrial species but faster than fully aquatic ones (Fig. 3). This pattern holds within semi-aquatic mammals, whereby species occupy different regions of the life history space depending on their degree of water-dependence. For example, species with greater aquatic involvement, such as seals and walruses (Davis 2019; Perrin *et al.* 2009), occupy portions of the life history space similar to fully aquatic species like manatees, beluga, or sperm whales. Conversely, species with lesser water-dependence, like rodents and otters (Davis 2019; Hood 2020; Perrin *et al.* 2009), are located in the hotspot typical of terrestrial mammals (Fig. 3). Interestingly, redundancy of semi-aquatic mammals was not significantly different from what expected by chance ($R_{semi-aquatic}$ = -0.89; p = 0.27; Supplementary material Table 2). We advocate for future research to investigate the evolution of aquatic and semi-aquatic adaptations, in order to understand the realized diversity of the mammalian return to water (Davis 2019).

The reported intermediate life history strategy location of semi-aquatic mammals is independent of taxonomic order. This robustness confirms the association between environmental realms and life history strategies even among closely related species. For example, terrestrial and semi-aquatic carnivores separate in the life history space following the same patterns seen in the global dataset with terrestrial carnivores being faster than their semi-aquatic counterparts (Supplementary material Fig. 6a-c). Moreover, the sea otter (*Enhydra lutris*), the only fully aquatic carnivore, exhibits a markedly different life history strategy compared to semi-aquatic otters (Supplementary material Fig. 6). Among

480 Cetartiodactyla, aquatic (dolphins, whales) and semi-aquatic (hippos) species have slower life

481 history strategies than terrestrial species (giraffe, bison, gazelles; Supplementary material Fig.

482 6d-f). Altogether, these findings suggest that species transitioning between aquatic and

483 terrestrial realms possess intermediate life history strategies.

Slow life history strategies in arboreal and fossorial species

Arboreal, semiarboreal, and fossorial mode-of-life are associated with distinct life history

strategies than land-confided species. Species confined on land mainly clustered in the fast

portion of the life history space, coinciding with the "fast hotspot" of mammalian global

structure (Supplementary material Fig. 7c). Conversely, arboreal, semiarboreal, and fossorial

species mainly cluster towards the slow (arboreal, Supplementary material Fig. 7a) and

intermediate (fossorial and semiarboreal; Supplementary material Fig. 7b-c) portions of the

life history space. These discrepancies between land-confined species and those using

specific mode-of-life to dwell in the terrestrial realm are confirmed by dissimilarities across

pairwise comparisons ($D_{land-arboreal} = 61\%$, $D_{land-semiarboreal} = 47\%$, $D_{land-fossorial} = 53\%$; $p < 0.001$

in all cases) and hold consistent across taxonomically related species (Fig. 4). Arboreal and

fossorial species are significantly skewed towards the slowest portion of the life history space

in comparison to representatives of the same order confined on land. Conversely, the pattern

for semiarboreal species is less clear and fluctuates among orders (Fig. 4, Supplementary

material Table 4), suggesting a role of arboreal (and fossorial) specialization in the selection

of slow life history strategies on land.

Arboreal and semiarboreal adaptations expand the portion of the life history space occupied

by the terrestrial realm towards life history strategies that are otherwise exclusive of the aerial

and aquatic realms. The overlap we detected between terrestrial realm and both aerial and

aquatic realms (Fig. 2) is primarily related to the overlap with arboreal and, to a lesser extent,

semiarboreal life history strategies (Supplementary material Fig. 8). Indeed, terrestrial and

aerial nestedness decreased by 55% (increasing the dissimilarity between the two realms by

10%) when considering only terrestrial species confined on land despite these species being

66.3% of terrestrial species. The dissimilarity between aquatic and land-confined species

increases by 15% if compared to the dissimilarity computed considering the full set of

terrestrial adaptation (Supplementary material Table 3b). Although with different extent,

arboreal, semiarboreal, and fossorial mode-of-life, allow species to escape mortality risks as

well as to explore different ecological opportunities (Bels & Russell 2023; Scheffers *et al.*

2017; Stroud & Losos 2016; Withers *et al.* 2016). Similarly to the aquatic and aerial realms,

the access of vertical movement may allow species to modify how the environmental

pressures are perceived in the terrestrial realms allowing for different life history strategies

otherwise unfavoured on land (Greenslade 1983; Grime & Pierce 2012).

Arboreal species exploit life history strategies that are characteristic of the aquatic realm

(Supplementary material Fig. 8). Interestingly, dissimilarities between arboreal and aquatic

species are lower than the one observed between arboreal and land-constrained species

($D_{water-arboreal}$ = 55%, $D_{land-arboreal}$ = 61%, p < 0.001). Accordingly, the most prevalent aquatic

life history strategies (50% of the probability distribution) are completely nested within

arboreal ones, overlapping in the portion of the life history space occupied by arboreal

primates and dolphins (Fig. 1,2a; Supplementary material Fig. 5,8). This shared set of life

history strategies among primates and dolphins, might be linked with complex social

structures and high encephalization in both groups (Perrin *et al.* 2009; Ridgway *et al.* 2017;

Worthy & Hickie 1986). Similarly to arboreal, semiarboreal, and fossorial mode-of-life, high

encephalization, which generally relates to both social and behavioural facets (Barton &

Capellini 2011; Pérez☐Barbería *et al.* 2007), allow species to exploit the environment in a

plastic way, avoiding hazards and stressful situations (González-Lagos *et al.* 2010; Zhu *et al.*

2023). We tested whether high encephalization may play a role in selecting terrestrial

strategies, by modelling the brain-mass ratio of terrestrial mammalian species (i.e. the ratio

between brain dimension and body mass (Benson-Amram *et al.* 2016; Worthy & Hickie

1986) as a function of species' position in the life history space (Supplementary methods 4).

We found a robust association between larger brain mass and slow life histories ($R^2 = 0.48$,

Supplementary material Fig. 9), suggesting that larger brain size may significantly influence

the diversity of terrestrial life history strategies. Consequently, while the environmental realm

in which species live seems to drive life history strategies, behavioural characteristics of

species seem to play a role in expanding the set of life history strategies exploitable in the

terrestrial realm (Cox *et al.* 2021). Specifically, we suggest that vertical movement (*e.g.*

arboreality, fossoriality) as well as plastic or social behaviours may allow species to modify

the environmental pressures perceived under specific environmental constraints (Greenslade

1983; Grime & Pierce 2012; Southwood 1988), playing a fundamental role in defining the set

of life history strategies realized under similar environmental requirements.

Conclusion

Our analyses of over 3,400 mammal species reveal that major environmental realm is a

strong predictor of mammalian life history strategies. Aerial (Babich Morrow *et al.* 2021;

Gaillard *et al.* 1989) and aquatic (Davis 2019; Gearty *et al.* 2018; Uhen 2007) realms require

specific adaptations, such as flight-capable skeletal structures (Barclay 1994; Maina 2000) or

enhanced thermoregulatory abilities for aquatic life (Davis 2019; Perrin *et al.* 2009). These

adaptations slow down and prolong lifespans in ways seemingly unexploited in the terrestrial

realm by mammals (Babich Morrow *et al.* 2021). However, despite each realm's unique set

of life history strategies, all of them overlapped to some degree within the life history space

due to species' adaptations that enable the exploitation of otherwise unsuccessful strategies in

each realm. Empirical research has shown that adaptations in the way species exploit the

environment (Babich Morrow *et al.* 2021; Healy *et al.* 2014; Shattuck & Williams 2010), as

well as social (Zhu *et al.* 2023), migratory (Soriano-Redondo *et al.* 2020), dietary

characteristics (Famoso *et al.* 2018), and cerebral development (Barton & Capellini 2011;

Seyfarth & Cheney 2002) can influence species' life histories. Consequently, variations in

these behavioural and ecological characteristics modulate species' responses to

environmental pressures (Grime & Pierce 2012) and expand the spectrum of life history

strategies available for exploitation within a realm, as observed for activity patterns (Cox *et*

al. 2021). For instance, differences in migratory, dietary, and social habits among baleen

whales and dolphins (Perrin *et al.* 2009) may facilitate distinct life history strategies in

aquatic environments, accounting for the rapid pace of life we observed in baleen whales.

Here we demonstrate that arboreal, semi-arboreal and fossorial mode-of-life play a

fundamental role in expanding the set of life history strategies in the terrestrial realms.

Confirming empirical evidence (Babich Morrow *et al.* 2021; Bels & Russell 2023; Healy *et*

al. 2014; Shattuck & Williams 2010), arboreal and fossorial mode-of-life favour slower life

history strategies within taxonomically related species, with arboreal mode-of-life allowing

terrestrial species to share life history strategies dominated by aquatic species. Similarly, the

strong relationship we uncovered between brain dimension and the position of terrestrial

species within the life history space (Supplementary material Fig. 9) suggest a role of

sociality and encephalization in allowing terrestrial mammals to exploit the slower end of the

life history spectrum. Accordingly, we advocate for future research to disentangle the

influence of different ecological and behavioural adaptations on life history variation among

terrestrial and aquatic mammals.

Our study provides a comprehensive assessment of global mammalian life history strategies,

revealing the fundamental role of environmental realm in shaping these patterns.

Remarkably, these patterns persist even among closely related species, highlighting the

profound relationship between environmental realms and life history evolution. Our findings

not only provide empirical insights into mammalian adaptation, but also establish a

foundation for further exploration of ecological and evolutionary processes (Burgin *et al.*

2018; Grossnickle *et al.* 2019). In addition, this foundation may allow to improves our

understanding of the drivers and consequences of ongoing global changes (Carmona *et al.*

2021b; Enquist *et al.* 2020) and future biodiversity patterns (Carmona *et al.* 2021b; Cooke *et*

al. 2019; Toussaint *et al.* 2021). As we confront these challenges, our study emphasises the

importance of understanding the intricate interplay between environment, ecology, and

evolution to disentangle, anticipate, and mitigate the impacts on mammalian life history

diversity.

589 Acknowledgements

590 C.P.C and EB were supported by the Estonian Research Council grant (PSG293) and the

591 European Regional Development Fund via the Mobilitas Pluss programme (MOBERC40).

592 P.C.L. was supported by the European Union-Next Generation EU Maria Zambrano Program

593 (ZAMBRANO 21-26). R. S-G. was supported by a NERC Independent Research fellowship

594 (NE/M018458/1).

595 Competing interests

596 All authors declare no competing interests.

597 References

Babich Morrow, C., Ernest, S.K.M. & Kerkhoff, A.J. (2021). Macroevolution of

dimensionless life-history metrics in tetrapods. *Philos. Trans. R. Soc. B: Biol. Sci.*,

288, 20210200.

Barclay, R.M.R. (1994). Constraints on Reproduction by Flying Vertebrates: Energy and

Calcium. *Am. Nat.*, 144, 1021–1031.

Barton, R.A. & Capellini, I. (2011). Maternal investment, life histories, and the costs of brain

growth in mammals. *Proc. Natl. Acad. Sci. U.S.A.*, 108, 6169–6174.

Bels, V.L. & Russell, A.P. (Eds.). (2023). *Convergent Evolution: Animal Form and Function*.

Fascinating Life Sciences. Springer International Publishing, Cham.

Benson-Amram, S., Dantzer, B., Stricker, G., Swanson, E.M. & Holekamp, K.E. (2016).

Brain size predicts problem-solving ability in mammalian carnivores. *Proc. Natl.*

Acad. Sci. U.S.A., 113, 2532–2537.

Bertrand, O.C., Shelley, S.L., Williamson, T.E., Wible, J.R., Chester, S.G.B., Flynn, J.J., *et*

al. (2022). Brawn before brains in placental mammals after the end-Cretaceous

extinction. *Science*, 376, 80–85.

Bielby, J., Mace, G.M., Bininda☐Emonds, O.R.P., Cardillo, M., Gittleman, J.L., Jones, K.E.,

et al. (2007). The Fast☐Slow Continuum in Mammalian Life History: An Empirical

Reevaluation. *Am. Nat.*, 169, 748–757.

Braithwaite, R.W. & Crowther, M. (2008). Brown antechinus Antechinus stuartii. In: *The*

mammals of Australia. Reed New Holland, pp. 94–96.

Bueno, C.G., Toussaint, A., Träger, S., Díaz, S., Moora, M., Munson, A.D., *et al.* (2023).

Reply to: The importance of trait selection in ecology. *Nature*, 618, E31–E34.

Burgin, C.J., Colella, J.P., Kahn, P.L. & Upham, N.S. (2018). How many species of

mammals are there? *J. Mammal.*, 99, 1–14.

Capdevila, P., Beger, M., Blomberg, S.P., Hereu, B., Linares, C. & Salguero-Gómez, R. (2020). Longevity, body dimension and reproductive mode drive differences in aquatic versus terrestrial life-history strategies. *Funct. Ecol.*, 34, 1613–1625.

Carmona, C.P. (2019). *TPD: Methods for Measuring Functional Diversity Based on Trait Probability Density*. Available at: https://CRAN.R-project.org/package=TPD. Last accessed .

Carmona, C.P., Bello, F. de, Mason, N.W.H. & Lepš, J. (2016). Traits Without Borders: Integrating Functional Diversity Across Scales. *Trends Ecol. Evol.*, 31, 382–394.

Carmona, C.P., de Bello, F., Mason, N.W.H. & Lepš, J. (2019). Trait probability density (TPD): measuring functional diversity across scales based on TPD with R. *Ecology*, 100, e02876.

Carmona, C.P., Bueno, C.G., Toussaint, A., Träger, S., Díaz, S., Moora, M., *et al.* (2021a). Fine-root traits in the global spectrum of plant form and function. *Nature*, 597, 683–687.

Carmona, C.P., Tamme, R., Pärtel, M., Bello, F. de, Brosse, S., Capdevila, P., *et al.* (2021b). Erosion of global functional diversity across the tree of life. *Sci. Adv.*

Chamberlain, S. (2020). *rredlist: "IUCN" Red List Client*. Available at: https://CRAN.R-project.org/package=rredlist. Last accessed .

Chamberlain, S.A. & Szöcs, E. (2013). *taxize: taxonomic search and retrieval in R. F1000Research, 2:191*. Available at: https://f1000research.com/articles/2-191/v2. Last accessed .

Cooke, R.S.C., Eigenbrod, F. & Bates, A.E. (2019). Projected losses of global mammal and bird ecological strategies. *Nat. Commun.*, 10, 2279.

Cox, D.T.C., Gardner, A.S. & Gaston, K.J. (2021). Diel niche variation in mammals associated with expanded trait space. *Nat. Commun.*, 12, 1753.

Davis, R.W. (2019). *Marine mammals: adaptations for an aquatic life*. Springer Cham.

Dembitzer, J., Castiglione, S., Raia, P. & Meiri, S. (2022). Small brains predisposed Late

Quaternary mammals to extinction. *Sci Rep*, 12, 3453.

Díaz, S., Kattge, J., Cornelissen, J.H.C., Wright, I.J., Lavorel, S., Dray, S., *et al.* (2016). The

global spectrum of plant form and function. *Nature*, 529, 167–171.

Duong, T. (2007). ks: Kernel Density Estimation and Kernel Discriminant Analysis for

Multivariate Data in R. *J. Stat. Softw.*, 21, 1–16.

Duong, T. (2022). *ks: Kernel Smoothing*. Available at: https://CRAN.R-

project.org/package=ks. Last accessed .

Enquist, B.J., Abraham, A.J., Harfoot, M.B.J., Malhi, Y. & Doughty, C.E. (2020). The

megabiota are disproportionately important for biosphere functioning. *Nat. Commun.*,

11, 699.

Famoso, N.A., Hopkins, S.S.B. & Davis, E.B. (2018). How do diet and body mass drive

reproductive strategies in mammals? *Biol. J. Linn.*, 124, 151–156.

Gaillard, J.-M., Pontier, D., Allainé, D., Lebreton, J.D., Trouvilliez, J. & Clobert, J. (1989).

An Analysis of Demographic Tactics in Birds and Mammals. *Oikos*, 56, 59–76.

Gearty, W., McClain, C.R. & Payne, J.L. (2018). Energetic tradeoffs control the size

distribution of aquatic mammals. *Proc. Natl. Acad. Sci. U.S.A.*, 115, 4194–4199.

Germain, R.R., Feng, S., Buffan, L., Carmona, C.P., Chen, G., Graves, G.R., *et al.* (2023).

Changes in the functional diversity of modern bird species over the last million years.

Proc. Natl. Acad. Sci. U.S.A., 120, e2201945119.

Goldbogen, J.A. (2018). Physiological constraints on marine mammal body size. *PNAS*, 115,

3995–3997.

González-Lagos, C., Sol, D. & Reader, S.M. (2010). Large-brained mammals live longer. *j.*

Evol. Biol., 23, 1064–1074.

Greenslade, P.J.M. (1983). Adversity Selection and the Habitat Templet. *The American*

673   *Naturalist*, 122, 352–365.

674   Grime, J.P. & Pierce, S. (2012). *The evolutionary strategies that shape ecosystems*. John

675   Wiley & Sons.

676   Grossnickle, D.M., Smith, S.M. & Wilson, G.P. (2019). Untangling the Multiple Ecological

677   Radiations of Early Mammals. *Trends Ecol. Evol.*, 34, 936–949.

678   Guigueno, M.F., Shoji, A., Elliott, K.H. & Aris-Brosou, S. (2019). Flight costs in volant

679   vertebrates: A phylogenetically-controlled meta-analysis of birds and bats.

680   *Comparative Biochemistry and Physiology Part A: Molecular & Integrative*

681   *Physiology*, 235, 193–201.

682   Healy, K., Ezard, T.H.G., Jones, O.R., Salguero-Gómez, R. & Buckley, Y.M. (2019). Animal

683   life history is shaped by the pace of life and the distribution of age-specific mortality

684   and reproduction. *Nat. Ecol. Evol.*, 3, 1217–1224.

685   Healy, K., Guillerme, T., Finlay, S., Kane, A., Kelly, S.B.A., McClean, D., *et al.* (2014).

686   Ecology and mode-of-life explain lifespan variation in birds and mammals. *Proc.*

687   *Natl. Acad. Sci. U.S.A.*, 281, 20140298.

688   Holstein, J. (2018). *worms: Retriving Aphia Information from World Register of Marine*

689   *Species*. Available at: https://CRAN.R-project.org/package=worms. Last accessed .

690   Hood, G.A. (2020). *Semi-aquatic Mammals: Ecology and Biology*. JHU Press.

691   IUCN. (2017). *The IUCN Red List of Threatened Species. Version 3*. Available at:

692   https://www.iucnredlist.org. Last accessed 23 September 2020.

693   Jeschke, J.M. & Kokko, H. (2009). The roles of body size and phylogeny in fast and slow life

694   histories. *Evol Ecol*, 23, 867–878.

695   Junker, R.R., Albrecht, J., Becker, M., Keuth, R., Farwig, N. & Schleuning, M. (2022).

696   Towards an animal economics spectrum for ecosystem research. *Funct. Ecol.*, 37.

Kuznetsova, A., Brockhoff, P.B. & Christensen, R.H.B. (2017). *lmerTest Package: Tests in*

Linear Mixed Effects Models. *Journal of Statistical Software*.

Maina, J.N. (2000). What it takes to fly: the structural and functional respiratory refinements

in birds and bats. *J. Exp. Biol.*, 203, 3045–3064.

Mason, N.W.H., Mouillot, D., Lee, W.G. & Wilson, J.B. (2005). Functional richness,

functional evenness and functional divergence: the primary components of functional

diversity. *Oikos*, 111, 112–118.

Micó, E., Ramilo, P., Thorn, S., Müller, J., Galante, E. & Carmona, C.P. (2020). Contrasting

functional structure of saproxylic beetle assemblages associated to different

microhabitats. *Sci. Rep.*, 10, 1520.

Mincer, S.T. & Russo, G.A. (2020). Substrate use drives the macroevolution of mammalian

tail length diversity. *Proceedings of the Royal Society B: Biological Sciences*, 287,

20192885.

Myhrvold, N.P., Baldridge, E., Chan, B., Sivam, D., Freeman, D.L. & Ernest, S.K.M. (2015).

An amniote life-history database to perform comparative analyses with birds,

mammals, and reptiles. *Ecology*, 96, 3109–3109.

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., *et al.* (2020).

vegan: Community Ecology Package. Available at: https://CRAN.R-

project.org/package=vegan. Last accessed .

Oli, M.K. (2004). The fast–slow continuum and mammalian life-history patterns: an

empirical evaluation. *Basic Appl. Ecol.*, 5, 449–463.

Penone, C., Davidson, A.D., Shoemaker, K.T., Di Marco, M., Rondinini, C., Brooks, T.M., *et*

al. (2014). Imputation of missing data in life-history trait datasets: Which approach

performs the best? *Methods Ecol. Evol.*, 5, 961–970.

Pérez‐Barbería, F.J., Shultz, S. & Dunbar, R.I.M. (2007). Evidence for coevolution of

sociality and relative brain size in three orders of mammals. *Evolution*, 61, 2811–

2821.

Perrin, W.F., Würsig, B. & Thewissen, J.G.M. (2009). *Encyclopedia of marine mammals*.

Elsevier Inc.

Pinsky, M.L., Comte, L. & Sax, D.F. (2022). Unifying climate change biology across realms

and taxa. *Trends in Ecology & Evolution*, 37, 672–682.

Pinsky, M.L., Eikeset, A.M., McCauley, D.J., Payne, J.L. & Sunday, J.M. (2019). Greater

vulnerability to warming of marine versus terrestrial ectotherms. *Nature*, 569, 108–

111.

R Core Team. (2020). R: A Language and Environment for Statistical Computing.

Reader, S.M. & Laland, K.N. (2002). Social intelligence, innovation, and enhanced brain size

in primates. *Proc. Natl. Acad. Sci. U.S.A.*, 99, 4436–4441.

Revell, L.J. (2009). Size-Correction and Principal Components for Interspecific Comparative

Studies. *Evolution*, 63, 3258–3268.

Revell, L.J. (2012). phytools: an R package for phylogenetic comparative biology (and other

things). *Methods Ecol. Evol.*, 3, 217–223.

Ridgway, S.H., Carlin, K.P., Van Alstyne, K.R., Hanson, A.C. & Tarpley, R.J. (2017).

Comparison of Dolphins' Body and Brain Measurements with Four Other Groups of

Cetaceans Reveals Great Diversity. *Brain Behav. Evol.*, 88, 235–257.

Salguero-Gómez, R., Jones, O.R., Jongejans, E., Blomberg, S.P., Hodgson, D.J., Mbeau-

Ache, C., *et al.* (2016). Fast–slow continuum and reproductive strategies structure

plant life-history variation worldwide. *Proceedings of the National Academy of*

Sciences, 113, 230–235.

Santini, L., Benítez-López, A., Dormann, C.F. & Huijbregts, M.A.J. (2022). Population

density estimates for terrestrial mammal species. *Global Ecology and Biogeography*,

31, 978–994.

Scheffers, Brett.R., Shoo, L., Phillips, B., Macdonald, S.L., Anderson, A., VanDerWal, J., *et*

al. (2017). Vertical (arboreality) and horizontal (dispersal) movement increase the

resilience of vertebrates to climatic instability. *Glob. Ecol. Biogeogr.*, 26, 787–798.

Schliep, K.P. (2011). phangorn: phylogenetic analysis in R. *Bioinformatics*, 27, 592–593.

Seyfarth, R.M. & Cheney, D.L. (2002). What are big brains for? *Proc. Natl. Acad. Sci.*

U.S.A., 99, 4141–4142.

Shattuck, M.R. & Williams, S.A. (2010). Arboreality has allowed for the evolution of

increased longevity in mammals. *PNAS*, 107, 4635–4639.

Sibly, R.M. & Brown, J.H. (2007). Effects of body size and lifestyle on evolution of mammal

life histories. *Proc. Natl. Acad. Sci. U.S.A.*, 104, 17707–17712.

Soria, C.D., Pacifici, M., Di Marco, M., Stephen, S.M. & Rondinini, C. (2021). COMBINE: a

coalesced mammal database of intrinsic and extrinsic traits. *Ecology*, 102, e03344.

Soriano-Redondo, A., Gutiérrez, J.S., Hodgson, D. & Bearhop, S. (2020). Migrant birds and

mammals live faster than residents. *Nat. Commun.*, 11, 5719.

Southwood, T.R.E. (1988). Tactics, Strategies and Templets. *Oikos*, 52, 3–18.

Stearns, S.C. (1992). *The evolution of life histories*. Oxford university press Oxford.

Steele, J.H. (1985). A comparison of terrestrial and marine ecological systems. *Nature*, 313,

355–358.

Stekhoven, D.J. (2022). *missForest: Nonparametric Missing Value Imputation using Random*

Forest. R package version 1.5. Available at: https://www.r-project.org. Last accessed .

Stewart, K., Carmona, C.P., Clements, C., Venditti, C., Tobias, J.A. & González-Suárez, M. (2023). Functional diversity metrics can perform well with highly incomplete data sets. *Methods Ecol. Evol.*

Stroud, J.T. & Losos, J.B. (2016). Ecological Opportunity and Adaptive Radiation. *Annu. Rev. Ecol. Evol. Syst.*, 47, 507–532.

Toussaint, A., Brosse, S., Bueno, C.G., Pärtel, M., Tamme, R. & Carmona, C.P. (2021). Extinction of threatened vertebrates will lead to idiosyncratic changes in functional diversity across the world. *Nat Commun*, 12, 5162.

Traba, J., Morales, M.B., Carmona, C.P. & Delgado, M.P. (2015). Resource partitioning and niche segregation in a steppe bird assemblage. *Community Ecol.*, 16, 178–188.

Tuljapurkar, S., Gaillard, J.-M. & Coulson, T. (2009). From stochastic environments to life histories and back. *Philos. Trans. R. Soc. B: Biol. Sci.*, 364, 1499–1509.

Uhen, M.D. (2007). Evolution of marine mammals: Back to the sea after 300 million years. *Anat.*, 290, 514–522.

University of Michigan. (2022). *Animal Diversity Web (ADW)*. Available at: https://animaldiversity.org. Last accessed .

Upham, N.S., Esselstyn, J.A. & Jetz, W. (2019). Inferring the mammal tree: Species-level sets of phylogenies for questions in ecology, evolution, and conservation. *PLOS Biol.*, 17, e3000494.

Webb, T.J. (2012). Marine and terrestrial ecology: unifying concepts, revealing differences. *Trends Ecol. Evol.*, 27, 535–541.

Wilhelm, S. & Manjunath, B., G. (2010). tmvtnorm: A Package for the Truncated Multivariate Normal Distribution. *R J.*, 2, 25.

Withers, P.C., Cooper, C.E., Maloney, S.K., Bozinovic, F. & Cruz-Neto, A.P. (2016). Physiological Adaptations to Extreme Environments. In: *Ecological and*

Environmental Physiology of Mammals (eds. Withers, P.C., Cooper, C.E., Maloney,

S.K., Bozinovic, F. & Cruz Neto, A.P.). Oxford University Press, p. 0.

WoRMS Editorial Board. (2020). *World Register of Marine Species*. Available at:

https://www.marinespecies.org. Last accessed 1 September 2020.

Worthy, G.A.J. & Hickie, J.P. (1986). Relative Brain Size in Marine Mammals. *Am. Nat.*

Zhu, P., Liu, W., Zhang, X., Li, M., Liu, G., Yu, Y., *et al.* (2023). Correlated evolution of

social organization and lifespan in mammals. *Nat. Commun.*, 14, 372.

801
Fig. 1 | Bimodal global mammal space occupation stems from differences among environmental realms. a,

Probabilistic species distributions in the space defined by the two first principal components (PC1 and PC2) of

PCA considering life history traits for all mammals. Arrows indicate the loadings of each trait in the PCA. The

legend shows the amount of functional space (i.e., functional richness) occupied at 50% and 99% probabilities

(indicated by thick contour lines). The red regions falling within the limits of 50% probability correspond to the

hotspots mentioned in the text. wea, weaning length; fmat, time to reach female maturity; ls, size of litter; gest,

gestation length; long, longevity; ly, number of litters yearly. **b – d,** Patterns in space occupation of aerial (**b**),

terrestrial (**c**), and aquatic (**d**) mammals. Legends show functional redundancy of the realm calculated as

absolute values of the Standardized Effect Size ($Red_{99\%}$). SES was calculated considering the amount of

functional space occupied by observed and randomized realm at 99% probability. * is SES p-value < 0.05; **

SES p-value < 0.01; *** SES p-value < 0.001. In all panels, the colour gradient (red, orange, and white) depicts

different density of species in the space (red areas have higher density of species). Silhouettes of characteristic

species from each realm were downloaded from PhyloPic (http://www.phylopic.org/).

Fig. 2 | Environmental realms select for distinct yet nested life history strategies. Overlap-based

dissimilarities between terrestrial (purple), aerial (light orange), and aquatic (light blue) species distributions in

the life history space. Specifically: **a,** dissimilarity between terrestrial and aerial realms; **b,** dissimilarity between

terrestrial and aquatic realms; **c,** dissimilarity between aerial and aquatic realms. Each realm is highlighted at

50% (fine lines) and 99% (thick lines) of total probability. SES of dissimilarity and nestedness values for each

pairwise comparisons are reported in the legend together with SES p-values (* SES p-value < 0.05; ** SES p-

value < 0.01; *** SES p-value < 0.001).

Fig. 3 | Semi-aquatic species transition between aquatic and terrestrial life histories depending on the evolutionary time they spent in water. **A,** Occupation of transitional mammals in the life history space. The colour gradient (red, orange, and white) depicts different density of species in the space (red areas have higher density of species). The legend shows the amount of functional space (*i.e.*, functional richness) occupied at 50% and 99% probabilities (indicated by thick contour lines). Dashed grey line show the portion of space occupied by global mammals. Legend show functional redundancy of transitional species calculated as absolute values of the Standardized Effect Size ($Red_{99\%}$). SES was calculated considering the amount of functional space occupied by observed and randomized realm at 99% probability. Silhouettes of characteristic semi-aquatic species were downloaded from PhyloPic (http://www.phylopic.org/). **B-c,** pairwise overlap-based dissimilarities between semi-aquatic (SA) and both terrestrial (**b**) and fully aquatic (**c**) mammals. In both panels, contours at 50% (fine lines) and 99% (thick lines) are highlighted for terrestrial (purple), aquatic (light blue), and semi-aquatic (orange) species. Legends show observed dissimilarities between realms and semi-aquatic species (Diss) and value of nestedness (Nest). P-values for the standardized effect size (SES) of dissimilarity and nestedness are present only for significant SES. ***: SES p-value < 0.001. All panels show the probabilistic species

 distributions in the spaces defined by the two first principal components (PC1 and PC2) of PCA considering life

 history traits for all mammals.

Fig. 4 | Arboreal and fossorial mode-of-life are associated with slower life histories in taxonomically related terrestrial species. Differences in centroid among the life history structures of species confined on land (terrestrial) and **(a)** arboreal, **(b)** semiarboreal, and **(c)** fossorial species of the same order. Each arrow connects the centroid of the terrestrial structure and mode-of-life structure present within the same order. Green dots are arboreal centroids, aquamarine dots are semiarboreal centroids, purple dots are fossorial centroids, and orange dots are centroids of species confined on land.