Mammal life history strategies are strongly associated with environmental realms

E. Beccari[1*], P. Capdevila Lanzaco[2,3], R. Salguero-Gómez[4,5], & C. Pérez Carmona[1]

[1] Department of Botany, Institute of Ecology and Earth Sciences, University of Tartu, J. Liivi 2, 50409 Tartu, Estonia

[2] Departament de Biologia Evolutiva, Ecologia i Ciències Ambientals, Facultat de Biologia, Universitat de Barcelona (UB), Barcelona 08028, Spain

[3] Institut de Recerca de la Biodiversitat (IRBio), Universitat de Barcelona (UB), Barcelona 08028, Spain

[4] Department of Biology, University of Oxford, 11a Mansfield Rd, Oxford, OX1, 3SZ, UK

[5] Evolutionary Demography laboratory, Max Plank Institute for Demographic Research, Konrad Zuße Straße1, Rostock 18057, Germany

*Author for correspondences: Eleonora Beccari (eleonora.beccari@ut.ee)

Summary

Mammalian investments in survival, development, and reproduction are constrained by physiological and evolutionary factors[1], so their life history strategies can be characterized by a few axes of variation[2–4]. These axes conform a space in which species are positioned according to which life history strategies are favoured in the environment they exploit[1,5,6]. Yet, a global synthesis quantifying the realized mammalian life history diversity is lacking and exploring how the environment shapes mammalian life history strategies is missing. Here, we used six life history traits to build a global life history space and explored how major environmental realms (land, air, water) influence mammalian life history strategies. We demonstrate that realms are tightly linked to distinct life history strategies. Predominantly, aquatic and aerial species adhere to slower life history strategies, while terrestrial species tend to exhibit faster life histories. Highly encephalized terrestrial species,

like monkeys, are a notable exception to these patterns. In addition, species transitioning

between the terrestrial and aquatic realms, such as seals and otters, show intermediate life

history strategies. Furthermore, time since evolutionary divergence from land is a strong

predictor of aquatic life history strategies. Our results provide compelling evidence linking

environmental realms to the diversity of life history strategies among mammals, potentially

informing assessments of species viability.

Main:

The diversity of mammalian adaptation, ranging from the fast and semelparous life histories

of antechinus[7] to the slow and long life of bowhead whales[8], have allowed mammals to

occupy all land[9,10], exploit air[11–13], and even return to the sea[14]. Despite this wide ecological

radiation[10], evolutionary history and energetic constrains limit the set of life history strategies

(*i.e.,* combinations of life history traits) that are viable in nature[1,5,6]. Mammalian investments

in survival, growth, and reproduction are summarised along two major axes of life history

trait variation[3,4,15,16]. A first dimension generally relates to the time scale of live (*i.e.,* fast-

slow continuum[1–4,15,17]). In contrast, the second dimension generally reflects the timing and

intensity of species' reproductive investment[2,3,18] (but see Healy et al., 2019). While these

two dimensions are consistent across empirical studies considering different subsets of

species and traits[2,3,17], species organization within the life history space remain largely

unknown, preventing the disentanglement of the drivers of mammalian life history diversity.

Environmental conditions profoundly influence species' life history strategies[5,6]. Variable

environments, for instance, affect the pace of life of species[19] selecting for delayed life cycles

and longer lives compared to constant and stable environments[5,19]. Thus, similar life history

strategies are selected by similar environmental pressures[5,6], resulting in a redundant

occupation of the life history space regardless of taxonomic affiliation[15,20,21]. However,

mammal's widespread distribution across major environmental realms (*i.e.*, land, water, air),

begs the question of whether their life history strategies are equally represented across these

three major realms. Indeed, distinct physical dimensions and environmental requirements

expose species to realm-specific pressures[22]. For instance, real-specific selection pressures

have led to slower-paced lives in aerial species[3,11] or more fecund life histories in the aquatic

realm[3,17]. Consequently, species inhabiting different environmental realms may possess

distinct life history strategies.

Time is a currency necessary to accumulate evolutionary adaptations needed to transgress

realms[23–26]. Consequently, time since evolutionary divergence to a new realm should reflect

the degree to which a species' life history strategies are adapted to the environment. For

instance, mammal lineages that later diverged from land back to the aquatic realm, like seals

or otters, generally have a lower degree of aquatic specialization[8,14,27] and higher genetic

similarities with terrestrial sister taxa[28]. Accordingly, mammalian species with early return

back to water may display a higher degree of life history specialization to the aquatic realm[14].

Yet, the association between time since evolutionary divergence to a different realm and

species' life history strategies remains unexplored.

Here, we use six life history traits representing investments in survival, growth, and

reproduction for 3,438 mammalian species[29] to examine the influence of major environmental

realms on mammalian life history strategies. In agreement with previous research[2,3], we first

show that two axes of variation relating to the fast-slow continuum and the reproductive

strategies continuum are sufficient to explain a large proportion mammalian life history

strategies (Fig. 1a). Based on this finding as a springboard, we then test three hypotheses:

(H1) Species' position in the life history space will associate with the environmental realm

they occupy, as a consequence of different environmental pressures acting among major

realms[5,19,22]. Specifically, we expect aerial species to exhibit slower life history strategies

compared to terrestrial species, and aquatic species to occupy areas corresponding to faster[3]

and more fecund strategies[17]. (H2) Species transitioning between aquatic and terrestrial

realms (*i.e.,* semi-aquatic species) will display intermediate life history strategies, given that

they experience selective pressures from both realms[8,27]. (H3) Aquatic history strategies will

be associated with the time since evolutionary divergence from land. Specifically, we expect

that mammalian species that have evolved for longer time in the aquatic realm will display

more distinct life history strategies (*i.e.*, faster pace in this case[3]) compared to species that

transgressed from land to water more recently[8,14,23,28].

Global life history strategies are constrained and profoundly divergent

Mammalian life history traits vary mostly along a two-dimensional space, which suggests a

strong trait covariation limiting viable life history strategies. We gathered trait data related to

survival, development, and reproduction from the AMNIOTE database[29] for 4,953 mammal

species. We excluded species without information for body mass and used a

phylogenetically-informed imputation to fill missing data for the other traits data[15,30],

resulting in a total of 3,438 species (see Supplementary Methods 1 for the accuracy of the

imputation procedure). Residuals from a linear regression of each trait against body mass

were then used as body-size corrected trait values for further analyses [2,3,31] (Extended Data

Fig. 1b). Finally, we conducted a principal component analysis (PCA) to discern the primary

dimensions of mammalian trait variation (Supplementary Methods 2 and Extended Data Fig.

2 for results about the robustness of the life history space).

The first two principal components (PC) of the PCA account for 75.1% of life history

variation, conforming a life history space characterised by two primary trade-offs: litter size

vs. gestation length, and number of litters per year *vs.* age at female maturity and weaning

length (Figure 1a). The configuration of this space is line with previous studies on smaller

sets of species (267 species in Bielby et al., 2007; 80 species in Gaillard et al., 1989). The

first dimension of this space (PC1) captures the fast-slow continuum[1], distinguishing fast

species with short lives, gestation and weaning, early maturity, and frequent litters, from slow

species with fewer and infrequent litters, longer lives, and later maturity (Fig. 1a). The second

dimension (PC2) reflects lifetime reproductive effort, illustrating the trade-off between litter

size and frequency, as well as offspring quality (*i.e.,* altricial *vs.* precocial), and reproductive

frequency (*i.e.,* semelparity *vs.* iteroparity). Species' positioning and life history trait

correlations in the life history space remain consistent after accounting for evolutionary

history ($r = 0.95$; $p < 0.001$ for species; $r = 0.85$, $p < 0.001$ for traits; see Supplementary

methods 2), suggesting that this two-dimensional life history strategy space applies to all

mammals, independent of shared ancestry (Extended Data Fig. 3).

Mammalian species exhibit a remarkably uneven, concentrated distribution within the life

history space. Comparing global mammal occupation patterns along this space to null models

representing bivariate normal distributions with identical mean and covariance[30], the

occupied space is smaller than expected by chance (Extended data Fig. 4a). For example, at

the 99% quantile of species probability distribution, observed functional richness (*i.e.*,

amount of space occupied by species) is 32% smaller than the simulated counterpart

($SES_{Fric99\%} = -11.79$, $p < 0.001$, $n = 999$ iterations). Additionally, species' distribution is

much more divergent than the simulated ($SES_{Fdiv} = 18.79$, $p < 0.001$; $n = 999$ iterations;

Extended data Fig.4b), primarily due to species clustering in two hotspots at opposite ends of

the fast-slow continuum. Our findings reveal a dichotomy in successful investments to

survival, development, and reproduction of global mammals: species investing on quantity of

reproductive investment contrast with those prioritizing quality[1,5,6]. Species in the "fast

hotspot" (Fig. 1), such as mice and rabbits, optimize fast turnover and high reproductive

output, enabling them to propagate their genes in unpredictable and unfavourable

environments[5,6]. Conversely, "slow hotspot" species (Fig. 1a), including primates, cetaceans

and bats, typically make costly parental investments to equip their few offspring with

advantageous adaptations for long-term persistence and environmental exploitation[5,6,32]. This

bimodal pattern mirrors that of plant traits[30,33], suggesting that, across the Tree of Life,

species tend to cluster around a few contrasting syndromes that lead to successful strategies[34].

Fig. 1 | Bimodal global mammal space occupation stems from differences among environmental realms. a,

Probabilistic species distributions in the space defined by the two first principal components (PC1 and PC2) of

PCA considering life history traits for all mammals. Arrows indicate the loadings of each trait in the PCA. The

legend shows the amount of functional space (i.e., functional richness) occupied at 50% and 99% probabilities

(indicated by thick contour lines). The red regions falling within the limits of 50% probability correspond to the

hotspots mentioned in the text. wea, weaning length; fmat, time to reach female maturity; ls, size of litter; gest,

gestation length; long, longevity; ly, number of litters yearly. **b – d,** Patterns in space occupation of aerial (**b**),

terrestrial (**c**), and aquatic (**d**) mammals. Legends show functional redundancy of the realm calculated as

absolute values of the Standardized Effect Size (SES). SES was calculated considering the amount of functional

space occupied by observed and randomized realm at 99% probability. * is SES p-value < 0.05; ** SES p-value

< 0.01; *** SES p-value < 0.001. In all panels, the colour gradient (red, orange, and white) depicts different

density of species in the space (red areas have higher density of species). Silhouettes of characteristic species

from each realm were downloaded from PhyloPic (http://www.phylopic.org/).

Divergent life history strategies correspond to different environmental realms

Major environmental realms select for distinct and highly redundant life history strategies

(Fig. 1b-d; Fig. 2). This finding is confirmed by a PERMANOVA analysis, which attributes

35% of variation in life history traits to differences among realms. The marked differences in

life history strategies among realms are further demonstrated by the dissimilarities between

pairs of realms ($D_{land-water} = 69\%$, $D_{land-air} = 86\%$, $D_{air-water} = 86\%$; $p < 0.001$ in all cases, n =

999 iterations, Fig. 2; Extended data Table 2), as well as by the high levels of species

redundancy within individual realms (*i.e.,* species within an environmental realm tend to have

similar trait values; $R_{air} = 17.42$; $R_{land} = 6.22$, $R_{air} = 17.42$, $R_{water} = 8.62$; $p < 0.001$ in all cases,

n = 999 iterations; Extended data Table 2). Terrestrial species disproportionally occupy the

"fast hotspot" (Fig. 1c), including representatives of marsupials as well as the Carnivora,

Lagomorpha, Rodentia, and Eulipotyphla orders (Extended data Fig. 5). Primates and

Perissodactyla are the only exceptions to this pattern since they cluster in the slowest portion

of the life history space. On the contrary, aerial (Fig. 1a) and aquatic species (Fig. 1d) are

confined to the slow end of the spectrum. Despite the similar pace of life of aquatic and aerial

species, their reproductive strategies differ substantially. Aerial species exhibit long gestation

times, while aquatic species display late female maturity and extended weaning (Fig. 1a). The

predominance of slow strategies in aquatic species contrasts with previous observations

mainly based on baleen whales[3], which are among the longest-lived aquatic species (Fig. 1d).

Our study, however, includes dolphins, displaying a more complete picture. Dolphins and

baleen whales exhibit distinct migratory, dietary, and social habits[8], influencing resource

perception and gathering within realms and affecting life history variation[35–37].

Despite marked differences and redundancy among realms, characteristic strategies of a given

environmental realm can be observed in others. This shared set of life history strategies is

particularly evident in the case of the aquatic realm, fully contained within the life history

space occupied by terrestrial species ($Nestedness_{land-water} = 1$, Fig. 2b). The aerial realm is also

highly overlapping with the terrestrial one ($Nestedness_{land-air} = 0.89$; Fig. 2a); primates, a

small subset of terrestrial species, are the main reason for such an overlap. For example, apes

(including humans), and dolphins have essentially the same life history traits, whereas

terrestrial Lemuriformes and Tarsius species are located in the part of the life history space

occupied by bats (Figure 1; Extended Data Fig. 5). The arboreal behaviour of lemurs[38] allows

them to escape from unfavourable situations, decreasing extrinsic mortality, extending

longevity, and selecting for slow lives[11,39]. Similar occupation of life history space between

higher primates and dolphins (Fig. 2a, Fig. 1) might be linked with complex social structures

and high encephalization in both groups[8,40,41]. The development of high encephalisation and

social behaviours requires vast parental investments[32,42] but, in turn, enables species to

exploit the environment in a plastic way, avoiding hazards and stressful situations[37,42].

Consequently, high encephalization, which generally relates with both social and behavioural

facets[32,43], may enable terrestrial species to adopt strategies that would be otherwise

unfavourable. We tested this hypothesis by modelling the brain-mass ratio of terrestrial

mammalian species (*i.e.* the ratio between brain dimension and body mass[41,44]) as a function

of species' position in the life history space (Supplementary methods 4). We found a robust

association between larger brain-mass ratios and slow life histories ($R^2 = 0.53$, Extended data

Fig. 6), suggesting that larger brain size may significantly influence the diversity of terrestrial

life history strategies. Consequently, while the environmental realm in which species live

seems to drive life history strategies, behavioural, and ecological characteristics of species

(*e.g.,* sociality, plastic thinking, arboreality) may allow them to successfully exploit parts of

the life history space that would otherwise not be viable in their specific realms[21].

Fig. 2 | Environmental realms select for distinct yet nested life history strategies. Overlap-based

dissimilarities between terrestrial (purple), aerial (light orange), and aquatic (light blue) species distributions in

the life history space. Specifically: **a,** dissimilarity between terrestrial and aerial structures; **b,** dissimilarity

between terrestrial and aquatic structures; **c,** dissimilarity between aerial and aquatic structures. Each system is

highlighted at 50% (dashed line) and 99% (full line) of total probability. SES of dissimilarity and nestedness

values for each pairwise comparisons are reported in the legend together with SES p-values (* SES p-value <

0.05; ** SES p-value < 0.01; *** SES p-value < 0.001).

Species transitioning between realms have intermediate life history

strategies

The life history strategies of semi-aquatic species provide further evidence for the profound

influence on life history variation by environment realms. We examined 58 semi-aquatic

mammal species that engage in activities such as feeding, mating, and offspring weaning in

aquatic environments, but are not exclusively aquatic like dolphins or whales (Supplementary

Methods 3). To understand how the life history strategies of these species differ from strictly

terrestrial and strictly aquatic species, we analysed their dissimilarity patterns. Semi-aquatic

species exhibit intermediate life history strategies, slower than terrestrial species but faster

than fully aquatic ones (Fig. 3). This pattern holds within semi-aquatic mammals, whereby

species occupy different regions of the life history space depending on their degree of water-

dependence. For example, species with greater aquatic involvement, such as seals and

walruses[8,14], occupy portions of the life history space similar to fully aquatic species like

manatees, beluga, or sperm whales. Conversely, species with lesser water-dependence, like

rodents and otters[8,14,27], are located in the hotspot typical of terrestrial mammals (Fig. 3).

The reported intermediate life history strategy location of semi-aquatic mammals is

independent of taxonomic order. This robustness confirms the association between

environmental realms and life history strategies even among closely related species. For

example, terrestrial and semi-aquatic carnivores separate in the life history space following

the same patterns seen in the global dataset with terrestrial carnivores being faster than their

semi-aquatic counterparts (Extended data Fig. 7a-c). Moreover, the sea otter (*Enhydra lutris*),

the only fully aquatic carnivore, exhibits a markedly different life history strategy compared

to semi-aquatic otters (Extended data Fig. 7). Among Cetartiodactyla, aquatic (dolphins,

whales) and semi-aquatic (hippos) species have slower life history strategies than terrestrial

species (giraffe, bison, gazelles; Extended data Fig. 7d-f). Altogether, these findings suggest

that species transitioning between aquatic and terrestrial realms possess intermediate life

history strategies.

Time since divergence to aquatic realm associates with species life history strategies

The amount of time a mammal has evolved in the aquatic realm is associated with its life history strategy. For the 143 aquatic-related species (*i.e.,* aquatic, and semi-aquatic) in our dataset that transgressed realms, back from the terrestrial to the aquatic realm, we extracted from the literature their time since divergence (in m.y.) from their last terrestrial ancestor. There is a significant association between the time since divergence to aquatic realms and the position of mammals in the life history space ($R^2 = 0.34$ in semi-aquatic species, Figure 3d; $R^2 = 0.40$ in water-related species, Figure 3e; Extended data Table 2). With the notable exception of the platypus, semi-aquatic species with longer evolution in the aquatic realm, like seals and hippos, overlap in the life history space with cetacean and sirenians, that transgress realms at earlier times (Fig. 3d-e). Conversely, species with shorter evolutionary history in water, like mustelids and otters[8,14,27], are clustered towards the regions dominated by terrestrial species (Figure 3 d-e). Accordingly, a longer evolution under aquatic selection pressures may foster the accumulation of necessary for aquatic specialization[14,23,28]. This process could cause early diverging semi-aquatic species to gradually align with the trait space occupied by fully aquatic ones. Our findings indicate a link between the evolutionary separation from a shared aquatic-terrestrial ancestor and the life history strategies of species, underscoring the role of time in shaping realm-specific adaptations.

Fig. 3 | Semi-aquatic species transition between aquatic and terrestrial life histories depending on the evolutionary time they spent in water. a, Occupation of transitional mammals in the life history space. The colour gradient (red, orange, and white) depicts different density of species in the space (red areas have higher density of species). The legend shows the amount of functional space (*i.e.*, functional richness) occupied at 50% and 99% probabilities (indicated by thick contour lines). Dashed grey line show the portion of space occupied by global mammals. Silhouettes of characteristic semi-aquatic species were downloaded from PhyloPic (http://www.phylopic.org/). b-c, pairwise overlap-based dissimilarities between semi-aquatic (SA) and both terrestrial (b) and fully aquatic (c) mammals. In both panels, contours at 50% (fine lines) and 99% (thick lines) are highlighted for terrestrial (purple), aquatic (light blue), and semi-aquatic (orange) species. Legends show

observed dissimilarities between realms and semi-aquatic species (Diss) and value of nestedness (Nest). P-

values for the standardized effect size (SES) of dissimilarity and nestedness are present only for significant SES.

***: SES p-value < 0.001. All panels show the probabilistic species distributions in the spaces defined by the

two first principal components (PC1 and PC2) of PCA considering life history traits for all mammals. **d-e,** time

since divergence from terrestrial ancestor mapped in the life history space. **d**; time since divergence modelled as

a function of the position of semi-aquatic species in the life history space. **e**; time since divergence modelled as

a function of the position of aquatic-related species in the life history space. Adjusted r-squared of each

generalized additive models (GAMs) are reported in the upper part of the panels. Darker tones represent long

time since divergence and lighter tones represent short time since divergence. White contour lines represent the

thresholds of different time since divergence.

Conclusion

Our analyses of over 3,000 mammal species reveal that major environmental realm is a

strong predictor of mammalian life history strategies. Aerial[3,11] and aquatic[14,45,46] realms

require specific adaptations, such as flight-capable skeletal structures[12,13] or enhanced

thermoregulatory abilities for aquatic life[8,14]. These adaptations slow down and prolong

lifespans in ways seemingly unexploited in the terrestrial realm by mammals[11]. However,

despite each realm's unique set of life history strategies, all of them overlapped within the life

history space due to species' adaptations that enable the exploitation of otherwise

unsuccessful strategies in each realm. Empirical research has shown that adaptations in

social[37], migratory[36], dietary characteristics[35], and cerebral[32,47] can influence species' life

histories. Consequently, variations in these behavioural and ecological characteristics

modulate species' responses to environmental pressures[5] and expand the spectrum of life

history strategies available for exploitation within a realm, as observed for activity patterns[21].

For instance, differences in migratory, dietary, and social habits among baleen whales and

dolphins[8] may facilitate distinct life history strategies in aquatic environments, accounting for

the rapid pace of life we observed in baleen whales. Similarly, the high sociality and

encephalization of primates[48] may allow these terrestrial mammals to exploit the slower end

of the life history spectrum. The strong relationship we uncovered between brain dimension

and the position of terrestrial species within the life history space (Extended data Fig. 4)

supports this hypothesis. Accordingly, we advocate for future research to investigate the

influence of ecological and behavioural adaptations on life history variation among terrestrial

and aquatic mammals.

The time elapsed since an extant aquatic mammal diverged from its last terrestrial ancestor

influences its life history strategy. The environmental pressures of a particular realm enable

species to develop the necessary adaptations for success in that environment[5,6]. Accordingly,

if the realm determines the type and magnitude of the environmental pressure acting on

species, the duration these pressures are experienced can be pivotal in accumulating the

evolutionary adaptations required for realm-specific specialization[23–26,28]. Our finding of a

correlation between the life history strategies of aquatic-related species and the duration of

aquatic-related evolution (Figure 3d-e) supports this hypothesis, suggesting a combined

impact of environmental pressures and exposure duration on realized mammalian life history

diversity. However, more research is needed to fully understand the process driving the

evolution of mammalian life history strategies under different environmental constraints.

Our study provides a comprehensive assessment of global mammalian life history strategies,

revealing the fundamental role of environmental realm in shaping these patterns.

Remarkably, these relations persist even among closely related species, highlighting the

profound relation between environmental realms and life history evolution. Our findings not

only provide empirical insights into mammalian adaptation, but also establish a foundation

for further exploration of ecological and evolutionary processes[9,10]. Simultaneously, this

foundation improves our understanding of the drivers and consequences of ongoing global

changes[15,49] and future biodiversity patterns[15,20,50]. As we confront these challenges, our study

emphasises the importance of understanding the intricate interplay between environment,

ecology, and evolution to disentangle, anticipate, and mitigate the impacts on mammalian life

history diversity.

Methods

Data collection and trait imputation

To characterise the mammals life history strategies, we downloaded the data from Amniote

database which comprises information for ca. 18.300 species of mammals, reptiles and

birds[29]. We extracted mammalian species (N = 4,953), and selected the life history traits with

the most complete information (*i.e.,* traits with records for at least 1000 species): litter size

(ls, number of offspring per reproductive event), litter per year (ly, number of litters per

year), age at female maturity (fmat, time needed for female individuals to reach maturity in

days), weaning length (wea, time needed to wean the offspring in days), gestation length

(gest, time passing between conception and birth in days), longevity (long, individual lifespan

in years) and adult body mass (bm, in grams).

Because no trait was fully informed for all species, we imputed missing life history trait

values. Recent studies have shown that imputation of individual traits allows to accurately

characterize the position of species in synthetic trait spaces that are the result of combining

different correlated traits space (see Carmona, Tamme et al., 2021 and Carmona, Bueno et al.,

2021 for examples). Further, recent simulations have shown that studying the patterns of

occupation of the synthetic space considering the imputed species yields results that are much

closer to the real patterns of occupation of the space than those reached when only those

species for which there is full trait information are considered[51]. Because accounting for

evolutionary relationships improves the accuracy of the imputation[52], we downloaded

mammalian phylogenies from Vertlife database[53]. We selected the complete fossil-based set

of 10.000 phylogenies and computed a single maximum clade credibility tree (MCC tree)

using the *maxCladeCred* function of *'phangorn'* R package[54]. Species absent in the

phylogeny but with life history information were added at the genus' root using

add.species.to.genus function of *'phytools'* R package[55]. Species that were missing in the life

history traits database were pruned using *drop.tip* function of *'phyltools'* R package.

For the imputation procedure, we selected mammalian species with information for body size

and at least one of the other considered life history traits (*i.e.,* ls, ly, fmat, wea, gest, long),

resulting in a subset of 3,438 species. Then, following Carmona, Tamme et al. 2021, we

included phylogenetic information in the imputation procedure using the first 10 phylogenetic

eigenvectors derived from the MCC tree. For the imputation, we used the *missForest* function

from the *'missForest'* R package[56]. After the imputation, we estimated the reliability of the

imputation procedure (Supplementary methods 1). The imputation procedure efficiently

retrieved the real position of a species, with an error of around 2.8% of the range for the first

component and 1.7% for the second component. The final procedure resulted in a dataset of

3,438 mammalian species (53.7% of currently recognized living mammals[9]) from 29 orders

in which 37.6% of traits information (1,293 species) derived from empirical measurement

and 62.4% (2,145 species) were imputed.

Ascribing species to environmental realms

In order to have information on species adscription to the terrestrial, aerial, and aquatic

realms, we downloaded information on the environment exploited by each mammal from the

Worms[57] and IUCN[58] databases using the R packages *'worms'*[59] and *'redlist'*[60], respectively.

We classified mammalian species as "terrestrial" (*i.e.,* land realm) in case of unambiguous

classification from both databases, whereas we limited "aerial" category (air) to powered

flight species (*i.e.,* Chiroptera order). Although there is no doubt on the classification of

"fully" aquatic species (*i.e.*, species that spent the entire life in water, *e.g.,* dolphins, whales,

dugongs[8]), there is not a strict classification of semi-aquatic species[27]. Due to this, the Worms

and IUCN databases classification did not match for many species. For instance, *Macaca*

mulatta is classified as semi-aquatic according to the Worms database, while IUCN classifies

it as terrestrial. To solve this problem, we performed a finer classification of aquatic species

by summarizing species' dependency on water. First, we selected all species classified as

freshwater and marine in at least one of the IUCN and Worms database. Then, we created an

Aquatic Dependency index (AD Index) to classify species as "fully" aquatic, "fully"

terrestrial, or semi-aquatic. We built the AD Index answering a series of binary questions

capable to summarize the amount of time spent in water during fundamental phases of

species life histories (*e.g.,* birth, mating; Supplementary methods 3). The AD Index was used

to refine aquatic classification by relabelling species scoring zero in AD Index (*i.e.,* no

important parts of life are strictly constrained by water presence) as terrestrial species. Then,

we classified as aquatic (hereafter aquatic) all species scoring 5 out of a maximum of 5 on the

AD Index (*i.e.,* species that spent their whole life in water, like dolphins), while we classified

as semi-aquatic all species scoring between 1 and 4 (Supplementary methods 3). Out of the

3,438 mammalian species considered in our analyses, 2,634 species were classified as

terrestrial (76.6% of total set of species), 661 species were classified as aerial (19.2% of the

total), 85 species were classified as aquatic (2.5% of the total), and 58 species were classified

as semi-aquatic (*i.e.,* transitional species, 1.7% of the total).

All previously mentioned taxonomies (*i.e.,* life history traits, phylogeny, Worms and IUCN

databases) were standardized using *gnr_resolve* function of the *'taxize'* R package[61] resolving

for the Global Biodiversity Information Facility (GBIF) Backbone Taxonomy.

Construction of the global life history traits space

Body mass is a strong allometric constraint that affects the values of many other life history

traits[2,31,62]. Mammals' body mass can span from the few grams of shrews, to the tons for

whales, and the aquatic realm strongly selects for bigger sizes compared to the terrestrial

realm[45]. Therefore, to address life history strategies without confounding effects from

different orders of magnitude, we accounted for the effect of body mass before computing the

mammalian life history space. For this, we performed a series of linear regressions between

single log-transformed life history traits and log-transformed species adult body mass. Then,

we extracted body mass–corrected residuals for each species from each regression[62,63]. Body

mass–corrected residuals were further centred, scaled, and used as size-corrected life history

traits (Extended data Fig. 1).

Mapping species position in multidimensional spaces based on trait information allows the

identification of major axis of trait variation[64,65], as well as summarizing species' ecological

strategies. We constructed mammalian life history space by performing principal component

analysis (PCA) on the complete set of mammalian species. We retained the first two axes

which together account for 75.1% of variance in mammalian life history traits. Then we

estimated the reliability of our methodological choices by comparing this life history space

(*i.e.,* the scores of species in the selected number of principal components) with three other

spaces: one built considering only complete set of records (*i.e.,* non-imputed data; N = 1,293;

Extended data Fig. 2); one built considering the imputed dataset without body mass

correction (N = 3,438); and one built considering the same body mass–corrected residuals

used to build the life history space (N = 3,438) but accounting for the phylogenetic history of

species (Extended data Fig. 3; Extended data Table 1). In all abovementioned cases, the

comparisons revealed strong correspondence between spaces, demonstrating that our

methodological choices do not influence the inferred relationships between life history traits

(Supplementary Methods 2).

Exploring patterns within the global life history space

We derived life history structure (*i.e.,* the patterns of organization of species in the life history

space) using the trait probability density framework (TPD)[66], which computes species'

probabilistic distribution in the life history space using multivariate kernel density estimation.

The kernel density estimation for each species was estimated as a multivariate normal

distribution centred at the species' coordinates in the PCA space[66,67], and with a standard

deviation computed using the unconstrained bandwidth selectors from the *'Hpi'* function in

the *'ks'* package[68,69]. To estimate the life history structure of the whole set of mammals, all

species' kernels were weighted and aggregated to form a single continuous trait probability

density (TPD) function across the whole mammalian life history space. The value of the TPD

function in any given point of the life history space reflects the density of species in that

particular area of the life history space[66,67]. We graphically represented the so built

mammalian life history structure by highlighting the contours containing 50% and 99% of the

total probabilistic distribution of species. We used the *'TPD'* R package[67,70].

We used mammalian global structure to test whether mammals distribute in the life history

space following a random assembly of life history traits (*i.e.,* life history strategies are

randomly selected) or whether certain life history strategies are more prevalent than

others[6,30,64]. First, we computed functional richness (the amount of functional space

occupied[66,71]) at increasing probability thresholds (*i.e.,* from 0.1% to 99.9% of the TPD

function, using 0.1% increments). In this way, we estimated a "profile" of the probabilistic

distribution of species, reflecting what proportion of the life history space is occupied at

different probability thresholds. We compared the observed functional richness profile with a

null model (999 repetitions) considering a bivariate normal distribution of species with the

same mean and variance as the observed data (see Carmona, Bueno et al., 2021).[30] This

analysis was performed by sampling 3,438 simulated species *(i.e.,* complete set of mammals)

from a bivariate normal distribution with the same mean and covariance of observed data

calculated using the *'rmvnorm'* function of the R package *'tmvtnorm'*[72]. In the same way as

done for the observed data, for each repetition of the null model we computed life history

structure and measured functional richness at increasing probability thresholds. We then

compared the observed and null functional richness profiles; functional richness values lower

than expected under the null model would suggest that the global set of mammals occupy a

smaller portion of the life history space than expected under a normal distribution and vice-

versa.

Then, using the same null model, we tested whether the portions of life history space that are

mainly exploited by mammalian species are closer or farther than expected from the centre of

the distribution. For this, we estimated functional divergence *(i.e.,* degree to which species'

density is skewed towards the extremes of the life history space) considering 99% of the total

species' probability[66,71] for both observed and simulated data. High functional divergence

would reveal that species are segregated towards the edges of the life history space, whereas

low divergence would indicate that species are clustered in the centre of the life history

space[30].

We estimated the standardized effect size (SES) of both the functional profiles (at each

probability threshold) and the functional divergence measures as: SES = (observed value –

mean (simulated values))/standard deviation (simulated values). SES values express the

number of SD units by which the observed value deviates from the mean of the simulated

ones. A SES value lower than 0 indicates that the observed metric is smaller than the average

of the simulated values and vice versa. Finally, we tested the statistical significance of

observed and simulated differences by estimating two-sided p-values computed by

confronting the SES values with a cumulative normal distribution with mean $= 0$ and

standard deviation $= 1^{30}$.

Lastly, to have a clearer picture of how different species are positioned in the life history

space, we computed TPD functions for each mammalian order with at least 6 species (total of

18 orders). We graphically represented the mammalian life history structure of each order by

highlighting the contours containing 50% and 99% of the total probabilistic distribution of

species order (Extended data Fig. 5).

Exploring environmental realms' patterns in the life history space

We produced environmental realm-specific TPD functions by aggregating the species that are

characteristic of each realm. We followed the same procedure explained above for the global

set of mammalian species. Similarly, to the complete mammalian life history structure, we

represented graphically the TPD for each realm and highlighted contours containing 50% and

99% of the total probability.

We compared several aspects of the structure of each of the environmental realms (*i.e.,* air,

land, water and transitional). First, we compared them in terms of the amount of life history

space that they occupy (*i.e.,* their functional richness). Because functional richness is known

to be positively related to the number of species present in a group, we built a null model

(999 repetitions) randomizing species assignment[30]. For each realm, we randomly extracted

the same number of observed species from the complete mammalian database and computed

TPD functions of the random community. We computed functional richness at the 99% of

total probability for both observed and randomized structures and calculated SES as

explained above. Under this null model, negative SES values would indicate that the species

present in the realm are more clustered in the life history space (*i.e.,* occupy a smaller portion

of the life history space, and therefore species are redundant in their life history strategy) than

expected given the same number of randomly selected species and vice versa. A highly

clustered realm would indicate high redundancy of life history strategies[30], suggesting that

the realms exert a strong selection on life history strategies. This procedure allows to

compare functional richness among realms with very different number of species (Extended

data Table 2a).

Then, we statistically tested whether dissimilarities in species positioning were a product of a

realm-driven selection of life history strategies. We estimated how much of the total variation

in the position of the species on life history space was explained by difference in realms. For

this, we first calculated the dissimilarities (using Euclidean distances) between all pairs of

species in the life history space. Then, we performed a PERMANOVA analysis considering

the three major environmental realms (*i.e.,* air, land, and water) as the explanatory variable (R

package *'vegan'*[73]). If the realm explains a large portion of life history variation, it means that

life history variability is mainly related to differences between realms rather than differences

among species within the realm[30].

We estimated pairwise dissimilarities in life history structures between pairs of realms (*i.e.,*

air, land, water, and transitional). Dissimilarities were calculated as 1 - overlap of the

corresponding TPD functions producing measures varying between 1, indicating a complete

overlap between the considered TPD functions, and 0, when the two functions are completely

disjoint. TPD-based dissimilarities consider both differences in the boundaries and

differences in the density of the occupation of the life history space[15,50,74]. These

dissimilarities can be further decomposed into two complementary components: one

quantifying up to what point the life history differences between realms are related to the

occupation of exclusive regions of the space (*i.e.,* turnover) and another reflecting how

differently the two realms occupy the shared parts of the life history space (*i.e.,* nestedness[67]).

We tested whether the observed dissimilarities between environmental realms were

significantly higher than expected by chance by means of null models[75,76]. In this null model,

for each pair of realms, we considered only the set of species belonging to both realms and

we randomized the realm classification. We maintained the same number of species observed

in each considered realm and avoided species assignment to multiple realms. For each

repetition of the null model (n = 999), we estimated the TPD functions of each realm and

estimated their dissimilarity, as well as its nestedness and turnover components. Finally, we

computed SES values for all mentioned metrics as described above (Extended Data Table

2b).

Then, we explored whether high encephalization[32,42,43,77] allows to exploit portion of the life

history space otherwise not successful for the terrestrial realm (Supplementary methods 4).

The final procedure resulted in generalized additive models (GAM) mapping species

encephalization (expressed as brain-mass ratio[41,44]; brain mass information were extracted

from Combine database[78]) in response to the species position in the life history space

(Extended data Fig. 6).

Lastly, we explored whether dissimilarities between environmental realms were maintained

also in closely related species. To do this, we computed TPD functions for orders having

species inhabiting in different realms. Because all aerial species belong to the Chiroptera

order, we decided to focus on orders having both terrestrial and aquatic species. We extracted

orders that had at least six species in both the aquatic (in this case identified by both aquatic

and semi-aquatic species) and terrestrial realms. For each of these orders, we calculated a life

history structure for terrestrial species and one for aquatic species. We represented

graphically structures highlighting 50 and 99% of the total probability and we computed

pairwise dissimilarities between the terrestrial and aquatic life history structure of each order

(Extended data Fig. 7).

Evolutionary time in the environmental realms and species life histories

While the aerial realm consists in the single Chiroptera order that diverged from a common terrestrial ancestor and diversified to all its modern form[79], aquatic-related (*i.e.,* aquatic and semi-aquatic) species have returned to water in multiple independent occasions across different lineages[8,80]. Consequently, we decided to focus on the subset of aquatic-related species and to test whether the length of evolution in relation to water is related to species life history strategies. For this, we performed a literature search to compile a database of time since divergence from the last terrestrial ancestor for the full set of aquatic and semi-aquatic species (N =143). It worth to mention that lineages diversified from a last common terrestrial ancestor, like seals or cetaceans[80,81], shared the same time since divergence.

We mapped squared root-transformed time since divergence from last terrestrial ancestor of the full set of aquatic-related species as a response to species positions in the life history space following Carmona, Tamme, et al., 2021. We used generalized additive models (GAM), where time was used as response variable, and the species coordinates in the life history space were used as explanatory variables. The latter was computed using 5 basis functions (k) of thin plate regression splines basis type, considering both main effect of each PCs composing the life history space as well as the interaction between them. As smoothing parameter estimation, we used the REML method, and we used a gamma family distribution (Figure 3e; Extended Data Table 3). Additionally, to test whether the findings were consistent also within semi-aquatic species, we performed another GAM model considering only the semi-aquatic species and using the same parameters reported above (Figure 3d; Extended Data Table 3). Both GAMs were computed using the r package *'mgcv'*[82].

All statistical analyses were performed using R version 4.0.3[83].

References

1. Stearns, S. C. *The evolution of life histories*. vol. 249 (Oxford university press Oxford, 1992).

2. Bielby, J. *et al.* The Fast☐Slow Continuum in Mammalian Life History: An Empirical Reevaluation. *Am. Nat.* **169**, 748–757 (2007).

3. Gaillard, J.-M. *et al.* An Analysis of Demographic Tactics in Birds and Mammals. *Oikos* **56**, 59–76 (1989).

4. Healy, K., Ezard, T. H. G., Jones, O. R., Salguero-Gómez, R. & Buckley, Y. M. Animal life history is shaped by the pace of life and the distribution of age-specific mortality and reproduction. *Nat. Ecol. Evol.* **3**, 1217–1224 (2019).

5. Grime, J. P. & Pierce, S. *The evolutionary strategies that shape ecosystems*. (John Wiley & Sons, 2012). doi:10.1002/9781118223246.

6. Southwood, T. R. E. Tactics, Strategies and Templets. *Oikos* **52**, 3–18 (1988).

7. Braithwaite, R. W. & Crowther, M. Brown antechinus Antechinus stuartii. in *The mammals of Australia* 94–96 (Reed New Holland, 2008).

8. Perrin, W. F., Würsig, B. & Thewissen, J. G. M. *Encyclopedia of marine mammals*. (Elsevier Inc., 2009). doi:10.1016/B978-0-12-373553-9.X0001-6.

9. Burgin, C. J., Colella, J. P., Kahn, P. L. & Upham, N. S. How many species of mammals are there? *J. Mammal.* **99**, 1–14 (2018).

10. Grossnickle, D. M., Smith, S. M. & Wilson, G. P. Untangling the Multiple Ecological Radiations of Early Mammals. *Trends Ecol. Evol.* **34**, 936–949 (2019).

11. Babich Morrow, C., Ernest, S. K. M. & Kerkhoff, A. J. Macroevolution of dimensionless life-history metrics in tetrapods. *Philos. Trans. R. Soc. B: Biol. Sci.* **288**, 20210200 (2021).

12. Barclay, R. M. R. Constraints on Reproduction by Flying Vertebrates: Energy and

Calcium. *Am. Nat.* **144**, 1021–1031 (1994).

13. Maina, J. N. What it takes to fly: the structural and functional respiratory refinements in

birds and bats. *J. Exp. Biol.* **203**, 3045–3064 (2000).

14. Davis, R. W. *Marine mammals: adaptations for an aquatic life*. (Springer Cham, 2019).

doi:10.1007/978-3-319-98280-9.

15. Carmona, C. P. *et al.* Erosion of global functional diversity across the tree of life. *Sci.*

Adv. (2021).

16. Oli, M. K. The fast–slow continuum and mammalian life-history patterns: an empirical

evaluation. *Basic Appl. Ecol.* **5**, 449–463 (2004).

17. Capdevila, P. *et al.* Longevity, body dimension and reproductive mode drive differences

in aquatic versus terrestrial life-history strategies. *Funct. Ecol.* **34**, 1613–1625 (2020).

18. Paniw, M., Ozgul, A. & Salguero-Gómez, R. Interactive life-history traits predict

sensitivity of plants and animals to temporal autocorrelation. *Ecol. Lett.* **21**, 275–286

(2018).

19. Tuljapurkar, S., Gaillard, J.-M. & Coulson, T. From stochastic environments to life

histories and back. *Philos. Trans. R. Soc. B: Biol. Sci.* **364**, 1499–1509 (2009).

20. Cooke, R. S. C., Eigenbrod, F. & Bates, A. E. Projected losses of global mammal and

bird ecological strategies. *Nat. Commun.* **10**, 2279 (2019).

21. Cox, D. T. C., Gardner, A. S. & Gaston, K. J. Diel niche variation in mammals associated

with expanded trait space. *Nat. Commun.* **12**, 1753 (2021).

22. Webb, T. J. Marine and terrestrial ecology: unifying concepts, revealing differences.

Trends Ecol. Evol. **27**, 535–541 (2012).

23. Chikina, M., Robinson, J. D. & Clark, N. L. Hundreds of Genes Experienced Convergent

Shifts in Selective Pressure in Marine Mammals. *Mol. Biol. Evol.* **33**, 2182–2192 (2016).

24. Foote, A. D. *et al.* Convergent evolution of the genomes of marine mammals. *Nat. Genet.*

47, 272–275 (2015).

25. Gingerich, P. D. Rates of evolution on the time scale of the evolutionary process. in

Microevolution Rate, Pattern, Process (eds. Hendry, A. P. & Kinnison, M. T.) 127–144

(Springer Netherlands, 2001). doi:10.1007/978-94-010-0585-2_9.

26. Stroud, J. T. & Losos, J. B. Ecological Opportunity and Adaptive Radiation. *Annu. Rev.*

Ecol. Evol. Syst. **47**, 507–532 (2016).

27. Hood, G. A. *Semi-aquatic Mammals: Ecology and Biology*. (JHU Press, 2020).

doi:10.1353/book.78278.

28. Yuan, Y. *et al.* Comparative genomics provides insights into the aquatic adaptations of

mammals. *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2106080118 (2021).

29. Myhrvold, N. P. *et al.* An amniote life-history database to perform comparative analyses

with birds, mammals, and reptiles. *Ecology* **96**, 3109–3109 (2015).

30. Carmona, C. P. *et al.* Fine-root traits in the global spectrum of plant form and function.

Nature **597**, 683–687 (2021).

31. Sibly, R. M. & Brown, J. H. Effects of body size and lifestyle on evolution of mammal

life histories. *Proc. Natl. Acad. Sci. U.S.A.* **104**, 17707–17712 (2007).

32. Barton, R. A. & Capellini, I. Maternal investment, life histories, and the costs of brain

growth in mammals. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 6169–6174 (2011).

33. Salguero-Gómez, R. *et al.* Fast–slow continuum and reproductive strategies structure

plant life-history variation worldwide. *Proceedings of the National Academy of Sciences*

113, 230–235 (2016).

34. Junker, R. R. *et al.* Towards an animal economics spectrum for ecosystem research.

Funct. Ecol. **37**, (2022).

35. Famoso, N. A., Hopkins, S. S. B. & Davis, E. B. How do diet and body mass drive

reproductive strategies in mammals? *Biol. J. Linn.* **124**, 151–156 (2018).

36. Soriano-Redondo, A., Gutiérrez, J. S., Hodgson, D. & Bearhop, S. Migrant birds and

mammals live faster than residents. *Nat. Commun.* **11**, 5719 (2020).

37. Zhu, P. *et al.* Correlated evolution of social organization and lifespan in mammals. *Nat.*

Commun. **14**, 372 (2023).

38. Gebo, D. L. Vertical clinging and leaping revisited: Vertical support use as the ancestral

condition of strepsirrhine primates. *Am. J. Phys. Anthropol.* **146**, 323–335 (2011).

39. Scheffers, Brett. R. *et al.* Vertical (arboreality) and horizontal (dispersal) movement

increase the resilience of vertebrates to climatic instability. *Glob. Ecol. Biogeogr.* **26**,

787–798 (2017).

40. Ridgway, S. H., Carlin, K. P., Van Alstyne, K. R., Hanson, A. C. & Tarpley, R. J.

Comparison of Dolphins' Body and Brain Measurements with Four Other Groups of

Cetaceans Reveals Great Diversity. *Brain Behav. Evol.* **88**, 235–257 (2017).

41. Worthy, G. A. J. & Hickie, J. P. Relative Brain Size in Marine Mammals. *Am. Nat.*

(1986) doi:10.1086/284579.

42. González-Lagos, C., Sol, D. & Reader, S. M. Large-brained mammals live longer. *j.*

Evol. Biol. **23**, 1064–1074 (2010).

43. Pérez☐Barbería, F. J., Shultz, S. & Dunbar, R. I. M. Evidence for coevolution of sociality

and relative brain size in three orders of mammals. *Evolution* **61**, 2811–2821 (2007).

44. Benson-Amram, S., Dantzer, B., Stricker, G., Swanson, E. M. & Holekamp, K. E. Brain

size predicts problem-solving ability in mammalian carnivores. *Proc. Natl. Acad. Sci.*

U.S.A. **113**, 2532–2537 (2016).

45. Gearty, W., McClain, C. R. & Payne, J. L. Energetic tradeoffs control the size

distribution of aquatic mammals. *Proc. Natl. Acad. Sci. U.S.A.* **115**, 4194–4199 (2018).

46. Uhen, M. D. Evolution of marine mammals: Back to the sea after 300 million years. *Anat.* **290**, 514–522 (2007).

47. Seyfarth, R. M. & Cheney, D. L. What are big brains for? *Proc. Natl. Acad. Sci. U.S.A.* **99**, 4141–4142 (2002).

48. Jones, J. H. Primates and the Evolution of Long-Slow Life Histories. *Curr. Biol.* **21**, R708–R717 (2011).

49. Enquist, B. J., Abraham, A. J., Harfoot, M. B. J., Malhi, Y. & Doughty, C. E. The megabiota are disproportionately important for biosphere functioning. *Nat. Commun.* **11**, 699 (2020).

50. Toussaint, A. *et al.* Extinction of threatened vertebrates will lead to idiosyncratic changes in functional diversity across the world. *Nat Commun* **12**, 5162 (2021).

51. Stewart, K. *et al.* Trait diversity metrics can perform well with highly incomplete datasets. *bioRxiv* (2022) doi:10.1101/2022.11.08.515633.

52. Penone, C. *et al.* Imputation of missing data in life-history trait datasets: Which approach performs the best? *Methods Ecol. Evol.* **5**, 961–970 (2014).

53. Upham, N. S., Esselstyn, J. A. & Jetz, W. Inferring the mammal tree: Species-level sets of phylogenies for questions in ecology, evolution, and conservation. *PLOS Biol.* **17**, e3000494 (2019).

54. Schliep, K. P. phangorn: phylogenetic analysis in R. *Bioinformatics* **27**, 592–593 (2011).

55. Revell, L. J. phytools: an R package for phylogenetic comparative biology (and other things). *Methods Ecol. Evol.* **3**, 217–223 (2012).

56. Stekhoven, D. J. missForest: Nonparametric Missing Value Imputation using Random Forest. *R package version 1.5* https://www.r-project.org (2022).

57. WoRMS Editorial Board. World Register of Marine Species. https://www.marinespecies.org (2020) doi:10.14284/170.

58. IUCN. The IUCN Red List of Threatened Species. Version 3. https://www.iucnredlist.org

(2017).

59. Holstein, J. worms: Retriving Aphia Information from World Register of Marine Species.

https://CRAN.R-project.org/package=worms (2018).

60. Chamberlain, S. rredlist: 'IUCN' Red List Client. https://CRAN.R-

project.org/package=rredlist (2020).

61. Chamberlain, S. A. & Szöcs, E. taxize: taxonomic search and retrieval in R.

F1000Research, 2:191 https://f1000research.com/articles/2-191/v2 (2013).

62. Jeschke, J. M. & Kokko, H. The roles of body size and phylogeny in fast and slow life

histories. *Evol Ecol* **23**, 867–878 (2009).

63. Revell, L. J. Size-Correction and Principal Components for Interspecific Comparative

Studies. *Evolution* **63**, 3258–3268 (2009).

64. Díaz, S. *et al.* The global spectrum of plant form and function. *Nature* **529**, 167–171

(2016).

65. Bueno, C. G. *et al.* Reply to: The importance of trait selection in ecology. *Nature* **618**,

E31–E34 (2023).

66. Carmona, C. P., Bello, F. de, Mason, N. W. H. & Lepš, J. Traits Without Borders:

Integrating Functional Diversity Across Scales. *Trends Ecol. Evol.* **31**, 382–394 (2016).

67. Carmona, C. P., de Bello, F., Mason, N. W. H. & Lepš, J. Trait probability density

(TPD): measuring functional diversity across scales based on TPD with R. *Ecology* **100**,

e02876 (2019).

68. Duong, T. ks: Kernel Density Estimation and Kernel Discriminant Analysis for

Multivariate Data in R. *J. Stat. Softw.* **21**, 1–16 (2007).

69. Duong, T. ks: Kernel Smoothing. https://CRAN.R-project.org/package=ks (2022).

70. Carmona, C. P. TPD: Methods for Measuring Functional Diversity Based on Trait

Probability Density. https://CRAN.R-project.org/package=TPD (2019).

71. Mason, N. W. H., Mouillot, D., Lee, W. G. & Wilson, J. B. Functional richness,

functional evenness and functional divergence: the primary components of functional

diversity. *Oikos* **111**, 112–118 (2005).

72. Wilhelm, S. & Manjunath, B., G. tmvtnorm: A Package for the Truncated Multivariate

Normal Distribution. *R J.* **2**, 25 (2010).

73. Oksanen, J. *et al.* vegan: Community Ecology Package. https://CRAN.R-

project.org/package=vegan (2020).

74. Germain, R. R. *et al.* Changes in the functional diversity of modern bird species over the

last million years. *Proc. Natl. Acad. Sci. U.S.A.* **120**, e2201945119 (2023).

75. Micó, E. *et al.* Contrasting functional structure of saproxylic beetle assemblages

associated to different microhabitats. *Sci. Rep.* **10**, 1520 (2020).

76. Traba, J., Morales, M. B., Carmona, C. P. & Delgado, M. P. Resource partitioning and

niche segregation in a steppe bird assemblage. *Community Ecol.* **16**, 178–188 (2015).

77. Reader, S. M. & Laland, K. N. Social intelligence, innovation, and enhanced brain size in

primates. *Proc. Natl. Acad. Sci. U.S.A.* **99**, 4436–4441 (2002).

78. Soria, C. D., Pacifici, M., Di Marco, M., Stephen, S. M. & Rondinini, C. COMBINE: a

coalesced mammal database of intrinsic and extrinsic traits. *Ecology* **102**, e03344 (2021).

79. Anderson, D. M. & Gillooly, J. F. Evaluating the tradeoff between offspring number and

survivorship across fishes, amphibians, reptiles and mammals. *Oikos* **130**, 798–807

(2021).

80. Davis, R. W. Return to the Sea: The Evolution of Marine Mammals. in *Marine*

Mammals: Adaptations for an Aquatic Life (ed. Davis, R. W.) 7–27 (Springer

International Publishing, 2019). doi:10.1007/978-3-319-98280-9_2.

81. Berta, A., Churchill, M. & Boessenecker, R. W. The Origin and Evolutionary Biology of

Pinnipeds: Seals, Sea Lions, and Walruses. *Annu. Rev. Earth Planet. Sci.* **46**, 203–228

(2018).

82. Wood, S. N. *Generalized Additive Models: An Introduction with R, Second Edition*.

(CRC Press, 2017).

83. R Core Team. R: A Language and Environment for Statistical Computing. (2020).

Acknowledgements

C.P.C and EB were supported by the Estonian Research Council grant (PSG293) and the
European Regional Development Fund via the Mobilitas Pluss programme (MOBERC40).
P.C.L. was supported by the European Union-Next Generation EU Maria Zambrano Program
(ZAMBRANO 21-26). R. S-G. was supported by a NERC Independent Research fellowship
(NE/M018458/1).

Author Contributions

P.C.L., R.S-G., and C.P.C. conceived the core ideas behind this paper, all authors provided
fundamental inputs for its final development. E.B collected all data and performed the
literature search needed to sort species in environmental realms. E.B performed the literature
search needed to collect data on evolutionary divergence, assisted by R.S-G. E.B. analysed
the data with fundamental inputs from C.P.C., P.C.L, and R.G-S. All authors contributed to
interpretation of the results. E.B. and C.P.C. led the writing of the manuscript which was
edited by all authors.