Behavioral and Neural Resilience to Isolation in a Eusocial Insect

Billie C. Goolsby[1,2]*, E. Jordan Smith[1], Isabella B. Muratore[1,3], Zach N. Coto[1], Mario L.

Muscedere[1], James F. A. Traniello[1]

[1] Department of Biology, Boston University, Boston, MA, 02215, USA

[2] Department of Biology, Stanford University, Stanford, CA, 94305, USA

[3] Department of Biological Sciences, New Jersey Institute of Technology, NJ, 07102, USA

*Corresponding authors:

Billie C. Goolsby
Department of Biology
Stanford University
371 Jane Stanford Way
Stanford, CA
bgoolsby@stanford.edu
23
24 James F. A. Traniello
25 Department of Biology
26 Boston University
27 Boston, MA 02215
28 jft@bu.edu
29
30

Abstract

Visual Abstract. Newly eclosed (callow) *Camponotus floridanus* minor workers were subjected to variable periods of isolation to determine whether and how diminished social experience and isolation influence the development of the brain, including compartment volumes, biogenic amine levels, and behavioral performance.

Social experience early in life appears to be necessary for the development of species-typical behavior in animals ranging from insects to primates. Isolation during critical periods of maturation has been shown to impact behavior, gene expression, and brain development in vertebrate and invertebrate clades, yet remarkable resilience to social deprivation, senescence, and loss of sensory input have been demonstrated in some ant species. We reared workers of *Camponotus floridanus* under conditions of social isolation for increasing time periods up to 45 days and evaluated their behavioral performance, quantified brain development and biogenic amine levels, and compared results of isolated workers with control workers that had natural social contact during development. We found that brood care and foraging performance by isolated workers were unaffected by a lack of social contact. Antennal lobe volume loss occurred in ants experiencing longer isolation periods, while the size of the mushroom bodies, which function in higher-order sensory processing, increased over time after eclosion and were not different from mature controls. Titers of the neuromodulators serotonin, dopamine, and octopamine remained

stable in isolated workers. Our results indicate that workers of *C. floridanus* are largely robust to

the deprivation of social contact early in life.

Introduction

Social experience shapes individual behavior. Exposure to social stimuli is critically

important to species-typical behavioral (Taborsky et al., 2012; Lévy et al., 2003), cognitive

(Amitai et al., 2014) and neural development (Smith et al., 2010; Wang et al., 2022), as well as

neuroimmune function (Donovan et al., 2020; Sallet et al., 2011; Serra et al., 2000; Mashoodh et

al., 2012; Hostinar et al., 2012). Across diverse clades, behavioral performance can be negatively

impacted if individuals are deprived of social contact when ontogenic trajectories are most

sensitive to receiving sensory inputs (Wang et al., 2022; Boyce et al., 2012). A lack of social

contact during these critical periods, therefore, can potently disrupt development across multiple

biological scales (Rodriguez-Romaguera & Stuber, 2018; Calati et al., 2019; Vora et al., 2022).

Although research on the effects of social isolation has historically focused on vertebrates,

eusocial insects have emerged as important models to examine the influences of social deprivation

in early adult life on behavior, brain development, and brain gene expression (Boulay and Lenoir,

2001; Chen, 1937; Seid and Junge, 2016; Scharf et al., 2021; Lihoreau et al., 2009; Muscedere et

al., 2013; Hewlett et al., 2018; Koto et al., 2015; Wang et al., 2022). Workers are deeply embedded

within the social organization of a colony, suggesting a high degree of developmental dependency.

Workers socially interact during physical encounters, trophallaxis (oral food exchange),

allogrooming, nursing, foraging, and defense. Our understanding of how the complex social fabric

of insect societies regulates behavioral and neuroanatomical development is limited (Kronauer and

Libbrecht, 2018; Godfrey and Gronenberg, 2019; Wang et al., 2022). Social isolation in ants, for

example, appears to result in significantly altered brain growth relative to that of nestmates

developing in typical social environments (Seid and Junge, 2016). Neuromodulators that function

in behavioral development may become dysregulated (Boulay et al., 2000) if workers are deprived

of social contact, potentially affecting age-related responsiveness to signals and cues that guide

the behavior of mature workers (Seid and Traniello, 2005; Muscedere et al., 2013).

Neuromodulators and receptor profiles that regulate individual and collective behavior (Kamhi

and Traniello, 2013; Kamhi et al., 2017; Friedman et al., 2018, 2020; Maloney, 2021) may also be

altered. Nevertheless, the brains of worker ants can be resilient to senescence (Giraldo et al., 2016)

and injury (Waxman et al., 2017). This robustness to aging and physical damage in ants raises

questions concerning their ability to compensate for stress due to social isolation during periods

that are likely critical for behavioral and neural development.

In most ant species, young adult workers (callows) eclose in a socially rich environment of

adult nestmates and immatures in their parent colony. Callows are distinguished from their mature

siblings by their light coloration due to unsclerotized cuticule and relatively low activity levels

(Morel, 1986; Muscedere et al., 2009). Species-typical behavioral maturation is usually incomplete

immediately after eclosion, with apparently few exceptions (Traniello, 1978), so manipulations of

callow social experiences can be used to experimentally examine the ontogeny of behavior. Social

interaction between callows and their siblings influences pheromone responsiveness and temporal

behavioral development (Topoff and Mirenda, 1978; McDonald and Topoff, 1985; Jaisson and

Fresneau, 1978; Morel, 1986; Richardson et al., 2021). The effects of being deprived of such inputs

are not consistently negative, as there is evidence of behavioral and neural robustness. For

example, 10-day old *Pheidole dentata* minor workers deprived of brood contact nursed as

efficiently as same-aged control workers and did not differ neuroanatomically or neurochemically

(Muscedere et al., 2013). The question nevertheless remains as to whether socially isolated ant

workers are neurally prepared for behavioral development independent of experience.

The genus *Camponotus* has served as a model to examine neuroethological impacts of

social deprivation in ants. In *C. floridanus* and *C. fellah*, neuroanatomical, neurochemical, and

behavioral changes in workers after periods of social isolation have been documented, including

decreases in brain volume and behavioral deficits (Seid and Junge, 2016; Boulay and Lenoir, 2001;

Boulay et al., 2000; Wada-Katsumata et al., 2011). Here, we integrate analyses of task

performance, brain volume, the growth of neuropils specialized for sensorimotor functions, and

titers of brain biogenic amine neuromodulators to examine how workers behaviorally and

neurobiologically develop during increased periods of isolation and determine if workers are

susceptible or resilient to a lack of social contact.

Methods

Colony collection and culture

Three queenright colonies of *C. floridanus* were collected in or near Gainesville, Florida

in 2012 and reared in Harris environmental chambers at 25°C and 40-55% relative humidity on a

12:12 light cycle. Parent colonies from which workers were sampled had a minimum of 100 minor

and 100 major workers. Artificial nests were made from test tubes 75% filled with water and fitted

with a tight cotton plug and placed in Fluon®-lined (Bioquip, Rancho Dominguez, CA) plastic

nest boxes 16x11x6 cm to 32x22x6 cm depending on colony size. Colonies were fed 1M sucrose

on saturated cotton balls and varied protein sources (mealworms, scrambled eggs, and fruit flies)

every other day in rotation.

Worker social isolation

Five groups of callow minor workers were collected from each of three parent colonies.

Randomly selected callows were isolated from nestmates for 2 [2 +/- day], 5 [5 +/- day], 20 [20

+/- 2 days], 30 [30 +/- 4 days], or 45 [45 +/- 8 days] days in separate Fluon®-lined boxes (16 x 11

x 6 cm). Isolated workers ("isolates") were provided similar conditions in smaller nest boxes and

the same diet as parent colonies. After the prescribed isolation period, isolates were assayed for

behavioral performance and subsequently sacrificed for immunohistochemical analysis of brain

structure or quantification of brain biogenic amine titers. Callow and mature minor workers

("matures") were collected and assayed using the same methods. We calculated a survival rate of

63.7% for isolated workers. Because of the mortality of workers experiencing longer periods of

isolation, sample sizes for 30- and 45- day isolates were generally smaller. Head width, measured

as the widest distance across the eyes, served as a proxy for body size. To control for body size,

which may correlate with brain size, we compared the head widths of workers across isolation

periods and control groups. We found no significant differences (p=0.429, R^2=0.116, F=1.132,

Z=0.206, SS type 3, **Supp Fig 1)** and our sampling of workers for assignment to isolation or control

treatments was thus unlikely to influence brain mass or volume.

Behavioral performance assays

Isolated and control workers (n = 16 [2 day], 19 [5 day], 20 [20 day], 6 [30 day], 5 [45

day], 8 [callow] and 13 [mature]) were exposed to a randomized series of assays to determine their

ability to recognize and care for brood, and foraging, tasks that affect colony fitness.

Brood recognition and care

To determine if isolates differ in the ability to recognize and care for immature siblings with similar efficacy as socialized matures and callows, we evaluated their interactions with pupae. Each focal worker was carefully placed in an arena (14 cm diameter Petri dish) beneath a 30 x 10 mm red plastic hollow tube that provided a dark area into which workers could move pupae. Three pupae were added at 3 locations equidistant from the darkened chamber before a worker was introduced to the arena. Workers relocated pupae either inside or outside of the dark chamber or otherwise moved alone without interaction. Pupae were considered to be aggregated ("piled") if workers moved two or more together. Workers that moved two or more pupae began doing so immediately. We did not assign lower scores to workers that clustered pupae but did not move them into the darkened area. Worker interactions with pupae were observed directly for 5 minutes, then after an additional 15 minutes the placement of pupae was recorded. This assay for brood recognition and care was measured on a 7-point scale: 1. no interaction or avoidance; 2. olfactory (antennal) investigation in the first 5 minutes; 3. mouthpart (mandibles, palps) contact in the first 5 minutes; 4. no pupae moved before 5 minutes but at least one pupa moved during the subsequent 15-minute period; 5. one pupa moved in the first 5 minutes; 6. two pupae moved within the entire trial; 7. three pupae moved during the entire trial. Workers received scores of 5-7 independent of where pupae were transported. Higher scores indicate brood recognition and movement of more pupae.

Predatory Behavior

A single worker was carefully placed in a Fluon®-lined Petri dish (9cm) and allowed to acclimate for 2 minutes prior to assay initiation. Using Dumont No. 5 fine forceps sanitized with

ethanol and dried, a live fly was then offered to the worker and the response was observed for 2

minutes. The degree of predatory behavior was scored on a 5-point scale: 1. no response or

avoidance; 2. olfactory (antennal) investigation; 3. mandibular flaring; 4. latent attack (delayed

and intermittent); and 5. immediate attack (persistent predatory attack from first encounter with

prey). The assay arena was also sanitized with ethanol between trials to minimize influence from

social signals or prey odors.

Locomotion

To examine if isolation affected the physical capability to pile pupae or forage, we used

locomotion as a simple and robust proxy to broadly assess neuromuscular impairment (Giraldo et

al., 2016). Focal workers were placed in the center of a 10 cm diameter Fluon®-lined Petri dish

divided into 4 equal-size quadrants with a crosshair pattern drawn beneath the bottom and

acclimatized for 2 minutes. After the acclimation period, worker movement was observed for 5

minutes, and the number of times a new quadrant was entered was recorded using a digit counter.

Movement between quadrants was only scored when the worker passed the division line by at least

one body length.

Neuroanatomical and neurochemical measurements

Brain mass

To quantify isolation-associated changes in brain size and to enable scaling of biogenic

amine levels, we recorded and compared brain mass among three isolate groups (n = 9 [2 day], 10

[5 day], 12 [20 day], 10 [callows], and 12 [matures]). Brains were dissected in ice-cold insect

Ringer's solution after head width was measured. Brains were placed in a tin capsule of known

weight, dried at 55°C, and weighed (+/- 1 microgram) after 48 hours using a Sartorius CP2P

microbalance (Sartorius, Goettingen, Germany). Brains were weighed three times and averaged to

estimate brain mass. Brain that were statistical outliers (Z>2.5) due to technical difficulties in the

process of weighing were excluded from this analysis.

Immunohistochemistry and confocal microscopy

Intact brains (n = 9 [2 day], 11 [5 day], 16 [20 day], 3 [30 day], 11 [40 day], 5 [matures],

and 9 [callows]) were dissected in ice-cold Ringer's solution, and then immediately placed in ice-

cold zinc-formaldehyde fixative with shaking overnight. Fixed brains were then washed in HEPES

buffered saline 6 times (10 minutes/wash) and fixed in Dent's Fixative (80% MeOH, 20% DMSO)

for 1 hour. Brains were placed in 100% methanol and stored at -17 C° for 1-3 weeks until

processed. Brains were next rinsed 6 times (10 minutes/rinse) with 3% Triton X-100 in 1X PBS

(3% PBST). Brains were stored overnight in 5% normal goat serum/0.2% PBST/0.005% sodium

azide for blocking. After blocking, brains were incubated for 4 days RT with an anti-synapsin

antibody to stain neuropils (1:30, Development Studies Hybridoma Bank, Iowa City, Iowa) in

constant nutation. Brains were washed with 0.2% PBST (6x10 minutes), then incubated in

secondary   antibody   (1:100   AlexaFluor   goat   α-mouse)   in   5%   normal   goat

serum/0.2%PBST/0.005% sodium azide for 4 days at RT with constant nutation. Brains were

washed in 0.2% PBST (6x10 minutes), then dehydrated through an ethanol and PBS series (30,

50, 70, 95, 100% ethanol in 1XPBS). Brains were cleared and immersed in methyl salicylate and

mounted on stainless steel glass-windowed slides for confocal imaging.

Brains were imaged with a Nikon C2 + Si spectral laser scanning confocal microscope

(Nikon, Melville, NY, USA) with sections at approximately 1.5 um throughout the entire brain.

Images were manually annotated using Amira 6.0 software to quantify neuropil volumes. The

individual who annotated brains was blind to brain sample category, thus minimizing human bias.

The margins of major functionally specialized neuropils were identified visually or with magic

wand tools in Amira. Every eighth frame was annotated manually. Frames without labels were

labeled using Biomedisa interpolation (Lösel et al., 2020), then evaluated and edited by the

annotator for accuracy. We labeled volumes of the optic lobe (OL), antennal lobe (AL), mushroom

body (MB), lateral calyx (MB-LC), and medial calyx (MB-MC), peduncles (MB-P), central

complex (CX), subesophogeal zone (SEZ), and rest of the central brain (ROCB). ROCB describes

tissue composed of the superior neuropils, ventromedial neuropils, ventrolateral neuropils, and the

lateral horn (Ito et al., 2014). For each brain compartment, volumes of only one intact hemisphere

were recorded. If both hemispheres were intact, the labeler chose randomly.

We compared the relative volumes of brains across all groups and scaling relationships

among brain compartments. A detailed description of brain anatomy and development in relation

to age and experience in *C. floridanus* is presented in Gronenberg et al. (1996). Relative volumes

were calculated by dividing the volume of each focal compartment by the ROCB volume.

Damaged compartments were excluded from analysis. If one or more compartments or the ROCB

was damaged, the sample was excluded. The MB-LC and MB-MC exhibited similar volumetric

trends and thus were combined to one metric (MB-C). As a sensitivity test for using the ROCB as

proxy for total brain volume, we ran additional analyses using total brain volume for scaling which

indicated that using the ROCB for scaling reflected the same trends as scaling to total brain size

(**Supplementary Methods**).

Quantification of brain biogenic amines

We quickly dissected workers (n = 11 [2 days], 11 [5 days], 13 [20 days], 3 [30 days], and

11 [mature workers]) to quantify levels of serotonin (5HT), dopamine (DA), and octopamine (OA)

using isocratic, reversed-phase high-performance liquid chromatography with electrochemical

detection (HPLC-ED). Monoamine quantification was performed blind to treatment. The HPLC-

ED system (formerly ESA, Inc., Chelmsford, MA, USA) included a model 584 pump, a model

MD-150 (3 x 150 mm) reversed-phase analytical column, a model 5011A dual-channel

coulometric analytical cell, and a Coulochem III electrochemical detector. Brains were dissected

in less than 5 minutes in ice-cold Ringer's solution, homogenized in mobile phase manually with

a plastic polypropylene pestle in a centrifuge tube, and centrifuged. The supernatant was then

injected into the HPLC column, and monoamine levels were quantified with reference to external

standards. Mobile phase was formulated as 50 mM citrate/acetate buffer, 1.5 mM sodium dodecyl

sulfate, 0.01% triethylamine, and 25% acetonitrile in MilliQ water (Muscedere et al., 2013).

Multiple daily runs of the external standards accounted for any changes in ambient conditions.

Statistical analysis and data visualization

All statistics analyses were performed in RStudio (RStudio version 4.2.3, Posit Team

2023). For generalized linear mixed models, we used the package '*RRPP*' (Collyer & Adams,

2019). We tested for statistical significance using ANOVA with residual randomization in a

permutation procedure of 1000 iterations and through the estimation method of ordinary least

squares, given combined sample size of 0 or more. Following ANOVAs, *post-hoc* tests were done

on pairwise comparisons using t tests with pooled SD with p value adjustment using the Bonferroni

method. Depending on the sample size of 30-day and 45-day isolates, we included type III error

to account for bias in sample size. Figures were created using the package '*ggplot2*' (Wickham,

2016). Color corrections and figure assembly were done in Adobe Illustrator (Adobe Illustrator

2023). Illustrations of ants in **Fig 1A** and **Supp Fig 1B** are courtesy of BioRender.com.

Results

263
Effect of Isolation on Predatory Behavior

Worker predatory behavior was significantly different between isolated groups (p=0.001,

R^2=0.294, F=5.540, Z=3.6525, SS type 1, **Fig 1A-B**). Mature workers preyed on fruit flies

significantly more than callows, 2-day isolates, 5-day isolates, 20-day isolates, 30-day isolates, but

not 45-day isolates (p<0.001, <0.005, <0.001, <0.05, <0.05, respectively; Bonferroni correction).

All other pairwise comparisons were not significantly different (p=0.149-1.0). Mature workers

moved significantly more than callows and isolates but did not show predatory behaviors

significantly different from those of 30- and 45-day isolates **(Fig 1B)**. Therefore, callows and 2-

day isolates interacted with prey in a similar manner, showing avoidance or olfactory investigation.

In contrast, older isolates (45-day, e.g.) had higher predatory scores resembling those of mature

worker controls, scoring on average 2.80 and 3.38, respectively. Workers that experience less than

45 days of isolation had significantly lower predatory behavior scores than mature workers, but

45-day isolates scored similarly to mature workers, suggesting that increased isolation did not

impair predatory behavior.

Effect of Isolation on Brood Recognition and Care

Isolated and control workers interacted with pupae to the same degree and were likely

recognized as immature nestmates by both groups. Brood-care performance was significantly

different among experimental groups (p=0.009, R^2=0.201, F=3.3518, Z=2.4572, SS type 1, **Fig**

1A, C), but in *post-hoc* tests, only mature workers and 2-day isolates differed significantly

(p=0.014, Bonferroni correction). All other pairwise comparisons were not significant (p=0.06-

1.0). With increased duration of isolation, brood-care scores became similar to those of callows

and mature worker controls. Isolated and control workers did not differ in their tendency to move

pupae into a darkened location.

288
Figure 1. Isolation effects on task performance. (A) Illustration of assays to assess brood-care and
predatory behaviors. **(B)** Predation scores of workers isolated for varying time periods compared to callows
and mature worker controls. Darkening shades of red correspond with increased time in isolation. **(C)**
Brood-care scores of isolates compared to callows and mature worker controls. Darkening shades of blue
correspond to increased age and time of isolation. 2-day and mature workers were the only groups that
differed significantly in brood care. X = average; horizontal line = median. Sample sizes are above box
plots. <0.001 = ***; 0.001-0.01 = **; 0.01-0.05 = *; >0.05 = ns. Socially typical mature workers and callow
controls are the darkest and lightest colors, respectively.

Mature workers had significantly higher levels of locomotion than isolate groups and callows, with

the exception of 30-day isolates (p=0.001, R^2=0.486, F=12.595, Z= 5.0508 SS type 1, [**Supp Fig**

1B-C, Supp Table 1-3]). Mature workers significantly differed from other experimental groups

(callow, 2, 5, 20, 30, 45: all p<0.001, Bonferroni correction) except for 30-day isolates. No other

pairwise comparisons were statistically significant (p>0.05).

Brain Size and Compartmental Allometry

Brain mass did not significantly differ between isolated and control workers (p=0.359,

R^2=0.092, F=1.09, Z=0.356, SS type 1, **Fig 2A)**. Size trajectories of brain compartments are

illustrated in **Fig 2B**.

Figure 2. Effects of isolation on brain growth. (A) Brain mass of workers isolated for 2, 5, and 20 days compared to callow and mature worker control groups. Sample sizes of each group are above box plots. X = average; horizontal line = median. **(B)** Spider plot of brain compartment volumes scaled to ROCB volume in callows, 2-day workers, 45-day workers, and mature workers. Numbers on spider plot are proportions relative to ROCB volume. * indicates p<0.01; p >0.05 = ns.

MB-C increased in volume with age and increased periods of isolation, becoming similar in size to the MB-C of mature workers. However, workers enduring longer periods of isolation had smaller AL volumes compared to those of mature workers. AL volumes were significantly different (p=0.003, R^2=0.323, F=4.539, Z=3.1128; **Fig 2A**): mature workers had significantly larger AL volumes relative to the ROCB than 30-day and 45-day isolates (p=0.017, 0.00033 respectively, Bonferroni correction). The average relative volume of the MB-P remained consistent throughout isolation periods with no significant changes, indicating that impact on the MB was limited to the calyces (p=0.295; R^2=0.127, F=1.2878, Z=0.544; **Fig 3B**), which showed modest growth during isolation (p=0.001, R^2=0.477, F=8.0699, Z=4.213; **Fig 3C**). Mature workers had significantly higher relative MB-C volumes than callows, 2-day isolates, and 5-day isolates (p<0.0001, <0.0001, <0.004, respectively; Bonferroni correction), but not 45-day isolates (p=0.0622). MB-C volumes of 45-day isolates were also significantly larger than callows (p=0.038, Bonferroni correction). The MB-C volumes of 20-day isolates were also significantly

larger than 2-day isolates and callows (p=0.0169, p=0.0107, Bonferroni correction). All pairwise

analyses for compartmental volumes can be found in **Supplementary Tables.** Other brain

compartments did not significantly change with isolation (p=0.228-0.871; **Fig 3D-F**).

Figure 3. Isolation and growth of brain compartments. A-E: brain compartment volumes (relative to ROCB) for isolated and control workers. Sample sizes of each group are above box plots. X in each box plot represents the average and horizontal line the median of each group **(A)** AL; **(B)** MB-P; **(C)** MB-C; **(D)** CX; **(E)** OL; **(F)** SEZ. p <0.001 = ***; p<0.001-0.01 = **; p<0.01-0.05 = *; p>0.05 = ns.

Biogenic amine levels

Brain titers of OA, DA, and 5HT significantly increased with longer periods of isolation

(**Fig 4A-C**); levels of 30-day isolates were similar to mature minor workers. DA titers significantly

differed across isolation treatments and controls (p=0.001, R^2=0.611, F=17.481, Z=5.340, SS type

3, **Fig 4A**) and significantly differed between mature worker controls and 2, 5, and 20-day isolates,

but not 30-day isolates (mature vs. 2, 5, 20-day isolates, p<0.00001, <0.00001, <0.0001,

respectively; Bonferroni correction). OA exhibited a similar pattern (p=0.001, R^2=0.723,

F=42.552, Z=7.844, SS type 3, **Fig 4B**). OA titers significantly increased between isolated worker

groups except for 2-day isolates vs. 5-day isolates, 5-day isolates vs. 20-day isolates, and 20-day

isolates vs. 30-day isolates. 5HT titers also significantly increased with longer periods of isolation

(p=0.001, R^2=0.744, F=109.136, Z=9.980, SS type 3, **Fig 4C**), except for 20-day isolates vs. 30-

day isolates.

While absolute levels of DA, OA, and 5HT increased over time, their ratios increased or

decreased compared to those recorded in mature workers (**Fig 4D-F**). OA:5HT significantly

decreased with time and extended periods of isolation (p=0.001, R^2=0.326, F=7.805, Z = 3.672)

with no significant difference between 30-day isolates and mature workers (**Fig 4D**). OA:DA did

not significantly change with isolation (p=0.989, R^2=0.00772, F=0.0855, Z=-2.220), nor did

5HT:DA (p=0.221, R^2=0.102, F=1.4831, Z=0.755) except 2-day isolates being having

significantly lower 5HT:DA ratios compared to mature workers (p=0.046, Bonferroni correction).

DA titers relative to total amines levels did not significantly differ across isolation groups

and mature workers (p=0.434, R^2=0.0705, F=0.968, Z=0.164). Relative OA levels, in contrast,

were significantly different (p=0.015, R^2=0.243, F=3.371, Z=2.126), including 2-day vs. 20-day

and 2-day vs. mature workers were significantly different (p=0.031, 0.034, respectively,

Bonferroni correction). Relative 5HT quantities were significantly different (p=0.015, R^2=0.194,

F=3.471, Z=2.083) between 2-day vs. 20-day isolates and 2-day vs. mature workers (p<0.001,

<0.005 respectively, Bonferroni correction).

363
364   **Figure 4. Isolation effects on monoamine levels (A-C) and ratios of DA, OA, and 5HT (D-E).** The
365   transition from green to blue corresponds to increasing isolation time periods. The deepest blue boxplot
366   corresponds to mature worker controls. Averages, medians, and samples sizes as in Figure 3. p <0.001 =
367   ***; p<0.001-0.01 = **; p< 0.01-0.05 = *; p>0.05 = ns.

368
Discussion

Overall, the behavior of *C. floridanus* workers isolated for varying durations after eclosion

was comparable to that of control workers that developed in a natural social context. Two-day

isolates showed predatory behavior similar to that of callow control workers, and predatory

behavior by 45-day isolates was similar to that of mature control workers, suggesting foraging was

not impaired by isolation. Social isolation resulted in a significant difference in brood care – as

measured in our assay – only between 2-day isolates and mature control workers, but otherwise

had no effect. This result may reflect the development of behavioral competencies associated with

biomechanical and physiological maturation (Muscedere et al., 2011). Workers in all isolation

treatment groups showed significantly decreased locomotion relative to mature workers except for

30-day isolates (**Supp Fig 1B-C**), consistent with Wang et al. (2022) for isolated bumblebees but

different from *Cataglyphis niger* ant workers (increased movement following isolation and then

decreased movement [Ahronberg and Scharf 2021]) and *Camponotus fellah* workers (increased

movement after isolation [Koto et al., 2015]). This decline in movement did not appear to influence

brood care or predatory behavior in isolated *C. floridanus* workers and thus did not appear to be a

cost of isolation.

C. floridanus worker brains normally increase in size during a 10-month period after adult

eclosion, the greatest growth rate (nearly 100%) occurring in the MBs and ALs (Gronenberg et al.,

1996). We found no significant difference in mass between isolates, control callows, and mature

workers, indicating that isolation did not significantly affect brain size during the 45-day

assessment period**.** Furthermore, older isolates and socially typical mature workers generally had

larger relative MB-C volumes compared to workers isolated for shorter time periods and socially

typical callows. MB-C size in 45-day isolates was not significantly different from that of mature

controls and 2-day isolates, suggesting gross neuroanatomical development remained stable under

conditions of isolation. These findings differ from the results of Seid et al. (2016), which showed

that MBs of isolated *C. floridanus* workers do not change in volume relative to total brain size,

whereas MBs increased in our study. MB-C resilience through 71 days of isolation has been found

in paper wasps (Jernigan et al., 2021). Using bumblebees as a model, Wang et al. (2022) associated

expression differences in juvenile hormone-related genes and transcriptomic network

dysregulation with isolation-induced variation in brain volume. Bumblebees isolated after eclosion

for 9 days showed significant differences in the variance of MB volume rather than average volume

compared to socially typical bees, suggesting growth destabilization in some, but not all, workers.

Our study identified resilience in brain growth, possibly attributable to different mechanisms of

developmental programming in *C. floridanus*. We note our findings do not demonstrate a complete

lack of influence of isolation on brain size and structure because we did not determine temporal

patterns of growth that could indicate the development of the MBs or other compartments

accelerated or slowed differentially over time in isolates and/or socially typical workers. Our

results nevertheless support the hypothesis that MB development is resilient to social isolation.

AL volume decreased with periods of isolation approximately 5 days or longer; 45-day

isolates had, on average, smaller ALs than control callows and mature workers. This decline in AL

size is similar to the findings of Wang et al. (2022) on bumblebees. Although AL investment

decreased with isolation duration in *C. floridanus,* workers still maintained olfactory abilities

necessary for performing the tasks we assayed. Isolation did not impact OL volume in our study

of *C. floridanus*, although it did reduce OL volume in paper wasps, which integrate visual

information used in conspecific facial recognition (Jernigan et al., 2021; Pardo-Sanchez, 2022).

The CX, which functions in sensorimotor processing in insects (Turner-Evans and Jayaraman,

2016; Pisokas et al., 2020), did not change significantly in volume in isolated workers. These

results also differ with those of Wang et al. (2022) for isolated bumblebees. The volume of the

SEZ, which regulates mouthparts (Miroschnikow et al., 2020) and is likely involved in the control

of brood-care motor functions, did not significantly change with isolation.

Worker behavioral maturation, including repertoire expansion and a corresponding

increase in brain size, correlates with increasing monoamine levels and the development of

serotonergic circuitry in some ants (Seid and Traniello, 2005; Seid et al., 2008; Muscedere and

Traniello 2012; Wnuk et al., 2011). Our *C. floridanus* data indicate that levels of OA, DA, and

5HT increase with age in the brains of isolated workers, despite a lack of social contact. Indeed,

30-day isolates had monoamine levels most like those of mature workers, whereas 2-day isolates

were least like mature workers. Ratios of biogenic amines can significantly change with age in

ants and may influence the development of age-related behavior (Seid and Traniello, 2005) by

modifying response thresholds. However, we found no significant differences between older

isolates and mature controls, indicating that isolation does not significantly alter neuromodulator

ratios. Boulay et al. (2000) and Wada-Katsumata (2011) found that 5HT and DA levels were robust

to social isolation. Boulay et al. (2000) hypothesized OA may reduce the effects of social

deprivation, acting homeostatically to maintain social cohesion. Whether octopaminergic

signaling has a stimulatory or inhibitory effect on prosocial behaviors such as trophallaxis, which

may be affected by periods of social isolation, remains unclear.

Although it may be generally assumed that smaller size reduces competence, we do not

know how whole-brain or compartmental volumes affect behavioral ability. Indeed, the

relationship between brain size and behavioral capacity is actively debated (e.g., Chittka and

Nevin, 2009). In ants, elaboration of the MB appears to represent increased demands for higher-

order processing (Kamhi et al., 2016; Godfrey and Gronenberg, 2019; Muratore et al., 2022) and

larger OLs are correlated with more complex visual ecologies (Sheehan et al., 2019; Arganda et

al., 2020). Larger ALs have been associated with age-related olfactory ability (Muscedere et al.,

2012), although Hart et al. (2023) showed that ant alarm pheromone response is encoded by less

than 10 of a total of approximately 500 AL glomeruli, a result that suggests that the relationship

of neuropil size and behavior is unclear. We also do not understand how circuit configuration,

activation, synaptic strength, or other neural processes in the ant brain connectome are influenced

by worker isolation.

Neuromodulation can restore sensorimotor functions following injury (Zhang et al., 2021).

Neuromodulators can alter synaptic functions to significantly change circuitry (Marder, 2012;

Sillar et al., 2014), increase the robustness of network functions (Haddad and Marder, 2018) and

their ability to recover from disruption and/or stress (Golowasch, 2019; Thuma et al., 2013). In the

ant *Pheidole dentata*, monoamine levels continue to increase in titer in aging workers, which do

not appear to decline in task performance capability (Giraldo et al., 2016). Given that DA, OA,

and 5HT can flexibly reshape circuits, neuromodulators may reduce isolation-induced

neuroanatomical deficits in ants, stabilizing circuit dynamics or returning them to a stable state if

disrupted. Although we did not identify an increase in monoamine levels in isolates, this does not

preclude neuromodulatory compensation to maintain circuitry and behavioral functions under

conditions of social deprivation.

Conclusion

The results of the present study and prior research on the impacts of senescence on

neurochemistry, neuroanatomy, apoptosis, and behavior (Giraldo et al., 2016), deprivation of

experience (Muscedere et al., 2013), and reduced olfactory processing ability (Waxman et al.,

2017), suggest that ant brains have an unusual capacity to buffer insults of aging, social isolation,

and physical damage to sensory appendages. The 36% mortality of isolated *C. floridanus* workers

we found in the present study indicates a significant negative effect that may, however, be due

more to social processes affecting nutrition and energy availability (Koto et al., 2015; Ahronberg

and Scharf 2021) than neurobiological deficits resulting from isolation. Workers that survive

neurobiological and physiological bottlenecks appear able to recover from challenges to brain

development, which in *C. floridanus* are dependent on age as well as experience (Gronenberg et

al., 1996). Age-related increases in brain size in ants appear to be due to stable programmed axonal

and dendritic growth processes. If sensory experience is lacking or temporally atypical, brain

development may still proceed in part in the absence of inputs or may be delayed (e.g.,

Mackevicius et al., 2023). In our study, the effect of social isolation in *C. floridanus* appears to

reflect such robust programming, perhaps due in part to stabilizing effects of neuromodulators.

The resilience of worker ants to challenges during development may functionally resemble

differences that enable some individuals – perhaps those with adaptive synaptic phenotypes

(Sakurai et al., 2014) or ion channel properties (Goaillard and Marder, 2021) – to withstand neural

damage and preserve behavior. We hypothesize that the ability to buffer stressors may involve

such variation in circuitry that characterizes clade-wide neuroarchitectures of the miniaturized

brains of ants, acknowledging that few species have been studied. Analyses of temporal patterns

of brain gene expression in *C. floridanus* isolated workers can identify transcriptomic changes

linked to brain development and influenced by social experience (Bonasio et al., 2010; Simola et

al., 2013) and possibly identify genetic mechanisms of resilience. Genomic studies of

phylogenetically and sociobiologically diverse ant species can also contribute to our understanding

of how workers are susceptible to and/or recover from social deprivation or are resilient to

isolation.

Acknowledgements

We thank Lloyd Davis for colony collection. This research was supported by National

Science Foundation grants IOS 1354291 and IOS 1953393 to JFAT, the Marion Kramer Award to

BCG, and the Boston University Undergraduate Research Opportunities Program. BCG was

supported by a HHMI Gilliam Fellowship (GT15685) and a National Institutes of Health Cellular

Molecular Biology Training Grant (T32GM007276). BCG thanks Dr. Katherine Fiocca for

manuscript feedback and Dr. Bryan Juarez for coding advice.

Author Contributions

BCG, IBM, and JFAT designed the study. BCG recorded behavior. BCG and IBM

immunohistologically prepared and imaged brains. EJS quantified brain volume. BCG, ZC, and

MM measured monoamine titers. BCG statistically analyzed volumetric data. BCG drafted the

manuscript. BCG, EJS, IBM, ZC, MM, and JFAT developed and edited the manuscript. JFAT

secured funding.

References

1. Ahronberg, A., & Scharf, I. (2021). Social isolation interaction with the feeding regime

differentially affects survival and results in a hump-shaped pattern in movement activity.

Behav Processes, *190*, 104460.

2. Amitai, N., Young, J. W., Higa, K., Sharp, R. F., Geyer, M. A., & Powell, S. B. (2014).

Isolation rearing effects on probabilistic learning and cognitive flexibility in rats. *Cogn Affect*

Behav Neurosci, 14(1), 388–406.

3. Arganda, S., Hoadley, A. P., Razdan, E. S., Muratore, I. B., & Traniello, J. F. A. (2020). The

neuroplasticity of division of labor: worker polymorphism, compound eye structure and brain

organization in the leafcutter ant *Atta cephalotes*. *J Comp Physiol A*, *206*(4), 651–662.

4. Bonasio, R., Zhang, G., Ye, C., Mutti, N. S., Fang, X., Qin, N., & Zhang, P. (2010). Genomic

comparison of the ants *Camponotus floridanus* and *Harpegnathos saltator*. *Science*, 329:

1068-1071.

5. Boulay, R., Soroker, V., Godzinska, E.J., Hefetz, A., & Lenoir, A. (2000). Octopamine reverse
the isolation-induced trophallaxis in the carpenter *Camponotus fellah*. *J Exp Biol,* 203: 513-
520.

6. Boulay, R., & Lenoir, A. (2001). Social isolation of mature workers affects nestmate
recognition in the ant *Camponotus fellah*. *Behav Processes*, *55*(2), 67–73.

7. Boyce, W., Sokolowski, M., & Robinson, G. (2012). Toward a new biology of social adversity.
PNAS, 109:17143-17148.

8. Calati, R., Ferrari, C., Brittner, M., Oasi, O., Olié, E., Carvalho, A. F., & Courtet, P. (2019).
Suicidal thoughts and behaviors and social isolation: A narrative review of the literature. *J*
Affect Disord, *245*, 653–667.

9. Chen, S. C. (1937). Social modification of the activity of ants in nest-building. *Physiol*
Zool, *10*(4), 420-436.

10. Chittka, L., & Niven, J. (2009). Are Bigger Brains Better? *Curr Biol*, *19*(21), R995–R1008.

11. Collyer ML, Adams DC (2019). "RRPP: Linear Model Evaluation with Randomized Residuals
in a Permutation Procedure. R package version 0.4.0." https://CRAN.R-
project.org/package=RRPP.

12. Donovan, M., Mackey, C. S., Platt, G. N., Rounds, J., Brown, A. N., Trickey, D. J., Liu, Y.,
Jones, K. M., & Wang, Z. (2020). Social isolation alters behavior, the gut-immune-brain axis,
and neurochemical circuits in male and female prairie voles. *Neurobiol Stress*, *13*, 100278.

13. Friedman, D. A., Pilko, A., Skowronska-Krawczyk, D., Krasinska, K., Parker, J. W., Hirsh,
J., & Gordon, D. M. (2018). The Role of Dopamine in the Collective Regulation of Foraging
in Harvester Ants. *IScience*, *8*, 283–294.

14. Friedman, D. A., York, R. A., Hilliard, A. T., & Gordon, D. M. (2020). Gene expression

variation in the brains of harvester ant foragers is associated with collective behavior. *Comm*

Biol, *3*(1), 100.

15. Giraldo, Y. M., Kamhi, J. F., Fourcassié, V., Moreau, M., Robson, S. K., Rusakov, A. &

Traniello, J. F.A. (2016). Lifespan behavioural and neural resilience in a social insect. *Proc R*

Soc B, 283(1822): 20152603.

16. Godfrey, R. & Gronenberg, W. (2019). Brain evolution in social insects: advocating for the

comparative approach. *J Comp Physiol A*, 205.1: 13-32.

17. Goaillard, J.-M., & Marder, E. (2021). Ion Channel Degeneracy, Variability, and Covariation

in Neuron and Circuit Resilience. *Annu Rev Neurosci, 44*(1), 335–357.

18. Golowasch, J. (2019). Neuromodulation of central pattern generators and its role in the

functional recovery of central pattern generator activity. *J Neurophysiol*, *122*(1), 300–315.

19. Gronenberg, W., Heeren, S., & Hölldobler, B. (1996). Age-dependent and task-related

morphological changes in the brain and the mushroom bodies of the ant *Camponotus*

floridanus. J Exp Bio, *199*(9)-2011-2019.

20. Haddad, S.A., & Marder, E. (2018). Circuit Robustness to Temperature Perturbation Is Altered

by Neuromodulators. *Neuron*, *100*(3), 609-623.e3.

21. Hart, T., Frank, D. D., Lopes, L. E., Olivos-Cisneros, L., Lacy, K. D., Trible, W., Ritger, A.,

Valdés-Rodríguez, S., & Kronauer, D. J. C. (2023). Sparse and stereotyped encoding

implicates a core glomerulus for ant alarm behavior. *Cell*. In press.

22. Hewlett, S.E., Wareham, D.M., & Barron, A.B. (2018). Honey bee (*Apis mellifera*) sociability

and nestmate affiliation are dependent on the social environment experienced post-eclosion. *J*

Exp Bio, 221(3).

23. Hostinar, C.E., Stellern, S.A., Schaefer, C., Carlson, S.M., & Gunnar, M.R. (2012).

Associations between early life adversity and executive function in children adopted

internationally from orphanages. *PNAS*, *109*, 17208–17212.

24. Ito, K., Shinomiya, K., Ito, M., Armstrong, J. D., Boyan, G., Hartenstein, V., Harzsch, S.,

Heisenberg, M., Homberg, U., Jenett, A., Keshishian, H., Restifo, L. L., Rössler, W., Simpson,

J. H., Strausfeld, N.J., Strauss, R., & Vosshall, L.B. (2014). A Systematic Nomenclature for

the Insect Brain. *Neuron*, *81*(4), 755–765. x

25. Jaisson, P., & Fresneau, D. (1978). The sensitivity and responsiveness of ants to their cocoons

in relation to age and methods of measurement. *Anim Behav*, *26*, 1064–1071.

26. Jernigan, C. M., Zaba, N. C., & Sheehan, M. J. (2021). Age and social experience induced

plasticity across brain regions of the paper wasp *Polistes fuscatus*. *Biol Lett*, *17*(4),

rsbl.2021.0073.

27. Kamhi, J.F. & Traniello, J.F.A. (2013). Biogenic amines and collective organization in a

superorganism: neuromodulatin of social behavior in ants. *Brain Behav Evol*, 82:220-236.

28. Kamhi, J. F., Gronenberg, W., Robson, S. K. A., & Traniello, J.F.A. (2016). Social complexity

influences brain investment and neural operation costs in ants. *Proc R Soc B*, *283*(1841),

20161949.

29. Kamhi, J.F., Arganda, S., Moreau, C.S., & Traniello J.F.A. (2017). Origins of aminergic

regulation of behavior in complex insect social systems. *Front Syst Neurosci*, 11:74.

30. Koto, A., Mersch, D., Hollis, B., & Keller, L. (2015). Social isolation causes mortality by

disrupting energy homeostasis in ants. *Behav Ecol and Sociobiol*, *69*(4), 583–591.

31. Kronauer, D., & Libbrecht, R. (2018). Back to the roots: the importance of using simple insects

societies to understand the molecular basis of complex social life. *Curr Opin Insect Sci,* 28:

33-39.

32. Lévy, F., Melo, A.I., Galef, B.G., Madden, M., & Fleming, A.S. (2003). Complete maternal

deprivation affects social, but not spatial, learning in adult rats. *Dev Psychobiol*, *43*(3), 177–

191.

33. Lihoreau, M., Brepson, L., & Rivault, C. (2009). The weight of the clan: Even in insects, social

isolation can induce a behavioural syndrome. *Behav Process*, *82*(1), 81–84.

34. Lösel, P. D., van de Kamp, T., Jayme, A., Ershov, A., Faragó, T., Pichler, O., Tan Jerome, N.,

Aadepu, N., Bremer, S., Chilingaryan, S.A., Heethoff, M., Kopmann, A., Odar, J., Schmelzle,

S., Zuber, M., Wittbrodt, J., Baumbach, T., & Heuveline, V. (2020). Introducing Biomedisa as

an open-source online platform for biomedical image segmentation. *Nat Commun, 11*(1), 5577.

35. Mackevicius, E.L., Gu, S., Denisenko, N.I., & Fee, M.S. (2023). Self-organization of songbird

neural sequences during social isolation. *eLife*, *12*.

36. Marder, E. (2012). Neuromodulation of Neuronal Circuits: Back to the Future. *Neuron*,

76(1), 1–11.

37. Mashoodh, R., Franks, B., Curley, J.P., & Champagne, F.A. (2012). Paternal social enrichment

effects on maternal behavior and offspring growth. *PNAS, 109*, 17232–17238.

38. McDonald, P., & Topoff, H. (1985). Social regulation of behavioral development in the ant,

Novomessor albisetosus (Mayr). *J Comp Physiol A*, *99*(1), 3-14.

39. Miroschnikow, A., Schlegel, P., & Pankratz, M. J. (2020). Making Feeding Decisions in the

Drosophila Nervous System. *Curr Biol*, *30*(14), R831–R840.

40. Morel, L. (1986). Ontogenesis of the antennal activity associated with food transfer in the

callow worker ant. *Dev Psychobiol*, *19*(5), 413–426.

41. Muratore I. B., Fandozzi E. M., & Traniello J.F.A. (2022). Behavioral performance and
division of labor influence brain mosaicism in the leafcutter ant *Atta cephalotes*. *J Comp*
Physiol A, 208: 325–44.

42. Muscedere, M. L., Willey, T. A., & Traniello, J. F. A. (2009). Age and task efficiency in the
ant Pheidole dentata: young minor workers are not specialist nurses. *Anim Behav*, *77*(4),
911–918.

43. Muscedere, M.L., Traniello, J.F.A., & Gronenberg, W. (2011). Coming of age in an ant colony:
cephalic muscle maturation accompanies behavioral development in *Pheidole dentata*.
Naturwissenschaften, *98*(9), 783–793.

44. Muscedere, M.L. & Traniello, J.F.A. (2012). Division of labor in the hyperdiverse ant genus
Pheidole is associated with distinct subcaste- and age-related patterns of worker brain
organization. *PLoS ONE,* 7.2:e31618.

45. Muscedere, M. L., Djermoun A., & Traniello, J. F. A. (2013). Brood-care experience, nursing
performance, and neural development in the ant *Pheidole dentata*. *Behav Ecol Sociobiol,* 67.5,
775-784.

46. Pardo-Sanchez, J., Kou, N., & Tibbetts, E. A. (2022). Type and amount of social experience
influences individual face learning in paper wasps. *Behav Ecol and Sociobiol*, *76*(11), 148.

47. Pisokas, I., Heinze, S., & Webb, B. (2020). The head direction circuit of two insect species.
eLife, *9*.

48. Posit team (2023). RStudio: Integrated Development Environment for R. Posit Software, PBC,
Boston, MA. URL http://www.posit.co/.

49. R Core Team (2023). R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

50. Richardson, T. O., Kay, T., Braunschweig, R., Journeau, O. A., Rüegg, M., McGregor, S., Los

Rios, P. de, & Keller, L. (2021). Ant behavioral maturation is mediated by a stochastic

transition between two fundamental states. *Curr Biol*, *31*(10), 2253-2260.e3.

51. Rodriguez-Romaguera, J., & Stuber, G.D. (2018). Social Isolation Co-opts Fear and

Aggression Circuits. *Cell*, *173*(5), 1071–1072.

52. Sakurai, A., Tamvacakis, A. N., & Katz, P. S. (2014). Hidden synaptic differences in a neural

circuit underlie differential behavioral susceptibility to a neural injury. *eLife*, *3*.

53. Sallet, J., Mars, R.B., Noonan, M.P., Andersson, J.L., O'Reilly, J.X., Jbabdi, S., Croxson, P.L.,

Jenkinson, M., Miller, K.L. and Rushworth, M.F. (2011). Social network size affects neural

circuits in macaques. *Science,* 334(6056): 697-700.

54. Scharf, I., Stoldt, M., Libbrecht, R., Höpfner, A.L., Jongepier, E., Kever, M., & Foitzik, S.

(2021). Social isolation causes downregulation of immune and stress response genes and

behavioural changes in a social insect. *Mol Ecol*, *30*(10), 2378–2389.

55. Seid, M., Harris K.M., & Traniello, J.F.A. (2005). Age-related changes in the number and

structure of synapses in the lip region of the mushroom bodies in the ant *Pheidole dentata*. *J

Comp Neurol*, 488.3: 269-277.

56. Seid, M., Goode, K., Li, C., & Traniello, J.F.A. (2008). Age-and subcaste-related patterns of

serotonergic immunoreactivity in the optic lobes of the ant *Pheidole dentata*. *Dev Neurobiol*,

68.11: 1325-1333.

57. Seid, M., & Junge, E. (2016). Social isolation and brain development in the ant *Camponotus

floridanus*. *Sci Nat-Heidelberg*, 103.5-6: 42.

58. Serra, M., Pisu, M. G., Littera, M., Papi, G., Sanna, E., Tuveri, F., & Biggio, G. (2000). Social

isolation-induced decreases in both the abundance of neuroactive steroids and GABAA

receptor function in rat brains. *J Neurochem,* 75(2): 732-740.

59. Sheehan, Z.B.V., Kamhi, J. F., Seid, M.A., & Narendra, A. (2019). Differential investment in

brain regions for a diurnal and nocturnal lifestyle in Australian *Myrmecia* ants. *J Comp Neurol*,

527(7), 1261–1277.

60. Sillar, K. T., Combes, D., & Simmers, J. (2014). Neuromodulation in developing motor

microcircuits. *Curr Opin in Neurobiol*, *29*, 73–81.

61. Simola, D.F., Chaoyang Y., Mutti, N.S., Dolezal, K., Bonasio, R., Liebig, J., Reinberg, D., &

Berger, S. (2013). A chromatin link to caste identity in the carpenter ant *Camponotus*

floridanus. Genome Res, 23:486-496.

62. Smith, A. R., Seid, M. A., Jiménez, L. C., & Wcislo, W. T. (2010). Socially induced brain

development in a facultatively eusocial sweat bee *Megalopta genalis* (Halictidae). *Proc R Soc*

B: Biol Sci, 277(1691), 2157–2163.

63. Taborsky, B.*,* Arnold, C., Junker, J., & Tschopp, A. (2012). The early social environment

affects social competence in a cooperative breeder. *Anim Behav,* 83, 1067-1074.

64. Thuma, J. B., Hobbs, K. H., Burstein, H. J., Seiter, N. S., & Hooper, S. L. (2013). Temperature

Sensitivity of the Pyloric Neuromuscular System and Its Modulation by Dopamine. *PLoS*

ONE, *8*(6), e67930.

65. Topoff, H. & Mirenda, J. (1978). Precocial behavior of callow workers of the army any

Neivamyrmex nigrescens: Importance of stimulation by adults during mass recruitment. *Anim*

Behav, 26(3): 698-700.

66. Traniello, J.F.A. (1978). Caste in a primitive ant: absence of age polyethism in *Amblyopone*.

Science, 202: 770-772.

67. Turner-Evans, D. & Jayaraman, V. (2016). The insect central complex. *Curr Biol*, 26:R445-

R460.

68. Vora, A., Nguyen, A. D., Spicer, C., & Li, W. (2022). The impact of social isolation on health

and behavior in *Drosophila melanogaster* and beyond. *Brain Sci Adv 8*(3), 183–196.

69. Wada-Katsumata, A., Yamaoka, R. and Aonuma, H. (2011). Social interactions influence

dopamine and octopamine homeostasis in the brain of the ant *Formica japonica*. *J Exp Biol,*

214(10): 1707-1713.

70. Wang, Z. Y., McKenzie-Smith, G. C., Liu, W., Cho, H. J., Pereira, T., Dhanerawala, Z.,

Shaevitz, J. W., & Kocher, S. D. (2022). Isolation disrupts social interactions and destabilizes

brain development in bumblebees. *Curr Biol*, *32*(12), 2754-2764.e5.

71. Waxman, H.K., Muscedere, M.L., & Traniello, J.F.A. (2017). Behavioral performance and

neural systems are robust to sensory injury in workers of the ant *Pheidole dentata*. *Brain Behav*

Evol, 89: 195-208.

72. Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New

York. ISBN 978-3-319-24277-4.

73. Wnuk., A., Wiater, M., & Godzińska, E. J. (2011). Effect of past and present behavioural

specialization on brain levels of biogenic amines in workers of the red wood ant Formica

polyctena. *Physiol Entomol*, *36*(1), 54–61.

74. Zhang, H., Liu, Y., Zhou, K., Wei, W., & Liu, Y. (2021). Restoring Sensorimotor Function

Through Neuromodulation After Spinal Cord Injury: Progress and Remaining Challenges.

Front Neurosci, 15.