Differential Neuroanatomical, Neurochemical, and Behavioral Impacts of

Early-Age Isolation in a Eusocial Insect

Billie C. Goolsby[1,2]*, E. Jordan Smith[1], Isabella B. Muratore[1,3], Zach N. Coto[1], Mario L.

Muscedere[1], James F. A. Traniello[1]

[1] Department of Biology, Boston University, Boston, MA, 02215, USA

[2] Department of Biology, Stanford University, Stanford, CA, 94305, USA

[3] Department of Biological Sciences, New Jersey Institute of Technology, NJ, 07102, USA

*Corresponding authors:

Billie C. Goolsby
Department of Biology
Stanford University
371 Jane Stanford Way
Stanford, CA
bgoolsby@stanford.edu
24
25   James F. A. Traniello
26   Department of Biology
27   Boston University
28   Boston, MA 02215
29   jft@bu.edu

Abstract

Social experience early in life appears to be necessary for the development of species-typical

behavior. Although isolation during critical periods of maturation has been shown to impact

behavior by altering gene expression and brain development in invertebrates and vertebrates,

workers of some ant species appear resilient to social deprivation and other neurobiological

challenges that occur during senescence or due to loss of sensory input. It is unclear if and to

what degree neuroanatomy, neurochemistry, and behavior will show deficiencies if social

experience in the early adult life of worker ants is compromised. We reared newly-eclosed adult

workers of *Camponotus floridanus* under conditions of social isolation for 2 to 53 days,

quantified brain compartment volumes, recorded biogenic amine levels in individual brains, and

evaluated movement and behavioral performance to compare the neuroanatomy, neurochemistry,

brood-care behavior, and foraging (predatory behavior) of isolated workers with that of workers

experiencing natural social contact after adult eclosion. We found that the volume of the

antennal lobe, which processes olfactory inputs, was significantly reduced in workers isolated for

an average of 40 days, whereas the size of the mushroom bodies, centers of higher-order sensory

processing, increased after eclosion and was not significantly different from controls. Titers of

the neuromodulators serotonin, dopamine, and octopamine remained stable and were not

significantly different in isolation treatments and controls. Brood care, predation, and overall

movement were reduced in workers lacking social contact early in life. These results suggest

that the behavioral development of isolated workers of *C. floridanus* is specifically impacted by

a reduction in the size of the antennal lobe. Task performance and locomotor ability therefore

appear to be sensitive to a loss of social contact through a reduction of olfactory processing

ability rather than change in the size of the mushroom bodies, which serve important functions in

learning and memory, or the central complex, which controls movement.

Introduction

Social experience shapes the nervous system and the behavior it generates. Exposure to social

stimuli and cues is critically important to neural development, behavioral performance and

cognition [1-5], neuroimmune function [6-10], and most significantly, survival [11, 12]. Across

diverse clades, behavioral capabilities can be negatively impacted if individuals are deprived of

social contact when ontogenic trajectories are most sensitive to receiving sensory inputs [13, 14].

A lack of social contact early in life can disrupt maturational processes required for feeding,

aggression, courtship, and interaction networks [12, 15].

Although research on the effects of social isolation has historically focused on humans, non-

human primates, and other mammals, eusocial insects have emerged as important systems to

examine the influences of social deprivation in early adult life on brain development, brain gene

expression, neuromodulators, and behavior [16-23]. Workers and immatures are

developmentally dependent [24]. Social interactions such as trophallaxis (oral food exchange),

allogrooming, and nursing influence social network structure and behavioral and

neuroanatomical development trajectories [23, 25-27]. Age-related changes in functionally

specialized brain compartments can be experience-expectant (growth programmed to occur

despite experiential input) or experience-dependent growth resulting from the performance of a

particular activity, such as foraging [28, 29]. Social isolation in ants appears to significantly alter

the growth of the mushroom body, a brain compartment involved in higher-order sensory

processing, learning, and memory, relative to that of nestmates developing in a natural social

environment [17]. If workers are deprived of social contact, neuromodulators and receptor

profiles that regulate behavior may be altered or become dysregulated [16, 30], potentially

affecting age-related responsiveness to signals and cues that guide neural and behavioral

development, and task performance inside and outside of the nest [20, 31-37]. Nevertheless,

worker brains and behavior may be resilient to senescence [38] and injury to sensory appendages

[39]. This robustness to aging and loss of olfactory input from the antennae raises questions

concerning the ability to compensate for stressors resulting from social isolation during early

adulthood and whether worker neurobiology and behavior are equally sensitive to such

disruption.

In ants, newly eclosed adult workers (callows) typically develop in a socially rich colony

environment of demographically variable nestmates and immatures. Callows are distinguished

from their mature siblings by their unsclerotized lightly pigmented cuticles and relatively low

activity [40, 41]. Behavioral repertoire formation is usually incomplete immediately following

adult eclosion, with apparently few exceptions [42]. Manipulations of the social experience of

callows therefore provides the ability to experimentally examine the ontogeny of behavior from

the earliest stage of adult life. Social interaction between callows and their siblings influences

pheromone responsiveness and temporal behavioral development [40, 43-46]. The effects of

being deprived of these and other inputs are not consistently negative: 10-day old *Pheidole*

dentata minor workers that lacked contact with brood nursed as efficiently as same-aged control

workers and did not differ in brain anatomy or neurochemistry [20]. The question remains as to

whether neuroanatomy, neuromodulators, behavioral responsiveness to social signals (such as

those regulating brood care), and stimuli involved in sensing the extranidal environment (such as

those involved in foraging) develop independently of worker interactions with colony members

early in life and are differentially influenced by social deprivation.

The ant genus *Camponotus* has served as a model to examine neuroanatomical plasticity during

development [28] and neuroethological and neurochemical impacts of social deprivation. In *C.*

floridanus and *C. fellah*, decreases in worker mushroom body size scaling, changes in levels of

biogenic amines, and deficits in social behavior have been documented in isolated individuals

[16, 17, 30, 47]. Here, we integrate analyses of brain size and the growth of neuropils

specialized for sensorimotor functions, titers of brain biogenic amine neuromodulators, and task

performance competence to examine how a lack of social contact during early life affects worker

neurobiology and behavior.

Methods

Colony collection and culture

Three queenright colonies of *C. floridanus* were collected in or near Gainesville, Florida in 2012

and reared in Harris environmental chambers at 25°C and 40-55% relative humidity on a 12:12

light cycle. Parent colonies from which workers were sampled had a minimum size of 100,

equally distributed between minors and majors. Artificial nests were made from test tubes 75%

filled with water and fitted with a tight cotton plug and placed in Fluon®-lined (Bioquip, Rancho

Dominguez, CA) plastic nest boxes 16x11x6 cm to 32x22x6 cm depending on colony size.

Colonies were fed 1M sucrose on saturated cotton balls and varied protein sources (mealworms,

scrambled eggs, and fruit flies) every other day in rotation.

Worker social isolation

C. floridanus has a dimorphic worker caste. We used minor workers because of their diverse

behavioral repertoire, which includes brood care and foraging, rather than majors, which are

limited to defense. Four groups of callow minors were collected from each of three parent

colonies. Randomly selected callows were isolated from nestmates for 2-3 days (n=29), 5-6 days

(n=36), 20-22 days (n=44), and 40 days (n=29, mean = 40.08 days, range = 30 – 53 days; herein

40 days) in separate Fluon®-lined boxes (16 x 11 x 6 cm). Socially isolated workers ("isolates")

were provided similar conditions in smaller nest boxes and the same diet as parent colonies.

After the prescribed isolation period, isolates were assayed for behavioral performance and

subsequently sacrificed for immunohistochemical analysis of brain structure, or for

quantification of brain biogenic amine titers. Callow (n = 29) and mature minor workers

("matures", n = 46) were collected and assayed using the same methods. Callow and mature

controls served as comparative groups for 2-day isolates and 40-day isolates, respectively, and

we report primarily on differences in behavior, neuroanatomy, and neurochemistry between

these groups. We compared brain volumes of callow controls estimated to be 2 days old and 2--

3-day isolates to ensure there were no intrinsic differences in brain size post eclosion. We used

5- and 20-day isolates lacking age-matched controls to estimate the trajectory of behavioral and

neural development from 2 days to 53 days of isolation (**Supp Fig 1-4**). Isolated workers had a

survival rate of 63.7%. Because of the relatively high mortality of workers experiencing longer

periods of isolation, sample sizes for 40-day isolates are generally lower. Head width, measured

as the widest distance across the eyes, served as a proxy for body size. To control for the effect

of body size on brain size, we compared the head widths of workers across isolation periods and

control groups and found no significant differences (p=0.40, R^2=0.098, F=1.18, Z=0.25, SS type

3, **Supp Fig 1a**). Our sampling of workers for assignment to isolation or control treatments was

thus unlikely to influence brain volume.

Behavioral performance assays

Isolated and control workers used for behavior (n = 16 [2 day], 19 [5 day], 20 [20 day], 11 [40

day], 8 [callow] and 13 [mature]) were exposed to a randomized series of assays to determine

foraging ability and ability to recognize and care for brood, tasks that affect colony fitness.

Simplified illustrations of the assays can be found in **Fig 1a.**

Predatory Behavior

A single worker was carefully placed in a Fluon®-lined Petri dish (9cm) and allowed to

acclimate for 2 minutes prior to assay initiation. Using Dumont No. 5 fine forceps sanitized with

ethanol and dried, a live fruit fly was offered to the worker and the response was observed for 2

minutes. The degree of predatory behavior was scored on a 5-point scale; higher scores indicate

greater predatory behavior. 1. no response or avoidance; 2. olfactory (antennal) investigation; 3.

mandibular flaring; 4. latent attack (delayed and intermittent); and 5. immediate attack (persistent

predatory attack from first encounter with prey). The assay arena was also sanitized with ethanol

between trials to minimize influence from social signals or prey odors.

Brood recognition and care

To determine if isolates differ in the ability to recognize and care for immature siblings with

similar efficacy as control matures and callows, we evaluated their social interactions with

pupae. Each focal worker was carefully placed in a 14 cm diameter Petri dish test arena where a

30 x 10 mm red plastic hollow tube provided a darkened area into which workers could move

pupae. Three pupae were added at three different locations equidistant from the darkened

chamber before a worker was introduced. Workers could relocate pupae either inside or outside

of the dark chamber or have no interaction. Pupae were recorded as aggregated ("piled") if

workers moved two or more together at any location. Workers showing this level of response

began pupal care immediately. We did not assign lower scores to workers that clustered pupae

without moving them into the darkened area. The darkened tube provided the opportunity for a

level of pupal care that would emulate natural placement of immatures within a colony.

However, no worker moved pupae into the tube and we therefore did not include this behavior in

the calculation of brood-care scores. Some workers did not move any pupae, but they

themselves moved into the darkened tube. This was scored as avoidance.

Worker interactions with pupae were observed directly for 5 minutes; after an additional 15

minutes , the final placement of pupae was recorded (total trial length = 20 minutes). Brood

recognition and care was measured on a 7-point scale: 1. no interaction or avoidance throughout

the duration of the trial; 2. olfactory (antennal) investigation; 3. mouthpart contact; 4. one pupa

moved during the last 15-minute period; 5. one pupa moved in the first five-minute period; 6.

two pupae moved beginning in the first five-minute period; 7. three pupae moved beginning in

the first five-minute period. Levels 4 and 5 distinguish the rate of response to pupae, which we

interpret as sensory processing ability that could be sensitive to social isolation. Higher scores

indicate greater brood-recognition ability (likely olfactory), restoration of brood clusters through

pupal transport, and sustained engagement with brood. Workers received scores of 5-7

independent of where pupae were transported.

Locomotion

To examine if isolation affected physiological and/or biomechanical capability to move pupae or

attack prey, we used a simple and robust proxy to broadly assess neuromuscular impairment

[38]. Focal workers were placed in the center of a 10 cm diameter Fluon®-lined Petri dish

divided into four equal quadrants by a crosshair pattern drawn beneath. After a two-minute

acclimation period, the number of times a worker moved among the four quadrants was recorded

during a five-minute period using a digital hand counter.   Movement between quadrants was

only scored when the worker passed a division line by at least one body length.

Neuroanatomical and neurochemical measurements

Immunohistochemistry and confocal microscopy

Intact brains (n = 8 [2 day], n = 11 [5 day], n = 15 [20 day], n = 14 [40 day], n = 5 [matures], and

n = 7 [callows]) were dissected in ice-cold Ringer's solution, and then immediately placed in ice-

cold zinc-formaldehyde fixative with shaking overnight. Fixed brains were then washed in

HEPES buffered saline 6 times (10 minutes/wash) and fixed in Dent's Fixative (80% MeOH,

20% DMSO) for 1 hour. Brains were placed in 100% methanol and stored at -17 C° for 1-3

weeks until processed. Brains were next rinsed 6 times (10 minutes/rinse) with 3% Triton X-100

in 1X PBS (3% PBST). Brains were stored overnight in 5% normal goat serum/0.2%

PBST/0.005% sodium azide for blocking. After blocking, brains were incubated for 4 days RT

with an anti-synapsin antibody to stain neuropils (1:30, Development Studies Hybridoma Bank,

Iowa City, Iowa) in constant nutation. Brains were washed with 0.2% PBST (6x10 minutes),

then incubated in secondary antibody (1:100 AlexaFluor goat □-mouse) in 5% normal goat

serum/0.2%PBST/0.005% sodium azide for 4 days at RT with constant nutation. Brains were

washed in 0.2% PBST (6x10 minutes), then dehydrated through an ethanol and PBS series (30,

50, 70, 95, 100% ethanol in 1XPBS). Brains were cleared and immersed in methyl salicylate and

mounted on stainless steel glass-windowed slides for confocal imaging.

Brains were imaged with a Nikon C2 + Si spectral laser scanning confocal microscope (Nikon,

Melville, NY, USA) with sections at approximately 1.5 um throughout the entire brain. Images

were manually annotated using Amira 6.0 software to quantify neuropil volumes. The individual

who annotated brains was blind to brain sample category, thus minimizing human bias. The

margins of neuropils were identified visually or with magic wand tools in Amira. Every eighth

frame was annotated manually. Biomedisa interpolation [48] was used on frames without labels,

then evaluated and edited by the annotator for accuracy. We labeled volumes of the optic lobe

(OL; vision), antennal lobe (AL; olfaction), mushroom body (MB; higher-order processing,

learning, and memory), lateral calyx (MB-LC), and medial calyx (MB-MC), peduncles (MB-P),

central complex (CX; movement and navigation), suboesophageal zone (SEZ; mouthpart

control), and rest of the central brain (ROCB). ROCB describes tissue composed of the superior

neuropils, ventromedial neuropils, ventrolateral neuropils, and the lateral horn [49]. For each

brain compartment, volumes of only one intact hemisphere were recorded. If both hemispheres

were intact, the labeler chose one randomly.

A detailed description of brain anatomy and development in relation to age and experience in *C.*

floridanus is presented in Gronenberg et al. [28]. We compared the relative volumes of brains

across all groups and scaling relationships among brain compartments. Relative volumes were

calculated by dividing the volume of each focal compartment by the ROCB volume. Damaged

compartments were excluded from analysis. If one or more compartments or the ROCB was

damaged, the sample was excluded. The MB-LC and MB-MC exhibited similar volumetric

trends and thus were combined to one metric (MB-C).

Quantification of brain biogenic amines

Isolated workers were quickly dissected (n = 11 [2 days], n = 10 [5 days], n = 12 [20 days], n = 3

[40 days], along with control mature workers (n = 11) to quantify levels of serotonin (5HT),

dopamine (DA), and octopamine (OA) using isocratic, reversed-phase high-performance liquid

chromatography with electrochemical detection (HPLC-ED). Monoamine quantification was

performed blind to treatment. The HPLC-ED system (formerly ESA, Inc., Chelmsford, MA,

USA) included a model 584 pump, a model MD-150 (3 x 150 mm) reversed-phase analytical

column, a model 5011A dual-channel coulometric analytical cell, and a Coulochem III

electrochemical detector. Brains were dissected in less than 5 minutes in ice-cold Ringer's

solution, homogenized in mobile phase manually with a plastic polypropylene pestle in a

centrifuge tube, and centrifuged. The supernatant was then injected into the HPLC column, and

monoamine levels were quantified with reference to external standards. Mobile phase was

formulated as 50 mM citrate/acetate buffer, 1.5 mM sodium dodecyl sulfate, 0.01%

triethylamine, and 25% acetonitrile in MilliQ water [20]. Multiple daily runs of the external

standards accounted for any changes in ambient conditions.

Statistical analysis and data visualization

All statistics analyses were performed in RStudio version 4.2.3 [50]. For generalized linear

mixed models, we used the package '*RRPP*' [51]. We tested for statistical significance using

ANOVA with residual randomization in a permutation procedure of 1000 iterations and the

estimation method of ordinary least squares, given combined sample size of 25 or more. We

included colony and head width as random effects in our behavior, brain volume, and biogenic

amine modeling; neither had a significant effect. Isolation significantly correlated with head

width only when head width was nested with the microscope and experimenter, rather than head

width alone, meaning that slight variation between ocular micrometers of the microscopes, rather

than treatment, influenced the effect of head width. ANOVA and modeling details can be found

in **Supp Tables Set 1 and 2.** Following ANOVA, *post-hoc* tests were done on pairwise

comparisons using Wilcoxon rank sum exact tests with a Benjamini-Hochberg (BH) procedure to

correct for multiple tests. Depending on the sample size of 40-day isolates, we included type III

error to account for sample size bias. Additional nonparametric statistical analyses including

Kruskal-Wallis H tests and *post-hoc* pairwise comparisons using t tests with a Bonferroni

correction, which agreed with our modeling, can be found in **Supp Tables Set 1 and 2.** We then

ran studentized Breusch-Pagan (BP) tests from the '*lmtest*' package [52] to confirm

homoscedasticity in our model. If heteroscedasticity was found in raw values, we log

transformed our data and reran the model and confirmed homoscedasticity. Heteroscedasticity

could not be removed from only our mixed model analysis of MB-P scaled volumes in

Supplementary Tables, Set 1. However, the model nonetheless agrees with our additional

nonparametric statistical tests. We confirmed that log-transformed data resulted in the same

pairwise comparison results as raw data, confirming that log transformation for model fitness did

not create false pairwise findings. Because 5-day isolates and 20-day isolates did not have

socially typical, age-matched controls and were instead meant to provide insight into the

trajectory of neurodevelopment and behavior, we excluded these groups from the analysis

reported in the manuscript. Our analyses including these excluded groups can be found in **Supp**

Figs 1-4 and in **Supp Tables, Set 2.** Figures were created using the package '*ggplot2*' [53].

Figure assembly was done in Adobe Illustrator (Adobe Illustrator 2023). Illustrations of ants in

Visual Abstract, Fig 1a, and **Supp Fig 1b** are courtesy of BioRender.com.

Results

Effect of Isolation on Predatory Behavior

Mature control workers preyed on fruit flies significantly more ($\bar{x}$ = 3.39) than workers in all

other experimental groups and callows (vs. callows, 2-day isolates, and 40-day isolates: $p <$

0.029; BH correction). Forty-day isolates exhibited significantly higher predatory behavior ($\bar{x}$

= 2.27) than callows ($\bar{x}$ =1.25; p=0.0093, BH correction). All other pairwise comparisons were

not significantly different (p=0.16-0.29). **(Fig 1a, b)**. Therefore, control callows and 2-day

isolates similarly avoided prey or showed olfactory investigation whereas older isolates (40-day,

e.g.) had higher predatory scores that resembled those of mature worker controls but nevertheless

were significantly lower.

Effect of Isolation on Brood Recognition and Care

Brood-care performance was not significantly different among treatments (p=0.071, R^2=0.17,

F=2.52, Z=1.43, SS type 1, **Fig 1a, c**), but in *post-hoc* tests, mature workers performed brood

care with significantly higher scores than 2-day isolates ($\bar{x}$ = 4.15 and 2.06, respectively;

p=0.0093, BH correction). Mature workers scored higher than 40-day isolates ($\bar{x}$ = 1.91;

p=0.0093, BH correction). Callows also had a significantly higher brood-care score (x□ score =

3.38) than both 2-day isolates and 40-day isolates (p=0.036 for each pairwise comparison, BH

correction). Socially typical callow and mature worker controls were not significantly different,

nor were 2-day isolates and 40-day isolates (p = 0.36 and 0.98, respectively).

Effect of Isolation on Locomotion

Mature controls had significantly higher movement (x□ = 77.84 times a worker moved among

quadrants) than workers than 2-day isolates, 40-day isolates, and callows (x□ = 26.38, 38.36,

and 18.38 respectively; p=0.004, R^2=0.35, F=5.97, Z= 2.74, SS type 1, [**Fig 1d, Supp Table Set**

1]); 40-day isolates had the next highest average score (x□ = 38.36). In pairwise comparisons,

mature controls were significantly higher than from all other experimental groups (callow, 2, 40:

all p<0.0057, BH correction). No other pairwise comparisons were statistically significant

(p>0.19).

Brain Compartment Allometries

Workers experiencing longer periods of isolation had significantly smaller AL volumes

compared to those of mature control workers (p=0.007, R^2=0.39, F=6.19, Z=2.56; **Fig 2a**).

Mature workers thus had significantly larger relative AL volumes (on average, 27% relative

volume) than 40-day isolates (17% relative volume; p=0.017, BH correction). Forty-day isolates

also had significantly lower AL volume than 2-day isolates (23% relative volume, p=0.048, BH

correction), suggesting that isolation reduced post-eclosion AL volume. Forty-day isolates had

significantly higher relative MB-C volumes (61% relative volume) than control callows (48%)

and 2-day isolates (49%; p<0.002, 0.005, respectively; BH correction); MB-C volumes were not

significantly different between callows and 2-day isolates (p=0.23; BH correction) or between

mature workers (76%) and 40-day isolates (p=0.13, BH correction). The average relative

volume of the MB-P showed significant increase between early isolates to older isolates

(p=0.029, R^2=0.21, F=3.74, Z=1.9; **Fig 2b**); however, there was no significant difference in

pairwise interactions, indicating that impact on the MB was limited to the calyces, which showed

modest growth during isolation (p=0.002, R^2=0.47, F=9.40, Z=3.52; **Fig 2c**). Other brain

compartments (OL, SEZ, CX) did not significantly change with isolation (p=0.25-0.93; **Fig 2d-**

f). All additional pairwise analyses of compartmental volumes can be found in **Supp Tables Set**

1.

Biogenic amine levels

Brain titers of OA, DA, and 5HT significantly increased with longer periods of isolation (**Fig 3a-**

c); levels of 40-day isolates were not significantly different from those of mature workers. DA

titers significantly increased with prolonged periods of isolation (p=0.001, R^2=0.47, F=11.86,

Z=3.38, SS type 3, **Fig 3a**) and significantly differed between mature control workers and 2-day

isolates, but not between mature controls and 40-day isolates (p < 0.0001, p > 0.05, respectively;

BH correction). OA exhibited a similar positive pattern with prolonged periods of isolation

(p=0.001, R^2=0.59, F=32.97, Z=4.96, SS type 3, **Fig 3b**); titers significantly increased between

isolated worker groups and with mature workers (p<0.006 for all groups, BH correction). 5HT

titers also significantly increased with longer periods of isolation (p=0.001, R^2=0.61, F=91.77,

Z=7.04, SS type 3, **Fig 3c**), with all pairwise comparisons being significantly different (p<0.006)

for all groups, BH correction).

While absolute levels of DA, OA, and 5HT increased from callows to 40-day isolates and

matures, their ratios in isolates increased or decreased compared to those recorded in mature

controls (**Fig 3d-f**). OA:5HT significantly decreased with time and extended periods of isolation

(p=0.004, R^2=0.21, F=6.37, Z = 3) with significant difference between 2-day isolates and mature

controls (p < 0.001, BH correction; **Fig 3d**). Forty-day isolates were not significantly different

from any treatment category. Neither OA:DA (p=0.96, R^2=0.0041, F=0.048, Z= -1.74) nor

5HT:DA (p=0.40, R^2=0.061, F=1.00 Z=0.30) significantly changed with isolation. However, 2-

day isolates had significantly lower 5HT:DA ratios than mature workers (p=0.02, respectively;

BH correction).

DA titers relative to total monoamine levels did not significantly differ across isolation treatment

groups and mature workers (p=0.47, R^2=0.050, F=0.77, Z=0.057). Relative OA levels, in

contrast, were significantly higher in younger isolates compared to mature workers (p=0.034,

R^2=0.26, F=4.14, Z=1.79); 2-day and mature controls were significantly different (p=0.002,

respectively, BH correction). Relative 5HT quantities did not change in isolation treatments and

in comparison, to mature controls (p=0.40, R^2=0.061, F=1.00 Z=0.30). However, 2-day isolates

had significantly lower 5HT:DA ratios compared to 20-day isolates and mature controls (p=0.02,

respectively; BH correction). Relative 5HT quantities were significantly different (p=0.094,

R^2=0.13, F=3.65, Z=1.32) between 2-day vs. mature controls (p<0.008 respectively, BH

correction). All additional pairwise analyses for biogenic amine levels can be found in **Supp**

Tables, Set 1.

Discussion

Brood-care and predatory behavior scores of *C. floridanus* workers isolated for varying durations

after eclosion were lower than those of control workers that developed in a natural social

environment. Two-day isolates showed a similar low level of predatory behavior as that of

callow controls, but the predation scores of 40-day isolates were significantly lower than those of

mature control workers, suggesting impairment due to isolation. Social isolation also resulted in

a significant difference in pupal care – as measured in our assay – between 2-day isolates and

callow control workers, and 40-day isolates and mature control workers. This result may reflect

the development of behavioral competencies associated with myological maturation [54] that

may be compromised by a lack of sensory inputs. Workers in all isolation treatment groups

showed significantly decreased locomotion relative to mature workers (**Fig 1d, Supp Fig 1c**),

consistent with Wang et al. [23] for isolated bumblebees but different from *Cataglyphis niger*

and *Camponotus fellah* ant workers, which increased movement following isolation and then

decreased movement [22, 55]. Older isolates (40-day isolates) generally scored higher than

workers isolated for a shorter period (2-day isolates), but were still significantly lower in their

activity, predatory, and brood-care scores than mature controls. This suggests that some

behaviors develop in an experience-expectant manner and require social contact for complete

development. *C. floridanus* may thus exhibit a combination of experience-expectant and

experience-dependent components. The latter has been found in honeybees, which show an

increase in mushroom body size with foraging experience [29].

C. floridanus worker brains normally increase in size during a 10-month period after adult

eclosion, the greatest growth rate (nearly 100%) occurring in the MBs and ALs [28]. Our

maximum early life isolation treatments were approximately 12-17% of this period of

maturation. Workers isolated for longer durations and socially typical mature workers generally

had larger relative MB-C volumes compared to workers isolated for shorter periods and socially

typical callows. MB-C size in 40-day isolates was not significantly different from that of mature

controls, suggesting gross neuroanatomical development remained stable under conditions of

isolation. These findings differ from the results of Seid and Junge [17], which showed that MBs

of isolated *C. floridanus* workers do not change in volume relative to total brain size. MB-C

resilience through 71 days of isolation has been found in paper wasps [56]. Using bumblebees as

a model, Wang et al. [23] associated expression differences in juvenile hormone-related genes

and transcriptomic network dysregulation with isolation-induced variation in brain volume.

Bumblebees isolated after eclosion for 9 days showed significant differences in the variance of

MB volume rather than average volume compared to socially typical bees, suggesting growth

destabilization in some, but not all, workers. On the other hand, our results agree with research

showing that the mushroom bodies of isolated honeybees continue to grow [57]. Our study

identified differential sensitivity and resilience in brain compartment growth. Although we did

not assess temporal patterns to determine if the MBs or other compartments accelerated or

decreased differentially in size over time in isolates and/or socially typical workers, results

nevertheless suggest that MB development is resilient to social isolation.

In contrast, AL volume decreased with periods of isolation approximately 5 days or longer; 40-

day isolates had, on average, smaller ALs than control callows and mature workers. This decline

in AL size is similar to the findings of Wang et al. [23] on bumblebees. Because AL size

decreased with isolation duration in *C. floridanus,* loss of olfactory ability could decrease brood-

care and predatory behavior by 40-day isolates compared to mature controls. Social contact

early in life thus appears to provide sensory inputs necessary to stimulate a level of AL

development required to perform the tasks we assayed. Isolation did not impact OL volume in

our study of *C. floridanus*, although it did reduce OL volume in paper wasps, which integrate

visual information used in conspecific facial recognition [56, 58]. The CX, which functions in

the control of movement [59] and sensorimotor processing in insects [60, 61], did not change

significantly in volume in isolated *C. floridanus* workers, a result inconsistent with Wang et al.

[23] for isolated bumblebees. The volume of the SEZ, which regulates mouthparts [62] and is

likely involved in the control of brood-care motor functions, did not significantly change with

isolation. This supports that the decline in brood-care behavior may have been due to a deficit in

olfactory processing rather than mouthpart motor deficits.

Although it may be generally assumed that smaller neuropil size reduces competence, we do not

know how whole-brain or compartmental volumes affect behavioral ability. Indeed, the

relationship between brain size and behavioral capacity is actively debated [72]. In ants,

elaboration of the MB appears to represent increased demands for higher-order processing [26,

73, 74] and larger OLs are correlated with more complex visual ecologies [75,76]. Larger ALs

have been associated with age-related olfactory ability [64]. However, Hart et al. [77] showed

that ant alarm pheromone response is encoded by less than 10 of a total of approximately 500 AL

glomeruli, a result suggesting that the relationship of neuropil size and behavior is unclear. We

also do not understand how circuit configuration, activation, synaptic strength, or other neural

processes in the ant brain connectome are influenced by worker isolation.

Worker behavioral maturation, including repertoire expansion and a corresponding increase in

brain size, correlates with increasing monoamine levels and the development of serotonergic

circuitry in some ants [32, 63-65]. Our *C. floridanus* data indicate that levels of OA, DA, and

5HT increase with age in the brains of isolated workers, despite a lack of social contact. Indeed,

40-day isolates had monoamine levels like those of mature workers. Ratios of biogenic amines

can significantly change with age in ants and may influence the development of age-related

behavior [32] by modifying response thresholds. However, we found no significant differences

in neuromodulator ratios between 40-day isolates and mature controls, indicating that aminergic

circuitry may be maintained under conditions of social deprivation. Boulay et al. [30] and

Wada-Katsumata [47] found that 5HT and DA levels were robust to social isolation and Boulay

et al. [30] hypothesized OA may reduce the effects of social deprivation, acting homeostatically

to maintain social cohesion. Whether octopaminergic signaling has a stimulatory or inhibitory

effect on prosocial behaviors such as trophallaxis, which may be affected by periods of social

isolation, requires further study.

Neuromodulators can restore sensorimotor functions following injury [72], alter synaptic

functions to significantly change circuitry [73, 74], and increase the robustness of network

functions [75] to recover from disruption and/or stress [76, 77]. In the ant *Pheidole dentata*,

monoamine levels continue to increase in titer in aging workers, which do not appear to decline

in task performance capability [38]. Given that DA, OA, and 5HT can flexibly reshape circuits,

neuromodulators may reduce isolation-induced neuroanatomical deficits in ants, stabilizing

circuit dynamics or returning them to a stable state if disrupted. Although we did not identify an

increase in monoamine levels in isolates, this does not preclude neuromodulatory compensation

to maintain circuitry and behavioral functions under conditions of social deprivation.

Conclusion

The results of prior research on the impacts of senescence on neurochemistry, neuroanatomy,

apoptosis, and behavior [38], as well as deprivation of experience [20] and reduced olfactory

processing ability [39], suggest that ant brains can have the capacity to buffer insults of aging

and physical damage to sensory appendages. The present study suggests brain compartments

vary in susceptibility and resilience to social deprivation and that aminergic modulatory systems

are robust to isolation. The 36% mortality of isolated *C. floridanus* workers we recorded

indicates a significant negative effect that may be due more to social processes affecting nutrition

and energy availability [22, 55] than neurobiological deficits resulting from isolation. Workers

that survive neurobiological and physiological "bottlenecks" appear able to recover from

challenges to some processes of brain development, which in *C. floridanus* are dependent on age

as well as experience [28]. Age-related increases in brain size in ants appear to be due to stable

programmed axonal and dendritic growth processes. If sensory experience is lacking or

temporally atypical, brain development may still proceed in part in the absence of inputs or may

be delayed (e.g., [78]). In our study, the effect of social isolation in *C. floridanus* appears to

reflect such robust programming, perhaps due in part to stabilizing effects of neuromodulators.

The resilience of worker ants to challenges during development may functionally resemble

differences that enable some individuals – perhaps those with adaptive synaptic phenotypes [79]

or ion channel properties [80] to withstand neural damage and preserve behavior. We

hypothesize that the ability to buffer stressors may involve such variation in circuitry that

characterizes clade-wide neuroarchitectures of the miniaturized brains of ants, acknowledging

that few species have been studied. Analyses of temporal patterns of brain gene expression in *C.*

floridanus isolated workers can identify transcriptomic changes linked to brain development and

influenced by social experience [81, 82] and possibly identify genetic mechanisms of resilience.

Genomic studies of phylogenetically and sociobiologically diverse ant species can also

contribute to our understanding of how workers are susceptible to and/or may recover from

social deprivation or are resilient to isolation.

Acknowledgements

499 We thank Lloyd Davis for colony collection. BCG thanks Dr. Katherine Fiocca for manuscript

500 feedback and Dr. Bryan Juarez for coding advice. We thank two anonymous reviewers for their

501 thoughtful comments.

Statement of Ethics

503 Ant colonies were cultured in an environment for optimal growth. Workers were cold

504 anaesthetized before dissection. Other issues of compliance with animal care regulations do not

505 apply. An ethics statement was not required for this study, as no humans nor vertebrates were

506 used in accordance with Boston University Institutional Animal Care and Use Committee.

Conflict of Interest Statement

508 The authors declare no conflict of interest.

Funding Sources

This research was supported by National Science Foundation grants IOS 1354291 and IOS

1953393 to JFAT, the Marion Kramer Award to BCG, and the Boston University Undergraduate

Research Opportunities Program to BCG during undergraduate study at Boston University.

BCG's graduate student funding during remote analysis and writing is supported by a Howard

Hughes Medical Institute Gilliam Fellowship (GT15685) and a National Institutes of Health

Cellular Molecular Biology Training Grant to Stanford University (T32GM007276).

Author Contributions Statement

BCG, IBM, and JFAT designed the study. BCG recorded behavior. BCG and IBM

immunohistologically prepared and imaged brains. EJS quantified brain volume. BCG, ZC, and

MM measured monoamine titers. BCG statistically analyzed volumetric data. BCG drafted the

manuscript. BCG, EJS, IBM, ZC, MM, and JFAT developed and edited the manuscript.

References

1. Amitai N, Young JW, Higa K, Sharp RF, Geyer MA, Powell SB. (2014). Isolation rearing
effects on probabilistic learning and cognitive flexibility in rats. *Cogn Affect Behav
Neurosci,* 14, 388-406.
2. Lévy F, Melo AI, Galef BG, Madden M, Fleming AS. (2003). Complete maternal deprivation
affects social, but not spatial, learning in adult rats. *Dev Psychobiol*, 43, 177-191.
3. Rosa-Salva O, Mayer U, Versace E, Hébert M, Lemaire BS, Vallortigara, G. (2021). Sensitive
periods for social development: Interactions between predisposed and learned
mechanisms. *Cognition*, 213, 104552.
4. Smith AR, Seid MA, Jiménez LC, Wcislo WT (2010). Socially induced brain development in
a facultatively eusocial sweat bee *Megalopta genalis* (Halictidae). *Proc R Soc B: Biol Sci,*
277, 2157-2163.
5. Taborsky B, Arnold C, Junker J, Tschopp A. (2012). The early social environment affects
social competence in a cooperative breeder. *Anim Behav,* 83, 1067-1074.
6. Donovan M, Mackey CS, Platt GN, Rounds J, Brown AN, Trickey DJ, et al. (2020). Social
isolation alters behavior, the gut-immune-brain axis, and neurochemical circuits in male
and female prairie voles. *Neurobiol Stress*, 13, 100278.
7. Sallet J, Mars RB, Noonan MP, Andersson JL, O'Reilly JX, Jbabdi S, et al. (2011). Social
network size affects neural circuits in macaques. *Science,* 334, 697-700.
8. Serra M, Pisu MG, Littera M, Papi G, Sanna E, Tuveri F, et al. (2000). Social
isolation☐induced decreases in both the abundance of neuroactive steroids and GABAA
receptor function in rat brains. *J Neurochem,* 75, 732-740.
9. Mashoodh R, Franks B, Curley JP, Champagne FA. (2012). Paternal social enrichment effects
on maternal behavior and offspring growth. *Proc Natl Acad Sci*, 109, 17232-17238.
10. Hostinar CE, Stellern SA, Schaefer C, Carlson SM, Gunnar MR. (2012). Associations
between early life adversity and executive function in children adopted internationally
from orphanages. *Proc Natl Acad Sci*, 109, 17208-17212.
11. Holt-Lunstad J, Smith TB, Baker M, Harris T, Stephenson D. (2015). Loneliness and social
isolation as risk factors for mortality. *Perspect Psychol Sci*, 10, 227-237.

12. Vora A, Nguyen AD, Spicer C, Li W. (2022). The impact of social isolation on health and
behavior in *Drosophila melanogaster* and beyond. *Brain Sci Adv*, 8, 183-196.
13. Boyce W, Sokolowski M, Robinson G. (2012). Toward a new biology of social adversity.
Proc Natl Acad Sci, 109, 17143-17148.
14. Cisneros-Franco JM, Voss P, Thomas ME, de Villers-Sidani E. (2020). Chapter 8 - Critical
periods of brain development. *Handb of Clin Neurol*, 173, 75-88.
15. Rodriguez-Romaguera J, Stuber GD. (2018). Social isolation co-opts fear and aggression
circuits. *Cell*, 173, 1071-1072.
16. Boulay R, Lenoir A. (2001). Social isolation of mature workers affects nestmate recognition
in the ant *Camponotus fellah*. *Behav Processes*, 55, 67-73.
17. Seid M, Junge E. (2016). Social isolation and brain development in the ant *Camponotus
floridanus*. *Sci Nat*, 103, 42.
18. Scharf I, Stoldt M, Libbrecht R, Höpfner AL, Jongepier E, Kever M, et al. (2021). Social
isolation causes downregulation of immune and stress response genes and behavioural
changes in a social insect. *Mol Ecol*, 30, 2378-2389.
19. Lihoreau M, Brepson L, Rivault C. (2009). The weight of the clan: Even in insects, social
isolation can induce a behavioural syndrome. *Behav Process*, 82, 81-84.
20. Muscedere ML, Djermoun A, Traniello JFA. (2013). Brood-care experience, nursing
performance, and neural development in the ant *Pheidole dentata. Behav Ecol Sociobiol,*
67, 775-784.
21. Hewlett SE, Wareham DM, Barron AB. (2018). Honeybee (*Apis mellifera*) sociability and
nestmate affiliation are dependent on the social environment experienced post-eclosion. *J
Exp Bio*, 221.
22. Koto A, Mersch D, Hollis B, Keller, L. (2015). Social isolation causes mortality by
disrupting energy homeostasis in ants. *Behav Ecol and Sociobiol*, 69, 583-591.
23. Wang ZY, McKenzie-Smith GC, Liu W, Cho HJ, Pereira T, Dhanerawala Z, et al. (2022).
Isolation disrupts social interactions and destabilizes brain development in bumblebees.
Curr Biol, 32, 2754-2764.
24. Snir O, Alwaseem H, Heissel S, Sharma A, Valdés-Rodríguez S, Carroll TS, et al. (2022).
The pupal moulting fluid has evolved social functions in ants. *Nature*, 612(7940), 488-
494.
25. Kronauer D, Libbrecht R. (2018). Back to the roots: the importance of using simple insects
societies to understand the molecular basis of complex social life. *Curr Opin Insect Sci,*
28, 33-39.
26. Godfrey R, Gronenberg W. (2019). Brain evolution in social insects: advocating for the
comparative approach. *J Comp Physiol A*, 205.1, 13-32.
27. Rittschof CC, Coombs CB, Frazier M, Grozinger CM, Robinson GE. (2015). Early-life
experience affects honeybee aggression and resilience to immune challenge. *Sci Rep*, 5,
15572.
28. Gronenberg W, Heeren S, Hölldobler B. (1996). Age-dependent and task-related
morphological changes in the brain and the mushroom bodies of the ant *Camponotus
floridanus. J Exp Bio*, 199, 2011-2019.
29. Withers GS, Fahrbach SE, Robinson GE. (1993). Selective neuroanatomical plasticity and
division of labour in the honeybee. *Nature,* 364, 238-240.
30. Boulay R, Soroker V, Godzinska EJ, Hefetz A, Lenoir, A. (2000). Octopamine reverses the
isolation-induced trophallaxis in the carpenter ant *Camponotus fellah*. *J Exp Biol,* 203,

513-520.

31. Seid M, Harris KM, Traniello JFA. (2005). Age☐related changes in the number and structure of synapses in the lip region of the mushroom bodies in the ant *Pheidole dentata*. *J Comp Neurol*, 488.3, 269-277.

32. Seid MA, Traniello JFA. (2005). Age-related changes in biogenic amines in individual brains of the ant *Pheidole dentata*. *Sci Nat,* 92, 198-201.

33. Kamhi JF, Traniello JFA. (2013). Biogenic amines and collective organization in a superorganism: neuromodulation of social behavior in ants. *Brain Behav Evol*, 82, 220-236.

34. Kamhi JF, Arganda S, Moreau CS, Traniello JFA. (2017). Origins of aminergic regulation of behavior in complex insect social systems. *Front Syst Neurosci*, 11, 74.

35. Friedman DA, Pilko A, Skowronska-Krawczyk D, Krasinska K, Parker JW, Hirsh J, et al. (2018). The role of dopamine in the collective regulation of foraging in harvester ants. *iScience*, 8, 283-294.

36. Friedman DA, York RA, Hilliard AT, Gordon DM. (2020). Gene expression variation in the brains of harvester ant foragers is associated with collective behavior. *Comm Biol*, 3, 100.

37. Maloney RT. (2021). Neuromodulation and individuality. *Front Behav Neurosci*, 15, 294.

38. Giraldo YM, Kamhi JF, Fourcassié V, Moreau M, Robson SK, Rusakov A, et al. (2016). Lifespan behavioural and neural resilience in a social insect. *Proc R Soc B*, 283, 20152603.

39. Waxman HK, Muscedere ML, Traniello JFA. (2017). Behavioral performance and neural systems are robust to sensory injury in workers of the ant *Pheidole dentata*. *Brain Behav Evol*, 89, 195-208.

40. Morel L. (1986). Ontogenesis of the antennal activity associated with food transfer in the callow worker ant. *Dev Psychobiol*, 19, 413-426.

41. Muscedere ML, Willey TA, Traniello JFA. (2009). Age and task efficiency in the ant *Pheidole dentata*: young minor workers are not specialist nurses. *Anim Behav*, 77, 911-918.

42. Traniello JFA. (1978). Caste in a primitive ant: absence of age polyethism in *Amblyopone*. *Science*, 202, 770-772.

43. Topoff H, Mirenda J. (1978). Precocial behavior of callow workers of the army ant *Neivamyrmex nigrescens:* Importance of stimulation by adults during mass recruitment. *Anim Behav*, 26: 698-700.

44. McDonald P, Topoff H. (1985). Social regulation of behavioral development in the ant, *Novomessor albisetosus (Mayr). J Comp Physiol A*, 99, 3-14.

45. Jaisson P, Fresneau D. (1978). The sensitivity and responsiveness of ants to their cocoons in relation to age and methods of measurement. *Anim Behav*, 26, 1064-1071.

46. Richardson TO, Kay T, Braunschweig R, Journeau OA, Rüegg M, McGregor S, et al. (2021). Ant behavioral maturation is mediated by a stochastic transition between two fundamental states. *Curr Biol*, 31, 2253-2260.

47. Wada-Katsumata A, Yamaoka R, Aonuma H. (2011). Social interactions influence dopamine and octopamine homeostasis in the brain of the ant *Formica japonica*. *J Exp Biol,* 214, 1707-1713.

48. Lösel PD, van de Kamp T, Jayme A, Ershov A, Faragó T, Pichler O, et al. (2020). Introducing Biomedisa as an open-source online platform for biomedical image segmentation. *Nat Commun,* 11, 5577.

49. Ito K, Shinomiya K, Ito M, Armstrong JD, Boyan G, Hartenstein V, et al. (2014). A
systematic nomenclature for the insect brain. *Neuron*, 81, 755-765.

50. Posit team (2023). RStudio: Integrated development environment for R. Posit Software,
PBC, Boston, MA. URL http://www.posit.co/.

51. Collyer ML, Adams DC. (2019). "RRPP: linear model evaluation with randomized residuals
in a permutation procedure. R package version 0.4.0." https://CRAN.R-
project.org/package=RRPP.

52. Zeileis A, Hothorn T. (2002). "Diagnostic checking in regression relationships." *R News*,
2(3), 7-10. https://CRAN.R-project.org/doc/Rnews/.

53. Wickham H. (2016). *ggplot2: elegant graphics for data analysis*. Springer-Verlag New
York.

54. Muscedere ML, Traniello JFA, Gronenberg W. (2011). Coming of age in an ant colony:
cephalic muscle maturation accompanies behavioral development in *Pheidole dentata*.
Sci Nat, 98, 783-793.

55. Ahronberg A, Scharf I. (2021). Social isolation interaction with the feeding regime
differentially affects survival and results in a hump-shaped pattern in movement activity.
Behav Processes, 190, 104460.

56. Jernigan CM, Zaba NC, Sheehan MJ. (2021). Age and social experience induced plasticity
across brain regions of the paper wasp *Polistes fuscatus*. *Biol Lett*, 17, 20210073.

57. Fahrbach SE, Moore D, Capaldi EA, Farris SM, Robinson GE. (1998). Experience-expectant
plasticity in the mushroom bodies of the honeybee. *Learn Mem,* 5:115-123.

58. Pardo-Sanchez J, Kou N, Tibbetts EA. (2022). Type and amount of social experience
influences individual face learning in paper wasps. *Behav Ecol and Sociobiol*, 76, 148.

59. Martin JP, Guo P, Mu L, Harley CM, Ritzmann RE. (2015). Central complex control of
movement in the freely walking cockroach. *Curr Biol*, 25, 2795-2803.

60. Turner-Evans D, Jayaraman V. (2016). The insect central complex. *Curr Biol*, 26, R445-
R460.

61. Pisokas I, Heinze S, Webb B. (2020). The head direction circuit of two insect species. *eLife*,
9, e53985.

62. Miroschnikow A, Schlegel P, Pankratz MJ. (2020). Making feeding decisions in the
Drosophila nervous system. *Curr Biol*, 30, R831-R840.

63. Chittka L, Niven J. (2009). Are bigger brains better? *Curr Biol*, *19*(21), R995-R1008.

64. Kamhi JF, Gronenberg W, Robson SKA, Traniello, JFA. (2016). Social complexity
influences brain investment and neural operation costs in ants. *Proc R Soc B*, 283,
20161949.

65. Muratore IB, Fandozzi EM, Traniello JFA. (2022). Behavioral performance and division of
labor influence brain mosaicism in the leafcutter ant *Atta cephalotes*. *J Comp Physiol A,*
208, 325-44.

66. Sheehan ZBV, Kamhi JF, Seid MA, Narendra A. (2019). Differential investment in brain
regions for a diurnal and nocturnal lifestyle in Australian *Myrmecia* ants. *J Comp Neurol*,
527, 1261-1277.

67. Arganda S, Hoadley AP, Razdan ES, Muratore IB, Traniello, JFA. (2020). The
neuroplasticity of division of labor: worker polymorphism, compound eye structure and
brain organization in the leafcutter ant *Atta cephalotes*. *J Comp Physiol A*, 206, 651-662.

68. Muscedere ML, Traniello JFA. (2012). Division of labor in the hyperdiverse ant genus *Pheidole* is associated with distinct subcaste- and age-related patterns of worker brain organization. *PLoS ONE,* 7.2, e31618

69. Hart T, Frank DD, Lopes LE, Olivos-Cisneros L, Lacy KD, Trible W, et al. (2023). Sparse and stereotyped encoding implicates a core glomerulus for ant alarm behavior. *Cell*, 186, 3079-3094.

70. Seid M, Goode K, Li C, Traniello JFA. (2008). Age□and subcaste□related patterns of serotonergic immunoreactivity in the optic lobes of the ant *Pheidole dentata*. *Dev Neurobiol*, 68.11: 1325-1333.

71. Wnuk A, Wiater M, Godzińska EJ. (2011). Effect of past and present behavioural specialization on brain levels of biogenic amines in workers of the red wood ant *Formica polyctena. Physiol Entomol*, 36, 54-61.

72. Zhang H, Liu Y, Zhou K, Wei W, Liu Y. (2021). Restoring sensorimotor function through neuromodulation after spinal cord injury: progress and remaining challenges. *Front Neurosci*, 15.

73. Marder E. (2012). Neuromodulation of neuronal circuits: back to the future. *Neuron*, 76, 1-11.

74. Sillar KT, Combes D, Simmers J. (2014). Neuromodulation in developing motor microcircuits. *Curr Opin in Neurobiol*, 29, 73-81.

75. Haddad SA, Marder E. (2018). Circuit robustness to temperature perturbation is altered by neuromodulators. *Neuron*, 100, 609-623.

76. Golowasch J. (2019). Neuromodulation of central pattern generators and its role in the functional recovery of central pattern generator activity. *J Neurophysiol*, 122, 300-315.

77. Thuma JB, Hobbs KH, Burstein HJ, Seiter NS, Hooper SL. (2013). Temperature sensitivity of the pyloric neuromuscular system and its modulation by dopamine. *PLoS ONE*, 8, e67930.

78. Mackevicius EL, Gu S, Denisenko NI, Fee MS. (2023). Self-organization of songbird neural sequences during social isolation. *eLife*, 12, e77262.

79. Sakurai A, Tamvacakis AN, Katz PS. (2014). Hidden synaptic differences in a neural circuit underlie differential behavioral susceptibility to a neural injury. *eLife*, 3, e02598.

80. Goaillard JM, Marder E. (2021). Ion channel degeneracy, variability, and covariation in neuron and circuit resilience. *Annu Rev Neurosci,* 44, 335-357.

81. Bonasio R, Zhang G, Ye C, Mutti NS, Fang X, Qin N, et al. (2010). Genomic comparison of the ants *Camponotus floridanus* and *Harpegnathos saltator*. *Science*, 329, 1068-1071.

82. Simola DF, Chaoyang Y, Mutti NS, Dolezal K, Bonasio R, Liebig J, et al. (2013). A chromatin link to caste identity in the carpenter ant *Camponotus floridanus. Genome Res*, 23, 486-496.

Visual Abstract. Newly eclosed (callow) *Camponotus floridanus* minor workers were subjected to variable periods of isolation to determine whether and how diminished social experience and isolation influence the development of the brain, including compartment volumes, biogenic amine levels, and behavioral performance.

Figure 1. Isolation effects on task performance. (a) Illustration of brood-care and predatory behavior assays. **(b)** Predation scores of workers isolated for varying time periods compared to control callow and mature workers. **(c)** Brood-care scores of isolates compared to callows and mature worker controls. **(d)**

733 Locomotion scores of isolates, callows and matures. X in box plot = average; horizontal line = median.
734 Sample sizes are above significance indicators. <0.001 = ***; 0.001-0.01 = **; 0.01-0.05 = *; >0.05 = ns.
735
736
737 **Figure 2. Isolation effects on size of brain compartments. (a-f)** brain compartment volumes (relative to
738 ROCB) for isolated and control workers. Averages, medians, and samples sizes as in Figure 1. **(a)** AL; **(b)**
739 MB-P; **(c)** MB-C; **(d)** CX; **(e)** OL; **(f)** SEZ. p <0.001 = ***; p<0.001-0.01 = **; p<0.01-0.05 = *; p>0.05
740 = ns.
741
742
743 **Figure 3. Isolation effects on monoamine levels (a-c), ratios of monoamines (d-e), and their**
744 **quantities relative to total biogenic amine levels (g-i).** . Averages, medians, and samples sizes as in
745 Figure 1. p <0.001 = ***; p<0.001-0.01 = **; p< 0.01-0.05 = *; p>0.05 = ns.
746
747
748 **Supplementary Figure 1. (a)** Head width variation of sampled workers **(B-C)** Principal component
749 analyses of behavior by **(b)** colony identity and **(c)** isolation duration**.** Colony identity and isolation
750 duration do not have significant effects. Significance level: <0.001 = ***; 0.001-0.01 = **; 0.01-0.05 = *;
751 >0.05 = ns. X = average; horizontal line = median. ns = not significant.
752
753 **Supplementary Figure 2. Task performance trajectories of isolated and control workers. (a)**
754 Illustration of assays to assess brood-care and predatory behaviors. **(b)** Predation scores of isolated and
755 control workers. **(c)** Brood-care scores of isolates, callows, and mature worker controls. **(d)** Locomotion
756 scores of isolates, callows and mature controls. **(b)-(d)**, X = average; horizontal line = median. Sample
757 sizes are above box plots. <0.001 = ***; 0.001-0.01 = **; 0.01-0.05 = *; >0.05 = ns.
758
759 **Supplementary Figure 3. Brain compartment growth trajectories for isolated workers. a-e**: brain
760 compartment volumes (relative to ROCB) for isolated and control workers. Sample sizes of each group
761 are above box plots. X in each box plot represents the average and horizontal line the median of each
762 group **(a)** AL; **(b)** MB-P; **(c)** MB-C; **(d)** CX; **(e)** OL; **(f)** SEZ. p <0.001 = ***; p<0.001-0.01 = **;
763 p<0.01-0.05 = *; p>0.05 = ns.
764
765 **Supplementary Figure 4. Trajectories and colony effects on biogenic monoamine levels. (a-b)**
766 Principal component analyses of biogenic amines clustered by **(a)** colony identity and **(b)** isolation
767 duration.
768
769 **Supplementary Figure 5. Biogenic amine levels across isolation trajectories (a-c), ratios of DA, OA,**
770 **and 5HT (d-f), and levels of DA, OA, and 5HT to total biogenic amines (g-i). (c-e)** Trajectories of
771 biogenic amine titers in response to isolation period. **(d-f)** Trajectories of biogenic amine titer ratios in
772 response to isolation period. **(h-i)** Relative biogenic amine to total amine titer ratios. Averages, medians,
773 and samples sizes as in Figure 3. p <0.001 = ***; p<0.001-0.01 = **; p< 0.01-0.05 = *; p>0.05 = ns.

Isolation Period

Parent Colony Increasing Time

Callow Mature

Neuroanatomy

Biogenic Amines

Behavior