**Pan-cancer characterization of ncRNA synergistic competition**

**uncovers potential carcinogenic biomarkers**

Junpeng Zhang[1*], Lin Liu[2], Xuemei Wei[1], Chunwen Zhao[1], Sijing Li[1], Jiuyong Li[2], and Thuc Duy Le[2*]

[1] School of Engineering, Dali University, Dali, Yunnan 671003, China

[2] UniSA STEM, University of South Australia, Mawson Lakes, SA 5095, Australia

*To whom correspondence should be addressed. Email: zhangjunpeng411@gmail.com

Correspondence may also be addressed to Thuc Duy Le. Email: thuc.le@unisa.edu.au

**Abstract**

Non-coding RNAs (ncRNAs) act as important modulators of gene expression and they have

been confirmed to play critical roles in the physiology and development of malignant tumors.

Understanding the synergism of multiple ncRNAs in competing endogenous RNA (ceRNA)

regulation can provide important insights into the mechanisms of malignant tumors caused by

ncRNA regulation. In this work, we present a framework, SCOM, for identifying ncRNA

synergistic competition. We systematically construct the landscape of ncRNA synergistic

competition across 31 malignant tumors, and reveal that malignant tumors tend to share hub

ncRNAs rather than the ncRNA interactions involved in the synergistic competition. In

addition, the synergistic competition ncRNAs (i.e. ncRNAs involved in the synergistic

competition) are likely to be involved in drug resistance, contribute to distinguishing

molecular subtypes of malignant tumors, and participate in immune regulation. Furthermore,

SCOM can help to infer ncRNA synergistic competition across malignant tumors and

uncover potential diagnostic and prognostic biomarkers of malignant tumors. Altogether, the

SCOM    framework    and    the    resulting    web-based    database    SCOMdb

(www.comblab.cn/SCOMdb/) serve as a useful resource for exploring ncRNA regulation and

to accelerate the identification of carcinogenic biomarkers.

**Author summary**

Abundant evidence reveals that ncRNAs are important modulators of gene expression, and it

is important for us to understand the regulation of ncRNAs in malignant tumors. In this work,

we hypothesize that ncRNAs acting as ceRNAs can crosstalk with each other in the form of

synergistic competition. We name the hypothesis as the *ncRNA synergistic competition*

*hypothesis*. To uncover potential carcinogenic biomarkers, we present a framework, SCOM,

for identifying ncRNA synergistic competition in malignant tumors. From the perspective of

synergistic competition, we have shown that the synergistic competition ncRNAs are likely to

involve in drug resistance, contribute to distinguishing molecular subtypes of malignant

tumors, and participate in immune regulation. In addition, the synergistic competition

ncRNAs are potential diagnostic and prognostic biomarkers of malignant tumors. We believe

that the proposed framework SCOM and the web-based database SCOMdb can contribute to

the design of reliable diagnosis and treatment biomarkers for malignant tumors.

## Introduction

Malignant tumors are a type of complex human diseases characterized by uncontrolled growth and spread of malignant cells and tissue infiltration [1]. According to the latest global cancer statistics from International Agency for Research on Cancer (IARC), malignant tumors are the second leading cause of death worldwide, and severely threaten human life and health [2]. Depending on the type of malignant tumor cells, malignant tumors are mainly divided into two categories: cancer and sarcoma [3]. Compared with healthy cells, the structure and function of genes in tumor cells have changed. It is found that polygenic synergy is one of the key characteristics of malignant tumors [4], suggesting that malignant tumors are usually polygenic disorders and their pathogenesis is caused by multi-gene synergy.

Non-coding RNAs (ncRNAs) are a class of various RNA transcripts that usually cannot be translated into proteins. Ordinarily, ncRNAs are categorized into two types: regulatory ncRNAs and structural ncRNAs [5]. For example, microRNAs (miRNAs), long non-coding RNAs (lncRNAs), circular RNAs (circRNAs) and pseudogenes belong to the type of regulatory ncRNAs, while ribosome RNAs (rRNAs) and transfer RNAs (tRNAs) are considered as structural ncRNAs. Existing studies [6,7] have demonstrated that ncRNAs are closely associated with the occurrence and development of malignant tumors, which opens a promising avenue for the diagnosis and preventive treatment of malignant tumors.

As regulators, ncRNAs can modulate the expression of thousands of genes in the manner

of trans-acting and cis-acting [8], thereby driving cell biological responses and fates [7]. In

addition to the role of regulators, by soaking up miRNAs, ncRNAs can act as competing

endogenous RNAs (ceRNAs) [9] which compete with the target messenger RNAs (mRNAs)

of the soaked miRNAs. In the ceRNA context, miRNAs serve as  mediators to connect non-

coding and coding RNAs, resulting in the ceRNA network (consisting of ceRNAs and

mRNAs), a more comprehensive regulatory network than the miRNA regulatory network

[10]. So far, three types of ncRNAs, including lncRNAs, circRNAs and pseudogenes have

been widely known to function as ceRNAs [11], and are regarded as potential carcinogenic

biomarkers in clinical applications [12,13]. For example, lncRNA *H19* serves as a ceRNA for

*LIN28* to promote breast cancer stem cell maintenance [14], circRNA *circTP63* functions as a

ceRNA for *FOXM1* to accelerate lung squamous cell carcinoma progression [15], and

pseudogene *PTENP1* acts as a ceRNA for *PTEN* in prostate cancer [16].

The ceRNA regulation of malignant tumors involves the interactions between multiple

non-coding and coding genes. In the ceRNA network in which ncRNAs are involved, the

competitive relationships between ncRNAs and target mRNAs are not one-to-one, but many-

to-many. For example, previous studies [17–25] have demonstrated that the three ncRNAs,

*H19*, *MALAT1* and *PVT1* synergistically compete with the four target genes,  *KLF4*, *VEGFA*,

*ZEB1* and *ZEB2*.  In this work, we call such collective competition by a group of ncRNAs

against a group of mRNAs as "ncRNA synergistic competition". It is observed that such

complicated gene regulation carries out important tasks in physiological and pathological

processes, and is considered to be a key component of cellular regulatory networks [7].

Furthermore, the entire cell system presents the characteristics of functional modularity, and

each gene functional module (i.e. a group of genes interacting or crosstalking with each other

frequently and not by chance) is responsible for specific biological functions in complex

human diseases, including malignant tumors [26]. These findings indicate that ncRNAs,

acting as ceRNAs, may synergistically rather than just independently play important roles in

malignant tumors.

In this work, to uncover potential carcinogenic biomarkers, we present the SCOM

(Synergistic COMpetition) framework to characterize ncRNA synergistic competition in Pan-

cancer. SCOM first systematically predicts ncRNA-related ceRNA networks across 31 tumor

types (containing 8001 tumor samples). Based on the proposed *ncRNA synergistic*

*competition hypothesis*, SCOM applies three criteria, (i) significant synergistic competition

for mRNAs, (ii) significant positive correlation, and (iii) significant sensitive correlation

conditioning on the mRNAs synergistically competed against by the ncRNAs, to predict the

ncRNA synergistic competition network (consisting of ncRNAs acting as ceRNAs) from the

gene expression data of ncRNAs and target mRNAs and the predicted ncRNA-related ceRNA

networks. By using SCOM, we provide a landscape of ncRNA synergistic competition in

Pan-cancer, contributing to the discovery of potential carcinogenic biomarkers. Ultimately,

SCOM provides a framework for discovering the relationships between the synergistic

competition ncRNAs (i.e. ncRNAs involved in the synergistic competition) and malignant

tumors, and can help to elucidate ncRNA synergistic competition mechanisms in malignant

tumors.

## Materials and methods

### Synergistic COMpetition (SCOM)

The proposed SCOM framework is based on the *ncRNA synergistic competition hypothesis*

(**Fig 1a**). For soaking up miRNAs, a pool of ncRNA transcripts (acting as ceRNAs) competes

with mRNAs through miRNA response elements (MREs). As a result, the competitive

interactions between ncRNAs and mRNAs form a ceRNA network (consisting of ncRNAs

acting as ceRNAs and target mRNAs). In the ceRNA network, we have observed that the

competitive relationships between ncRNAs and mRNAs are not one-to-one but many-to-

many, revealing synergistic competition of ncRNAs. Based on this observation, we

hypothesize that ncRNAs acting as ceRNAs can crosstalk with each other in the form of

synergistic competition (i.e. ncRNAs synergistically compete with mRNAs). We name the

hypothesis as the *ncRNA synergistic competition hypothesis*.

112
113 **Fig 1. Schematic illustration of SCOM.** (a) An illustration of the ncRNA synergistic competition
114 hypothesis. ncRNA *A* and ncRNA *B* acting as ceRNAs collectively compete with mRNAs for soaking up
115 miRNAs. In the context of ceRNA competition, the crosstalk between ncRNA *A* and ncRNA *B* is in the
116 form of synergistic competition. (b) For each tumor type in Pan-cancer, SCOM firstly extends the
117 SPONGE [33] method to predict ncRNA-related (including lncRNA- and pseudogene-related) ceRNA

network by incorporating gene (including miRNAs, lncRNAs, pseudogenes, and mRNAs) expression data in matched tumor tissues and priori information of miRNA-target (including miRNA-lncRNA, miRNA-pseudogene and miRNA-mRNA) interactions. By integrating gene (including lncRNAs and pseudogenes acting as ceRNAs, and target mRNAs) expression data in matched tumor tissues and predicted ncRNA-related ceRNA networks, SCOM utilise three criteria to infer ncRNA synergistic competition network for each tumor type in Pan-cancer. Based on the predicted ncRNA synergistic competition networks across 31 malignant tumors, SCOM further uncovers potential carcinogenic biomarkers.

The first step of SCOM (**Fig 1b**) is to predict ncRNA-related ceRNA network of each

malignant tumor type. The matched ncRNA (including miRNA, lncRNA and pseudogene)

and mRNA expression data in Pan-cancer (consisting of 8001 matched tumor samples) are

obtained from The Cancer Genome Atlas (TCGA) [27] project. The information of miRNA

targets is obtained by integrating five well-known databases, including miRTarBase v9.0 [28],

TarBase v8.0 [29], LncBase v2.0 [30], NPInter 4.0 [31] and ENCORI [32]. By integrating

gene (including ncRNAs and mRNAs) expression data and putative miRNA-target

interactions, SCOM extends SPONGE [33] (a three-criterion evaluation method) to predict

ncRNA-related ceRNA networks across 31 malignant tumors. The three-criterion evaluation

method predicts ceRNA interactions by integrating sample-matched ncRNA (miRNA,

lncRNA and pseudogene) and mRNA expression data, and putative miRNA-target

interactions. We reason that if a candidate ncRNA-mRNA pair meets the three criteria

(significant sharing of miRNAs, significantly positive correlation, and significantly sensitive

correlation conditioning on shared miRNAs) with $p$-value $< 0.05$, the candidate ncRNA-

mRNA pair is regarded as a ceRNA interaction. All of the predicted ceRNA interactions form

a large-scale ncRNA-related ceRNA network, with genes (including lncRNAs, pseudogenes

and mRNAs) denoted as nodes and the ncRNA-mRNA competition relationships as

undirected edges.

The second step of SCOM (**Fig 1b**) is to infer ncRNA synergistic competition network of

each malignant tumor type. For a malignant tumor type, to predict the ncRNA synergistic

competition network, we use the expression data of ncRNAs acting as ceRNAs and target

mRNAs and the predicted ncRNA-related ceRNA networks. Following the *ncRNA*

*synergistic competition hypothesis*, we consider a ncRNA-ncRNA pair having a synergistic

competition relationship if the pair have (i) significant synergistic competition with mRNAs,

(ii) significant positive correlation, and (iii) significant sensitive correlation conditioning on

their competing mRNAs. The synergistic competition network of a malignant tumor type is

then formed by integrating all of ncRNA synergistic competition relationships.

**Pan-cancer transcriptomics data**

After removing non-matching tumor samples between ncRNAs (including miRNAs,

lncRNAs and pseudogenes) and mRNA expression data, the Pan-cancer transcriptomics data

includes the expression data of 17,212 ncRNAs (including 894 miRNAs, 4366 lncRNAs, and

11,952 pseudogenes) and 19,068 mRNAs in 8001 matched tumor samples covering 31 tumor

types (**S1 Data**), and the clinical information of 8001 matched tumor samples (**S2 Data**).

**Putative miRNA-target interactions**

Three types of putative miRNA-target interactions, including miRNA-mRNA, miRNA-lncRNA, and miRNA-pseudogene are used for predicting ncRNA-related ceRNA networks. The putative miRNA-mRNA interactions are acquired from miRTarBase v9.0 [28] and TarBase v8.0 [29], the putative miRNA-lncRNA interactions are obtained from LncBase v2.0 [30] and NPInter 4.0 [31], and the putative miRNA-pseudogene interactions are from ENCORI [32] (the previous version is starBase). In total, we have 762,540 miRNA-mRNA interactions, 225,063 miRNA-lncRNA interactions, and 91,619 miRNA-pseudogene interactions as priori information of miRNA-target interactions.

**Predicting ceRNA networks in Pan-cancer**

Given the gene expression data of miRNAs, lncRNAs, pseudogenes, and mRNAs, and miRNA-target interactions, three criteria, (i) significant sharing of miRNAs, (ii) significant positive correlation, and (iii) significant sensitive correlation conditioning on shared miRNAs, are used to predict ceRNA network (see **Supplementary Methods** in **S1 File**).

In this work, a candidate ncRNA-mRNA pair meeting the above three criteria with $p$-value $< 0.05$ is regarded as a ceRNA interaction. For each of the 31 tumor types, we integrate all of ceRNA interactions to form the ceRNA network of the tumor type.

**Constructing ncRNA synergistic competition networks in Pan-cancer**

Given the gene expression data of lncRNAs and pseudogenes acting as ceRNAs, and miRNA

targets mRNAs and the predicted ncRNA-related ceRNA networks, three criteria, (i)

significant synergistic competition for mRNAs, (ii) significant positive correlation, and (iii)

significant sensitive correlation conditioning on synergistically competed mRNAs are used to

construct ncRNA synergistic competition network.

The first criterion uses a hypergeometric test to assess the significance of synergistic

competition of ncRNA $i$ and ncRNA $j$ for mRNAs. The significance $p$-value of the test is

computed as:

184
$$p_{ij} = 1 - \sum_{x_{ij}=0}^{L_{ij}-1} \frac{\binom{M_{ij}}{x_{ij}}\binom{N_{ij}-M_{ij}}{K_{ij}-x_{ij}}}{\binom{N_{ij}}{K_{ij}}} \tag{1}$$

where $N_{ij}$ is the number of all mRNAs in a Pan-cancer dataset, $M_{ij}$ and $K_{ij}$ represent the total

numbers of mRNAs competing with ncRNA $i$ and ncRNA $j$ respectively, and $L_{ij}$ is the

number of mRNAs synergistically competed by ncRNA $i$ and ncRNA $j$.

The second criterion requires that ncRNA $i$ and ncRNA $j$ are positively correlated. The

Pearson [34] method is also utilized to evaluate whether the expression level of ncRNA $i$ and

the expression level of ncRNA $j$ are positively correlated at a significant level (e.g. 0.05). The

correlation $cor_{ij}$ between ncRNA $i$ and ncRNA $j$ is computed as:

192
$$cor_{ij} = cor(p, q) = \frac{\sum_{k=1}^{s}(p_k - \overline{p})(q_k - \overline{q})}{\sqrt{\sum_{k=1}^{s}(p_k - \overline{p})^2}\sqrt{\sum_{k=1}^{s}(q_k - \overline{q})^2}}$$ (2)

193  where $p = (p_1, p_2, ..., p_s)$ and $q = (q_1, q_2, ..., q_s)$ denote the expression level of ncRNA $i$ and

194  ncRNA $j$ respectively, $\overline{p}$ and $\overline{q}$ represent the average expression level of ncRNA $i$ and ncRNA

195  $j$ respectively, and $s$ is the number of samples for each tumor type. Accordingly, the

196  significance $p$-value $pc_{ij}$ is calculated as:

197
$$pc_{ij} = 2pt\left(cor_{ij}\sqrt{\frac{s-2}{1-(cor_{ij})^2}}\right)$$ (3)

where the $pt$ function is used to calculate the probability of $cor_{ij}\sqrt{\frac{s-2}{1-(cor_{ij})^2}}$.

The third criterion is used to evaluate the influence of synergistically competed mRNAs

on ncRNA $i$ and ncRNA $j$. The sensitive correlation [35] is also employed to calculate the

influence of the mRNAs synergistically competed by ncRNA $i$ and ncRNA $j$, and the null

model [33] is also applied to evaluate the significance of this influence. The sensitive

correlation $sc_{ij}$ is calculated as:

204
$$sc_{ij} = cor_{ij} - pcor_{ij}$$ (4)

205  where $pcor_{ij}$ is the partial correlation between ncRNA $i$ and ncRNA $j$, i.e. the correlation

206  conditioning on the mRNAs synergistically competed by the ncRNAs, which is the

207  correlation between ncRNA $i$ and ncRNA $j$ when the influence of the synergistically

208  competed mRNAs is eliminated. $pcor_{ij}$ is calculated as:

$$pcor_{ij} = cor(p,q\,|\,M) = cor(p,q\,|\,(M_1,M_2,...,M_n))$$

209
$$= \frac{cor(p,q\,|\,(M_1,M_2,...,M_{n-1})) - cor(p,M_n\,|\,(M_1,M_2,...,M_{n-1}))cor(q,M_n\,|\,(M_1,M_2,...,M_{n-1}))}{\sqrt{1 - cor(p,M_n\,|\,(M_1,M_2,...,M_{n-1}))^2}\,\sqrt{1 - cor(q,M_n\,|\,(M_1,M_2,...,M_{n-1}))^2}} \quad (5)$$

where $p$, $q$ and $M$ denote the expression level of ncRNA $i$, ncRNA $j$, and the synergistically

competed $n$ mRNAs respectively, $cor(p,q\,|\,(M_1,M_2,...,M_n))$ denotes the partial correlation

between $p$ and $q$ conditioning on $(M_1,M_2,...,M_n)$, $cor(p,q\,|\,(M_1,M_2,...,M_{n-1}))$ represents the

partial correlation between $p$ and $q$ conditioning on $(M_1,M_2,...,M_{n-1})$ ,

$cor(p,M_n\,|\,(M_1,M_2,...,M_{n-1}))$ is the partial correlation between $p$ and $M_n$ conditioning on

$(M_1,M_2,...,M_{n-1})$, $cor(q,M_n\,|\,(M_1,M_2,...,M_{n-1}))$ is the partial correlation between $q$ and $M_n$

conditioning on $(M_1,M_2,...,M_{n-1})$.

To evaluate the significance of $sc_{ij}$, the null model hypothesizes that the synergistically

competed mRNAs do not affect the correlation between ncRNA $i$ and ncRNA $j$, i.e. the

sensitive correlation (the difference between $cor_{ij}$ and $pcor_{ij}$) between ncRNA $i$ and ncRNA $j$

is 0. The number of datasets sampled is set to 1E+03 for the null model, and the pre-

computed covariance matrices in the R package SPONGE [33] are used to build the null

model. Based on the constructed null model, the significance $p$-value of $sc_{ij}$ is calculated.

In this study, a candidate ncRNA-ncRNA pair meeting the three criteria for synergistic

competition with $p$-value < 0.05 is considered as a ncRNA synergistic competition

relationship. For each tumor type, we combine all of the ncRNA synergistic competition

relationships to create the ncRNA synergistic competition network for the tumor type.

**Network topological analysis**

Network topological features yield useful insights into biological networks, e.g. degree, path and density. If the node degree of a biological network obeys a power law distribution, the biological network is regarded as a scale-free network. The path and density features are used to measure the tightness of a biological network. Compared with the random networks, a shorter characteristic path length or higher density of a biological network means that the biological network is more likely to be a small-world network [36,37].

The topological features (degree, path and density) of ncRNA-related ceRNA networks and ncRNA synergistic competition networks are analyzed using the R package igraph [38], a useful tool for network analysis and visualization. The ncRNA-related ceRNA networks and ncRNA synergistic competition networks are undirected networks, and the degree of a node is the number of edges connected with the node. The Kolmogorov-Smirnov (KS) test [39] is used to evaluate whether the node degree of a network follows a power law distribution. If the $p$-value of the KS test is smaller than a cutoff (e.g. 0.05), the node degree of a ncRNA-related ceRNA network or a ncRNA synergistic competition network does not follow a power law distribution, then these indicate that the ncRNA-related ceRNA network or the ncRNA synergistic competition network is not a scale-free network. To determine whether the ncRNA-related ceRNA networks and the ncRNA synergistic competition networks are small-world networks, we generate 100 random networks (the same number of

nodes and edges with real network) for each ncRNA-related ceRNA network or ncRNA

synergistic competition network. The Student's *t*-test [40] is applied to evaluate whether the

characteristic path length (or density) of a ncRNA-related ceRNA network or a ncRNA

synergistic competition network is smaller (or higher) than that of its corresponding random

networks at a significance level. For a ncRNA-related ceRNA network or a ncRNA

synergistic competition network, if the *p*-value of the Student's *t*-test is smaller than a cutoff

(e.g. 0.05), the characteristic path length (or density) of a ncRNA-related ceRNA network or

a ncRNA synergistic competition network is significantly smaller (or higher) than that of its

corresponding random networks, then the network is a small-world network.

**Network and hub similarity calculation**

In terms of ncRNA synergistic competition networks or hub ncRNAs, the network or hub

similarity between tumor *c* and tumor *d* is computed as follows.

258
$$sim_{cd} = \frac{|SCR_c \cap SCR_d|}{min(|SCR_c|, |SCR_d|)} \tag{6}$$

259 where $SCR_c$ and $SCR_d$ denote the identified ncRNA synergistic competition regulation

260 (networks or hub ncRNAs) in tumor *c* and tumor *d*, respectively, $|SCR_c \cap SCR_d|$ is the

261 number of common ncRNA synergistic competition relationships or hub ncRNAs between

262 $SCR_c$ and $SCR_d$, and $min(|SCR_c|, |SCR_d|)$ denotes the smaller number of ncRNA

263 synergistic competition relationships or hub ncRNAs between $SCR_c$ and $SCR_d$.

**Hub ncRNA identification**

Hub ncRNAs are defined as highly connected nodes in a ncRNA synergistic competition network. Here, we use the cumulative probability of Poisson distribution to decide whether a ncRNA is a hub ncRNA:

$$p(x \geq k) = 1 - \sum_{i=0}^{k-1} \frac{\lambda^i e^{-\lambda}}{i!} \tag{7}$$

where $\lambda = np$, $p = \frac{m}{A_n^2}$, $n$ is the number of ncRNAs, $m$ is the number of ncRNA synergistic competition relationships in a ncRNA synergistic competition network, and $A_n^2$ is the number of all possible ncRNA synergistic competition pairs. Smaller $p$-value of a ncRNA indicates that the ncRNA is more likely to be a hub ncRNA. In this work, the cutoff of the $p$-value is set to 0.05.

**Uncovering conserved and rewired interactions and hubs**

The ncRNA synergistic competition interaction or hub ncRNA existing in at least two malignant tumors is defined as a conserved ncRNA synergistic competition interaction or hub ncRNA, whereas the ncRNA synergistic competition interaction or hub ncRNA existing in only one malignant tumor is defined as a rewired ncRNA synergistic competition interaction or hub ncRNA. In the case of the identified ncRNA synergistic competition networks and hub ncRNAs across 31 malignant tumors, we can uncover conserved and rewired ncRNA synergistic competition interactions and hub ncRNAs.

**Multi-class classification analysis**

To understand the classification performance of the discovered conserved ncRNA synergistic

competition interactions and hub ncRNAs, we train a classification model to evaluate the

discriminating ability of the features in classifying tumor types, and also train a model to

classify BRCA subtypes (LumA, LumB, Her2, Basal and Normal) to understand the ability of

the features in discriminating cancer subtypes. We carry out multi-class classification

analysis by using the Binary Relevance (BR) [41] strategy in the R package utiml [42] to

perform multi-class classification analysis. In the BR strategy, the Support Vector Machine

(SVM) [43] implemented in the R package e1071 [44] is selected as the base classifier to

create the multi-class model. We use 22 metrics (accuracy, average-precision, clp, coverage,

F1, hamming-loss, macro-AUC, macro-F1, macro-precision, macro-recall, margin-loss,

micro-AUC, micro-F1, micro-precision, micro-recall, mlp, one-error, precision, ranking-loss,

recall, subset-accuracy and wlp as described in **S1 File**) and conduct 10-fold cross-validation

to evaluate the performance of ncRNAs in conserved ncRNA synergistic competitions and

hub ncRNAs. Also, we apply a commonly used baseline method [45] for comparison in the

evaluation of the classification performance.

**Survival analysis**

To compute the risk score of each malignant tumor sample, the multivariate Cox model [46]

in the R package survival [47] is fitted by using ncRNAs in conserved ncRNA synergistic

301 competition and hub ncRNAs as features. According to the risk scores of each malignant

302 tumor sample, the top one-third of the malignant tumor samples are put in the high risk group

303 and the remaining two thirds are put in the low risk group. In terms of the gene expression

304 levels of the ncRNAs, the Log-rank test is applied to assess the distinction of the feature

305 ncRNAs between the high and the low risk groups or between different molecular subtypes.

306 In addition, the proportional hazard ratio (HR) between the high and the low risk groups or

307 between different molecular subtypes is also computed.

308 **Enrichment analysis**

309 To understand the biological significance of hub ncRNAs across the 31 malignant tumors, we

310 conduct enrichment analysis in three categories, disease, cell marker and cancer hallmark

311 using LncSEA [48]. The hub ncRNAs are enriched in significant disease, cell marker and

312 cancer hallmark terms with an adjusted $p$-value (adjusted by Benjamini-Hochberg method)

313 cutoff of 0.05. Moreover, LncSEA is also used to uncover whether the synergistic

314 competition ncRNAs are involved in drug resistance or not. The synergistic competition

315 ncRNAs are significantly enriched in drug terms with an adjusted $p$-value (adjusted by

316 Benjamini-Hochberg method) less than 0.05.

317 **Molecular subtyping of malignant tumors**

318 Within the same type of malignant tumor, malignant tumors have great phenotypic

319 differences and genetic heterogeneity. Because of this, molecular subtyping is usually applied

to divide malignant tumors into subtypes with distinct molecular and clinical characteristics.

To assess the performance of the synergistic competition ncRNAs in molecular subtyping, we

use an ensemble method implemented in the CancerSubtypes [49] R package (combining the

Similarity Network Fusion (SNF) [50] and Consensus Clustering (CC) [51] methods) to

generate malignant tumor subtypes. For each malignant tumor, the number of molecular

subtypes is empirically set according to the previous molecular subtyping studies (**S3 Data**).

**Immune regulation analysis**

Numerous ncRNAs have been shown to play important roles in immune regulation. For the

immune regulation analysis, we obtain the significant immune-related ncRNA-pathway and

ncRNA-cell pairs from ImmReg [52]. Based on the significance of the enrichment of

ncRNAs with immune pathways and the significance of the correlation between ncRNAs and

immune cells, we further determine which synergistic competition ncRNAs are significantly

enriched in immune pathways and correlated with immune cells in each malignant tumor.

**Results**

**The ceRNA regulation landscape in Pan-cancer**

Using the transcriptomics data in TCGA, we apply SCOM for detecting two types of ceRNA

regulation landscape across the 31 malignant tumors. In the ceRNA regulation landscape, the

two types of ceRNA networks (lncRNA- and pseudogene-related ceRNA networks) have

different numbers of ceRNA interactions across the 31 malignant tumors. We have found that

the number of ceRNA interactions of the pseudogene-related ceRNA networks is

significantly larger than that of the lncRNA-related ceRNA networks (**Fig 2a**, *p*-value =

1.84E-13 with paired Student's *t*-test). Network topological analysis reveals that 5 out of the

31 (16.13%) lncRNA-related ceRNA networks and 16 out of the 31 (51.61%) pseudogene-

related ceRNA networks follow a power law distribution (**Fig 2b**). After merging the lncRNA

and pseudogene related ceRNA networks of each tumor, 14 out of the 31 (45.16%) merged

ncRNA-related ceRNA networks follow a power law distribution (**Fig S1** in **S1 File**).

Moreover, 7 out of the 31 (22.58%) lncRNA-related ceRNA networks and none of the

pseudogene-related ceRNA networks have shorter characteristic path lengths than their

corresponding random networks, and 30 out of the 31 (96.77%) lncRNA (and pseudogene

related) ceRNA networks have higher densities than their corresponding random networks

(**Fig 2b**). In the merged ncRNA-related ceRNA networks of each tumor, we have found that

none of the ncRNA-related ceRNA networks have shorter characteristic path lengths than

their corresponding random networks, and all of the ncRNA-related ceRNA networks have

higher densities than their corresponding random networks (**Fig S1** in **S1 File**). These

observations indicate that less than or nearly half of the ceRNA networks across the 31

malignant tumors tend to be scale-free networks, and most or all of the ceRNA networks

across the 31 malignant tumors are likely to be small-world networks with higher densities

than their corresponding random networks.

With the lncRNA and pseudogene related ceRNA networks, the similarity intervals

between the 31 malignant tumors are [0.13 0.52] and [0.10 0.44], respectively (**Fig 2c**).

Moreover, in terms of the ncRNA (including both lncRNA and pseudogene) related ceRNA

networks, the similarity interval between the 31 malignant tumors is [0.12 0.47] (**Fig S2** in **S1**

**File**). This result shows that the malignant tumors may have common ncRNA-related ceRNA

regulation. We have also discovered that each tumor ceRNA network varies in similarity to

other tumor ceRNA networks, revealing known and hidden relationships among these

malignant tumors. In addition, we have observed that ~16.45% lncRNA-related ceRNA

regulations and ~28.58% pseudogene-related ceRNA regulations only exist in one malignant

tumor (**Fig 2d**). Overall, ~24.80% ncRNA-related ceRNA regulations are occurred in one

malignant tumor (**Fig S3** in **S1 File**). For malignant tumors, the percentages of the rewired

ncRNA-related ceRNA regulations are different, demonstrating different degrees of

heterogeneity across malignant tumors (**Fig S4** in **S1 File**). Particularly, 106 (less than 0.01%)

ncRNA-related ceRNA regulations are conserved in the 31 malignant tumors (**Fig 2e**),

indicating a low conservation of ceRNA regulation in Pan-cancer. By conducting the multi-

class classification and survival analysis, we have shown that the conserved ncRNA-related

ceRNA regulations in the 31 malignant tumors have better performance than the baseline

method in classifying 31 malignant tumors, and can distinguish the high and the low risk

groups of each malignant tumor at a significant level (Log-rank $p$-value < 0.05 and HR > 2)

(**Fig S5** in **S1 File**). This result reveals that the conserved ncRNA-related ceRNA regulations

may act as a common core of ceRNA regulation across malignant tumors.

**Fig 2. The ceRNA regulation landscape across malignant tumors.** (a) The number of ncRNA-related

ceRNA interactions in each malignant tumor. (b) The topological characteristics of ncRNA-related ceRNA

networks for the 31 malignant tumors. The Degree column is the *p*-value of fitting power-law distribution.

The *p*-values equal to or more than 0.05 indicate that the degree of ncRNA-related ceRNA networks obeys

the power-law distribution. The Path and Density columns represent the $-\log10$(*p*-value) of Student's *t*-test,

and the *p*-values less than 0.05 display that the characteristic path lengths (or densities) of ncRNA-related

ceRNA networks are significantly shorter (or higher) than those of their corresponding random networks.

(c) The similarity matrix shows the similarity between each pair of lncRNA (upper triangle) and

pseudogene (lower triangle) related ceRNA networks across 31 malignant tumors. (d) The radar chart

shows the percentage of ceRNA regulations predicted in different numbers of malignant tumors. (e) The

conserved ceRNA regulation predicted in 31 malignant tumors.

**The ncRNA synergistic competition landscape in Pan-cancer**

Taking advantage of the transcriptomics data in TCGA and the predicted ncRNA-related

ceRNA interactions, we utilise SCOM for uncovering ncRNA (including lncRNAs and

pseudogenes) synergistic competition landscape across 31 malignant tumors. Similar to the

number of ceRNA interactions, the number of ncRNA synergistic competition interactions

(including lncRNA-lncRNA, lncRNA-pseudogene and pseudogene-pseudogene synergistic

competition pairs) also varies across the 31 malignant tumors. The largest and least numbers

of ncRNA synergistic competition interactions are inferred in Stomach adenocarcinoma

(STAD) and Cholangiocarcinoma (CHOL), respectively (**Fig 3a**). This result may be

explained in part by the different number of ceRNA interactions predicted in the 31

malignant tumors. Network topological analysis shows that 27 out of the 31 (87.10%)

ncRNA synergistic competition networks follow a power law distribution (**Fig 3b**).

Furthermore, 19 out of the 31 (61.29%) ncRNA synergistic competition networks have

shorter characteristic path lengths than their corresponding random networks, and all of the

ncRNA synergistic competition networks have higher densities than their corresponding

random networks (**Fig 3b**). These findings indicate that most of the ncRNA synergistic

competition networks across the 31 malignant tumors are scale-free networks, and small-

world networks with shorter characteristic path lengths and higher densities than their

corresponding random networks.

With the ncRNA synergistic competition networks, the similarity among the 31 malignant

tumors is less than 6% (**Fig 3c**). This low similarity indicates that the malignant tumors may

share a small portion of ncRNA synergistic competition with each other. In addition, ~94.06%

ncRNA synergistic competition interactions only occur in one malignant tumor, and none of

the ncRNA synergistic competition interactions exist in more than six malignant tumors (**Fig**

**3d**). Specifically, we have observed that the percentage of the conserved ncRNA synergistic

competition interactions in at least two malignant tumors is ~5.94%. This result reveals that

the ncRNA synergistic competition interactions of different malignant tumors tend to be

rewired in their respective tumor microenvironments. To understand whether the ncRNAs of

the conserved synergistic competition in at least two malignant tumors are biologically

meaningful, we further perform enrichment analysis in three categories including diseases,

cell markers and cancer hallmarks. The functional enrichment analysis has revealed that the

ncRNAs of the conserved synergistic competition in at least two malignant tumors are

significantly enriched in diseases (e.g. Cancer), cell markers (e.g. Exhausted CD8+ T cell),

and cancer hallmarks (e.g. Epithelial-mesenchymal transition) common to multiple malignant tumors (**S4 Data**). Moreover, the percentage of the rewired ncRNA synergistic competition for each malignant tumor is all more than 80%, demonstrating a great heterogeneity across malignant tumors in terms of ncRNA synergistic competition (**Fig 3e**).

**Fig 3. Network analysis of ncRNA synergistic competition across malignant tumors.** (a) The number of ncRNA synergistic competition interactions in each malignant tumor. (b) The topological characteristics

of ncRNA synergistic competition networks for the 31 malignant tumors. The Degree column is the *p*-value of fitting power-law distribution. The *p*-values equal to or more than 0.05 indicate that the degree of the ncRNA synergistic competition networks obeys the power-law distribution. The Path and Density columns represent the –log10(*p*-value) of Student's *t*-test, and the *p*-values less than 0.05 suggest that the characteristic path lengths (or densities) of ncRNA synergistic competition networks are significantly shorter (or higher) than those of their corresponding random networks. (c) The similarity matrix showing the similarity between each pair of ncRNA synergistic competition networks across 31 malignant tumors. (d) The radar chart showing the percentages of ncRNA synergistic competition interactions predicted in different number of malignant tumors. (e) The bar chart showing the percentages of ncRNA synergistic competition interactions predicted in one malignant tumor.

Based on the topological features of the ncRNA synergistic competition networks, we further identify hub ncRNAs associated with malignant tumors. We have discovered that the number of hub ncRNAs vary across different malignant tumors (**Fig 4a**). In terms of the hub ncRNAs, the similarity between the 31 malignant tumors is [0.07 0.79] (**Fig S6** in **S1 File**). This result shows that the malignant tumors may have common hub ncRNAs. In addition, we have observed that the hub ncRNAs can significantly distinguish the high risk and the low risk groups of each malignant tumor (**Fig S7a** in **S1 File**), and the hub ncRNAs covering 96.77% malignant tumors are significantly enriched at least one of Disease, Cell markers and Cancer hallmarks terms (**Fig S7b** in **S1 File**, **S5 Data**). Different from the ncRNA synergistic competition networks, less than 13% hub ncRNAs are only in one malignant tumor but nearly 90% hub ncRNAs exist in at least two malignant tumors (**Fig 4b**). In particular, ~7.63% hub ncRNAs are even conserved in at least 16 malignant tumors. This result reveals that the hub ncRNAs of different malignant tumors tend to be conserved in their respective tumor microenvironments.

To understand whether there is a common core of hub ncRNAs across malignant tumors,

we investigate the conserved hub ncRNAs occurred in at least 16 malignant tumors.

Functional analysis indicates that the conserved hub ncRNAs have better performance than

the baseline method in classifying the 31 malignant tumors (**Fig 4c**), and can significantly

distinguish the high risk and the low risk groups of each malignant tumor (**Fig 4d**). This

result shows that the conserved hub ncRNAs may act as potential diagnostic and prognostic

biomarkers across malignant tumors. Moreover, the conserved hub ncRNAs are significantly

enriched in diseases (**Fig 4e**), cell markers (**Fig 4f**) and cancer hallmarks (**Fig 4g**) common to

several malignant tumors. This result suggests that the malignant tumors may have common

cell markers and cancer hallmarks.

In addition to the conserved hub ncRNAs, the rewired hub ncRNAs switching from tumor

to tumor are important to understand the heterogeneity of malignant tumors. We have found

that the percentage of the rewired hub ncRNAs for each malignant tumor is less than 7%,

revealing a small heterogeneity across malignant tumors in terms of hub ncRNAs (**Fig 4h**).

Multi-class classification analysis shows that the rewired hub ncRNAs have better

performance than the baseline method in classifying the 31 malignant tumors (**Fig S7c** in **S1**

**File**). Moreover, survival analysis displays that the rewired hub ncRNAs can significantly

distinguish the high risk and the low risk groups of each malignant tumor (**Fig S7d** in **S1**

**File**). This result suggests that the rewired hub ncRNAs may also act as potential diagnostic

and prognostic biomarkers across malignant tumors.

**Fig 4. Hub analysis of ncRNA synergistic competition across malignant tumors.** (a) The number of

hub ncRNAs in each malignant tumor. (b) The radar chart displaying the percentage of hub ncRNAs

predicted in different number of malignant tumors. (c) Multi-class classification analysis of conserved hub

ncRNAs in at least 16 malignant tumors. (d) Survival analysis of conserved hub ncRNAs in at least 16

malignant tumors. The symbol "/" stands for an infinite value. (e-g) Enrichment analysis (three categories

including disease, cell marker and cancer hallmark) of conserved hub ncRNAs in at least 16 malignant

tumors. (h) The bar chart indicating the percentage of hub ncRNAs predicted in one malignant tumor.

**Synergistic competition ncRNAs are involved in drug resistance**

It is reported that ncRNAs play important roles in drug resistance, and the dysregulation of

them can induce resistance to anti-cancer therapy and cause a failure of malignant tumor

clinical treatment [53,54]. Drug enrichment analysis shows that the ncRNA synergistic

competition networks of ~96.77% of the 31 malignant tumors and the hub ncRNAs of ~64.52%

of the 31 malignant tumors are significantly enriched in at least one drug resistance term (**Fig**

**5a**, **S6 Data and S7 Data**). We have found that the ncRNA synergistic competition networks

of 29 malignant tumors are involved in the drug *cisplatin* resistance (**Fig 5b**). The drug

*cisplatin* is a well-known chemotherapy medication, and can be used to treat a number of

cancers including bladder cancer, brain tumors, cervical cancer, esophageal cancer, head and

neck cancer, lung cancer, mesothelioma, ovarian cancer, etc [55]. Moreover, the resistance of

two drugs *L-685458* and *LBW242* is closely associated with the ncRNA synergistic

competition networks of 4 malignant tumors (**Fig 5b**). The drug *L-685458* is a potent

transition state analog (TSA) γ-secretase inhibitor (GSI) and is an attractive therapeutic

candidate for Alzheimer Disease (AD) and cancers [56], and the drug *LBW242* is a potent and

orally active proapoptotic IAP inhibitor and potentiates treat multiple myeloma and ovarian

cancer [57,58]. Similar to the ncRNA synergistic competition networks, the top three

enriched drugs of hub ncRNAs across malignant tumors are also *cisplatin*, *L-685458* and

*LBW242* (**Fig 5c**). This result suggests that the synergistic competition ncRNAs (i.e. ncRNAs

involved in the synergistic competition) across malignant tumors are involved in drug

resistance, and may act as targets for anti-cancer therapy.

**Fig 5. Drug resistance analysis of ncRNA synergistic competition networks and hub ncRNAs across**
**malignant tumors.** (a) The number of significantly enriched drugs for each malignant tumor. (b)
Relationships between ncRNA synergistic competition networks and enriched drugs. For example, the

node "Network_ACC" denotes the ncRNA synergistic competition network in ACC. Larger drug nodes
denote more links with ncRNA synergistic competition networks in malignant cancers. (c) Relationships
between hub ncRNAs and enriched drugs. For example, the node "Hub_ACC" denotes the hub ncRNAs in
ACC. Larger drug nodes denote more links with hub ncRNAs in malignant cancers.

**Synergistic competition ncRNAs contribute to distinguishing molecular subtypes of**

**malignant tumors**

Malignant tumors are a collection of heterogeneous diseases with both phenotypic and

genetic heterogeneity, and a malignant tumor usually possess multiple distinct molecular

subtypes [59]. Molecular subtyping analysis of the synergistic competition ncRNAs (i.e.

ncRNAs involved in the synergistic competition) shows that the average silhouette width of

the molecular subtyping of ~90.32% of the 31 malignant tumors is equal to or greater than

0.60 (**Fig 6a**), indicating that the synergistic competition ncRNAs can well assign malignant

tumor samples into distinct molecular subtypes. In the case of using hub ncRNAs, the

molecular subtypes of ~32.26% of the 31 malignant tumors can be significantly distinguished

(Log-rank test $p$-value $< 0.05$, **Fig 6a**). Moreover, in the case of using ncRNA synergistic

competition networks, the molecular subtypes of ~25.81% of the 31 malignant tumors can be

significantly distinguished (Log-rank test $p$-value $< 0.05$, **Fig 6a**). For seven malignant tumor

types, including Adrenocortical carcinoma (ACC), Kidney renal clear cell carcinoma (KIRC),

Brain Lower Grade Glioma (LGG), Liver hepatocellular carcinoma (LIHC), Pancreatic

adenocarcinoma (PAAD), Skin Cutaneous Melanoma (SKCM) and Uveal Melanoma (UVM),

the hub ncRNAs and ncRNA synergistic competition networks all have contributions to

distinguish molecular subtypes of them. For example, in the case of using hub ncRNAs and

ncRNA synergistic competition networks, LGG can be divided into three distinct molecular

subtypes (**Fig 6b** and **Fig 6c**). This result reveals that the synergistic competition ncRNAs

contribute to distinguishing molecular subtypes of malignant tumors, and may act as

prognostic biomarkers in several malignant tumors. Since the molecular subtypes (including

Luminal A, Luminal B, HER-2, Basal, Normal-like) of Breast invasive carcinoma (BRCA)

are mostly accepted by researchers, we focus on studying the BRCA subtype classification

performance by using hub ncRNAs and ncRNA synergistic competition networks in BRCA.

Multi-class classification analysis shows that the synergistic competition ncRNAs have better

performance than the baseline method in classifying BRCA subtypes (**Fig 6d**), suggesting

that the synergistic competition ncRNAs may act as potential diagnostic biomarkers of

BRCA.

**Fig 6. Molecular subtype analysis of malignant tumors.** (a) Molecular subtyping of the 31 malignant tumors using ncRNA synergistic competition networks and hub ncRNAs. Higher average silhouette width indicates that a malignant tumor sample is better matched with the identified molecular subtypes. (b) Molecular subtyping of LGG using ncRNA synergistic competition network. (c) Molecular subtyping of LGG using hub ncRNAs. (d) Classification analysis of BRCA subtypes.

**Synergistic competition ncRNAs participate in immune regulation**

As regulatory elements in immune system, ncRNAs can influence immune development and

function, leading to immune diseases including autoimmunity and haematological cancers

[60,61]. Immune regulation analysis reveals that the ncRNA synergistic competition

networks of all the 31 malignant tumors and the hub ncRNAs of 30 malignant tumors are

significantly involved in at least one immune-related pathway (**Fig 7a**, **S8 Data and S9**

**Data**). For instance, the ncRNA synergistic competition networks of the 31 malignant tumors

are closely associated with six immune-related pathways (*Antigen Processing and*

*Presentation*, *Cytokine Receptors*, *Antimicrobials*, *Cytokines*, *Natural Killer Cell Cytotoxicity*,

*TCR signaling Pathway*) (**Fig 7c**). In addition, the hub ncRNAs of 29 malignant tumors are

closely related to two immune-related pathways (*Antigen Processing and Presentation*,

*Cytokine Receptors*) (**Fig 7d**). Moreover, the ncRNA synergistic competition networks and

the hub ncRNAs of 31 malignant tumors are significantly associated with at least 40 immune

cells (**Fig 7b**, **S10 Data and S11 Data**). This result suggests that the synergistic competition

ncRNAs (i.e. ncRNAs involved in the synergistic competition) take part in immune

regulation, and the dysregulation of them may influence the occurrence and development of

immune diseases including malignant tumors.

**Fig 7. Immune regulation analysis of synergistic competition ncRNAs across malignant tumors.** (a)

The number of significant immune-related pathways associated with each malignant tumor. (b) The

number of significant immune cells associated with each malignant tumor. (c) Relationships between

ncRNA synergistic competition networks and immune-related pathways. For example, the node

"Network_ACC" denotes the ncRNA synergistic competition network in ACC. Larger immune-related

pathway nodes denote more links with ncRNA synergistic competition networks in malignant cancers. (d)

Relationships between hub ncRNAs and immune-related pathways. For example, the node "Hub_ACC"

denotes the hub ncRNAs in ACC. Larger immune-related pathway nodes denote more links with hub

ncRNAs in malignant cancers.

**SCOMdb: a web-based resource for ncRNA synergistic competition in Pan-cancer**

To provide a convenient and web-accessible resource of ncRNA synergistic competition

landscape across malignant tumors for biologists, we have developed a comprehensive and

interactive database, SCOMdb (www.comblab.cn/SCOMdb/). By using this web-based

resource, users can browse the ceRNA interactions in specific malignant tumor of interest

(**Fig 8a**). It also provides the ncRNA synergistic competition interactions in malignant tumors

(**Fig 8b**). In addition, users can query the hub ncRNAs of interest across malignant tumors

(**Fig 8c**). All the data in this work can be downloaded for downstream analysis (**Fig 8d**). To

help users apply SCOM to new datasets, we have also provided the SCOM R package for

inferring ncRNA synergistic competition by integrating sample-matched gene (ncRNA and

mRNA) expression data and putative miRNA-target interactions (**Fig 8e**). The provided

resources will be continuously updated, and should serve as a valuable resource for biologists

interested in investigating ncRNA synergistic competition across malignant tumors.

**Fig 8. Diagram of SCOMdb database.** (a) The search of ceRNA regulation in Pan-cancer. (b) The search of ncRNA synergistic competition in Pan-cancer. (c) The search of hub ncRNAs in Pan-cancer. (d) All resources are provided to be downloaded for further analysis. (e) The SCOM method is available for users to study ncRNA synergistic competition by using their own datasets.

## Conclusions and discussion

Abundant evidence reveals that ncRNAs can serve as important regulators across malignant tumors. However, from the perspective of synergistic competition, ncRNA regulation is still unearthed. In this work, we present the *ncRNA synergistic competition hypothesis* and the SCOM framework, and further elaborate the application of the SCOM framework to

comprehensively infer the synergistic competition ncRNAs that potentially act as

carcinogenic biomarkers. From the perspective of synergistic competition, we have shown

that the synergistic competition ncRNAs are potential diagnostic and prognostic biomarkers

of malignant tumors. Moreover, the synergistic competition ncRNAs are likely to involve in

drug resistance, contribute to distinguishing molecular subtypes of malignant tumors, and

participate in immune regulation. The web-based database SCOMdb

(www.comblab.cn/SCOMdb/) provides a valuable resource for exploring ncRNA regulation

across malignant tumors.

As important regulators of gene expression, ncRNAs play important roles in the

physiology and development of malignant tumors. Yet, the experimentally validated

carcinogenic biomarkers are far from complete. Thus, it is challenging to confirm the

accuracy of the findings by the SCOM framework. To show the effectiveness of SCOM, we

conduct a series of functional analyses including enrichment analysis, molecular subtyping

analysis, multi-class classification analysis, survival analysis, and immune regulation analysis

for the synergistic competition ncRNAs. We have found that the conserved ncRNA

synergistic competition ncRNAs and hub ncRNAs are significantly enriched in diseases, cell

markers and cancer hallmarks. In addition, the conserved and rewired hub ncRNAs may act

as potential diagnostic and prognostic biomarkers across malignant tumors. Moreover, the

synergistic competition ncRNAs are likely to be involved in drug resistance, contribute to

distinguishing molecular subtypes of malignant tumors, and participate in immune regulation.

These results indicate that the SCOM framework can effectively infer the synergistic

competition ncRNAs with biological significance. Although experimentally validated

carcinogenic biomarkers are limited, we have discovered that a significant number of

synergistic competition ncRNAs are confirmed as carcinogenic biomarkers by the literature

(**Fig S8** in **S1 File**, **S12 Data**).

The findings in this paper have provided several biological insights although the identified

synergistic competition ncRNAs need to be further confirmed by subsequent biological

experiments. Firstly, not all the ncRNA synergistic competition networks follow a power law

distribution. For example, the ncRNA synergistic competition networks in Kidney

Chromophobe (KICH), Lung adenocarcinoma (LUAD), Lung squamous cell carcinoma

(LUSC) and Rectum adenocarcinoma (READ) do not obey the power-law distribution. This

result indicates that the synergistic competition ncRNAs in these four malignant tumors

might equally interact with each other. Secondly, a majority of the ncRNA interactions

involved in the synergistic competition are rewired, but a minority of hub ncRNAs is rewired.

This result suggests that malignant tumors are likely to share hub ncRNAs rather than the

ncRNA interactions involved in the synergistic competition. Thirdly, the similarity of hub

ncRNAs between the 31 malignant tumors is significantly larger than that of ncRNA

synergistic competition networks ($p$-value $<$ 2.20E-16 with paired Student's $t$-test).

Moreover, the percentage of rewired ncRNA synergistic competition interactions is

significantly larger than that of rewired hub ncRNAs ($p$-value $<$ 2.20E-16 with paired

Student's $t$-test). This result reveals that although malignant tumors have great differences in

ncRNA synergistic competition networks, the hub ncRNAs between malignant tumors tend to

be similar. Fourthly, the synergistic competition ncRNAs can distinguish five BRCA

subtypes with accuracy equal to or more than 70%. This result shows that the synergistic

competition ncRNAs might be good diagnostic biomarkers in BRCA. Fifthly, we have used

in this study the transcriptomics data in TCGA [27], which is one of the most well-known

cancer genomics programs. However, the SCOM framework can also be applied to the

transcriptomics data in other sources (e.g. CCLE [62] and MET500 [63]) to investigate

ncRNA synergistic competition. Finally, SCOM is not limited to the exploration of lncRNA

and pseudogene synergistic competition. It can also be used to infer the synergistic

competition of other ncRNAs (e.g. circRNAs) that are important modulators to malignant

tumors.

Prediction of ncRNA-related ceRNA network is a critical step to infer ncRNA synergistic

competition. Yet, there is still a lack of study on the prediction of ncRNA-related ceRNA

networks. In the current work, we apply three criteria (significant sharing of miRNAs,

significant positive correlation, and significant sensitive correlation conditioning on shared

miRNAs) to predict ncRNA-related ceRNA networks in 31 malignant tumors. Since the three

criteria have been successfully used to investigate ceRNA regulation [33,35,64–66], the

ncRNA-related ceRNA network predictions are promising. However, evaluating whether a

ncRNA can act as a ceRNA to compete with mRNAs is still challenging. For example, the

stoichiometry of miRNAs and ncRNAs is considered as a vital factor to predict ceRNA

interactions of a ncRNA of interest [67]. With deeper understanding of ceRNA mechanisms,

we can more accurately predict ncRNA-related ceRNA networks to improve the inference of

ncRNA synergistic competition and accelerate the discovery of carcinogenic biomarkers.

Given gene expression data and miRNA-target interactions, SCOM constructs the

landscape of ncRNA synergistic competition across malignant tumors at the bulk level.

Recently, increasingly advanced single-cell and spatial RNA sequencing technology has

opened a way for exploring ncRNA synergistic competition at the single-cell level. With the

advance of single-cell and spatial RNA sequencing technology, SCOM will contribute to

uncover more carcinogenic biomarkers.

ncRNAs are emerging as potential carcinogenic biomarkers. It is important for us to

understand the regulation of ncRNAs in malignant tumors. In-depth study of synergistic

competition ncRNAs inferred in this study would contribute to the design of reliable

diagnosis and treatment biomarkers for human malignant tumors.

**References**

1. Hanahan D. Hallmarks of cancer: new dimensions. Cancer Discov. 2022;12: 31–46.
doi:10.1158/2159-8290.CD-21-1059

2. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global
cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for
36 cancers in 185 countries. CA Cancer J Clin. 2021;71: 209–249.
doi:10.3322/caac.21660

3. Weinberg RA. The biology of cancer, 2nd Edition. Second Edition. New York;

4. Sparmann A, van Lohuizen M. Polycomb silencers control cell fate, development and
cancer. Nat Rev Cancer. 2006;6: 846–856. doi:10.1038/nrc1991

5. Dahariya S, Paddibhatla I, Kumar S, Raghuwanshi S, Pallepati A, Gutti RK. Long non-
coding RNA: Classification, biogenesis and functions in blood cells. Mol Immunol.
2019;112: 82–92. doi:10.1016/j.molimm.2019.04.011

6. Esteller M. Non-coding RNAs in human disease. Nat Rev Genet. 2011;12: 861–874.
doi:10.1038/nrg3074

7. Anastasiadou E, Jacob LS, Slack FJ. Non-coding RNA networks in cancer. Nat Rev
Cancer. 2018;18: 5–18. doi:10.1038/nrc.2017.99

8. Fu XD. Non-coding RNA: a new frontier in regulatory biology. Natl Sci Rev. 2014;1:
190–204. doi:10.1093/nsr/nwu008

9. Salmena L, Poliseno L, Tay Y, Kats L, Pandolfi PP. A ceRNA hypothesis: the Rosetta
Stone of a hidden RNA language? Cell. 2011;146: 353–358.
doi:10.1016/j.cell.2011.07.014

10. Smillie CL, Sirey T, Ponting CP. Complexities of post-transcriptional regulation and the
modeling of ceRNA crosstalk. Crit Rev Biochem Mol Biol. 2018;53: 231–245.
doi:10.1080/10409238.2018.1447542

11. Tay Y, Rinn J, Pandolfi PP. The multilayered complexity of ceRNA crosstalk and
competition. Nature. 2014;505: 344–352. doi:10.1038/nature12986

12. Wang Y, Hou J, He D, Sun M, Zhang P, Yu Y, et al. The emerging function and
mechanism of ceRNAs in cancer. Trends Genet. 2016;32: 211–224.
doi:10.1016/j.tig.2016.02.001

13. Qi X, Zhang D-H, Wu N, Xiao J-H, Wang X, Ma W. ceRNA in cancer: possible
functions and clinical implications. J Med Genet. 2015;52: 710–718.
doi:10.1136/jmedgenet-2015-103334

14. Peng F, Li T-T, Wang K-L, Xiao G-Q, Wang J-H, Zhao H-D, et al. H19/let-7/LIN28
reciprocal negative regulatory circuit promotes breast cancer stem cell maintenance.
Cell Death Dis. 2017;8: e2569. doi:10.1038/cddis.2016.438

15. Cheng Z, Yu C, Cui S, Wang H, Jin H, Wang C, et al. circTP63 functions as a ceRNA
to promote lung squamous cell carcinoma progression by upregulating FOXM1. Nat
Commun. 2019;10: 3200. doi:10.1038/s41467-019-11162-4

16. Poliseno L, Salmena L, Zhang J, Carver B, Haveman WJ, Pandolfi PP. A coding-
independent function of gene and pseudogene mRNAs regulates tumour biology. Nature.
2010;465: 1033–1038. doi:10.1038/nature09144

17. Arun K, Arunkumar G, Bennet D, Chandramohan SM, Murugan AK, Munirajan AK.
Comprehensive analysis of aberrantly expressed lncRNAs and construction of ceRNA
network in gastric cancer. Oncotarget. 2018;9: 18386–18399.
doi:10.18632/oncotarget.24841

18. Xiong JM, Liu H, Chen J, Zou QQ, Wang YYJ, Bi G. Curcumin nicotinate suppresses
abdominal aortic aneurysm pyroptosis via lncRNA PVT1/miR-26a/KLF4 axis through
regulating the PI3K/AKT signaling pathway. Toxicol Res (Camb). 2021;10: 651–661.
doi:10.1093/toxres/tfab041

19. Huang M, Zhong Z, Lv M, Shu J, Tian Q, Chen J. Comprehensive analysis of
differentially expressed profiles of lncRNAs and circRNAs with associated co-
expression and ceRNA networks in bladder carcinoma. Oncotarget. 2016;7: 47186–
47200. doi:10.18632/oncotarget.9706

20. Sun X, Luo L, Li J. LncRNA MALAT1 facilitates BM-MSCs differentiation into
endothelial cells via targeting miR-206/VEGFA axis. Cell Cycle. 2020;19: 3018–3028.
doi:10.1080/15384101.2020.1829799

21. Lai SW, Chen MY, Bamodu OA, Hsieh M-S, Huang T-Y, Yeh C-T, et al. Exosomal
lncRNA PVT1/VEGFA axis promotes colon cancer metastasis and stemness by
downregulation of tumor suppressor miR-152-3p. Oxid Med Cell Longev. 2021;2021:
9959807. doi:10.1155/2021/9959807

22. Chen L, Yao H, Wang K, Liu X. Long non-coding RNA MALAT1 regulates ZEB1
expression by sponging miR-143-3p and promotes hepatocellular carcinoma progression.
J Cell Biochem. 2017;118: 4836–4843. doi:10.1002/jcb.26158

23. Yang T, Zhou H, Liu P, Yan L, Yao W, Chen K, et al. lncRNA PVT1 and its splicing
variant function as competing endogenous RNA to regulate clear cell renal cell
carcinoma progression. Oncotarget. 2017;8: 85353–85367.
doi:10.18632/oncotarget.19743

24. Wang Y, Zhou Y, Yang Z, Chen B, Huang W, Liu Y, et al. MiR-204/ZEB2 axis
functions as key mediator for MALAT1-induced epithelial-mesenchymal transition in
breast cancer. Tumour Biol. 2017;39: 1010428317690998.
doi:10.1177/1010428317690998

25. Chen SW, Zhu J, Ma J, Zhang J, Zuo S, Chen G-W, et al. Overexpression of long non-
coding RNA H19 is associated with unfavorable prognosis in patients with colorectal
cancer and increased proliferation and  migration in colon cancer cells. Oncol Lett.
2017;14: 2446–2452. doi:10.3892/ol.2017.6390

26.   Choobdar S, Ahsen ME, Crawford J, Tomasoni M, Fang T, Lamparter D, et al.
748           Assessment of network module identification across complex diseases. Nat Methods.
749           2019;16: 843–852. doi:10.1038/s41592-019-0509-5

750    27.   Weinstein JN, Collisson EA, Mills GB, Shaw KRM, Ozenberger BA, Ellrott K, et al.
751           The Cancer Genome Atlas Pan-Cancer analysis project. Nat Genet. 2013;45: 1113–1120.
752           doi:10.1038/ng.2764

753    28.   Huang H, Lin Y-C, Cui S, Huang Y, Tang Y, Xu J, et al. miRTarBase update 2022: an
754           informative resource for experimentally validated miRNA-target interactions. Nucleic
755           Acids Res. 2022;50: D222–D230. doi:10.1093/nar/gkab1079

756    29.   Karagkouni D, Paraskevopoulou MD, Chatzopoulos S, Vlachos IS, Tastsoglou S,
757           Kanellos I, et al. DIANA-TarBase v8: a decade-long collection of experimentally
758           supported miRNA-gene interactions. Nucleic Acids Res. 2018;46: D239–D245.
759           doi:10.1093/nar/gkx1141

760    30.   Paraskevopoulou MD, Vlachos IS, Karagkouni D, Georgakilas G, Kanellos I, Vergoulis
761           T, et al. DIANA-LncBase v2: indexing microRNA targets on non-coding transcripts.
762           Nucleic Acids Res. 2016;44: D231-238. doi:10.1093/nar/gkv1270

763    31.   Teng X, Chen X, Xue H, Tang Y, Zhang P, Kang Q, et al. NPInter v4.0: an integrated
764           database of ncRNA interactions. Nucleic Acids Res. 2020;48: D160–D165.
765           doi:10.1093/nar/gkz969

766    32.   Li JH, Liu S, Zhou H, Qu LH, Yang JH. starBase v2.0: decoding miRNA-ceRNA,
767           miRNA-ncRNA and protein-RNA interaction networks from large-scale CLIP-Seq data.
768           Nucleic Acids Res. 2014;42: D92-97. doi:10.1093/nar/gkt1248

769    33.   List M, Dehghani Amirabad A, Kostka D, Schulz MH. Large-scale inference of
770           competing endogenous RNA networks with sparse partial correlation. Bioinformatics.
771           2019;35: i596–i604. doi:10.1093/bioinformatics/btz314

772    34.   Pearson K. Notes on the history of correlation. Biometrika. 1920;13: 25–45.
773           doi:10.2307/2331722

774    35.   Paci P, Colombo T, Farina L. Computational analysis identifies a sponge interaction
775           network between long non-coding RNAs and messenger RNAs in human breast cancer.
776           BMC Syst Biol. 2014;8: 83. doi:10.1186/1752-0509-8-83

777    36.   Telesford QK, Joyce KE, Hayasaka S, Burdette JH, Laurienti PJ. The ubiquity of small-
778           world networks. Brain Connect. 2011;1: 367–375. doi:10.1089/brain.2011.0038

779    37.   Zaidi F. Small world networks and clustered small world networks with random
780           connectivity. Social Network Analysis and Mining. 2013;3: 51–63. doi:10.1007/s13278-
781           012-0052-1

782    38.   Csardi G, Nepusz T. The igraph software package for complex network research.
783           InterJournal, Complex Systems. 2006;1695: 1–9.

39.  Massey FJ. The Kolmogorov-Smirnov test for goodness of fit. Journal of the American
Statistical Association. 1951;46: 68–78. doi:10.2307/2280095

40.  Student. The probable error of a mean. Biometrika. 1908;6: 1–25. doi:10.2307/2331554

41.  Tsoumakas G, Katakis I, Vlahavas I. Mining multi-label data. Data mining and
knowledge discovery handbook Springer, Boston, MA. Springer, Boston, MA; 2009. pp.
667–685.

42.  Rivolli A, Carvalho ACPLF de. The utiml package: multi-label classification in R. The
R Journal. 2018;10: 24–37. doi:10.32614/RJ-2018-041

43.  Chang CC, Lin CJ. LIBSVM: a library for support vector machines. ACM Transactions
on Intelligent Systems and Technology. 2011;2: 1–27.

44.  Meyer D, Dimitriadou E, Hornik K, Weingessel A, Leisch F, Chang C-C, et al. e1071:
misc functions of the department of statistics, probability theory group (Formerly:
E1071), TU Wien. Available: https://CRAN.R-project.org/package=e1071

45.  Metz J, de Abreu LF, Cherman EA, Monard MC. On the estimation of predictive
evaluation measure baselines for multi-label learning. Ibero-American Conference on
Artificial Intelligence, Cartagena de Indias, Colombia. Springer; 2012. pp. 189–198.

46.  Andersen PK, Gill RD. Cox's regression model for counting processes: a large sample
study. Ann Statist. 1982;10: 1100–1120. doi:10.1214/aos/1176345976

47.  Terry M. Therneau, Patricia M. Grambsch. Modeling survival data: extending the Cox
model. New York: Springer; 2000.

48.  Chen J, Zhang J, Gao Y, Li Y, Feng C, Song C, et al. LncSEA: a platform for long non-
coding RNA related sets and enrichment analysis. Nucleic Acids Res. 2021;49: D969–
D980. doi:10.1093/nar/gkaa806

49.  Xu T, Le TD, Liu L, Su N, Wang R, Sun B, et al. CancerSubtypes: an R/Bioconductor
package for molecular cancer subtype identification, validation and visualization.
Bioinformatics. 2017;33: 3131–3133. doi:10.1093/bioinformatics/btx378

50.  Wang B, Mezlini AM, Demir F, Fiume M, Tu Z, Brudno M, et al. Similarity network
fusion for aggregating data types on a genomic scale. Nat Methods. 2014;11: 333–337.
doi:10.1038/nmeth.2810

51.  Monti S, Tamayo P, Mesirov J, Golub T. Consensus clustering: a resampling-based
method for class discovery and visualization of gene expression microarray data.
Machine Learning. 2003;52: 91–118. doi:10.1023/A:1023949509487

52.  Jiang T, Zhou W, Chang Z, Zou H, Bai J, Sun Q, et al. ImmReg: the regulon atlas of
immune-related pathways across cancer types. Nucleic Acids Res. 2021;49: 12106–
12118. doi:10.1093/nar/gkab1041

53.  Si W, Shen J, Zheng H, Fan W. The role and mechanisms of action of microRNAs in
cancer drug resistance. Clin Epigenetics. 2019;11: 25. doi:10.1186/s13148-018-0587-8

54.  Smallegan MJ, Rinn JL. Linking long noncoding RNA to drug resistance. Proc Natl
Acad Sci U S A. 2019;116: 21963–21965. doi:10.1073/pnas.1915690116

55.  Wishart DS, Feunang YD, Guo AC, Lo EJ, Marcu A, Grant JR, et al. DrugBank 5.0: a
major update to the DrugBank database for 2018. Nucleic Acids Res. 2018;46: D1074–
D1082. doi:10.1093/nar/gkx1037

56.  Yang G, Zhou R, Guo X, Yan C, Lei J, Shi Y. Structural basis of γ-secretase inhibition
and modulation by small molecule drugs. Cell. 2021;184: 521-533.e14.
doi:10.1016/j.cell.2020.11.049

57.  Weisberg E, Kung AL, Wright RD, Moreno D, Catley L, Ray A, et al. Potentiation of
antileukemic therapies by Smac mimetic, LBW242: effects on mutant FLT3-expressing
cells. Mol Cancer Ther. 2007;6: 1951–1961. doi:10.1158/1535-7163.MCT-06-0810

58.  Petrucci E, Pasquini L, Bernabei M, Saulle E, Biffoni M, Accarpio F, et al. A small
molecule SMAC mimic LBW242 potentiates TRAIL- and anticancer drug-mediated cell
death of ovarian cancer cells. PLoS One. 2012;7: e35073.
doi:10.1371/journal.pone.0035073

59.  Zhao L, Lee VHF, Ng MK, Yan H, Bijlsma MF. Molecular subtyping of cancer: current
status and moving toward clinical applications. Brief Bioinform. 2019;20: 572–584.
doi:10.1093/bib/bby026

60.  Mehta A, Baltimore D. MicroRNAs as regulatory elements in immune system logic. Nat
Rev Immunol. 2016;16: 279–294. doi:10.1038/nri.2016.40

61.  Peltier DC, Roberts A, Reddy P. LNCing RNA to immunity. Trends Immunol. 2022;43:
478–495. doi:10.1016/j.it.2022.04.002

62.  Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S, et al. The
Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug
sensitivity. Nature. 2012;483: 603–607. doi:10.1038/nature11003

63.  Robinson DR, Wu Y-M, Lonigro RJ, Vats P, Cobain E, Everett J, et al. Integrative
clinical genomics of metastatic cancer. Nature. 2017;548: 297–303.
doi:10.1038/nature23306

64.  Le TD, Zhang J, Liu L, Li J. Computational methods for identifying miRNA sponge
interactions. Brief Bioinform. 2017;18: 577–590. doi:10.1093/bib/bbw042

65.  Li Y, Jin X, Wang Z, Li L, Chen H, Lin X, et al. Systematic review of computational
methods for identifying miRNA-mediated RNA-RNA crosstalk. Brief Bioinform.
2019;20: 1193–1204. doi:10.1093/bib/bbx137

66.  Zhang J, Liu L, Zhang W, Li X, Zhao C, Li S, et al. miRspongeR 2.0: an enhanced R
package for exploring miRNA sponge regulation. Bioinformatics Advances. 2022;2:
vbac063. doi:10.1093/bioadv/vbac063

67.  Thomson DW, Dinger ME. Endogenous microRNA sponges: evidence and controversy.
Nat Rev Genet. 2016;17: 272–283. doi:10.1038/nrg.2016.20

**Supporting information**

**S1 Data. The matched tumor types in Pan-cancer dataset.**

**S2 Data. Clinical information of the matched 8001 tumor samples.**

**S3 Data. The molecular subtypes of malignant tumors in Pan-cancer dataset.**

**S4 Data. Enrichment analysis of the conserved ncRNA synergistic competition in at**

**least two malignant tumors.**

**S5 Data. Enrichment analysis of hub ncRNAs across malignant tumors.**

**S6 Data. Drug enrichment of ncRNA synergistic competition networks.**

**S7 Data. Drug enrichment of hub ncRNAs.**

**S8 Data. Relationships between ncRNA synergistic competition networks and immune-**

**related pathways across malignant tumors.**

**S9 Data. Relationships between hub ncRNAs and immune-related pathways across**

**malignant tumors.**

**S10 Data. Relationships between ncRNA synergistic competition networks and immune**

**cells across malignant tumors.**

**S11 Data. Relationships between hub ncRNAs and immune cells across malignant**

**tumors.**

**S12 Data. Experimentally validated synergistic competition ncRNAs as biomarkers**

**across malignant tumors.**

**S1 File. Supporting file.**

**a**

mRNAs

ceRNA network

ncRNA A

ncRNA B

ncRNA A

Common mRNAs

ncRNA A

ncRNA B

ncRNA B

--- Competition
····· Synergistic competition

**b**

Pan-cancer

LGG

HNSC
ESCA
THCA
THYM
PCPG
BRCA
LUAD
LUSC
LIHC
PAAD
STAD
CHOL
KICH
ACC
KIRP
KIRC
SARC
COAD
UVM
UCEC
READ
UCS
OV
BLCA
CESC
TGCT
PRAD
DLBC
MESO
SKCM

ncRNAs          Priori information          mRNAs

Pan-cancer tissues          Pan-cancer tissues

miRNA-target interactions

Predicting ceRNA network

**Predicted ncRNA-related ceRNA network**

Tumor type 1          Tumor type 2          Tumor type C

ceRNA interactions    ceRNA interactions    ...    ceRNA interactions

**Inferring ncRNA synergistic competition network**

Criterion I:
Synergistic competition

Criterion II:
Positive correlation

Criterion III:
Sensitive correlation

ncRNA A          Common mRNAs

ncRNA B

ncRNA A          ncRNA A          ncRNA A          ncRNA A

ncRNA B          ncRNA B          ncRNA B          ncRNA B

**Predicted ncRNA synergistic competition network**

Tumor type 1          Tumor type 2          Tumor type C

● ncRNAs acting as ceRNAs          Synergistic competition

**a**

Number of ncRNA synergistic competition interactions

Tumors

**b**

p-value

−log10 (p-value)

Degree | Path | Density

**c**

**d** The radar chart of conserved ncRNA synergistic competition

**e** The bar chart of rewired ncRNA synergistic competition

Percentage of rewired synergistic competition (%)

**a** — Number of hub ncRNAs

| Tumor | Value |
|---|---|
| UVM | 346 |
| UCS | 305 |
| UCEC | 1246 |
| THYM | 867 |
| THCA | 1215 |
| TGCT | 979 |
| STAD | 2357 |
| SKCM | 1236 |
| SARC | 770 |
| READ | 208 |
| PRAD | 1317 |
| PCPG | 352 |
| PAAD | 618 |
| OV | 1143 |
| MESO | 165 |
| LUSC | 962 |
| LUAD | 1054 |
| LIHC | 768 |
| LGG | 1238 |
| KIRP | 1264 |
| KIRC | 1490 |
| KICH | 129 |
| HNSC | 1405 |
| ESCA | 1448 |
| DLBC | 460 |
| COAD | 517 |
| CHOL | 17 |
| CESC | 628 |
| BRCA | 1402 |
| BLCA | 905 |
| ACC | 95 |

**b** The radar chart of conserved hub ncRNAs

**c** Classification analysis of conserved hub ncRNAs

| Metrics | Conserved | Baseline |
|---|---|---|
| accuracy | **0.91** | 0.09 |
| average-precision | **0.95** | 0.17 |
| clp | **0.00** | 0.03 |
| coverage | **0.21** | 14.42 |
| F1 | **0.91** | 0.09 |
| hamming-loss | **0.01** | 0.06 |
| macro-AUC | **0.99** | 0.50 |
| macro-F1 | **0.89** | 0.01 |
| macro-precision | **0.89** | 0.00 |
| macro-recall | **0.90** | 0.03 |
| margin-loss | **0.21** | 14.42 |
| micro-AUC | **1.00** | 0.53 |
| micro-F1 | **0.91** | 0.09 |
| micro-precision | **0.90** | 0.09 |
| micro-recall | **0.92** | 0.09 |
| mlp | **0.00** | 0.97 |
| one-error | **0.09** | 0.91 |
| precision | **0.91** | 0.09 |
| ranking-loss | **0.01** | 0.48 |
| recall | **0.92** | 0.09 |
| subset-accuracy | **0.89** | 0.09 |
| wlp | **0.00** | 0.97 |

**d** Survival analysis of conserved hub ncRNAs

| Tumors | p-value | HR | HRlow95 | HRup95 |
|---|---|---|---|---|
| ACC | 5.33E-15 | 9.79 | 3.33 | 28.81 |
| BLCA | <2.22E-16 | 10.15 | 6.53 | 15.77 |
| BRCA | <2.22E-16 | 14.78 | 8.13 | 26.88 |
| CESC | <2.22E-16 | 8.80 | 3.99 | 19.42 |
| CHOL | 4.63E-09 | 9.31 | 2.08 | 41.74 |
| COAD | <2.22E-16 | 8.44 | 3.66 | 19.45 |
| DLBC | 6.51E-08 | 18.77 | 3.12 | 113.01 |
| ESCA | <2.22E-16 | 7.50 | 3.18 | 17.72 |
| HNSC | <2.22E-16 | 12.06 | 7.80 | 18.66 |
| KICH | 6.19E-07 | / | / | / |
| KIRC | <2.22E-16 | 15.26 | 7.05 | 33.00 |
| KIRP | <2.22E-16 | 11.18 | 4.04 | 30.94 |
| LGG | <2.22E-16 | 10.38 | 6.16 | 17.48 |
| LIHC | <2.22E-16 | 14.06 | 8.49 | 23.30 |
| LUAD | <2.22E-16 | 10.85 | 6.92 | 17.03 |
| LUSC | <2.22E-16 | 12.98 | 7.80 | 21.62 |
| MESO | <2.22E-16 | 7.31 | 2.99 | 17.90 |
| OV | <2.22E-16 | 7.49 | 4.75 | 11.81 |
| PAAD | <2.22E-16 | 7.69 | 3.83 | 15.44 |
| PCPG | 1.64E-05 | / | / | / |
| PRAD | 3.74E-10 | 23.38 | 5.30 | 103.27 |
| READ | 3.05E-10 | 10.00 | 2.38 | 42.11 |
| SARC | <2.22E-16 | 8.68 | 4.58 | 16.43 |
| SKCM | <2.22E-16 | 8.50 | 5.44 | 13.28 |
| STAD | <2.22E-16 | 10.95 | 6.16 | 19.47 |
| TGCT | 2.36E-03 | / | / | / |
| THCA | 7.27E-14 | / | / | / |
| THYM | 6.66E-09 | 20.61 | 3.23 | 131.69 |
| UCEC | <2.22E-16 | 64.84 | 35.96 | 116.91 |
| UCS | 1.11E-16 | 9.22 | 3.04 | 27.95 |
| UVM | 2.16E-14 | 11.05 | 3.19 | 38.26 |

**e** Diseases associated with conserved hub ncRNAs

**f** Cell markers associated with conserved hub ncRNAs

**g** Cancer hallmarkers associated with conserved hub ncRNAs

**h** The bar chart of rewired hub ncRNAs — Percentage of rewired hub ncRNAs (%)

| Tumor | Value |
|---|---|
| UVM | 1.16 |
| UCS | 0 |
| UCEC | 1.61 |
| THYM | 1.5 |
| THCA | 0.58 |
| TGCT | 2.76 |
| STAD | 6.66 |
| SKCM | 1.78 |
| SARC | 0.52 |
| READ | 0.96 |
| PRAD | 0.84 |
| PCPG | 1.42 |
| PAAD | 0.32 |
| OV | 1.49 |
| MESO | 0 |
| LUSC | 1.04 |
| LUAD | 1.42 |
| LIHC | 0.65 |
| LGG | 0.81 |
| KIRP | 1.66 |
| KIRC | 2.42 |
| KICH | 0.78 |
| HNSC | 1.57 |
| ESCA | 2.14 |
| DLBC | 1.09 |
| COAD | 0.58 |
| CHOL | 0 |
| CESC | 0 |
| BRCA | 2.07 |
| BLCA | 0.88 |
| ACC | 1.05 |

## a
### Drug enrichment

## c
### Visualization of the relationships between hub ncRNAs and enriched drugs

## b
### Visualization of the relationships between ncRNA synergistic competition networks and enriched drugs

**a** Molecular subtyping

**b** Molecular subtyping of LGG using ncRNA synergistic competition network

**c** Molecular subtyping of LGG using hub ncRNAs

**d** BRCA subtype classification

| Metrics | Network | Hub | Baseline |
|---|---|---|---|
| accuracy | 0.70 | 0.79 | 0.40 |
| average-precision | 0.83 | 0.89 | 0.60 |
| clp | 0.06 | 0.00 | 0.20 |
| coverage | 0.48 | 0.26 | 1.38 |
| F1 | 0.71 | 0.81 | 0.40 |
| hamming-loss | 0.12 | 0.08 | 0.24 |
| macro-AUC | 0.72 | 0.94 | 0.50 |
| macro-F1 | 0.59 | 0.74 | 0.11 |
| macro-precision | 0.61 | 0.75 | 0.08 |
| macro-recall | 0.62 | 0.75 | 0.20 |
| margin-loss | 0.48 | 0.26 | 1.38 |
| micro-AUC | 0.88 | 0.96 | 0.63 |
| micro-F1 | 0.70 | 0.80 | 0.40 |
| micro-precision | 0.68 | 0.77 | 0.40 |
| micro-recall | 0.72 | 0.83 | 0.40 |
| mlp | 0.24 | 0.02 | 0.80 |
| one-error | 0.30 | 0.20 | 0.60 |
| precision | 0.70 | 0.79 | 0.40 |
| ranking-loss | 0.12 | 0.06 | 0.34 |
| recall | 0.72 | 0.83 | 0.40 |
| subset-accuracy | 0.67 | 0.75 | 0.40 |
| wlp | 0.24 | 0.04 | 0.80 |

**a** Immune-related pathways

**b** Immune cells

**c** Relationships between ncRNA synergistic competition networks and immune-related pathways

**d** Relationships between hub ncRNAs and immune-related pathways