Diagnosis of planktonic trophic network dynamics with sharp qualitative changes

ABSTRACT

Trophic interaction networks are notoriously difficult to understand and to diagnose (i.e., to identify contrasted network functioning regimes). Such ecological networks have many direct and indirect connections between species, and these connections are not static but often vary over time. These topological changes, as opposed to a dynamic on a static (frozen) network, can be triggered by natural forcings (e.g., seasons) and/or by human influences (e.g., nutrient or pollution inputs). Aquatic trophic networks are especially dynamic and versatile, thus suggesting new approaches for identifying network structures and functioning in a comprehensive manner.

In this study, a qualitative model was devised for this purpose. Applying discrete-event models from theoretical computer science, a mechanistic and qualitative model was developed that allowed computation of the exhaustive dynamics of a given trophic network and its environment. Once the model definition is assumed, it provides all possible trajectories of the network from a chosen initial state. In a rigorous and analytical approach, for the first time, we validated the model on one theoretical and two observed trajectories recorded at freshwater stations in the La Rochelle region (Western France). The model appears to be easy to build and intuitive, and it provides additional relevant trajectories to the expert community. We hope this formal approach will open a new avenue in identifying and predicting trophic (and non-trophic) ecological networks.

Keywords: Interaction network; Freshwater ecosystem; Qualitative model; Discrete-event model, Plankton

INTRODUCTION

Trophic networks (TNs) form the backbone of ecosystem functioning, as they simultaneously condition food acquisition, prey and predator demography, individual and population behaviors, and phenotype selection, among other consequences (Lindeman 1942, Johnson 2000, Majdi et al. 2018). Trophic processes are responsible for most matter and energy fluxes within ecosystems, but the fates and properties of ecosystems are hard to predict, mainly due to the lack of knowledge (Mouquet et al. 2015). Trophic processes have been extensively studied in ecology, although mostly considered as frozen in time, i.e., with a fixed (or static) topology and fixed signed interactions. This simplification may be due to a lack of long-term data as well as to equation-based models dedicated to flux and abundance variations on a frozen network (e.g., Thébault and Fontaine 2010, Kéfi et al. 2015). In this study, we provide an original framework to handle TNs with sharply changing structures and to model their possible dynamics.

To date, TNs have been difficult to understand and handle, in other words, hard to *diagnose* between contrasted functioning under changing environmental conditions. Any new functioning involves specific ecosystem components and interactions, thus justifying why we have focused on qualitative functioning *regimes* rather than quantitative component abundances and interaction intensities. In addition, TNs usually gather a large number of populations or species in an even larger number of trophic interactions. Hence, understanding trophic dynamics would require not only modeling a large and realistic number of components but also being able to calibrate the weights (coefficients) of each component and each interaction involved (Ings et al. 2009, Wallach et al. 2017, Majdi et al. 2018). For this reason, most trophic models to date have focused on wide categories of populations, with functional categories, such as carnivores, herbivores, and/or detritivores (e.g., Thébault and Fontaine 2010), and approximate their trophic parameters. Even powerful models aimed at bypassing such limitations, such as qualitative models based on differential equation systems, are limited in size (May 1973, Dambacher et al. 2003).

There is an even more pronounced limitation of trophic studies in ecology, as they mostly assume a frozen network of interaction (Thébault and Fontaine 2010, Kéfi et al. 2016). Not only is it harder to handle a network that changes in terms of topology (structure), but it is also not known how such a network may change over time and, thus, how to model it. Indeed, as soon as the study covers several generations of some of the species involved in the network, other species may invade and/or become extinct (Mooney and Hobbs 2001, Warren et al. 2005). Hence, these events greatly modify the network structure and, in turn, the system dynamics. Equation-based models are not well suited to handle dynamic systems *on* dynamic structures (sometimes called DS², Giavitto and Michel 2003), whereas certain tools developed in theoretical computer science are perfectly adapted to this task. In particular, discrete-event systems such as graph transformations or Petri nets are able to handle sharply changing networks by formalizing the way components and interactions can appear or disappear (König et al. 2018, Gaucherel and Pommereau 2019). While graph transformations directly add/remove some nodes and edges, Petri nets only mimic such addition/removal by marking the presence/absence of the handled nodes and edges with some tokens. In this study, we developed a Petri net to model any interaction network topological change, and we illustrate its use on a realistic planktonic TN.

Planktonic TN models are usually composed of a fixed number of functional nodes that gather groups of individuals sharing the same ecological function. Mass fluxes (usually in carbon or nitrogen) between nodes are predefined according to trophic interactions. In a context of an emerging biological oceanography discipline and considering the limited computing resources, the first planktonic TN (or food web) models simply consisted of mass fluxes between nutrients, phytoplankton, and zooplankton nodes (Steele 1958, 1974). These so-called NPZ models (NPZD, NPZDB, or even NPZF when detritus, bacteria, or fishes are comprised, respectively) coupled to observed or simulated physical conditions have demonstrated their predictive ability to capture bulk system properties (biomass and primary production) at both regional and global scales (Mitra et al. 2007, Kriest et al. 2010, Oke et al. 2013, Hernández-Carrasco et al. 2014, Turner et al. 2014, Kumar and Kumari 2015).

To better understand biogeochemical cycling (e.g., export fluxes, carbon sequestration, organic matter recycling, microbial loop), planktonic TN should be delineated, and planktonic compartments in models could thus be refined into Planktonic Functional Types (PFTs). Plankton groups are thus defined according to

common ecological functions (e.g., nitrogen fixers, calcifiers, and silicifiers), sizes (e.g., picophytoplankton,
nanophytoplankton, microphytoplankton), and/or key taxonomic groups (e.g., diatoms, flagellates) (Le Fouest
et al. 2013, Villaescusa et al. 2016, Kerimoglu et al. 2017, Petersen et al. 2017, Maar et al. 2018, Meddeb et al.
2019). However, refinements of planktonic TN models greatly complicate model formalization and
parametrization, as well as requiring more data, which increases uncertainties in terms of the model outcomes
and fluxes between defined groups (Anderson 2005).

To address gaps in ecological knowledge and lacking data, inverse modeling is aimed at deriving flows of
energy within TNs from simple biomass estimates and rate measurements. Vézina and Platt (1988) were the
first to use this for inferring mass fluxes through a planktonic TN in the English Channel. Inverse modeling is,
therefore, advantageous when dealing with underdetermined systems and results in a space of possible
solutions that fulfill a set of linear equalities and inequalities. A preferred solution is then selected by
optimization or statistical methods. While vital rates and biomass can be readily measured for high trophic
levels (e.g., fishes), their quantification for low trophic levels (e.g., bacteria, autotrophic plankton) remains
uncertain and questions the robustness of inverse modeling for the study of planktonic compartments (Vernet
et al. 2017, Saint-Béat et al. 2018). Overall, biological constants (production, consumption, assimilation),
biomass, and ecological interactions are, therefore, not easy to measure in planktonic TNs, resulting in an
oversimplification of planktonic TN models (Anderson 2005, Flynn 2006). For all of these reasons, our main
objective in this study consisted of developing a model able to identify (diagnose) any qualitative functioning
regimes of the same TN under changing environmental conditions.

We here addressed the leading question: what are all the possible trajectories (pathways) of such an
aquatic TN? A trajectory is defined here as a sequence of TN states (regimes) and transitions in time, possibly
exhibiting bifurcations and not necessarily being quantitative. More specifically, we aimed to identify the
various qualitative regimes the system can reach between winter and summer environmental conditions. As a
second subquestion, we assessed whether a detailed model could exhibit new or counter-intuitive TN
trajectories. We assumed that the system may be qualified and thus exhibit a finite number of states,
computed and gathered into a Petri net *state space* (Pommereau 2010, Reisig 2013). A detailed and automatic
analysis of this state space then exhaustively provides the possible fates (e.g., stabilities, collapses, if any) of
the studied system. We chose to illustrate this original method with a well-studied plankton TN in wetlands,
namely freshwater marshes of the Charente Maritime region (Western France, Tortajada et al. 2011). Such a
system is well instrumented (measured) and will provide an expected theoretical trajectory of changing TN, as
well as two observed trajectories at different stations (Masclaux et al. 2014). The succession of planktonic TN
and the different regimes of the planktonic TN are well known according to the season (Masclaux et al. 2014).
We developed the corresponding Petri net of this system and then validated it, for the first time, on theoretical
and observed trajectories. We finally discuss the power and drawbacks of such discrete and qualitative models
for trophic ecology.

MATERIALS AND METHODS

Aquatic trophic networks
The Charente-Maritime marshes of the French Atlantic coast (Fig. 1) are the second largest French wetland
zone (more than 1000 km²). The type of freshwater marshes is unreplenished drained marshes, which
constitute a significant artificial hydrographic network of channels and ditches. To mitigate and prevent drying
of the marshes, locks control the channels and ditches (for more details, see Masclaux et al. 2014). Samples of
the planktonic TN were recorded at two stations (stations A and B) on successive dates (eight weeks during
winter and spring 2012) to reconstruct the network trajectories and their environment over time (Masclaux et
al. 2014). All of the details have been presented in the publication by Masclaux et al. (2014). Briefly, the water
parameters studied were the temperature, nutrients (nitrates, etc.), the dissolved organic matter (*DOC*)
concentration, the biomass and production of bacteria, the biomass and primary production of phytoplankton
by size class (microphytoplankton for > 20 μm; nanophytoplankton for 3–20 μm; and picophytoplankton for <
3 μm), the bacterial biomass, the protozoa biomass, and the metazoan microzooplankton and
mesozooplankton biomasses. Different fluxes between preys and predators were measured: micro- and
mesozooplankton grazing rates on the three phytoplankton size classes, as well as mesozooplankton grazing

rates on protozoa (Masclaux et al. (2014). The TN regimes were determined with a hierarchical ascendancy classification (HAC, Euclidean distance, and Ward method). The planktonic TN regime changed during the winter to spring transition, from biological winters, followed by herbivorous TNs, to finally reach TNs qualified as multivorous and distinguishing three levels of multivory (weakly multivorous, multivorous, and highly multivorous) (Masclaux et al. 2014).

Figure 1

Figure 1 - Location of the study site and the two sampled stations (A and B, inset) along the Atlantic coast of France.

The model was built with several categories of variables (Fig. 2): Phytoplankton, Zooplankton, Resources, Microbes, and Abiotic components characterizing the environment. The main functional groups were: 1) phytoplankton by size class (microphytoplankton for > 20 μm: *MicroP*; nanophytoplankton for 3–20 μm: *NanoP;* and picophytoplankton for < 3 μm: *PicoP*) all in green (Fig. 2); 2) metazoan microzooplankton (*MicroZ*) and mesozooplankton (*MesoZ*) in red; 3) resources as nitrates (*Nit*) and Dissolved Organic Matter (*DOC*) in brown; 4) microbes such as bacteria (*Bact*) and protozoa (*Proto*) in blue; and 5) abiotic variables in grey with the component *envir*, which corresponds to the temperature and light conditions and *renew* which corresponds to the possible renewal of water (i.e., flush) in the marshes depending on the rainfall and water usages (agriculture, breeding, etc.). In more detail, the planktonic TN and all the possible fluxes (interactions) between components concern grazing fluxes with some preferential predation, and potential competitions between organisms (Fig. 2). Protozoa graze on bacteria, *PicoP* and *NanoP*, and are grazed by *MicroZ* and *MesoZ*. *MicroZ* graze on bacteria, *PicoP*, *NanoP*, and *Proto*. *MicroP* is grazed mainly by *MesoZ*, which used *MicroZ*, *NanoP*, and *Proto* as preys. Two preferential interactions force the model: i) competition between bacteria and *PicoP*, suggesting that each may survive and be detrimental to the second one, and ii) preferential grazing of *MicroP* by *MesoZ* and preferential grazing of *NanoP* by *MicroZ*. In brief, plain upward edges correspond to prey-predator interactions, while dashed downward edges are the resulting detritus (toward the *DOC* variable).

Figure 2

Figure 2 - The detailed interaction network used in this study. The trophic and non-trophic components
are displayed as nodes with various colors (Phytoplankton: green, Zooplankton: red, Resources: brown,
Microbes: blue, and Abiotic components characterizing the environment: grey). Predation interactions
are displayed as plain (upward for trophic) and dashed (downward for degradation) edges, with preferred
prey populations highlighted with red doted-dashed edges.

Petri Nets and a simplistic Prey-Predator model

We first summarize the successive steps required to build any ecosystem Petri net and we then illustrate
these steps with a toy model. Our generic approach to model ecosystems has been called the EDEN (Ecological
Discrete-Event Network) modeling framework and is specifically applied here to an aquatic trophic TN. Any
ecosystem Petri net is developed in three successive steps (Fig. 3): i) an intuitive graph (i.e., a set of nodes and
edges) is built to represent the studied ecosystem with its components and their related processes, focusing
on the leading question addressed by the model (Fig. 3a); ii) this ecosystemic graph, now called the *interaction
network*, is then transformed into a formal model based on a discrete-event Petri net and its associated rules
(as explained in the next paragraph, Fig. 3b); iii) the Petri net is run (computed, Fig. 3c) and analyzed (Fig. 3d)
to determine the entire range of the ecosystem dynamics. However, the Petri net (steps i and ii) is hidden from
the (ecologist) user and is automatically built (in Python language, see Suppl. Mat.) once the ecosystem
components and processes have been chosen by the ecologists. Additional technical details regarding the
principle and use of Petri nets, in particular the way they are computed, can be found in the literature
(Pommereau 2010, Reisig 2013, Gaucherel and Pommereau 2019).

We here illustrate the basic functioning of the model using a simplistic prey-predator system (Fig. 3). Any
ecosystem network can be represented as a directed graph (with parallel edges). In this graph, every material
component of the ecosystem (e.g., abiotic: temperature; biotic: a population; anthropogenic: nitrate inputs)
is represented by a node (or variable), with two Boolean states: "present" (the component is functionally
present in the system, or above a chosen threshold, also denoted "+") and "absent" (functionally absent from

the system, or below the same threshold, denoted "-"). In the prey-predator toy model, only two nodes are defined: the prey and the predator populations. Any state of the ecosystem is defined by the set of "+" and "-" nodes (Fig. 3b), and "±" in the tables of this paper, when they may oscillate between successive states. The state of a node depends on the nodes to which it is connected, while a connection between nodes is assumed as soon as one process explicitly connects the different components (Figs. 3a-b). The rules correspond to any physicochemical and/or bio-ecological processes (e.g., if the prey population is below a chosen threshold (-), the predator population also ends up below its associated threshold), and thus represent all possible interactions between the components comprising the studied ecosystem. In the prey-predator system, only two rules are defined: R1, the predation itself: the predator eats the prey, and R2, the mortality: without prey, the predator dies (Figs. 3a and c). In the Petri net language, nodes are called places, and rules are called transitions, both being connected through oriented arcs (Fig. 3b).

Figure 3

Figure 3 - Illustration of a simplistic prey-predator system (a), with its associated Petri net (b), its qualitative dynamics (c), and the computed marking graph, also called state space (d). The system comprises two components, the prey (N) and predator (P) populations, and two interactions connecting them (rules R1 and R2, (a)). The corresponding Petri net comprises four places (P+, P-, N+, and N-) and two transitions, R1 and R2, linked by unlabeled and unweighted arcs (b). Starting with the presence of both populations, it is possible to list all system states encountered {d_1, d_2, and d_3} (c), and to connect them with the rules (absent nodes and inactivated rules are displayed in grey). The net is depicted in the initial state (c), and the successive states can be deduced from the token (black dot in (a)) movements between places (b). The marking graph of the Petri net (d) is depicted with each state number {d_0, d_1, and d_2} referring to the dynamics described above (b). It should be noted that a specific state of the system {d_4} may not be reached from this initial condition and with these rules (d).

Discrete and qualitative dynamics

Any rule of such discrete-event models combines the left-hand *condition* and right-hand *realization* as: "transition's name: condition >> realization". For a rule to be applied, the state of the node (variable) must

satisfy its application condition; the rule is then designated as *enabled*. If so, the application of the rule modifies the state of the node as stipulated in its realization part; the rule is then *fired* (i.e., executed or applied). In the prey-predator system (Fig. 3), the rules are written as R1: P+, N+ >> N- and R2: N-, and P+ >> P-. Since the rules modify node states, they change the overall system state (i.e., the state of the system aggregates all node states). Therefore, the entire system shifts from one state to another through the successive applications of the enabled rules (Fig. 3c). Computation of the defined Petri net produces the *state space*, which provides the set of all system states reachable by the rules defined (Fig. 3d). As a corollary, the system states are also connected to each other by some of these rules in the state space. The size of this state space is usually much smaller than the number of possible system states (2^n, where n is the number of components or nodes/variables), because the computation starts from a specific initial condition and because the rules have specific application conditions. Following the computer science community, we developed certain tools to automatically divide large state spaces into *merged* (simplified) state spaces, as explained in the next subsection.

Firing a rule independently of some others often leads to unrealistic trajectories (e.g., flushing water without removing the plankton in it). Therefore, we defined *constraints*, which prevent the model from simulating such unrealistic trajectories. Constraints have a condition and a realization part, just as rules *stricto sensu* do, and the model inevitable (mandatory) transitions given the system state. The sole difference between rules and constraints is that constraints have priority over stricto sensu rules. In the prey-predator system, the system state d_1 = {N-, P+} is unrealistic; so, the rule R2 has to be transformed into a constraint C1: N-, P+ >> P-. From a given state, the model first simulates all trajectories opened up by the defined constraints, and then, when all the system states obtained are realistic (i.e., there is no longer any enabled constraint), only the enabled rules are fired (Fig. 3d). It should be noted that the modeled system can remain an indefinite time in any of the computed states, as no rule forces it to leave the qualitative state (i.e., the system can experience quantitative dynamics, yet without sharp qualitative changes). In brief, the discrete model proposed here is qualitative, mechanistic (the processes are explicit), non-deterministic (no stochasticity yet several possible outcomes from each state), and asynchronous (i.e., all rules are applied as soon as possible, no rule conflict) (Reisig 2013, Gaucherel and Pommereau 2019). The EDEN models are also causal and chronological yet non-temporized, i.e., transitions and time steps are not quantified (and not probabilized).

TN trajectories and validation methodology

The theoretical plankton TN modeled here combines nine different components, associated with the dominant functional groups that may be present in the channel freshwater marshes, and two additional components featuring environmental conditions (Table 1, Fig. 2). To connect them, we defined 34 processes and seven constraints encompassing at least four trophic levels (Tables 2-3, Fig. 2). To validate the TN model, one theoretical trajectory was defined and two observed trajectories were recorded at two distant stations (Supplementary Materials, Tables SM1-2). For the model to be validated, we expect not only to detect these successive states (e.g., {S0, S1, S2, S3}) in the modeled state space but also to detect them in the correct succession order. To determine whether the model was able to recover the expected trophic regimes, we tested two variants of the model: i) the full model intending to encompass the TN functioning, and ii) a similar model (called *seasonal*) yet without a return to winter conditions (R0, Table 2), thus resulting in the model being stuck in summer conditions. The model starts in winter conditions or with a flush in summer conditions, with only the node *Renew* present, which returns a source of inorganic nitrogen to the system (Table 1).

For the full and seasonal models, we computed the state space and the *merged* state space, and we checked whether the observed trajectories were correctly recovered. A merged state space is a simplified state space gathering the sets of mutually reachable states of the modeled system, a topological structure called a *structural stability*, into the same nodes. This type of structure is interpreted as a stable regime as, by definition, any state in this stability can be reached by any other state belonging to it. Other stabilities can be identified automatically, such as *terminal stabilities*, from which the system can no longer exit, *basins* gathering states that have the same following states, and *deadlocks*, which are single states from which the system can no longer exit. Merged state spaces are much more compact than full state spaces, and summarized trajectories are readily revealed.

263 **Table 1** - The plankton TN components and their associated modeled variables, with their abbreviations
264 and descriptions. Whether these ecological components are present (+) or absent (-) in the initial state is
265 also indicated (second column).

Acronym	In initial state	Description
Bact	-	Bacteria
PicoP	-	Picophytoplankton
NanoP	-	Nanophytoplankton
MicrP	-	Microphytoplankton
Proto	-	Protozoa
MicrZ	-	Microzooplankton (metazoa)
MesoZ	-	Mesozooplankton
DOC	-	Dissolved organic matter
Nit	-	Nitrates, linked to seasons or flush
Envir	-	Environmental and climate conditions
Renew	+	Water and Nitrate inputs (rain or humans)

267 **Table 2** - Trajectories of the theoretical expert elicitation and observed at stations A and B. For each
268 trajectory, theoretical and observed regimes are listed in columns, and present (+)/absent (-) components
269 of the trophic network are listed in rows. The corresponding regimes displayed in Figs. 5a-c are listed in
270 the last row of each trajectory, with a single index A1 to A3 and B1 to B4 for successive regimes.

THEORY	Stage 1	Stage 2	Stage 3	Stage 4	Stage 5	Stage 6
Bact	-	-	-	+	+	+
PicoP	-	-	-	+	+	+
NanoP	-	+	+	+	-	-
MicrP	-	+	+	-	-	-
Proto	-	-	-	+	+	-
MicrZ	-	-	+	+	+	-
MesoZ	-	-	+	+	-	-
Nit	+	+	+	-	-	-
DOC	-	-	+	+	+	+
Envir	-	+	+	+	+	+
Renew	+	+	-	-	-	-
Regimes (Fig. 5)	Biological winter	Low herbivorous TN	Herbivorous TN	Multivorous/Highly multivorous TN	Microbial TN	Microbial loop

271
272

RESULTS

State spaces and computed dynamics
The full model provides a state space comprising 765 states gathered into a single dynamic structure (a so-
called structural stability, Fig. SM1a). The seasonal model becomes stuck in a high number (12, plus two basins)
of successive structural stabilities of various sizes (Figs. 4 and SM1b). When oriented downward in time; i.e.,
following causality and chronology, the whole system inevitably converges toward a small terminal stability
(made up of four states) in which the system is in a biological winter (i.e., few living species, in green, Fig.
SM1b). Some of the stabilities that are reached exhibit a large number of states and may keep the system into
such specific stabilities for an indefinite time (in purple, Figs. 4 and SM1b). In brief, the seasonal model displays

the same state space as the full model, but with a possible return to the initial biological winter regime, thus
connecting the bottom states (in red, Fig. 4a) to the top states (in pink, Fig. 4a). This is why we observed a
single cycling stability in the full model state space (Fig. SM1a).

Table 3 - Rules and constraints used in both the full and seasonal models, with their formal expression
(first column) and descriptions (second column). Only the rule N°0 (in italics and bold) is discarded in the
seasonal version of the model.

CONSTRAINTS	Descriptions
DOC- >> Bact-	C1 Without organic matter (DOC), Bacteria disappear
Nit- >> NanoP-, MicrP-, DOC+	C2 Without Nitrates, all types of phytoplankton except Pico disappear (and produce organic matter)
Bact+, PicoP-, NanoP- >> Proto-, DOC+	C3 Without all its preys (but Bacteria), Protozoa disappear (and produce organic matter)
Bact-, Proto-, PicoP-, NanoP- >> MicrZ-, DOC+	C4 Without all its preys (but Bacteria), Microzooplankton disappear (and produces organic matter)
Proto-, MicrZ-, NanoP-, MicrP- >> MesoZ-, DOC+	C5 Without all its preys, Mesozooplankton disappear (and produces organic matter)
Envir- >> PicoP-, NanoP-, MicrP-	C6 In winter, all Phytoplankton disappear (and do not produce organic matter)
PicoP+, NanoP+, MicrP+ >> Nit-	C7 With all Phytoplankton (present), Nitrates disappear

RULES	
Envir+ >> Envir-, Renew+	***R0 After summer (and spring) comes winter***
Envir- >> Envir+	R1 After winter comes summer (and spring), with a bloom
Envir+, Renew+ >> Nit+, Renew-	R2 Nitrate input is due to water flush from human management or rainfall
Envir-, Renew+ >> Nit+, DOC-, Bact-, Proto-, PicoP-, NanoP-, MicrP-, MicrZ-, MesoZ-	R3 In winter, the Reset of the system is due to water flush from rainfall (but there is a Nitrate input)
Envir-, Renew+ >> Renew-	R4 In winter, we stop the flush after its action (reset)
DOC+ >> Bact+	R5 Bacteria use organic matter, without removing it
Bact+, DOC+ >> DOC-	R6 Bacteria use organic matter and remove it
Envir+, Nit+ >> MicrP+, NanoP+	R7 Microphytoplankton and Nanophytoplankton use Nitrates in summer, without removing them
Envir+, Nit+ >> PicoP+	R8 Picophytoplankton use Nitrates, without removing them
Bact+ >> Proto+	R9 Protozoa graze on Bacteria, without removing them
Bact+, Proto+ >> Bact-, DOC+	R10 Protozoa graze on Bacteria, which disappear and produce organic matter
PicoP+ >> Proto+	R11 Protozoa graze on Picophytoplankton, without removing it
PicoP+, Proto+ >> PicoP-, DOC+	R12 Protozoa graze on Picophytoplankton, which disappears and produce organic matter
NanoP+ >> Proto+	R13 Protozoa graze on Nanophytoplankton, without removing it
NanoP+, Proto+ >> NanoP-, DOC+	R14 Protozoa graze on Nanophytoplankton, which disappears and produce organic matter
MicrP+ >> MesoZ+	R15 Mesozooplankton graze on Microphytoplankton (its PREFERED prey), without removing it

MicrP+, MesoZ+ >> MicrP-, DOC+	R16 Mesozooplankton graze on Microphytoplankton, which disappears and produces organic matter
MicrP-, NanoP+ >> MesoZ+	R17 Mesozooplankton graze on Nanophytoplankton, without removing it
MicrP-, NanoP+, MesoZ+ >> NanoP-, DOC+	R18 Mesozooplankton graze on Nanophytoplankton (secondary preferential prey), which disappears and produces organic matter
MicrP-, Proto+ >> MesoZ+	R19 Mesozooplankton graze on Protozoa, without removing them
MicrP-, Proto+, MesoZ+ >> Proto-, DOC+	R20 Mesozooplankton graze on Protozoa, which disappears and produces organic matter
NanoP+ >> MicrZ+	R21 Microzooplankton graze on Nanophytoplankton (its PREFERED prey), without removing it
NanoP+, MicrZ+ >> NanoP-, DOC+	R22 Microzooplankton graze on Nanophytoplankton, which disappear and produce organic matter
NanoP-, PicoP+ >> MicrZ+	R23 Microzooplankton graze on Picophytoplankton (secondary preferential prey), without removing it
NanoP-, PicoP+, MicrZ+ >> PicoP-, DOC+	R24 Microzooplankton graze on Picophytoplankton, which disappear and produce organic matter
NanoP-, Proto+ >> MicrZ+	R25 Microzooplankton graze on Protozoa, without removing them
NanoP-, Proto+, MicrZ+ >> Proto-, DOC+	R26 Microzooplankton graze on Protozoa, which disappear and produce organic matter
NanoP-, Bact+ >> MicrZ+	R27 Microzooplankton graze on Bacteria, without removing them
NanoP-, Bact+, MicrZ+ >> Bact-, DOC+	R28 Microzooplankton graze on Bacteria, which disappear and produce organic matter
MicrZ+ >> MesoZ+	R29 Mesozooplankton graze on Microzooplankton, without removing it
MicrZ+, MesoZ+ >> MicrZ-, DOC+	R30 Mesozooplankton graze on Microzooplankton, which disappears and produces organic matter

Four TN regimes were revealed by the full and seasonal models (Fig. 4a and Table SM1): the S0 biological
winter regime, without planktonic bloom, with oscillations of both zooplankton feeding on bacteria and
organic matter; the S1 regime, in which all components are present because the environment is favorable to
the development of organisms with many fluctuations of them; the S2 multivorous regime, with a mix of preys
and various predators such as protozoa and both zooplankton (MicrZ and MesoZ), and finally, the S3 regime,
centered on protozoa and mesozooplankton with a few preys but oscillation of Bacteria (Fig. 4a and Table
SM1). The trajectories passing through different regimes were diverse and depended on the rules fired from
the initial state (Fig. 4b): either the TN system shifts between various types of S0 regimes, or it successively
crosses regimes S0 to S1, S2, and S3 (with possible ways back) and then back to S0.

Figure 4

Figure 4 - The merged state space of the seasonal model (a, as in Fig. SM1a), in which each node corresponds to a structural stability (i.e., a set of mutually reachable states), and each edge corresponds to irreversible transitions between successive stabilities. Here, structural stabilities are labeled with system components that are systematically present (+) in their associated states (see Fig. SM2). To see components that are systematically absent in stabilities). This figure helps identify the various regimes (b, and Table SM1) reached by the TN system along to the (downward) trajectories computed.

Model validation

305
All states of the theoretical trajectory were correctly predicted by the model and, as the model exhibited
a single stability, the system is certain to successively reach all these states (although we do not know by which
complicated trajectories, Fig. 4a, Table SM1). This observation definitely validates the model. The theoretical
planktonic TN trajectory (Table 2 and Fig. 5a) started by an (immature) regime found during the biological
winter. It then evolved toward low herbivorous TN, herbivorous TN, and variable multivorous TN (from weak
to highly multivorous regimes, with protozoa, microzooplankton, or mesozooplankton, respectively), and a
microbial TN regime, to finally reach a microbial loop regime. When the water in the marshes was renewed
(*Renew*+), the TN returned to the biological winter regime, as can the herbivorous regimes as well. All these
TN regimes were recovered by the model (Fig. 5a), yet with slightly different stabilities. Figure 5 is not meant
to demonstrate this validation stage (already achieved by identification of the successive trajectory states),
rather than *showing* that all predicted states (colored nodes) were correctly recovered in the computed state
space, and indeed connected through successive transitions (bold edges). Note that this modeled trajectory
crossed many other intermediate states (Fig. 5a) not found in the observations of (Masclaux et al. 2014). In
the seasonal model (rule R0 deactivated), the theoretical trajectory was also predicted (colored states, Fig. 5a),
yet with the last two states (blue states at the bottom) in the reverse order, as the fourth multivorous regime
can directly reach the last depleted one.

322
323   The data recorders at stations A and B, and associated with observed trajectories, were also correctly
324 recovered. At station A, three regimes succeeded over time, from A1 to A3 (Table SM2, and Fig. 5b). The TN

started with biological winter for four weeks with the presence of nitrates and organic matter, but it did not
reach favorable conditions for biological development. Then, the favorable conditions at week five allowed the
development of phytoplankton (micro- and pico-plankton), and thus their zooplankton predators with
bacteria. This situation was typical of situations between herbivorous and multivorous TNs. A multivorous
regime of TN then took hold until week eight. At station B, the observed succession displayed four regimes,
from B1 to B4 (Table SM2 and Fig. 5c). The TN started with biological winter for three weeks and favorable
conditions occurring at week four, which allowed the presence of multivorous TN ("weak multivorous TN"
according to (Masclaux et al. 2014)) and at week five an herbivorous TN. Then, a multivorous regime of TN
took hold from week six to week eight.

Figure 5

Figure 5 - The full (not merged) seasonal state space highlighting the three trajectories used in this study
(Table 2), namely the theoretical (a), station A (b), and station B (c) ones (Table 2 Suppl.). Here, each node
corresponds to a TN state, connected to each other by downward transitions. The specific states
underlying the three trajectories are highlighted by node colors other than red and identifiers
corresponding to their numbers (last rows of trajectories in Table 2) and by bold edges.

DISCUSSION

The discrete-event and qualitative model of trophic networks (TN) presented here can be computed
instantaneously (< 0.01 s) and provided, once the model was defined and assumed, all possible trajectories
of this system (Fig. 4). To our knowledge, this is the first attempt to exhaustively model a detailed TN (11
components, Table 1) and to accurately validate its qualitative dynamics.

Complex dynamics of aquatic trophic networks
In the Charente-Maritime trophic system, we discovered that this TN may have followed other
trajectories than the one identified by experts in the theoretical model and in the ones observed (Fig. 5
and Supplementary Materials tables). First, station B showed that *DOC* may be present in winter, thus with
the TN fluctuating in intermediary states before reaching the usual trajectory observed in Masclaux et al.
(2014). Indeed, *DOC* in winter could be an allochthonous input from the terrestrial environment (Del Gorgio
and Davis 2003). After winter (i.e., when *Renew+* and *Envir+* were present, Table 1), all the modeled
trajectories and all the TN regimes appeared at reach. The TN can return to biological winter system states
due to the nitrate inputs (*Nit+*, with R2) and to anthropogenic activities (Tortajada et al. 2011). This
situation occurs when the water renewal is substantial and no planktonic biomass accumulation is possible
(David et al. 2020). Moreover, rainfall could occur and favor nitrate leaching (R3), thus pushing back the
planktonic TN to biological winter system states. The model confirmed the key role of organic matter (*DOC*),
as the system trajectories differed depending on whether or not organic matter was present at the
beginning of winter.
359

From the initial state of the TN, the trajectories could pass through slightly different biological winter
regimes (Figs. 4a-b) with oscillations in organic matter, bacteria, and micro- and mesozooplankton.
Similarly, Masclaux et al. (2014) found two types of Biological winter regimes, mainly depending on the
presence or absence of bacteria, and on some prey and predator combinations. The model correctly
recovered different states of biological winters. The regime of multivorous TN was also well recovered by
the model (Masclaux et al. 2014). The multivorous TN is known to be highly stable (Legendre and
Rassoulzadegan 1995). However, the microbial loop, which has a transient nature (Legendre and
Rassoulzadegan 1995) did not appear as a structural stability in the model either.
368

The regime gathering protozoa and mesozooplankton (*Proto/MesoZ* cycles) characterized by the
presence of predators with a few preys but oscillation of bacteria was not found in the observations
(Masclaux et al. 2014). The modeled trajectory crossed many intermediate states (Figs. 5a-c) not sampled
in the field. The field sampling frequency or the structural characteristics of the sampled wetlands likely
did not allow capture of all the possible states of TN: this reveals the ability of the model to explore many
other possible states of the planktonic TNs and other trajectories of TN. In particular, the predicted
Proto/MesoZ regime has not yet been identified at the Charente Maritime sites, but work in progress at
other Atlantic arc territories has identified related TNs (F.-X. Robin, pers. comm.). Finally, bacteria were
frequently present in the ecosystem, and they occupied a prominent place in the model (Table SM1–2,
Table 2). Bacteria appeared to oscillate frequently (Fig. 4a), although this was not visible in the merged
state space (i.e., bacteria frequently appear and disappear within structural stabilities). The model
confirmed that bacteria are frequently grazed by their grazers, as are small protists (Pernthaler 2005, Šimek
et al. 2013). Indeed, the high level of control of bacteria by the protozoa in freshwater ecosystems is already
known.

Power and drawbacks of discrete-event models
An increasing number of TN models have been developed (Mitra et al. 2007, Kriest et al. 2010, Thébault
and Fontaine 2010, Turner et al. 2014, Kéfi et al. 2016, Hansen and Visser 2016, Kloosterman et al. 2016).
But they still suffer from three main limitations: limited size and complexity, and a frozen (static) network
with frozen (i.e., topology) interactions. In this study, we proposed a novel model family (called the EDEN
framework (Gaucherel and Pommereau 2019, Cosme et al. 2022)) to bypass these limitations. Our model
is based on a discrete-event system, well-known to computer scientists and more recently also some

molecular biologists (Thomas and Kaufman 2001, Reisig 2013). The price to pay for using our qualitative model is that no quantitative and detailed dynamics are available; but in turn, no difficult parameterization and construction are required. Consequently, such an approach is fully complementary to already existing models in (trophic) ecology. Here, to the best of our knowledge, we provide for the first time a discrete and qualitative model of TN to bypass such limitations. Of note, in continuity with previous theoretical attempts (May 1973, Dambacher et al. 2003), we here open a new avenue for the use of such novel qualitative models in (ecosystem and trophic) ecology. The foundations of this proposition, yet beyond the present scope, are based on a theoretical consideration that assumes that ecosystems are informational systems comprised of material components and immaterial processes (Gaucherel 2019) summarized into their interaction networks.

Such a model is intuitive, easy to build, tractable, and rigorous (i.e., no trajectories have been forgotten or added according to the mathematical Petri net engine). In addition, we claim that it does not require any detailed or quantitative calibration, as no parameter is required. The central assumption of this approach is that it is possible to summarize ecological processes into qualitative rules, possibly interpreted as long-term and discrete events. Other studies have shown that this approach is not limited to trophic processes and can be applied to a high diversity of social-ecosystems (Gaucherel and Pommereau 2019, Mao et al. 2021). In this study, we were fortunate enough to collate several theoretical and observed trajectories with which to validate the model, thus confirming that it is conform and accurate (Fig. 5). Another quality of this type of model is to be heuristic, to force scientists questioning the knowledge they have regarding the studied system and to collate it into a single coherent framework.

As perspectives, it appears suitable to model many TN stressors such as pollution, cleaning, drought, invasive species, and/or climate changes (Mooney and Hobbs 2001, Mouquet et al. 2015). Any complexification of the studied social-ecosystem is also possible, in theory, as the model is still far from reaching its limits in terms of components, processes, and their nature diversity. It may then be used in a more applied manner, for exploration of other scenarios by changing the initial conditions. Coupling this model with other components describing the mechanisms behind these stressors would provide a relevant territorialized model to anticipate trends in a context of global warming and coastline changes. In the near future, it would be relevant not only to improve the model's realism but also to develop analysis tools already used in similar studies focusing on social-ecological systems (Mao et al. 2021, Cosme et al. 2022). Additionally, it would be relevant to complexify our discrete and qualitative approach by using quantitative and multivalued schemes, to bridge the gap with more traditional (e.g., equation-based or individual-based) models (Vézina and Platt 1988, Kéfi et al. 2016).

In brief, by modeling trophic networks with a novel (EDEN) framework, we recovered theoretical as well as observed trajectories. With such qualitative models, the dynamics and predicted new states and new trophic network functioning regimes that may be observed in the field can be better understood. We illustrated these with a specific and well-documented freshwater trophic network. Such models provide an intuitive and robust approach to diagnosing any trophic (and non-trophic) network by computing all possible trajectories it can reach from a given chosen initial state. The known processes at play in the system help identify all of the possible dynamics and thus counter-intuitive trajectories of such complex (social-eco-)systems. Connecting such biotic dynamics to human-related activities can be expected to provide additional insightful understanding of trophic systems.

APPENDICES

Additional Tables and Figures (Appendix 1)

ACKNOWLEDGMENTS

We thank François-Xavier Robin for his relevant comments on an earlier draft of this paper.

DATA SCRIPTS CODE AND SUPPLEMENTARY MATERIAL AVAILABILITY

Data are available in this article (Tables and Figures) (*citation of the data* Gaucherel et al, 2023);

Scripts and code are available online: DOI:10.1111/2041-210X.13242 of the webpage hosting the data https://github.com/fpom/ecco (*citation of the scripts and code* Pommereau et al., 2022);

Supplementary information is available online: XXXXDOI of the webpage hosting the data https://doi.org/10.5802/fake3.doi (*citation of the scripts and code* Gaucherel et al, 2023);

[The references of the datasets, scripts and codes should also be present in the reference list and cited in the text.]

CONFLICT OF INTEREST DISCLOSURE

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article. *[C. Gaucherel is a recommender PCI ecology]*

FUNDING

We also thank the INRA ECOSERV Meta-program for his financial support on wetland studies and modeling (Project SERVICESCALES), now part of the INRAE ECODIV department.

REFERENCES

Anderson, T. R. 2005. Plankton functional type modelling: running before we can walk? Journal of plankton research 27:1073-1081.

Baldan, P., M. Bocci, D. Brigolin, N. Cocco, and M. Simeoni. 2015. Petri nets for modelling and analysing trophic networks.in BioPPN 2015, a satellite event of PETRI NETS 2015. CEUR Workshop Proceedings.

Cosme, M., C. Hély, F. Pommereau, P. Pasquariello, C. Tiberi, A. C. Treydte, and C. Gaucherel. 2022. Qualitative Modeling for Bridging Expert-Knowledge and Social-Ecological Dynamics of an East African Savanna. Land 11:https://doi.org/10.3390/land11010042.

Dambacher, J. M., H.-K. Luh, H. W. Li, and P. A. Rossignol. 2003. Qualitative Stability and Ambiguity in Model Ecosystems. The American Naturalist 161:876–888.

David, V., S. Tortajada, N. Savoye, M. Bréret, N. Lachaussée, O. Philippine, F. X. Robin, and C. Dupuy. 2020. Impact of human activities on the spatio-seasonal dynamics of plankton diversity in drained marshes and consequences on eutrophication. Water Research 170:115287.

Del Gorgio, P. A. and J. Davis. 2003. Patterns in Dissolved Organic Matter Lability and Consumption across Aquatic Ecosystems Aquatic Ecosystems Aquatic Ecology 2003:399-424.

Flynn, K. J. 2006. Reply to Horizons Article 'Plankton functional type modelling: running before we can walk'Anderson (2005): II. Putting trophic functionality into plankton functional types. Journal of plankton research 28:873-875.

Gaucherel, C. 2019. The Languages of Nature. When nature writes to itself. Lulu editions, Paris, France.

Gaucherel, C. and F. Pommereau. 2019. Using discrete systems to exhaustively characterize the dynamics of an integrated ecosystem. Methods in Ecology and Evolution 00:1–13.

Giavitto, J. L. and O. Michel. 2003. Modeling the topological organization of cellular processes. Biosystems 70:149-163.

Hansen, A. N. and A. W. Visser. 2016. Carbon export by vertically migrating zooplankton: an optimal behavior model. Limnology and Oceanography 61, 701–710.

Hernández-Carrasco, I., V. Rossi, E. Hernández-García, V. Garçon, and C. López. 2014. The reduction of plankton biomass induced by mesoscale stirring: A modeling study in the Benguela upwelling. Deep sea research part I. Oceanographic research papers 83:65-80.

Ings, T. C., J. M. Montoya, J. Bascompte, N. Blüthgen, L. Brown, C. F. Dormann, F. Edwards, D. Figueroa, U. Jacob, J. I. Jones, R. B. Lauridsen, M. E. Ledger, H. M. Lewis, J. M. Olesen, F. J. F. van Veen, P. H. Warren, and G. Woodward. 2009. Ecological networks – beyond food webs. Journal of Animal Ecology 78:253–269.

Johnson, K. H. 2000. Trophic-dynamic considerations in relating species diversity to ecosystem resilience.
Biological Reviews 75:347-376.
Kéfi, S., E. L. Berlow, E. A. Wieters, L. N. Joppa, S. A. Wood, U. Brose, and S. A. Navarrete. 2015. Network
structure beyond food webs: mapping non-trophic and trophic interactions on Chilean rocky shores.
Ecology 96:291-303.
Kéfi, S., V. Miele, E. Wieters, S. Navarrete, and E. Berlow. 2016. How Structured Is the Entangled Bank? The
Surprisingly Simple Organization of Multiplex Ecological Networks Leads to Increased Persistence and
Resilience. PLoS Biology 14:e1002527.
Kerimoglu, O., S. Jacquet, B. Vinçon-Leite, B. J. Lemaire, F. Rimet, F. Soulignac, and O. Anneville. 2017.
Modelling the plankton groups of the deep, peri-alpine Lake Bourget. Ecological Modelling 359:415-
433.
Kloosterman, M., S. A. Campbell and F. J. Poulin. 2016. An NPZ model with state-dependent delay due to
size-structure in juvenile zooplankton. Siam Journal of Applied Mathematics 76 (2), 551–577.
König, B., D. Nolte, J. Padberg, and A. Rensink. 2018. A Tutorial on Graph Transformation. Pages 83–104 in
R. Heckel and G. Taentzer, editors. Ehrig Festschrift Springer International Publishing AG, part of
Springer Nature.
Kriest, I., S. Khatiwala, and A. Oschlies. 2010. Towards an assessment of simple global marine
biogeochemical models of different complexity. Progress in Oceanography 86:337-360.
Kumar, V. and B. Kumari. 2015. Mathematical modelling of the seasonal variability of plankton and forage
fish in the Gulf of Kachchh. Ecological Modelling 313:237-250.
Le Fouest, V., B. Zakardjian, H. Xie, P. Raimbault, F. Joux, and M. Babin. 2013. Modeling plankton ecosystem
functioning and nitrogen fluxes in the oligotrophic waters of the Beaufort Sea, Arctic Ocean: a focus on
light-driven processes. ????
Legendre, L. and F. Rassoulzadegan. 1995. Plankton and nutrient dynamics in marine waters. Ophelia
41:153-172.
Lindeman, R. 1942. The trophic-dynamic aspect of ecology. ecology 23:199-418.
Maar, M., M. Butenschön, U. Daewel, A. Eggert, W. Fan, S. S. Hjøllo, and M. A. Peck. 2018. Responses of
summer phytoplankton biomass to changes in top-down forcing: Insights from comparative modelling.
Ecological Modelling 376:54-67.
Majdi, N., N. Hette-Tronquart, E. Auclair, A. Bec, T. Chouvelon, B. Cognie, M. Danger, P. Decottignies, A.
Dessier, C. Desvilettes, S. Dubois, C. Dupuy, C. Fritsch, C. Gaucherel, M. Hedde, F. Jabot, S. Lefebvre, M.
P. Marzloff, B. Pey, N. Peyrard, T. Powolny, R. Sabbadin, E. Thébault, and M.-E. Perga. 2018. There's no
harm in having too much: A comprehensive toolbox of methods in trophic ecology. Food webs
17:e00100.
Mao, Z., J. Centanni, F. Pommereau, A. Stokes, and C. Gaucherel. 2021. Maintaining biodiversity promotes
the multifunctionality of social-ecological systems: Holistic modelling of a mountain system. Ecosystem
services 47:101220.
Masclaux, H., S. Tortajada, O. Philippine, F.-X. Robin, and C. Dupuy. 2014. Planktonic food web structure
and dynamic in freshwater marshes after a lock closing in early spring. Aquatic Sciences:DOI
10.1007/s00027-00014-00376-00021.
May, R. M. 1973. Qualitative Stability in Model Ecosystems. Ecology 54:638-641.
Meddeb, M., N. Niquil, B. Grami, K. Mejri, M. Haraldsson, A. Chaalali, and A. S. Hlaili. 2019. A new type of
plankton food web functioning in coastal waters revealed by coupling Monte Carlo Markov chain linear
inverse method and ecological network analysis. Ecological Indicators 104:67-85.
Mitra, A., K. J. Flynn, and M. J. R. Fasham. 2007. Accounting for grazing dynamics in nitrogen-
phytoplankton-zooplankton (NPZ) models. Limnology and Oceanography 52.
Mooney, H. A. and R. J. Hobbs. 2001. Invasive Species in a Changing World. Island Press, Washington D.C.,
USA.
Mouquet, N., Y. Lagadeuc, V. Devictor, L. Doyen, A. Duputié, D. Eveillard, D. Faure, E. Garnier, O. Gimenez,
P. Huneman, F. Jabot, P. Jarne, D. Joly, R. Julliard, S. Kéfi, G. J. Kergoat, S. Lavorel, L. L. Gall, L. Meslin, S.
Morand, X. Morin, H. Morlon, G. Pinay, R. Pradel, F. M. Schurr, W. Thuiller, and M. Loreau. 2015.
Predictive ecology in a changing world. Journal of Applied Ecology 52:1293-1310.
Oke, P. R., D. A. Griffin, A. Schiller, R. J. Matear, R. Fiedler, J. Mansbridge, and K. Ridgway. 2013. Evaluation
of a near-global eddy-resolving ocean model. Geoscientific model development 6:591.

Pernthaler, J. 2005. Predation on prokaryotes in the water column and its ecological implications. Nat Rev
Microbiol 3:37-46.
Petersen, M. E., M. Maar, J. Larsen, E. F. Møller, and P. J. Hansen. 2017. Trophic cascades of bottom-up and
top-down forcing on nutrients and plankton in the Kattegat, evaluated by modelling. Journal of Marine
Systems 169:25-39.
Pommereau, F. 2010. Algebras of coloured Petri nets. Lambert Academic Publishing (LAP).
Reisig, W. 2013. Understanding Petri Nets. Springer Berlin Heidelberg, Berlin, Heidelberg.
Saint-Béat, B., F. Maps, and M. Babin. 2018. Unraveling the intricate dynamics of planktonic Arctic marine
food webs. A sensitivity analysis of a well-documented food web model. Progress in Oceanography
160:167-185.
Šimek, K., V. Kasalický, J. Jezbera, K. Horňák, J. Nedoma, M. W. Hahn, D. Bass, S. Jost, and J. Boenigk. 2013.
Differential freshwater flagellate community response to bacterial food quality with a focus on
Limnohabitans bacteria. ISME 7:1519–1530.
Steele, J. H. 1958. Plant production in the northern North Sea. Scottish Home Dept. Marine Res. 7:1- 36.
Steele, J. H. 1974. The Structure of Marine Ecosystems. Harvard University Press, Cambridge,
Massachusetts.
Thébault, E. and C. Fontaine. 2010. Stability of Ecological Communities and the Architecture of Mutualistic
and Trophic Networks. Science 329:853 -856.
Thomas, R. and M. Kaufman. 2001. Multistationarity, the basis of cell differentiation and memory. II. Logical
analysis of regulatory networks in terms of feedback circuits. Chaos 11:180-195.
Tortajada, S., V. David, A. Brahmia, C. Dupuy, T. Laniesse, B. Parinet, F. Pouget, F. Rousseau, B. Simon-
Bouhet, and F.-X. Robin. 2011. Variability of fresh- and salt-water marshes characteristics on the west
coast of France: A spatio-temporal assessment. Water Research 45:4152-4168.
Turner, E. L., D. A. Bruesewitz, R. F. Mooney, P. A. Montagna, J. W. McClelland, A. Sadovski, and E. J. Buskey.
2014. Comparing performance of five nutrient phytoplankton zooplankton (NPZ) models in coastal
lagoons. Ecological Modelling 277:13-26.
Vernet, M., T. L. Richardson, K. Metfies, E. M. Nöthig, and I. Peeken. 2017. Models of plankton community
changes during a warm water anomaly in arctic waters show altered trophic pathways with minimal
changes in carbon export. Frontiers in Marine Science 4:160.
Vézina, A. F. and T. Platt. 1988. Food web dynamics in the ocean. 1. Best-estimates of flow networks using
inverse methods. Marine ecology progress series. Oldendorf 42:269-287.
Villaescusa, J. A., S. E. Jørgensen, C. Rochera, D. Velazquez, A. Quesada, and A. Camacho. 2016. Carbon
dynamics modelization and biological community sensitivity to temperature in an oligotrophic
freshwater Antarctic lake. Ecological Modelling 319:21-30.
Wallach, A. D., A. H. Dekker, M. Lurgi, J. M. Montoya, D. A. Fordham, and E. G. Ritchie. 2017. Trophic
cascades in 3D: network analysis reveals how apex predators structure ecosystems. Methods in Ecology
and Evolution 8:135-142.
Warren, P. H., R. Law, and A. J. Weatherby. 2005. Invasion biology as a community process: messages from
microbial microcosms. Pages 353–377 in M. W. Cadotte, S. M. McMahon, and T. Fukami, editors.
Conceptual ecology and invasions biology. Springer, Printed in Great Britain.

579