Molecular quantitative trait loci in reproductive tissues impact male fertility

in a large mammal

Xena Marie Mapel[1,*], Naveen Kumar Kadri[1,*], Alexander S. Leonard[1], Qiongyu He[1], Audald Lloret-

Villas[1], Meenu Bhati[1,$], Maya Hiltpold[1,&], Hubert Pausch[1,#]

[1] Animal Genomics, ETH Zurich, Universitatstrasse 2, 8092 Zurich, Switzerland

[$] current address: Roslin Institute, The University of Edinburgh, Easter Bush Campus, Midlothian

EH25 9RG, United Kingdom

[&] current address: GenPhySE, Université de Toulouse, INRAE, ENVT, 31326, Castanet Tolosan, France

[*] equal contribution

[#] Corresponding author

Abstract

Cattle are well suited to investigate inherited variation in male fertility because insemination success

is recorded in thousands of artificial insemination bulls. However, functional data from relevant

tissues are lacking, which prevents fine-mapping fertility-associated genomic regions to reveal

functional links between genotype and phenotype. Here, we profile testis, epididymis, and vas

deferens transcriptomes in 118 mature bulls and conduct association tests between 417k molecular

phenotypes and 21M genome-wide variants to identify 41k regulatory loci. We show broad

consensus in tissue-specific and tissue-enriched gene expression between the three bovine tissues

and their human and murine counterparts. Expression- and splicing-mediating variants are more

than three times as frequent in testis than epididymis and vas deferens, highlighting the

transcriptional complexity of testis. We identify genes (*WDR19*, *SPATA16*, *KCTD19*, *ZDHHC1*) and

molecular phenotypes that are associated with male fertility through transcriptome-wide association

and colocalization analyses, revealing molecular-genetic underpinnings of quantitative variation in

male fertility.

Introduction

Male fertility varies considerably between individuals, but it remains difficult to identify underlying

genetic mechanisms (Tüttelmann et al., 2018; Xavier et al., 2021). Exploring the genetic basis of

reproductive success requires separating genetic from environmental effects and differentiating

between male and female factors that contribute to fertilization; this is only possible when repeated

phenotypic measurements are available for large cohorts. While such cohorts are unavailable for

nearly all species, including humans, the beef and dairy industries' reliance on artificial insemination

has generated comprehensive catalogs of semen quality and male fertility records for thousands of

genotyped artificial insemination bulls (Thibier & Wagner, 2002). Indeed, studies investigating bull

fertility have high translational value as they can elucidate evolutionarily conserved functional

mechanisms that contribute to undisturbed mammalian fertilization (Lamas-Toranzo et al., 2020;

Pausch et al., 2014; Tang et al., 2022), making cattle an unconventional—but ideally suited—species

to study the genetic architecture of male reproductive ability.

Bovine male reproductive traits have low-to-moderate heritability, suggesting they are amenable to

association testing with dense molecular markers (Butler et al., 2020; Gebreyesus et al., 2021; Nagai

et al., 2022; Pausch & Mapel, 2023). Datasets from artificial insemination bulls provide enough

statistical power to identify variants that contribute to inherited variation in male fertility through

genome-wide association studies (GWAS). Several thousand quantitative trait loci (QTL) for male

reproductive traits are listed in the AnimalQTL Database (https://www.animalgenome.org/cgi-

bin/QTLdb/index); many of these loci are in non-coding regions of the genome and segregate within

populations at relatively high frequencies (Han & Peñagaricano, 2016; Nicolini et al., 2018; Pacheco

et al., 2022; Rezende et al., 2018). Though numerous male fertility QTL have been identified in cattle,

a lack of functional data has caused the molecular underpinnings for most of these loci to remain

elusive (Druet et al., 2009; Hiltpold et al., 2021; Rezende et al., 2018).

Variants that regulate gene expression and splicing influence complex traits and diseases (Garrido-

Martín et al., 2021; Y. I. Li et al., 2016; Walker et al., 2019; Yamaguchi et al., 2022). Several

Genotype-Tissue Expression (GTEx) projects identified thousands of regulatory molecular

quantitative trait loci (molQTL; specifically expression and splicing QTL) throughout mammalian

genomes for dozens of tissues (GTEx Consortium, 2020; Liu et al., 2022; The FarmGTEx-PigGTEx

Consortium et al., 2022). Integration of molQTL and phenotypic records, or summary statistics from

GWAS, through transcriptome-wide association studies (TWAS) can identify regulatory variants that

are associated with complex trait variation (Barbeira et al., 2018; Wainberg et al., 2019; Wu et al.,

2019). However, since this requires uniform cohorts with detailed trait records and transcriptomics

data for functionally relevant tissues, TWAS for male fertility have not been feasible thus far.

Here, we establish a large and homogeneous cohort of bulls that contains both DNA sequencing and

transcriptome data from testis, epididymis, and vas deferens. We characterize the transcriptomic

profile of these three reproductive tissues and identify thousands of loci that influence gene

expression and splicing. In doing so, we discover an excess of regulatory small-effect variants in

testis that are often tissue specific. Lastly, we conduct transcriptome-wide association studies and

colocalization analyses to pinpoint molecular phenotypes that impact bovine male fertility.

Results

We sampled testis, epididymis (caput), and vas deferens tissue from 118 bulls of Braunvieh (BV)

ancestry (i.e., Brown Swiss (BS), Original BV (OB), or a cross between one of these and a different

breed; Figure 1A;1B;1C; Figure S1). All bulls were considered post-pubertal, although the age at

sampling varied from 10 to 37 months. Whole-genome sequencing (WGS) of DNA extracted from

testis tissue yielded between 65,207,590 and 461,834,248 cleaned paired-end reads per sample

(mean = 208,761,950 ± 63,260,972). The subsequent alignment of these reads against the *Bos taurus*

reference genome achieved an average coverage of 12.34 ± 3.71 -fold. Variant calling identified

29,660,795 polymorphic sites (single nucleotide polymorphisms (SNPs), insertions and deletions

(indels) smaller than 50 bp), of which 21,501,032 met our quality criteria for downstream analyses

(Table S1).

Total RNA from testis, epididymis, and vas deferens tissue was extracted from the same bulls and

deeply sequenced with stranded paired-end libraries. We used a splice-aware alignment tool to map

an average of 257,050,199, 283,746,666, and 262,089,072 reads per sample for testis, epididymis,

and vas deferens, respectively, to the bovine reference genome and the Ensembl gene annotation

(Table S2). An additional set of alignments that included WASP filtering (Van De Geijn et al., 2015) to

mitigate reference allele bias contained an average of 240,544,139 mapped reads in testis,

267,645,035 mapped reads in epididymis, and 251,031,216 mapped reads in vas deferens (Table S2).

Our final dataset contained 117, 103, and 84 samples for testis, epididymis, and vas deferens,

respectively, 74 of which had data of sufficient quality for all three tissues (Figure 1A; 1B).

Gene expression and splicing variation is pervasive in male reproductive tissues

We estimated gene expression in transcripts per million (TPM) and, after filtering, detected 21,844

expressed genes (≥ 0.1 TPM and ≥ 6 reads in ≥ 20% of samples) across the three tissues. When

considering the three tissues individually, we identified slightly more expressed genes in epididymis

(20,376 genes) than in testis and vas deferens (20,222 and 19,051 expressed genes, respectively;

Table 1). A majority of expressed genes were found in all three tissues (17,737 genes; Figure 1E) and

were protein coding (87.4% – 89.7%; Tables S3). Testis had the most tissue-specific expressed genes

(1,106 genes), followed by epididymis (443 genes), and vas deferens (227 genes; Figure 1E).

Figure 1. Overview of the molQTL cohort for three male reproductive tissues. A.) The three male

reproductive tissues used in our study and the number of samples considered for each tissue. B.) Sample

overlap across tissue types. Bar height represents the number of individuals. C.) PCA of LD-pruned sequence

variant genotypes (366,090 variants total), with colors corresponding to the breed assigned by the Swiss

Braunvieh herd book or "Cross". D.) Scatter plot of the top two principal components of PCAs of normalized

expression values (TPM; top panel) and normalized splicing phenotypes (PSI; bottom panel) for all tissues. E).

Overlap of expressed and spliced genes across the three tissues (based off the TPM and PSI matrices).

Table 1. Results for e/sQTL mapping across tissues with a MAF filter of ≥ 1% and FDR of 5%.

	Testis	Epididymis	Vas deferens
Samples	117	103	84
Expressed genes	20,222	20,376	19,063
eQTL	15,642	4,768	4,211
eGenes	11,164	4,347	3,889
eGenes with >1 eQTL	3,647	402	310
eVariants	3,687,265	925,916	673,696
Spliced genes	14,243	13,558	12,332
sQTL	11,450	3,165	1,920
sGenes	7,000	2,662	1,718
sGenes with >1 sQTL	2,849	425	182
sVariants	2,901,402	683,747	313,771

We examined the tissue specificity of expressed genes following an approach proposed by

Djureionovic et al. (2014), which considers genes that are at least 50-fold higher expressed in one

tissue compared to other tissues as 'highly tissue-enriched' (Supporting File 1). Genes with critical

roles in sperm maturation and the epididymal innate immune response against bacteria (e.g.,

DEFB110, *DEFB124*, *DEFB121*, *DEFB114*, *WFDC8*, *RNASE10*, *ADAM7*, *ADAM28*, *LCN8*, *LCN10*) were

among the 50 most abundant tissue-specific (i.e., not expressed in testis and vas deferens) or highly

tissue-enriched transcripts in epididymis. A gene ontology (GO) term enrichment analysis

(http://geneontology.org/) revealed that this gene set was 20.57-fold enriched (p = 4.97e-12) for

genes related to the GO term "defense response to bacterium". The same gene set enrichment

analysis on the 50 most abundant genes that were either specific or enriched in testis (e.g., *INSL3*,

FATE1, *DEFB123*, *DAZL*, *ARHGAP36*) and vas deferens (e.g., *CD52*, *TNC*, *IGFBP5*, *MMP7*, *HOXD10*,

HOXD11) did not reveal significant GO terms for either tissue.

The cattle GTEx consortium analyzed 60 testis transcriptomes (Liu et al., 2022). However, most of

them lacked metadata, were collected from pre-pubertal animals, or were from individuals of non-

taurine ancestry, precluding an immediate comparison with those from our homogeneous cohort.

The abundance of 20,221 testis-expressed genes shared between ten cattle GTEx testis

transcriptomes from post-pubertal (1.5 – 9 years) *Bos taurus taurus* bulls and our dataset was highly

correlated (Pearson's r = 0.84). The correlation was higher (Pearson's r = 0.90) for 20,002 genes that

were expressed with less than 500 TPM in our cohort. The expression of some genes differed

between our cohort and the samples included in cattle GTEx (Figure S2), likely due to different RNA

sequencing strategies applied to each dataset. The cattle GTEx dataset contained only one vas

deferens and two epididymis transcriptomes from adult bulls of taurine ancestry, preventing a

meaningful comparison with our data.

We compared gene expression in bovine testis, epididymis, and vas deferens with male reproductive

tract-specific gene expression in humans and mice (Robertson et al., 2020). Using Ensembl BioMART,

we retrieved bovine orthologs for 380 and 329 human and mice reproductive-tract specific genes,

respectively, of which the overwhelming majority (92.1% for human; 97.9% for mouse) were also

expressed in the three bovine reproductive tissues. A large fraction of the commonly expressed

genes was expressed in all three bovine tissues, but testis contributed the most and vas deferens the

least number of genes to reproductive tract-specific expression (Figure S3). We observed several

differences in gene expression between the species; for instance, *NANOS2,* which is moderately

expressed in human testis (13.84 TPM), was barely expressed in bovine reproductive tissue and thus

filtered from our dataset. We examined the expression pattern of the bovine orthologs for 220

genes that are testis-specific expressed in humans (Djureinovic et al., 2014). The overwhelming

majority (218/220) of human testis-specific genes were expressed in bovine reproductive tissue and

largely showed testis-biased expression (213/218), although, most were also expressed in bovine

epididymis (n=22), vas deferens (n=2), or both other tissues (n=192). We again identified noteworthy

differences between the species. Bovine orthologs of two highly expressed testis-specific genes in

humans, *DEFB119* encoding defensin beta 119 and *EPPIN* encoding epididymal peptidase inhibitor,

were also highly expressed in bovine testis (median TPM *DEFB119* (ENSBTAG00000003364): 64.5;

median TPM *EPPIN* (ENSBTAG00000001495): 498.2); however, both genes were three-fold higher

expressed in bovine epididymis (median TPM: 217.4 and 1524.3). We detected low and moderate

expression of *RNASE11* encoding ribonuclease A family member 11 in epididymis (median TPM:

2.18) and vas deferens (median TPM: 19.06) but did not detect any expression in bovine testis tissue

despite reports that it is highly tissue-enriched in human testis (Djureinovic et al., 2014).

We used LeafCutter to detect RNA splicing variation and calculated 'percent spliced-in' (PSI) values

for excised-intron clusters. Across the three tissues we observed 15,493 alternatively spliced genes,

with 95% of these being protein-coding genes (14,719 genes; Table S3). We detected 11,087 genes

that were alternatively spliced in all three tissues. Genes that were alternatively spliced in all three

tissues were higher expressed than those for which we did not detect alternative isoforms (Table

S4). We identified 139,186 splice junctions and 46,417 intron clusters in testis, 114,343 splice

junctions and 44,891 intron clusters in epididymis, and 103,809 splice junctions and 40,480 intron

clusters in vas deferens. The total number of genes with alternative splicing events was similar

across the three tissues, with 14,243, 13,558, and 12,332 alternatively spliced genes for testis,

epididymis, and vas deferens, respectively.

The similarity of the transcriptional profile of the three male reproductive tissues was assessed

through cluster analyses. Principal component analysis (PCA) of normalized expression values (TPM)

and splicing phenotypes (PSI) separated samples by tissue type (Figure 1C; 1D), which was further

supported by hierarchical clustering (Figure S4). In both expression and splicing PCAs, the first PC

separated testis from epididymis and vas deferens, suggesting that the transcriptional profile of

testis is distinct from other male reproductive tissues. To further explore the similarity between

tissues, we inferred pairwise similarity of gene expression and splice-junction counts with Spearman

correlation. For both expression and splicing, vas deferens and epididymis were more similar to each

other than to testis (epididymis x vas deferens – expression: ρ = 0.92, splicing: ρ = 0.83; epididymis x

testis – expression: ρ = 0.82, splicing: ρ = 0.67; vas deferens x testis – expression: ρ = 0.74, splicing: ρ

= 0.57).

Thousands of variants impact gene expression and splicing

To identify loci that influence gene expression (expression QTL, hereafter referred to as eQTL),

normalized and standardized TPM values were regressed on variants that had minor allele frequency

greater than 1.0% and were within 1 Mb of the annotated transcription start site (TSS). We

accounted for covariates such as age, RNA integrity, and hidden confounders that were estimated

directly from the data (PEER factors; Figure S5). We identified eQTL in at least one of the three

tissues for 12,950 genes (hereafter referred to as eGenes) at a false discovery rate (FDR) of 5%. A

majority of eGenes were protein-coding (11,951 eGenes; Table S3). Testis had considerably more

eGenes than epididymis and vas deferens (Table 1)—over 55% of genes expressed in testis were also

eGenes. Epididymis had 4,347 eGenes (21% of genes expressed in epididymis) and vas deferens had

3,889 eGenes (20% of genes expressed in vas deferens). One-third of testis eGenes had more than

one independent-acting eQTL (Figure 2A), corresponding to 15,642 eQTL total and 3,687,265 unique

significant variants (eVariants). We identified 4,768 independent eQTL (9.2% eGenes having > 1

eQTL; Figure 2A) and 925,916 eVariants in epididymis. Vas deferens had the fewest eQTL, with 4,211

independent eQTL and 673,696 eVariants. For all tissues, approximately half of all eQTL were within

100 kb of the TSS (8,000 testis eQTL; 2,802 epididymis eQTL; 2,423 vas deferens eQTL; Figure S6).

The effect of each eQTL was quantified as the allelic fold change (slope_aFC) in expression of the

associated eGene. The slope_aFC value was highly correlated (Spearman's $\rho \geq$ 0.84) with the beta

coefficient from the linear model used for eQTL mapping (Figure 2B). Across the three tissues, only

2,321 eQTL (9.4%) had large effects (defined as |slope_aFC $\geq$ 1|), with 65% of these loci residing

within 100 kb of the eGene's TSS. Notably, testis had the smallest proportion of large effect eQTL

(7.58%; 1,186 eQTL; Figure S6). The proportion of large effect eQTL in both epididymis and vas

deferens was approximately 12% (599 and 536 eQTL in epididymis and vas deferens, respectively).

Large effect eQTL were enriched for low-frequency alleles (Figure S7). We observed that eQTL

annotated as non-coding transcript variants had a stronger effect on expression (median slope_aFC

= 0.93; Figure 2D).

Figure 2. **Properties of eQTL and sQTL in three male reproductive tissues.** A.) Proportion of independent-

acting eQTL (top) and sQTL (bottom) across tissues. B.) Correlation (Spearman's ρ) between the β-coefficient of

the linear model (used for eQTL identification) and slope_aFC of each eQTL's top eVariant. The linear

relationship between the two variables is shown with dotted lines. Colors correspond to the tissue type. C.)

Cumulative proportion (%) of eQTL showing significant allelic imbalance in the expression of the associated

eGene grouped (in bins of 5 Kb) by their proximity to the transcription start site (TSS). Data for large

($|\text{slope\_aFC}| \geq 1$) and small ($|\text{slope\_aFC}| < 1$) effect eQTL are presented separately. Similar data for eQTL not

showing significant allelic imbalance are shown for comparison D.) Annotation class of eQTL and rank-

normalized absolute effect size for eQTL identified in all three tissues. The number of eQTL within an

annotation class is listed in parentheses. E.) Distribution of sQTL (top independent-acting sVariants) from the

ICSS for all three tissues. Darker colors represent large effect sQTL ($|\text{β-coefficient} \geq 1|$). F.) Annotation class of

sQTL and rank-normalized absolute effect size for sQTL identified in all three tissues. The number of sQTL

within an annotation class is listed in parentheses.

eQTL are expected to cause an imbalanced expression of the associated gene if they are cis-acting

(Aguet et al., 2017; Pirinen et al., 2015). In this scenario, the expression of the associated eGene is

influenced only by the linked eQTL allele leading to unbalanced expression of the eGene. Thus, to

assess and independently quantify putative cis-regulatory effects of the eQTL, we investigated allelic

imbalance in the expression of associated eGenes. In the three tissues between 78 and 80% of the

variant-gene pairs were available to this evaluation, of which approximately a quarter (23 – 28%)

showed significant evidence of allelic imbalance (FDR < 5%; Wilcoxon rank sum test). The proportion

of eGenes that showed allelic imbalance was nearly twice as high (49.8, 46.8, and 47.2% in testis,

epididymis, and vas deferens, respectively) for large effect eQTL ($|slope\_aFC| \geq 1$; n = 1156, 669, 545

in testis, epididymis, and vas deferens, respectively (Figure S8B; Figure S8C)). Approximately three

quarters (76.53, 76.53, and 74.49% for testis, epididymis, and vas deferens, respectively) of the

eGenes with significant allelic imbalance were within 100 kb of the eQTL (Figure 2C). This is between

1.26- and 1.44-fold enrichment (p < 2.16e-16; Fisher's exact test) compared to random expectation

(Figure S8A). The eQTL effect estimated as the magnitude of allelic imbalance (AI_aFC) was strongly

correlated with slope_aFC (Spearman's ρ = 0.69, 0.73, 0.75 in testis, epididymis, and vas deferens,

respectively). The correlation was even stronger (Spearman's ρ = 0.87, 0.88 and 0.88 for testis,

epididymis and vas deferens, respectively) for a subset of the eQTL where the associated eGenes

showed significant allelic imbalance (Figure S8C).

To identify variants that influence splicing, we conducted sQTL mapping between variants within 1

Mb of the intron cluster start site (ICSS) and intron-excision ratios of inferred intron clusters, which

were normalized and standardized within and across samples, respectively. We identified 7,777

spliced genes across the three tissues with at least one sQTL (sGenes; Table 1) at an FDR of 5%—a

majority of which were protein-coding genes (7,471 protein-coding genes; 96% of sGenes). Testis

had the most sQTL of the three tissues, further corroborating its transcriptional complexity. We

detected 7,000 sGenes in testis, which accounted for nearly half of its alternatively spliced genes. For

epididymis and vas deferens, we identified 2,662 and 1,718 sGenes, respectively, corresponding to

20% and 14% of their spliced genes. In testis, 40.7% of sGenes had more than one independent-

acting sQTL, resulting in 11,450 independent sQTL and 2,901,402 unique significant variants

(sVariants; Figure 2A). Fewer sGenes had multiple independent-acting sQTL in epididymis and vas

deferens (approximately 16% in epididymis and 11% in vas deferens; Figure 2A); in total, epididymis

contained 3,165 sQTL and 683,747 sVariants while vas deferens contained 1,920 sQTL and 313,771

sVariants. In all tissues, between 60 and 70% of sQTL were located within 100 kb of the ICSS,

corresponding to 6,822, 2,190, and 1,339 sQTL for testis, epididymis, and vas deferens, respectively

(Figure 2E).

sQTL effects were quantified as the regression coefficient from the linear model used for sQTL

mapping. There were few large effect sQTL (3,793 sQTL with |β-coefficient| ≥ 1) across the three

tissues. Most large effect sQTL were within 100 kb of the ICSS (65%, or 2,476 sQTL across all tissues;

Figure 2E). The proportion of large effect sQTL was similar across the three tissues, corresponding to

22.7%, 22.5%, and 24.9% of sQTL in testis, epididymis, and vas deferens, respectively (2,603 sQTL in

testis, 712 sQTL in epididymis, and 478 sQTL in vas deferens; Figure 2E). We observed that top

sVariants within splicing regions—such as splice donor and splice acceptor sites—had the largest

effect sizes (median effect size = 1.14; Figure 2F), corroborating that variants overlapping these

nucleotides have pronounced impacts on pre-mRNA splicing.

Identifying distant—or trans—eQTL and sQTL that lie > 5 Mb from a gene's TSS requires larger

sample sizes, as they typically have small effects and an increased multiple-testing burden. Thus, we

only conducted trans eQTL and sQTL analyses for testis, for which we had the largest number of

samples. We observed 76 trans eGenes and 6,038 trans eVariants (FDR 5%) in testis. Most trans

eQTL were located on a different chromosome than the target trans eGene (73 trans eGenes and

6,024 trans eVariants; Figure S9A). Nearly half (46%; 35 total trans eQTL) of trans eQTL were

colocalized (PP4 > 0.8) with at least one cis eQTL, suggesting a complex network of cis mediation. We

identified 33 intron clusters with trans sQTL, corresponding to 25 trans sGenes and 1,023 trans

sVariants (Figure S9B).

Testis molQTL effects are unique and small, while molQTL effects in other male reproductive

tissues are shared

Tissues with similar biological functions often share molQTL (GTEx Consortium, 2020; Liu et al., 2022;

Urbut et al., 2019). Thus, we sought to determine if eQTL and sQTL in male reproductive tissues

were shared or tissue-specific. We used Mashr to compare molQTL effects for genes that were

expressed in all three tissues and considered 11,311 and 5,012 top eVariants and sVariants,

respectively. For eQTL, we identified 10,738 variant-gene pairs that were significant in at least one

tissue (local false sign rate (LFSR) < 0.05). A large proportion of eQTL were tissue-specific (4,760

eQTL; 44% of the eQTL considered), while there was a similar number of eQTL identified in either

two or three tissues (two tissues: 2,949 eQTL, 27% of eQTL; three tissues: 3,029 eQTL, 28% of eQTL;

Figure 3A). Testis had the most tissue-specific eQTL, followed by vas deferens, and epididymis

(Figure 3B). eQTL effects were most similar between vas deferens and epididymis (Spearman's ρ =

0.58), and clearly less similar between testis and epididymis (Spearman's ρ = 0.32) and between

testis and vas deferens (Spearman's ρ = 0.28). Conversely, most sQTL were ubiquitous—73% of the

4,737 variant-gene pairs were significant (LFSR < 0.05) in all tissues (3,451 sQTL; Figure 3A). Only 648

sQTL were tissue-specific (14% of sQTL considered; Figure 3A), most of which were in testis (505

sQTL; Figure 3B). We observed high pairwise similarity of sQTL effects across the three tissues,

particularly between epididymis and vas deferens (testis x epididymis: Spearman's ρ = 0.75; testis x

vas deferens: Spearman's ρ = 0.69; epididymis x vas deferens: Spearman's ρ = 0.88).

Figure 3. Tissue-specific and shared effects of eQTL and sQTL. A.) Proportion of eQTL and sQTL that are

significant (LFSR < 0.05) in one, two, or three tissues. B.) Overlap of significant (LFSR < 0.05) eQTL and sQTL

across the three tissues. Bar height represents the number of QTL. C.) Boxplots of the eQTL effect size across

tissue specificity classes (significant in one, two, or all three tissues at a threshold of LFSR < 0.05) for testis

(blue), epididymis (green), and vas deferens (purple) tissues. Absolute effect size corresponds to |aFC_slope|.

Median effect size of each category is included. D.) Boxplots demonstrating the effect size of sQTL across tissue

specificity classes for testis, epididymis, and vas deferens tissues. Absolute effect size corresponds to the β-

coefficient of the linear model. Median effect size of each category is included.

Tissue-specific molQTL typically have larger effects than those identified in many, or all, tissues (The

FarmGTEx-PigGTEx Consortium et al., 2022; Urbut et al., 2019). We confirmed this phenomenon for

epididymis- and vas deferens-specific eQTL, which had larger effects (slope_aFC) than eQTL found in

all three tissues (vas deferens: Wilcoxon rank-sum p = 2.48e-16; epididymis: Wilcoxon rank-sum p =

0.003; Figure 3C). Vas deferens-specific sQTL also had larger effects than ubiquitous sQTL (Wilcoxon

rank-sum p = 7.92e-11), though there was no statistical difference in sQTL effect size across tissue

specificity classes for epididymis (Wilcoxon rank-sum p = 0.39). Alternatively, testis specific eQTL and

sQTL had smaller effects than those shared across all three tissues (eQTL: Wilcoxon rank-sum p =

3.93e-30; sQTL: Wilcoxon rank-sum p = 1.40e-16; Figure 3D), suggesting that testis is composed of

many tissue-specific small effect molQTL.

Integration of molecular phenotypes, molQTL, and GWAS summary statistics reveals genes

associated with male fertility

We sought to investigate the impact of molQTL on male reproductive ability. First, we conducted

GWAS for male fertility with a cohort of 3,736 bulls that had imputed genotypes for 14,587,859

SNPs, and separately tested additive and non-additive inheritance. The fertility of the bulls was

quantified through the proportion of successful artificial inseminations and accounted for genetic

and environmental factors (see Hiltpold et al., 2021). While the GWAS cohort had no overlap with

the molQTL cohort, both predominantly consisted of Brown Swiss bulls. Non-additive association

testing revealed four male fertility QTL on chromosomes 1, 6, 18, and 26 (significance threshold of

5e-08) that contained 4,890 variants (Figure 4A). Additive GWAS did not detect any additional

significant associations and failed to identify the QTL on chromosomes 1 and 18. The QTL on

chromosomes 6 and 26 were detected by the additive GWAS, but with lower significance (Figure

S10).

We implemented the MetaXcan framework to establish if variation in molecular phenotypes is

associated with variation in male fertility. Briefly, S-PrediXcan was used to integrate summary

statistics from the non-additive GWAS with expression and splicing phenotypes from the three

reproductive tissues. Signals were subsequently filtered based on the colocalization probability

between male fertility QTL and molQTL. We identified two and seven unique genes for expression

and splicing, respectively, that overlapped three of the male fertility QTL (Figure 4, Table 2). The

most striking association was on chromosome 6; *WDR19* encoding WD Repeat Domain 19 was the

most significantly associated gene in epididymis and vas deferens (p = 3.17e-43 and 1.04e-48). An

intron cluster encompassing a canonical (58373894:58374821) and a cryptic (58373885:58374821)

splice junction was the top fertility-associated molecular phenotype in both tissues (Figure S11). A

highly significant S-PrediXcan signal (p = 6.10e-38) was also obtained for this *WDR19* intron cluster in

testis, however the evidence for colocalization with the GWAS peak was low (PP.H4.abf = 0.39).

Figure 4. Genes associated with male fertility. Manhattan plots showing genome-wide association between

(A) male fertility and imputed sequence variants from non-additive association models. Variants significantly (p

< 5e-8) associated with male fertility are highlighted in orange. Transcriptome-wide association with S-

PrediXcan between (B) expression values or (C) splicing phenotypes and male fertility. Significantly associated

genes after the stringent filtering based on the MetaXcan framework are marked with their names.

The splicing MetaXcan in testis revealed an association (p = 1.71e-15) between *SPATA16* encoding

spermatogenesis associated protein 16 and male fertility (Figure 4C). *SPATA16* mRNA was highly

expressed in testis (270.57 ± 32.26 TPM), lowly expressed in epididymis (15.28 ± 9.90 TPM), and

barely expressed in vas deferens (1.05 ± 1.18 TPM). Only one *SPATA16* transcript is annotated in

Ensembl, but the Refseq annotation suggests three isoforms, of which the canonical isoform

(XM_002684936.6) was 12 and 17 times more abundant in testis than two alternative isoforms

XM_024995427.1 and XM_005201702.4 (Figure S12A). The fertility-associated *SPATA16* intron

cluster comprised four splice junctions that span the first four exons, including a non-coding exon

that is only part of the Refseq annotation (Figure 5C; Figure 5D; Figure 5F). The fertility and splicing

QTL were colocalized with a probability of 0.86 and a variant (Chr1:93,954,637 bp) 365 kb upstream

the annotated *SPATA16* transcription start site was the most probable causal variant (Figure 5A). An

association analysis conditional on Chr1:93,954,637 did not identify any additional variants as being

associated with male fertility on chromosome 1 (Figure 5B).

Table 2. Male fertility-associated genes prioritized by MetaXcan. P-value is the gene's p-value from S-

PrediXcan. PP.H4 is the probability of a shared causal variant for molQTL and QTL from colocalization analysis.

Colocalized variant is the most likely colocalized variant from coloc.abf (H4.PP.SNP with largest posterior

probability). β–coefficient is the effect estimate of the most likely colocalized variant from the linear model

used for molQTL mapping.

Gene	Expression (TPM)			Splice junction and intron cluster	Molecular phenotype	Tissue	P-value	PP.H4	Colocalized variant	β–coefficient
	Testis	Epididymis	Vas deferens							
SPATA16	270.57 ± 32.26	15.28 ± 9.90	1.05 ± 1.18	1:94345963-94396748:clu_3038	Splicing	Testis	1.71E-15	0.86	Chr1: 93,954,637	0.78
WDR19	20.09 ± 2.65	29.46 ± 19.57	36.26 ± 4.82	6:58373894-58374821:clu_63814	Splicing	Epididymis	3.17E-43	0.96	Chr6: 57,900,948	-1.17
				6:58373894-58374821:clu_46646		Vas deferens	1.04E-48	0.99	Chr6: 57,900,948	-1.34
KCTD8	2.71 ± 0.73	0.73 ± 1.00	0.68 ± 0.93	6:62885148-62886027:clu_81370	Splicing	Testis	8.10E-14	0.81	Chr6: 62,206,554	-0.73
KCTD19	238.01 ± 56.20	9.05 ± 6.26	0.58 ± 1.42	--	Expression	Testis	2.27E-07	0.96	Chr18: 34,914,479	1.12
				18:34910787-34910930:clu_25862	Splicing	Epididymis	2.28E-07	0.87	Chr18: 34,914,479	0.67
ZDHHC1	21.96 ± 3.04	24.77 ± 17.29	21.00 ± 3.32	18:35019435-35026730:clu_33937	Splicing	Testis	8.11E-07	0.94	Chr18:34,815,920	-1.12
				18:35019435-35026730:clu_25875		Epididymis	9.44E-07	0.92	Chr18: 34,914,479	-0.81
FBXL8	3.80 ± 1.51	2.05 ± 0.72	1.52 ± 0.46	--	Expression	Epididymis	7.86E-07	0.87	Chr18:35,059,627	0.96
EXOC3L1	13.50 ± 2.15	1.34 ± 0.55	1.73 ± 0.50	18:34825696-34825972:clu_33908	Splicing	Testis	9.91E-07	0.93	Chr18: 34,914,479	0.87
ENSBTAG00000001287	7.17 ± 0.97	3.33 ± 1.00	3.26 ± 0.81	18:34842264-34842709:clu_33910	Splicing	Testis	1.73E-07	0.93	Chr18: 34,815,920	-0.91

The Chr1:93,954,637 T allele compromises male fertility (p = 4.15e-16; Figure 5E) but is associated

with increased *SPATA16* mRNA expression (β = 0.71 ± 0.17 TPM_{norm}; p = 6.41e-05)—albeit not at the

stringent significance threshold applied to our eQTL analysis (Figure S12B; Figure S12C). Eleven of

the twelve *SPATA16* exons were expressed at higher levels in animals carrying the T allele, though

this difference was only significant (p = 3.7e-06) for the fourth exon (94,396,748 bp – 94,396,893

bp). Interestingly, the second exon (94,323,018 bp – 94,323,140 bp) is the only exon for which

abundance is reduced (β = 0.25 ± 0.09 TPM$_{norm}$; p = 0.0057) by the Chr1:93,954,637 T allele (Figure

S12E). This exon is part of SPATA16 isoform XM_024995427.1 but is absent in both the canonical

(XM_002684936.6) and another alternative isoform (XM_005201702.4). While the Chr1:93,954,637

T allele has no effect on the abundance of the predominant canonical isoform (XM_002684936.6, p =

0.15), it increases (β = 3.6 ± 0.68 TPM; p = 6.7e−07) abundance of XM_005201702.4 and reduces

abundance of XM_024995427.1 (β = −1.1 ± 0.42 TPM; p = 0.015), suggesting that it promotes

differential isoform usage (Figure S12D). This is further supported by an inverse association between

inclusion ratios of the second and third exon with the abundance of two alternative SPATA16

isoforms (XM_005201702.4 and XM_024995427.1; Figure S13).

The non-additive GWAS for male fertility revealed a QTL on chromosome 18 (p = 6.41e-09, lead SNP

at 36,480,384) that coincides with a previously localized 3 Mb region (between 34 and 37 Mb), for

which no compelling candidate causal variants were found (Hiltpold et al., 2021). This 3 Mb window

contains 97, 27, and 12 eQTL and sQTL in testis, epididymis and vas deferens, respectively. The

MetaXcan and colocalization analyses revealed five significantly associated genes

(*ENSBTAG00000001287*, *EXOC3L1*, *FBXL8*, *KCTD19*, *ZDHHC1*) at this QTL, of which *KCTD19* encoding

the potassium channel tetramerization domain containing protein 19 shows a highly testis-biased

expression (Table 2). Two genes (*KCTD19* and *ZDHHC1* encoding the zinc finger DHHC-type

containing protein 1) were each associated with two molecular phenotypes in two tissues. Three

variants (Chr18:34,914,479, Chr18:34,815,920, Chr18:35,059,627), which are in moderate LD

(between 0.54 and 0.6), are colocalized for the male fertility QTL and seven distinct molecular

phenotypes (Table 2). Association analyses conditional on any of these variants did not identify

additional variants associated with male fertility on chromosome 18. The Chr18:34,914,479 T allele is

associated with higher fertility, an increased KCTD19 mRNA abundance in testis, and the expression

of a KCTD19 isoform in the epididymis that has an additional 30 bp coding sequence added to the

14th exon which results in a protein that is 10 amino acids longer than that encoded by the canonical

form (Figure S14). Alternative *KCTD19* splicing was also highly significant (p = 3.13e–15) in testis, but

MetaXcan did not reveal association of this molecular phenotype with male fertility. This variant

(Chr18:34,914,479) was also associated with differential *ZDHHC1* splicing in epididymis and

differential *EXOC3L1* splicing in testis.

A QTL (p = 7.59e-23, lead SNP at Chr26:50,145,932) for male fertility was detected on chromosome

26. Neither the splicing nor the expression MetaXcan revealed any significant gene-trait associations

for this QTL.

Figure 5. A molecular *SPATA16* phenotype overlaps with a male fertility QTL. A.) Mirrored plots of –log$_{10}$(p)-

values from association testing between imputed sequence variants and bull fertility (top) and splicing

phenotypes for intron cluster 3038 of the gene *SPATA16* (bottom). The inset shows the correlation between

GWAS and sQTL –log$_{10}$(p)-values. Color indicates the linkage disequilibrium (R^2) between the most likely

colocalized variant (1:93954637_C_T, red colored) and all other variants. B.) Association testing between

imputed sequence variants and male fertility before (gray) and after (maroon) conditioning on the genotype of

the most likely colocalized variant (1:93954637_C_T). C.) Gene structure of bovine *SPATA16*, showing positions

of exons (black lines) and alternative splicing events (arcs). D.) Splicing within *SPATA16* intron cluster 3038 for

different genotypes of the top colocalized variant. Red lines show the different splice junctions with the values

representing "percent spliced in" estimates within the intron cluster. E.) Standardized bull fertility across the

genotypes of the colocalized variant. F.) PSI values for different splicing events within Cluster 3038 across

different genotypes of the most likely colocalized variant.

Discussion

We characterized the transcriptomic profile of testis, epididymis, and vas deferens in a cohort of 118

bulls. To the best of our knowledge, a cohort of this size has not been established for male

reproductive tissues thus far, though smaller cohorts are available for several species (Djureinovic et

al., 2014; Gamble et al., 2020; Zhang et al., 2023). Our molQTL cohort contained bulls within a

narrow post-pubertal age range that predominantly belonged to one breed of European taurine

cattle. We demonstrated the benefit of establishing such a homogeneous cohort for transcriptome

analyses, as we detected 14-times (11,164 vs. 809) more eGenes and 7-times more sGenes (11,450

vs. 1,573) in testis than cattle GTEx (Liu et al., 2022) despite our molQTL cohort being only twice as

large (117 individuals vs. 60 individuals in cattle GTEx). This suggests that the heterogeneity of cattle

GTEx leads to substantial underestimation of the transcriptional complexity of bovine testis and

likely other tissues. The number of eGenes and sGenes identified in testis in our study is similar to

what was reported in a heterogeneous—but three-times larger—human GTEx cohort (GTEx

Consortium, 2020). Comparing our findings with similar-sized cohorts sampled at other

developmental stages and the analysis of single-cell transcriptomes will reveal spatiotemporal

transcriptomic changes in male reproductive tissue, such as those occurring during testicular

development and puberty (Guo et al., 2018, 2020; Liu et al., 2023).

Our comprehensive set of expressed and spliced genes enabled us to thoroughly characterize and

compare the transcriptional complexity of three reproductive tissues in a large mammal, which was

so far not possible due to a lack of epididymis and vas deferens data in all existing GTEx cohorts

(Robertson et al., 2020). We revealed nearly 20,000 expressed and 14,000 spliced genes within each

tissue. Expressed and spliced genes were generally shared across the three reproductive tissues;

however, the transcriptional profiles of vas deferens and epididymis were similar, while that of testis

was distinct. We found good consensus between pairs of bovine-mouse and bovine-human

orthologous genes with male reproductive tract-specific expression, thus demonstrating that our

molQTL cohort is a valuable resource for translational research in mammalian male reproductive

biology. We observed a similar pattern for eQTL and sQTL—the effects of regulatory variants in vas

deferens and epididymis were similar and often shared, whereas testis had a distinct and

exceptionally large array of eQTL and sQTL. When compared to the other tissues, we found between

3- and 6-times the number of eGenes and sGenes in testis, many of which were regulated by

multiple eQTL and sQTL, respectively, which emphasizes its transcriptional complexity that has been

reported previously (Ramsköld et al., 2009; Soumillon et al., 2013). Testis contained thousands of

tissue-specific, small effect molQTL. An abundance of molQTL in testis was observed in numerous

mammalian species (GTEx Consortium, 2020; The FarmGTEx-PigGTEx Consortium et al., 2022),

though the reason for this remains unclear. The accumulation of small effect molQTL in testis may be

a result of male sexual selection and sperm competition, which is driven by a more permissive

transcription (Soumillon et al., 2013), relaxed pleiotropic constraints, and other selective

mechanisms (Kaessmann, 2010; Murat et al., 2023). In humans, when compared to other tissues,

testis eQTL were more likely to contain signatures of positive selection (Quiver & Lachance, 2022);

furthermore, tissue specific eQTL that showed evidence of positive selection had small effect sizes

(Quiver & Lachance, 2022). Thus, functional variants in testis may experience unique selection

pressures that are essential for male reproduction. Further research is warranted to fully understand

the genomic basis of testis's complex functional patterns.

By leveraging our comprehensive array of molecular phenotypes from reproductive tissues and

fertility records from genotyped artificial insemination bulls through gene-based association testing

and colocalization analyses, we identified candidate causal regulatory mechanisms underpinning

three of the four male fertility QTL. Among them is a previously suspected candidate causal

molecular phenotype of *WDR19* (Hiltpold et al., 2020), which adds additional confidence that this

approach provides biologically meaningful associations. Our results corroborate that gene splicing

and expression variation have substantial effects on complex traits (Xiang et al., 2022) and highlight

that regulatory variants need to be considered thoroughly as candidate causal variants. The male

fertility QTL were associated with two large effect and one small effect molQTL, thereby

demonstrating that molQTL effect size may be a poor predictor of impact on complex trait variation.

Spermatogenesis, sperm maturation, and sperm transport are pivotal functions of the three

reproductive tissues considered. Gene-based association testing with traits such as sperm

morphology or sperm motility may reveal additional gene-trait associations, as they are closer

related to the biological function of these tissues than male fertility. However, establishing cohorts

with repeated semen quality measurements that are large enough to identify loci that explain only a

small proportion of trait variation is challenging in cattle—and almost impossible in most other

species.

We identified an alternative splicing event in testis for *SPATA16* that was significantly associated

with male fertility. The colocalized variant fell within an intergenic region between *SPATA16* and

NLGN1. Previously, a GWAS-only approach prioritized a missense variant of *SPATA16* as a putative

causal variant (Hiltpold et al., 2021). However, by combining molecular phenotypes, molQTL

analyses, and GWAS—which were appropriately modeled to QTL inheritance—we identified an

alternative splicing and regulatory mechanism that is significantly more likely to underpin the QTL.

While loss-of-function alleles of *SPATA16* lead to severe sperm defects that prevent fertilization

(Dam et al., 2007; Fujihara et al., 2017), our results demonstrate that variants mediating *SPATA16*

expression and splicing can contribute to quantitative variation in male fertility as they are less

deleterious to protein function. We also identified several gene-trait associations that showed

colocalization between molQTL and a non-additive male fertility QTL on chromosome 18. Our

inability to resolve this QTL may be due to high linkage disequilibrium or possibly indicates that the

causal variant affects multiple genes, as may be the case for large structural variants (Leonard et al.,

2023). However, MetaXcan prioritized genes that were previously implicated in male fertility

disorders, including *KCTD19* and *ZDHHC1* (Oura et al., 2021; S. Wang et al., 2021; W. Wang et al.,

2023). Future fine-mapping efforts that consider additional phenotypic observations and complex

genotype variations, such as structural variants called from long read cohorts, will be required to

fully describe the molecular underpinnings of this QTL (Leonard et al., 2023).

Methods

Ethics statement

Tissue was sampled at a commercial abattoir. The decision to slaughter the bulls was independent

from our study. None of the authors of the present study were involved in the decision to slaughter

the bulls. No ethics approval was required for this study.

Tissue sampling and DNA/RNA extraction

Testes from 128 bulls were collected at a commercial abattoir in Zürich, Switzerland after regular

slaughter, then transported to a laboratory for preparation. Testis, epididymis (caput), and vas

deferens tissue samples were flash-frozen in liquid nitrogen and stored at -80°C until DNA and RNA

extraction. The time from slaughter to freezing ranged from 40 minutes to 270 minutes (average:

126 minutes ± 47 minutes). Sample collection took place during the fall, winter, and spring of 2019,

2020, and 2021. Breed was assigned according to entries from the Swiss Braunvieh herdbook and

the animal traffic database.

We extracted DNA and total RNA from testis tissue with Qiagen AllPrep Mini kits (Qiagen, Hilden,

Germany). Frozen testis tissue was homogenized with a MagNA Lyser (Roche, Basel, Switzerland) at

6000 rpm for 50 seconds (x2) in RTL Plus Buffer with β-mercaptoethanol. DNA extraction followed

standard manufacturer protocols. Concentration was estimated with a Qubit 2.0 fluorometer

(Thermo Fisher Scientific) prior to submission for whole genome sequencing (WGS). Extraction of

testis total RNA followed manufacturer protocols. Epididymis and vas deferens total RNA was

extracted from frozen tissue with Qiagen RNeasy Mini Kits (Qiagen, Hilden, Germany) and included

an on-column DNase digestion with Qiagen RNase-Free DNase (Qiagen, Hilden, Germany). Frozen

epididymis tissue was homogenized with the same conditions as described for testis tissue. We

optimized the homogenization and lysis of vas deferens tissue by adding an additional

homogenization period at 6000rpm for 50 seconds and a 60-minute waiting period. We assessed the

quality of testis, epididymis, and vas deferens total RNA with an RNA integrity number (RIN) inferred

from a Bioanalyzer RNA 600 Nano assay (Agilent Technologies). For subsequent analyses, we only

considered samples with RIN > 4.0.

DNA and RNA sequencing

Genomic libraries (paired-end, 150 bp) for WGS were prepared with the Illumina TruSeq DNA PCR-

Free protocol and sequenced on two Illumina NovaSeq 6000 S4 flowcells. Paired-end total RNA

libraries (150 bp) were prepared with the TruSeq Stranded Total RNA protocol and included rRNA

depletion with Ribo-Zero Plus. Libraries were sequenced on four S4 flowcells and one S2 flowcell of

the Illumina Novaseq 6000.

DNA alignment and variant calling

We used fastp (v0.19.4; Chen et al., 2018) with default parameters to remove adapter sequences

and low-quality bases, and trim poly-G tails from raw DNA sequence data. After quality control, we

aligned reads to the ARS-UCD1.2 reference genome (Rosen et al., 2020) with the mem-algorithm

from BWA (v0.717; H. Li, 2013) and the -M flag. We sorted the aligned reads by coordinates with

Sambamba (v0.6.6; Tarasov et al., 2015), then combined the read-group specific BAM files to create

sample-specific sorted BAM files. We marked duplicate reads in the sample specific sorted BAM files

with the MarkDuplicates module from Picard tools (https://broadinstitute.github.io/picard/). We

used mosdepth (v0.6.6; Pedersen & Quinlan, 2018) to infer the sequencing coverage at a given

genomic position, which was then used to estimate the average coverage per sample. Only high-

quality reads (mapping quality > 10 or without SAM flag 1796) were considered when calculating the

average coverage. We called variants with DeepVariant (v1.3; (Poplin et al., 2018) using the WGS

mode. Called samples were merged using GLnexus (v1.4.1; Yun et al., 2020) with the

"DeepVariantWGS" configuration. Variants that had a missingness rate greater than 50% and

variants for which the genotypes deviated from Hardy-Weinberg proportions ($P < 1e-08$) were

removed. We applied Beagle (v4.1; Browning & Browning, 2016) to impute sporadically missing

genotypes and infer haplotypes, then removed sites with model-based imputation accuracy < 0.5. To

assess relationships among individuals, we constructed a genomic relationship matrix with 366,090

LD-pruned variants (obtained from Plink v1.9 with --indep-pairwise 1000 5 0.2; Chang et al., 2015)

that had MAF > 0.5% and performed a PCA with the "pca" function in QTLtools (Delaneau et al.,

2017).

RNA alignment

We removed adapter sequences, low-quality bases, poly-A tails, and poly-G tails from the RNA

sequence data with fastp. We split cleaned reads into read groups with gdc-fastq-splitter

(https://github.com/kmhernan/gdc-fastq-splitter) then aligned them to the cattle reference genome

(ARS-UCD1.2) and the Ensembl gene annotation (release 104) with STAR (version 2.7.9a; Dobin et al.,

2013). To account for allelic mapping bias in the sQTL and allele specific expression (ASE) analyses,

we produced additional alignments with the flag --waspOutputMode and heterozygous SNPs. We

performed coordinate sorting and added read groups with Picard

(https://broadinstitute.github.io/picard/). We merged each sample's BAM files with Sambamba

(v0.6.6; Tarasov et al., 2015) and marked duplicate reads with the MarkDuplicates module from

Picard (https://broadinstitute.github.io/picard/). Statistics on the alignment quality were inferred

with the flagstat function from Sambamba for each sample.

Expression and splicing quantification

We quantified gene-level expression with the "quan" function in QTLtools. We produced gene-level

TPM values with high quality reads that passed all alignment quality filters and only included reads

that were uniquely mapped and properly paired. Gene-level read counts were inferred with

featureCounts (Liao et al., 2014). We filtered expression values for each tissue to include genes with

expression ≥ 0.1 TPM and ≥ 6 reads in ≥ 20% of samples. The filtered expressed genes were inverse

normal transformed and quantile normalized for subsequent analyses (Robinson & Oshlack, 2010).

To observe clustering within and across tissue types, we used the filtered, normalized TPM values to

perform a PCA and hierarchical clustering with the dendextend (v1.16.0) package in R.

To infer splicing events, we quantified intron-excision values for identified intron clusters. We

extracted exon-exon junctions from reads that passed WASP filtering with Regtools (Cotto et al.,

2023) and considered a minimum anchor length of 8 bp, minimum intron size of 50 bp, and

maximum intron size of 500,000 bp. Intron clustering, calculation of intron excision ratios, filtering,

and preparation for sQTL mapping was performed with Leafcutter (Y. I. Li et al., 2018) and executed

with the cluster_prepare_fastqtl.py wrapper provided by the human GTEx consortium

(https://github.com/broadinstitute/gtex-pipeline/tree/master/qtl/leafcutter). We assigned clusters

to annotated genes with an altered map_clusters_to_genes.R script that accounted for strandedness

and gene coordinates from a collapsed gene annotation for cattle (Ensembl release 104; generated

with scripts from https://github.com/broadinstitute/gtex-pipeline/tree/master/gene_model). We

limited our dataset to high-quality introns by imposing stringent filters that removed introns without

read counts in > 50% of samples, introns with low variability across samples, and introns with fewer

than max (10, 0.1n) unique values (where n is sample size). To normalize the filtered counts and

produce files for sQTL mapping, we used the prepare_phenotype_table.py script from Leafcutter.

We performed hierarchical clustering and conducted a PCA on the resulting normalized intron

excision phenotypes and hierarchical clustering with the dendextend package in R.

eQTL mapping, allelic imbalance, and sQTL mapping

eQTL mapping was performed for all three tissues with QTLtools. To account for population

structure and other covariates that influence gene expression, normalized expression phenotypes

were corrected with the first three genotype PCs, ten PEER (Stegle et al., 2012) factors (estimated

for each tissue of interest; Figure S5), the age of the individual, and RIN value of the sample. We

considered sequence variants with MAF > 1% and included variants 1 Mb up- and downstream of the

transcription start site (TSS). To account for multiple testing, we used the "permute" function in

QTLtools to conduct 1000 permutations and produce beta corrected p-values. We used the adjusted

p-values and the qtltools_runFDR_cis.R script to apply a false discovery rate (FDR) threshold of 5%

and identify expressed genes with at least one significant eQTL. To identify all significant variant-

gene pairs, we followed the procedure detailed in Delaneau et al., (2017). Specifically, each gene's

beta p-values and most-likely beta parameters were used with the FDR level to feed the beta

quantile function. This generated nominal p-value thresholds for each expressed gene, and all

variants below this threshold were deemed significant. To identify genes with multiple independent

acting eQTL, we implemented the conditional analysis from QTLtools. Briefly, this process uses the

aforementioned nominal p-value thresholds to identify all significant eVariants, then implements

forward and backward stepwise linear regression to identify independent acting eQTL. The

significant eVariants are then assigned to the independent signals.

To estimate the effect size of significant eQTL, we computed the log allelic fold change (slope_aFC;

Mohammadi et al., 2017) with aFC.py (https://github.com/secastel/aFC) for the top variant of each

independent eQTL. We used gene counts from featureCounts, which were normalized with DESeq2

(Love et al., 2014) and log transformed. We compared the normalized eQTL effect sizes to the

annotation category of the top variant (obtained with the Variant Effect Predictor (VEP) from

Ensembl; McLaren et al., 2016 et al., 2016) to investigate functional consequences.

We studied allelic imbalance using RNA sequencing reads filtered with WASP (Van De Geijn et al.,

2015) to mitigate reference allele bias. Whole genome DNA sequence data phased with beagle 5.4

(Browning et al., 2021) were complemented with the "read backed phasing" of the RNA seq data

using phASER (Castel et al., 2016) with the flag --baseq 10. Sequencing reads overlapping

heterozygous sites within protein coding genes (Ensembl version 104) were used to produce gene

level haplotypic expression counts using phASER_gene_AE.py with the –min_haplo_maf 0.005 flag. A

median number of 3,834, 3,345, and 1,353 RNA sequencing reads per gene, and 6,352,369,

8,453,661 and 6,739,602 reads per sample covering the heterozygous sites of the protein coding

genes were available in testis, epididymis, and vas deferens, respectively. Haplotype expression

counts across all samples were compiled using phaser_expr_matrix.py to validate cis-regulatory

effects of eQTL identified in the three tissues and to independently estimate their effects. eGenes

harboring heterozygous sites covered with at least 8 sequencing reads and top eVariants with at

least 10 homozygous and 10 heterozygous individuals were considered for this assessment, resulting

in a subset of 12,261, 3,832, and 2,871 variant-gene pairs in testis, epididymis, and vas deferens,

respectively. Gene level haplotypes were then phased with their associated top eVariant using

phaser_cis_var.py. Individual level magnitude of allelic imbalance was estimated as log allelic fold

change (AI_aFC) using phaser_cis_var.py. The AI_aFC is the ratio of gene haplotype counts linked to

alternative allele to the gene haplotype counts linked to the reference allele of the top eVariant on

log2 scale. The median of AI_aFC in individuals heterozygous for the top eVariant is presented as an

independent estimate of the cis regulatory effect of that eQTL. Comparison of AI_aFCs in individuals

heterozygous for the top eVariant allelic imbalance expected) and homozygous for the top eVariant

(allelic imbalance not expected) was used to test the significance of allelic imbalance using a

Wilcoxon ranksum test implemented in phaser_cis_var.py.

sQTL mapping was performed with QTLtools and generally followed the approach used for eQTL

discovery. However, we used the "grp-best" option to adjust the permutation scheme to correct for

multiple splice junctions and intron clusters within a gene. Mapping was conducted with variants 1

Mb up and downstream of the TSS and we included three genotype PCs, age, RIN, and ten PEER

factors as covariates. We conducted the QTLtools conditional analysis to identify all significant

variants (sVariants) and independent acting sQTL. We considered the β–coefficient from the

QTLtools linear model for the top sVariant as the sQTL effect size. Normalized sQTL effect sizes were

compared across VEP functional annotation categories using the previously described variant

annotation classes.

Trans eQTL and sQTL were only mapped for testis. We considered 5,685,508 variants that had MAF >

5%, were outside of repetitive regions (identified by RepeatMasker (v4.1.4);

http://www.repeatmasker.org), had imputation accuracy of 1.0, and had mappability > 1. Variant

mappability was determined with GenMap (v1.3.0; Pockrandt et al., 2020) using kmers with lengths

of 36 for UTRs and 75 for exons (based on the ARS-UCD1.2 reference) and allowed for two

mismatches. We mapped trans QTL for protein coding genes that had a mappability > 0.80 (18,876

genes). Gene mappability was estimated with crossmap (https://github.com/battle-lab/crossmap;

Saha & Battle, 2019). Phenotypes (either TPM values or PSI values) were corrected for the same

covariates used in the cis analysis and rank normal transformed. We used the "trans" function from

QTLtools to identify QTL > 5Mb from a gene's TSS and performed 100 permutations, which were

then ranked and used to estimate FDR. We considered a trans eVariant or sVariant significant if it

had an FDR < 0.05. We removed trans QTL that were cross-mapped (https://github.com/battle-

lab/crossmap; Saha & Battle, 2019). Due to excessive linkage observed in cattle, we removed trans

QTL with variants that had LD > 0.01 with eQTL or sQTL for those eGenes or sGenes, respectively.

Shared eQTL and sQTL

We identified shared and tissue specific QTL with Mashr (v0.2.69; Urbut et al., 2019). We included top eVariants and sVariants from genes that were expressed in all three tissues. To prevent the double-counting of molQTL that are in linkage disequilibrium and impact the same gene or intron cluster, we selected a single top molQTL that had the largest effect across tissues (as described in Urbut et al., 2019). This resulted in 11,311 eVariants and 5,012 sVariants. We considered a subset of 200,000 randomly selected variants and invoked a threshold of LFSR < 0.05 to establish if an effect was significant in a tissue. For significant effects, we assessed pairwise tissue similarity with Spearman's correlation. We compared the magnitude of molQTL effect sizes across specificity classes (tissue specific, in two tissues, or in all tissues) for each tissue with a Wilcoxon signed rank test.

GWAS of male fertility

Genome-wide association tests assuming additive and non-additive modes of inheritance were carried out between imputed genotypes and estimates of sire insemination success (which we used as a proxy for "male fertility"; Hiltpold et al., 2021) in 3,736 bulls using GCTA (Yang et al., 2011). The additive association testing was carried out using additively coded SNP genotypes. For the non-additive association testing, SNP genotypes were coded assuming recessive inheritance to the reference (R) allele and alternate (A) allele. Genotypes were coded as RR = 1, RA = 0, AA = 0 (M1) in the former case and as AA = 1, RA = 0, RR = 0 (M2) in the latter case. Only SNPs with MAF > 0.5% and where the frequency of 1's was between 0.5% and 0.95% (N = 14,587,856) were retained for association testing. The smallest P value from M1 and M2 was considered as the strength of non-additive association between genotypes and phenotypes for each of the tested variant, and variants with $P < 5e-08$ were considered significant. A genomic relationship matrix constructed from 18,187,234 autosomal SNPs with MAF > 0.005 was included in the association model to control for relatedness among samples.

Integrating QTL and molQTL in MetaXcan framework

We used S-PrediXcan (Barbeira et al., 2018) to test for an association between molecular phenotypes (expression and splicing phenotypes) and male fertility. The summary statistic-based approach was applied to include results from non-additive association testing. Briefly, we trained elastic-net based prediction models for expression values (TPM) and splicing phenotypes (PSI), respectively, for all genes and intronic clusters separately in testis, epididymis, and vas deferens.

SNPs located within 1 Mb of the TSS of the genes were considered for training with the script

gtex_tiss_chrom_training.R. In total, 18,053,560 sequence variants (MAF > 1%) called in the molQTL

cohort were available for training the prediction models. Covariates used in the eQTL and sQTL scans

were included in training of respective prediction models to account for confounding factors. High-

performance prediction models (zscore_pval < 0.05 and rho_avg > 0.1) were available to predict

expression levels of 15,810 genes and to predict splicing phenotypes for 46,426 intron clusters

across three tissues (77, 80, and 58% of eGenes and for 92, 94, and 80% of intron clusters in testis,

epididymis, and vas deferens respectively). Using weights for each SNPs in these high-performance

prediction models, we then carried out gene level and cluster level association test for expression

and splicing phenotype respectively using the script SPrediXcan.py. Following the recommendations

in Barbeira et al., (2018), Bonferroni-corrected P values of 3.16e-06 (0.05/15,810) and 1.08e-06

(0.05/46,426), respectively considering the number of genes and intron clusters tested for

expression and splicing phenotypes, were set as a significant threshold. Further, a filter was applied

on the prediction performance of the model with the threshold set at 0.05/number of significant

results after the previous filtering.

Next, to avoid capturing LD-contaminated associations, as recommended in the MetaXcan

framework (Barbeira et al., 2018), we carried out Bayes Factor colocalization analyses with the R

package coloc (Giambartolomei et al., 2014). For every significant GWAS signal (p < 5e-08), we

searched for evidence of its colocalization with molQTL using the "coloc.abf" function. The search

space was limited to 5 Mb up- and downstream of the significant GWAS peaks. Colocalizations with

PP.H4.abf (probability of a shared causal variant) > 0.8 and PP.H3.abf (probability of different causal

variants) < 0.5 were retained and used to filter TWAS results.

Characterization of a SPATA16 sQTL

The Ensembl annotation of *SPATA16* contains only one transcript, whereas the Refseq annotation

suggests three isoforms (XM_002684936.6, XM_024995427.1, XM_005201702.4) that differ in both

the number of coding and the number of non-coding exons. The abundance of the three isoforms

was quantified with kallisto (Bray et al., 2016). Testis RNA sequencing data was processed and

aligned to the cattle reference genome (ARS-UCD1.2) and the Refseq gene annotation (version 106)

with STAR (see above). Exon-specific SPATA16 expression was quantified with QTLtools "quan" on

the STAR-aligned bam files (see above). Association tests between exon expression and

Chr1:93,954,637 were conducted with a linear model in R that considered three PCs, age, and RIN as

covariates. Bonferroni correction was applied to determine a significance threshold with 0.05/n,

where n is the number of exons tested.

Data availability

DNA and RNA sequencing data of 118 bulls are available in the ENA database at the study accessions PRJEB28191 and PRJEB46995. Gene expression and splicing matrices, a VCF file of genome-wide genotypes used for e/sQTL mapping, a cross-table to link genotype and transcriptome data, as well as summary statistics used for transcriptome-wide association testing have been archived at zenodo (https://doi.org/10.5281/zenodo.8093287).

Author contributions

XMM sampled tissue and established the molQTL cohort, aligned RNA sequencing data, quantified gene expression and splicing variation, developed workflows and applied them to map e/sQTL, contributed to the TWAS, interpreted results, and drafted the manuscript; NKK imputed sequence variant genotypes, established workflows for GWAS, conducted ASE analyses and TWAS, contributed to eQTL analysis, interpreted results, and contributed to the writing of the manuscript; NKK and XMM performed colocalization analyses; MH coordinated the sampling of tissue with the abattoir staff, sampled tissue, and prepared phenotype data; ALV contributed to short read alignment; ASL performed variant discovery and genotyping, and contributed to e/sQTL mapping; QH contributed to GWAS; MB performed the WASP alignment of the RNA-seq data and contributed to discussions about the molQTL mapping workflows; HP conceptualized the study, contributed to the molQTL fine-mapping, interpreted results, and contributed to the writing of the manuscript; all authors read and approved the final version of the manuscript.

Acknowledgements

We appreciate support from the abattoir veterinarians and staff as well as Till Graf, Patrick Fitzi, and Remo Hengartner for assistance with tissue sampling. We thank Dr. Fritz-Schmitz Hsu and Dr. Ulrich Witschi (Swissgenetics) for providing phenotype data. We thank Braunvieh Schweiz for providing genotype data of Swiss BSW bulls. We acknowledge the Arbeitsgemeinschaft Deutsches Braunvieh, Braunvieh Austria, Tierzuchtforschung Grub, the Chair of Animal Breeding of TU München, the Institute of Animal Breeding from Bayerische Landesanstalt für Landwirtschaft and ZuchtData EDV Dienstleistungen GmbH for providing genotype and phenotype data of Austrian and German BSW bulls. We thank the Functional Genomics Center Zurich (Dr. Catharine Aquino) for generating DNA and RNA sequencing data. We thank Dr. Jeremie Vandenplas (Wageningen University & Research, Wageningen, the Netherlands) for providing a modified version of the GCTA tool that allows considering allele dosage data.

Funding

This study was supported by grants from the Swiss National Science Foundation, an ETH Research

Grant, Swissgenetics, and the European Union's Horizon 2020 research and innovation programme

under Grant Agreement No. 815668 (BovReg). The funding bodies were neither involved in the

design of the study and collection, analysis, and interpretation of data nor in writing the manuscript.

References

Aguet, F., Brown, A. A., Castel, S. E., Davis, J. R., He, Y., Jo, B., Mohammadi, P., Park, Y. S., Parsana, P.,

Segrè, A. V., Strober, B. J., Zappala, Z., Cummings, B. B., Gelfand, E. T., Hadley, K., Huang, K. H.,

Lek, M., Li, X., Nedzel, J. L., … Zhu, J. (2017). Genetic effects on gene expression across human

tissues. *Nature*, *550*(7675), 204–213. https://doi.org/10.1038/nature24277

Barbeira, A. N., Dickinson, S. P., Bonazzola, R., Zheng, J., Wheeler, H. E., Torres, J. M., Torstenson, E.

S., Shah, K. P., Garcia, T., Edwards, T. L., Stahl, E. A., Huckins, L. M., Aguet, F., Ardlie, K. G.,

Cummings, B. B., Gelfand, E. T., Getz, G., Hadley, K., Handsaker, R. E., … Im, H. K. (2018).

Exploring the phenotypic consequences of tissue specific gene expression variation inferred

from GWAS summary statistics. *Nature Communications*, *9*(1), 1–20.

https://doi.org/10.1038/s41467-018-03621-1

Bray, N. L., Pimentel, H., Melsted, P., & Pachter, L. (2016). Near-optimal probabilistic RNA-seq

quantification. *Nature Biotechnology*, *34*(5), 525–527. https://doi.org/10.1038/nbt.3519

Browning, B. L., Tian, X., Zhou, Y., & Browning, S. R. (2021). Fast two-stage phasing of large-scale

sequence data. *American Journal of Human Genetics*, *108*(10), 1880–1890.

https://doi.org/10.1016/j.ajhg.2021.08.005

Butler, M. L., Bormann, J. M., Weaber, R. L., Grieger, D. M., & Rolf, M. M. (2020). Selection for bull

fertility: A review. *Translational Animal Science*, *4*(1), 423–441.

https://doi.org/10.1093/TAS/TXZ174

Castel, S. E., Mohammadi, P., Chung, W. K., Shen, Y., & Lappalainen, T. (2016). Rare variant phasing

and haplotypic expression from RNA sequencing with phASER. *Nature Communications*, *7*.

https://doi.org/10.1038/ncomms12817

Chang, C. C., Chow, C. C., Tellier, L. C. A. M., Vattikuti, S., Purcell, S. M., & Lee, J. J. (2015). Second-

generation PLINK: Rising to the challenge of larger and richer datasets. *GigaScience*, *4*(1).

https://doi.org/10.1186/s13742-015-0047-8

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). Fastp: An ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics, *34*(17), i884–i890. https://doi.org/10.1093/bioinformatics/bty560

Cotto, K. C., Feng, Y.-Y., Ramu, A., Richters, M., Freshour, S. L., Skidmore, Z. L., Xia, H., McMichael, J.

F., Kunisaki, J., Campbell, K. M., Chen, T. H.-P., Rozycki, E. B., Adkins, D., Devarakonda, S.,

Sankararaman, S., Lin, Y., Chapman, W. C., Maher, C. A., Arora, V., … Griffith, M. (2023).

Integrated analysis of genomic and transcriptomic data for the discovery of splice-associated

variants in cancer. *Nature Communications*, *14*(1), 1589. https://doi.org/10.1038/s41467-023-

37266-6

Dam, A. H. D. M., Koscinski, I., Kremer, J. A. M., Moutou, C., Jaeger, A. S., Oudakker, A. R., Tournaye,

H., Charlet, N., Lagier-Tourenne, C., Van Bokhoven, H., & Viville, S. (2007). Homozygous

mutation in SPATA16 is associated with male infertility in human globozoospermia. *American*

Journal of Human Genetics, *81*(4), 813–820. https://doi.org/10.1086/521314

Delaneau, O., Ongen, H., Brown, A. A., Fort, A., Panousis, N. I., & Dermitzakis, E. T. (2017). A

complete tool set for molecular QTL discovery and analysis. *Nature Communications*, *8*.

https://doi.org/10.1038/ncomms15452

Djureinovic, D., Fagerberg, L., Hallström, B., Danielsson, A., Lindskog, C., Uhlén, M., & Pontén, F.

(2014). The human testis-specific proteome defined by transcriptomics and antibody-based

profiling. *Molecular Human Reproduction*, *20*(6), 476–488.

https://doi.org/10.1093/molehr/gau018

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., &

Gingeras, T. R. (2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*, *29*(1), 15–21.

https://doi.org/10.1093/bioinformatics/bts635

Druet, T., Fritz, S., Sellem, E., Basso, B., Gérard, O., Salas-Cortes, L., Humblot, P., Druart, X., & Eggen,

A. (2009). Estimation of genetic parameters and genome scan for 15 semen characteristics

traits of Holstein bulls. *Journal of Animal Breeding and Genetics*, *126*(4), 269–277.

https://doi.org/10.1111/j.1439-0388.2008.00788.x

Fujihara, Y., Oji, A., Larasati, T., Kojima-Kita, K., & Ikawa, M. (2017). Human globozoospermia-related

gene spata16 is required for sperm formation revealed by CRISPR/Cas9-mediated mouse

models. *International Journal of Molecular Sciences*, *18*(10).

https://doi.org/10.3390/ijms18102208

Gamble, J., Chick, J., Seltzer, K., Graber, J. H., Gygi, S., Braun, R. E., & Snyder, E. M. (2020). An

expanded mouse testis transcriptome and mass spectrometry defines novel proteins.

Reproduction, 159(1), 15–26. https://doi.org/10.1530/REP-19-0092

Garrido-Martín, D., Borsari, B., Calvo, M., Reverter, F., & Guigó, R. (2021). Identification and analysis

of splicing quantitative trait loci across multiple tissues in the human genome. *Nature*

Communications, *12*(1). https://doi.org/10.1038/s41467-020-20578-2

864 Gebreyesus, G., Lund, M. S., Kupisiewicz, K., & Su, G. (2021). Genetic parameters of semen quality
865   traits and genetic correlations with service sire nonreturn rate in Nordic Holstein bulls. *Journal*
866   *of Dairy Science*, *104*(9), 10010–10019. https://doi.org/10.3168/jds.2021-20403

867 Giambartolomei, C., Vukcevic, D., Schadt, E. E., Franke, L., Hingorani, A. D., Wallace, C., & Plagnol, V.
868   (2014). Bayesian Test for Colocalisation between Pairs of Genetic Association Studies Using
869   Summary Statistics. *PLoS Genetics*, *10*(5). https://doi.org/10.1371/journal.pgen.1004383

870 GTEx Consortium. (2020). The GTEx Consortium atlas of genetic regulatory effects across human
871   tissues. *Science*, *369*(6509), 1318–1330. www.gtexportal.org

872 Guo, J., Grow, E. J., Mlcochova, H., Maher, G. J., Lindskog, C., Nie, X., Guo, Y., Takei, Y., Yun, J., Cai, L.,
873   Kim, R., Carrell, D. T., Goriely, A., Hotaling, J. M., & Cairns, B. R. (2018). The adult human testis
874   transcriptional cell atlas. *Cell Research, 28*(12), 1141–1157.

875 Guo, J., Nie, X., Giebler, M., Mlcochova, H., Wang, Y., Grow, E. J., Kim, R., Tharmalingam, M.,
876   Matilionyte, G., Lindskog, C., Carrell, D. T., Mitchell, R. T., Goriely, A., Hotaling, J. M., & Cairns,
877   B. R. (2020). The Dynamic Transcriptional Cell Atlas of Testis Development during Human
878   Puberty. *Cell Stem Cell*, *26*(2), 262-276.e4. https://doi.org/10.1016/j.stem.2019.12.005

879 Hiltpold, M., Kadri, N. K., Janett, F., Witschi, U., Schmitz-Hsu, F., & Pausch, H. (2021). Autosomal
880   recessive loci contribute significantly to quantitative variation of male fertility in a dairy cattle
881   population. *BMC Genomics*, *22*(1). https://doi.org/10.1186/s12864-021-07523-3

882 Hiltpold, M., Niu, G., Kadri, N. K., Crysnanto, D., Fang, Z. H., Spengeler, M., Schmitz-Hsu, F., Fuerst, C.,
883   Schwarzenbacher, H., Seefried, F. R., Seehusen, F., Witschi, U., Schnieke, A., Fries, R., Bollwein,
884   H., Flisikowski, K., & Pausch, H. (2020). Activation of cryptic splicing in bovine WDR19 is
885   associated with reduced semen quality and male fertility. *PLoS Genetics*, *16*(5).
886   https://doi.org/10.1371/journal.pgen.1008804

887 Kaessmann, H. (2010). Origins, evolution, and phenotypic impact of new genes. In *Genome Research*
888   (Vol. 20, Issue 10, pp. 1313–1326). https://doi.org/10.1101/gr.101386.109

889 Lamas-Toranzo, I., Hamze, J. G., Bianchi, E., Fernández-Fuertes, B., Pérez-Cerezales, S., Laguna-
890   Barraza, R., Fernández-González, R., Lonergan, P., Gutiérrez-Adán, A., Wright, G. J., Jiménez-
891   Movilla, M., & Bermejo-álvarez, P. (2020). TMEM95 is a sperm membrane protein essential for
892   mammalian fertilization. *ELife*, *9*, 1–18. https://doi.org/10.7554/eLife.53913

893 Leonard, A. S., Crysnanto, D., Mapel, X. M., Bhati, M., & Pausch, H. (2023). Graph construction
894   method impacts variation representation and analyses in a bovine super-pangenome. *Genome*
895   *Biology*, *24*(1), 124. https://doi.org/10.1186/s13059-023-02969-y

Leonard, A. S., Mapel, X. M., & Pausch, H. (2023). Pangenome genotyped structural variation

improves molecular phenotype mapping in cattle. *BioRxiv*.

https://doi.org/10.1101/2023.06.21.545879

Li, H. (2013). *Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM*.

http://arxiv.org/abs/1303.3997

Li, Y. I., Knowles, D. A., Humphrey, J., Barbeira, A. N., Dickinson, S. P., Im, H. K., & Pritchard, J. K.

(2018). Annotation-free quantification of RNA splicing using LeafCutter. *Nature Genetics*, *50*(1),

151–158. https://doi.org/10.1038/s41588-017-0004-9

Li, Y. I., Van De Geijn, B., Raj, A., Knowles, D. A., Petti, A. A., Golan, D., Gilad, Y., & Pritchard, J. K.

(2016). RNA splicing is a primary link between genetic variation and disease. *Science*,

352(6285), 600–604. https://www.science.org

Liao, Y., Smyth, G. K., & Shi, W. (2014). FeatureCounts: An efficient general purpose program for

assigning sequence reads to genomic features. *Bioinformatics*, *30*(7), 923–930.

https://doi.org/10.1093/bioinformatics/btt656

Liu, S., Gao, Y., Canela-Xandri, O., Wang, S., Yu, Y., Cai, W., Li, B., Xiang, R., Chamberlain, A. J., Pairo-

Castineira, E., D'Mellow, K., Rawlik, K., Xia, C., Yao, Y., Navarro, P., Rocha, D., Li, X., Yan, Z., Li,

C., … Fang, L. (2022). A multi-tissue atlas of regulatory variants in cattle. *Nature Genetics*, *54*(9),

1438–1447. https://doi.org/10.1038/s41588-022-01153-5

Liu, S., Ma, X., Wang, Z., Lin, F., Li, M., Li, Y., Yang, L., Rushdi, H. E., Riaz, H., Gao, T., Yang, L., Fu, T., &

Deng, T. (2023). MAEL gene contributes to bovine testicular development through the m5C-

mediated splicing. *IScience*, *26*(2). https://doi.org/10.1016/j.isci.2023.105941

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biology*, *15*(12). https://doi.org/10.1186/s13059-014-

0550-8

McLaren, W., Gil, L., Hunt, S. E., Riat, H. S., Ritchie, G. R. S., Thormann, A., Flicek, P., & Cunningham,

F. (2016). The Ensembl Variant Effect Predictor. *Genome Biology*, *17*(1).

https://doi.org/10.1186/s13059-016-0974-4

Mohammadi, P., Castel, S. E., Brown, A. A., & Lappalainen, T. (2017). Quantifying the regulatory

effect size of cis-acting genetic variation using allelic fold change. *Genome Research*, *27*(11),

1872–1884. https://doi.org/10.1101/gr.216747.116

Murat, F., Mbengue, N., Winge, S. B., Trefzer, T., Leushkin, E., Sepp, M., Cardoso-Moreira, M.,

Schmidt, J., Schneider, C., Mößinger, K., Brüning, T., Lamanna, F., Belles, M. R., Conrad, C.,

Kondova, I., Bontrop, R., Behr, R., Khaitovich, P., Pääbo, S., … Kaessmann, H. (2023). The

929 molecular evolution of spermatogenesis across mammals. *Nature*, *613*(7943), 308–316.

930 https://doi.org/10.1038/s41586-022-05547-7

931 Nagai, R., Kinukawa, M., Watanabe, T., Ogino, A., Kurogi, K., Adachi, K., Satoh, M., & Uemoto, Y.

932 (2022). Genome-wide detection of non-additive quantitative trait loci for semen production

933 traits in beef and dairy bulls. *Animal*, *16*(3). https://doi.org/10.1016/j.animal.2022.100472

934 Oura, S., Koyano, T., Kodera, C., Yuki Horisawa-Takada, Matsuyama, M., Ishiguro, K. I., & Ikawa, M.

935 (2021). KCTD19 and its associated protein ZFP541 are independently essential for meiosis in

936 male mice. *PLoS Genetics*, *17*(5 May). https://doi.org/10.1371/journal.pgen.1009412

937 Pausch, H., Kölle, S., Wurmser, C., Schwarzenbacher, H., Emmerling, R., Jansen, S., Trottmann, M.,

938 Fuerst, C., Götz, K. U., & Fries, R. (2014). A Nonsense Mutation in TMEM95 Encoding a

939 Nondescript Transmembrane Protein Causes Idiopathic Male Subfertility in Cattle. *PLoS*

940 *Genetics*, *10*(1). https://doi.org/10.1371/journal.pgen.1004044

941 Pausch H, Mapel XM. Review: Genetic mutations affecting bull fertility. *Animal*, in press

942 Pirinen, M., Lappalainen, T., Zaitlen, N. A., Dermitzakis, E. T., Donnelly, P., McCarthy, M. I., & Rivas,

943 M. A. (2015). Assessing allele-specific expression across multiple tissues from RNA-seq read

944 data. *Bioinformatics*, *31*(15), 2497–2504. https://doi.org/10.1093/bioinformatics/btv074

945 Pockrandt, C., Alzamel, M., Iliopoulos, C. S., & Reinert, K. (2020). GenMap: Ultra-fast computation of

946 genome mappability. *Bioinformatics, 36*(12), 3687–3692.

947 https://doi.org/10.1093/bioinformatics/btaa222

948 Poplin, R., Chang, P. C., Alexander, D., Schwartz, S., Colthurst, T., Ku, A., Newburger, D., Dijamco, J.,

949 Nguyen, N., Afshar, P. T., Gross, S. S., Dorfman, L., McLean, C. Y., & Depristo, M. A. (2018). A

950 universal snp and small-indel variant caller using deep neural networks. *Nature Biotechnology*,

951 *36*(10), 983. https://doi.org/10.1038/nbt.4235

952 Ramsköld, D., Wang, E. T., Burge, C. B., & Sandberg, R. (2009). An abundance of ubiquitously

953 expressed genes revealed by tissue transcriptome sequence data. *PLoS Computational Biology*,

954 *5*(12). https://doi.org/10.1371/journal.pcbi.1000598

955 Rezende, F. M., Dietsch, G. O., & Peñagaricano, F. (2018). Genetic dissection of bull fertility in US

956 Jersey dairy cattle. *Animal Genetics*, *49*(5), 393–402. https://doi.org/10.1111/age.12710

957 Robertson, M. J., Kent, K., Tharp, N., Nozawa, K., Dean, L., Mathew, M., Grimm, S. L., Yu, Z., Légaré,

958 C., Fujihara, Y., Ikawa, M., Sullivan, R., Coarfa, C., Matzuk, M. M., & Garcia, T. X. (2020). Large-

959 scale discovery of male reproductive tract-specific genes through analysis of RNA-seq datasets.

960 *BMC Biology, 18*(1). https://doi.org/10.1186/s12915-020-00826-z

961 Rosen, B. D., Bickhart, D. M., Schnabel, R. D., Koren, S., Elsik, C. G., Tseng, E., Rowan, T. N., Low, W.

962 Y., Zimin, A., Couldrey, C., Hall, R., Li, W., Rhie, A., Ghurye, J., McKay, S. D., Thibaud-Nissen, F.,

Hoffman, J., Murdoch, B. M., Snelling, W. M., … Medrano, J. F. (2020). De novo assembly of the

cattle reference genome with single-molecule sequencing. *GigaScience*, *9*(3).

https://doi.org/10.1093/gigascience/giaa021

Saha, A. & Battle, A. (2019) False positives in trans-eQTL and co-expression analyses arising from

RNA-sequencing alignment errors [version 2; peer review: 3 approved]. *F1000Research*, 7:1860

https://doi.org/10.12688/f1000research.17145.2

Smit, AFA., Hubley, R. & Green, P. *RepeatMasker Open-4.0*. 2013-2015.

http://www.repeatmasker.org

Soumillon, M., Necsulea, A., Weier, M., Brawand, D., Zhang, X., Gu, H., Barthès, P., Kokkinaki, M.,

Nef, S., Gnirke, A., Dym, M., deMassy, B., Mikkelsen, T. S., & Kaessmann, H. (2013). Cellular

Source and Mechanisms of High Transcriptome Complexity in the Mammalian Testis. *Cell*

Reports, *3*(6), 2179–2190. https://doi.org/10.1016/j.celrep.2013.05.031

Stegle, O., Parts, L., Piipari, M., Winn, J., & Durbin, R. (2012). Using probabilistic estimation of

expression residuals (PEER) to obtain increased power and interpretability of gene expression

analyses. *Nature Protocols*, *7*(3), 500–507. https://doi.org/10.1038/nprot.2011.457

Tang, S., Lu, Y., Skinner, W. M., Sanyal, M., Lishko, P. V, Ikawa, M., & Kim, P. S. (2022). Human sperm

TMEM95 binds eggs and facilitates membrane fusion*. Biophysics and computational biology*

open access. https://doi.org/10.1073/pnas

Tarasov, A., Vilella, A. J., Cuppen, E., Nijman, I. J., & Prins, P. (2015). Sambamba: fast processing of

NGS alignment formats. *Bioinformatics*, *31*(12), 2032–2034.

https://doi.org/10.5281/zenodo.13200

The FarmGTEx-PigGTEx Consortium, Teng, J., Gao, Y., Yin, H., Bai, Z., Liu, S., Zeng, H., Bai, L., Cai, Z.,

Zhao, B., Xu, Z., Lin, Q., Pan, Z., Yang, W., Yu, X., Hou, Y., Keel, B. N., Rohrer, G. A., Lindholm-

Perry, A. K., … Fang, L. (2022). A compendium of genetic regulatory effects across pig tissues.

BioRxiv. https://doi.org/10.1101/2022.11.11.516073

Thibier, M., & Wagner, H.-G. (2002). World statistics for artificial insemination in cattle. In Livestock

Production Science (Vol. 74). www.elsevier.com/locate/livprodsci

Tüttelmann, F., Ruckert, C., & Röpke, A. (2018). Spermatogenesestörungen: Perspektiven für

erweiterte genetische Diagnostik nach 20 Jahren unveränderter Routine. Medizinische Genetik,

30(1), 12–20. https://doi.org/10.1007/s11825-018-0181-7

Urbut, S. M., Wang, G., Carbonetto, P., & Stephens, M. (2019). Flexible statistical methods for

estimating and testing effects in genomic studies with multiple conditions. *Nature Genetics*,

51(1), 187–195. https://doi.org/10.1038/s41588-018-0268-8

Van De Geijn, B., Mcvicker, G., Gilad, Y., & Pritchard, J. K. (2015). WASP: Allele-specific software for
robust molecular quantitative trait locus discovery. In *Nature Methods* (Vol. 12, Issue 11, pp.
1061–1063). Nature Publishing Group. https://doi.org/10.1038/nmeth.3582
Wainberg, M., Sinnott-Armstrong, N., Mancuso, N., Barbeira, A. N., Knowles, D. A., Golan, D., Ermel,
R., Ruusalepp, A., Quertermous, T., Hao, K., Björkegren, J. L. M., Im, H. K., Pasaniuc, B., Rivas,
M. A., & Kundaje, A. (2019). Opportunities and challenges for transcriptome-wide association
studies. *Nature Genetics*, *51*(4), 592–599. https://doi.org/10.1038/s41588-019-0385-z
Walker, R. L., Ramaswami, G., Hartl, C., Mancuso, N., Gandal, M. J., de la Torre-Ubieta, L., Pasaniuc,
B., Stein, J. L., & Geschwind, D. H. (2019). Genetic Control of Expression and Splicing in
Developing Human Brain Informs Disease Mechanisms. *Cell*, *179*(3), 750-771.e22.
https://doi.org/10.1016/j.cell.2019.09.021
Wang, S., Qiao, H., Wang, P., Wang, Y., & Qin, D. (2021). ZDHHC19 is dispensable for
spermatogenesis, but is essential for sperm functions in mice. *International Journal of*
Molecular Sciences, *22*(16). https://doi.org/10.3390/ijms22168894
Wang, W., Su, L., Meng, L., He, J., Tan, C., Yi, D., Cheng, D., Zhang, H., Lu, G., Du, J., Lin, G., Zhang, Q.,
Tu, C., & Tan, Y.-Q. (2023). Biallelic variants in KCTD19 associated with male factor infertility
and oligoasthenoteratozoospermia. *Human Reproduction*.
https://doi.org/10.1093/humrep/dead095
Wu, Y., Broadaway, K. A., Raulerson, C. K., Scott, L. J., Pan, C., Ko, A., He, A., Tilford, C., Fuchsberger,
C., Locke, A. E., Stringham, H. M., Jackson, A. U., Narisu, N., Kuusisto, J., Pajukanta, P., Collins, F.
S., Boehnke, M., Laakso, M., Lusis, A. J., … Mohlke, K. L. (2019). Colocalization of GWAS and
eQTL signals at loci with multiple signals identifies additional candidate genes for body fat
distribution. *Human Molecular Genetics*, *28*(24), 4161–4172.
https://doi.org/10.1093/hmg/ddz263
Xavier, M. J., Salas-Huetos, A., Oud, M. S., Aston, K. I., & Veltman, J. A. (2021). Disease gene
discovery in male infertility: past, present and future. In Human Genetics (Vol. 140, Issue 1, pp.
7–19). Springer Science and Business Media Deutschland GmbH.
https://doi.org/10.1007/s00439-020-02202-x
Xiang, R., Fang, L., Liu, S., Macleod, I. M., Liu, Z., Breen, E. J., Gao, Y., Liu, G. E., Tenesa, A.,
Consortium, C., Mason, B. A., Chamberlain, A. J., Wray, N. R., & Goddard, M. E. (2022). Gene
expression and RNA splicing explain large proportions of the heritability for complex traits in
cattle. *BioRxiv*. https://doi.org/10.1101/2022.05.30.494093
Yamaguchi, K., Ishigaki, K., Suzuki, A., Tsuchida, Y., Tsuchiya, H., Sumitomo, S., Nagafuchi, Y., Miya, F.,
Tsunoda, T., Shoda, H., Fujio, K., Yamamoto, K., & Kochi, Y. (2022). Splicing QTL analysis

focusing on coding sequences reveals mechanisms for disease susceptibility loci. *Nature*

Communications, *13*(1). https://doi.org/10.1038/s41467-022-32358-1

Yang, J., Lee, S. H., Goddard, M. E., & Visscher, P. M. (2011). GCTA: A tool for genome-wide complex

trait analysis. *American Journal of Human Genetics*, *88*(1), 76–82.

https://doi.org/10.1016/j.ajhg.2010.11.011

Yun, T., Li, H., Chang, P. C., Lin, M. F., Carroll, A., & McLean, C. Y. (2020). Accurate, scalable cohort

variant calls using DeepVariant and GLnexus. *Bioinformatics*, *36*(24), 5582–5589.

https://doi.org/10.1093/bioinformatics/btaa1081

Zhang, Y., Liu, Z., Yun, X., Batu, B., Yang, Z., Zhang, X., Zhang, W., & Liu, T. (2023). Transcriptome

Profiling of Developing Testes and First Wave of Spermatogenesis in the Rat. *Genes, 14*(1).

https://doi.org/10.3390/genes14010229

1041